Base editors for citrus gene editing

Xiaoen Huang, Yuanchun Wang, Nian Wang*

Citrus Research and Education Center, Department of Microbiology and Cell Science, Institute

of Food and Agricultural Sciences, University of Florida.

*Correspondence: Nian Wang (nianwang@ufl.edu)

Abstract

Base editors, such as adenine base editors (ABE) and cytosine base editors (CBE), provide

alternatives for precise genome editing without generating double-strand breaks (DSBs), thus

avoiding the risk of genome instability and unpredictable outcomes caused by DNA repair.

Precise gene editing mediated by base editors in citrus has not been reported. Here, we have

successfully adapted the ABE to modify the TATA box in the promoter region of the canker

susceptibility gene LOB1 from TATA to CACA in grapefruit and Hamlin sweet orange.

Inoculation of the TATA-edited plants with the canker pathogen Xanthomonas citri subsp. citri

(*Xcc*) demonstrated that the TATA-edited plants were resistant to *Xcc*. In addition, CBE was

successfully used to edit the *acetolactate synthase (ALS)* gene of Carrizo citrange, a hybrid of

Citrus sinensis 'Washington' sweet orange X *Poncirus trifoliata*. Editing the ALS genes

conferred resistance of Carrizo to the herbicide chlorsulfuron. Two *ALS*-edited Carrizo plants did

not show green florescence although the starting construct for transformation contains a GFP

expression cassette. We performed PCR amplification for *Cas9* gene in the mutant plants and

found that *Cas9* gene was undetectable in the herbicide resistant citrus plants. This indicates that

the *ALS* edited plants are transgene-free, representing the first transgene-free gene-edited citrus

using the CRISPR technology. In summary, we have successfully adapted the base editors for

precise citrus gene editing. The CBE base editor has been used to generate transgene-free citrus

via transient expression.

Nickase Cas9 (nCas9) derived base editors, such as adenine base editors (ABE) and cytosine

base editors (CBE), are a useful tool for precise genome editing (Kluesner et al., 2021). Precise

gene editing mediated by base editors in citrus has not been reported. Here, we first tested if

ABE, such as ABE8e which can mediate A•T-to-G•C base changes (Richter et al., 2020), can

edit citrus genes. A citrus optimized ABE8e vector was constructed (Figure 1a). TadA8e

(Richter et al., 2020) was fused with Cas9^{D10A} nickase at the N-terminal. There is a nucleus

localization signal (NLS) at each end of TadA8e- Cas9^{D10A} for nucleus transportation. NLS-

TadA8e-Cas9^{D10A}-NLS is driven by the CmYLCV promoter, which confers high gene editing

efficiency in citrus. We aimed to edit the TATA box (Figure 1b) located upstream of the effector

binding elements (EBE) of the promoter of the citrus canker susceptible (S) gene *LOB1* with

ABE8e. The EBE in the promoter of susceptibility genes usually overlaps with or locates

immediately downstream of the TATA box of the S gene promoter (Huang et al., 2017). PthA4,

a TAL effector from citrus canker pathogen *Xanthomonas citri* subsp. *citri* (Xcc), binds to the

EBE region of *LOB1* to activate its expression. The TATA box of the *LOB1* promoter overlaps

with the EBE region (Figure 1d). We hypothesized that editing of the EBE-associated TATA box

may abolish or reduce the induction of *LOB1* by *Xcc*, thus generating canker resistant citrus

plants. The transient assay (Figure 1c) through *Xcc*-facilitated agroinfiltration of citrus leaves

showed that the ABE construct edited the TATA box as anticipated (Figure 1d). The ABE

construct mutated T<u>A</u>T<u>A</u> to C<u>A</u>C<u>A</u> (from T<u>A</u>T<u>A</u> to T<u>G</u>T<u>G</u> for the complementary strand). Test

on another gene *CsTub* (Cs1g21050) through transient assay also demonstrated precise editing

(Figure 1e). Next, *Agrobacterium*-mediated stable transformation of grapefruit and Hamlin sweet

orange epicotyl tissues was conducted to edit the TATA box in the *LOB1* promoter of both

varieties. Two transgenic grapefruit and one transgenic sweet orange plants were obtained.

Genotyping showed the TATA box was successfully edited in the three transformed plants

(Figure 1f). The T<u>A</u>T<u>A</u> in the *LOB1* promoter was edited into C<u>A</u>C<u>A</u>. By using the ABE system,

we achieved 100% biallelic/homozygous editing of the TATA box in the *LOB1* promoter. After

micro-grafting, one transgenic grapefruit plant and one transgenic sweet orange plant survived.

Inoculation of the TATA-edited plants with *Xcc* demonstrated that the TATA-edited plants were

resistant to *Xcc* (Figure 1g). On the contrary, all plants were susceptible to *Xcc pthA4*:Tn5

dLOB2, which carries a designer TAL dLOB2 for *LOB2* induction to cause canker disease in

citrus plants. These results showed that base editor ABE8e can efficiently and precisely edit

citrus genome.

Next, we tested if CBE, which mediates C•G-to-T•A base changes, can edit citrus genes.

A citrus optimized CBE vector was constructed (Figure 1h). We chose APOBEC3A (A3A)

deaminase for citrus CBE considering its wide deamination window and high editing efficiency

(Zhang et al., 2020; Zong et al., 2018). This CBE system contains CsU6-tRNA-gRNA-scaffold

unit for multiplex editing. The final CBE construct also contains a GFP expression cassette for

direct visualization of transgene integration. We first determined if citrus *acetolactate synthase*

(ALS) gene can be edited with this CBE. Acetolactate synthase (ALS) is an enzyme required for

biosynthesis of some branched-chain amino acids, such as valine, leucine, and isoleucine. A

single amino acid mutation in *ALS* genes in diverse plants confers resistance to herbicides such

as chlorsulfuron (Kuang et al., 2020; Malabarba et al., 2021). The growth of Carrizo citrange, a

hybrid of *Citrus sinensis* 'Washington' sweet orange X *Poncirus trifoliata*, was completely

inhibited in the presence of chlorsulfuron (Figure 1i). We designed two gRNAs targeting the two

ALS alleles of Carrizo (Figure 1j). The transient expression result showed that the construct

edited both alleles of the *ALS* gene of Carrizo (Figure 1j). *Agrobacterium*-mediated stable

transformation of Carrizo epicotyl tissues was performed with herbicide chlorsulfuron selection

in culture media. Two chlorsulfuron-resistant plants grew on the chlorsulfuron containing

medium while most of the transformed epicotyl tissues did not grow (Figure 1k). Genotyping of

the chlorsulfuron-resistant plants showed that both alleles of the *ALS* gene were edited in the two

plants (Figure 1l and m). Intriguingly, these two plants did not show green florescence although

the starting construct for transformation contains a GFP expression cassette. We performed PCR

amplification for *Cas9* gene in the mutant plants and found that *Cas9* gene was undetectable in

the herbicide resistant citrus plants. This indicates that the *ALS* edited plants are transgene-free,

which were obtained from transient expression of the CBE construct, representing the first

transgene-free gene-edited citrus using the CRISPR technology.

In summary, we have successfully adapted the base editors for precise citrus gene editing.

The CBE base editor has been used to generate transgene-free citrus via transient expression.

Conflicts of interest

The authors declare no competing financial interests.

Author contributions

XH designed and performed the experiments. YW performed micro-grafting. XH and NW wrote

the paper. NW supervised the study.

References

Huang, R.Y., Hui, S.G., Zhang, M., Li, P., Xiao, J.H., Li, X.H., Yuan, M. and Wang, S.P. (2017)
A Conserved Basal Transcription Factor Is Required for the Function of Diverse TAL
Effectors in Multiple Plant Hosts. *Frontiers in Plant Science* **8**, 1919.
Kluesner, M.G., Lahr, W.S., Lonetree, C.L., Smeester, B.A., Claudio-Vazquez, P.N., Pitzen, S.P.,
Vignes, M.J., Lee, S.C., Bingea, S.P., Andrews, A.A., Webber, B.R., Moriarity, B.S., Qiu,
X., Slipek, N.J. and Pomeroy, E.J. (2021) CRISPR-Cas9 cytidine and adenosine base
editing of splice-sites mediates highly-efficient disruption of proteins in primary and
immortalized cells. *Nature Communications* **12**, 2437.
Kuang, Y.J., Li, S.F., Ren, B., Yan, F., Spetz, C., Li, X.J., Zhou, X.P. and Zhou, H.B. (2020)
Base-Editing-Mediated Artificial Evolution of OsALS1 In Planta to Develop Novel
Herbicide-Tolerant Rice Germplasms. *Molecular Plant* **13**, 565-572.
Liu, L., Kuang, Y.J., Yan, F., Li, S.F., Ren, B., Gosavi, G., Spetz, C., Li, X.J., Wang, X.F., Zhou,
X.P. and Zhou, H.B. (2021) Developing a novel artificial rice germplasm for
dinitroaniline herbicide resistance by base editing of OsTubA2. *Plant Biotechnology*
Journal **19**, 5-7.
Malabarba, J., Chevreau, E., Dousset, N., Veillet, F., Moizan, J. and Vergne, E. (2021) New
Strategies to Overcome Present CRISPR/Cas9 Limitations in Apple and Pear: Efficient
Dechimerization and Base Editing. *International Journal of Molecular Sciences* **22**, 319.
Richter, M.F., Zhao, K.V.T., Eton, E., Lapinaite, A., Newby, G.A., Thuronyi, B.W., Wilson, C.,
Koblan, L.W., Zeng, J., Bauer, D.E., Doudna, J.A. and Liu, D.V.R. (2020) Phage-assisted
evolution of an adenine base editor with improved Cas domain compatibility and activity.
Nature Biotechnology **38**, 883-891.
Zhang, X.H., Chen, L., Zhu, B.Y., Wang, L.R., Chen, C.Y., Hong, M.J., Huang, Y.F., Li, H.Y.,
Han, H.H., Cai, B.L., Yu, W.S., Yin, S.M., Yang, L., Yang, Z.Z., Liu, M.Z., Zhang, Y.,
Mao, Z.Y., Wu, Y.X., Liu, M.Y. and Li, D.L. (2020) Increasing the efficiency and
targeting range of cytidine base editors through fusion of a single-stranded DNA-binding
protein domain. *Nature Cell Biology* **22**, 740-750.
Zong, Y., Song, Q.N., Li, C., Jin, S., Zhang, D.B., Wang, Y.P., Qiu, J.L. and Gao, C.X. (2018)
Efficient C-to-T base editing in plants using a fusion of nCas9 and human APOBEC3A.
Nature Biotechnology **36**, 950-953.

Figure legend

Figure 1 Highly efficient base editors for citrus gene editing. (a) Illustration of ABE8e adenine

base editing (ABE) system for citrus. CmYLCV, *Cestrum yellow leaf curling virus* promoter;

TadA8e, evolved *Escherichia coli* tRNA adenosine deaminase; Cas9^{D10A}, Cas9 nickase. (b) The

TATA box located upstream of the *LOB1* EBE region is associated with general transcription

factors and TAL effector PthA4. (c) *Xcc*-facilitated transient expression of ABE8e construct in

citrus leaf. The construct carries a GFP expression cassette. (d) Editing of TATA in the TATA

box of the *LOB1* promoter into CACA in transient assay from (c). Underlined nucleotides were

the target region for gRNA design. (e) Editing of the *CsTub* gene using transient assay. (f)

Chromatograms for the TATA-edited grapefruit mutant line 1 (upper), line 2 (middle) and

Hamlin sweet orange mutant line (lower). (g) Inoculation of WT and TATA-edited mutants with

Xcc or *Xcc pthA4*:Tn5 dLOB2. *Xcc pthA4*:Tn5 is a Tn5 insertion mutant of *pthA4* gene of *Xcc*.

Leaf area 1 inoculated with *Xcc*; leaf area 2 inoculated with *Xcc pthA4*:Tn5 dLOB2. (h)

Illustration of cytosine base editing system (CBE). (i) Growth of citrus seedlings (Carrizo

citrange) was inhibited by herbicide chlorsulfuron. (j) CBE base editing of the *ALS* gene via

transient assay. The restriction enzyme *Dra*II-resistant PCR amplicon was subject to cloning and

sequencing. (k) Selection of herbicide resistant Carrizo citrange on chlorsulfuron-containing

media (150 nM). (l) Sequencing results from chlorsulfuron-resistant mutant plants. (m)

Chromatograms for (l). (n) PCR of *Cas9* and *ALS* for control plant (GFP positive) and herbicide

chlorsulfuron-resistant citrus plants.