Title: *Antennapedia* regulates metallic silver wing scale development and cell

shape in *Bicyclus anynana* butterflies

Authors: Anupama Prakash[1*], Cédric Finet[2], Vinodkumar Saranathan[2], Antónia Monteiro[1,2*]

Affiliations:

[1] Department of Biological Sciences, National University of Singapore, Singapore.

[2] Division of Science, Yale-NUS College, National University of Singapore, Singapore.

*Correspondence to: anupama@u.nus.edu or antonia.monteiro@nus.edu.sg

Abstract: Butterfly wing scale cells can develop very intricate cuticular nanostructures that

interact with light to produce structural colors such as silver, but the genetic basis of such

nanostructures is mostly unexplored. Here, we address the genetic basis of metallic silver scale

development by leveraging existing crispants in the butterfly *Bicyclus anynana*, where knockouts

of five genes – *apterous A*, *Ultrabithorax*, *doublesex*, *Antennapedia* and *optix* – either led to

ectopic gains or losses of silver scales. Most wildtype silver scales had low amounts of

pigmentation and exhibited a common ultrastructural modification for metallic broadband

reflectance, i.e., an undulatory air layer enclosed by an upper and lower lamina. Crispant brown

scales differed from wildtype silver scales via the loss of the continuous upper lamina, increased

lower lamina thickness, and increased pigmentation. The reverse was seen when brown scales

became silver. On the forewings, we identified *Antennapedia* as a high-level selector gene, acting

through *doublesex* to induce silver scale development in males and having a novel, post-embryonic

role in the determination of ridge and crossrib orientation and overall scale cell shape in both sexes.

We propose that *apterous A* and *Ultrabithorax* repress *Antennapedia* on the dorsal forewings and

ventral hindwings, respectively, thereby repressing silver scale development, whereas *apterous A*

activates the same GRN on the dorsal hindwings, promoting silver scales.

<u>Introduction</u>

Silver or gold colors in insect cuticles (1–3), fish scales (4), and the eyes of cephalopods (1, 4–6)

are all examples of naturally occurring broadband structural coloration. These colors/structures

serve multiple ecological functions such as in promoting vision, in serving as inter- or intra-

specific signals, or in thermoregulation (5, 7–12) and arise from the interaction of light with

specific classes of broadband reflectors found in the animal integument (13–15).

Broadband metallic reflectors are highly reflective across a broad range of wavelengths and are often thick structures. The most common type of reflector involves thin film or multi-layer interference reflectors made of alternating materials with different refractive indices and varying thicknesses (1, 16). Chirped multilayer reflectors, which vary the optical thickness of each layer systematically with depth, are found in the exocuticle of gold beetles (16, 17) or the endocuticle of pupal cases in some butterflies (3, 18, 19). When the optical thickness of each layer changes randomly with depth, like in the alternating layers of guanine crystals and cytoplasm in fish scales, they form chaotic multilayer stacks that have silver reflectances (6, 20). Due to the small difference in refractive indices of biological materials, broadband multilayer reflectors with high reflectivity usually require a minimum of 10-20 alternating layers of high- and low-refractive index materials, leading to very thick (tens of microns) multilayer reflectors (19).

In contrast, broadband metallic reflectors found in lepidopteran wing scales are anatomically constrained by the thickness of the scales. These metallic reflectors are ultra-thin, with an overall thickness of a few microns (10, 21–24). It has been suggested that broadband reflectance is achieved by additive color mixing that occurs due to local spatial variation or disorder in the scale ultrastructure (10, 21, 23–25). An essential modification of the basic scale Bauplan to produce such broadband reflectors appears to be the consistent presence of a contiguous upper lamina that closes the normally "open" windows seen in a typical scale (10, 21, 24, 26). This creates an undulatory air layer sandwiched by the lower and upper laminas whose thicknesses also spatially vary. The broadband metallic reflectors, seen in fossil (23, 27) and extant moths (22), and more basal springtails (28), however, utilize thin film interference from a single chitin layer, resulting from fused scales. The lower lamina of an open-type plan scale in butterflies is often also tuned to produce broadband colors (29, 30). Therefore, both the necessity and the optical role of the upper lamina and the air layer in broadband color generation are enigmatic.

The production of metallic broadband reflectors may also depend on the presence or absence of pigments embedded in the scale cuticle. For instance, the brown ground scales in *Hypolimnas salmacis* or the yellow scales in *Heliconius* butterflies have a closed upper lamina, but these scales do not exhibit broadband reflectance due to the high concentration of pigments (31, 32).

Furthermore, the retention of an upper lamina in a *yellow* mutant in *Bicyclus anynana* butterflies

(33), does not lead to metallic scales, either due to the presence of pigments or incorrect thicknesses

of both the laminas and air gap in these scales. These examples suggest that the GRN that creates

a metallic scale type must involve regulation of both scale ultrastructure and pigmentation. Yet,

despite numerous studies having addressed the structural origins of biological broadband

reflections across arthropods (3, 10, 19, 21, 23, 25) and its evolution within butterflies (24), the

optical basis, the genetic circuits, and developmental mechanisms that create these broadband

metallic reflectors remain sparse and unexplored.

We have begun to explore the genetic basis of silver scale development with a series of CRISPR

experiments performed in the nymphalid butterfly *B. anynana*. This species exhibits different types

of broadband reflecting metallic scales on both fore- and hindwings (34). These scales are usually

associated with the sex pheromone producing regions in males (the androconia), except for the

coupling scales near the base of the wings that are present in both sexes. Previously, the knockout

of five genes including *apterous A* (*apA*) (35), *Ultrabithorax* (*Ubx*) (36), *doublesex* (*dsx*) (34),

Antennapedia (*Antp*) (36) and *optix* (37) led to phenotypic effects on metallic scale development

in *B. anynana* (Fig 1A). Some knockouts produced ectopic brown to silver scale transformations

(*Ubx* and *apA*), whereas other knockouts led to the loss of silver broadband reflection and turned

the scales brown (*Antp*, *dsx*, *apA* and *optix*) (Fig 1A). These modified scales provide a great

opportunity to investigate the structural basis of the different metallic colors in *B. anynana* scales

and understand the roles of these five genes in creating broadband reflectors in lepidopterans.

Here, we asked whether these genes modified different structural and pigmentary aspects and

components of the scale to create a silver color, and vice versa, when silver scales became brown.

We also investigated whether transformations of different structures and levels of pigmentation

within a scale, with the manipulation of each gene in isolation, is complete or gradual. We

hypothesized that more up-stream genes regulating the silver scale GRN will produce more

complete and extreme transformations of the scales, making them resemble a wildtype silver scale

(or a brown scale) more closely. We also posited that each gene may be involved in regulating a

subset of the characteristics of silver or brown scales, or, alternatively, may be regulating all of its

distinctive traits but only partially.

To answer these questions, we used light, conventional and focused ion beam (FIB) scanning

electron microscopy (SEM), UV-VIS-NIR microspectrophotometry, and systematic optical

modeling, to understand different aspects of the wildtype (WT) and transformed scales. We show

that the variable air gap thickness in silver scales is an important determinant of the broadband

reflectance. Transformation of silver to brown scales is accompanied by the loss of the contiguous

upper lamina, increases in lower lamina thickness and gains in pigmentation. The opposite occurs

when brown scales become silver scales. In addition, we identify *Antp* as a top regulator of the

silver scale GRN along with a novel role of this gene in determining scale shape.

<u>Results</u>

Ultrastructural and optical origin of broadband silver reflections in *Bicyclus anynana*

butterflies

We first investigated how the silver color is produced in five types of silver scales present on the

wings of *B. anynana* using light reflection and absorbance measures, as well as SEM (Fig. 1B-F).

On the ventral forewing, we focused on the silver and the gland scales on the androconial patch of

males (Fig 1A, WT-FW arrows). On the dorsal hindwing, we sampled the silver and grey-silver

scales near the androconial patches of males. Finally, we sampled coupling scales near the base of

the dorsal hindwing in females (Fig 1A, WT-HW arrows). All silver scales generally feature

rounded edges. The coupling scales, however, have a characteristic trowel-head shape (Fig 1F, F′),

and their ridges are oriented at an angle to the proximo-distal axis of the scale as compared to the

parallel arrangement of ridges in most other scale types. The abwing and adwing surfaces of the

scales exhibited an oil-slick like multi-hued appearance, typical of a thin film with varying

thickness (Fig 1B-F). The reflectance spectra of all silver scales exhibited broadband reflectance,

with higher intensities from the adwing surface (Fig 1B‴-F‴). Pigmentation levels were low in all

scales, as measured from low light absorbance (Fig 1B⁗- F⁗), with the exception of the grey-

silver scales which absorbed more light, indicating the presence of pigments. All silver scales

exhibited a closed upper lamina (Fig 1B″-F″). The different silver scale types exhibited varying

levels of perforations in the total area of upper lamina (Fig 1H, Supplementary Table S5 – source

data): from ~0% (fore- and hindwing silver scales) and 2.35% (hindwing coupling scales), to 5.4%

open (hindwing grey-silver scales) (Supplementary Tables S1,2). This correlation between the

integrity of the upper lamina coverage to the production of broadband reflectance affirms the

important role attributed to the upper lamina and/or the enclosed air layer.

Figure 1: *B. anynana* crispants sampled in this study and broadband reflectors in wildtype *B. anynana*. (A) WT dorsal and ventral wing surfaces with magnified views of the silver scales alongside ectopic silver scales or silver to brown transformed scales in crispants of the five genes. The colored box outlines indicate the position on the different wing surfaces of the WT and crispant scales. White arrows in the WT panels indicate the five different silver scale types characterized. Optical microscopy images of the abwing (upperside) (B-F) and adwing (underside) (B′-F′) surfaces of single scales, SEM images of the abwing surface (B″-F″), measured and modeled reflectance (B‴-F‴) and absorbance spectra (B⁗-F⁗) of the WT forewing silver and gland scales and the hindwing silver, grey-silver and coupling scales respectively. (G) Violin plots of the upper and lower lamina thicknesses of the different forewing and hindwing silver scales. (H) Area of the open upper lamina of the different silver scales on the forewings and hindwings. Boxplots show the median, inner and outer quartiles and whiskers up to 1.5 times the inter-quartile range. Means sharing the same letter are not significantly different (Tukey-adjusted comparisons). Scale bars: B,B′-F,F′ are 20 µm and B″-F″ are 2 µm.

In order to theoretically model these broadband metallic colors, we measured the air gap, lower and upper lamina thicknesses from FIB-SEM cross-sections of the different silver scale types (Fig 1G, Supplementary Table S5 – source data). The mean upper and lower lamina thicknesses of the different metallic scale types ranged from approximately 55-120 nm, with the coupling scales having the thickest lower and upper laminas (Fig 1G). We modeled the theoretical adwing and abwing spectra, the latter by incorporating the measured absorptivity curves (Fig 1B⁗-F⁗), which produced broadband reflectance with a series of very shallow peaks, whose positions are largely consistent with those in the corresponding measured spectra (Fig 1B‴-F‴, Supp Fig S1). A lower lamina thin film of an appropriate thickness could, alone, produce a broadband silver color (Supp Fig S2A). However, our results indicated that the presence of the additional upper lamina enclosing an air gap with varying thickness increased the mean broadband reflectivity by around 11% (Supp Fig S2A). Furthermore, our hierarchical modeling (Supp Figs S2 and S3) suggests that the observed variation in the upper and lower lamina thicknesses cannot reproduce the relatively flat broadband reflectance even when the sclerotized ridges are included (Supp Figs S2A-C), whereas the variation in the air gap layer, alone, can sufficiently account for the broadband silvery

reflectance, including the patches of distinct colors (e.g. pink and green) seen in between ridges in

high magnification light micrographs (Fig 1B,B′-F,F′, Supp Fig S2C) (24). Finally, our systematic

parametric modeling (Supp Fig S3) revealed that the thicknesses of both upper and lower laminas

for each of the silver scale types, except coupling, are more or less optimized to produce the

maximum broadband reflectivity. By contrast, the air gap layer is sufficiently thick (> 750 nm) to

ensure the broadband nature of the reflectivity, for a thickness of less than 500 nm will produce

strong peak(s) with chromatic effects destroying the silvery appearance (Supp Fig S3).

Interestingly, the modeling results predicted lower reflectivity at short wavelengths below 450 nm

for the coupling scales (Supp Fig S2n) similar to their measured spectra (Fig 1F‴). Given that

coupling scales are among the least pigmented (Fig. 1F⁗), our results indicate that this lower

reflectivity is a previously undocumented structural rather than a pigmentary effect. Further

corroborating these modeling results, a longitudinal cross-section of the coupling scale showed a

decrease in the thickness of the lower lamina from base to tip (Supp Fig S4) and a concomitant

change in the color of the scale from bronze/brown in the base/center to silver at the tips. This

increased thickness near the base of the scale suppresses shorter wavelengths of light and leads to

a greyish darker color (Fig 1F′), while optimum thickness towards the tip reinforces broadband

silver reflectance.

Antp is a selector gene for rounded scale morphology and acts upstream of _dsx_ in the silver

scale GRN

Both _Antp_ and _dsx_ male crispants exhibited a silver to brown color transformation on the forewing

but had different effects on scale shape (Fig 2A-C). The rounded edges of the distal end of the

silver scales of _Antp_ crispants changed into serrated finger-like projections, similar to other brown

scales on the same wing surface (Fig 2D, Supp Fig S5A). This was true in females also, where the

normal rounded brown scales on the posterior region of the ventral forewings became brown scales

with dentate margins in the crispants (Supp Fig S5B). In contrast, the loss of _dsx_ expression in the

same region produced brown scales with a rounded distal edge (Fig 2E). Furthermore, some of the

Antp crispant brown scales exhibited unusual ridge and crossrib orientations near the distal tips of

the scales, often lying perpendicular to the normal parallel arrangement of ridges (Fig 2F). In _Antp_

male crispants, we also noticed small clonal patches of brown scales within the hindwing silver

scale area (Fig 2G). Such clonal patches were also visible on female hindwings in homologous

regions of the wing, though not as contrasting (Supp Fig 5D). The brown scales in these clonal
patches had a dentate morphology similar to other dorsal hindwing scales, in contrast to the
rounded scales seen in the hindwing silver scale region in males or the homologous regions in
females (Supp Fig S5C, D).

Ultrastructure and pigmentation modifications accompanied the loss of silver reflectance in male
scales of both crispant types. The brown crispant scales had lower reflectivity in comparison with
the wildtype control scales (Fig 2A‴-C‴) and this was accompanied by an increase in absorbance
due to greater pigmentation (Fig 2A⁗-C⁗). The crispant brown scales also lost their continuous
upper lamina which now exhibited perforated windows (Fig 2A″-C″, I, Supplementary Tables
S1,2). Other ultrastructural modifications included an increase in lower lamina thickness (Fig 2H)
(Posthoc tests from LME, adjusted P-values <0.001; Supplementary Tables S3,4). *Antp* brown
scales had a greater lower lamina thickness (125.82 nm ± 19.7 nm) compared to the *dsx* brown
scales (91.9 nm ± 17.6nm) and both were significantly different from wildtype silver lower lamina
thickness (68.8 ± 14.75). The increase in pigmentation, loss of the upper lamina and increased
lower lamina thickness led to the loss of broadband metallic silver reflectance and created a brown
color due to the combination of pigments and the now dominant lower lamina reflectance (Fig
2A‴-C‴).

These results suggest that *Antp* is an upstream selector gene of silver scale color and morphology
in males and scale shape in females. *Antp* likely acts upstream of *dsx*, which appears to have similar
effects on color and upper lamina morphology but does not impact scale shape. To investigate the
extent to which Antp expression maps to silver scales, we examined the expression of Antp in 24-
hour male and female pupal wings. Antp protein was visible in punctate nuclei in the forewing
(Supp Fig S6A) and hindwing silver scale regions in males, especially around the future hindwing
gland (Supp Fig S6C), and also in female wings at homologous locations (Supp Fig S6B, D). Antp
protein expression was lower in the grey-silver scales on male hindwings (Supp Fig S6C). This
pattern on male hindwings was especially clear in a 48-hour pupal hindwing (Supp Fig S7). These
results indicate that Antp protein maps precisely to the area where silver scales develop on both
wings.

Figure 2: Characterization of the ultrastructure and pigmentation of the silver to brown scales in *Antp* and *dsx* B. anynana crispants. Optical microscopy images of the abwing (A-C) and adwing (A′-C′) surfaces, SEM images of the abwing surface (A″-C″), reflectance (A‴-C‴) and absorbance spectra (A⁗-C⁗) of the WT forewing control silver scales and the mutant brown scales of *Antp* and *dsx* crispants respectively. (D) Mosaic scale phenotype in *Antp* crispant forewing illustrating the mutant brown scales with dentate sculpting in the distal tips. (E) Mosaic scale phenotype in *dsx* crispant forewing illustrating the mutant brown scales with rounded distal tips. (F) SEM image of an *Antp* crispant forewing brown scale showing a nearly orthogonal ridge and crossrib orientation in the midst of the normal ridge and crossrib orientation parallel to the scale. These variations always occur at the distal tips of the crispant scales (N=4). (G) Mosaic scale phenotype in male *Antp* crispant hindwing illustrating the mutant brown scales with dentate sculpting in the distal tips. (H) Violin plots of the lower lamina thicknesses of the control silver scales and the crispant brown scales of *Antp* and *dsx* crispants (I) Total area of open upper lamina of the different control and mutant scales. Boxplots show the median, inner and outer quartiles and whiskers up to 1.5 times the inter-quartile range. Means sharing a letter are not significantly

different (Tukey-adjusted comparisons). Scale bars: A,A′-C,C′ and F are 20 μm, A″-C″ are 2 μm,

D, E,G are 200 μm.

Ectopic silver scales gain an upper lamina, decrease the thickness of their lower lamina and

lose pigmentation

We next investigated the ectopic forewing silver scales produced in the *apA* and *Ubx* crispants by

the transformation of non-reflective brown scales (Fig 1A). We characterized both ectopic silver

and gland scale types in *apA* crispants but only the ectopic silver scale type in the single *Ubx*

crispant due to size limitation of the mutant clone in that individual. Ectopic silver scales in both

apA and *Ubx* crispants looked similar to wildtype silver scales, including changes in cell shape

(Fig 3). The *Ubx* crispant scales exhibited an extensive transformation, from scalloped brown

scales to rounded silvery scales (Fig 3D, E), while the *apA* crispant scales changed primarily in

color and less extensively in shape (Fig 3A-C). The abwing and adwing surfaces of the ectopic

scales were metallic silvery thin films (Fig 3B, C, E). These scales were dramatically different

from the control brown scales from the same crispant regions which were brown in color from the

abwing side (Fig 3A, D) and had a bronze-golden adwing thin film (Fig 3A′, D′). Reflectance

spectra corresponded with the optical images, with the silver scales reflecting broadly (Fig 3A‴-

E‴). Ectopic metallic scales of both *apA* and *Ubx* crispants also lost pigmentation as compared to

the control brown scales, though there was variation between the different scale types (Fig 3A″″-

E″″).

Figure 3: Characterization of ectopic silver scale ultrastructure and pigmentation in *apA* and *Ubx* *B. anynana* crispants. Optical microscopy images of the abwing (A-E) and adwing (A′-E′) surfaces, SEM images of the abwing surface (A″-E″), reflectance (A‴-E‴) and absorbance spectra (A⁗-E⁗) of the *apA* forewing control brown scales, ectopic silver and gland scales and the *Ubx* hindwing control brown scales and ectopic silver scales respectively. The control scales for these

crispants are from the same wings as the ectopic scales, close to the mosaic crispant patches. (F)

Violin plots of the upper and lower lamina thickness of the control brown scales and the ectopic

silver and gland scales of *apA* and *Ubx* crispants. (G) Area of open upper lamina of the different

control and ectopic scales. Boxplots show the median, inner and outer quartiles and whiskers up

to 1.5 times the inter-quartile range. Means sharing a letter are not significantly different (Tukey-

adjusted comparisons). Scale bars: A,A′-E,E′ are 20 μm and A″-E″ are 2 μm.

Structural modifications accompanying brown to silver scale transformations, in *apA* and *Ubx*

crispants, were opposite to those seen when silver scales became brown. Ectopic silver scales

gained a closed upper lamina and an enclosed air layer, reduced their lower lamina thickness, and

lost pigmentation in comparison to control brown scales (Fig 3A″-E″, F, G). The upper lamina

transformation was complete for the *apA* and *Ubx* silver scale types, whereas many of the *apA*

ectopic gland scales exhibited a partial transformation, having a significantly greater percentage

of upper lamina open (Posthoc tests from LME, adjusted p-values <0.01; Supplementary Tables

S1,2) (Fig 3G). All ectopic silver scale types exhibited a significant decrease in lower lamina

thickness when compared to the control brown scales (Posthoc tests from LME: adjusted P-values

<0.0001; Supplementary Tables S3,4). Similar to the wildtype forewing silver scales, the ectopic

forewing silver scales in both *apA* and *Ubx* crispants also had thicker lower laminas as compared

to the respective upper laminas (Supplementary Tables S3,4). The greater variability in the

ultrastructural transformations and pigmentation of the *apA* ectopic silver scale types explains the

greater variability in the abwing reflectance spectra measured from these crispant scales (Fig 3B‴,

C‴).

apA, dsx and optix promote silver coloration via the gain of an upper lamina, decrease in

lower lamina thickness and loss of pigmentation

On the hindwings, crispants of *apA*, *dsx*, and *optix* showed loss of metallic silver scales and

transformation of these scales into brown scales. For *apA* these are the exact opposite phenotypes

from those observed on the forewing, where *apA* transformed brown scales into ectopic silver

scales. In *apA* and *dsx* crispants, hindwing silver and grey-silver scales were transformed into

brown scales (Fig 4A-F) while in the *optix* crispants, the coupling silver scales became brown

scales (Fig 4G-H). Brown crispant scales of *apA* individuals exhibited a dentate morphology while

those in *dsx* individuals exhibited some variation in morphology, from rounded scales in the silver

region to sometimes dentate scales in the grey-silver region. Brown *optix* coupling scales displayed

a noticeable change in morphology from the characteristic trowel-head shape of the control scales

to an oblong morphology in the crispant brown scales (Fig 4G-H). Reflectance spectra reflected

the change in coloration (Fig 4A‴ – H‴). In all instances of transformation, the adwing sides of

the crispant scales were thin films reflecting a deep bronze-golden color instead of the silver to

gold reflectance of the wildtype control scales (Fig 4A′-H′). In addition, in line with the silver to

brown transformation, there was an increase in pigmentation in most scales (Fig 4A″″-C″″, G″″-

H″″), except in the grey-silver region, where crispant brown scales, in both *apA* and *dsx* crispants,

had decreased pigmentation as compared to the controls (Fig 4D″″-F″″), despite control scales

displaying a lot of variation in pigmentation levels (Fig 4 D″″).

Similar to the forewing silver to brown transformations, crispant brown scales on the hindwing,

produced by all three genes, also exhibited a loss of the upper lamina (Fig 4A″-H″, J,

Supplementary Tables S1,2). This was not a clean morphological change, with many scales

exhibiting a gradualism in transformation, often having remnants of the lamina attached to the

crossribs and ridges (Fig 4A″-H″, J). The lower lamina thickness of these scales also increased

compared to wildtype silver controls (Posthoc tests from LME, adjusted P-values <0.001;

Supplementary Tables S3,4). Both *apA* and *dsx* hindwing brown scales in the silver and grey-silver

regions had significantly thicker lower laminas than control scales, while there was no change of

lamina thickness in *optix* crispant brown scales (Fig 4I, Supplementary Tables S3,4).

Figure 4: Characterization of the ultrastructure and pigmentation of transformed silver to brown scales in *apA*, *dsx* and *optix B. anynana* crispants. Optical microscopy images of the

abwing (A-H) and adwing (A'-H') surfaces, SEM images of the abwing surface (A″-H″), reflectance (A‴-H‴) and absorbance spectra (A⁗-H⁗) of the control WT silver scales and transformed crispant brown scales in the hindwing silver region (A-C), the hindwing grey-silver region (D-F) and the hindwing coupling scales (G-H). The control scales are WT silver scales from homologous regions on the wing. (I) Violin plots of the lower lamina thicknesses of control WT silver scales and the crispant brown scales of *apA, dsx* and *optix* crispants (G) Total area of open upper lamina of the different control and crispant brown scales. Boxplots show the median, inner and outer quartiles and whiskers up to 1.5 times the inter-quartile range. Means sharing a letter are not significantly different (Tukey-adjusted comparisons). Scale bars: A,A′-H,H′ are 20 µm and A″-H″ are 2 µm.

Discussion and conclusion

Metallic scales in *Bicyclus anynana* are undulatory broadband reflectors

We characterized five different metallic reflectors in *B. anynana* butterflies. Four out of the five scale types are present only in males, associated with the androconial regions that produce and secrete sex pheromones, while the fifth, the coupling scales, are found in both sexes near the base of the wings. All five metallic scale types investigated in this study produced broadband silver reflectance from an undulatory thin film achieved by the closure of the "open" windows present in typical wing scales. The sandwiched air gap layer of varying thickness between two chitinous laminas reflects spatially different wavelengths of light, resulting in a specular metallic color at the far-field due to additive color mixing. Such an ultrastructural modification involving the gain of a contiguous upper lamina leading to metallic coloration has been well-documented in multiple butterfly species across different families (21, 24), suggesting that the convergent evolution of metallic reflectors in butterflies can occur repeatedly with a few simple modifications to the basic scale architecture. Our modeling demonstrates that while butterflies can achieve broadband metallic colors by tuning just the lower lamina, the three-layer system comprising a sandwiched air layer between chitinous laminas is substantially brighter, perhaps explaining its repeated evolutionary origin (24). Moreover, the varying thickness of the air gap was the most important structural parameter to produce a nearly flat (smoothed out) broadband reflectance, while the upper and lower lamina thicknesses were more optimized for enhancing the broadband reflectivity. Pigmentation levels were low overall but varied among the different silver scales along with slight

variations in the total open area of the upper lamina. This intriguingly suggests that the amount of
pigmentation may be directly impacting the formation of the upper lamina, as previously
demonstrated in other scale types of *B. anynana* (33). We were unable to characterize the exact
pigment composition in these scales, but the absorbance spectra of the grey-silver scales suggests
that the pigments are not purely melanins but could instead be a combination of melanins and
additional pigments such as ommochromes (29).

Each of five genes that led to gains and losses of metallic reflectance also changed
pigmentation levels and scale ultrastructure
We discovered that disruptions to the five genes investigated (*Antp, dsx, apA, Ubx and optix*) lead
to simultaneous change in three scale features, suggestive of a switch-like or pleiotropic role for
these genes in determining the development of metallic scales. Transformation of metallic silver
scales into brown, non-reflecting scales in *Antp*, *dsx*, *apA* and *optix* crispants involved loss of an
upper lamina, gain in pigmentation levels, and an increase in lower lamina thickness. This was
generally true except in the case of the grey-silver scales of the hindwing where transformation to
brown scales led to a decrease in pigmentation. Likewise, gain of silver reflectance in scales of
apA and *Ubx* crispants was accompanied by the appearance of a continuous upper lamina that
closed the windows of the typical scale ground plan, a decrease in pigmentation levels, and a
decrease in the thickness of the lower lamina. The breadth of variation seen in individual crispant
brown scales and ectopic silver scales was potentially due to incomplete knockouts, and variation
in the protein function of each mutant allele. This was especially the case of *apA* crispants, possibly
due to the incomplete knockout of these genes in individual scales cells.

The gain and loss of an upper lamina with concurrent changes in the thickness of the lower lamina
suggests a possible conservation of total chitin production within a cell, but variable deposition
between the two laminas. There is some evidence for this in the forewing silver and gland scales
but not in the hindwing silver scale types. The summed thicknesses of the two laminas in the
wildtype forewing silver scales and ectopic silver scales was similar to the lower lamina thickness
of the forewing control brown scales and crispant brown scales. Though there is some indication
of the conservation of chitin production within a single cell, testing this proposition in the future
will require finer measurements of chitin production within different scale color types. Further,

Matsuoka and Monteiro (33) speculated that downstream effector genes such as *yellow* could affect
cuticle polymerization around crossribs to create windows because *yellow B. anynana* crispant
black and brown scales exhibited an upper lamina covering the windows. Given the low amounts
of pigmentation seen in the different silver scale types, it is possible that the gene *yellow* is not
highly expressed in silver scales, mirroring *yellow* crispant scales. *yellow* would then be a potential
downstream target, that would be repressed by the silver scale GRN. This hypothesis could be
tested by measuring levels of *yellow* expression in the different colored vs silver scales of *B.*
anynana, as well as in crispant individuals during development.

Antp is a selector gene for rounded scales and acts through DsxM to determine silver scales
Investigation of the mutant brown scales in *Antp* crispants uncovered an important role for this
gene in determining a rounded scale morphology. Scales on the posterior ventral forewing and
anterior dorsal hindwing of both sexes of *B. anynana*, a region where Antp is strongly expressed
(Supp Fig S6), normally have rounded distal ends, appearing silver in males and shades of brown
in females. In *Antp* crispants, the rounded scales developed serrated distal tips in both sexes,
concurrent with the transformation of silver to brown color in males only. These crispant brown
scales with serrated edges were similar to the scales usually found on the rest of the ventral
forewing (Supp Fig S5A, B) or dorsal hindwing (Supp Fig S5C, D), indicating that *Antp* is essential
for converting a serrated distal scale edge into a rounded scale. Furthermore, this result suggests
that *Antp* acts upstream of *dsx* in the forewing and hindwing silver scale GRN because male *dsx*
crispants exhibited a more restricted silver to brown transformation, still displaying the rounded
distal edge of a typical silver scale, instead of the dentate morphology. We propose that the protein
Antp activates *dsx* in the posterior ventral forewing or anterior dorsal hindwing (Fig 5). In females,
Antp would lead to the production of the DsxF protein splice variant, resulting in brown scales
with a rounded morphology since DsxF has no function in silver scale development (34). In males,
Antp would activate DsxM protein splice variant, which leads to the development of rounded silver
scales in both regions (34). Interestingly, the metallic silver scales on male forewing and hindwings
fall into two populations with different aspect ratios: the shorter silver scales covering most of the
region and the longer, thin gland scales that cover only the underlying pheromone glands (38).
The spatially defined occurrence of the long, thin gland scales is likely specified by the spatially
restricted expression of some unknown gland-specific gene, downstream of DsxM in the silver

scale GRN.

Figure 5: Proposed gene regulatory network (GRN) controlling silver scale development

and scale shape in different wing regions of a male *B. anynana*. For each wing region, GRNs

are indicated within boxes. The scale schematics inside the boxes represent the scale shape when

the 'rounded scales' part of the network is active (dark lines). Repressed parts of the network are

in grey. The scale schematics outside the boxes, next to each wing surface, reflect the scale shape

generally found on that surface otherwise.

The unusual ridge and crossrib orientation, almost orthogonal to the normal orientation, seen in a

few *Antp* crispant brown scales, hinted at a role for this gene in ridge patterning during the early

stages of scale development. In developing scales, the positioning and orientation of ridges is

determined by the spacing of F-actin filaments that emanate from the base of the scale, parallel to

the scale's long axis (39, 40). The disrupted and abrupt change in orientation of the ridges and

their corresponding crossribs in the *Antp* crispant scales suggests a potential disruption of the

orientation of F-actin filaments. Based on the distal domain of occurrence of the unusual ridge

patterns, along with the rounded to serrated distal edge change seen in *Antp* crispants, we speculate

that *Antp* regulates the distal expression of an unknown factor, possibly a cytoskeletal component,

that determines scale morphology and ridge/crossrib orientation. Hox genes such as *Abd-B* have

been identified to activate cytoskeletal components like cadherins in the development of posterior spiracles of *Drosophila* (41). Furthermore, *Ubx* directly regulates cytoskeletal components such as actin and contributes to cytoskeletal reorganization in haltere cells (42). In that context, our results identify a novel post-embryonic role of a Hox gene in the determination of the shape of a single cell. How *Antp* directs scale shape and ridge orientation, and which genes are its downstream targets, remains an exciting avenue for future work.

The silver scale GRN is modulated in various ways across the fore- and hindwings

The appearance of forewing silver and gland scales in ectopic dorsal locations on the forewings of *apA* crispants and ventral locations on the hindwings of *Ubx* crispants, lead us to propose that both genes are repressing the forewing silver scale GRN (Fig 5). *Ubx* is a well-known homeotic gene whose expression is restricted to the hindwings of many insects and which functions in determining hindwing shape and color patterns in butterflies (36, 43–45). Knockout of *Ubx* in *B. anynana* converts hindwing patterns into forewing patterns (36). Since *Antp* is a top regulator of the silver scale GRN, these results suggest that *Ubx* represses *Antp* expression on the ventral hindwings. Recent work showed that upon loss of *Ubx*, *Antp* is strongly expressed on the hindwings (Matsuoka and Monteiro, in prep). We propose that the silver scales observed on the ventral hindwing of the *Ubx* male crispant resulted from the de-repression of *Antp* in that region, followed by the activation of DsxM, and the development of rounded silver scales typical of the ventral forewing (Fig 5). Along similar lines, *apA* is a dorsal surface-specific gene and in *apA* crispants, forewing ventral silver and gland scales appeared on the dorsal forewing surface (35). This is again suggestive of *apA* repressing *Antp* on the dorsal forewing surface. Upon de-repression, Antp activates DsxM and this leads to the development of the metallic silver scales on the dorsal surface of males (Fig 5). This regulatory circuit appears to be different from the development of somatic mesodermal muscle tissues in *Drosophila* where *Antp* directly and positively regulates *apterous* (46). Our proposed forewing silver scale GRN thus identifies *Antp* as a top regulator, directing metallic scale development via *dsx*. *apA* and *Ubx* are repressors of this network on the dorsal forewings and ventral hindwings, respectively (Fig 5). However, the molecular basis of the direct or indirect regulatory interactions between these different genes remains to be identified.

On the hindwings, the silver and grey-silver scale GRN retains the same topology, with *Antp*

determining a rounded scale shape and acting via *dsx* to determine silver scale development (Fig

5). In contrast to its repressive role on the dorsal forewing silver scale GRN, *apA* acts as an

activator of the GRN on the hindwings. The difference in levels of Antp protein expression

between the silver and grey-silver scale regions suggests that varying thresholds of this protein are

functionally relevant in the two scale types. Lastly, *optix* is necessary to create a silver coupling

scale but it also modulates scale shape from an oblong to a trowel-shaped morphology. Its

expression in coupling scales of other butterflies such as *Heliconius* implicates a conserved role

for *optix* in the development of trowel-shaped coupling scales (47, 48).

In conclusion, this study identified the structural and photonic origin of broadband silver coloration

by comparing silver scales in WT and crispants for five genes in the butterfly *B. anynana*. Based

on the epistasis between the investigated genes, we proposed a GRN for metallic scale

development in butterflies. We identified *Antp* as an important top-level gene in the regulatory

network for silver scale development and a novel role for this Hox gene in determining the shape

of single cells.

<u>Materials and methods</u>

<u>Crispant individuals and sampled scale types</u>

All crispant individuals used in this study were previously generated in our lab by CRISPR Cas9-

mediated gene editing in *B. anynana. B. anynana* crispants were for the following genes: *apterous*

A (*apA*) (35), *Ultrabithorax* (*Ubx*) (36), *doublesex* (*dsx*) (34), *Antennapedia* (*Antp*) (36) and *optix*

(37). Five different silver scale types were sampled, namely, the silver and the gland scales on the

androconial patch of male ventral forewings, the silver and grey-silver scales near the androconial

patches of male dorsal hindwings and the coupling scales near the base of the dorsal hindwing in

females (Fig 1A).

<u>Scanning electron microscopy</u>

Five to six scales from different regions of the crispant individuals (Fig 1A) were individually

mounted onto carbon tape and sputter coated with gold using a JFC-1100 Fine Coat Ion Sputter

(JEOL Ltd. Japan). Images were obtained using a JEOL JSM-6510LV scanning electron

microscope (JEOL Ltd. Japan). For the wildtype, *apA* and *dsx* crispants, three different individuals

were sampled for each case. Two individuals were sampled for *Antp* and *optix* crispants while only

one crispant was obtained for *Ubx.*

<u>Percentage area of open upper lamina measurements</u>

From the SEM images, the percentage of area of open upper lamina was calculated using ImageJ

1.52a (Java 1.8.0_112) (49). A 20 μm^2 (10 μm^2 for gland scales) region of interest was defined

approximately at the center of the scale and the region outside cleared. A thresholding was applied

based on the values of the bright (ridge and crossribs) and dark (windows) regions of the SEM.

The dark areas were selected and added to the ROI Manager in ImageJ using the Analyze Particles

function. The selected regions were combined using the OR function in the ROI Manager. The

combined area of the open upper lamina within the defined regions was measured and converted

into a percentage.

<u>Focused ion beam scanning electron microscopy (FIB-SEM)</u>

FIB-SEM was used to measure the lower lamina, upper lamina, and air gap thicknesses for all

scale types in our analysis and corrected for tilted perspective (measured thickness / sin 52°).

Briefly, samples were prepared by sputter-coating with platinum to increase conductivity. The

scales were milled using a gallium ion beam on a FEI Versa 3D with the following settings: beam

voltage −8kV, beam current −12pA at a 52° tilt. Image acquisition was performed in the same

equipment with the following settings: beam voltage −5kV, beam current −13pA. Milling was

done at the center of each scale. Thickness measurements were done in ImageJ. For each scale

type, ten measurements were taken per scale with 3-16 scales sampled from 1-2 individuals.

Measurements were made along most of the lower lamina which is uniform, excluding the region

around the ridge base, where the thickness is highly variable.

<u>Optical imaging and UV-VIS-NIR microspectrophotometry</u>

Light microscope images of individual scales were recorded using the 20X lens of a uSight-2000-

Ni microspectrophotometer (Technospex Pte. Ltd., Singapore) and a Touptek U3CMOS-05

camera. Scales were individually mounted on a glass slide or in a refractive index matching

medium (clove oil) and multiple images at different focal planes (z-stack) were obtained. Stacking

was done in Adobe Photoshop v 22.5.1 (Adobe, California, USA).

Normal-incidence UV-VIS-NIR reflectance spectra of scales were acquired using the same

microspectrophotometer setup but with a 100x objective. Spectra with usable range between 335-

950 nm were collected using a high NA 100x objective from a ~2 μm sized spot (100 ms integration

time, 10x averaging) and calibrated using an Avantes WS-2 reference tile made of white diffuse

polytetrafluoroethylene. Individual scales were mounted on a black carbon tape and illuminated

with a Mercury-Xenon lamp (ThorLabs Inc., New Jersey, USA). Measurements were taken from

both abwing and adwing surfaces. For each scale type, measurements from five to ten individual

scales from one individual were averaged. Absorbance spectrum was measured for individual

scales immersed in a refractive index matching liquid (clove oil) using a 20X objective. Six to

eight individual measures from one individual were averaged for each crispant type and wildtype.

Analysis and spectral plots were done in R Studio 1.4.1106 with R 4.0.4 (50) using the R-package

pavo (v 2.7) (51).

Acknowledgements

We thank Tirtha Das Banerjee for providing us the *optix* crispants, Yuji Matsuoka for providing

us the *Antp* and *Ubx* crispants and Emilie Dion for helpful discussions on the statistics. We thank

Sree Vaishnavi Sundararajan and Gianluca Grenci (MBI) for access and help with SEM, and the

Pennycook group (MSE) for use of FIB-SEM. This research was supported by the National

Research Foundation (NRF) Singapore under the Competitive Research Programme (NRF-CRP20-

2017-0001 Award) and the National University of Singapore.

Author Contributions

A.P and A.M conceived and designed the study. A.P, C.F and V.S collected the spectral

measurements. A.P and C.F collected the SEM data and C.F collected the FIB-SEM data. V.S

performed the theoretical modeling. A.P analyzed all the data and did all the immunostainings.

A.P wrote the manuscript with inputs from all the authors.

References

1.   M. F. Land, The physics and biology of animal reflectors. *Prog. Biophys. Mol. Biol.* **24,**

75–106 (1972).

2.   A. E. Seago, P. Brady, J.-P. Vigneron, T. D. Schultz, Gold bugs and beyond: A review of

iridescence and structural colour mechanisms in beetles (Coleoptera). *J. R. Soc. Interface*

6 Suppl 2, S165–S184 (2009).

3.   A. C. Neville, Metallic gold and silver colours in some insect cuticles. *J. Insect Physiol.*

23, 1267–1274 (1977).

4.   E. J. Denton, M. F. Land, Mechanism of reflexion in silvery layers of fish and

cephalopods. *Proc. R. Soc. London. Ser. B, Biol. Sci.* **178**, 43–61 (1971).

5.   A. L. Holt, A. M. Sweeney, S. Johnsen, D. E. Morse, A highly distributed Bragg stack

with unique geometry provides effective camouflage for Loliginid squid eyes. *J. R. Soc.*

Interface **8**, 1386–1399 (2011).

6.   D. Gur, *et al.*, The Dual Functional Reflecting Iris of the Zebrafish. *Adv. Sci.* **5**, 1800338

(2018).

7.   E. J. Denton, Review lecture: on the organization of reflecting surfaces in some marine

animals. *Philos. Trans. R. Soc. London. Ser. B, Biol. Sci.* **258**, 285–313 (1970).

8.   S. Johnsen, Hide and Seek in the Open Sea: Pelagic Camouflage and Visual

Countermeasures. *Ann. Rev. Mar. Sci.* **6**, 369–392 (2014).

9.   H.-J. Wagner, R. H. Douglas, T. M. Frank, N. W. Roberts, J. C. Partridge, A Novel

Vertebrate Eye Using Both Refractive and Reflective Optics. *Curr. Biol.* **19**, 108–114

(2009).

10. B. D. Wilts, P. Pirih, K. Arikawa, D. G. Stavenga, Shiny wing scales cause spec(tac)ular

camouflage of the angled sunbeam butterfly, Curetis acuta. *Biol. J. Linn. Soc.* **109**, 279–

289 (2013).

11. D. G. Stavenga, H. L. Leertouwer, D. C. Osorio, B. D. Wilts, High refractive index of

melanin in shiny occipital feathers of a bird of paradise. *Light Sci. Appl.* **4**, e243–e243

(2015).

12. S. N. Nan, *et al.*, Keeping cool: Enhanced optical reflection and radiative heat dissipation

in Saharan silver ants. *Science (80-.).* **349**, 298–301 (2015).

13. P. Vukusic, J. R. Sambles, Photonic structures in biology. *Nature* **424**, 852–5 (2003).

14. S. Kinoshita, S. Yoshioka, J. Miyazaki, Physics of structural colors. *Reports Prog. Phys.*

71 (2008).

15. M. Srinivasarao, Nano-Optics in the Biological World: Beetles, Butterflies, Birds, and
Moths. *Chem. Rev.* **99**, 1935–1962 (1999).

16.   a R. Parker, D. R. McKenzie, M. C. J. Large, Multilayer reflectors in animals using green
and gold beetles as contrasting examples. *J. Exp. Biol.* **201**, 1307–1313 (1998).

17. E. D. Finlayson, L. T. McDonald, P. Vukusic, Optically ambidextrous circularly polarized
reflection from the chiral cuticle of the scarab beetle Chrysina resplendens. *J. R. Soc.*
Interface **14** (2017).

18. R. A. Steinbrecht, Fine structure and development of the silver and golden cuticle in
butterfly pupae. *Tissue Cell* **17**, 745–762 (1985).

19. R. a. Steinbrecht, W. Mohren, H. K. Pulker, D. Schneider, Cuticular Interference
Reflectors in the Golden Pupae of Danaine Butterflies. *Proc. R. Soc. B Biol. Sci.* **226**,
367–390 (1985).

20. D. R. McKenzie, Y. Yin, W. D. McFall, Silvery fish skin as an example of a chaotic
reflector. *Proc. R. Soc. London. Ser. A Math. Phys. Sci.* **451**, 579–584 (1995).

21. P. Vukusic, R. Kelly, I. Hooper, A biological sub-micron thickness optical broadband
reflector characterized using both light and microwaves. *J. R. Soc. Interface* **6 Suppl 2**,
S193-201 (2009).

22. C. Kilchoer, U. Steiner, B. D. Wilts, Thin-film structural coloration from simple fused
scales in moths. *Interface Focus* **9** (2019).

23. L. D'Alba, B. Wang, B. Vanthournout, M. D. Shawkey, The golden age of arthropods:
Ancient mechanisms of colour production in body scales. *J. R. Soc. Interface* **16** (2019).

24. A. Ren, *et al.*, Convergent Evolution of Broadband Reflectors Underlies Metallic
Coloration in Butterflies . *Front. Ecol. Evol.* **8**, 206 (2020).

25. A. Dolinko, *et al.*, Analysis of the optical properties of the silvery spots on the wings of
the Gulf Fritillary, Dione vanillae. *Sci. Rep.* **11**, 19341 (2021).

26. T. J. Simonsen, Comparative morphology and evolutionary aspects of the reflective under
wing scale-pattern in Fritillary butterflies (Nymphalidae: Argynnini). *Zool. Anz.* **246**, 1–10
(2007).

27. Z. Qingqing, *et al.*, Fossil scales illuminate the early evolution of lepidopterans and
structural colors. *Sci. Adv.* **4**, e1700988 (2021).

28. B. Vanthournout, *et al.*, Springtail coloration at a finer scale: mechanisms behind vibrant

collembolan metallic colours. *J. R. Soc. Interface* **18**, 20210188 (2021).

29.   B. R. Wasik, *et al.*, Artificial selection for structural color on butterfly wings and
comparison with natural evolution. *Proc. Natl. Acad. Sci.* (2014)
https:/doi.org/10.1073/pnas.1402770111 (August 5, 2014).

30.   R. C. Thayer, F. I. Allen, N. H. Patel, Structural color in Junonia butterflies evolves by
tuning scale lamina thickness. *Elife* **9**, e52187 (2020).

31.   R. H. Siddique, S. Vignolini, C. Bartels, I. Wacker, H. Hölscher, Colour formation on the
wings of the butterfly Hypolimnas salmacis by scale stacking. *Sci. Rep.* **6**, 36204 (2016).

32.   L. Livraghi, *et al.*, Cortex cis-regulatory switches establish scale colour identity and
pattern diversity in heliconius. *Elife* **10**, 1–31 (2021).

33.   Y. Matsuoka, A. Monteiro, Melanin Pathway Genes Regulate Color and Morphology of
Butterfly Wing Scales. *Cell Rep.* **24**, 56–65 (2018).

34.   A. Prakash, A. Monteiro, Doublesex Mediates the Development of Sex-Specific
Pheromone Organs in Bicyclus Butterflies via Multiple Mechanisms. *Mol. Biol. Evol.* **37**,
1694–1707 (2020).

35.   A. Prakash, A. Monteiro, *apterous A* specifies dorsal wing patterns and sexual traits in
butterflies. *Proc. R. Soc. B Biol. Sci.* (2018) https:/doi.org/10.1098/rspb.2017.2685.

36.   Y. Matsuoka, A. Monteiro, Hox genes are essential for the development of eyespots in
Bicyclus anynana butterflies. *Genetics* **217** (2021).

37.   T. Das Banerjee, S. K. Shan, A. Monteiro, optix is involved in
eyespot development via a possible positional information mechanism. *bioRxiv*,
2021.05.22.445259 (2021).

38.   E. Dion, A. Monteiro, J. Y. Yew, Phenotypic plasticity in sex pheromone production in
Bicyclus anynana butterflies. *Sci. Rep.* **6**, 1–13 (2016).

39.   A. Dinwiddie, *et al.*, Dynamics of F-actin prefigure the structure of butterfly wing scales.
Dev. Biol. **392**, 404–418 (2014).

40.   C. R. Day, J. J. Hanly, A. Ren, A. Martin, Sub-micrometer insights into the cytoskeletal
dynamics and ultrastructural diversity of butterfly wing scales. *Dev. Dyn.* **248**, 657–670
(2019).

41.   B. Lovegrove, *et al.*, Coordinated Control of Cell Adhesion, Polarity, and Cytoskeleton
Underlies Hox-Induced Organogenesis in Drosophila. *Curr. Biol.* **16**, 2206–2216 (2006).

42. A. Pavlopoulos, M. Akam, Hox gene Ultrabithorax regulates distinct sets of target genes
at successive stages of Drosophila haltere morphogenesis. *Proc. Natl. Acad. Sci.* **108**,
2855–2860 (2011).

43. X. Tong, S. Hrycaj, O. Podlaha, A. Popadic, A. Monteiro, Over-expression of
Ultrabithorax alters embryonic body plan and wing patterns in the butter fl y Bicyclus
anynana. *Dev. Biol.* **394**, 357–366 (2014).

44. Y. Tomoyasu, S. R. Wheeler, R. E. Denell, Ultrabithorax is required for membranous
wing identity in the beetle Tribolium castaneum. *Nature* **433**, 643–647 (2005).

45. S. D. Weatherbee, *et al.*, Ultrabithorax function in butterfly wings and the evolution of
insect wing patterns. *Curr. Biol.* **9**, 109–115 (1999).

46. M. Capovilla, Z. Kambris, J. Botas, Direct regulation of the muscle-identity gene apterous
by a Hox protein in the somatic mesoderm. *Development* **128**, 1221–1230 (2001).

47. A. Martin, *et al.*, Multiple recent co-options of Optix associated with novel traits in
adaptive butterfly wing radiations. *Evodevo* **5**, 1–13 (2014).

48. L. Zhang, A. Mazo-Vargas, R. D. Reed, Single master regulatory gene coordinates the
evolution and development of butterfly color and iridescence. *Proc. Natl. Acad. Sci.* **114**,
10707 LP – 10712 (2017).

49. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* **9**, 671–675 (2012).

50. R Core Team, R: A Language and Environment for Statistical Computing (2021).

51. R. Maia, H. Gruson, J. A. Endler, T. E. White, pavo 2: New tools for the spectral and
spatial analysis of colour in r. *Methods Ecol. Evol.* **10**, 1097–1107 (2019).

681