Communication

Major complex trait for early de novo programming 'CoV-MAC-TED' detected in human nasal epithelial cells infected by two SARS-CoV-2 variants is promising for designing therapeutic strategies

José Hélio Costa[1,2]*, Shahid Aziz[1,2], Birgit Arnholdt-Schmitt[1,2]*

1. Functional Genomics and Bioinformatics Group, Department of Biochemistry and Molecular Biology, Federal University of Ceará, Fortaleza, Brazil

2. Non-Institutional Competence Focus (NICFocus) 'Functional Cell Reprogramming and Organism Plasticity' (FunCROP), coordinated from Foros de Vale de Figueira, Alentejo, Portugal

*Corresponding authors:

Birgit Arnholdt-Schmitt
Non-Institutional Competence Focus (NICFocus) 'Functional Cell Reprogramming and Organism Plasticity' (FunCROP), coordinated from Foros de Vale de Figueira, Alentejo, Portugal
Email: biarnaflora@gmail.com

José Hélio Costa
Functional Genomics and Bioinformatics
Group, Department of Biochemistry and
Molecular Biology, Federal University of Ceará,
Fortaleza, Brazil
Email: helio.costa@ufc.br

Abstract

Background: Early metabolic reorganization was only recently recognized as essentially integrated part of immunology. In this context, unbalanced ROS/RNS levels that connected to increased aerobic fermentation, which linked to alpha-tubulin-based cell restructuration and control of cell cycle progression, was identified as *ma*jor *c*omplex *t*rait for *e*arly *d*e novo programming ('CoV-MAC-TED') during SARS-CoV-2 infection. This trait was highlighted as critical target for developing early anti-viral/anti-SARS-CoV-2 strategies. To obtain this result, analyses had been performed on transcriptome data from diverse experimental cell systems. A call was released for wide data collection of the defined set of genes for transcriptome analyses, named 'ReprogVirus', which should be based on strictly standardized protocols and data entry into the 'ReprogVirus Platform'. This platform is currently under development. However, so far an *in vitro* cell system from primary target cells for virus attacks that could ideally serve for standardizing data collection of early SARS-CoV-2 infection responses was not defined. *Results*: Here, we demonstrate transcriptome level profiles of the most critical 'ReprogVirus' gene sets for identifying 'CoV-MAC-TED' in cultured human nasal epithelial cells. Our results (a) validate 'Cov-MAC-TED' as crucial trait for early SARS-CoV-2 reprogramming and (b) demonstrate its relevance in cultured human nasal epithelial cells. *Conclusion*: *In vitro*-cultured human nasal epithelial cells proved to be appropriate for standardized transcriptome data collection in the 'ReprogVirus Platform'. Thus, this cell system is highly promising to advance integrative data analyses by help of Artificial Intelligence methodologies for designing anti-SARS-CoV-2 strategies.

Keywords: immunology paradigm shift, melatonin, ADH5, E2F1, cell cycle, SARS-CoV-2 Δ382, SARS-CoV-2 helicase, SARS-CoV-2 RdRp, anti-viral diagnosis and strategies, repurposing drugs

1. Background

Is there a paradigm shift in understanding immunology?

It is increasingly understood that plants and animals have similar responses and cell memory mechanisms to manage immunology (1, 2). Effective immunologic protection requires a diversity of innate and adaptive cell responses and cell memory tools (1-4). However, immunologic responses are energy consuming and require efficient metabolic reprogramming, but metabolic reorganization is not fully recognized as an integrated part of immunology (5-8). Viruses have comparatively low Gibbs energy due to their chemical compositions (9, 10). This makes their replication highly competitive and is the basis for their 'structural violence' against the host cell metabolism [see discussion on this term in Costa et al., 2021 (11)]. Virus-induced host cell reprogramming that is intended to support host defense is turned ad absurdum for the host and favors virus replication as the driving force. As a consequence of this conflict, any viral infection provokes struggling for commanding coordination of host cell program and this starts in the initially infected cells. Competing for bioenergy and for 'territories' is decisive for the success of virus reproduction and evolution.

These insights stimulated our group to take profit of mixed plant and human cell-based experimental systems for gaining general, relevant knowledge on early reprogramming during SARS-CoV-2 infection. Our complex approach was explained on the basis of extensive literature reviews in a recent paper entitled 'From plant survival under severe stress to anti-viral human defense – a perspective that calls for common efforts' (11, 12). Having applied our theoretical concept by comparing transcriptome data from a resilient plant system with coronavirus-infected human cells of diverse origins, a *ma*jor *c*omplex *t*rait for *e*arly *d*e novo programming upon SARS-CoV-2, named 'CoV-MAC-TED', was identified that should help tracing critical virus footprints. This trait covers unbalanced levels of ROS/RNS (i.e. reactive oxygen species in relation to reactive nitrogen species), which connects to temporarily increased aerobic fermentation that links to α-tubulin-based cell restructuration and cell cycle regulation (11).

Resilience can depend on the capacity for efficient early reprogramming – learning from plants

Plants as settled organisms are especially challenged to rapidly confront highly diverse and complex abiotic and biotic environmental constraints, including virus threats. Marker development is essentially required to permanently advance breeding strategies that can timely cope with ever changing environmental conditions, such as climate changes. Consequently, prediction of adaptive robustness that provides resilience is best explored in plants. In this context, early reprogramming in target cells and tissues that impacts final agronomic or quality characteristics (such as yield stability or richness in secondary metabolites) was raised by our group as a trait *per se* (13-19). We developed unique concepts and tool kits that allows predicting plant robustness in the field from respiration traits as early as several hours during seed germination (20-25). This innovative approach was tested and preliminary validated by using diverse plant species (24-26). This observation suggests common mechanisms for resilient life performance across plant species. As a hypothesis, we expect similar performance across oxygen-dependent, respiring eukaryotic organisms. Applying a higher degree of abstraction, we argued that these results might also be promising to identify critical mechanisms in human cells under virus stress including SARS-CoV-2, which might help to pave the way for designing early combating strategies.

Early reprogramming can link to ROS/RNS equilibration and sugar-dependent fermentation

We found that adaptive early reprogramming in plants can be critically connected to temporarily enhanced, sugar-dependent fermentation (11, 25). This process was essentially regulated by an enzyme named alternative oxidase (AOX), which maintains ROS and RNS levels equilibrated, regulates metabolic and energetic homeostasis and adjusts respiration overload and fermentation, which at the same time associates to induction and regulation of cell division growth (11, 25). These characteristics make AOX a highly promising functional marker resource for improving general plant resilience. Its role in early reprogramming was indicated in several applied systems cited in Arnholdt-Schmitt et al. (12). However, humans do not possess this enzyme (see discussions in 11, 12). However, understanding the functional importance of AOX in plants can guide research strategies to identify marker gene candidates for resilience in human target cells against virus attacks appropriate to characterize resilient traits early at reprogramming. Costa et al. (11) suggested that melatonin could at least in part substitute the early role of AOX during reprogramming. Melatonin is a natural hormone in humans, which has also been recognized as a phytohormone (27). It is produced in most organs and cells (27-31), including human salivary gland cells (32). Costa et al. (11) observed that *ASMT* transcript levels were increasing dependent on MOI level and infection time in MERS-CoV-infected MRC5 cells, which encouraged studying markers for melatonin metabolism in nose cells. Melatonin is known to possess anti-oxidant properties and shows high fluctuation in its cellular concentration (27, 33-35). Furthermore, melatonin is since long proposed as anti-viral agent (36) and promising as repurposing drug to treat SARS-CoV-2 infections (37, 38).

Driving a standardized collection of data on virus-induced early reprogramming

Arnholdt-Schmitt et al. (12) initiated common efforts for a standardized collection of transcriptome data during early reprogramming after virus infection from a defined set of genes, named 'ReprogVirus'. In the present communication, we reduced the number of genes from ReprogVirus to promising core markers for CoV-MAC-TED components (11). Thus, genes were selected to identify a shift in ROS/RNS (ASMTL, SOD1, SOD2, ADH5, NOS2), represent glycolysis (PFK, GAPDH, Eno) and lactic fermentation (LDH) as well as structural cell organization (α-Tub). We assumed that ASMTL could indicate oxidative stress equilibration, while SOD1 and SOD2 genes mark anti-oxidative activities and were selected to indicate oxidative stress. ASMTL is a paralog of ASMT, which is involved in melatonin synthesis in human cells (28). However, we could not find ASMT gene transcripts in collected human nasal epithelial cells. ADH5 is known to be involved in ROS/RNS equilibration through NO homeostasis regulation (39-41) and the inducible NOS gene, NOS2, relates to the induction of NO production (25, 42). NOS1 and NOS3 had not been encountered in sufficient quantities in the collected epithelial nose cell data. Further, we selected SNRK and mTOR to highlight cell energy-status signaling (43-46). The mTOR is activated when there is excess of energy in contrast to SNRK, where higher expression indicates energy depletion. Genes for E2F1 together with mTOR were included to indicate changes in cell cycle regulation, namely cell cycle progression (G1/S and G2/M transitions) (46). E2F1 belongs to the transcription factor family E2F and is known as cell cycle activator. The interferon regulator factor, IRF9, demonstrated early transcription in SARS-CoV-2 infected human lung adenocarcinoma cells and was therefore proposed as functional marker candidate that could signal initiation of the classical immune system (11).

2. Material and Methods

2.1 Gene expression analyses of RNA-seq data from SARS-CoV2 infected human nasal epithelial cells

In this work we analyzed the expression of the main ReprogVirus genes (11, 12) in transcriptomic data from human nasal epithelial cells infected with two SARS-CoV2 variants (wild type and mutant Δ382) at times 0, 8, 24 and 72 hpi (hours post infection). Transcriptomic data (RNA-seq data) are available in in SRA database at GenBank (NCBI) under the bioproject PRJNA680711 previously published by Gamage et al., (47). The ReprogVirus genes analyzed were involved in shift in ROS/RNS (ASMTL, SOD1, SOD2, ADH5, NOS2), glycolysis [Total PFK (PFKM, PFKL and PFKP), GAPDH, Total Eno (Eno1, Eno2 and Eno3)], lactic fermentation [Total LDH (LDH-A, LDH-B, LDH-C, LDH-AL6A, LDH-AL6B)] structural cell organization [Total α-Tub (TUB-A1B, TUB-A1C, TUB-A4A)], cell cycle activator (E2F1), cell energy-status (SNRK, mTOR) and immune response (IRF9). The accession numbers of these genes are available in Costa et al. (11). The SARS-CoV2 proliferation was monitored evaluating the transcript levels of virus helicase (YP_009725308.1) and virus RNA-dependent RNA-polymerase (RdRp) (YP_009725307.1) genes.

The gene expression was evaluated mapping, quantifying and normalizing the reads of each ReprogVirus gene in the RNA-seq data with three biological replicates. For this, the ReprogVirus cDNAs were aligned against the RNA-seq data using the Magic-Blast software (48). Specific parameters in Magic-Blast as word size of 64 was included to certificate specific read detection for each gene. The number of mapped reads was obtained using the HTSeq (49) program exploring an alignment file (in SAM format) derived from Magic-Blast. The normalization of reads was performed using the RPKM (Reads Per Kilobase of transcript per Million of mapped reads) method (50) according to the following equation: RPKM = (number of mapped reads X 10^9) / (number of sequences in each database X number of nucleotides of each gene).

3. Results

Figure 1 shows relative transcript level changes of selected ReprogVirus genes in epithelial nasal cells infected with SARS-CoV-2 or SARS-CoV-2 mutant Δ382 at 8 hpi, 24 hpi and 72 hpi. Transcript level changes are expressed as % of 0 hpi. In **supplementary table S1**, standard errors (SE) are listed for all genes.

Transcript profile level changes for SARS-CoV-2 infection are given in Fig.1A and demonstrate similar increases for ASMTL and ADH5 (207% and 197%) at 8 hpi and indicate unbalanced ROS/RNS at 24 hpi (94% ASMTL and 159% ADH5), which is supported by near-basal SOD1 and SOD2 transcript levels (112% and 108%) in relation to an increased transcript level of NOS2 (163%). At 72 hpi, transcript levels of all ROS/RNS related genes were simultaneously reduced to below basal values observed at 0 hpi (57% ASMTL, 76% SOD1, 68% SOD2, 73% ADH5 and 61% NOS2). Transcription of glycolysis-related enzymes was rapidly increased at 8 hpi with highest transcript accumulation for enolase (267%). This high increase at 8 hpi was associated to a similar increase in LDH transcription (251%) linked to highly enhanced transcript levels also for α-tubulin (238%). SARS-CoV-2 Δ382 (Fig. 1B) shows a very similar profile for ROS/RNS signaling and metabolism-and structural cell organization-related traits. They also display almost identical signaling for a change in cell cycle regulation and the lack of strong stimulation of the classical immune system represented by the missing of IRF9 transcript accumulation level changes. Both genotypes demonstrate high increase in E2F1 transcript stimulation at 8 hpi, which signals rapid cell cycle activation. E2F1 transcript levels decreased slightly at 24 hpi, but fall rapidly below the basal values observed at 0 hpi to 62% respectively 72%. These observations together with increased transcript levels of SNRK and mTOR at 8 hpi (126% and 162%) and 24 hpi (131% and 182%) for the original SARS-CoV-2, and for SARS-CoV-2 Δ382 at 8 hpi (111% and 150%) and at 24 hpi (149% and 205%) point for both virus variants to early energy depletion and rapidly driven cell cycle progression plus cell cycle arrest at 72 hpi. At 72 hpi, no energy depletion is signaled anymore for both virus variants (SNRK: 76%, 83%). However, SARS-CoV-2 Δ382 shows at 72 hpi a slower decrease of mTOR transcript accumulation (109% vs 86%), and this together with the slower decrease also for E2F1 transcription (72% vs. 62%) might indicate a slightly delayed cell cycle progression and arrest. However, the effect of such difference did not influence the initiation kinetics of IRF9 transcription. Transcript level increase for IRF9 is only marginally seen at 72 hpi though noticeable for both virus variants (>110% from 24 hpi). The infection trials had been performed in unsynchronized nasal cell cultures (47, 51). It cannot be excluded that these observed small differences in E2F1 and mTOR transcript levels between both virus variants may be due to experiment-dependent, differentially non-synchronized cell cycle phases.

Figure 1: Transcript accumulation of selected *ReprogVirus* marker genes in human nasal epithelial cells infected with two SARS-CoV-2 variants at 8 hours post infection (hpi), 24 hpi and 72 hpi. Transcript levels are averages from three cell origins (donators/cell cultures) given in % of 0 hpi. A – SARS-CoV-2 (virus originally discovered); B – SARS-CoV-2 Δ382

Figure 2 shows transcript accumulation in RPKM for both virus variants from nasal epithelial cells separated for the three cell origins that resulted in the averages shown in Figure 1. Additionally, this figure integrates transcript accumulation of virus helicase and virus RNA-dependent RNA-polymerase (RdRp) as markers for virus proliferation. In **Figure 2A**, it can be seen that cells from origin 1 demonstrate lower transcript levels for enolase and LDH at 24 hpi. This was observed after infection by both SARS-CoV-2 variants. Also for both virus variants, virus helicase and RdRp indicate the start of virus replication at around 24 hpi though to a slightly lower degree for the SARS-CoV-2 Δ382. In **Figure 2B**, transcript accumulation of virus helicase and virus RdRp point to a burst of virus replication at 72 hpi. The level of virus transcripts is lower for the mutant than for the original SARS-CoV-2 virus. However, cells that originated from donor 1 that showed lower transcript accumulation for enolase and LDH at 24 hpi in comparison to cells from donors 2 and 3 in response to both virus variants did not indicate a burst of virus replication at 72 hpi, but only a slow increase from 24 hpi. Nevertheless, cells of donor 1 also demonstrated lower virus RdRp transcript levels at 72 hpi (Fig. 3). This differential performance of cells from donor 1 connects to higher transcription levels of SNRK at 72 hpi in cells from donor 1 after infection with both, the original SARS-CoV-2 and SARS-CoV-2 Δ382 (Fig. 2 and 3). It signals higher depletion of energy for donor 1 cells at 72 hpi and confirms critical energy-dependency of SARS-CoV-2 replication for both variants. However, the higher values for enolase and LDH observed at 24 hpi for cells infected by the mutant, the differential development of SNRK over time and higher values at 72 hpi observed for mTOR from donor 1 in the mutant (Fig. 3) argues for prolonged cell cycle progression respectively a delay in cell cycle arrest related to both, the origin of cells from donor 1 and infection by the mutant. Nevertheless, since longer observation times and technical repetitions in synchronized cultures are missing, it cannot be concluded whether individual origin or differentially unsynchronized cell cultures caused these differences in the emergence of a burst in virus replication.

Figure 2: Burst of SARS-CoV-2 virus proliferation at 72 hpi in nasal epithelial cells from three origins infected by two SARS-CoV-2 variants indicates an energy-dependent link to aerobic glycolysis and fermentation at 24 hpi.

Figure 3: Transcript accumulation of E2F1, SNRK, and mTOR in human nasal epithelial cells infected by two SARS-CoV-2 variants at 8 hpi, 24 hpi and 72 hpi and transcript levels of RdRp (x0,5) at 72 hpi for cell origin 1

4. Discussion

This is the first time that relevance of the marker system 'CoV-MAC-TED', which is based on a relatively small set of key genes, was validated in primary human target nose cells for respiratory SARS-CoV-2 infections.

In **Figure 4**, we present a simplified scheme from the results here obtained in human nasal epithelial cells for the five main components of CoV-MAC-TED (1) ROS/RNS balance shift, represented by ASMTL, SOD1, SOD2, ADH5, and NOS2, (2) glycolysis and fermentation, represented by enolase and LDH, (3) structural reorganization and cell cycle progress and arrest, represented by E2F1, mTOR and α-tubulin, (4) energy status signaling, represented by SNRK and mTOR, and (5) initiation of the classical immune system, represented by IRF9. For our conclusions and hypothetical inferences, we consider the dynamic interplay between virus infection, ROS/RNS signaling, carbohydrate stress metabolism, aerobic fermentation and cyt respiration based on knowledge related to our recently published insights (11, 25) and state-of-the-art knowledge on SARS-CoV-2 infection and general stress biology. Thus, we expect that the indicated increase of oxidative stress signaled at 8 hpi by ASMTL and ADH5 upon SARS-CoV-2 infection associates with rapidly increased sucrose/glucose cell levels that stimulate the Cyt pathway via enhanced glycolysis, pyruvate production and increased TCA cycling in a way that the respiration chain gets overloaded by electrons. Consequently, ROS and RNS concentrations might increase. On the other hand, the Cyt pathway will temporarily be restricted due to rapidly consumed oxygen. In turn, aerobic lactic fermentation will be activated. Depending on stress level and the amount of sugar and duration of the situation of high-sugar level, anaerobic glycolysis can reach high turnover during cell reprogramming and a level of ATP production corresponding to the Warburg effect. Warburg effects are increasingly recognized as being part of normal

physiology (52, 53) that enables host cells to rapidly mobilize energy for host maintenance and stress escape. Results of Bharadwaj et al. (25) suggests that microbiota can help stress alleviation through forming a sink for sucrose that supports their own growth, and, depending on their nature, this could support symbiotic or parasitic performance. Viruses are parasitic structures. These structures are supposed to profit from the mobilized high energy favored by the thermodynamic conditions of their replication. However, in case of SARS-CoV-2 our results indicate that during the first hours post infection (8 hpi to 24 hpi) rapidly available energy is consumed first for cell cycle progression (E2F1, mTOR and α-tubulin signaling) and only from around 24 hpi energy is increasingly used for virus replication (Figure 2). This might signal during early hours after infection stress (observed at 8 hpi by highly increased ASMTL and ADH5 transcription) relieve to the host cell (low oxidative stress indicated by ASMTL, SOD1 and SOD2), lower sugar concentration (less glycolysis combined with lower degree of fermentation observed from 24 hpi) and, thus, enable normalization of TCA cycling and decrease of energy depletion as it is SNRK-signaled after 24 hpi at 72 hpi before the expected burst of the classical immune system. However, the change for ROS/RNS in favor of NO production at 24 hpi, when virus replication starts, points to a 'non-normal' unbalanced situation.

The lower values for enolase and LDH observed at 24 hpi for cells/cell cultures from donator 1 linked to delayed virus replication (Figure 2) might point to a potential strategy target for combating early initiation of SARS-CoV-2 replication by blocking the Warburg effect and its link to cell cycle progression.

Our analyses have been performed in public RNA sequencing data provided from experiments with human nasal epithelial cells described in Tan et al. (51) and Gamage et al. (47). Thus, results and conclusions of these authors helped us additionally to validate relevance our key marker candidate approach: Transcriptome accumulation of the selected set of 'ReprogVirus' genes showed similar performance of 'CoV-MAC-TED' in human nasal epithelial cells infected by the originally discovered SARS-CoV-2 virus compared to infection by SARS-CoV-2 Δ382. These results are in conformity with the conclusions drawn by Gamage et al. (47). In addition, our observation that both virus variants induced cell cycle activator E2F1 to highest transcript levels at 8 hpi among all tested genes and times is in good agreement with results of Gamage et al. (47). These authors highlighted enhanced numbers of transcription factor E2F targets during early virus infections and a decrease during time combined with high activity related to cell cycle checkpoint G2M. Further, we could validate IRF9 as a good marker gene candidate for the initiation of the classical immune system at 72 hpi, since Gamage et al. (47) had in the same data intensively studied the immune response of SARS-CoV-2 infected cells compared to cells infected by influenza H3N2, which showed a burst for virus replication already at 48 hpi.

Furthermore, ASMTL and ADH5 could be identified as functional marker genes that can trace changes in early ROS/RNS balances in virus-infected human nasal epithelial cells.

Conclusion

In vitro-cultured human nasal epithelial cells proved to be appropriate for standardized transcriptome data collection in the 'ReprogVirus Platform'. Thus, this cell system is highly promising to advance integrative data analyses by help of Artificial Intelligence methodologies for designing anti-SARS-CoV-2 strategies.

Figure 4: Validating CoV-MAC-TED as crucial trait for early SARS-CoV-2-induced reprogramming in human nasal epithelial cells – A simplified scheme for hypothetical metabolic principles and validation of markers.

Author Contribution

JHC & BA-S conceived the basic idea and plan the study. SA performed BLAST and helped JHC in transcriptome analyses, and manuscript preparation. BA-S reviewed the data analyses, interpretation and wrote the manuscript.

Funding

JHC is grateful to CNPq for the Researcher fellowship (CNPq grant309795/2017-6). SA is grateful to CAPES for the Doctoral fellowship.

Acknowledgements

The authors are grateful to Isabel Velada from the University of Évora, Portugal, for help in figure improvement.

Conflicts of Interest

The authors declare that there is no conflict of interest.

References

1. Haney CH, Ausubel FM, Urbach JM. Innate Immunity in Plants and Animals: Differences and Similarities. Biochem (Lond) (2014) 36(5):40–5. doi: 10.1042/BIO03605040
2. Gourbal B, Pinaud S, Beckers GJM, Van Der Meer JWM, Conrath U, Netea MG. Innate Immune Memory: An Evolutionary Perspective. Immunol Rev (2018) 283(1):21–40. doi: 10.1111/imr.12647
3. Nejat N, Mantri N. Plant Immune System: Crosstalk Between Responses to Biotic and Abiotic Stresses the Missing Link in Understanding Plant Defence. Curr Issues Mol Biol (2017) 23:1–16. doi: 10.21775/cimb.023.001
4. Kirman JR, Quinn KM, Seder RA. Immunological Memory. Immunol Cell Biol (2019) 97(7):615–6. doi: 10.1111/imcb.12280
5. Priyadarshini S, Aich P. Effects of Psychological Stress on Innate Immunity and Metabolism in Humans: A Systematic Analysis. PLoS One (2012) 7(9): e43232. doi: 10.1371/journal.pone.0043232
6. Delmastro-Greenwood MM, Piganelli JD. Changing the Energy of an Immune Response. Am J Clin Exp Immunol (2013) 2(1):30–54.
7. Ganeshan K, Nikkanen J, Man K, Leong YA, Sogawa Y, Maschek JA, et al. Energetic Trade-Offs and Hypometabolic States Promote Disease Tolerance. Cell (2019) 177(2):399–413.e12. doi: 10.1016/j.cell.2019.01.050
8. O'Sullivan D. The Metabolic Spectrum of Memory T Cells. Immunol Cell Biol (2019) 97(7):636–46. doi: 10.1111/imcb.12274
9. Popovic M, Minceva M. A Thermodynamic Insight Into Viral Infections: Do Viruses in a Lytic Cycle Hijack Cell Metabolism Due to Their Low Gibbs Energy? Heliyon (2020) 6(5):e03933. doi: 10.1016/j.heliyon.2020.e03933
10. Popovic M, Minceva M. Thermodynamic Insight Into Viral Infections 2: Empirical Formulas, Molecular Compositions and Thermodynamic Properties of SARS,

MERS and SARS-CoV-2 (Covid-19) Viruses. Heliyon (2020) 6(9):e04943. doi: 10.1016/j.heliyon.2020.e04943

11. Costa, J. H., Mohanapriya, G., Revuru, B., Noceda, C., Thiers, K. L. L., Aziz, S., ... & Arnholdt-Schmitt, B. (2021). ROS/RNS balancing, aerobic fermentation regulation and cell cycle control a complex early trait ('CoV-MAC-TED') for combating SARS-CoV-2-induced cell reprogramming. Frontiers Imm. https://doi.org/10.3389/fimmu.2021.673692.

12. Arnholdt-Schmitt, B., Mohanapriya, G., Bharadwaj, R., Noceda, C., Santos Macedo, E., & Sathishkumar, R. (2021). From plant survival under severe stress to anti-viral human defense–a perspective that calls for common efforts. Frontiers Imm. https://doi.org/10.3389/fimmu.2021.673723

13. Arnholdt-Schmitt B. Stress-Induced Cell Reprogramming. A Role for Global Genome Regulation? Plant Physiol (2004) 136(1):2579–86. doi: 10.1104/pp.104.042531

14. Arnholdt-Schmitt B (2005) Efficient cell reprogramming as a target for functional marker strategies? Towards new perspectives in applied plant nutrition research. J. Plant Nutr. Soil Sci. (Special Issue) 168 (4) 617-624

15. Arnholdt-Schmitt B (2005) Functional markers and a systemic strategy: Convergency between plant breeding, plant nutrition and molecular biology. Plant Physiology and Biochemistry 43 (9), 817-820

16. Arnholdt-Schmitt B, Costa JH, de Melo DF. AOX–a Functional Marker for Efficient Cell Reprogramming Under Stress? Trends Plant Sci (2006) 11 (6):281–7. doi: 10.1016/j.tplants.2006.05.001

17. SircarD, Cardoso HG, Mukherjee C, Mitra A, Arnholdt-Schmitt A (2012). Alternative oxidase (AOX) and phenolicmetabolism in methyl-jasmonate-treated hairy root cultures of Daucus carota L. Journal of Plant Physiology 169(7):657-63.doi:
10.1016/j.jplph.2011.11.019.http://www.ncbi.nlm.nih.gov/pubmed/22326792.

18. Cardoso HG, Arnholdt-Schmitt B. (2013). Functional Marker Development Across Species in Selected Traits. In Diagnostics in Plant Breeding, eds. Lübberstedt T & Varshney RK, 467-515. Springer Netherlands. doi: 10.1007/978-94-007-5687-8_21.

19. Orfanoudakis M, Sinapidou E, Arnholdt- Schmitt B (2015) Can AOX gene diversity mark herbal tea quality? A proposal. Gupta KJ, MuR LAJ, NeelwarnE B (eds.) Alternative respiratory pathways in higher plants. John Wiley & Sons, Inc. Oxford. pp: 311-314. ISBN: 978-1-118-79046-5

20. Arnholdt-Schmitt B, Hansen LD, Nogales A (2016). Calorespirometry, oxygen isotope analysis and functional-marker-assisted selection ('CalOxy-FMAS') for genotype screening: A novel concept and tool kit for predicting stable plant growth performance and functional marker identification. Briefings in Functional Genomics. Doi: 10.1093/bfgp/elv008.

21. Arnholdt-Schmitt B, Ragonezi C, Cardoso H. "A Central Role of Mitochondria for Stress-Induced Somatic Embryogenesis," In: MA Germanà and M Lambardi, editors. In Vitro Embryogenesis in Higher Plants. New York, NY: Humana Press (2016) p. 87–100. doi: 10.1007/978-1-4939-3061-6_4

22. Arnholdt-Schmitt B and Vinod Kumar Patil (2017) Calorespirometry – a novel tool in functional hologenomics to select 'green' holobionts for biomass production. Editor-in-chief: Walker, John M. Editor: Dr Kapuganti Jagadis Gupta Methods in Molecular Biology: ISSN: 1064-3745. Indexed in PubMed and ISI web of Science; Published online Springer protocols and hard cover by Humana Press

23. Arnholdt-Schmitt B and Vinod Kumar Patil (2017) Studying individual plant AOX gene functionality in early growth regulation – a new approach. Chapter in 'Methods in in Molecular Biology: Plant Respiration& Internal Oxygen: Methods and Protocols', Editor-in-chief: Walker, John M. Editor: Dr Kapuganti Jagadis Gupta Methods in Molecular Biology: ISSN: 1064-3745. Indexed in PubMed and ISI web of Science; Published online Springer protocols and hard cover by Humana Press

24. Mohanapriya G, Bharadwaj R, Noceda C, Costa JH, Kumar SR, Sathishkumar R, Thiers KLL, Santos Macedo E, Silva S, Annicchiarico P, Groot SPC, Kodde J, Kumari A, Gupta KJ and Arnholdt-Schmitt B (2019) Alternative Oxidase (AOX) Senses Stress Levels to Coordinate Auxin-Induced Reprogramming From Seed Germination to Somatic Embryogenesis—A Role Relevant for Seed Vigor Prediction and Plant Robustness. Front. Plant Sci. 10:1134. doi: 10.3389/fpls.2019.01134

25. Bharadwaj R, B., Noceda, C., Gunasekaran, M., Kumar, S. R., Thiers, K. L. L., Costa, J. H., ... & Arnholdt-Schmitt, B. (2021). Adaptive reprogramming during early seed germination requires temporarily enhanced fermentation–a critical role for alternative oxidase (AOX) regulation that concerns also microbiota effectiveness. Front. Plant Sci. | doi: 10.3389/fpls.2021.686274.

26. Arnholdt-Schmitt B, Gunasekaran Mohanapriya, Ramalingam Sathishkumar, Elisete Santos Macedo, José Hélio Costa (2018) Predicting biomass production from plant robustness and germination efficiency by calorespirometry. Chapter 5 in: Kumar, Ashwani, Ogita, Shinjiro, Yau, Yuan-Yeu (Eds.) Biofuels: Greenhouse Gas Mitigation and Global Warming. Next Generation Biofuels and Role of Biotechnology. Springer Nature DOI10.1007/978-81-322-3763-1, Hardcover ISBN978-81-322-3761-7

27. Zia SF, Berkowitz O, Bedon F, Whelan J, Franks AE, Plummer KM. Direct Comparison of Arabidopsis Gene Expression Reveals Different Responses to Melatonin Versus Auxin. BMC Plant Biol (2019) 19(1):567. doi: 10.1186/s12870-019-2158-3

28. Zhao D, Yu Y, Shen Y, Liu Q, Zhao Z, Sharma R, et al. Melatonin Synthesis and Function: Evolutionary History in Animals and Plants. Front Endocrinol (Lausanne) (2019) 10:249. doi: 10.3389/fendo.2019.00249

29. Hakkaart GA, Dassa EP, Jacobs HT, Rustin P. Allotopic Expression of a Mitochondrial Alternative Oxidase Confers Cyanide Resistance to Human Cell Respiration. EMBO Rep (2006) 7(3):341–5. doi: 10.1038/sj.embor.7400601

30. Silvestri M, Rossi GA. Melatonin: Its Possible Role in the Management of Viral Infections–a Brief Review. Ital J Pediatr (2013) 39:61. doi: 10.1186/ 1824-7288-39-61

31. Oláh J, Lehotzky A, Szunyogh S, Szénási T, Orosz F, Ovádi J. Microtubule Associated Proteins With Regulatory Functions by Day and Pathological Potency at Night. Cells (2020) 9(2):357. doi: 10.3390/cells9020357

32. Isola M, Lilliu MA, Loy F, Isola R. Diabetic Status Influences the Storage of Melatonin in Human Salivary Glands. Anat Rec (Hoboken) (2018) 301 (4):711–6. doi: 10.1002/ar.23750

33. Liu L, Labani N, Cecon E, Jockers R. Melatonin Target Proteins: Too Many or Not Enough? Front Endocrinol (Lausanne) (2019) 10:791. doi: 10.3389/ fendo.2019.00791

34. Zhang R, Wang X, Ni L, Di X, Ma B, Niu S, et al. COVID-19: Melatonin as a Potential Adjuvant Treatment. Life Sci (2020) 250:117583. doi: 10.1016/j.lfs.2020.117583

35. Zhou Y, Hou Y, Shen J, Huang Y, Martin W, Cheng F. Network-Based Drug Repurposing for Novel Coronavirus 2019-Ncov/Sars-Cov-2. Cell Discovery (2020) 6:14. doi: 10.1038/s41421-020-0153-3

36. Huang, S. H., Liao, C. L., Chen, S. J., Shi, L. G., Lin, L., Chen, Y. W., ... & Lin, G. J. (2019). Melatonin possesses an anti-influenza potential through its immune modulatory effect. Journal of Functional Foods, 58, 189-198.

37. Juybari, K. B., Pourhanifeh, M. H., Hosseinzadeh, A., Hemati, K., & Mehrzadi, S. (2020). Melatonin potentials against viral infections including COVID-19: Current evidence and new findings. Virus research, 198108.

38. Fernandes, P. A., Kinker, G. S., Navarro, B. V., Jardim, V. C., Ribeiro-Paz, E. D., Córdoba-Moreno, M. O., ... & Markus, R. P. (2021). Melatonin-Index as a biomarker for predicting the distribution of presymptomatic and asymptomatic SARS-CoV-2 carriers. Melatonin Research, 4(1), 189-205.

39. Shi YF, Wang DL, Wang C, Culler AH, Kreiser MA, Suresh J, et al. Loss of GSNOR1 Function Leads to Compromised Auxin Signaling and Polar Auxin Transport. Mol Plant (2015) 8(9):1350–65. doi: 10.1016/j.molp.2015.04.008

40. Rizza S, Filomeni G. Role, Targets and Regulation of (De)Nitrosylation in Malignancy. Front Oncol (2018) 8:334. doi: 10.3389/fonc.2018.00334

41. Barnett SD, Buxton ILO. The Role of S-Nitrosoglutathione Reductase (GSNOR) in Human Disease and Therapy. Crit Rev Biochem Mol Biol (2017) 52(3):340–54. doi: 10.1080/10409238.2017.1304353

42. Akaike T, Maeda H. Nitric Oxide and Virus Infection. Immunology (2000) 101:300–8. doi: 10.1046/j.1365-2567.2000.00142.

43. Bailey-Serres J, Pierik R, Ruban A, Wingler A. The Dynamic Plant: Capture, Transformation, and Management of Energy. Plant Physiol (2018) 176 (2):961–6. doi: 10.1104/pp.18.00041

44. Sakr S, Wang M, Dédaldéchamp F, Perez-Garcia MD, Ogé L, Hamama L, et al. The Sugar-Signaling Hub: Overview of Regulators and Interaction With the Hormonal and Metabolic Network. Int J Mol Sci (2018) 19(9):2506. doi: 10.3390/ijms19092506

45. Wurzinger B, Nukarinen E, Nägele T, Weckwerth W, Teige M. The SnRK1 Kinase as Central Mediator of Energy Signaling Between Different Organelles. Plant Physiol (2018) 176(2):1085–94. doi: 10.1104/pp.17.01404

46. McNulty S, Flint M, Nichol ST, Spiropoulou CF. Host Mtorc1 Signaling Regulates Andes Virus Replication. J Virol (2013) 87(2):912–22. doi: 10.1128/JVI.02415-12

47. Gamage AM, Tan KS, Chan WOY, Liu J, Tan CW, Ong YK, Thong M, Andiappan AK, Anderson DE, Wang Y, Wang LF. Infection of human Nasal Epithelial Cells with SARS-CoV-2 and a 382-nt deletion isolate lacking ORF8 reveals similar viral kinetics and host transcriptional profiles. PLoS Pathog. 2020 Dec 7;16(12):e1009130. doi: 10.1371/journal.ppat.1009130.

48. Boratyn GM, Thierry-Mieg J, Thierry-Mieg D, Busby B, Madden TL. Magic-BLAST, an accurate RNA-seq aligner for long and short reads. BMC Bioinformatics. 2019 Jul 25;20(1):405. doi: 10.1186/s12859-019-2996-x.

49. Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput sequencing data. Bioinformatics. 2015 Jan 15;31(2):166-9. doi: 10.1093/bioinformatics/btu638.

50. Mortazavi A, Williams BA, Mccue K, Schaeffer L, Wold B. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods* (2008) 5:621-628.

51. Tan KS, Andiappan AK, Lee B, Yan Y, Liu J, Tang SA, Lum J, He TT, Ong YK, Thong M, Lim HF, Choi HW, Rotzschke O, Chow VT, Wang Y. RNA Sequencing of H3N2 Influenza Virus-Infected Human Nasal Epithelial Cells from Multiple Subjects Reveals Molecular Pathways Associated with Tissue Injury and Complications. Cells. 2019 Aug 27;8(9):986. doi: 10.3390/cells8090986.

52. Melkonian EA, Schury MP. Biochemistry, Anaerobic Glycolysis. 2021 Aug 9. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan–. PMID: 31536301.

53. Kutschera U, Pieruschka R, Farmer S, Berry JA. The Warburg-effects: basic metabolic processes with reference to cancer development and global photosynthesis. Plant Signal Behav. 2020 Jul 2;15(7):1776477. doi: 10.1080/15592324.2020.1776477.