Specificities and commonalities of carbapenemase producing *Escherichia coli* isolated in France from
2012 to 2015.

[1,2]Rafael Patiño-Navarrete[#], [1,2]Isabelle Rosinski-Chupin[#], [1,2]Nicolas Cabanel, [1,2,3]Pengdbamba
Dieudonné Zongo, [1,4]Mélanie Héry, [1,4]Saoussen Oueslati, [1,4]Delphine Girlich, [1,4,5,6]Laurent
Dortet, [1,4,6]Rémy A Bonnin, [1,4,5,6]Thierry Naas[$] and [1,2,*]Philippe Glaser[$].

[$ and #] Contributed equally

[1]Unité EERA, Institut Pasteur, APHP, Université Paris Saclay, [2]UMR3525, CNRS, 28 rue du Dr Roux,
75015, Paris, France; [3]Sorbonne Université, Paris, France, [4] Team "Resist" INSERM U1184 "Immunology
of Viral, Auto-Immune, Hematological and Bacterial diseases (IMVA-HB),"; [5]Department of
Bacteriology-Hygiene, Bicêtre Hospital, APHP, [6]Associated French National Reference Center for
Antibiotic Resistance, Le Kremlin-Bicêtre, France.
15

17  *Corresponding author:
18  Philippe GLASER,
19  Institut Pasteur, 28 Rue du Dr Roux, 75724 Paris Cedex 15,
20  Tel + 33 1 45 68 89 96 - E-mail: pglaser@pasteur.fr

22  Running title: Carbapenemase producing *E. coli* from the French NRC
23

**ABSTRACT**

Carbapenemase-producing *Escherichia coli* (CP-*Ec*) represent a major public health threat with a risk of dissemination in the community as it has occurred for lineages producing extended spectrum ß-lactamases. To characterize the extend of CP-*Ec* spread in France, isolates from screening and infection samples received at the French National Reference Centre laboratory (F-NRC) for carbapenemase-producing Enterobacterales were investigated. Six hundred and ninety one CP-*Ec* isolates collected between 2012 and 2015 and 22 before were fully sequenced. Analysis of their genome sequences revealed some disseminating multidrug resistant (MDR) lineages frequently acquiring diverse carbapenemase genes mainly belonging to clonal complex (CC) 23 (ST 410) and CC10 (ST10, ST167) and sporadic isolates including rare ST131 isolates (n=17). However, the most represented ST was ST38 (n=92) with four disseminated lineages carrying $bla_{OXA-48-like}$ genes inserted in the chromosome. Globally, the most frequent carbapenemase gene (n=457) was $bla_{OXA-48}$. It was also less frequently associated with MDR isolates being the only resistance gene in 119 isolates. Thus, outside the ST38 clades, its acquisition was frequently sporadic with no sign of dissemination, reflecting the circulation of the IncL plasmid pOXA-48 in France and its high frequency of conjugation. In contrast $bla_{OXA-181}$ or $bla_{NDM}$ genes were often associated with the evolution of MDR *E. coli* lineages characterized by mutations in *ftsI* and *ompC.*

**IMPORTANCE**

Carbapenemase-producing *Escherichia coli* (CP-*Ec*) might be difficult to detect, as minimal inhibitory concentrations can be very low. However, their absolute number and their proportion among carbapenem-resistant *Enterobacterales* have been increasing, as reported by WHO and national surveillance programs. This suggests a still largely uncharacterized community spread of these isolates. Here we have characterized the diversity and evolution of CP-*Ec* isolated in France before 2016. We show that carbapenemase genes are associated with a wide variety of *E. coli* genomic backgrounds and a small number of dominant phylogenetic lineages. In a significant proportion of CP-*Ec*, the most frequent carbapenemase gene $bla_{OXA-48}$, was detected in isolates lacking any other resistance gene, reflecting the dissemination of pOXA-48 plasmids, likely in the absence of any antibiotic pressure. In contrast carbapenemase gene transfer may also occur in multi-drug resistant *E. coli*, ultimately giving rise to at-risk lineages encoding carbapenemases with a high potential of dissemination.

**INTRODUCTION**

*Escherichia coli* is one of the first causes of diverse bacterial infections in the community and in the hospital. In particular, it is the most frequent cause of urinary tract infection (UTI) and 50-60% of women will suffer at least once a UTI during her life (1). Therefore, multidrug resistance (MDR) in *E. coli* is a major public health issue making *E. coli* infections more difficult to treat. In addition, as carbapenemase-producing *Enterobacterales* (CPE) are increasingly detected (2, 3) and *E. coli* is an ubiquitous member of the normal gut microbiome in humans, carbapenemase-producing *E. coli* (CP-*Ec*) are also becoming a major actor for the global dissemination of carbapenemase genes (4).

The emergence and spread of carbapenem-resistant Gram-negative bacteria is mainly linked to the widespread dissemination through horizontal gene transfer (HGT) of mobile genetic elements (MGEs) encoding carbapenemases. These carbapenemases belong to Ambler class A (mainly KPC-type), class B (metallo-β-lactamases IMP-, VIM- and NDM-types) or class D (OXA-48-like enzymes) of beta-lactamases (5). The global epidemiology of extended spectrum ß-lactamases (ESBL) producing *E. coli* has been characterized through multiple studies, revealing in Western countries the major contribution of the sequence type (ST)131 lineage in the high prevalence of $bla_{CTX-M}$ family ESBL genes (6). Much less is known with respect to CP-*Ec*.

Since in 2012, the French National Reference Centre Laboratory for carbapenemase-producing *Enterobacterales* (F-NRC) has experienced a steady increase in the number of CP-*Ec* isolates received each year (2). A multi locus sequence typing (MLST) analysis of isolates received in 2012 and 2013 revealed a broad diversity of STs, as the 140 analyzed isolates belonged to 50 different STs. However, a few STs were over-represented (7), such as the ST38 (24 isolates) and the ST410 (10 isolates). In that study, only one isolate belonged to ST131, contrasting with the situation in the UK where ST131 isolates represented 10% of the CP-*Ec* isolates received between 2014 and 2016 by Public Health England (8). On the other hand, a genome based survey of CPE in the Netherlands between 2014 and 2018, revealed that the 264 received *E. coli* isolates belonged to 87 different STs, with three dominant lineages, ST38 (n=46), ST167 (n=22) and ST405 (n=16) representing 32 % of the isolates (9). F-NRC isolates also showed a predominance of isolates producing OXA-48-like carbapenemases followed by NDM-producing isolates and suggested clonally related isolates among ST38 OXA-48-producing isolates and ST410 OXA-181-producing isolates, respectively (7).

As whole-genome sequencing (WGS) significantly increases our ability to infer phylogenetic relationships between isolates, we recently sequenced 50 ST410 CP-*Ec* isolates received by the F-NRC between 2011 and 2015 (10) and found that 72% of them belonged to a newly described ST410 lineage producing OXA-181 (11). We showed that this clade is characterized by mutations in the two porin genes *ompC* and *ompF* leading to a decreased outer membrane permeability to certain ß-lactams and by a four-codon duplication (YRIN) in the *ftsI* gene encoding the penicillin binding protein 3 (PBP3)

leading to a decreased susceptibility to ß-lactams targeting this PBP (10). After a thorough analysis of

CP-*Ec* genome sequences from public databases for mutations in these three genes, we proposed that

CP-*Ec* followed three different evolutionary trajectories. In some lineages which are enriched in CP-*Ec*

isolates and have disseminated globally, acquisition of carbapenem resistance genes might have been

facilitated by the mutations in porin genes and in *ftsI*. In ST38, the genetic background and in particular

a specific *ompC* allele with reduced permeability to some ß-lactams may have similarly facilitated the

acquisition of carbapenemase genes. In contrast, other CP-*Ec* isolates including from ST131 might have

occurred sporadically following the acquisition of plasmids encoding carbapenemase and with no clue

of dissemination (10).

Here we thoroughly characterized the diversity of CP-*Ec* circulating in France by sequencing the

genome of almost all isolates received by the F-NRC from its creation until 2015 (Table S1). By

combining whole genome phylogeny with the addition of *E. coli* genome sequences publicly available

(Table S2), and tracking mutations in candidate genes, we show that three different situations are

encountered. The high transmission potency of the IncL pOXA-48 plasmids has led to a high frequency

of OXA-48-producing isolates, often characterized by susceptibility to most non-ß-lactam antibiotics.

On the other hand, an increasing number of CP-*Ec* lineages, mainly producing OXA-181 and NDM

carbapenemases, are observed in France as in other Eurpean countries. These lineages are multi-drug

(MDR) or extensively drug resistant (XRD) lineages. They are strongly mutated in quinolone resistance

determinants and are often mutated in *ftsI* and in porin genes. Finally, the rapid dissemination of four

OXA-48/OXA-244 ST38 lineages might have been favored by the chromosomal integration of the

carbapenemase gene.

**RESULTS**

**CP-*Ec* isolates collected until 2015 by the French National Reference Center**.

Isolates analyzed in this work were sent to the F-NRC laboratory on a voluntary basis from private and

public clinical laboratories from different parts of France mainly between the years 2012 to 2015.

During this period, we encountered a regular increase in the number of isolates received each year

(Fig 1A) corresponding to an increasing number of isolated strains and an increasing frequency of

isolates sent to the F-NRC. 713 CP-*Ec* isolates, including 22 collected from 2001 to 2011 were submitted

to WGS. The 691 sequenced isolates of the 2012-2015 period represented 87.5 % of the 790 CP-*Ec*

isolates received by the F-NRC during this period.

The majority of the sequenced isolates (66.5%, 474/713) were from rectal swab screening of patients

suspected of carrying CPE (patients repatriated from an hospital abroad, patients that have visited a

foreign country within the last six months, contact patients of a former carrier, or a previously known

carrier), 24% (172/713) were considered to be responsible for infection (isolated from clinical

samples), and the source was unknown for 67 isolates (Fig. 1A, Table S1). During the four-year period,

The number of infection-related isolates was found to increase more slowly than the number of

screening or of unknown source isolates. A nearly two-fold decrease in the proportion of clinical versus

total isolates was observed between 2013 and 2014 (Fig. 1B). Among the 172 isolates associated with

disease, 122 were isolated from urine (71%), 15 from blood samples, eight from deep site samples,

eight from wound samples, nine from the respiratory tract, three from vaginal samples, three from

pus, three from bile, and one from the skin of a newborn. All these isolates were previously identified

as carbapenemase producers by PCR.

**Diversity of CP-*Ec* isolates as assessed by whole genome sequencing.**

The genome sequences of the 713 CP-*Ec* isolates were first analyzed following *de novo* assembly. For

each isolate, we determined its MLST type (Achtman scheme), its phylogroup (ClermonTyping) and its

antibiotic resistance genes (ARG) content. We also searched for mutations in the quinolone resistance

determining regions (QRDRs) of *gyrA*, *parC* and *parE,* and for mutations in *ftsI*, *ompC* and *ompF* we

previously identified as associated with CP-*Ec* disseminated lineages (10) (table S1). F-NRC CP-*Ec* were

assigned to 168 different STs including six new allelic combinations. ST38 was the most prevalent ST

(n=92, 12.9 %), followed by ST10 (n=67, 9.4 %), ST410 (n=64, 9 %), and ST167 (n=34, 4.8 %). Ten

additional STs were represented by at least 10 isolates (Fig. S1A, table S1) and 154 STs with less than

ten isolates including 102 STs with a single isolate.

We next performed a core genome phylogeny following read mapping using strain MG1655

(NC_000913.3) as reference genome sequence. The phylogenetic tree, estimated from 372,238 core

SNPs was consistent with the results of phylogroup determination by using *in silico* ClermonTyping (12)

except for a few isolates (Fig. 2). In agreement with the MLST-based analysis, this tree showed a broad

diversity of CP-*Ec* isolates belonging to the eight phylogroups and three dominant clades

corresponding to CC10 (phylogroup A; including ST10, ST167 and ST617), CC23 (phylogroup C;

including ST410, ST88 and ST90) and ST38 (phylogroup D) with 161, 97, and 98 isolates respectively,

accounting for 49.9 % of the F-NRC CP-*Ec* isolates analyzed (Fig. 2). Phylogroup B2 isolates represented

11 % of the total isolates (n=80) and only 17 CP-ST131 isoaltes were identified. Fluctuations in the

proportions of the main STs could be observed during the four years of the analysis but no clear

tendency could be identified (Fig. S1A).

Infection-related and screening isolates were intermixed throughout the phylogeny (Fig. 2). However,

an enrichment in infection-related isolates was observed in phylogroup C (Pearson's Chi-squared test,

p<0.02, ddl1) and phylogroup B2 (Pearson's Chi-squared test p<0.0005, ddl1) (Fig. S1B). In phylogroup

B2, 5 out of 9 ST127 isolates, 5 out of 17 ST131 isolates and 7 out of 8 ST636 were responsible for UTIs

(Table S1).

**Diversity of antibiotic resistance genes carried by CP-*Ec* isolates**

The number of acquired ARGs was found to vary between 1 and 26 among the 713 CP-*Ec* isolates. The

median was higher in phylogroup C isolates (m=16) and lower in isolates from phylogroup B2 (m=3)

compared to other phylogroups (Medians: A: 9; B1: 9; D: 11) (Fig. 3A). An ESBL of the CTX-M family

was present in 40.7% (N=290) of the isolates, with a predominance of $bla_{CTX-M-15}$ gene (N=205) (Table

S1). Mutations in *gyrA*, *parC* and/or *parE* potentially leading to fluoroquinolone (FQ) resistance

occurred in 425 CP-*Ec* isolates (59.6%), with mutations in *gyrA*, *parC* and *parE* identified in 412 isolates,

309 and 261 isolates respectively (Table S1). Up to five mutations in QRDRs were identified in 19

isolates and 250 isolates had four mutations in QRDR, suggesting they have been submitted to a long-

term evolution under antibiotic pressure including FQ (Table S1). Globally a higher number of

mutations in QRDR was associated with a higher number of resistance genes (Fig. 3B), furthering the

link between the number of QRDR mutations and a likely evolution under antibiotic selective pressure

for CP-*Ec* isolates.

Four isolates from 2015, two ST648 collected in a same hospital at one-week interval, one ST216 and

one ST744, encoded a *mcr-9* gene conferring resistance to colistin (Table S1). Although not associated

with infection, the four isolates were MDR, carrying 13 to 20 acquired ARGs in addition to *mcr-9* and

carbapenemase genes ($bla_{VIM-1}$ for three of them, $bla_{NDM-1}$ and $bla_{OXA-48}$ for one). Determination of the

MIC for colistin for these four isolates were at the resistance breakpoint (MIC=2 mg/l).

**Diversity of carbapenemase genes**

Carbapenemase genes identified by WGS (Table S1) were in agreement with molecular data collected

by the F-NRC laboratory. The $bla_{OXA-48}$ and $bla_{OXA-181}$ genes were the most frequent carbapenemase

genes detected in 464 (65%) and 101 (14.1%) *E. coli* isolates respectively. Twenty-nine other isolates

carried minor $bla_{OXA-48-like}$ genes (Table S1). $bla_{NDM}$ family genes were detected in 14.9% (106/713) of

the CP-*Ec* isolates ($bla_{NDM-5}$ in 49 isolates and $bla_{NDM-1}$ in 41). Three additional $bla_{NDM}$ alleles were

identified: $bla_{NDM-7}$ (n=10) including a variant coding for a NDM7-like carbapenemase with a S24G

mutation, $bla_{NDM-4}$ (n=5), and $bla_{NDM-6}$ (n=1). Four different $bla_{VIM}$ alleles were detected in 18 isolates:

$bla_{VIM-1}$ (n=8), $bla_{VIM-4}$ (n=8), $bla_{VIM-2}$ (n=1), and $bla_{VIM-29}$ (n=1). Only five *E. coli* isolates expressed $bla_{KPC-2}$

(n=2) or $bla_{KPC-3}$ (n=3) alleles. In twelve isolates, two different carbapenemase genes were found;

$bla_{OXA-48}$ / $bla_{NDM-1}$ was the most frequent co-occurrence (n=5) (Table S1).

The analysis of the number of additional ARGs and of mutations in QRDR regions showed that,

compared with other carbapenemase genes the presence of $bla_{OXA-48}$ was frequently associated with

less resistant isolates (Fig. 4A, 4B). In particular it was the only ARG in 117 isolates among the 457

(24.5%) encoding this carbapenemase, including 23 infection-related isolates. Among those isolates,

only five showed mutations in QRDR. Conversely, only two $bla_{NDM-1}$ gene carrying isolates among 42

and one $bla_{NDM-5}$ gene carrying isolates among 49 carried no other ARG.

A temporal analysis of the proportion of isolates displaying $bla_{OXA-48}$ did not reveal significant variations

(Pearson's Chi-squared test, p=0.05, ddl= 3) during the 2012-2015 period (Fig. 4C). In contrast, the

frequency of $bla_{OXA-181}$ relatively to other alleles was found to significantly increase from 0 % in 2012

to 18.9% in 2015 (Pearson's Chi-squared test, p < 0.0002 ddl=3) (Fig. 4C).

We observed some association between the carbapenemase gene and the ST. Among the 22 STs with

at least seven F-NRC CP-*Ec* isolates, fourteen predominantly displayed $bla_{OXA-48}$ (Fig. 5A); $bla_{OXA-48}$ was

also predominant in STs represented by six or less isolates. In three lineages, ST410, ST940, and

ST1284, $bla_{OXA-181}$ was the predominant allele. These three STs grouped 74% of the $bla_{OXA-181}$ gene

carrying *E. coli* isolates from the F-NRC, and ST410 accounted for 50.5% of them. Finally, $bla_{OXA-204}$ was

the most prevalent allele in ST90 and $bla_{NDM-5}$ the predominant allele in ST636.

**Characteristics of the STs preferentially associated with F-NRC CP-*Ec* isolates.**

Eight among the 14 STs with more than 10 CP-*Ec* isolates (ST410, ST167, ST940, ST405, ST617, ST90,

ST1284, ST101) were characterized by a larger number of ARGs (median ≥ 10) and a larger number of

mutations in QRDR as compared to ST38, ST10, ST131, ST69, ST88 and ST represented by less than

seven isolates (Fig. 5 B, C). ST648 isolates have an intermediate position, being highly mutated in QRDR

but less rich in ARGs. Analysis of CP-*Ec* isolates from the F-NRC for polymorphisms in *ftsI* revealed that

131 (18.4%) had a four-AA-insertion between positions 333 and 334 of PBP3, the insertion being

particularly prevalent in ST101, ST167, ST410 and ST1284 (Sup. Table 1 and Fig. 5D). Similarly, *ompC*

alleles we previously characterized as modifying susceptibility to antibiotics were more prevalent in

some STs: an ST38-like *ompC* allele resulting from a recombination event or a G138D mutation were

present in 33 and 26 isolates belonging to six different ST, whereas the R195L mutation was observed

in 50 isolates all from ST410 (Table S1 and Fig. 5E). In addition, 144 isolates belonging to phylogroups

D and F possessed the ST38-like allele inherited vertically. On the other hand the *ompF* gene was

pseudogeneized in 47 isolates (Table S1).

To analyze the phylogenetic relationships between F-NRC Cp-*Ec* and other Cp-*Ec* isolates collected

worldwide, we built ST-based maximum-likelihood trees for the 14 STs with more than 10 isolates and

included to this analysis genome sequences publicly available (Table S2). Together with the analysis of

QRDR mutations and mutations in *ftsI* and *ompC*, this showed that 51 (80%) of the ST410 (Fig.S2A), 17

(49%) of the ST167 (Sup Fig.S2B), 3 (33%) of the ST405 (Sup Fig.S2C) and 8 (62%) of the ST101 (Fig.

S2D) F-NRC CP-*Ec* belonged to internationally disseminating MDR subclades enriched in CP we

previously identified (10). Subclades characterized by mutations in *ftsI* and *ompC,* mainly expressed

*bla*$_{OXA-181}$ (ST410) or *bla*$_{NDM}$ (other lineages). Other isolates belonging to these ST were dispersed on

their respective phylogenies and with no sign of clonal dissemination for most of them.

ST38 corresponded to the most represented ST (n=92) among the studied CP-*Ec*. The phylogenetic

reconstruction together with 314 additional non redundant CP-*Ec* genome sequences retrieved from

Enterobase (http://enterobase.warwick.ac.uk/) and 150 non-CP isolates from NCBI provided

evidence that 80.4% (n=67) of the French isolates clustered into four clades (Fig. 6). Three of them, G1

(n=35), G2 (n=25) and G4 (n=4) only contained isolates encoding *bla*$_{OXA-48}$, while G3 (n=7), included

isolates expressing *bla*$_{OXA-48}$ or its single nucleotide derivative *bla*$_{OXA-244}$. All but one isolates in G1

expressed *bla*$_{CTX-M-24}$, while isolates of G3 expressed *bla*$_{CTX-M-14b}$. The phylogenetic analysis provided

evidence of worldwide dissemination of G1, G3 and G4 clades and multiple introduction in France.

Strickingly, among the 28 isolates of the G2 clade 24 were collected in France and four in the

Netherlands, suggesting at this time a more regional dissemination. In none of the isolates of the four

clades, an IncL plasmid, that generally encode *bla*$_{OXA-48}$ (13), could be identified by PlasmidFinder (14).

It suggests a chromosomal integration as previously shown among ST38 isolates collected in the UK

(15). Analysis of two isolates of the G1 and G3 cluster (G1: GCA_900607445.1; G3: GCA_004759025.1),

whose genome sequences were completely assembled, confirmed that *bla*$_{OXA-48}$ and *bla*$_{OXA-244}$, in these

two isolates respectively, were chromosomally integrated. To assess the genetic support of *bla*$_{OXA-48}$ in

the two other clades, we have determined the complete genome sequence of two isolates belonging

to G2 (CNR65D6) and G4 (CNR85I8) by combining long-read Pacbio and Illumina sequencing and found

that both possess as chromosomally inserterted *bla*$_{OXA-48}$ gene. A fifth clade, composed of seven closely

related French isolates corresponded to a possible outbreak in the East of Paris between December

2014 and December 2015. These isolates are predicted to be highly resistant as they are carrying in

addition to *bla*$_{VIM-4,}$ *bla*$_{CTX-M-15}$, as well as 15 to 18 additional ARGs and five mutations in QRDRs.

ST10 *E. coli* isolates are commensals of a variety of mammals and bird species (16) and studies in

different contexts have shown that this ST is also associated with ARGs carriage (17, 18). The 67 F-NRC

ST10 CP-*Ec* isolates were distributed throughout the ST10 phylogeny. Nonetheless 31 isolates

belonged to a clade enriched with carbapenemase producers (Fig. S3A, in green) that mainly expressed

*bla*$_{OXA-48}$, with the exception of three isolates expressing *bla*$_{OXA-181}$, three expressing *bla*$_{NDM-5}$ and one

and *bla*$_{VIM-4}$. Two of the *bla*$_{OXA-181}$ and the three *bla*$_{NDM-5}$ expressing isolates were closely related, shared

a CTX-M-27 ESBL, four mutations in QRDR and a *ftsI* allele with a four AA YRIN insertion (Fig. S3A, in

red).

ST131 isolates are major contributors for extra-intestinal infections and the main responsible for the

dissemination of CTX-M-15 ESBL (19, 20). ST131 evolution has been thoroughly analyzed and four main

lineages A, B, C1 and C2 have been described (21). We identified only 17 ST131 CP-*Ec* isolates received

by the F-NRC (Fig. 2), ten belonged to lineage A, while the others were scattered in the three other

lineages (Fig. S3B). The main carbapenemase associated with these isolates was OXA-48 (N=10). We

also did not observe any association with the carriage of $bla_{CTX-M}$ genes, as only four ST131 CP-*Ec*

isolates carryied a $bla_{CTX-M-15}$ gene. These results are in accordance with the UK survey (8) and our

previous observations from sequences retrieved from public databases (10).

ST940 (phylogroup B1) and ST1284 (phylogroup A) represented 2.4% (n=17) and 1.9% (n=14) of the

sequenced isolates while they were poorly represented in sequence databases. For instance, in May,

2021, only 83 and 103 sequences genome sequences could be retrieved from Enterobase for these ST

respectively, but with 43% (n=36) and 22% (n=22) carrying a carbapenemase gene respectively. This

suggested that ST940 and ST1284 were associated with the dissemination of carbapenemase genes.

Phylogenetic trees were drawn by adding the non-redundant sequences from enterobase to those of

the F-NRC and NCBI. For ST940 (Fig. S4A) this revealed a well-differentiated sub-clade of 18 isolates

from Asia, Australia and Europe carrrying $bla_{NDM-5}$ and sharing mutations in *ftsI* and *ompC*. Isolates

from the F-NRC were distributed on the tree, showing that the over-representation of ST940 in France

did not result from local outbreaks. Three F-NRC isolates belonged to a second smaller sub-clade of

$bla_{NDM}$ expressing isolates, also mutated in *ftsI* and o*mpC*. In contrast, in ST1284, thirteen encoding

$bla_{OXA-181}$, were found to be closely related (Fig. S4B). Metadata analysis showed that 12 out of them

were recovered from the same health facility in the Paris suburb during a one-month period in 2015,

demonstrating a local outbreak origin. A fourteenth unrelated isolate expressed $bla_{NDM-5}$. The F-NRC

isolates belonged to a clade characterized by four mutations in QRDR determinants, a YRIN duplication

in PBP3, a G137D mutation in OmpC and containing additional OXA-181- or NDM-producing isolates.

Among the fifteen ST90 CP-*Ec* isolates, nine encoding the $bla_{OXA-204}$ allele were isolated in hospitals East

of Paris between August 2012 to April 2013 and suspected to be associated with the use of a

contaminated endoscope (22). Isolates closely related to this outbreak reappeared in 2014 (three

times), and in 2015 (twice), also in East of Paris located hospitals. They belonged to a MDR clade

characterized by four mutations in QRDR (Fig. S5A). Finally, in the four last STs with more than 10 CP-

*Ec* isolates (ST69, ST88, ST617, and ST648), French CP-*Ec* isolates were distributed throughout their

respective phylogenetic trees (FigS5B and C,S6A and B) with no sign of clonal dissemination, except for

a small cluster of five ST69 from two geographical regions and expressing *bla*OXA-48.

In addition to potential outbreaks detected in ST38 ($bla_{VIM-4}$), in ST90 ($bla_{OXA-204}$) and ST1284 ($bla_{OXA-}$

$_{181}$), we also found evidence for another potential outbreak among isolates belonging to the ST359.

The five isolates encoded $bla_{OXA-48}$ and $bla_{CTX-M-32}$ and had three mutations in QRDR. They were isolated

during January 2014, four of them in the south-eastern part of France and one in the Parisian suburbs.

**DISCUSSION**

The prevalence of CP-*Ec* is increasing worldwide. However whether this reveals conjugation event in

*E. coli* isolates by circulating CP-encoding plasmids or the emergence and dissemination of at-risk

clones is still largely unknown. Here, we have analyzed, by using WGS, 691 (87.8%) of the CP-*Ec* isolates

received by the F-NRC during the period 2012-2015 and 22 isolates from the Bicetre hospital strain

collection (2001-2011) to characterize both the diversity of carbapenemase genes and of CP-*Ec* isolates

in France. Altogether 713 CP-*Ec* were sequenced, representing to our knowledge the most extended

collection of CP-*Ec* sequenced, published to date.

The number of CP-*Ec* isolates sent to the F-NRC was found to strongly increase during the four-year

period of analysis, which might be a consequence of an increased circulation of CP-*Ec* in France but

also of an increased screening of potential CPE carriers at their admission at hospital. Indeed, while

the number of infection-related CP-*Ec* isolates was regularly increasing, their proportion compared to

the total number of received and sequenced isolates decreased by two-fold, with a clear change

observed between 2013 and 2014 (Fig. 1). This is likely a consequence of the implementation of the

recommendations on MDR screening of the French Public Health Advisory Board 2013 (23). However,

as CP-*Ec* isolates were sent on a voluntary basis by clinical laboratories, we cannot exclude some

sampling bias.

The most predominant carbapenemase allele was $bla_{OXA-48}$ (65%) followed by $bla_{OXA-181}$ (14.1%) which

detection constantly increased from no cases in 2012 to 73 in 2015. Next was $bla_{NDM-5}$ that was found

to progressively substitute to the $bla_{NDM-1}$ gene in frequency of detection (Fig. 4). The predominance

of $bla_{OXA-48-like}$ genes and particularly of $bla_{OXA-48}$, among CP-*Ec* was also noted in other studies (9, 24,25)

although in lower proportions. The small number of $bla_{KPC}$ genes detected in these studies is in contrast

with the situation reported in *K. pneumoniae* for many European countries (26). However, $bla_{OXA-48}$

was found the most prevalent allele in *K. pneumoniae* in France (2) and in the Netherlands (9). This

difference might result from a lower capacity for $bla_{KPC}$ encoding plasmids to conjugate to *E. coli* as

compared to pOXA-48 or to a higher fitness cost. It might also be linked to sampling differences as

different thresholds of carbapenem susceptibility might have been used to collect CRE isolates among

studies. Indeed OXA-48-like carbapenemases are generally associated with lower levels of carbapenem

resistance than KPC.

The number of ARGs identified in the CP-*Ec* isolates from the F-NRC varied from one (the

carbapenemase gene only) to 26, showing that carbapenemase genes were acquired not only in MDR

isolates but also in an *E. coli* population that can be considered as "naïve" relative to an evolution

under antibiotic selective pressure and resident of the intestinal microbiota. This was particularly true

for isolates producing $bla_{OXA-48}$ that were generally associated with a lower number of resistance genes

than isolates producing other carbapenemases, irrespectively of their clinical status (Fig. 4A). In

particular 118 out of the 121 (97.5%) isolates with no other ARG than the carbapenemase gene carried

$bla_{OXA-48}$. Therefore, the predominance of $bla_{OXA-48}$ among the carbapenemase genes identified in the

F-NRC collection might mainly rely on the higher conjugative transfer rate of pOXA-48 related plasmids

compared to those encoding other carbapenemases (27). pOXA48 has indeed been shown to rapidly

conjugate among *Enterobacterales* in hospitalized patients (28). This could also explain the high

frequency of $bla_{OXA-48}$ carrying isolates belonging to ST10, ST69, ST88, ST127, ST12 and ST58 (Fig. 5),

that carried a small number of resistance genes and QRDR mutations and among which four were

previously identified among the most frequent *E. coli* ST characterized from fecal samples (29).

On the other hand, MDR CP-*Ec* isolates may result from the transfer of carbapenem resistance genes

into isolates already selected through multiple exposures to antibiotic treatments that have already

acquired multi-resistance plasmids or chromosomal mutations increasing their intrinsic drug

resistance, such as mutations in QRDR, in *ftsI* and/or in porin genes. Alternatively, the circulation of

carbapenem-resistant lineages, such as the OXA-181-producing ST410 *E. coli* lineage, showing a stable

association between the lineage and the resistance gene, may also occur and significantly contribute

to the number of CP-*Ec* isolates collected. Discriminating between both alternatives would require a

more extended comparison of plasmids carried in these lineages after long-read sequencing similarly

to what been done in *K. pneumoniae* (30). Strikingly, the analysis of *ftsI* alleles coding for a PBP3 with

the four amino-acid duplication revealed a contrasted situation among CP-*Ec* isolates collected in

France, with the duplication in *ftsI* found in 72.3%, 77.8% and 67% of the isolates carrying $bla_{NDM-5}$

(34/47), $bla_{NDM-7}$ (7/9) and $bla_{OXA-181}$ (68/101) genes but in only 0.7% (4/458) of the isolates carrying

$bla_{OXA-48}$. We previously proposed that the fixation of mutations reducing the intrinsic susceptibiliy to

carbapenems might have favored the efficient conjugative transfer of plasmids carrying the

carbapenemase genes from other CPE species in the gut (10). The transconjugant would be selected

in a context of low biliary excretion of carbapenems or other ß-lactams after parenteral administration.

Increasing the proportion of the donor and recipient bacteria would therefore be less necessary for

plasmids with high conjugative rates, such as pOXA-48 (27). Alternatively, these mutations might also

have favored the plasmid maintenance by increasing the resistance level and selection of the isolates

during ß-lactam or carbapenem treament. Of particular concern in France are the OXA-181 producing

ST410 and the NDM (NDM-1, -5 and -7) ST167 lineages that significantly contribute to the circulation

of carbapenem resistance. However our study also reveals smaller lineages, that although less

frequently encountered in international studies, are nevertheless circulating. In contrast we did not

obtain evidence for clonal dissemination of CP-*Ec* ST131 lineages derived from the B2 clade responsible

for $bla_{CTX-M-15}$ gene dissemination (19, 20).

Despite their high frequency among the F-NRC collection, the ST38 isolates do not enter into one of

the two previous categories. Indeed, while they mainly express $bla_{OXA-48}$ or its single-point-mutant

derivative $bla_{OXA-244}$ (31), our phylogenetic analysis revealed that a majority of them belong to four

different lineages, including one mostly associated with a rapid dissemination in France. In contrast

with OXA-181 and NDM-producing lineages previously described, these lineages are associated with a

moderate number of ARGs and QRDR mutations. None of the ST38 isolates is mutated in *ftsI*, however

we previously demonstrated that all are characterized by an *ompC* allele that encodes a porin with a

reduced permeability to certain ß-lactams and has disseminated into unrelated lineages by

homologous recombination (10). A common feature of the four lineages is the integration of part of

the pOXA-48 plasmid carrying $bla_{OXA-48}$. This could have reduced a fitness cost of this plasmid and

facilitated the clonal expansion of these lineages. However, ST38 isolates were identified as

unfrequent colonizer of the GI tract (29, 32). Therefore additional features of this CP-*Ec* ST38 lineages

might have contributed to its dissemination.

In conclusion, by analysing genome sequences of CP-*Ec* isolates collected by the F-NRC we showed that

MDR lineages, enriched in carbapenemase-producing isolates are circulating in France, some of them,

such as the ST1284 isolates being associated with outbreaks. It also suggests that the evolutionary

trajectory may depend on the carbapenemase gene, $bla_{OXA-181}$ or $bla_{NDM}$ genes being more frequently

associated with the evolution of MDR *E. coli* lineages characterized by mutations in *ftsI* and *ompC*.

Surveillance of these mutations may be an important parameter in controling the circulation of MDR

lineages. On the other hand, carbapenemase genes are also frequently acquired through plasmid

dissemination from other bacterial species. Depending on the resistance background of the receiver

*E. coli*, this may lead to XDR or to isolates sensitive to a broad range of antibiotics. Finally, we also

observed a strong and rapid dissemination of ST38 isolates that might have been favored by a reduced

susceptibility to carbapenems linked to the ST38 *ompC* allele and by the chromosomal integration of

the carbapenemase gene. These results strengthen our model of different evolutionary trajectories

associated with the gain of carbapenemase genes (10). They also show that systematic genome

sequencing of CP-*Ec* and at a larger of CPE isolates, irrespective of their clinical or resistance status is

able to provide useful information not only on the circulation of MDR lineages, but also on the

propagation of resistance genes through horizontal gene transfer.

**MATERIAL AND METHODS**

**Isolate collection and sequencing.**

CP-*Ec* isolates analyzed in this study were collected by the F-NRC, mainly between the years 2012 to

2015. Twenty-two additional isolates we have received before the creation of the F-NRC in 2012, were

included in the analysis. Information on these isolates as the year of isolation, the region and

department in France and summary of their genomic features are reported in Table S1.

**Whole genome sequencing and analyses.**

DNA were extracted by using the Qiagen Blood and Tissue DNA easy kit. Sequencing libraries were

constructed by using the Nextera XT kit (Illumina) following the manufacturer instruction, and

sequenced with the Illumina HiSeq2500 or NextSeq500. FastQ files were trimmed for adaptors and

low-quality bases (setting the minimum base quality threshold to 25) with the Cutadapt fork Atropos

(33). *De novo* assemblies were generated from the trimmed reads with SPAdes v3.12.0 (34), using k-

mer sizes of 51, 71, 81, and 91, the coverage cut-off option was set to "auto" and the "careful" option

was activated. QUAST v2.2 (35) was used to assess the assembly quality and contigs shorter than 500

bp were filtered out for phylogenetic analyses. The ST38 isolates CNR65D6 and CNR85I8 were

sequenced to completion by using the long-read Pacbio technology. PacBio reads were assembled with

the RS_HGAP_Assembly.3 protocol from the SMRT analysis toolkit v2.3 (36) and with Canu (37),

polished with Quiver (36) and manually corrected by mapping Illumina reads using Breseq (38).

Assembled genomes were annotated with Prokka v1.9 (39). To analyze the F-NRC CP*Ec* isolates

belonging to the main ST recovered during the analysis in a more global context, their genome

sequences were combined to the assembled genome sequences from the same ST retrieved from the

NCBI database (July, 2019) and in some STs from Enterobase (http://enterobase.warwick.ac.uk/, April

2021). Retrieved genomes were annotated in the same way as the genome sequences of the F-NRC

isolates (Table S2).

To generate a core-genome phylogeny tree of all isolates, the best reference genome was first selected

among a set of 18 genomes analyzed by Touchon et al. (40) using the software refRank (41) and three

subsets of 100 randomly selected sequences from our study as an input. This led to the selection of

the genome sequence of strain MG1655-K12 (NC_000913.3) as the reference for read mapping and

SNP identification. Sequence reads for the 17 reference genomes (40) as well as for the genome

sequence of *Escherichia fergunsonii* strain ATCC 35469 (NC_011740.1), used as an outgroup were

simulated with ART (42). Trimmed sequencing reads were mapped against the MG1655-K12 genome

with BWA-MEM algorithm of the BWA v0.7.4 package (43). For SNP calling the Genome Analysis Toolkit

(GATK) v3.6.0 (44) was used with the following criteria, a minimum depth coverage (DP) of 10, a quality

by depth (QD) bigger than 2, a fisher strain bias (FS) below 60, a root mean square of the mapping

quality (MQ) above 40, and the mapping quality rank sum test (MQRankSum) and the read position

rank sum test (ReadPosRankSum) greater than -12.5 and -8 respectively. A Maximum-Likelihood tree

was estimated with RAxML v8.28 (45) using core-genome SNPs after removing positions in the

accessory genome identified with the filter_BSR_variome.py script from the LS-BSR pipeline (46).

For ST with more than 10 CP-*Ec* isolates from the F-NRC, a core genome alignment was generated with

Parsnp (47), by using a finished genome sequence as reference. A closely related isolate outside the

ST lineage was selected from the global phylogenetic tree including all F-NRC isolates and used as

outgroup to root the ST phylogenetic trees. Maximum-Likelihood (ML) trees were generated with

RAxML v8.28 (45) after removing regions of recombination with Gubbins (48). All graphic

representations were performed by using ITOL (49).

**MLST type, phylogroups, resistance gene and plasmid replicon identification.**

Sequence type was assigned to each assembly through a python script that relies on BLAST (10).

Antibiotic resistance genes (ARGs) and mutations in *gyrA, parC* and *parE* quinolone resistance

determining regions (QRDR) were identified with Resfinder 4.0 and PointFinder (50) run in local

respectively. The scripts and database (Retrieved: January 4, 2021) were downloaded from the

repositories of the Centre for Genomic Epidemiology (https://bitbucket.org/genomicepidemiology/).

The identified ARGs were manually reviewed to eliminate potential redundant ARG predicted at the

same genomic position. *mdf*(A) that is present in *E. coli* core genome was not taken into account.

Phylogroups were assigned by using EzClermont (51) run in local. Plasmid replicons were identified

with plasmidfinder run on each assembly (14). *ftsI, ompC and ompF* CDS sequences and *ompF*

promoter sequences were identified by using BlastN. Translated or nucleotide sequences were

clustered by cd-hit (cd-hit-v4.8.1) with an amino acid (FtsI, OmpC, OmpF) or nucleotide sequence

(*ompF* promoter) identity threshold of 1. Sequences of each cluster were aligned to detect mutations

in regions of interest: four amino-acid insertions between P333 and Y334 and E349K and I532L

mutations (FtsI), mutation modifying charge in L3 constriction loop, R195L mutation, nonsense or

frameshift mutations or OmpC sequence clustering with ST-38 OmpC sequences (OmpC), nonsense

mutations (OmpF) or mutations affecting one of the OmpR-boxes: mutation -46T/C in OmpR-F3 box

and mutation -75ΔT in OmpR-F2 box (*ompF* promoter).

**Statistical analysis.**

Pearson's chi-squared tests were performed by using standard libraries contained within the R

statistics package (http://www.R-project.org)

**Availability of data.**

All sequence data have been deposited at DDBJ/EMBL/GenBank (BioProject PRJEB46636) and

bioSample identifiers for the Illumina sequence data are listed in Table S1. Complete genome

sequences of CNR65D6 and CNR85I8 and the long-read sequencing data have been deposited at

DDBJ/EMBL/GenBank with the accession number ERZ3517884 (PacBio reads, ERR6414227) and

ERZ3518335 (ERS6682837 PacBio reads) respectively.

**ACKNOWLEDGMENTS**

This work was supported by grants from the French National Research Agency (ANR-10-LABX-62-IBEID, ANR-10-LABX-33, INCEPTION project (PIA/ANR-16-CONV-0005)) and from the European Union's Horizon 2020 research and Innovation Program under grant agrement No 773830 (Project ARDIG, One Health EJP). Pengdbamba Dieudonné Zongo is a scholar of Ed525 CDV, Sorbonne Université, Paris.

**REFERENCES**

1.  Medina M, Castillo-Pino E. 2019. An introduction to the epidemiology and burden of urinary tract infections. Ther Adv Urol 11:1756287219832172.
2.  Dortet L, Cuzon G, Ponties V, Nordmann P. 2017. Trends in carbapenemase-producing Enterobacteriaceae, France, 2012 to 2014. Euro Surveill 22.
3.  French National Reference Centre for antibiotic resistance, activity report 2019 - 2020. https://wwwcnr-resistance-antibiotiquesfr/bilans-dactivitehtml.
4.  Saleem AF, Allana A, Hale L, Diaz A, Salinas R, Salinas C, Qureshi SM, Hotwani A, Rahman N, Khan A, Zaidi AK, Seed PC, Arshad M. 2020. The Gut of Healthy Infants in the Community as a Reservoir of ESBL and Carbapenemase-Producing Bacteria. Antibiotics (Basel) 9:286.
5.  Bush K, Jacoby GA. 2010. Updated functional classification of beta-lactamases. Antimicrob Agents Chemother 54:969-76.
6.  Stoesser N, Sheppard AE, Pankhurst L, De Maio N, Moore CE, Sebra R, Turner P, Anson LW, Kasarskis A, Batty EM, Kos V, Wilson DJ, Phetsouvanh R, Wyllie D, Sokurenko E, Manges AR, Johnson TJ, Price LB, Peto TE, Johnson JR, Didelot X, Walker AS, Crook DW. 2016. Evolutionary History of the Global Emergence of the *Escherichia coli* Epidemic Clone ST131. MBio 7:e02162.
7.  Gauthier L, Dortet L, Cotellon G, Creton E, Cuzon G, Ponties V, Bonnin RA, Naas T. 2018. Diversity of carbapenemase-producing *Escherichia coli* isolates, France 2012-2013. Antimicrob Agents Chemother doi:10.1128/aac.00266-18.
8.  Ellaby N, Doumith M, Hopkins KL, Woodford N, Ellington MJ. 2019. Emergence of diversity in carbapenemase-producing *Escherichia coli* ST131, England, January 2014 to June 2016. Euro Surveill 24:1800627
9.  van der Zwaluw K, Witteveen S, Wielders L, van Santen M, Landman F, de Haan A, Schouls LM, Bosch T. 2020. Molecular characteristics of carbapenemase-producing *Enterobacterales* in the Netherlands; results of the 2014-2018 national laboratory surveillance. Clin Microbiol Infect 26:1412.e7-1412.e12.
10. Patino-Navarrete R, Rosinski-Chupin I, Cabanel N, Gauthier L, Takissian J, Madec JY, Hamze M, Bonnin RA, Naas T, Glaser P. 2020. Stepwise evolution and convergent recombination underlie the global dissemination of carbapenemase-producing *Escherichia coli*. Genome Med 12:10.
11. Roer L, Overballe-Petersen S, Hansen F, Schønning K, Wang M, Røder BL, Hansen DS, Justesen US, Andersen LP, Fulgsang-Damgaard D, Hopkins KL, Woodford N, Falgenhauer L, Chakraborty T, Samuelsen Ø, Sjöström K, Johannesen TB, Ng K, Nielsen J, Ethelberg S, Stegger M, Hammerum AM, Hasman H. 2018. *Escherichia coli* Sequence Type 410 Is Causing New International High-Risk Clones. mSphere 3:e00337-18
12. Beghain J, Bridier-Nahmias A, Le Nagard H, Denamur E, Clermont O. 2018. ClermonTyping: an easy-to-use and accurate *in silico* method for *Escherichia* genus strain phylotyping. Microb Genom 4:0.000192
13. Hendrickx APA, Landman F, de Haan A, Witteveen S, van Santen-Verheuvel MG, Schouls LM, The Dutch Cpe Surveillance Study G. 2021. bla (OXA-48)-like genome architecture among carbapenemase-producing *Escherichia coli* and *Klebsiella pneumoniae* in the Netherlands. Microb Genom 7:000512

14.  Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, Moller Aarestrup
F, Hasman H. 2014. In silico detection and typing of plasmids using PlasmidFinder and plasmid
multilocus sequence typing. Antimicrob Agents Chemother 58:3895-903.
15.  Turton JF, Doumith M, Hopkins KL, Perry C, Meunier D, Woodford N. 2016. Clonal expansion
of *Escherichia coli* ST38 carrying a chromosomally integrated OXA-48 carbapenemase gene. J
Med Microbiol 65:538-46.
16.  Leverstein-van Hall MA, Dierikx CM, Cohen Stuart J, Voets GM, van den Munckhof MP, van
Essen-Zandbergen A, Platteel T, Fluit AC, van de Sande-Bruinsma N, Scharinga J, Bonten MJ,
Mevius DJ. 2011. Dutch patients, retail chicken meat and poultry share the same ESBL genes,
plasmids and strains. Clin Microbiol Infect 17:873-80.
17.  Oteo J, Diestra K, Juan C, Bautista V, Novais A, Pérez-Vázquez M, Moyá B, Miró E, Coque TM,
Oliver A, Cantón R, Navarro F, Campos J. 2009. Extended-spectrum beta-lactamase-producing
*Escherichia coli* in Spain belong to a large variety of multilocus sequence typing types, including
ST10 complex/A, ST23 complex/A and ST131/B2. Int J Antimicrob Agents 34:173-6.
18.  Manges AR, Mende K, Murray CK, Johnston BD, Sokurenko EV, Tchesnokova V, Johnson JR.
2017. Clonal distribution and associated characteristics of *Escherichia coli* clinical and
surveillance isolates from a military medical center. Diagn Microbiol Infect Dis 87:382-385.
19.  Price LB, Johnson JR, Aziz M, Clabots C, Johnston B, Tchesnokova V, Nordstrom L, Billig M,
Chattopadhyay S, Stegger M, Andersen PS, Pearson T, Riddell K, Rogers P, Scholes D, Kahl B,
Keim P, Sokurenko EV. 2013. The Epidemic of Extended-Spectrum-beta-Lactamase-Producing
*Escherichia coli* ST131 Is Driven by a Single Highly Pathogenic Subclone, H30-Rx. MBio
4:e00377-13
20.  Nicolas-Chanoine MH, Bertrand X, Madec JY. 2014. *Escherichia coli* ST131, an intriguing clonal
group. Clin Microbiol Rev 27:543-74.
21.  Decano AG, Downing T. 2019. An *Escherichia coli* ST131 pangenome atlas reveals population
structure and evolution across 4,071 isolates. Sci Rep 9:17394.
22.  Potron A, Bernabeu S, Cuzon G, Pontiès V, Blanchard H, Seringe E, Naas T, Nordmann P, Dortet
L. 2017. Analysis of OXA-204 carbapenemase-producing Enterobacteriaceae reveals possible
endoscopy-associated transmission, France, 2012 to 2014. Euro Surveill 22:17-00048
23.  Lepelletier D, Berthelot P, Lucet JC, Fournier S, Jarlier V, Grandbastien B. 2015. French
recommendations for the prevention of 'emerging extensively drug-resistant bacteria' (eXDR)
cross-transmission. J Hosp Infect 90:186-95.
24.  Ortega A, Sáez D, Bautista V, Fernández-Romero S, Lara N, Aracil B, Pérez-Vázquez M, Campos
J, Oteo J. 2016. Carbapenemase-producing *Escherichia coli* is becoming more prevalent in
Spain mainly because of the polyclonal dissemination of OXA-48. J Antimicrob Chemother
71:2131-8.
25.  Grundmann H, Glasner C, Albiger B, Aanensen DM, Tomlinson CT, Andrasević AT, Cantón R,
Carmeli Y, Friedrich AW, Giske CG, Glupczynski Y, Gniadkowski M, Livermore DM, Nordmann
P, Poirel L, Rossolini GM, Seifert H, Vatopoulos A, Walsh T, Woodford N, Monnet DL. 2017.
Occurrence of carbapenemase-producing *Klebsiella pneumoniae* and *Escherichia coli* in the
European survey of carbapenemase-producing Enterobacteriaceae (EuSCAPE): a prospective,
multinational study. Lancet Infect Dis 17:153-163.
26.  David S, Reuter S, Harris SR, Glasner C, Feltwell T, Argimon S, Abudahab K, Goater R, Giani T,
Errico G, Aspbury M, Sjunnebo S, Feil EJ, Rossolini GM, Aanensen DM, Grundmann H. 2019.
Epidemic of carbapenem-resistant *Klebsiella pneumoniae* in Europe is driven by nosocomial
spread. Nat Microbiol 4:1919-1929.
27.  Potron A, Poirel L, Nordmann P. 2014. Derepressed transfer properties leading to the efficient
spread of the plasmid encoding carbapenemase OXA-48. Antimicrob Agents Chemother
58:467-71.
28.  León-Sampedro R, DelaFuente J, Díaz-Agero C, Crellen T, Musicha P, Rodríguez-Beltrán J, de la
Vega C, Hernández-García M, López-Fresneña N, Ruiz-Garbajosa P, Cantón R, Cooper BS, San

Millán Á. 2021. Pervasive transmission of a carbapenem resistance plasmid in the gut
microbiota of hospitalized patients. Nat Microbiol 6:606-616.
29.  Matsui Y, Hu Y, Rubin J, de Assis RS, Suh J, Riley LW. 2020. Multilocus sequence typing of
*Escherichia coli* isolates from urinary tract infection patients and from fecal samples of healthy
subjects in a college community. Microbiologyopen 9:1225-1233.
30.  David S, Cohen V, Reuter S, Sheppard AE, Giani T, Parkhill J, Rossolini GM, Feil EJ, Grundmann
H, Aanensen DM. 2020. Integrated chromosomal and plasmid sequence analyses reveal
diverse modes of carbapenemase gene spread among *Klebsiella pneumoniae*. Proc Natl Acad
Sci U S A 117:25043-25054.
31.  Emeraud C, Girlich D, Bonnin RA, Jousset AB, Naas T, Dortet L. 2021. Emergence and Polyclonal
Dissemination of OXA-244-Producing *Escherichia coli*, France. Emerg Infect Dis 27:1206-1210.
32.  Massot M, Daubié AS, Clermont O, Jauréguy F, Couffignal C, Dahbi G, Mora A, Blanco J, Branger
C, Mentré F, Eddi A, Picard B, Denamur E, The Coliville G. 2016. Phylogenetic, virulence and
antibiotic resistance characteristics of commensal strain populations of *Escherichia coli* from
community subjects in the Paris area in 2010 and evolution over 30 years. Microbiology
(Reading) 162:642-650.
33.  Didion JP, Martin M, Collins FS. 2017. Atropos: specific, sensitive, and speedy trimming of
sequencing reads. PeerJ 5:e3720.
34.  Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,
Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA.
2012. SPAdes: a new genome assembly algorithm and its applications to single-cell
sequencing. J Comput Biol 19:455-77.
35.  Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome
assemblies. Bioinformatics 29:1072-5.
36.  Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A,
Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome
assemblies from long-read SMRT sequencing data. Nat Methods 10:563-9.
37.  Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res
27:722-736.
38.  Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved microbes
from next-generation sequencing data using breseq. Methods Mol Biol 1151:165-88.
39.  Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068-9.
40.  Touchon M, Hoede C, Tenaillon O, Barbe V, Baeriswyl S, Bidet P, Bingen E, Bonacorsi S,
Bouchier C, Bouvet O, Calteau A, Chiapello H, Clermont O, Cruveiller S, Danchin A, Diard M,
Dossat C, Karoui ME, Frapy E, Garry L, Ghigo JM, Gilles AM, Johnson J, Le Bouguenec C, Lescat
M, Mangenot S, Martinez-Jehanne V, Matic I, Nassif X, Oztas S, Petit MA, Pichon C, Rouy Z, Ruf
CS, Schneider D, Tourret J, Vacherie B, Vallenet D, Medigue C, Rocha EP, Denamur E. 2009.
Organised genome dynamics in the *Escherichia coli* species results in highly diverse adaptive
paths. PLoS Genet 5:e1000344.
41.  Becker L, Fuchs S, Pfeifer Y, Semmler T, Eckmanns T, Korr G, Sissolak D, Friedrichs M, Zill E,
Tung ML, Dohle C, Kaase M, Gatermann S, Rüssmann H, Steglich M, Haller S, Werner G. 2018.
Whole Genome Sequence Analysis of CTX-M-15 Producing *Klebsiella Isolates* Allowed
Dissecting a Polyclonal Outbreak Scenario. Front Microbiol 9:322.
42.  Huang W, Li L, Myers JR, Marth GT. 2012. ART: a next-generation sequencing read simulator.
Bioinformatics 28:593-4.
43.  Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics 25:1754-60.
44.  McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler
D, Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. Genome Res 20:1297-303.

45.  Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30:1312-3.
46.  Sahl JW, Caporaso JG, Rasko DA, Keim P. 2014. The large-scale blast score ratio (LS-BSR)
pipeline: a method to rapidly compare genetic content between bacterial genomes. PeerJ
2:e332.
47.  Treangen TJ, Ondov BD, Koren S, Phillippy AM. 2014. The Harvest suite for rapid core-genome
alignment and visualization of thousands of intraspecific microbial genomes. Genome Biol
15:524.
48.  Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris SR. 2015.
Rapid phylogenetic analysis of large samples of recombinant bacterial whole genome
sequences using Gubbins. Nucleic Acids Res 43:e15.
49.  Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. Nucleic Acids Res 47:W256-w259.
50.  Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A, Allesoe RL, Rebelo
AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B, Werner G, Bender JK, Stingl K,
Nguyen M, Coppens J, Xavier BB, Malhotra-Kumar S, Westh H, Pinholt M, Anjum MF, Duggett
NA, Kempf I, Nykäsenoja S, Olkkola S, Wieczorek K, Amaro A, Clemente L, Mossong J, Losch S,
Ragimbeau C, Lund O, Aarestrup FM. 2020. ResFinder 4.0 for predictions of phenotypes from
genotypes. J Antimicrob Chemother doi:10.1093/jac/dkaa345.
51.  Waters NR, Abram F, Brennan F, Holmes A, Pritchard L. 2020. Easy phylotyping of Escherichia
coli via the EzClermont web app and command-line tool. Access Microbiol 2:acmi000143.
642

**Figure legends**

**Figure 1. Origin of the isolates received by the F-NRC during the 2012-2015 period**. **A**. Absolute

number of isolates per year from infection, screening or unknown origins, differentiating between

isolates that have or not been sequenced. **B**. Proportions of isolates from infection or from screening

or unknown origins showing that isolates from infection origin represent the same proportions among

total isolates that have been received by the F-NRC and isolates that have sequenced.

**Figure 2: Core genome phylogeny of CP-*Ec* isolates received by the F-NRC**. ML phylogeny of the 713

CP-*Ec* isolates was built with RAxML (45) from 372,238 core SNPs after sequence alignment on

MG1655-K12 (NC_000913.3) selected as the best reference. The genome sequence of *Escherichia*

*fergunsonii* strain ATCC 35469 (NC_011740.1) was used as an outgroup for the phylogenetic analysis.

Genomes from Touchon et al. (40) were also incorporated into the analysis. Genomic features are

indicated as in the figure key (left) from the inside to the outside circles: carbapenemases of the OXA,

NDM and other types, CTX-M ESBL, mutations in *gyrA* and *parC* QRDR region (FQ resistance), origin,

main ST, phylogroups. Reference genome from Touchon *et al*. are indicated as Touchon.

**Figure 3. Resistance gene content of the F-NRC CP-*Ec* isolates**. **A**. Box plot representation of the ARGs

content according to the phylogroup. The number of isolates belonging to each phylogroup is indicated

between brackets. The limits of the box indicate the lower and upper quartile. Outliers are indicated

by points above the maximal value. **B**: Relationships between the ARGs number and the number of

mutations in QRDR. In the box plot representations, the median is indicated by an horizontal bar and
the mean by a cross.

**Figure 4. Carbapenemase gene content of the F-NRC CP-*Ec* isolates**. **A**. Relationship between the
carbapenemase allele and the ARGs content. In the box plot representation, the median is indicated
by an horizontal bar and the mean by a cross. The limits of the box indicate the lower and upper
quartile. Outliers are indicated by points above the maximal value. **B**. Number of isolates carrying a
specific carbapenemase allele according to the number of QRDR mutations in their genomes. **C**. Per-
year evolution of the percentage of isolates carrying the different carbapenemase alleles in the 2012-
2015 period. Year of isolation are indicated as in the figure key (right). Given their small number (n=22),
strains isolated before 2012 are not indicated**.**

**Figure 5**. Features of the main ST associated with CP-*Ec* isolates in France. **A**. Proportion of the isolates
according to the carbapenemase allele in each ST; **B**. Box plot representation of the ARGs content
according to the ST; **C.** Proportion of isolates according to the number of mutations in QRDR; **D** :
Proportion of isolates with a specific PBP3 allele. PBP3 characterized by a YRIP, YRIK or YRIN insertion
between positions 333 and 334, the YRIN insertion is generally associated with a A413V mutation
(YRIN-L) and less frequently with a E349K mutation (YRIN-K-L) (10); **E.** Proportion of isolates carrying
specific mutations in *ompC:* R195L, G138D or acquisition of a ST38-like *ompC* allele by recombination
(10). For each ST, the phylogroup, the total number of isolates and the number of isolates associated
with urinary tract infections are indicated in the upper panel. Intermediate (Inter) and minor ST
indicate ST represented by 4-6 and 1-3 isolates, respectively.

**Figure 6. Core genome phylogeny of ST38 isolates.** ML Phylogeny was based on genome sequences
of 92 CP-*Ec* from the F-NRC and 464 genome sequences retrieved from Enterobase and from the NCBI
database including 331 carrying a carbapenemase gene. A core genome (2,900,000 nt) alignment of
the *de novo* assemblies on the sequence of GCA_005886035.1 used as a reference was performed by
using Parsnp (47); ML phylogeny was built with RAxML (45) from 6,170 core SNPs after removing
recombined regions with Gubbins (48). The genome sequence of CNRC6O47 (ST963) was used as an
outgroup for the phylogenetic analysis. F-NRC isolates are indicated by red triangles (inner circle).
Other genomic features are indicated as indicated in the figure key (left) from the inside to the outside
circles: carbapenemases of the OXA, NDM and other types, number of mutations in *gyrA* and *parC*
QRDR region (FQ resistance), number of ARGs, mutations in *ftsI, ompC, ompF*, geographical origin. The
four OXA-48-like clades (G1, G2, G3, G4) clustering most French isolates are colored in blue, red, violet

and green. A fifth clade (G5) corresponding to a possible outbreak of VIM-4 isolates in the East of Paris

is colored in brown

**Supplementary figures**

**Figure S1**. Per-year analysis of the origin of the isolates received by the F-NRC. **A**. Proportions of the

isolates belonging to the different ST. The total number of isolates is indicated between brackets. The

large proportion of ST90 isolates collected in 2012 is linked to a local outbreak. **B**. Proportions of the

isolates collected in screening and infection situations as a function of the year and the phylogroup.

The absolute number of isolates for each class is indicated inside the bars.

**Figure S2**. Core genome phylogenies of the main ST characterized by clades disseminating

internationally. A. ST410; B. ST167; C. ST405; D. ST101. Phylogenies were based on genome sequences

of A: 64 CP-*Ec* from the F-NRC and 146 genome sequences from the NCBI database;B: 35 CP-*Ec* from

the F-NRC and 134 genome sequences from the NCBI database; C. 15 CP-*Ec* from the F-NRC and 145

genome sequences from the NCBI database; D. 13 CP-*Ec* from the F-NRC and 194 genome sequences

from the NCBI database. Core genome (ST410: 3,522,000 nt; ST167: 3,276,000 nt; ST405: 3,685,000

nt; ST101: 3,774,000 nt) alignments of the *de novo* assemblies on the sequences of GCA_001442495.1

(ST410), GCA_003028815.1 (ST167), GCA_002142675.1 (ST405), GCA_002163655.1 (ST101) used as

reference were performed by using Parsnp (47); ML phylogeny was built with RAxML (45) from 6,176

(ST410), 10,393 (ST167), 37,482 (ST405), 19,967(ST101) core SNPs after removing recombined regions

with gubbins (48). The genome sequences of CNR93E7 (ST88), CNR93D10 (ST746), CNR73I9 (ST115),

CNR93I2 (ST906) were used as outgroups for the phylogenetic analyses of ST410, ST167, ST405 and

ST101 respectively. The origin from the F-NRC is indicated by red triangles close to the isolate name.

Other genomic features are indicated as indicated in the figure key (left) from the inside to the outside

lines: carbapenemases of the OXA, NDM and other types, number of mutations in *gyrA* and *parC* QRDR

region (FQ resistance), number of ARGs, mutations in *ftsI, ompC, ompF*, geographical origin.

**Figure S3**. Core genome phylogenies of ST10 and ST131 isolates. **A.** ST10 phylogeny based on the

genome sequences of 67 CP-*Ec* from the F-NRC and 153 genome sequences from the NCBI. **B.** ST131

phylogeny based on the genome sequences of 17 CP-*Ec* from the F-NRC and 462 genome sequences

from the NCBI. Core genome (ST10: 813,000 nt; ST131: 1,641,000 nt) alignments of the *de novo*

assemblies on the sequence of MG1655 (ST10) or GCA_000285655.3 (ST131) used as references were

performed by using Parsnp (47); ML phylogeny was built with RAxML (45) from 20,245 (ST10) and

14,366 (ST131) core SNPs after removing recombined regions with Gubbins (48). The genome

sequences of CNR93D10 (ST746) and CNRAL47G10 (ST640) were used as outgroups for the

phylogenetic analyses of ST10 and ST131, respectively. F-NRC isolates are indicated by red triangles

(inner circle). Other genomic features are indicated as indicated in the figure key (left) from the inside

to the outside circles: carbapenemases of the OXA, NDM and other types, number of mutations in *gyrA*

and *parC* QRDR region (FQ resistance), number of ARGs, mutations in *ftsI*, *ompC*, *ompF*, geographical

origin. The ST10 clade enriched in CP-*Ec* is indicated in green and the sub-clade with the YRIN insertion

in PBP3 in red. Note that the circle for *ftsI* mutations is absent in ST131, as no YRIN-like duplication in

PBP3 was identified among the analyzed isolates.

**Figure S4.** Core genome phylogeny of ST940 and ST1284 isolates. **A:** ST940 phylogeny based on

genome sequences of 17 CP-*Ec* from the F-NRC and 88 genome sequences from Enterobase and the

NCBIdatabase **B:** ST1284 phylogeny based on genome sequences of 14 CP-*Ec* from the F-NRC and 60

genome sequences from Enterobase and the NCBI database. A core genome (ST940: 3,639,000 nt;

ST1284: 3,779,000 nt) alignment of the *de novo* assemblies on the sequence of ST940: ESC_LB2149

and ST1284: ESC_LB2152AA (from Enterobase) used as references was performed by using Parsnp

(47); ML phylogeny was built with RAxML (45) from 13,859 (ST940) and 2,009 (ST1284) core SNPs after

removing recombined regions with Gubbins (48). The genome sequences of CNR98G1 (ST3022) and

MG1655 (ST10) were used as outgroups for the phylogenetic analyses of ST940 and ST1284,

respectively. F-NRC isolates are indicated by red triangles first column on the left. Other genomic

features are as indicated in the figure key (left) from the left to the right columns: carbapenemases of

the OXA, NDM and other types, number of mutations in *gyrA* and *parC* QRDR region (FQ resistance),

number of ARGs, mutations in *ftsI*, *ompC*, *ompF*, geographical origin.

**Figure S5**. Core genome phylogenies of ST69, ST88 and ST90. isolates. **A:** ST90 phylogeny based on the

15 CP-*Ec* from the F-NRC and 38 genome sequences from the NCBI database**; B:** ST69 phylogeny based

on the 16 CP-*Ec* from the F-NRC and 244 genome sequences from the NCBI database; **C.** ST88

phylogeny based on the 15 CP-*Ec* from the F-NRC and 99 genome sequences from the NCBI database.

Core genome (ST69: 2,735,000 nt; ST88: 3,472,000 nt; ST90: 3,801,000nt) alignments of the *de novo*

assemblies on the sequence of GCA_002443135.1 (ST69), GCA_002812685.1 (ST88) or

GCA_001900635.1 (ST90) used as references were performed by using Parsnp (47); ML phylogeny was

built with RAxML (45) from 4,596 (ST69), 12,557 (ST88) or 2,878 (ST90) core SNPs after removing

recombined regions with Gubbins (48). The genome sequences of CNR33D9 (ST394), CNR83B9 (ST410)

and CNR88B9 (ST847) were used as outgroups for the phylogenetic analyses of ST69, ST88 and ST90,

respectively. F-NRC isolates are indicated by red triangles (first column on the left). Other genomic

features are indicated as indicated in the figure key (left) from the left to the right columns:

carbapenemases of the OXA, NDM and other types, number of mutations in *gyrA* and *parC* QRDR

region (FQ resistance), number of ARGs, mutations in *ftsI*, *ompC*, *ompF*, geographical origin.

**Figure S6**. Core genome phylogeny of ST617 and ST648 isolates based on genome sequences of **A**:

ST617 genome sequences of 15 CP-*Ec* from the F-NRC and 65 genome sequences from the NCBI

database; **B**: ST648 genome sequences of 14 CP-*Ec* from the F-NRC and 126 genome sequences from

the NCBI database. A core genome (ST617: 3,407,000; ST648: 3,501,000 nt) alignment of the *de novo*

assemblies on the sequences of GCA_002142695.1 (ST617) or GCA_004138645.1 (ST648) used as

references was performed by using Parsnp (47); ML phylogeny was built with RAxML (45) from 15,030

(ST617) or 3,782 (ST648) core SNPs after removing recombined regions with Gubbins (48). The genome

sequences of CNR93D10 (ST746) and CNR71A8 (ST1485) were used as outgroups for the phylogenetic

analyses of ST617 and ST648, respectively. F-NRC isolates are indicated by red triangles close to the

isolate name. Other genomic features are indicated as indicated in the figure key (left) from the left to

the right columns: carbapenemases of the OXA, NDM and other types, number of mutations in *gyrA*

and *parC* QRDR region (FQ resistance), number of ARGs, mutations in *ftsI*, *ompC*, *ompF*, geographical

origin.

Supplementary Tables

**Table S1**: Main characteristics of the F-NRC CP-Ec

**Table S2**: Main characteristics of genome sequences used for phylogenetic reconstructions

Fig. 1

## OXA
- OXA-181
- OXA-204
- OXA-232
- OXA-244
- OXA-48
- NA

## NDM
- NDM-1
- NDM-4
- NDM-5
- NDM-6
- NDM-7
- NA

## Other CPase
- KPC-2
- KPC-3
- VIM-1
- VIM-2
- VIM-29
- VIM-4
- NA

## CTX-M
- CTX-M-15
- CTX-M-24
- CTX-M-14b
- Other
- NA

## FQDR
- GyrA/ParC mutation
- NA

## Origin
- Clinical
- Screening
- Touchon
- unknown

## STSimplified
- ST10
- ST101
- ST1284
- ST131
- ST167
- ST38
- ST405
- ST410
- ST617
- ST648
- ST69
- ST88
- ST90
- ST940
- NA

## Phylogroups
- A
- B1
- B2
- C
- D
- E
- F
- G

Tree scale: 0.1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

**Study**
- ◄ Fr-NRC

**OXA-like**
- OXA-181
- OXA-48
- OXA-244

**NDM-like**
- NDM-1
- NDM-5

**Other CPase**
- KPC-2
- KPC-3
- VIM-4
- IMP-4

**Plasmids**
- ● No plasmid

**CTX-M**
- CTX-M-14
- CTX-M-14b
- CTX-M-15
- CTX-M-24
- CTX-M-27
- Other CTX-M
- CTX-M-15+other CTX-M

**QRDR**
- ▮ QRDR mutations

**ARGs**
- ▮ ARGs

**FtsI insertion**
- YRIK
- YRIN-1
- YRIN-2

**OmpF mutation**
- pseudogene
- promoter: (ompR-F3: -46C/T)
- promoter: (ompR-F2: -62T/C)
- promoter: (OmpR-F2: -75 insT )
- promoter: (ompR-F2: -75ΔT )

**Zone**
- East Asia
- Europe
- Middle East
- North Africa
- North America
- Oceania
- South America
- South Asia
- South-East Asia
- Sub-Saharan Africa

G1

G2

G3

G4

G5

Tree scale: 0.001

A

B

Fig. S1

**Study**
▶ Fr-NRC

**OXA-like**
- OXA-48
- OXA-181
- OXA-232
- OXA-244
- OXA-204
- Other OXA-like

**NDM-like**
- NDM-1
- NDM-5
- NDM-7
- NDM-like

**other CPase**
- KPC-2
- KPC-3
- KPC-like
- VIM-1
- VIM-4
- VIM-like
- IMP-4
- IMP-like

**QRDR mutations**
- QRDR mutations

**ARGs**
- ARGs

**Insertion in PBP3**
- YRIN-K-L
- YRIN-L
- YRIK
- YRIP
- YTIP

**OmpC mutation**
- ST38-like
- G137D
- R195L
- pseudogene

**OmpF mutation**
- pseudogene
- promoter: (OmpR-F3: -46C/T)
- promoter: (OmpR-F2: -75 insT )

**Zone**
- East Asia
- Europe
- Middle East
- North Africa
- North America
- Oceania
- South America
- South Asia
- South-East Asia
- Sub-Saharan Africa
- NA

A

ST410

Tree scale: 0.01

B

ST167

Tree scale: 0.01

C

ST405

Tree scale: 0.001

D

ST101

Tree scale: 0.001

Fig. S2

**Study**
- ▶ Fr-NRC

**OXA-like**
- OXA-48
- OXA-181
- OXA-232
- OXA-244
- OXA-204
- Other OXA-like

**NDM-like**
- NDM-1
- NDM-5
- NDM-7
- NDM-like

**other CPase**
- KPC-2
- KPC-3
- KPC-like
- VIM-1
- VIM-4
- VIM-like
- IMP-4
- IMP-like

**QRDR mutations**
- QRDR mutations

**ARGs**
- ARGs

**Insertion in PBP3**
- YRIN-K-L
- YRIN-L
- YRIK
- YRIP
- YTIP

**OmpC mutation**
- ST38-like
- G137D
- R195L
- pseudogene

**OmpF mutation**
- pseudogene
- promoter: (OmpR-F3: -46C/T)
- promoter: (OmpR-F2: -75 insT )

**Zone**
- East Asia
- Europe
- Middle East
- North Africa
- North America
- Oceania
- South America
- South Asia
- South-East Asia
- Sub-Saharan Africa
- NA

A

ST10

Tree scale: 0.001

B

ST131

Tree scale: 0.001

Fig. S3

**Study**
▶ Fr-CNR

**OXA-like**
■ OXA-181
■ OXA-48

**NDM-like**
■ NDM-5
■ NDM-like
■ NDM-1
■ NDM-7

**QRDR mutations**
■ QRDR mutations

**ARGs**
■ ARGs

**Insertion in PBP3**
■ YRIK
■ YRIN-K-L
■ YRIN-L

**OmpC**
■ mut G137D
■ ompC-ST38

**OmpF mutation**
■ pseudogene
■ promoter (ompR-F2: ΔA68)

**Zone**
■ East Asia
■ Europe
■ Middle East
■ North America
■ Oceania
■ South America
■ South Asia
■ South-East Asia
■ Sub-Saharan Africa
■ North Africa
□ NA

A

Tree scale: 0.001

ST940

B

Tree scale: 0.001

ST1284

Fig. S4

## Study
▶ Fr-NRC

## OXA-like
- OXA-48
- OXA-181
- OXA-204

## NDM-like
- NDM-1

## other CPase
- KPC-3

## QRDR mutations
- QRDR mutations

## ARGs
- ARGs

## Zone
- East Asia
- Europe
- Middle East
- North Africa
- North America
- Oceania
- South America
- South Asia
- South-East Asia
- Sub-Saharan Africa
- NA

A

ST69          Tree scale: 0.001

B

ST88          Tree scale: 0.001

C

ST90          Tree scale: 0.001

Fig. S5

Fig. S6