Global diversity and distribution of prophages is lineage-specific
within the *Ralstonia solanacearum* plant pathogenic bacterium
species complex

Samuel T. E. Greenrod[1,*], Martina Stoycheva[1], John Elphinstone[2], Ville-Petri Friman[1,*]

[1]Department of Biology, University of York, York, UK

[2] Fera Science Ltd, National Agri-Food Innovation Campus, Sand Hutton, York, UK

*Corresponding authors

10
Abstract

Ralstonia solanacearum is a destructive plant pathogenic bacterium and the causative agent of
bacterial wilt disease, infecting over 200 plant species worldwide. In addition to chromosomal
genes, its virulence is mediated by mobile genetic elements including integrated DNA of
bacteriophages, i.e. prophages, which may carry fitness-associated auxiliary genes or modulate
host gene expression. Although experimental studies have characterised several prophages that
shape *R. solanacearum* virulence, the global diversity, distribution, and wider functional gene
content of *R. solanacearum* prophages is unknown. In this study, prophages were identified in a
diverse collection of 192 *R. solanacearum* draft genome assemblies originating from six
continents. Prophages were identified bioinformatically and their diversity investigated using
genetic distance measures, gene content, GC, and total length. Prophage distribution was
characterised using metadata on *R. solanacearum* geographic origin and lineage classification
(phylotypes), and their functional gene content was assessed by identifying putative prophage-
encoded auxiliary genes. In total, 343 intact prophages were identified, forming ten genetically
distinct clusters. These included five prophage clusters belonging to the *Inoviridae*, *Myoviridae*,
and *Siphoviridae* phage families, and five uncharacterised clusters, possibly representing novel,
previously undescribed phages. The prophages had broad geographical distribution being
present across multiple continents. However, they were generally host phylogenetic lineage-
specific, and overall, prophage diversity was proportional to the genetic diversity of their hosts.
The prophages contained a myriad of auxiliary genes involved in metabolism and virulence of
both phage and bacteria. Our results show that while *R. solanacearum* prophages are highly

diverse globally, they make lineage-specific contributions to the *R. solanacearum* accessory
genome, which could result from shared coevolutionary history.

Keywords: *Ralstonia solanacearum*, prophage, plant pathogenic bacterium, mobile genetic
element, coevolution, diversity

Introduction

Bacteriophages, or phages for short, are viruses that infect bacteria. They outnumber bacteria by
up to six orders of magnitude (1) and through mutualistic and antagonistic interactions are thought
to have a significant impact on bacterial population dynamics and evolution (2,3). While some
phages are obligately lytic, killing their host after successful infection (4), other phages have a
lysogenic life cycle where phage genetic material integrates into the host chromosome forming a
prophage. Lytic phages can impose strong bottom-up density regulation of bacteria across
different ecosystems, driving nutrient turnover by infecting their host bacteria (5,6). They can also
drive bacterial diversification through frequency-dependent selection (5), whilst selecting for
phage resistance evolution via the acquisition of phage defence systems (7) and cell membrane
alterations that disrupt phage infection (8). As phage resistance mechanisms are often associated
with fitness trade-offs (9,10), these evolutionary changes can further alter phage-bacteria
population dynamics leading to eco-evolutionary feedbacks in microbial communities (for review
see (11)). In contrast, temperate prophages tend to have a more modest effect on bacterial
population dynamics as they often replicate during bacterial cell division and become lytic only
when induced by environmental stresses such as UV irradiation or antibiotic treatment (12,13).
Temperate phages can drive bacterial evolution by facilitating the lateral transfer of auxiliary
genes which are expressed in the prophage state (14). These genes have been found to be
associated with bacterial antibiotic resistance (15,16), competitiveness (17), and virulence (18,19),
being responsible for the disease severity of important bacterial pathogens including
shigatoxigenic *Escherichia coli* (20) and *Vibrio cholerae* (21). Temperate phages can also affect
host fitness by changing gene expression or knocking out host genes after inserting into the
genome (22), providing resistance to secondary phage infection, termed "superinfection immunity"
(23), and by acting as hotspots of recombination (24). Prophages are thus often beneficial for their
host bacteria and overrepresented in the genomes of pathogenic bacteria (18).

While prophages have been studied extensively with human opportunistic bacteria, they are also
important for the fitness and evolution of plant pathogenic bacteria (19,25). For example,

prophages of some of the most destructive plant pathogens including *Pseudomonas* (26), *Xylella*
(27), and *Xanthomonas* (28) spp., have been associated with auxiliary genes that encode plant
immune response inhibitors (29,30), secretion system proteins, degradative enzymes, and toxin
exporters (31–33). Plant pathogen competitiveness can also be mediated by prophages that
encode competitor-repressing bacteriocins (34), provide resistance to environmental stresses
such as toxic metal ions (35) and antimicrobials (36), or encourage survival during nutrient scarcity
by increasing metabolic potential (37). However, the distribution, diversity, and functional potential
of prophages is still relatively understudied at the pangenome-level with plant pathogenic bacteria.
Prophage pangenome studies based on a representative set of genomes of one species are
useful for determining the total prophage diversity and auxiliary gene content potentially linked to
host bacterium fitness. They can also help us to understand if certain prophages can be
considered to belong to core (shared by all host strains) or accessory genome of their hosts
(shared only by a subset of strains). Whilst this information could provide an insight into prophage
geographical distribution, it could also reveal whether certain prophages are evenly spread
throughout their host phylogeny, or if they show lineage-specific patterns, potentially due to local
adaptation or shared coevolutionary history.

Prophage pangenome studies in plant pathogenic bacteria have been primarily carried out with
Dickeya spp. and *Pectobacterium* spp. from the soft rot Pectobacteriaceae family. These studies
have focused on prophage diversity and their auxiliary gene content, identifying prophage clusters
harbouring bacterial genes involved in ecological fitness and virulence (35,38,39). Notably, whilst
prophages are generally only present in approximately half of Pectobacteriaceae strains, nearly
all prophages contain fitness-associated bacterial ORFs (35,38,39). Subsequently, variability in
prophage presence, movement, and auxiliary gene content has been linked to variation in
virulence, swimming motility, and cellulase production (38). Recently, prophage diversity was also
assessed in the plant pathogenic bacterium *Ralstonia solanacearum*, a causative agent of
bacterial wilt (40), through an analysis of 120 *R. solanacearum* genomes available in the NCBI
database (41). This study revealed many characterised and novel prophages, the latter of which
belonged to either the virulence-associated *Inoviridae* family or had no similarity to known phages.
It also highlighted prophage-encoded auxiliary genes with potential roles in host virulence, cellular
metabolism, environmental stress resistance, and antibiotic resistance. By analysing multiple
bacterial strains, these studies suggest that prophages may be fundamental drivers of phenotypic
diversity in bacterial populations. However, our understanding is still based on relatively small
number of genomes (35,38,39) derived either from local (38) or publicly available databases
(35,39,41), which are often susceptible to sampling biases. As a result, the total diversity,

distribution, and auxiliary gene content of prophages in plant pathogenic bacteria is likely
underestimated.

Here, we conducted a comprehensive pangenome analysis of the prophages within the *Ralstonia*
solanacearum species complex (RSSC), which has a broad host range of over 200 plant species
within 50 families (42,43). *R. solanacearum* strains are genetically diverse and are classified into
four lineages, termed phylotypes (44,45), which generally follow their geographical location of
isolation: Phylotype I includes strains originating primarily from Asia, Phylotype II from America,
Phylotype III from Africa and surrounding islands in the Indian ocean, and Phylotype IV from
Indonesia, Japan, and Australia (44). Recently, the four phylotypes have been redefined as three
separate species, including *R. solanacearum sensu stricto* (Phylotype II), *R.*
pseudosolanacearum (Phylotypes I and III) and an array of *R. syzygii* subspecies (Phylotype IV)
(46). Considerable variation exists between and within *R. solanacearum* lineages regarding their
metabolic versatility (47) and tolerance to environmental stresses, including starvation and low
temperatures (48,49). *R. solanacearum* virulence and competitiveness are largely determined by
a diverse accessory genome. RSSC strains contain an arsenal of host immunity-modifying type
III effectors (50), many of which are phylotype-specific (51) and linked to variation in disease
severity ranging from 100% mortality to symptomless infections (52). Furthermore, *R.*
solanacearum strains contain a high diversity of *Inoviridae* and *Myoviridae* prophages which are
known to have a direct influence on their virulence. For example, temperate phages of the
Inoviridae family, within two clades labelled RSS-type and RSM-type, can differentially affect host
virulence by up or downregulating the expression of virulence factors. Infection with RSS-type
phages, such as φRSS1, increases host virulence factor expression by inducing cell aggregation,
activating the quorum sensing-mediated master regulator PhcA (53). However, infection with
RSM-type phages, such as φRSM3, knocks out pathogenicity due to the activity of a prophage-
encoded repressor which inhibits the expression of virulence factors (54). Recent metabolic
models have suggested that the production of virulence factors restricts bacterial proliferation and
reduces metabolic versatility (55). Therefore, virulence-reducing prophages such as φRSM3
could also affect bacterial fitness by increasing bacterial growth rate and the range of metabolic
substrates it can utilize. This is supported by competitiveness assays where infection with the
RSS-RSM intermediate φRS551, which reduces virulence, increased *R. solanacearum*
competitiveness (34). In contrast to *Inoviridae* phages, infection with *Myoviridae* phages, such as
φRSA1, φRsoM1USA, and φRSY1, appears to have a limited impact on host virulence (56–58),
despite φRSY1 lysogens showing increased twitching motility and aggregation frequency (57),
typically indicative of enhanced virulence (53). However, sequence homology between *Myoviridae*

phage genomes is disrupted by low GC content "AT islands", which are hotspots of recombination and have been shown to contain virulence factors in other prophages (59). As experimental studies investigating temperate *Myoviridae* phages in *R. solanacearum* have only analysed isolates from single locations, such as fields in Hiroshima, Japan (φRSA1) (58) and Florida, USA (φRsoM1USA) (56), *Myoviridae* phages may contribute to *R. solanacearum* fitness through a diverse accessory genome that has remained undetected during single sample experimental studies.

The large diversity of the strains within the complex and the economic losses associated with the disease make RSSC a salient target for prophage pangenome analyses (34,53,54). While a recent study made a significant contribution on understanding RSSC prophages using publicly available *R. solanacearum* genomes (41), it had a sampling bias with low representation of strains originating from Africa and Europe, missing a subset of hosts which are cold-adapted (48). In our study, *R. solanacearum* prophages were investigated using a new representative collection of 192 *R. solanacearum* draft genome sequences. These included isolates from all four phylotypes and six continents, with extensive sampling from Africa and Europe. We specifically aimed to: i) characterise the global diversity and distribution of prophages in *R. solanacearum,* ii) assess whether prophages are spread throughout their host phylogeny or show host lineage-specific distribution; and iii) investigate prophage auxiliary gene content. Prophages were identified using the PHASTER web server (https://phaster.ca/) and their diversity assessed by calculating genetic distances using Mash and by comparing gene content, GC content, and lengths. Their distribution was characterised by assessing continent and lineage-specificity, with potential prophage-host coevolution determined by comparing prophage and host genetic dissimilarities. Finally, prophage auxiliary gene content was investigated through gene annotation and comparisons with bacterial metabolism and virulence databases.

Materials and methods

R. solanacearum hosts, sequencing, and genome assembly

R. solanacearum prophage hosts were selected from Protect and the National Collection of Plant Pathogenic Bacteria (NCPPB) and other reference strains maintained at Fera Science Ltd. Genomic DNA extraction was performed on 384 isolates using Qiagen DNeasy Blood and Tissue Kit (DNeasy® Blood & Tissue Handbook, Qiagen, Hilden, Germany, 2020) followed by quantification of double stranded DNA product using Quantit dsDNA Assay Kit Broad range and

Nanodrop (Thermo Fisher Scientific, Waltham, MA, USA). Host DNA was sequenced using
Illumina MiSeq at the Earlham Institute, UK. Sequence reads quality was assessed using FastQC
(60) and trimming of adapters and low-quality ends was performed using Trimmomatic (v0.39)
(61). Genomes were then assembled into draft assemblies using Unicycler (v0.4.8) on strict mode
(62). To classify the genomes, a pangenome analysis was performed on 357 high quality genome
assemblies plus 48 complete genomes downloaded from NCBI Genbank (Accessions available
in Supplementary Data) and *R. picketti* 12b used as an outgroup. Using Panaroo (v1.2.4) (63)
with strict mode and MAFFT aligner (64), a core genome alignment was generated. Phylogenetic
tree was then constructed with IQ-TREE (65) and GTR+G4 model and genomes forming clusters
were assigned to the phylotype of strains with known phylotype (Fig. S1 and Supplementary
Data). As more than half of the available strains were from phylotype IIB that shared high genetic
similarity, a subsample of 192 *R. solanacearum* draft genome sequences representing all four
phylotypes from six continents were chosen for prophage analysis (Table S1).

Prophage sequence identification

Putative prophage regions were identified in *R. solanacearum* draft genomes using the web
server PHASTER (PHAge Search Tool Enhanced Release) (66) which automatically classified
them into intact (score > 90), questionable (score 70-90) and incomplete (score < 70) prophages
based on their size and the presence of phage-like and phage cornerstone genes (for example,
'capsid', 'head', 'plate', 'tail', 'coat', 'portal' and 'holin'). Only intact prophages were used in
downstream analysis.

Prophage diversity analysis

Initially, intact prophage diversity was investigated by analysing prophage gene content to search
for shared phage core genes, which could have been used in phylogenetic tree construction.
However, no shared core genes were identified, and so prophage diversity was instead assessed
using a multifaceted, alignment-free approach (Mash) (67). Firstly, prophage genetic distances
were calculated based on the presence of shared k-mers using Mash v.2.2 (67). Mash distance
matrices were generated using the "mash triangle" command with sketch size = 10,000, with
similar prophages identified using Euclidean clustering. A Mash distance neighbour-joining (NJ)
tree was constructed using Mashtree v.1.2.0 (68) with default parameters. Secondly, prophage
gene content profiles were compared through gene annotation using VIGA v.2.7.16 (69) with
parameters E-value < e^{-5} and amino acid identity > 30%. Prophage pangenome analysis was

carried out using Roary v.3.13.0 (70). Finally, prophage GC content and lengths were determined
and compared using SeqKit v.0.12.0 (71).

Prophage taxonomic identities were determined using three different approaches. Firstly, Mash
distances were calculated between identified prophages and known phages in the Mash RefSeq
and NCBI Virus RefSeq databases, with successful hits determined using a p-value $< e^{-50}$
threshold. Secondly, BLASTn was used to compare identified prophages with a local blast
database, constructed using all available NCBI Virus RefSeq genomes, using BLAST+ v.2.10.0
(72) with an E-value $< e^{-50}$ threshold. Thirdly, known *R. solanacearum* phage genomes were
downloaded from the NCBI Virus RefSeq database and included in the prophage Mash distance
NJ tree to determine relatedness between known prophages and ones identified in this study
based on their clustering.

Determination of global prophage distribution and diversity in *R. solanacearum*

Prophage global and pangenome-level distribution was assessed using metadata on *R.*
solanacearum host geographical origin and phylotype classification (Table S1). Phylotype
classification was determined based on the whole genome phylogeny clustering rather than egl
sequence similarity as described earlier (45). Due to low sample sizes for phylotypes III and IV,
prophage host phylotype distributions were only investigated using hosts from phylotypes I, IIA,
and IIB (174/192 hosts). Prophage host phylotype distributions were determined by comparing
prophage presence and absence against a maximum likelihood (ML) phylogenetic tree, which
was constructed using concatenated host core genes (found in > 90% of hosts) identified with
Panaroo v.1.2.3 (63) and aligned and visualised using ClustalO v.1.2.4 (73) and IQTREE v.1.6.6
(65), respectively.

The potential historical signal of prophage-host coevolution was assessed by comparing genetic
dissimilarity between *R. solanacearum* host and its prophages, using the *R. solanacearum* ML
tree for host dissimilarity and Bray-Curtis dissimilarity measure for prophage content dissimilarity,
which accounted for presence, absence, and relative abundance of prophages in host genomes.
A pairwise prophage Bray-Curtis dissimilarity matrix was generated using the QIIME
beta_diversity script and was used to construct a UPGMA tree with the QIIME upgma_cluster
script. A tanglegram between the *R. solanacearum* ML tree and the prophage Bray-Curtis
UPGMA tree was generated with functions in the R 'ape' package, using the 'phytools' package
to rotate the *R. solanacearum* ML tree to minimise connected lines crossing between the trees.
Congruence between the *R. solanacearum* ML tree and the prophage Bray-Curtis UPGMA tree

was assessed using Procrustes Approach to Cophylogenetic Analysis (PACo) (74) in R. Briefly,
cophenetic distance matrices were constructed using the prophage Bray-Curtis and *R.*
solanacearum phylogenetic trees. The distance matrices were then compared using PACo to
determine the statistical significance of tree congruence using a Procrustean super-imposition of
the sum of squared 10,000 network randomizations under the "r0" randomization model.

R. solanacearum genetic dissimilarity was further compared with prophage dissimilarity with a
linear mixed model using functions in the R 'nlme' package. The model used average host genetic
dissimilarity as predictor variable, the average prophage dissimilarity as response variable, and
host phylotype as a random effect. Average *R. solanacearum* genetic dissimilarity was
determined by generating a pairwise Mash distance matrix of hosts within each phylotype and
calculating the average pairwise Mash distance for each host. Average prophage dissimilarity
was determined by generating a pairwise prophage Bray-Curtis dissimilarity matrix of hosts within
each phylotype and calculating the average pairwise Bray-Curtis dissimilarity for each host. To
remove non-linearity, the average *R. solanacearum* genetic dissimilarity was log-transformed.

Putative auxiliary gene analysis

Putative auxiliary genes were initially identified through manual curation of prophage annotations
generated using VIGA (v.2.7.16) (69) with parameters E-value $< e^{-5}$ and amino acid identity $> 30\%$
and Roary (v.3.13.0) (70). They were further investigated through BLASTn of prophage genomes
against the CAZy (http://www.cazy.org/) (75), the Ralsto T3E
(https://iant.toulouse.inra.fr/bacteria/annotation/site/prj/T3Ev3/) (51), and the PHI-base
databases (http://www.phi-base.org/) (76) with parameters E-value $< e^{-5}$. Comparisons within
databases generally provided multiple overlapping hits for each putative auxiliary gene.
Therefore, in all searches, only the top hit was used if multiple hits overlapped. Prophage
disruption of type III effectors was determined by generating a circular genome map of
comprehensively annotated UY031 strain chromosome (NCBI accession: NZ_CP012687) with
GView (77) and mapping prophages using the GView Blast Atlas function. UY031 gene product
labels were determined using the NCBI GenBank annotation. UY031 was chosen as a refence
genome as it has the closest complete genome to phylotype IIB strains that only showed potential
disruption of type III effectors by prophages.

Data visualisation and statistical analysis

Statistical analyses and data visualisation were carried out using Microsoft Excel v.2102, R v.4.0.3
and RStudio v1.4.1103. Putative prophage genome sizes and GC content were compared using

a Mann-Whitney *U* test. Comparisons of intact prophage number, *R. solanacearum* genetic

dissimilarity, and prophage dissimilarity between phylotypes were compared using Kruskal-Wallis

tests (with Dunn's test for pairwise comparisons) and linear regression. Graphs and heatmaps

were made using the R 'ggplot2' v.3.3.3 and R 'pheatmap' v.1.0.12 packages, respectively. The

Mash NJ and ML phylogenetic trees were visualised using the R 'ggtree' package v.2.1.4.

Results

Prophages are commonly found in *R. solanacearum* genomes

Prophages were identified in 192 *R. solanacearum* hosts sampled from six continents,

representing phylotypes I (56), IIA (22), IIB (96), III (8), and IV (10) (Table S1). In total, 532

prophage regions were identified across all hosts, including 343 intact prophages, 72

questionable regions, and 117 incomplete prophages as determined based on PHASTER criteria

(66) (Table S2). Intact prophages, which are predicted to contain cornerstone phage genes and

are hence likely inducible, were found in 173 host genomes (90.1%). Intact prophages had

significantly lower GC content than incomplete prophages (Mann-Whitney: w = 45803; n1 = 189;

n2 = 343; $p < 0.001$) with an average GC content of 63.0% (± 0.15, se) compared to 64.9% (±

0.17, se) (Fig. 1a). Incomplete prophage GC content was also closer to the average host GC

content of 66.8% (± 0.012, se). Moreover, intact prophages had significantly greater genome sizes

than incomplete prophages (combined questionable regions and incomplete prophages) (Mann-

Whitney: w = 14581; n1 = 189; n2 = 343; $p < 0.001$) with an average length of 24.4kb (± 0.6kb,

se) compared to 14.3kb (± 0.5kb, se) (Fig. 1b). Prophage GC content and length distributions

were multimodal, possibly indicating that *R. solanacearum* prophages include genetically distinct

groups. Polylysogeny, where more than one intact prophage is integrated into a genome, was

found in 139 hosts (72.3%). On average, *R. solanacearum* hosts contained 2.8 putative prophage

regions and 1.8 intact prophages per genome. The number of intact prophages per genome was

significantly different between phylotypes (Kruskal-Wallis: x^2 = 14.26; d.f = 4, $p < 0.05$). However,

pairwise comparisons using Dunn's test suggested that only one comparison was significantly

different with phylotype IIB having more prophages per genome than phylotype IIA (Fig. S2, $p <$

0.05). In contrast, the number of incomplete prophages was significantly different between

phylotypes (Kruskal-Wallis: x^2 = 98.85; d.f = 4, $p < 0.001$) with significant differences observed in

all pairwise comparisons ($p < 0.05$).

Figure 1. *R. solanacearum* **prophages have a multimodal length and GC content distribution.** (**a**) GC content and (**b**) length distributions of incomplete (combined incomplete and questionable) and intact *R. solanacearum* prophages. In (**a**), numbers in legend brackets show the total number of identified prophages.

***R. solanacearum* genomes contain ten genetically distinct prophage clusters**

The diversity and similarity of intact prophages was assessed by calculating prophage genetic

distances based on the presence of shared k-mers using Mash, where genetically similar

prophages were expected to have lower Mash distances. *R. solanacearum* prophages formed

ten clusters (labelled A-J and four singletons, Fig. 2), each representing genetically similar phage

groups. Two clusters, A and B, were largely clonal, predominantly containing prophages with

Mash distances equalling zero. However, the remaining clusters were more diverse containing

prophages with a range of Mash distances. Cluster E had low Mash distances with prophages in

cluster B, indicating they may be related. Prophage clusters were further visualised using a Mash

distance neighbour-joining tree, which supported the clusters identified with Euclidean distances,

and were further verified by comparing their gene content, GC content, and length (Fig. 3).

Prophage pangenome content was determined through gene annotation, identifying a total of

1,649 unique genes. Each prophage cluster had a unique gene content profile (Fig. 3), containing

between 89-100% cluster-specific genes (Table 1). However, no core prophage genes (found in

> 90% of isolates) were identified. Most prophage clusters also had distinct GC content and length

boundaries (Fig. 3; Table 1), except for cluster D which had high gene content, GC content, and

length variation.

Figure 2. Prophages are clustered into ten separate groups and four singletons (one phage per group). Pairwise Mash distance heatmap of *R. solanacearum* prophages, clustered using Euclidean distances. Clusters are labelled with both letters (A-J) and coloured bands. Singletons are combined into one white band. Prophage similarity (dark grey) decreases with increasing Mash distances (white).

Prophage clusters include both known and potentially novel, uncharacterised phages

The taxonomic identities of the prophage clusters were determined using the NCBI Virus RefSeq

and Mash RefSeq databases. A total of 163 prophages (47.5%) from five clusters (B, D, E, H, J)

belonged to the *Inoviridae*, *Myoviridae*, and *Siphoviridae* phage families within the Caudovirales

order (Fig. 3; Table S2). Clusters B and E represented the *Inoviridae* family; Cluster B contained

the RSM-type phages with a large clonal group identified as φRS551, and a smaller group

identified as φRSM3. Cluster E contained RSS-type prophages identified as φRSS0, φRSS1,

φRSS30, in addition to φPE226, and φp12J. Cluster D represented the *Myoviridae* family

containing prophages identified as φRSA1, φRSY1, and φRsoM1USA. Cluster H and J

prophages, identified as φRS138 and Unclassified J respectively, belonged to the *Siphoviridae*

family. Known *R. solanacearum* temperate phages aggregated with clusters in the same families

(Fig. S3), supporting prophage identity determination. The remaining 184 prophages (52.5%) from

five clusters (A, C, F, G, I) had no sequence similarity with phages present in the NCBI Virus

RefSeq or Mash RefSeq databases (last accessed April 2021), showing no clear clustering with any known phages. Therefore, these clusters may represent novel, uncharacterised prophage groups.

Table 1. Genetic characteristics of identified prophage clusters.

Prophage cluster	Proportion of genes that are cluster-specific	Average GC content (% ± se)	Average length (kbp ± se)	Taxonomic classification
A	166/194 (86%)	62.96 ± 0.04	35.19 ± 0.16	Uncharacterised
B	92/95 (97%)	59.35 ± 0.17	13.41 ± 0.37	*Inoviridae*
C	273/299 (91%)	66.11 ± 0.07	22.65 ± 0.56	Uncharacterised
D	323/323 (100%)	65.56 ± 0.17	31.51 ± 1.38	*Myoviridae*
E	67/68 (99%)	62.26 ± 0.24	8.97 ± 1.55	*Inoviridae*
F	86/86 (100%)	66.41 ± 0.14	15.17 ± 1.67	Uncharacterised
G	220/220 (100%)	62.75 ± 0.31	48.37 ± 2.71	Uncharacterised
H	36/36 (100%)	64.90 ± 0	31.80 ± 0	*Siphoviridae*
I	27/27 (100%)	60.00 ± 0.04	14.48 ± 1.41	Uncharacterised
J	85/85 (100%)	64.25 ± 0.55	31.75 ± 6.25	*Siphoviridae*

Prophages have broad geographical distributions reflecting the host evolutionary history

The global distribution of *R. solanacearum* prophages was determined by assessing their presence and absence across different continents, including Africa, Asia, Europe, North America, South America, and Oceania (Fig. 4). Phage families had broad geographical distributions; *Inoviridae* and *Myoviridae* prophages were distributed across all six continents, and *Siphoviridae* prophages were found in the Americas and Asia. Individual prophages were generally found in 2-3 continents, while some prophages were more widespread. The *Inoviridae* phage φRS551, *Myoviridae* phage φRSA1, and novel prophages Unclassified A and Unclassified C, were each found in all six continents. Only two low abundance prophages, Unclassified B and φRSS0, were continent-specific, being found exclusively in Asia.

Despite broad geographical distributions, prophages tended to have higher abundances in certain continents; φRS551 and Unclassified A were primarily found in hosts from Africa (φRS551, 34%;

Figure 3. Prophage clusters show unique gene content, GC content, and length profiles. From top-down: Prophage Mash distance neighbour-joining tree (see Appendix Fig. A1), coloured and labelled with phage families. Letters and coloured bars refer to prophage clusters from Figure 2. Blue and grey panel shows the presence (blue) and absence (grey) of genes within the prophage pangenome, shown as rows in order of gene prevalence. Bottom panels show prophage length (blue) and GC content (orange).

Unclassified A, 33%) and Europe (φRS551, 46%; Unclassified A, 45%), and φRSY1 and

φRsoM1USA were primarily found in North (φRSY1, 45%; φRsoM1USA, 56%), and South

America (φRSY1, 27%; φRsoM1USA, 22%). Prophages were also absent from certain

continents, and for example, φRSY1 and φRsoM1USA were not found in any host bacteria from

Africa or Europe, whilst Unclassified C was abundant in all continents except for Europe.

Therefore, *R. solanacearum* prophages appear to mainly follow continent borders.

As *R. solanacearum* phylotypes tend to have different geographical origins (44), we investigated

if the presence and absence of prophages was associated with specific *R. solanacearum*

phylotypes (Fig. 5a). *Inoviridae* prophages were predominantly found in phylotype I genomes,

which exclusively contained φPE226, φRSS0, φRSS1, φRSS30, and 5/6 φRSM3 prophages.

However, φRS551 was predominantly found in phylotype IIB and φp12J was found in IIA hosts.

Moreover, the *Myoviridae* prophages φRSY1 and φRSA1 were primarily found in phylotype I,

whilst φRsoM1USA was found in phylotype IIB and IIA hosts. The *Siphoviridae* prophages

φRS138 and Unclassified J were only found in phylotype IIA. Prophages within the same family

tended to be mutually exclusive; only two *R. solanacearum* hosts contained multiple *Inoviridae* or

Figure 4. Prophages have broad geographical distributions. World map overlaid with pie charts of prophage relative abundances in six continents (from left to right: North America, South America, Europe, Africa, Oceania, and Asia). Black lines link pie charts to continents. Numbers in brackets refer to the number of prophages found in hosts sampled from each continent.

multiple *Myoviridae* prophages. Unclassified A, similar to φRS551, was found predominantly in

phylotype IIB with only low abundances in phylotypes I and IIA. However, it was also present in

phylotype III (3/8) and IV (4/10) hosts and so may be more widespread. Unclassified G was

exclusively found in phylotype I and Unclassified F and I were primarily found in phylotype IIB and

IIA hosts, respectively. Notably, Unclassified C was generalist being found more evenly

distributed between phylotypes and with six hosts containing two prophage copies. *R.*

solanacearum prophages therefore appear to be mainly phylotype-specific, with only a few

prophages being associated with multiple host phylotypes.

Genetically similar host bacteria are associated with similar prophage profiles

In addition to harbouring unique prophages, *R. solanacearum* phylotypes also differed in their

overall prophage profiles (Fig. 5a; Fig. S4). For example, phylotype I genomes contained 14

different prophage types, with over 75% being found in less than 15% of phylotype I hosts (Fig.

S4a) in addition to highly prevalent prophages such as φRSY1, φRSA1, and Unclassified C group.

Phylotype IIA hosts only contained 9 prophage types, with 67% found in less than 15% of hosts

and only φRsoM1USA and Unclassified C groups had relatively high prevalence (Fig. S4b). In

contrast, Phylotype IIB hosts predominantly contained two high abundance prophages, φRS551

and Unclassified A, which were present in 81% and 83% of hosts, respectively (Fig. S4c).

Figure 5. Prophages are phylotype-specific and have diversity proportional to the genetic diversity of their hosts. (**a**) Maximum likelihood tree of *R. solanacearum* isolates from phylotypes I, IIA, and IIB (with phylotype III and IV clades collapsed – see Fig. S1) midpoint rooted and annotated with prophage presence (dark grey) and absence (white). Coloured bars on left show phylotype clustering within *R. solanacearum* tree. Coloured bars on top show prophage clusters, labelled with phage families. (**b**) Average *R. solanacearum* dissimilarity (log-transformed), measured using average pairwise Mash distances, versus average prophage dissimilarity, measured using average pairwise prophage Bray-Curtis distances. Points are coloured by phylotype. Bottom-right box shows regression equation, R^2 statistic, and p-value. Regression line is plotted. (**c**) Tanglegram of *R. solanacearum* maximum likelihood tree and prophage Bray-Curtis UPGMA tree. Blue lines connect the same labels on each tree with horizontal lines supporting tree congruence and crossed lines indicating tree incongruence.

Interestingly, the diversity of prophages within each phylotype appeared to reflect the branching on the *R. solanacearum* phylogenetic tree (Fig. 5a), indicating that there may be an association between *R. solanacearum* host and prophage genetic dissimilarities. We tested this by i) a linear regression and ii) congruence analysis between the *R. solanacearum* phylogenetic maximum-likelihood and prophage dissimilarity UPGMA trees. In support of our hypothesis, phylotype IIB hosts had both significantly lower prophage dissimilarity (Kruskal-Wallis: x^2 = 66.250.9; df = 2; p < 0.001) (Fig. S5a), and significantly lower genetic diversity than phylotypes IIA and I (Kruskal-Wallis: x^2 = 100.6; df = 2; p < 0.001) (Fig. S5b). As a result, host genetic dissimilarity explained a significant amount of variation in prophage dissimilarity based on linear regression (R^2 = 0.81, p < 0.001, Fig. 5b). This association was further verified by comparing the *R. solanacearum* phylogenetic and a prophage dissimilarity UPGMA trees, which showed significant congruence (M^2_{xy} = 0.296, p < 0.001, N = 10,000, Figure 5c). Therefore, prophages appear to be diverging in tandem with their hosts. Importantly, we found that tree congruence was not driven by individual clades as significant congruence was also present in phylotype I (M^2_{xy} = 0.001, p < 0.001, N = 10,000), IIA (M^2_{xy} = 0.003 p = 0.0176, N = 10,000), and IIB strains (M^2_{xy} = 0.001, p < 0.001, N = 10,000) when tested independently. These results suggest that genetically similar hosts had more similar prophage profiles, possibly reflecting a shared coevolutionary history or local adaptation.

Prophages encode various putative auxiliary genes linked with metabolism and virulence

Prophages are known to encode various auxiliary genes with potential benefits for bacterial fitness (18–22). Therefore, prophage gene content was analysed for auxiliary genes through gene annotation using VIGA and comparisons with bacterial metabolic (CAZy) and virulence gene databases (T3E and PHI). Most prophage genes encoded unannotated hypothetical proteins (56.8%) and cornerstone phage proteins related to structural components (25.8%), replication (7.8%), and lysis (3.4%) (Fig 6). Prophages also contained many genes with potential auxiliary functions, which made up 4% of the prophage pangenome, including transcriptional regulators, DNA methyltransferases, bacterial cell division proteins, and stress tolerance proteins. Moreover, several prophage genes with potential roles in bacterial metabolism and virulence were identified.

Putative auxiliary metabolic genes primarily encoded glycosyl hydrolases and glycosyl transferases, the latter of which were exclusively found in *Inoviridae* prophages. Glycosyl hydrolases tended to be specific to phage family or phage type, with *Inoviridae* RSM-type phages exclusively encoding proteins with GH1|GH95 domains and *Myoviridae* prophages all encoding proteins containing LysM domains, as well as domains involved in alkaline and neutral invertase

and lysozyme proteins. However, *Siphoviridae* and novel, uncharacterised prophages all encoded
proteins with either peptidoglycan lytic transglycosylase or GH13_30 domains.

Putative virulence genes predominantly encoded type III effectors, transcriptional regulators, and
membrane transporters. These were mainly found in *Inoviridae* prophages although the
Myoviridae prophage φRSY1 was found to contain two type VI secretion system proteins and the
type III effector RipS4. Further, the *Siphoviridae* prophage φRS138 was found to carry the type
III effector RipH3 and Unclassified J contained a histone-like DNA binding protein. In the novel,
uncharacterised prophages, putative virulence genes were only found in Unclassified A,
Unclassified F, and Unclassified G groups. Unclassified F and Unclassified G encoded an HNS-
like regulatory protein and an acyl hydroxylase, respectively. Moreover, Unclassified A, like
Unclassified J, encoded a histone-like DNA binding protein in addition to the type III effector
RipS5. As RipS5 is pseudogenised in phylotype IIB strains (78), which predominantly contain
Unclassified A, the position of Unclassified A relative to the RipS5 coding sequence was assessed
in the *R. solanacearum* strain UY031. Unclassified A was found to be flanked by two type III
effector regions, one of which was annotated as skwp5, otherwise known as RipS5 (79) (Fig. S6).
This suggests that, in phylotype IIB hosts, RipS5 may be disrupted by an Unclassified A prophage.

Similar to the auxiliary metabolic genes, some putative virulence genes were also phage family
or phage type specific, with the *Inoviridae* RSM-type phages all encoding the replication-
associated recombination protein RarA, whilst the RSS-type phages all encoded a zot-like toxin.
Moreover, all *Myoviridae* phages encoded a DNA methyltransferase. These genes were found in
multiple phylotypes and may be under more general selection than other genes which were phage
cluster specific. Nonetheless, prophages appear to encode many putative auxiliary genes, with
potential lineage-specific functional contributions to the *R. solanacearum* accessory genome.

Figure 6. Prophages encode many putative auxiliary genes involved in different bacterial processes. Top bar chart: stacked bar chart showing prophage gene content as a proportion of total prophage genes, coloured by putative function. Numbers above bars represent total number of genes in each phage species. Heatmaps: putative auxiliary genes identified through comparisons of prophage CDS with bacterial metabolism and virulence databases coloured by absolute abundance. Different colours reflect different databases used (labelled on the right). Putative gene functions are labelled on the left. Bottom bar chart: stacked bar chart showing the phylotype distribution of isolates containing each prophage.

Discussion

While prophages are known to affect plant pathogen fitness by mediating host growth, competitiveness, and virulence (29–37,41,80), only very little is known about their diversity and distribution at the pangenome-level. In this study, we analysed the prophage content of the plant pathogenic *R. solanacearum* bacterium using a representative, global collection of new 192 draft genome assemblies. Prophages were found in all screened host genomes, forming ten genetically distinct prophage clusters. While some of these could be assigned to known prophages based on databases, no matches were found for five clusters, which could represent novel prophages. Interestingly, while prophages had broad geographical distributions, they showed phylotype-specific associations and genetically similar hosts had similar prophage profiles over the whole RSSC. Several potential auxiliary genes potentially linked to *R. solanacearum* metabolism and virulence were also identified, and some of these were unique to specific prophage clusters and host phylotypes. Together, our results advance knowledge on the global *R. solanacearum* prophage diversity and distribution at the pangenome level.

Putative prophage elements were identified in all hosts and included both intact and incomplete prophages, the latter of which were shorter and had higher GC content that was closely matched with the GC content of their bacterial hosts. These incomplete prophages may represent "grounded" prophages which have lost the ability to excise themselves from the bacterial chromosome (24), becoming truncated and ameliorating to their hosts' nucleotide usage (81,82). On average, *R. solanacearum* hosts contained 1.8 intact prophages per genome, which is an intermediate number compared to other plant pathogens, ranging from less than one per genome in soft rot Pectobacteriaceae (35) to more than five in *Xylella fastidiosa* (27). These results generally support a previous analysis of *R. solanacearum* prophages which found that strains contained 1.2 intact prophages per genome (41). However, the slightly higher average number of intact prophages in our study could reflect higher sampling of phylotype IIB strains which we found had a higher average number of intact prophages than strains from all other phylotypes except for phylotype IV.

Intact prophages had a multimodal length and GC content distribution, suggestive of multiple genetically distinct prophage groups. Indeed, ten prophage clusters were identified which had different gene content profiles, GC content, and lengths. Whilst five clusters belonged to the *Inoviridae*, *Myoviridae*, and *Siphoviridae* families, over half of the prophages identified were novel, uncharacterised prophages with no sequence similarity to known phages in public databases. These results are consistent with a previous genomic analysis of *R. solanacearum* prophages

(41) which identified both known and novel prophages and demonstrates that *R. solanacearum*
has high prophage diversity. Most of the prophages we identified, except for φRsoM1USA, were
the same as those found previously. This was surprising given that previous analyses have relied
on publicly available databases which predominantly contain hosts from South America and Asia,
whilst we used a global genome collection representing all six continents. Therefore, most of the
R. solanacearum prophage diversity appears to be represented in South America and Asia.
However, whether this is due to prophage diversity being particularly high in these continents, or
due to prophages being ubiquitous, remains unclear.

Prophages within the same family tended to be mutually exclusive, with intra-family polylysogeny
only observed in two hosts. This contradicts previous studies on *Myoviridae* phages which have
found polylysogeny-promoting homology disruption at attachment sites (58), and the co-existence
of both φRSA1 and φRSY1 prophages within the genome of *R. solanacearum* strain EJAT-1458
(57). Polylysogeny could be inhibited by prophage-mediated superinfection immunity which
represses secondary infection by similar phages (23). Indeed, superinfection immunity has
previously been observed in both *Inoviridae* and *Myoviridae* lysogens (83,84). Importantly,
superinfection immunity could repress infection by lytic phages from the same family, potentially
changing bacteria-phage population dynamics and disrupting the evolution of costly phage
resistance mechanisms (8,10). In addition, it could also restrict the efficacy of phage therapies,
where lytic phages are used to treat bacterial infections (85). Therefore, prophage-mediated
superinfection immunity should be further investigated, for example, by assessing the rates of
lysogeny and cell lysis of *Inoviridae* and *Myoviridae* lysogens following infection with different
temperate and lytic phages.

The total geographical distribution of *R. solanacearum* prophages remains unclear. This is mainly
due to lack of research into it as only one genomic study investigating *R. solanacearum* prophages
has been published before this (41). Here, we show that *R. solanacearum* prophages have broad
geographical distributions, each being found in multiple continents, with continent-specificity only
observed in two low abundance prophages. Whilst these findings contrast with analyses by
Gonçalves et al (41) which suggested that *R. solanacearum* prophages are continent-specific,
this is likely due to previous studies relying on geographically unrepresentative publicly available
genomes. For example, φRS551 and φRSA1 were previously thought to be exclusive to South
America and Korea, respectively, yet we found that they were present in all six continents, with
particularly high prevalence in Africa and Europe. Therefore, by using a more diverse,

representative *R. solanacearum* genome collection, our results suggest that prophages are more
widespread than previously thought and prevalent in all six sampled continents.

Despite being widespread, prophages generally followed continent borders reticent of the distinct
geographical distributions found between *R. solanacearum* phylotypes (44). Yet, due to an under-
representation of phylotype IIA and IIB hosts in publicly available genomes, previous analyses
were unable to assess prophage phylotype distributions. We found that prophages tended to be
phylotype-specific with the *Inoviridae* and *Myoviridae* prophages almost exclusively found in
phylotype I and the RSM-RSS intermediate φRS551 and novel prophage Unclassified A primarily
found in phylotype IIB. These findings support previous studies which have observed prophage
lineage-specificity in other, primarily human, pathogens (86–91), in addition to free living bacteria
(92,93), resulting in their investigation as potential molecular markers of bacterial genomic
diversity (94–96). Therefore, this may be a widespread phenomenon. However, some prophages
were found to be more generalist, being more evenly distributed across multiple phylotypes and
occasionally in multiple copies (Unclassified C group). These may represent highly transmissible
prophages that have been acquired independently by each phylotype. Alternatively, they may
have been acquired by an ancestral *R. solanacearum* strain prior to phylotype divergence.

Prophage dissimilarity varied between phylotypes with phylotype IIB hosts generally containing
similar prophage contents and phylotype I and IIA containing more dissimilar prophages.
Prophage dissimilarity was found to be associated with host genetic dissimilarity with genetically
similar hosts tending to harbour similar prophages. Further, the *R. solanacearum* phylogenetic
tree was congruent with a UPGMA tree constructed based on prophage dissimilarity. Combined
with prophage phylotype-specificity, this suggests that, historically, prophages and their hosts
may have maintained a stable lineage-specific association and have coevolved in tandem. The
co-occurrence of prophage and host evolution within phylotypes indicates that there may have
been ancestral transmission barriers preventing prophages from moving between phylotypes.
These may have been geographical barriers as the phylotypes are thought to have arisen from
geographical isolation (97) although prophages were found to have widespread, often
overlapping, geographical distributions. Alternatively, inter-phylotype prophage transmission may
be limited by biological transmission barriers, such as phage defence systems, which are
abundant in *R. solanacearum* (7). Furthermore, it is possible that prophages could encode specific
auxiliary genes that lead to phylotype-specific ecological differences, which reduce the likelihood
of strains coexistence and horizontal movement of prophages. These hypotheses should be
assessed experimentally in future studies.

To characterise prophage auxiliary gene content, putative prophage-encoded genes were annotated and investigated. Auxiliary genes were found to comprise approximately 4% of the prophage pangenome and were potentially involved in a variety of different functions including transcriptional regulation, DNA methylation, bacterial metabolism, and virulence. Prophage-encoded metabolic genes primarily included glycosyl hydrolases and glycosyl transferases which facilitate the degradation of carbohydrates and mediate pathogen virulence (98). In *R. solanacearum*, glycosylation of type IV pilin proteins by glycosyl transferases is required for biofilm formation and pathogenicity (99). Glycosyl transferases were exclusively found in *Inoviridae* prophages, many of which are known to affect host virulence and competitiveness (34,53,54). Although the virulence reduction by *Inoviridae* prophages is typically mediated by transcriptional regulators (34,54), these results suggest that prophage-encoded auxiliary metabolic genes should be considered when assessing prophage effects on *R. solanacearum* virulence and competitiveness.

Prophages also encoded a variety of putative virulence genes including type III effectors, transcriptional regulators, and membrane transporters. Notably, transcriptional regulators were identified in the Inoviridae prophages φRS551 and φRSM3, supporting experimental studies which have attributed reduced virulence (*i.e.*, hypovirulence) in φRS551 and φRSM3 lysogens to transcriptional repressors (34,54). Transcriptional regulators were also identified in φRS138, Unclassified J, and Unclassified A, indicating these previously uncharacterised prophages may also affect host virulence. Interestingly, despite reducing host virulence in experimental assays, φRSM3 was found to contain many virulence genes, including the type III effectors RipS5 and RipTAL. This possibly reflects a variable auxiliary gene repertoire missed during experimental studies and suggests φRSM3 may also affect evasion of plant immunity. Despite their high genetic variation, *Myoviridae* prophages all appeared to have very similar gene contents with few auxiliary genes beyond a DNA methyltransferase, possibly explaining their lack of fitness effects in experimental studies (56–58). However, φRSY1 contained two type IV secretion system proteins and a low abundance type III effector RipS4. φRSY1 lysogens exhibit higher twitching motility and aggregation than non-lysogens (57) and φRSY1 virulence assays may have been impacted by using very low virulence *R. solanacearum* strains (57). Therefore, φRSY1 prophages may affect host fitness and should be re-examined using higher virulence strains. Novel, uncharacterised prophages also contained virulence genes with Unclassified A encoding the type III effector RipS5 and a histone-like DNA binding protein. Interestingly, phylotype IIB hosts, which predominantly harbour φRS551 and Unclassified A, typically contain an inactive, pseudogenised copy of RipS5 (78). Previously RipS5 pseudogenisation has been attributed to a transposon due to the presence

of an intragenic transposase (100). However, we found that, in the phylotype IIB strain UY031,

RipS5 appears to be disrupted by an Unclassified A prophage. Notably, Unclassified A contains

a Mu-like transposase and so may have previously been mislabelled as a transposon. Therefore,

prophage fitness effects may be attributed to both encoding of auxiliary gene content and host

gene disruption.

Some prophage-encoded auxiliary genes were phage family or phage-type specific, including a

glycosyl hydrolase and replication-associated recombination protein in *Inoviridae* RSM-type

phages, and a zot-like toxin in RSS-type prophages. Moreover, *Myoviridae* prophages all encoded

three glycosyl hydrolases and a DNA methyltransferase. The ubiquity of these genes suggests

that they may be under strong selection, and their presence in prophages from different

phylotypes indicates that selection is phylotype-independent. As these genes have all been

implicated in bacterial fitness (101–103), they may have been selected for by providing general

selective advantage for their hosts. Alternatively, they may provide fitness benefits to the phages

themselves by encouraging phage replication and transmission. Glycosyl hydrolases have

previously been implicated in bacterial cell wall degradation, therefore promoting cell lysis (104),

and the zot-like toxin has N-terminus similarity to the phage assembly-associated pI protein (105).

Moreover, phage-encoded DNA methyltransferases are often used by phages to evade bacterial

restriction-modification systems (106). This exposes the limitations of using bacterial databases

to identify bacterial fitness-associated auxiliary genes in prophages and highlights that, where

possible, future studies should combine bioinformatic auxiliary gene identification with

experimental and structural analyses when assessing prophage function.

In conclusion, this study provides an insight into the global diversity and distribution of *R.*

solanacearum prophages at pangenome-level. Our results highlight that while *R. solanacearum*

prophages are highly diverse and widespread, their prevalence and distribution is proportional to

their host phylotype genetic similarity. Prophages thus make a lineage-specific contribution to *R.*

solanacearum accessory genome, potentially affecting the fitness of their hosts.

Acknowledgements

586 S.G. was funded by Microbiology Society Harry Smith Vacation Studentship. M.S. is funded by a

587 NERC iCASE PhD studentship jointly with Fera Ltd. V-P.F is funded by the Royal Society

588 (RSG\R1\180213 and CHL\R1\180031) and jointly by a grant from UKRI, Defra, and the Scottish

589 Government, under the Strategic Priorities Fund Plant Bacterial Diseases program

590 (BB/T010606/1) at the University of York.

References

1.   Parikka KJ, Romancer ML, Wauters N, Jacquet S. Deciphering the virus-to-prokaryote ratio (VPR):
insights into virus–host relationships in a variety of ecosystems. Biol Rev. 2017;92(2):1081–100.

2.   Bohannan BJM, Lenski RE. Linking genetic change to community evolution: insights from studies of
bacteria and bacteriophage. Ecol Lett. 2000;3(4):362–77.

3.   Harrison E, Brockhurst MA. Ecological and Evolutionary Benefits of Temperate Phage: What Does
or Doesn't Kill You Makes You Stronger. BioEssays. 2017;39(12):1700112.

4.   Doss J, Culbertson K, Hahn D, Camacho J, Barekzi N. A Review of Phage Therapy against Bacterial
Pathogens of Aquatic and Terrestrial Organisms. Viruses. 2017 Mar;9(3):50.

5.   van Hannen EJ, Zwart G, van Agterveld MP, Gons HJ, Ebert J, Laanbroek HJ. Changes in Bacterial
and Eukaryotic Community Structure after Mass Lysis of Filamentous Cyanobacteria Associated
with Viruses. Appl Environ Microbiol. 1999 Feb;65(2):795–801.

6.   Thingstad TF, Lignell R. Theoretical models for the control of bacterial growth rate, abundance,
diversity and carbon demand. Aquat Microb Ecol. 1997 Jul 24;13:19–27.

7.   Castillo JA, Secaira-Morocho H, Maldonado S, Sarmiento KN. Diversity and Evolutionary Dynamics
of Antiphage Defense Systems in *Ralstonia solanacearum* Species Complex. Front Microbiol. 2020
[Internet]. 2020 [cited 2021 May 12];11. Available from:
https://www.frontiersin.org/articles/10.3389/fmicb.2020.00961/full

8.   Lenski RE, Levin BR. Constraints on the Coevolution of Bacteria and Virulent Phage: A Model, Some
Experiments, and Predictions for Natural Communities. Am Nat. 1985 Apr 1;125(4):585–602.

9.   Meaden S, Paszkiewicz K, Koskella B. The cost of phage resistance in a plant pathogenic bacterium
is context-dependent. Evol Int J Org Evol. 2015 May;69(5):1321–8.

10.   Wang X, Wei Z, Yang K, Wang J, Jousset A, Xu Y, et al. Phage combination therapies for bacterial
wilt disease in tomato. Nat Biotechnol. 2019 Dec;37(12):1513–20.

11.   Koskella B, Brockhurst MA. Bacteria–phage coevolution as a driver of ecological and evolutionary
processes in microbial communities. FEMS Microbiol Rev. 2014 Sep 1;38(5):916–31.

12.   Filipiak M, Łoś JM, Łoś M. Efficiency of induction of Shiga-toxin lambdoid prophages in Escherichia
coli due to oxidative and antibiotic stress depends on the combination of prophage and the
bacterial strain. J Appl Genet. 2020;61(1):131–40.

13.   Osterhout RE, Figueroa IA, Keasling JD, Arkin AP. Global analysis of host response to induction of a
latent bacteriophage. BMC Microbiol. 2007 Aug 31;7(1):82.

14.   Juhala RJ, Ford ME, Duda RL, Youlton A, Hatfull GF, Hendrix RW. Genomic sequences of
bacteriophages HK97 and HK022: pervasive genetic mosaicism in the lambdoid bacteriophages11.
Edited by M. Gottesman. J Mol Biol. 2000 May 26;299(1):27–51.

15. Billard-Pomares T, Fouteau S, Jacquet ME, Roche D, Barbe V, Castellanos M, et al. Characterization
of a P1-Like Bacteriophage Carrying an SHV-2 Extended-Spectrum β-Lactamase from an Escherichia
coli Strain. Antimicrob Agents Chemother. 2014 Nov;58(11):6550–7.

16. Iannelli F, Santagati M, Santoro F, Oggioni M, Stefani S, Pozzi G. Nucleotide Sequence of
Conjugative Prophage Φ1207.3 (formerly Tn1207.3) carrying the mef(A)/msr(D) genes for efflux
resistance to macrolides in *Streptococcus pyogenes*. Front Microbiol. 2014;5:687.

17. Frazão N, Sousa A, Lässig M, Gordo I. Horizontal gene transfer overrides mutation in *Escherichia*
coli colonizing the mammalian gut. Proc Natl Acad Sci. 2019 Sep 3;116(36):17906–15.

18. Busby B, Kristensen DM, Koonin EV. Contribution of phage-derived genomic islands to the
virulence of facultative bacterial pathogens. Environ Microbiol. 2013 Feb;15(2):307–12.

19. Hacker J, Carniel E. Ecological fitness, genomic islands and bacterial pathogenicity. EMBO Rep.
2001 May 1;2(5):376–81.

20. Wagner PL, Livny J, Neely MN, Acheson DWK, Friedman DI, Waldor MK. Bacteriophage control of
Shiga toxin 1 production and release by *Escherichia coli*. Mol Microbiol. 2002;44(4):957–70.

21. Das B, Bischerour J, Barre F-X. VGJφ integration and excision mechanisms contribute to the genetic
diversity of *Vibrio cholerae* epidemic strains. Proc Natl Acad Sci. 2011 Feb 8;108(6):2516–21.

22. Feiner R, Argov T, Rabinovich L, Sigal N, Borovok I, Herskovits AA. A new perspective on lysogeny:
prophages as active regulatory switches of bacteria. Nat Rev Microbiol. 2015 Oct;13(10):641–50.

23. Bondy-Denomy J, Qian J, Westra ER, Buckling A, Guttman DS, Davidson AR, et al. Prophages
mediate defense against phage infection through diverse mechanisms. ISME J. 2016
Dec;10(12):2854–66.

24. Ramisetty BCM, Sudhakari PA. Bacterial 'Grounded' Prophages: Hotspots for Genetic Renovation
and Innovation. Front Genet. 2019;10:65.

25. Varani AM, Monteiro-Vitorello CB, Nakaya HI, Van Sluys M-A. The role of prophage in plant-
pathogenic bacteria. Annu Rev Phytopathol. 2013;51:429–51.

26. Buell CR, Joardar V, Lindeberg M, Selengut J, Paulsen IT, Gwinn ML, et al. The complete genome
sequence of the Arabidopsis and tomato pathogen *Pseudomonas syringae* pv. tomato DC3000.
Proc Natl Acad Sci. 2003 Sep 2;100(18):10181–6.

27. de Mello Varani A, Souza RC, Nakaya HI, de Lima WC, Paula de Almeida LG, Kitajima EW, et al.
Origins of the *Xylella fastidiosa* prophage-like regions and their impact in genome differentiation.
PloS One. 2008;3(12):e4059.

28. da Silva ACR, Ferro JA, Reinach FC, Farah CS, Furlan LR, Quaggio RB, et al. Comparison of the
genomes of two *Xanthomonas* pathogens with differing host specificities. Nature. 2002 May
23;417(6887):459–63.

29. Jain M, Fleites LA, Gabriel DW. Prophage-Encoded Peroxidase in 'Candidatus Liberibacter asiaticus'
Is a Secreted Effector That Suppresses Plant Defenses. Mol Plant-Microbe Interactions®. 2015 Aug
27;28(12):1330–7.

30. Jain M, Munoz-Bodnar A, Zhang S, Gabriel DW. A Secreted 'Candidatus Liberibacter asiaticus'
Peroxiredoxin Simultaneously Suppresses Both Localized and Systemic Innate Immune Responses
In Planta. Mol Plant-Microbe Interactions®. 2018 Jun 28;31(12):1312–22.

31. Bellieny-Rabelo D, Tanui CK, Miguel N, Kwenda S, Shyntum DY, Moleleki LN. Transcriptome and
Comparative Genomics Analyses Reveal New Functional Insights on Key Determinants of
Pathogenesis and Interbacterial Competition in *Pectobacterium* and *Dickeya* spp. Appl Environ
Microbiol. 2019 Jan 15;85(2).

32. Slater SC, Goldman BS, Goodner B, Setubal JC, Farrand SK, Nester EW, *et al*. Genome Sequences of
Three *Agrobacterium* Biovars Help Elucidate the Evolution of Multichromosome Genomes in
Bacteria. J Bacteriol. 2009 Apr 15;191(8):2501–11.

33. Summer EJ, Gill JJ, Upton C, Gonzalez CF, Young R. Role of phages in the pathogenesis of
Burkholderia, or 'Where are the toxin genes in *Burkholderia* phages?' Curr Opin Microbiol. 2007
Aug 1;10(4):410–7.

34. Ahmad AA, Stulberg MJ, Mershon JP, Mollov DS, Huang Q. Molecular and biological
characterization of φRs551, a filamentous bacteriophage isolated from a race 3 biovar 2 strain of
Ralstonia solanacearum. PLOS ONE. 2017 Sep 21;12(9):e0185034.

35. Czajkowski R. May the Phage be With You? Prophage-Like Elements in the Genomes of Soft Rot
Pectobacteriaceae: *Pectobacterium* spp. and *Dickeya* spp. Front Microbiol. 2019 Feb 14;10:138.

36. Roszniowski B, McClean S, Drulis-Kawa Z. *Burkholderia cenocepacia* Prophages-Prevalence,
Chromosome Location and Major Genes Involved. Viruses. 2018 May 31;10(6).

37. Kumar R, Kumar Yadav S, Swain DM, Jha G. *Burkholderia gladioli* strain NGJ1 deploys a prophage
tail-like protein for mycophagy. Microb Cell. 5(2):116–8.

38. Zoledowska S, Motyka-Pomagruk A, Sledz W, Mengoni A, Lojkowska E. High genomic variability in
the plant pathogenic bacterium *Pectobacterium parmentieri* deciphered from de novo assembled
complete genomes. BMC Genomics. 2018 Oct 16;19(1):751.

39. Alič Š, Pédron J, Dreo T, Van Gijsegem F. Genomic characterisation of the new *Dickeya
fangzhongdai* species regrouping plant pathogens and environmental isolates. BMC Genomics.
2019 Jan 11;20(1):34.

40. Mansfield J, Genin S, Magori S, Citovsky V, Sriariyanum M, Ronald P, *et al*. Top 10 plant pathogenic
bacteria in molecular plant pathology. Mol Plant Pathol. 2012;13(6):614–29.

41. Gonçalves OS, Souza F de O, Bruckner FP, Santana MF, Alfenas-Zerbini P. Widespread distribution
of prophages signaling the potential for adaptability and pathogenicity evolution of *Ralstonia
solanacearum* species complex. Genomics. 2021 May 1;113(3):992–1000.

42. Genin S. Molecular traits controlling host range and adaptation to plants in *Ralstonia*
solanacearum. New Phytol. 2010;187(4):920–8.

43. Hayward AC. Biology and Epidemiology of Bacterial Wilt Caused by *Pseudomonas Solanacearum*.
Annu Rev Phytopathol. 1991;29(1):65–87.

44. How complex is the *Ralstonia solanacearum* species complex. In: Bacterial wilt disease and the
Ralstonia solanacearum species complex [Internet]. Saint Paul: APS Press; 2005. Available from:
http://publications.cirad.fr/une_notice.php?dk=524964

45. Remenant B, Coupat-Goutaland B, Guidot A, Cellier G, Wicker E, Allen C, *et al*. Genomes of three
tomato pathogens within the *Ralstonia solanacearum* species complex reveal significant
evolutionary divergence. BMC Genomics. 2010 Jun 15;11(1):379.

46. Safni I, Cleenwerck I, De Vos P, Fegan M, Sly L, Kappler U 2014. Polyphasic taxonomic revision of
the *Ralstonia solanacearum* species complex: proposal to emend the descriptions of *Ralstonia*
solanacearum and *Ralstonia syzygii* and reclassify current *R. syzygii* strains as *Ralstonia syzygii*
subsp. syzygii subsp. nov., *R. solanacearum* phylotype IV strains as *Ralstonia syzygii* subsp.
indonesiensis subsp. nov., banana blood disease bacterium strains as *Ralstonia syzygii* subsp.
celebesensis subsp. nov. and *R. solanacearum* phylotype I and III strains as *Ralstonia*
pseudosolanacearum sp. nov. Int J Syst Evol Microbiol. 64(Pt_9):3087–103.

47. Lowe-Power TM, Hendrich CG, von Roepenack-Lahaye E, Li B, Wu D, Mitra R, et al. Metabolomics
of tomato xylem sap during bacterial wilt reveals *Ralstonia solanacearum* produces abundant
putrescine, a metabolite that accelerates wilt disease. Environ Microbiol. 2018 Apr;20(4):1330–49.

48. Williamson L, Nakaho K, Hudelson B, Allen C. *Ralstonia solanacearum* Race 3, Biovar 2 Strains
Isolated from Geranium Are Pathogenic on Potato. Plant Dis. 2002 Sep;86(9):987–91.

49. Colburn-Clifford JM, Scherf JM, Allen C. *Ralstonia solanacearum* Dps contributes to oxidative stress
tolerance and to colonization of and virulence on tomato plants. Appl Environ Microbiol. 2010
Nov;76(22):7392–9.

50. Landry D, González-Fuente M, Deslandes L, Peeters N. The large, diverse, and robust arsenal of
Ralstonia solanacearum type III effectors and their *in planta* functions. Mol Plant Pathol.
2020;21(10):1377–88.

51. Sabbagh CRR, Carrere S, Lonjon F, Vailleau F, Macho AP, Genin S, *et al*. Pangenomic type III effector
database of the plant pathogenic *Ralstonia* spp. PeerJ. 2019 Aug 6;7:e7346.

52. Tjou-Tam-Sin NNA, van de Bilt JLJ, Westenberg M, Gorkink-Smits PPMA, Landman NM, Bergsma-
Vlami M. Assessing the Pathogenic Ability of *Ralstonia pseudosolanacearum* (Ralstonia
solanacearum Phylotype I) from Ornamental *Rosa* spp. Plants. Front Plant Sci. 2017 Nov 2;8:1895.

53. Addy HS, Askora A, Kawasaki T, Fujie M, Yamada T. The filamentous phage φRSS1 enhances
virulence of phytopathogenic *Ralstonia solanacearum* on tomato. Phytopathology. 2012
Mar;102(3):244–51.

54. Addy HS, Askora A, Kawasaki T, Fujie M, Yamada T. Loss of Virulence of the Phytopathogen
Ralstonia solanacearum Through Infection by φRSM Filamentous Phages. Phytopathology. 2012
Feb 21;102(5):469–77.

55. Peyraud R, Cottret L, Marmiesse L, Gouzy J, Genin S. A Resource Allocation Trade-Off between
Virulence and Proliferation Drives Metabolic Versatility in the Plant Pathogen *Ralstonia*
solanacearum. PLOS Pathog. 2016 Oct 12;12(10):e1005939.

56. Addy HS, Ahmad AA, Huang Q. Molecular and Biological Characterization of *Ralstonia* Phage
RsoM1USA, a New Species of P2virus, Isolated in the United States. Front Microbiol. 2019 Feb
19;10:267.

57. Askora A, Kawasaki T, Fujie M, Yamada T. Lysogenic Conversion of the Phytopathogen *Ralstonia*
solanacearum by the P2virus phi RSY1. Front Microbiol [Internet]. 2017 Nov 14;8. Available from:
http://dx.doi.org/10.3389/fmicb.2017.02212

58. Fujiwara A, Kawasaki T, Usami S, Fujie M, Yamada T. Genomic Characterization of *Ralstonia*
solanacearum Phage φRSA1 and Its Related Prophage (φRSX) in Strain GMI1000. J Bacteriol. 2008
Jan 1;190(1):143–56.

59. Nakayama K, Kanaya S, Ohnishi M, Terawaki Y, Hayashi T. The complete nucleotide sequence of
φCTX, a cytotoxin-converting phage of *Pseudomonas aeruginosa*: implications for phage evolution
and horizontal gene transfer via bacteriophages. Mol Microbiol. 1999;31(2):399–419.

60. Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010; Available from:
http://www.bioinformatics.babraham.ac.uk/projects/fastqc

61. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinforma Oxf Engl. 2014 Aug 1;30(15):2114–20.

62. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies from short
and long sequencing reads. PLOS Comput Biol. 2017 Jun 8;13(6):e1005595.

63. Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, Lees JA, et al. Producing polished
prokaryotic pangenomes with the Panaroo pipeline. Genome Biol. 2020 Jul 22;21(1):180.

64. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence
alignment based on fast Fourier transform. Nucleic Acids Res. 2002 Jul 15;30(14):3059–66.

65. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015 Jan;32(1):268–74.

66. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster version of the
PHAST phage search tool. Nucleic Acids Res. 2016 Jul 8;44(W1):W16-21.

67. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, et al. Mash: fast genome
and metagenome distance estimation using MinHash. Genome Biol. 2016 Jun 20;17(1):132.

68. Katz LS, Griswold T, Morrison SS, Caravas JA, Zhang S, Bakker HC den, et al. Mashtree: a rapid
comparison of whole genome sequence files. J Open Source Softw. 2019 Dec 10;4(44):1762.

69. González-Tortuero E, Sutton TDS, Velayudhan V, Shkoporov AN, Draper LA, Stockdale SR, et al.
VIGA: a sensitive, precise and automatic de novo VIral Genome Annotator. bioRxiv. 2018 Mar
7;277509.

70. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary: rapid large-scale
prokaryote pan genome analysis. Bioinforma Oxf Engl. 2015 Nov 15;31(22):3691–3.

71. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File
Manipulation. PLOS ONE. 2016 Oct 5;11(10):e0163962.

72. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture
and applications. BMC Bioinformatics. 2009 Dec 15;10:421.

73. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-
quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011 Oct
11;7:539.

74. Hutchinson MC, Cagua EF, Balbuena JA, Stouffer DB, Poisot T. paco: implementing Procrustean
Approach to Cophylogeny in R. Methods Ecol Evol. 2017;8(8):932–40.

75. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The carbohydrate-active
enzymes database (CAZy) in 2013. Nucleic Acids Res. 2014 Jan;42(Database issue):D490-495.

76. Urban M, Cuzick A, Seager J, Wood V, Rutherford K, Venkatesh SY, et al. PHI-base: the pathogen–
host interactions database. Nucleic Acids Res. 2020 Jan 8;48(D1):D613–20.

77. Petkau A, Stuart-Edwards M, Stothard P, Van Domselaar G. Interactive microbial genome
visualization with GView. Bioinforma Oxf Engl. 2010 Dec 15;26(24):3125–6.

78. Prokchorchik M, Pandey A, Moon H, Kim W, Jeon H, Jung G, et al. Host adaptation and microbial
competition drive *Ralstonia solanacearum* phylotype I evolution in the Republic of Korea. Microb
Genom 2020 Nov;6(11).

79. Peeters N, Carrère S, Anisimova M, Plener L, Cazalé A-C, Genin S. Repertoire, unified nomenclature
and evolution of the Type III effector gene set in the *Ralstonia solanacearum* species complex.
BMC Genomics. 2013 Dec 6;14(1):859.

80. Glickman C, Kammlade SM, Hasan NA, Epperson LE, Davidson RM, Strong M. Characterization of
integrated prophages within diverse species of clinical nontuberculous mycobacteria. Virol J. 2020
Aug 17;17(1):124.

81. Almpanis A, Swain M, Gatherer D, McEwan N 2018. Correlation between bacterial G+C content,
genome size and the G+C content of associated plasmids and bacteriophages. Microb Genomics.
4(4):e000168.

82. Bobay L-M, Touchon M, Rocha EPC. Pervasive domestication of defective prophages by bacteria.
Proc Natl Acad Sci U S A. 2014 Aug 19;111(33):12127–32.

83. Bleriot I, Trastoy R, Blasco L, Fernández-Cuenca F, Ambroa A, Fernández-García L, et al. Genomic
analysis of 40 prophages located in the genomes of 16 carbapenemase-producing clinical strains of
Klebsiella pneumoniae. Microb Genomics. 6(5):e000369.

84. Roux S, Krupovic M, Daly RA, Borges AL, Nayfach S, Schulz F, et al. Cryptic inoviruses revealed as
pervasive in bacteria and archaea across Earth's biomes. Nat Microbiol. 2019 Nov;4(11):1895–906.

85. Fujiwara A, Fujisawa M, Hamasaki R, Kawasaki T, Fujie M, Yamada T. Biocontrol of *Ralstonia*
solanacearum by Treatment with Lytic Bacteriophages. Appl Environ Microbiol. 2011
Jun;77(12):4155–62.

86. Mottawea W, Duceppe M-O, Dupras AA, Usongo V, Jeukens J, Freschi L, *et al. Salmonella enterica*
Prophage Sequence Profiles Reflect Genome Diversity and Can Be Used for High Discrimination
Subtyping. Front Microbiol. 2018;9:836.

87. Brooks MR, Padilla-Vélez L, Khan TA, Qureshi AA, Pieper JB, Maddox CW, et al. Prophage-Mediated
Disruption of Genetic Competence in *Staphylococcus pseudintermedius*. mSystems. 2020 Feb
18;5(1).

88. Goerke C, Pantucek R, Holtfreter S, Schulte B, Zink M, Grumann D, *et al*. Diversity of prophages in
dominant *Staphylococcus aureus* clonal lineages. J Bacteriol. 2009 Jun;191(11):3462–8.

89. Salloum M, van der Mee-Marquet N, Valentin-Domelier A-S, Quentin R. Diversity of prophage DNA
regions of *Streptococcus agalactiae* clonal lineages from adults and neonates with invasive
infectious disease. PloS One. 2011;6(5):e20256.

90. Vu HTK, Benjakul S, Vongkamjan K. Characterization of *Listeria* prophages in lysogenic isolates
from foods and food processing environments. PLOS ONE. 2019 Apr 1;14(4):e0214641.

91. Zamudio R, Haigh RD, Ralph JD, De Ste Croix M, Tasara T, Zurfluh K, *et al*. Lineage-specific evolution
and gene flow in *Listeria monocytogenes* are independent of bacteriophages. Environ Microbiol.
2020 Dec;22(12):5058–72.

92. Forcone K, Coutinho FH, Cavalcanti GS, Silveira CB. Prophage Genomics and Ecology in the Family
Rhodobacteraceae. Microorganisms. 2021 Jun;9(6):1115.

93. Claisse O, Chaïb A, Jaomanjaka F, Philippe C, Barchi Y, Lucas PM, et al. Distribution of Prophages in
the *Oenococcus oeni* Species. Microorganisms. 2021 Apr 16;9(4).

94. Brüssow H, Canchaya C, Hardt W-D. Phages and the Evolution of Bacterial Pathogens: from
Genomic Rearrangements to Lysogenic Conversion. Microbiol Mol Biol Rev. 2004 Sep;68(3):560–
602.

95. Cooke FJ, Wain J, Fookes M, Ivens A, Thomson N, Brown DJ, et al. Prophage Sequences Defining
Hot Spots of Genome Variation in *Salmonella enterica* Serovar Typhimurium Can Be Used To
Discriminate between Field Isolates. J Clin Microbiol. 2007 Aug 1;45(8):2590–8.

96.   Feasey NA, Dougan G, Kingsley RA, Heyderman RS, Gordon MA. Invasive non-typhoidal *Salmonella*
835   disease: an emerging and neglected tropical disease in Africa. Lancet. 2012 Jun
836   30;379(9835):2489–99.

837   97.   Castillo JA, Greenberg JT. Evolutionary Dynamics of *Ralstonia solanacearum*. Appl Environ
838   Microbiol. 2007 Feb 15;73(4):1225–38.

839   98.   Li J, Wang N. The gpsX gene encoding a glycosyltransferase is important for polysaccharide
840   production and required for full virulence in *Xanthomonas citri* subsp. citri. BMC Microbiol. 2012
841   Mar 9;12(1):31.

842   99.   Elhenawy W, Scott NE, Tondo ML, Orellano EG, Foster LJ, Feldman MF. Protein O-linked
843   glycosylation in the plant pathogen *Ralstonia solanacearum*. Glycobiology. 2016 Mar;26(3):301–
844   11.

845   100.   Castillo JA, Agathos SN. Candidate genes under balancing selection in a plant bacterial pathogen.
846   bioRxiv. 2018 Aug 9;388207.

847   101.   Stephenson SA-M, Brown PD. Epigenetic Influence of Dam Methylation on Gene Expression and
848   Attachment in Uropathogenic *Escherichia coli*. Front Public Health. 2016 Jun 27;4:131.

849   102.   Romero H, Serrano E, Hernández-Tamayo R, Carrasco B, Cárdenas PP, Ayora S, *et al. Bacillus
850   subtilis* RarA Acts as a Positive RecA Accessory Protein. Front Microbiol. 2020;11:92.

851   103.   Pérez-Reytor D, Pavón A, Lopez-Joven C, Ramírez-Araya S, Peña-Varas C, Plaza N, *et al*. Analysis of
852   the Zonula occludens Toxin Found in the Genome of the Chilean Non-toxigenic *Vibrio
853   parahaemolyticus* Strain PMC53.7. Front Cell Infect Microbiol. 2020 Sep 24;10:482.

854   104.   Vermassen A, Leroy S, Talon R, Provot C, Popowska M, Desvaux M. Cell Wall Hydrolases in
855   Bacteria: Insight on the Diversity of Cell Wall Amidases, Glycosidases and Peptidases Toward
856   Peptidoglycan. Front Microbiol. 2019 Feb 28;10:331.

857   105.   Hay ID, Lithgow T. Filamentous phages: masters of a microbial sharing economy. EMBO Rep. 2019
858   Jun;20(6).

859   106.   Kossykh VG, Schlagman SL, Hattman S. Comparative studies of the phage T2 and T4 DNA (N6-
860   adenine) methyltransferases: amino acid changes that affect catalytic activity. J Bacteriol. 1997
861   May 1;179(10):3239–43.

Supplementary figures

Figure S1. Phylogeny of *Ralstonia solanacearum* species complex. Maximum Likelihood phylogeny was constructed based on the genomes of 192 *Ralstonia solanacearum* species complex strains from Protect and the National Collection of Plant Pathogenic Bacteria (NCPPB) and other reference strains maintained at Fera Science Ltd, along with 5 previously phylotyped and sequenced strains from NCBI Genbank (names shown at the tips of tree). Phylogenetic relationships between known phylotypes was used to assign the 192 strains sequenced in this study to given phylotype clusters

Figure S2. Prophage number per genome is similar between phylotypes for intact but not incomplete prophages.

Boxplot and violin plot of the number of prophages per genome for isolates from each phylotype. Intact prophages (PHASTER score > 90) are shown in red and incomplete prophage (combined questionable and incomplete, PHASTER score < 90) are shown in blue. Box width varies with number of isolates.

Figure S3. Known *R. solanacearum* phages cluster with prophages from the same family.

Prophage neighbour-joining tree based on Mash distances. Numbers at nodes are from 1000 bootstrap replicates. Red labels are known *R. solanacearum* phages downloaded from NCBI Virus RefSeq database.

Figure S4. *R. solanacearum* **phylotypes have different prophages contents.**

Plots showing the percent of isolates in each phylotype that contain each prophage. Phylotypes analysed include phylotype I (**a**), IIA (**b**), and IIB (**c**).

Figure S5. *R. solanacearum* **phylotype IIB isolates have lower prophage dissimilarity and host genetic diversity than phylotypes I and IIA.**

Boxplots of (**a**) Average prophage dissimilarity of each phylotype, measured using average pairwise prophage Bray-Curtis distances, and (**b**) Average *R. solanacearum* genetic diversity of phylotype I, IIA, and IIB isolates, measured using average pairwise Mash distances. Box width varies with sample size.

Figure S6. **RipS5 type III effector in phylotype IIB hosts may be disrupted by the novel prophage Unclassified A.**

Circular genome visualisation of *R. solanacearum* UY031 chromosome (NCBI accession: NZ_CP012687). Inner purple ring shows GC skew and black ring shows GC content. Orange ring shows bacterial coding sequences with type III effectors highlighted in red. Outer rings show the positions of Unclassified A and RS551 prophages in the chromosome.