Immune, endothelial and neuronal network map in human lymph node and spleen

Seth Currlin[1,2], Harry S. Nick[1], Marda Jorgensen[3], Jerelyn A. Nick[3], Maigan A. Brusko[3], Hunter

Hakimian[2], Jesus Penaloza-Aponte[3], Natalie Rodriguez[3], Miguel Medina-Serpas[3], Mingder

Yang[3], Robert P. Seifert[3], Irina Kusmartseva[3], Todd M. Brusko[3,4], Kevin Otto[2], Amanda L.

Posgai[3], Clive H. Wasserfall[3], Mark A. Atkinson[3,4]*

[1]Department of Neuroscience, College of Medicine, University of Florida, 1149 Newell Drive,

Room L1-100, Gainesville, FL 32601.

[2]J. Crayton Pruitt Family Department of Biomedical Engineering, Herbert Wertheim College of

Engineering, University of Florida, 1275 Center Drive, Biomedical Sciences Building, 3[rd] Floor,

Gainesville, FL 32611.

[3]Department of Pathology, Immunology, and Laboratory Medicine, Diabetes Institute, College of

Medicine, University of Florida, 1275 Center Drive, Biomedical Sciences Building, 5[th] Floor,

Gainesville, FL 32611.

[4]Department of Pediatrics, College of Medicine, University of Florida, 1600 SW Archer Road,

Gainesville, FL 32610.

Words: 2,730

Display items: 5 Figures

References: 78

Methods References: 3

Extended Data: 11 Figures, 1 Table

Supplementary Information: 1 Table, 6 Videos

Summary

The spleen and lymph node represent important hubs for both innate and adaptive immunity[1,2].

Herein, we map immune, endothelial, and neuronal cell networks within these tissues from

"normal"/non-diseased organ donors, collected through the NIH Human BioMolecular Atlas

Program (HuBMAP)[3], using highly multiplexed CODEX (CO-Detection by indEXing) imaging

and 3D light sheet microscopy of cleared tissues. Building on prior reports[4-6], we observed the

lymph node subcapsular sinus expressing podoplanin, smooth muscle actin, and LYVE1. In the

spleen, LYVE1 was expressed by littoral cells lining venous sinusoids, whereas podoplanin was

restricted to arteries and trabeculae. 3D visualization of perivascular innervation revealed a

subset of axonal processes expressing choline acetyl transferase in both tissues, in contrast

with prior literature on human spleen[7]. We further report our novel observations regarding the

distinct localization of GAP43 and β3-tubulin within the vascular anatomy of both lymph node

and spleen, with Coronin-1A+ cells forming a dense cluster around β3-tubulin positive GAP43

low/negative segments of large vessels in spleen. These data provide an unprecedented 2D

and 3D visualization of cellular networks within secondary lymphoid tissues, laying the

groundwork for future disease-specific and system-wide studies of neural regulation of immunity

in human lymphatics.

Main

With similarities to the circulation, the lymphatic system stretches across the entire human body[8,9] to provide the critical functions of conducting lymphatic fluid, filtering blood, mounting defenses against infections and cancer, and shaping immune cell responses to protect against pathogens and autoimmune diseases[8,9]. Tissue organization and function are directly linked in spleen (SPL; blood filtration) and lymph nodes (LN; interstitial fluid filtration) facilitating immune cell priming and activation within follicles as well as afferent and efferent migration through vessels, ducts, and sinusoids[9]. The autonomic nervous system provides control of host immune defense mechanisms through innervation from sympathetic postganglionic neurons, specifically the splenic nerve for the SPL and anatomically paired nerve fibers for respective LNs[10]. However, much of our knowledge regarding the architecture and function of the lymphatic system derives from rodents[11], with recent studies noting a series of key architectural differences compared to humans[12-14].

Indeed, recent 3D imaging and topological mapping of the murine LN[15,16] have described a mesh-like network composed of fibroblastic reticular cells (FRCs) that provide the homing zone for T cells and may mediate B cell homeostasis[17]. Moreover, Huang and colleagues recently demonstrated the murine LN capsule to be encased by a network of sensory neurons capable of neuroimmune transcriptional regulation[18]. The delivery of antigens[19], inflammatory mediators, and even larger molecules[20] to the LN T cell zone involves the transport of lymph initially filtered through the subcapsular sinus, followed by passage along an elaborate lymphatic conduit system[21]. Similarly, Ogembo et al.[13] and Buckley et al.[22] have elegantly described SIRPα, FHOD1, HLA-DR, CD36 (platelet glycoprotein IV), CD71 (transferrin receptor) and CD8α as cell markers expressed by littoral cells (LCs) lining the splenic sinusoids. LCs exist exclusively in human SPL and are not observed in rodents or even non-human primates[13,22]. However, human tissue studies have historically been limited to two-dimensional (2D) tissue

cross-sections stained for low-parameter combinations of markers, highlighting the need to

characterize the human SPL and LN using a combination of highly multiplexed 2D and three-

dimensional (3D) modalities.

Multiplexed 2D-imaging of immune, lymphatic and blood vessel markers in human SPL

and LN

To extensively profile the cellular architecture of immune cells along with lymphatic and blood

endothelial cells (LECs and BECs, respectively) within the human SPL and LN, antibodies were

first validated by immunohistochemistry (IHC), demonstrating the expected organ-specific

staining patterns within formalin fixed paraffin embedded (FFPE) tissue sections (Extended

Data Fig. 1). SPL and LN tissue sections were then stained with a 29-marker panel

(Supplemental Table 1) for highly multiplexed imaging using CODEX (CO-Detection by

indexing)[23,24]. Visualizing the expression patterns (Extended Data Fig. 2) for immune cell

type/subset-defining markers[25,26] [B cell (CD20, CD21), follicular dendritic cell (FDC; CD21,

CD35), T cell (CD3ε, CD4, CD8α, CD45RO), neutrophil (CD15), and macrophage (CD68,

CD163)] delineates the organ-specific structures comprised of repeating functional units (i.e.,

lymphoid follicles). When combined with the endothelial markers [LYVE1, podoplanin (PDPN),

and PROX1 for LECs[25,27]; CD31 and CD34 for BECs[28,29]], one is able to appreciate the intricate

cellular architecture of "normal"/non-diseased LN and SPL (Extended Data Fig. 3-5).

Contrary to observations from single cell transcriptomic profiling of murine LN[4,5], our

results demonstrate that LYVE1 and PDPN expression do not overlap in the human LN or SPL

(Fig. 1). To substantiate these observations, a pair-wise pixel-based Pearson's correlation

analysis was utilized to generate averaged R values, providing a measure of signal correlation

in LN and SPL (Extended Data Fig. 5e-f). Indeed, the LYVE1/PDPN marker pair had little to no

overlap (LN, R = 0.028; SPL R = 0.0), in contrast with marker pairs such as CD4/CD3 (LN, R =

0.69; SPL, R = 0.48) and CD21/CD35 (LN, R = 0.57; SPL, R = 0.61) which were moderately

correlated, as expected[30].

PDPN and smooth muscle actin (SMA) appear to define the LN subcapsular sinus[31]

(SCS), as depicted in Fig. 1a with annotations (inset) of the LN including lymphoid follicles and

lymphatic endothelium[6,31]. Indeed, our results illustrate the expected expression patterns for

LYVE1[32,33], which identifies the complex network of LN LECs[15,16] (Fig.1a,b,c,d,e). Consistent

with previous studies[34,35], PDPN+ cells are observed within LN follicles (Fig. 1a,e,f; PDPN/CD21

R = 0.21, PDPN/CD35 R = 0.24, Extended Data Fig. 5e). Within SPL, we are able to visualize

follicles, periarteriolar lymphoid sheath (PALS), trabeculae, and associated trabecular arteries

positioned amongst the intricate network of venous sinuses, as shown in Fig. 1h-i and

annotated within the inset. Interestingly, LYVE1 labels CD8α+ LCs lining the venous

sinusoids[13,36] (Fig. 1j,k). In contrast with LN, PDPN expression is absent in SPL follicles (Fig

1h,l; PDPN/CD21 R = 0.02, PDPN/CD35 R = 0.08, Extended Data Fig. 5f). Finally, PDPN is

observed surrounding large arteries in both organs (Fig. 1a,h), and specifically in splenic

trabeculae (Fig. 1i,m).

Vimentin is observed throughout the LN sinuses and endothelium labeled by CD34,

SMA, and CD31 (Fig. 1c,f,g). In the SPL, vimentin partially overlaps with LYVE1 (R = 0.44),

most notably in splenic sinusoids identifying CD8α+ LCs[13] (Fig. 1j). The CD8$\alpha\beta$ heterodimer is a

canonical T cell marker[37] and as expected, CD8α labels individual T cells scattered across the

LN (Fig. 1c,d,e and Extended Data Fig. 2b). However, within the SPL, CD8α also defines the

LCs of the splenic sinuses, in line with previous reports[13], overlapping with vimentin and LYVE1

(Fig. 1j,k,l). CD107a (lysosome-associated membrane glycoprotein 1 [LAMP1]), which marks

degranulation on Natural Killer (NK) cells and CD8+ T cells[38] along with leukocyte adhesion to

vascular endothelium[39], and the monocyte/M2 macrophage marker CD163 are observed

throughout the LN sinus network (Fig. 1d,g) while in SPL, both markers localized to the cords of

Billroth between venous sinuses within the red pulp (Fig. 1k,n). To corroborate the

representative images in Figure 1, we provide additional CODEX datasets both from

independent donors and within multiple common coordinate framework (CCF)[40,41]-defined tissue

regions for both LN and SPL (Extended Data Fig. 3-5), and complete CODEX datasets are

publicly available on the HuBMAP data portal (Extended Data Table 1).

Cell clustering and quantitation in 2D

For a subset of donors, we utilized a 16 marker spatial X-shift clustering[42] to segregate and

classify various cell populations in both LN and SPL (Fig. 2a). We then annotated cell clusters

based on canonical marker expression and tissue localization. As anticipated[43], T cells account

for approximately one-half of all cells in LN; this, in contrast with approximately 10% of all cells

in SPL. The majority of T cells in either tissue are CD8$^+$ (Fig. 2b), as expected[44-46]. The CD15$^+$

neutrophil population provides a further distinction between the two organs comprising

approximately 1% of cells in LN versus 30% in SPL, which is in line with the literature[47-49]. A

color coordinated Voronoi tessellation image provides a visual illustration of annotated cell

populations organized within and around a follicle in each tissue (Fig. 2c). Specifically, B cells

and FDCs localized within the follicle, surrounded by marginal zone B cells, which were more

pronounced in LN than in SPL, in agreement with prior reports[50]. Follicles were further

surrounded by a distributed mixture of the other cell clusters.

137   We next performed a nonlinear high dimensionality reduction using t-Distributed

Stochastic Neighbor Embedding (t-SNE)[51] to project all clustered populations defined in Fig. 2a

into low-dimensional space (Figs. 2d). Foremost, these renderings highlight the dramatic

differences in T cell and neutrophil population frequencies within the two organs, which is further

illustrated by the gated graphs displaying individual total T cell, B cell, and

neutrophil/macrophage populations in the LN and SPL (Fig. 2e). Fig. 2e (far right) also

demonstrates the proximity of the macrophage population in the cords of Billroth (Fig.1k,n) with

the LCs lining the splenic sinuses. Cellular clustering, Voronoi images and global as well as

gated t-SNE images for four additional LN and SPL are provided in Extended Data Figures 6

and 7, respectively.

3D-imaging of perivascular innervation in SPL and LN

To visualize vascular innervation of the human LN and SPL in 3D, we employed multiplexed,

three-color light sheet fluorescence microscopy (LSFM)[52,53] on tissues cleared using CLARITY-

based methodology[54,55]. An example of a pre- and post-cleared intact human LN is shown in

Extended Data Figure 8a. Extensive 3D perivascular innervation is observed on CD31+

endothelial cells within vessels of human LN (Fig. 3a-d) and SPL (Fig. 3e-h). CD31 specificity is

visualized at the individual endothelial cell level in the SPL (Extended Data Fig. 8b,c). Staining

with the pan-neuronal marker PGP9.5 reveals a complex matrix of neural processes

surrounding CD31+ vasculature of the LN (Fig. 3b-d) and SPL (Fig. 3f-h) located within the *nervi*

vasorum of the *tunica externa* layer[56] of the vessel wall (Extended Data Fig. 8d). *Nervi vasorum*

staining is also observed in SPL for the peripheral sympathetic neuronal marker tyrosine

hydroxylase (TH) (Extended Data Fig. 8e, Supplemental Video 1). Interestingly, choline

acetyltransferase (ChAT), an enzyme required for production of the neurotransmitter

acetylcholine within parasympathetic neurons[57], colocalizes with a subset of the PGP9.5+

perivascular axons in the LN (Fig. 3a-d) and SPL (Fig. 3e-h). This is in contrast with the

previous literature[58], including recent IHC data from human cadaveric specimens where

PGP9.5[+] staining was documented in the absence of ChAT[7] as well as 3D imaging of cleared

SPL from mice where ChAT[+] nerve fibers were not detected[59]. However, our intensity-based

volumetric segmentation demonstrates that 21% of the PGP9.5+ neuronal processes also

contain ChAT in the LN evaluated, and similarly, 22% contain ChAT in SPL (Fig. 3c,g).

Renderings of sequential optical slices through the Z dimension provide 3D visualization of

perivascular innervation within human LN (Supplemental Video 2) and SPL (Supplemental

Video 3), demonstrating that PGP9.5, ChAT, and TH primarily label neuronal processes on

large vessels in both organs.

LSFM imaging of LN further identifies perivascular innervation of large vessels

expressing the pan-neuronal marker β3-tubulin as well as an extensive mesh-like vessel

network expressing both β3-tubulin and neuronal growth-associated protein (GAP43, or

neuromodulin)[15,16,60,61] (Fig. 4a-c; Supplemental Video 4). Indeed, in comparison to 2D analysis,

LSFM affords the ability to better appreciate these complex structures in 3D. Indeed, in

Extended Data Fig. 9a, we provide a close examination of high-resolution 3D max-projection

images. We also observe β3-tubulin+ nerve bundles adjacent to blood vessels (Fig. 4b;

Supplemental Video 5), along with a single GAP43[+] axon within a vessel labeled by β3-tubulin

(Extended Data Fig. 9b).

As expected, the leukocyte marker Coronin-1A[62] is detected throughout the LN (Fig. 4a-

c; Supplemental Video 5). In contrast, Coronin-1A expression is less homogenous in SPL,

observed both on single cells and on cells clustered around large vascular structures identified

by perivascular nerves expressing β3-tubulin (Fig. 4e). Within the SPL, β3-tubulin identifies

large vascular processes, whereas GAP43 identifies both large and small vessels (Fig. 4e-i),

corresponding either to BEC or efferent LECs[63]. Interestingly, immunolabeling with GAP43

demonstrated staining localized to the internal elastic membrane or smooth muscle layer of

large blood vessels in both LN (Fig. 4d) and SPL (Fig. 4h). These GAP43 data clearly define

filamentous neuronal processes existing on the internal aspect of large vessels (Extended Data

Fig. 8d), unexpected from conventional *nervi vasorum* location of vascular innervation[7]. To our

knowledge, this localization of GAP43 expressing cells has not previously been reported in

human LN or SPL vessels.

Differential expression of perivascular GAP43 and β3-tubulin in LN and SPL

A detailed review of our 3D LSFM data reveals a novel observation involving differential

expression of β3-tubulin and GAP43 along discrete segments of large vessels in both LN and

SPL (Fig. 5a-g). Specifically, within LN, select segments of large vessels display a reduction or

absence of protein expression for either GAP43 (Fig. 5a) or β3-tubulin (Fig. 5b,c) as quantified

using an intensity-based volumetric segmentation (Extended Data Fig. 10a,b).

Similarly, GAP43 protein expression is notably diminished along segments of large

vessels within the SPL, in the context of contiguous β3-tubulin expression (Fig. 5d,e). Figure 5d

also reaffirms the observation that within the SPL, GAP43 detects neuronal elements on both

large and small vessels, while β3-tubulin is primarily found on cells along large vessels (Fig.

4e,i). Notably, diminished expression of GAP43 along segments of large vessels in the SPL

appears to associate with densely clustered populations of Coronin-1A+ cells (Fig. 5f,g,

Supplemental Video 6), as supported by quantitative volumetric segmentation (Extended Data

Fig. 10c,d). To further this observation, we utilized CODEX to examine a region of the SPL that

included a PALS with CD45RO+ T cells surrounding blood vessels (Fig. 5h and Extended Data

Fig. 11) and noted that these T cell clusters may potentially correspond to the Coronin-1A+ cells

observed in LSFM.

Discussion

Herein we provide novel 2D and 3D characterization of the immune, endothelial, and neuronal

cell networks within the normal human LN and SPL. Among key observations, LYVE1 and

PDPN do not overlap in either tissue, as supported by pairwise pixel-based quantitation.

Specifically, PDPN and SMA label the LN subcapsular sinus with LYVE1 also staining the

lymphatic endothelium within the LN interior. Importantly, studies using a PDPN+ cell sorting

strategy to isolate LEC from murine LN prior to single cell transcriptomics, may fail to observe

the LYVE1+ population of cells[4,5]. Building on a seminal studies of LCs[13] and endothelial cells[27],

we report that LYVE1 labels LCs lining the SPL sinusoids, along with vimentin and CD8a, while

PDPN is expressed in central arteries and trabeculae. LCs do not exist in rodents or non-human

primates, emphasizing the importance of performing these cell mapping studies in human

tissue[13]. The abundance of macrophages expressing CD163 (hemoglobin scavenger receptor)

within the splenic cords of Billroth is in line with the existing literature and supports this organ's

well-established function as the primary site for filtration of aged or damaged red blood cells

from circulation[13,64]. Neighborhood analysis using our CODEX data, coupled with population cell

counts, delineates the disparate composition of T cell and CD15+ neutrophil populations in the

LN versus the SPL, in line with prior reports[44-49]. In particular, we observe a much higher

proportion of CD3+CD8a+ T cells in the LN compared with SPL. In addition, we note a smaller

population of CD15+ neutrophils in the LN versus the SPL, corroborating the existing

literature[65,66].

Our LSFM data highlight the power and utility of 3D imaging to observe intact tissue

volumes, consisting of thousands of optical sections from any vantage point, providing new

insights into human tissue organization that may be masked by max projection, single-section

2D views, or 3D stitching of 2D images. This effort, which applied three-color

immunofluorescent staining to human secondary lymphoid organs, represents an advance on

prior LSFM imaging studies[52,53,67-69]. Moreover, given HuBMAP's use of a CCF[40,41], the images

presented herein can be mapped back to their physical location within the intact organ. Within

both LN and SPL, the PGP9.5, β3-tubulin, ChAT, and TH markers identify an extensive network

of perivascular neuronal processes localized to the *nervi vasorum* of the *tunica externa*, while

GAP43 appears to localize to the internal elastic membrane or smooth muscle layers of the

artery[70]. We further observe that nerves expressing GAP43 and β3-tubulin associate with an

extensive network of small vessels within the LN. Moving forward, there is a need to corroborate

our interpretation that LN LYVE1 staining, visualized in 2D via CODEX, may correspond with

this vessel network.

The leukocyte marker Coronin-1A identifies scattered single cells in LN imaged by

LSFM. In contrast, Coronin-1A+ cells are both scattered and clustered around vascular

structures in SPL. Interestingly, accumulation of leukocytes around SPL blood vessels (PALS)

appears associated with reduced expression of GAP43. In support of this observation, 2D SPL

CODEX images reveal similar clustering of CD4+ T cells in PALS. Future work will require

functional assays to determine if regions lacking GAP43 correspond to sites for neurovascular

regulation of immune cell extravasation and navigation[71,72].

The data presented herein showcase the unique staining patterns of immune,

endothelial, and neural markers within human secondary lymphoid organs. Given that most

human immune responses occur in secondary lymphatic organs and not peripheral blood[30],

there is a clear need for additional studies using human tissues to gain insights on how

neuroimmune interactions modulate cellular activity across the lymphatic system. Importantly,

the current 3D cell mapping effort, together with complementary datasets and tools generated

across the HuBMAP Consortium[41,73-77], among others (e.g., Human Cell Atlas[78]), also set the

stage for future studies to determine how these pathways may be dysregulated in various

disease settings, including autoimmunity, sepsis, cancer, and viral infection. We envision that

novel therapies targeting the neurovascular sites identified herein could potentially be

developed to specifically modulate immune function within the human SPL and LN.

Main References

Krishnamurty, A. & Turley, S. Lymph node stromal cells: cartographers of the immune system. *Nature Immunology* **21**, 369-380 (2020). https://doi.org:10.1038/s41590-020-0635-3

Bronte, V. & Pittet, M. The spleen in local and systemic regulation of immunity. *Immunity* **39**, 806-818 (2013). https://doi.org:10.1016/j.immuni.2013.10.010

HuBMAP Consortium. The human body at cellular resolution: the NIH Human Biomolecular Atlas Program. *Nature* **574**, 187-192 (2019). https://doi.org:10.1038/s41586-019-1629-x

Fujimoto, N. *et al.* Single-cell mapping reveals new markers and functions of lymphatic endothelial cells in lymph nodes. *PLoS Biology* **18**, e3000704 (2020). https://doi.org:10.1371/journal.pbio.3000704

Berendam, S. *et al.* Comparative Transcriptomic Analysis Identifies a Range of Immunologically Related Functional Elaborations of Lymph Node Associated Lymphatic and Blood Endothelial Cells. *Frontiers in Immunology* **10**, 816 (2019). https://doi.org:10.3389/fimmu.2019.00816

Louie, D. & Liao, S. Lymph Node Subcapsular Sinus Macrophages as the Frontline of Lymphatic Immune Defense. *Frontiers in Immunology* **10**, 347 (2019). https://doi.org:10.3389/fimmu.2019.00347

Verlinden, T. *et al.* Innervation of the human spleen: A complete hilum-embedding approach. *Brain, Behavior, and Immunity* **77** (2019). https://doi.org:10.1016/j.bbi.2018.12.009

Betterman, K. L. & Harvey, N. L. The lymphatic vasculature: development and role in shaping immunity. *Immunol Rev* **271**, 276-292 (2016). https://doi.org:10.1111/imr.12413

Randolph, G. J., Ivanov, S., Zinselmeyer, B. H. & Scallan, J. P. The Lymphatic System: Integral Roles in Immunity. *Annu Rev Immunol* **35**, 31-52 (2017). https://doi.org:10.1146/annurev-immunol-041015-055354

Bachmann, S. B. *et al.* A Distinct Role of the Autonomic Nervous System in Modulating the Function of Lymphatic Vessels under Physiological and Tumor-Draining Conditions. *Cell Rep* **27**, 3305-3314 e3313 (2019). https://doi.org:10.1016/j.celrep.2019.05.050

Xiang, M. *et al.* A Single-Cell Transcriptional Roadmap of the Mouse and Human Lymph Node Lymphatic Vasculature. *Front Cardiovasc Med* **7**, 52 (2020). https://doi.org:10.3389/fcvm.2020.00052

Steiniger, B., Timphus, E. M. & Barth, P. J. The splenic marginal zone in humans and rodents: an enigmatic compartment and its inhabitants. *Histochem Cell Biol* **126**, 641-648 (2006). https://doi.org:10.1007/s00418-006-0210-5

Ogembo, J. G. *et al.* SIRPalpha/CD172a and FHOD1 are unique markers of littoral cells, a recently evolved major cell population of red pulp of human spleen. *J Immunol* **188**, 4496-4505 (2012). https://doi.org:10.4049/jimmunol.1103086

Cleypool, C. *et al.* Age-Related Variation in Sympathetic Nerve Distribution in the Human Spleen. *Frontiers in Neuroscience* **15**, 726825 (2021). https://doi.org:10.3389/fnins.2021.726825

Kelch, I. D. *et al.* High-resolution 3D imaging and topological mapping of the lymph node conduit system. *PLoS Biol* **17**, e3000486 (2019). https://doi.org:10.1371/journal.pbio.3000486

Novkovic, M., Onder, L., Bocharov, G. & Ludewig, B. Topological Structure and Robustness of the Lymph Node Conduit System. *Cell Rep* **30**, 893-904 e896 (2020). https://doi.org:10.1016/j.celrep.2019.12.070

17 Fletcher, A., Acton, S. & Knoblich, K. Lymph node fibroblastic reticular cells in health and
disease. *Nature Reviews Immunology* **15**, 350-361 (2015).
https://doi.org:10.1038/nri3846
18 Huang, S. *et al.* Lymph nodes are innervated by a unique population of sensory neurons
with immunomodulatory potential. *Cell* **184**, 441-459.e425 (2021).
https://doi.org:10.1016/j.cell.2020.11.028
19 Sixt, M. *et al.* The conduit system transports soluble antigens from the afferent lymph to
resident dendritic cells in the T cell area of the lymph node. *Immunity* **22**, 19-29 (2005).
https://doi.org:10.1016/j.immuni.2004.11.013
20 Reynoso, G. V. *et al.* Lymph node conduits transport virions for rapid T cell activation.
Nat Immunol **20**, 602-612 (2019). https://doi.org:10.1038/s41590-019-0342-0
21 Roozendaal, R., Mebius, R. & Kraal, G. The conduit system of the lymph node.
International Immunology **20**, 1483-1487 (2008). https://doi.org:10.1093/intimm/dxn110
22 Buckley, P. J., Dickson, S. A. & Walker, W. S. Human splenic sinusoidal lining cells
express antigens associated with monocytes, macrophages, endothelial cells, and T
lymphocytes. *J Immunol* **134**, 2310-2315 (1985).
23 Goltsev, Y. *et al.* Deep Profiling of Mouse Splenic Architecture with CODEX Multiplexed
Imaging. *Cell* **174**, 968-981 e915 (2018). https://doi.org:10.1016/j.cell.2018.07.010
24 Schurch, C. M. *et al.* Coordinated Cellular Neighborhoods Orchestrate Antitumoral
Immunity at the Colorectal Cancer Invasive Front. *Cell* **182**, 1341-1359 e1319 (2020).
https://doi.org:10.1016/j.cell.2020.07.005
25 Kong, L. L. *et al.* The optimum marker for the detection of lymphatic vessels. *Mol Clin
Oncol* **7**, 515-520 (2017). https://doi.org:10.3892/mco.2017.1356
26 Ribatti, D., Tamma, R., Ruggieri, S., Annese, T. & Crivellato, E. Surface markers: An
identity card of endothelial cells. *Microcirculation* **27**, e12587 (2020).
https://doi.org:10.1111/micc.12587
27 Breiteneder-Geleff, S. *et al.* Angiosarcomas express mixed endothelial phenotypes of
blood and lymphatic capillaries: podoplanin as a specific marker for lymphatic
endothelium. *The American Journal of Pathology* **154**, 385-394 (1999).
https://doi.org:10.1016/S0002-9440(10)65285-6
28 Lertkiatmongkol, P., Liao, D., Mei, H., Hu, Y. & Newman, P. J. Endothelial functions of
platelet/endothelial cell adhesion molecule-1 (CD31). *Curr Opin Hematol* **23**, 253-259
(2016). https://doi.org:10.1097/MOH.0000000000000239
29 Fina, L. *et al.* Expression of the CD34 gene in vascular endothelial cells. *Blood* **75**, 2417-
2426 (1990).
30 Kumar, B., Connors, T. & Farber, D. Human T Cell Development, Localization, and
Function throughout Life. *Immunity* **48**, 202-213 (2018).
https://doi.org:10.1016/j.immuni.2018.01.007
31 Jalkanen, S. & Salmi, M. Lymphatic endothelial cells of the lymph node. *Nat Rev
Immunol* **20**, 566-578 (2020). https://doi.org:10.1038/s41577-020-0281-x
32 Park, S. *et al.* Mapping the distinctive populations of lymphatic endothelial cells in
different zones of human lymph nodes. *PloS one* **9**, e94781 (2014).
https://doi.org:10.1371/journal.pone.0094781
33 Jackson, D. The lymphatics revisited: new perspectives from the hyaluronan receptor
LYVE-1. *Trends in Cardiovascular Medicine* **13**, 1-7 (2003).
https://doi.org:10.1016/s1050-1738(02)00189-5
34 Marsee, D., Pinkus, G. & Hornick, J. Podoplanin (D2-40) is a highly effective marker of
follicular dendritic cells. *Applied immunohistochemistry & molecular morphology : AIMM*
17, 102-107 (2009). https://doi.org:10.1097/PAI.0b013e318183a8e2

35 Xie, Q. *et al.* Podoplanin (d2-40): a new immunohistochemical marker for reactive
follicular dendritic cells and follicular dendritic cell sarcomas. *International journal of*
clinical and experimental pathology **1**, 276-284 (2008).
36 Steiniger, B., Stachniss, V., Schwarzbach, H. & Barth, P. J. Phenotypic differences
between red pulp capillary and sinusoidal endothelia help localizing the open splenic
circulation in humans. *Histochem Cell Biol* **128**, 391-398 (2007).
https://doi.org:10.1007/s00418-007-0320-8
37 Reina-Campos, M., Scharping, N. E. & Goldrath, A. W. CD8(+) T cell metabolism in
infection and cancer. *Nat Rev Immunol* (2021). https://doi.org:10.1038/s41577-021-
00537-8
38 Aktas, E., Kucuksezer, U., Bilgic, S., Erten, G. & Deniz, G. Relationship between
CD107a expression and cytotoxic activity. *Cellular Immunology* **254**, 149-154 (2009).
https://doi.org:10.1016/j.cellimm.2008.08.007
39 Kannan, K. *et al.* Lysosome-associated membrane proteins h-LAMP1 (CD107a) and h-
LAMP2 (CD107b) are activation-dependent cell surface glycoproteins in human
peripheral blood mononuclear cells which mediate cell adhesion to vascular
endothelium. *Cellular Immunology* **171**, 10-19 (1996).
https://doi.org:10.1006/cimm.1996.0167
40 Rood, J. *et al.* Toward a Common Coordinate Framework for the Human Body. *Cell* **179**,
1455-1467 (2019). https://doi.org:10.1016/j.cell.2019.11.019
41 Weber, G., Ju, Y. & Börner, K. Considerations for Using the Vasculature as a Coordinate
System to Map All the Cells in the Human Body. *Frontiers in Cardiovascular Medicine* **7**,
29 (2020). https://doi.org:10.3389/fcvm.2020.00029
42 Samusik, N., Good, Z., Spitzer, M., Davis, K. & Nolan, G. Automated mapping of
phenotype space with single-cell data. *Nature Methods* **13**, 493-496 (2016).
https://doi.org:10.1038/nmeth.3863
43 Bousso, P. T-cell activation by dendritic cells in the lymph node: lessons from the
movies. *Nature Reviews Immunology* **8**, 675-684 (2008). https://doi.org:10.1038/nri2379
44 Poppema, S., Bhan, A., Reinherz, E., McCluskey, R. & Schlossman, S. Distribution of T
cell subsets in human lymph nodes. *The Journal of Experimental Medicine* **153**, 30-41
(1981). https://doi.org:10.1084/jem.153.1.30
45 Sathaliyawala, T. *et al.* Distribution and compartmentalization of human circulating and
tissue-resident memory T cell subsets. *Immunity* **38**, 187-197 (2013).
https://doi.org:10.1016/j.immuni.2012.09.020
46 Langeveld, M., Gamadia, L. & ten Berge, I. T-lymphocyte subset distribution in human
spleen. *European Journal of Clinical Investigation* **36**, 250-256 (2006).
https://doi.org:10.1111/j.1365-2362.2006.01626.x
47 Puga, I. *et al.* B cell–helper neutrophils stimulate the diversification and production of
immunoglobulin in the marginal zone of the spleen. *Nature Immunology* **13**, 170-180
(2011). https://doi.org:doi:10.1038/ni.2194
48 Meinderts, S. *et al.* Human and murine splenic neutrophils are potent phagocytes of IgG-
opsonized red blood cells. *Blood Advances* **1**, 875-886 (2017).
https://doi.org:10.1182/bloodadvances.2017004671
49 Lok, L. *et al.* Phenotypically distinct neutrophils patrol uninfected human and mouse
lymph nodes. *Proceedings of the National Academy of Sciences of the United States of*
America **116**, 19083-19089 (2019). https://doi.org:10.1073/pnas.1905054116
50 Palm, A. & Kleinau, S. Marginal zone B cells: From housekeeping function to
autoimmunity? *Journal of Autoimmunity* **119**, 102627 (2021).
https://doi.org:10.1016/j.jaut.2021.102627
51 van der Maaten, L. & Hinton, G. Visualizing Data using t-SNE. *Journal of Machine*
Learning Research **9**, 2579-2605 (2008).

52 Power, R. M. & Huisken, J. A guide to light-sheet fluorescence microscopy for multiscale
imaging. *Nat Methods* **14**, 360-373 (2017). https://doi.org:10.1038/nmeth.4224
53 Wan, Y., McDole, K. & Keller, P. J. Light-Sheet Microscopy and Its Potential for
Understanding Developmental Processes. *Annu Rev Cell Dev Biol* **35**, 655-681 (2019).
https://doi.org:10.1146/annurev-cellbio-100818-125311
54 Chung, K. *et al.* Structural and molecular interrogation of intact biological systems.
Nature **497**, 332-337 (2013). https://doi.org:10.1038/nature12107
55 Gradinaru, V., Treweek, J., Overton, K. & Deisseroth, K. Hydrogel-Tissue Chemistry:
Principles and Applications. *Annu Rev Biophys* **47**, 355-376 (2018).
https://doi.org:10.1146/annurev-biophys-070317-032905
56 *Structure and Function of Blood Vessels | Anatomy and Physiology II*,
<https://courses.lumenlearning.com/suny-ap2/chapter/structure-and-function-of-blood-
vessels/> (
57 Bellier, J. & Kimura, H. Peripheral type of choline acetyltransferase: biological and
evolutionary implications for novel mechanisms in cholinergic system. *J Chem
Neuroanat* **42**, 225-235 (2011). https://doi.org:10.1016/j.jchemneu.2011.02.005
58 Nance, D. & Sanders, V. Autonomic innervation and regulation of the immune system
(1987-2007). *Brain, Behavior, and Immunity* **21**, 736-745 (2007).
https://doi.org:10.1016/j.bbi.2007.03.008
59 Ding, X. *et al.* Panicle-Shaped Sympathetic Architecture in the Spleen Parenchyma
Modulates Antibacterial Innate Immunity. *Cell Reports* **27**, 3799-3807.e3793 (2019).
https://doi.org:10.1016/j.celrep.2019.05.082
60 Shaikh, H. *et al.* Mesenteric Lymph Node Transplantation in Mice to Study Immune
Responses of the Gastrointestinal Tract. *Frontiers in Immunology* **12**, 689896 (2021).
https://doi.org:10.3389/fimmu.2021.689896
61 Matryba, P. *et al.* Systematic Evaluation of Chemically Distinct Tissue Optical Clearing
Techniques in Murine Lymph Nodes. *Journal of Immunology* **204**, 1395-1407 (2020).
https://doi.org:10.4049/jimmunol.1900847
62 Ferrari, G., Langen, H., Naito, M. & Pieters, J. A coat protein on phagosomes involved in
the intracellular survival of mycobacteria. *Cell* **97**, 435-447 (1999).
https://doi.org:10.1016/s0092-8674(00)80754-0
63 Mebius, R. E. & Kraal, G. Structure and function of the spleen. *Nat Rev Immunol* **5**, 606-
616 (2005). https://doi.org:10.1038/nri1669
64 Nagelkerke, S. *et al.* Red pulp macrophages in the human spleen are a distinct cell
population with a unique expression of Fc-γ receptors. *Blood Advances* **2**, 941-953
(2018). https://doi.org:10.1182/bloodadvances.2017015008
65 Jakovija, A. & Chtanova, T. Neutrophil Interactions with the Lymphatic System. *Cells* **10**,
2106 (2021). https://doi.org:10.3390/cells10082106
66 Maletto, B. *et al.* Presence of neutrophil-bearing antigen in lymphoid organs of immune
mice. *Blood* **108**, 3094-3102 (2006). https://doi.org:10.1182/blood-2006-04-016659
67 Luna-Palacios, Y. *et al.* Multicolor light-sheet microscopy for a large field of view
imaging: A comparative study between Bessel and Gaussian light-sheets configurations.
Journal of Biophotonics **15**, e202100359 (2022). https://doi.org:10.1002/jbio.202100359
68 Schoppmeyer, R., Zhao, R., Hoth, M. & Qu, B. Light-sheet Microscopy for Three-
dimensional Visualization of Human Immune Cells. *Journal of Visualized Experiments*,
57651 (2018). https://doi.org:10.3791/57651
69 Dekkers, J. F. *et al.* High-resolution 3D imaging of fixed and cleared organoids. *Nature
Protocols* **14**, 1756-1771 (2019). https://doi.org:doi:10.1038/s41596-019-0160-8
70 Yang, H., Wang, L., Huang, C. & Ju, G. Plasticity of GAP-43 innervation of the spleen
during immune response in the mouse. Evidence for axonal sprouting and redistribution

of the nerve fibers. *Neuroimmunomodulation* **5**, 53-60 (1998).
https://doi.org:10.1159/000026326
71   Stark, K. *et al.* Capillary and arteriolar pericytes attract innate leukocytes exiting through
venules and 'instruct' them with pattern-recognition and motility programs. *Nature*
Immunology **14**, 41-51 (2013). https://doi.org:10.1038/ni.2477
72   Iadecola, C. The Neurovascular Unit Coming of Age: A Journey through Neurovascular
Coupling in Health and Disease. *Neuron* **96**, 17-42 (2017).
https://doi.org:10.1016/j.neuron.2017.07.030
73   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587.e3529 (2021). https://doi.org:10.1016/j.cell.2021.04.048
74   Missarova, A. *et al.* geneBasis: an iterative approach for unsupervised selection of
targeted gene panels from scRNA-seq. *Genome Biol* **22**, 333 (2021).
https://doi.org:10.1101/2021.08.10.455720
75   Bueckle, A., Buehling, K., Shih, P. & Börner, K. 3D virtual reality vs. 2D desktop
registration user interface comparison. *PLoS One* **16**, e0258103 (2021).
https://doi.org:10.1371/journal.pone.0258103
76   Hasanaj, E., Wang, J., Sarathi, A., Ding, J. & Bar-Joseph, Z. Interactive single-cell data
analysis using Cellar. *Nature Communications* **13**, 1998 (2022).
https://doi.org:10.1038/s41467-022-29744-0
77   Börner, K. *et al.* Anatomical structures, cell types and biomarkers of the Human
Reference Atlas. *Nature Cell Biology* **23**, 1117-1128 (2021).
https://doi.org:doi:10.1038/s41556-021-00788-6
78   Regev, A. *et al.* The Human Cell Atlas. *eLife* **6**, e27041 (2017).
https://doi.org:10.7554/eLife.27041
486

Figures

Figure 1. Immune, LEC, BEC, and stromal cell organization in human LN and SPL.

Representative CODEX images of **a-g**, LN and **i-n**, SPL. **a**, LN cross-section (scale bar 690µm)

with annotated inset (scale bar 180µm) showing capsule, subcapsular sinus (SCS), follicles (Fo;

CD21+PDPN+), and lymphatic endothelium within the sinus (LYVE1+). The LN showed normal

microarchitecture with respect to sinus distribution and follicular meshwork formation. **b**,

CD31+CD34+SMA+ high endothelial venule (HEV, white arrow) within the LYVE1+ sinus along

with PDPN+ veins (white arrowheads). **c**, CD4+ (left) and CD8a+ (right) interfollicular T cell

zone with Vimentin+ stroma. **d**, CD107a cytotoxic lymphocytes and CD163 macrophages within

the interfollicular space. **e,** CD20+ B cells enrich the PDPN+ follicular dendritic cell (FDC)

meshwork within a Fo with CD8a identifying the T cells within the interfollicular space along with

CD163+ macrophages. **f**, CD20+ B cells enrich the PDPN+ Fo with SMA highlighting proximal

arteries and CD34 highlighting venules. Vimentin illustrates background stromal architecture. **g**,

CD20 identifies a Fo with CD163 highlighting interfollicular macrophages and with SMA,

Vimentin, and CD31 highlighting stroma and endothelium. **h**, SPL cross-section (scale bar

690µm) with annotated inset (scale bar 180µm) showing CD21+PDPN- Fo, LYVE1+ venous

sinusoids (Sinus), with CD31+CD34+SMA+PDPN+ highlighting the arterioles and periarteriolar

lymphoid sheaths (PALS). **i**, CD34+PDPN+SMA+ trabecula surrounded by LYVE1+ venous

sinusoids. **j**, CD4+ T cells (left) within the cords of Billroth; CD8a+ LCs with LYVE1+ lymphatic

endothelium and LYVE1-CD8a+ T cells within the cords of Billroth (right). **k**, Cords of Billroth

with CD107a+, CD8a+/- cytotoxic lymphocytes and CD163+ macrophages arranged between

the CD8a+ LCs and LYVE1+ sinusoidal lymphatic endothelium. **l**, CD20+PDPN- Fo surrounded

by CD8a+ LC and LYVE1+ sinusoidal lymphatic endothelium with CD163+ macrophages in the

cords of Billroth. **m**, SMA+CD34+ trabecula with Vimentin+ highlighting background stroma and

only rare CD20+ B cells and absence of PDPN+ FDCs. **n**, White pulp with tangentially sectioned

CD20+ Fo and adjacent SMA+ PALS. Vimentin highlights stroma and CD163+ macrophages in

cords of Billroth. **b-g and i-n**, Scale bars, 180μm. Donor ID, tissue, and region defined to the

right of each panel.

Figure 2. Cell clustering, segmentation, and t-SNE visualization. Analysis of CODEX data

from LN (*left*, n=5) and SPL (*right*, n=5) with identical color coding for each cluster throughout

and Littoral cells included as an additional cluster for SPL. **a,** Supervised X-shift clustering with

heatmap illustrating protein markers that define each cluster and cell counts for each cluster

with donor IDs in upper left corner. **b,** Cell counts for each supervised cluster are plotted as the

percent of the total cell population for each donor image. **c,** Clustered Voronoi plot of follicles

prominently highlighting the three supervised B cell populations in the LN and SPL. **d,** Global

high dimensionality reduction of clustered data set using t-SNE (t-Distributed Stochastic

Neighbor Embedding). **e** Three gated t-SNE plots for T (upper), B (middle), as well as

neutrophil, macrophage, and/or Littoral cell (lower) populations. Analyses performed with Akoya

CODEXMav plugin to NIH Fiji using default algorithm parameters.

Figure 3. Perivascular innervation of human LN and SPL imaged with 3D LSFM. Multicolor

LSFM imaging of CD31+ vessels with ChAT and PGP9.5 innervation in **a-d**, LN and **e-h**, SPL.

a, e, 3D renderings of 0.5 mm^3 (LN) and 1 mm^3 (SPL) tissue volumes. **b, f**, Max intensity

projection of vessels expressing CD31 surrounded by extensive PGP9.5-positive innervation

with partial ChAT-positive neuronal processes in LN (100 μm) and SPL (50 μm). **c, g**, 3D

renderings of perivascular neuronal network illustrating a subset of ChAT+ axons with the

percent of total PGP9.5+ axons expressing ChAT determined using semi-automated, intensity-

based volumetric segmentation with the Arivis software package. **d**, **h**, Cross-section through

LN (50 μm) and SPL vessels (10 μm) demonstrating co-localization of ChAT and PGP9.5 neural

components within the *tunica externa*. Donor ID, tissue, and region are defined at the top.

Figure 4. Neural markers along vessels of human LN and SPL. Multicolor LSFM imaging of β3-tubulin and GAP43 innervation with Coronin-1A+ leukocytes in **a-d**, LN and **e-i**, SPL. **a,** 3D rendering of 0.5 mm^3 LN tissue. **b**, Max intensity projection (*left*) of LN large vessel and nerve bundles expressing β3-tubulin woven between an extensive β3-tubulin+GAP43+ small vessel network (100 µm) with inset showing zoomed in cross-section (*right*) with blood vessel diameter annotated (60 µm). **c**, Wide-field max intensity projection showing Coronin-1A+ leukocytes among a complex network of β3-tubulin+GAP43+ small diameter vessels (100 µm). **d,** LN vessel cross-section demonstrating localization of β3-tubulin to the *tunica externa* and GAP43 to the *tunica interna* of the vascular wall (50 µm). **e**, 3D rendering of 1 mm^3 SPL tissue. **f, g,** Max intensity projection of SPL vessels expressing β3-tubulin within the wall and GAP43 within the internal lining (20 µm). **h,** Max intensity projection showing β3-tubulin+GAP43+ large and GAP43+ small vessels (100 µm). **i,** SPL vessel cross-section demonstrating localization of β3-tubulin to the *tunica externa* and GAP43 to the *tunica interna* of the vascular wall (20 µm). Donor ID, tissue, and region are defined at the top.

Figure 5. Differential expression of β3-tubulin and GAP43 along discrete segments of large blood vessels in human LN and SPL. Multicolor LSFM imaging of β3-tubulin and GAP43 innervation in LN (**a-c**) and SPL with Coronin-1A+ leukocytes (**d-g**). **a**, Max intensity projection showing large β3-tubulin+ vascular segment with a region of decreased GAP43 expression (yellow bracket; 100 µm). **b**, 3D rendering (40 µm) and **c,** longitudinal cross-section of LN vessel (20 µm) demonstrating absence of β3-tubulin (dashed circle) where GAP43 expression is present. **d, e**, Max projections (50 µm) show a β3-tubulin+GAP43- vessel (**d**) and a β3-tubulin-GAP43+ vessel (**e**) indicated with yellow brackets in SPL. **f, g,** Max intensity projection of SPL vessels showing regions of continuous β3-tubulin, reduced GAP43, and clustered Coronin-1A expression (dashed outlines; 50 µm). **h,** Multiplexed CODEX image of two

565   CD20+ splenic follicles with a CD31+ central artery passing through a PALS region

566   demonstrating a clustering of CD4+CD45RO+ memory T cells around the artery, in contrast to a

567   perifollicular distribution of CD8a+ T cells (180 µm). Donor ID, tissue, and region are defined to

568   the right.

Methods

Donor Acceptance Criteria

SPL and LN were acquired from 14 non-diseased organ donors (Extended Data Table 1) through the Human BioMolecular Atlas Program (HuBMAP)[3] Lymphatic System Tissue Mapping Center (TMC) at the University of Florida, via a nationwide organ procurement network. Prior to organ recovery, written informed consent for research is provided by the donor families. Criteria for donor acceptance by the HuBMAP Lymphatic System TMC at the University of Florida are available on the HuBMAP site (https://www.protocols.io/view/donor-acceptance-criteria-for-tmc-florida-zurich-h-bipykdpw) and as follows: donors less than 7 days on a ventilator, tissue received within 24 hours of recovery, and donor body mass index (BMI) between the 5[th] and 95[th] percentile as defined by the United States Centers for Disease Control and Prevention (CDC). Contraindications for acceptance include: more than four blood transfusions, autoimmune disease, chromosomal abnormalities, positive culture for bacterial or viral meningitis, or methicillin-resistant *Staphylococcus aureus* (MRSA), sepsis, observable infarction, rupture, SLP or LN deflation or gross abnormality, immune deficiency disease, cancer, infection including severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2), and splenomegaly.

SPL and LN Case Processing and Quality Control (QC)

Human SPL and mesenteric LN were processed within 16 hours of cross clamp. Residual fat or connective tissues were removed from each LN and size (mm) recorded. LN > 1 cm^2 were bisected to 1 cm^2. LN < 1 cm^2 were left intact. Intact spleens were photo documented with hilum to left, inferior end at top. Cross-sectional slabs were cut 1 cm thick, and a 1 cm wide strip was cut through the center of each slab with the superior pole surface facing up and the hilum serving as the point of origin. 1 cm^3 blocks were dissected from each strip with all cuts photo documented for location and orientation. Tissue cassettes were placed in at least 20 volumes of 4% PFA (i.e., 20 mL 4% PFA per 1 mL tissue) for 20-24 hours, transferred to 70% ethanol, and

paraffin embedded in an automated tissue processor (Sakura VIP) within 3 days. SPL and LN

were sectioned to 5 μm and stained with hematoxylin and eosin (H&E) and evaluated by an

independent pathologist to assess organ normality.

Immunohistochemistry

IHC was performed according to HuBMAP Lymphatic System TMC standard operating

procedures (SOPs) as detailed (https://www.protocols.io/view/dab-staining-of-ffpe-slides-

bdiei4be). Formalin fixed, paraffin embedded (FFPE) human SPL and LN were sectioned to 5

μm, mounted on SuperFrost Plus (Fisher Scientific) microscope slides and air dried overnight.

Slides were deparaffinized twice in xylene and 100% ethanol for 5 minutes, peroxidase blocked

with 3% peroxide in methanol for 10 minutes followed by serial ethanol washes (95%, 75% and

50%) each for 5 minutes. Finally, slides were washed in deionized water twice for 5 minutes.

Antigen retrieval involved microwaving at 50% power for 7 minutes in 10 mM sodium citrate

buffer pH 6.0 followed by an 18 minutes incubation in the hot citrate. Slides were then

rehydrated with 50mM Tris buffered saline pH 7.6 with 0.1% Tween (TBST) for 5 minutes in a

humidified chamber. Diaminobenzidine (DAB) staining was performed using VECTASTAIN Elite

ABC Kit (peroxidase-HRP) kit (Vector Labs) and IMPACT DAB (peroxidase substrate). Avidin

and biotin were sequentially blocked using Avidin/Biotin Blocking Kit (Vector Labs) for 20

minutes, followed by TBST rinse and incubation in 2% normal serum block (in TBST) for 1 hour.

Primary antibodies were diluted in Antibody Diluent Reagent Solution (Life Technologies;

Supplemental Table 1), applied to the slide, covered with a plastic coverslip and incubated

overnight at 4°C in humidified chamber. Slides were incubated with diluted biotinylated

secondary antibodies (5 μl antibody, 15 μl serum and 1 mL TBST) for 30 minutes at room

temperature followed by addition of ABC Reagent for 30 minutes. Diluted DAB reagent was then

added, development observed microscopically and reaction stopped in tap water. Slides were

then counterstained with hematoxylin followed by alcohol and xylene dehydration and cover-

slipped.

CODEX Antibody Conjugation and Validation

Commercially available oligonucleotide conjugated antibodies were purchased from Akoya

Biosciences and used at a standard dilution of 1/200 (Supplemental Table 1). Akoya CODEX

antibody validation consists of comparing its oligo-conjugated antibodies with dye-conjugated

antibodies to assure equivalent staining pattern, antibody titration for optimal signal, and single

and multiplex staining for consistent staining patterns (https://www.akoyabio.com/wp-

content/uploads/2022/01/Phenocycler_Technical-Note_Validation-of-Commercial_DN-

00140.pdf). For custom conjugation, purified, carrier-free, antibodies are purchased from

reputable vendors, providing validation data including IHC-Paraffin and western blots. All

antibodies, including commercially available CODEX antibodies are validated by IHC in our lab

using the tissue of interest. Barcodes are assigned to antibodies based on the abundance and

intensity of the protein expression. Less abundant antigens are assigned to fluorescence

channels having lower natural autofluorescence, namely Cy5 and AF750. Conversely,

antibodies for highly expressed antigens can be assigned to ATTO550 where tissue

autofluorescence is higher. Conjugations were performed with commercially available CODEX

Conjugation Kits (Akoya Biosciences) according the manufacturer's recommendation with the

following modifications. Briefly, for each antibody, a 50 kDa molecular weight cut-off (MWCO)

filter (Amicon Ultra) was washed with Filter Blocking Solution, centrifuged at 12,000 xg for 2

minutes and emptied of all liquid. To concentrate the antibody, 50 µg of protein, as determined

by Nanodrop absorbance, was diluted in 100 µl PBS, added to the filter and centrifuged at

12,000 g for 8 minutes. To initiate the disulfide reduction reaction, Antibody Reduction Master

Mix was added to the filter, vortexed for 2-3 seconds, incubated at room temperature for no

more than 25 minutes followed by buffer exchange. A unique barcode was prepared for each

antibody (Supplemental Table 1) by reconstituting in 10 µl distilled water, adding 210 µl of the

Conjugation Solution with incubation at room temperature for 2 hours. 5-10 µl of antibody

solution was reserved for protein electrophoresis, and the filter washed three times with

Purification Solution. Newly conjugated antibodies were collected by centrifugation. Custom

conjugated antibodies were titrated (1/200, 1/100, 1/50) and validated by comparing the staining

patterns of each conjugated antibody alone in CODEX against IHC DAB staining before being

included in multiplex experiments. Newly conjugated antibodies were also stained alongside

positive   and   negative   controls   in   the   CODEX   platform   as   described

(https://www.protocols.io/view/hubmap-tmc-uf-validation-of-custom-conjugated-anti-bkpzkvp6).

All antibodies, including commercially available CODEX antibodies are validated by IHC in our

lab using the tissue of interest. Barcodes are assigned to antibodies based on the abundance

and intensity of the protein expression. Less abundant antigens are assigned to fluorescence

channels having lower natural autofluorescence, namely Cy5 and AF750. Conversely,

antibodies for highly expressed antigens can be assigned to ATTO550 where tissue

autofluorescence is higher.

CODEX Staining

Oligonucleotide barcoded antibody staining of tissue sections mounted on cover slips (n=1

section per tissue region per donor) was performed using a commercially available CODEX

Staining Kit according to the manufacturer's instructions for FFPE tissue (Akoya Biosciences)

and   as   recorded   for   HuBMAP   Lymphatic   System   TMC   SOPs

(https://www.protocols.io/view/codex-antibody-staining-protocol-for-ffpe-tissues-bbsdina6).   In

brief, sample coverslips were heated to 55°C, cooled, deparaffinized and rehydrated. Antigen

retrieval was performed using 1X Citrate pH 6.0 in a pressure cooker. Tissue coverslips were

washed and equilibrated in CODEX Staining Buffer. Samples were incubated with barcoded

antibodies (diluted as described in Supplemental Table 1) in CODEX blocking buffer for 3 hours

in a humidity chamber at room temperature. After PBS washes, tissue sections were

sequentially fixed with 1.6% paraformaldehyde (PFA), methanol (on ice) and Codex Fixative

Reagent. Stained sample coverslips were stored for up to 2 weeks in CODEX Storage Buffer at

4°C. Barcoded fluorescent reporters corresponding to the barcoded primary antibodies were

added to 96-well plates containing Codex Reporter Stock solution and nuclear stain, in groups

of three (one of each wavelength) for each multiplex cycle (https://www.protocols.io/view/codex-

preparation-of-reporter-96-well-plates-bc2riyd6).

CODEX Image Acquisition

CODEX data were collected using CODEX Instrument Manager (CIM) software version 1.29,

which automates image acquisition. During cycling, the CODEX instrument automatically adds

the reporter solution to the coverslip, acquires fluorescent images, removes the reporters

(DMSO/1X CODEX Buffer wash), and prepares the tissue for the next cycle. Images were

acquired at 20X magnification using a Keyence BZ-X810 microscope with a metal halide light

source, a Plan Apo λNA 0.75 20X air objective (Nikon), and the following emission filters

(Chroma) and acquisition times: DAPI (358 nm, 10ms), TRITC (550 nm), CY5 (647 nm) and

CY7 (750 nm) Alexa Fluor 750 (500ms), Atto 550 (350ms), and Alexa Fluor 647 (500ms).

Keyence and image settings were multi-color z-stack, excitation light 100%, low photo bleach,

Shadow 0, Highlight 255, gamma 1.0. The camera was set to mono and high resolution.

Images were acquired in 13 cycles with the appropriate number of z-planes for the tissue of

interest at 1.5 pitch: spleen 17 and lymph node 11. To address the consequences of and

changes in autofluorescence during a CODEX experiment, a blank sample was imaged for each

cycle providing a direct assessment of all background signal (autofluorescence). Images were

7x9 (3.77 x 3.58 mm). Raw images were saved in 14bit TIFF (*.tif) format.

CODEX Image Processing

CODEX datasets were processed using the Akoya CODEX Processor 1.7.0.6. Multi-color z

stacks acquired during each cycle were aligned by 3D drift compensation, and the best focal

plane from each z stack was identified and used for quantitation. Specifically, there were four

fluorescent channels with channel 1 (DAPI) used as the reference in cycle 2. Tiling was snaked

by rows with 30% X and Y overlap, and 25 deconvolution iterations using the vectorial model.

The segmentation parameters included: Anisotropic Region Growth, nuclear segmentation

(DAPI), with a radius of 6, concentric circles 0, minimum/maximum cutoffs 0.02/0.99, relative

cutoff 0.01, size cutoff factor 0.01 and inner ring size of 1.0. Advanced parameters included: tile

processing (cycle alignment, background subtraction, deconvolution, extended depth of field),

region processing (shading correction, tile registration, overlap cropping), stitching, full stitching,

watershed segmentation, t-SNE report generation. Fluidic, temperature, and time dependent

effects could occur between imaging cycles and were addressed with a using 3D phase

correlation function. The calculated offsets were used in the alignment of all cycles as

determined from the comparison of the reference cycle to the nuclear DAPI stain from the

second cycle, for each cycle. Tile registration, from inaccuracies in camera positioning, were

addressed using a Microscopy Image Stitching Tool (MIST) library[79] that finds offsets by utilizing

CUDA FFT phase correlation. Given the potential for variation of background signal across

cycles, the normalized data were derived by subtracting background from each channel using

both the first and last blank cycles. To address potential effects of exposure time, the Akoya

software provides a ratio of signal exposure time to blank exposure time. For visualization

purposes, some images had brightness subject to adjustment.

Cell-Based and Neighborhood Analyses

We utilized the Akoya CODEX Multiplex Analysis Viewer (CODEX MAV; version 1.5.0.8 9)

plugin in ImageJ (version1.53f51) for image analysis. The MAV software allows for image

navigation, cell segmentation and biomarker quantification with clustering of cellular phenotypes

accomplished by an X-shift clustering algorithm[42] based on nearest neighbor density estimation.

The cell types/clusters were manually annotated based on phenotype marker expression and

displayed as an intuitive heatmap. Cell counts from each cluster from 5 LN and 5 SPL were then

displayed as the percent of the total cell population. Voronoi analyses implemented as part of

CODEX MAV were employed to illustrate separate B cell populations derived from marker-

based annotation coupled with cellular localization. Neighborhood analyses for total data sets as

well gated data subsets were displayed using dimensionality reduction and t-SNE visualization

via the CODEX MAV software plugin.

Pixel-based pair-wise colocalization analyses were performed using the Coloc2 plugin

(ImageJ/Fiji)[80] to obtain Pearson's correlation coefficients for respective protein marker pairings.

Pearson correlation analysis, with R values near 1 indicating complete colocalization and R

values near 0 indicating no colocalization of pixels above threshold intensity.

Tissue Clearing

A CLARITY[81] based protocol was used to clear tissues. All incubation and wash steps were

performed with gentle rocking. Samples (< 1 cm^3) were fixed in 40 mL of 4% PFA (Electron

Microscopy Sciences) for 24 hours at 4°C, incubated for five days at 4°C in 40 mL of hydrogel

monomer solution [4% acrylamide (Bio-Rad), 0.05% bis-acrylamide (Bio-Rad), 0.25% VA-044

Initiator (Wako Specialty Chemicals), 4% PFA, in PBS], then polymerized for 5 hours in 50 mL

PBS at 37°C. Hydrogel embedded tissues were then cleared of lipids by incubating in 50 mL of

clearing solution (Logos Bio) at 60°C, replacing clearing solution weekly until optically

transparent. Cleared samples were washed in PBS with 0.1% Triton X-100 (PBST) for 48 hours

at 37°C, replacing PBST wash buffer every 12 hours. The HuBMAP Lymphatic System TMC

receives 1 cm^3 pieces of SPL and LNs of varying sizes, ranging from 1 mm^3 to 0.5 cm^3.

Samples are further portioned to test multiple immunolabeling combinations, generally we

image 0.25 cm^3 pieces of SPL, which is ~1% of the total spleen volume; whereas we image

~50% of each intact LN.

Immunolabeling and LSFM 3D Imaging/Analysis

All incubation and wash steps were performed with gentle rocking. Cleared samples were

immunolabeled by first incubating in 10 mL RTU Blocker and Diluent (Vector Laboratories) for

48 hours at room temperature. Primary antibodies (Supplemental Table 1) were diluted (1/200)

in RTU Blocker and Diluent for 3 mL total volume added to samples, then incubated for five

days at 37°C, then two days at 4°C. Samples were washed in 50 mL PBST at room temperature

for five days, exchanging PBST wash buffer each day. Secondary antibodies (Supplemental

Table 1) were diluted (1/200) in 5 mL RTU Blocker and Diluent, and samples were protected

from light and incubated for five days at 37°C, then two days at 4°C. Samples were washed in

50 mL PBST at room temperature for five days, exchanging PBST wash buffer daily.

Immunolabeled samples were incubated in 20%, 40%, then 63% 2,2'-thiodiethanol

(Sigma Aldrich) diluted with PBS at room temperature for two hours each to attain a refractive

index of n = 1.45. Samples were then adhered to a glass capillary using cyanoacrylate to mount

into the light sheet imaging chamber. Images were obtained with a Zeiss Z.1 Lightsheet

microscope using a 20x objective. Images were processed with Arivis Vision4D software

(v.3.0.1) and are presented as single optical sections, max intensity projections, and 3D

renderings. Quantification of 3D volumes was conducted with Arivis Vision4D in the following

manner: image volumes were cropped to isolate structural features of interest, then a

supervised intensity-based segmentation was performed using the 'magic-wand' feature to

isolate channel specific objects, from which volumetric output was derived.

Data availability statement

The data generated and/or analyzed during the current study are publicly available in the

HuBMAP repository. Specifically, the complete CODEX datasets (unprocessed, processed data,

along with blank and DAPI channels) with all markers are available on the HuBMAP data portal

(https://portal.hubmapconsortium.org/search?entity_type[0]=Dataset).

Methods References

779   79   Chalfoun, J. *et al.* MIST: Accurate and Scalable Microscopy Image Stitching Tool with

780   Stage Modeling and Error Minimization. *Scientific Reports* **7**, 1-10 (2017).

781   https://doi.org:doi:10.1038/s41598-017-04567-y

782   80   Rajani, S., Gell, C., Abakir, A. & Markus, R. Computational Analysis of DNA

783   Modifications in Confocal Images. *Methods in Molecular Biology* **2198**, 227-254 (2021).

784   https://doi.org:10.1007/978-1-0716-0876-0_19

785   81   Chung, K. & Deisseroth, K. CLARITY for mapping the nervous system. *Nature Methods*

786   **10**, 508-513 (2013). https://doi.org:10.1038/nmeth.2481

Acknowledgements

We thank the families of the organ donors for the gift of tissues. This work was supported by a grant from the NIH NIAID (U54AI142766) to MAA, HSN, TMB, and CHW as well as (P01AI042288) to MAA, The Leona M. and Harry B. Helmsley Charitable Trust (G-2004-03813) to MAA, and JDRF (5-SRA-2018-557-Q-R) to MAA.

Author Contributions

SC, MJ, JAN, MAB, HH, and CHW performed the experiments and/or analyzed the data. SC, MJ, MAB, JPA, NR, MMS, MY, IK contributed to donor organ acquisition and processing. SC, HSN, CHW, and MAA designed the research. HSN, KO, TMB, CHW, and MAA supervised the research. SC, HSN, and CHW integrated the data and interpreted the results. SC, HSN, CHW, ALP, and MAA wrote the manuscript. All authors discussed the results and edited the manuscript.

Competing Interest Declaration

The authors declare no competing interests.

Additional Information

Supplementary information is available for this paper.

Correspondence and requests for materials should be addressed to Mark A. Atkinson, PhD.

Extended Data Figure and Table Legends

Extended Data Table 1. Donor demographics, clinical information, and tissue sample

identification numbers for SPL and LN examined by IHC, CODEX, and LSFM.

Extended Data Figure 1. Immunohistochemical assessment of immune, nervous,

endothelial, and stromal markers in human LN and SPL. CD3ε, CD4, CD8α, CD20, CD21,

CD15, Coronin-1A, CD31, CD34, Vimentin, CD107a, CD163, PGP9.5, β3-tubulin, GAP43,

ChAT, TH, LYVE1, PDPN and PROX1 were visualized by chromogen-based IHC in human LN

(left) and SPL (right) to confirm anticipated staining pattern and determine appropriate antibody

dilutions. DAB (brown) indicates a positive signal. Red arrows identify fine neural processes of

PGP9.5 within insets of LN and SPL (100 µm).

Extended Data Figure 2. Immune cell organization within human LN and SPL. CODEX

images of **a,b,c**, LN and **d,e,f**, SPL wide-field (top left, 560 µm) along with high resolution

images (30 µm) depicting immune cell distribution throughout tissue parenchyma. **a, d**, CD20

highlights B cells in follicles supported by CD21+, CD35+ FDC meshwork. **b,e**, CD20+ B cells

are organized in follicles with admixed CD4+ and CD8+ T cells in the interfollicular space. The

periphery of select follicles demonstrates enrichment for CD4+CD45RO+ memory helper T

cells, which may represent T follicular helper cells. In the SPL, CD3-CD8α+ cells defining the

LCs line the venous sinusoids. **c,** CD15+ neutrophils and CD68+ macrophages are sparsely

distributed throughout LN cortex. **f**, In SPL, CD15+ neutrophils are highly abundant both

circulating in sinuses and concentrated around follicles throughout the red pulp while CD68+

macrophages are found primarily in the red pulp cords. Donor ID, tissue, and region are defined

to the right of each row.

Extended Data Figure 3. Representative organization of LEC, BEC, and stromal cells in human LN. Representative CODEX images from five LNs of four independent donors (120 μm). **a**, LYVE1 highlights sinus lymphatics with PROX1 labeling LEC nuclei. LYVE1+ sinusoids, CD4+ T cells admixed with CD8a+ T cells in the perisinusoidal space and Vimentin staining the background stroma (*middle*). CD31 and CD34 identify BECs with SMA accentuating arterioles (*left*). CD163+ macrophages in the perisinusoidal space, and CD15+ neutrophils can be seen emerging from the LYVE1+ sinus (*right*). **b-d**, Additional regions from three donor LNs show the distribution of LYVE1 and PDPN expression within perivascular lymphatic sinuses as identified by CD31, CD34, and SMA. **e-g**, Three Additional LN regions from two donors with LYVE1+ sinusoids, CD4+ T cells admixed with CD8a+ T cells in the perisinusoidal space and Vimentin staining the background stroma. **h-j**, Additional LN regions from two donors showing a similar pattern as described in e-g with CD8+CD107a+ T cells and CD163+ macrophages in the perisinusoidal space. Donor ID, tissue, and region are defined to the right of each panel.

Extended Data Figure 4. Representative organization of LEC, BEC, and stromal cells in human SPL. Representative CODEX images of five SPL regions from three independent donors (120 μm). **a-c**, LYVE1 highlights splenic lymphatics while SMA and PDPN highlight a trabecula. A CD31+SMA- venules are present and PDPN co-expression is also seen in other smaller SMA+ splenic arterioles. **d-f**, LYVE1+CD8a+ LCs are present with Vimentin highlighting background stroma. Note the distribution of CD4+ T cells around a PALS (unstained, dashed oval, d), early follicle (e), and within the cords of Billroth (f) versus CD8+ T cells. **g-i**, LYVE1 again highlights LCs with CD8a coexpression, and the cords of Billroth show a regular distribution of CD8a+CD107a+ T cells and CD163+ macrophages. Donor IDs and regions are defined to the right of each panel.

Extended Data Figure 5. Highly multiplexed CODEX images of immune, stromal, and
myeloid cells in human LN and SPL along with marker colocalization analysis. **a**. The LN
shows CD20+ B cells organized into follicles with perifollicular CD4+CD3e+CD45RO+ memory
T cells (likely T follicular helper) that distribute to nearby CD31+ vasculature. CD3+CD8+ T cells
show a somewhat patchy distribution closer to the vasculature and sparing the follicles. HLA-
DR+ antigen presenting cells are scattered adjacent to the follicle and vasculature with signal
from the CD35+/CD21+ FDCs appearing in the follicles. **b.** The SPL shows a CD31+ arteriole
with a white-pulp periarteriolar follicle containing CD20+ B cells and CD21+/CD35+ FDCs. The
follicle also contains HLA-DR+ antigen presenting cells, likely dendritic cells. CD8 finely
highlights the LCs near which lie CD31+ venules. CD4+CD45RO+ memory T helper cells
appear distributed throughout the follicle, rather than the follicle periphery as seen in the LN (a)
which implies a difference in follicle age between the two sections. CD3e+CD4-CD8- NK cells
are better appreciated in this SPL section than in the LN (a). **c**. LYVE1 highlights LN sinusoidal
LECs with focal PDPN expression. PDPN is also identified in the CD21+ follicle. CD31+CD34+
venules and CD31+CD34+SMA+ arterioles are noted. CD68 and CD163 highlight macrophages
mostly within the lymphatic sinus and a few CD107a+ interfollicular cytotoxic lymphocytes are
noted. Vimentin demonstrates the background stromal architecture (120 µm). **d.** SPL sinuses
are demonstrated by CD31, CD34 and LYVE1 with PDPN highlighting a component of an SMA+
arteriole. CD68+CD163+ macrophages and CD107a+ cytotoxic lymphocytes are distributed in
the cords of Billroth with a small periarteriolar follicle identified by CD21+ follicular dendritic
meshwork. Donor ID and tissue region are defined to the right of each row. **e,f**, Pearson's R
values derived from pair-wise pixel-based colocalization analyses (Coloc2, ImageJ) for multiple
protein pairs (N=15 SPL and N=14 LN samples, mean and SD shown in red).

Extended Data Figure 6. Cell clustering, segmentation, and t-SNE visualization of high-
parameter imaging data from human LN. Parallel analysis of CODEX images from LN (N=4)

with **a-d,** donor ID and tissue region defined to the right of each panel. Each panel displays

supervised X-shift clustering with identical coloring for each cluster used throughout (*upper left*),

heat maps illustrating the protein markers that define each cluster and respective cell counts

(*upper middle*), as well as wide-angle and zoomed-in Voronoi diagrams pseudo-colored by

cluster assignment (*upper right*). Corresponding t-SNE plots (*lower*) demonstrate global

clustering as well as t-SNEs gated for T cells, B cells, neutrophils/macrophages, and endothelial

cells. All analyses were performed with the Akoya CODEXMav plugin to NIH Fiji using native

algorithm parameters.

Extended Data Figure 7. Cell clustering, segmentation, and t-SNE visualization of high-

parameter imaging data from human SPL. Parallel analysis of CODEX images from SPL

(N=4) with **a-d,** donor ID and tissue region defined to the right of each panel. Each panel

displays supervised X-shift clustering with identical coloring for each cluster used throughout

(*upper left*), heat maps illustrating the protein markers that define each cluster and respective

cell counts (*upper middle*), as well as wide-angle and zoomed-in Voronoi diagrams pseudo-

colored by cluster assignment (*upper right*). Corresponding t-SNE plots (*lower*) demonstrate

global clustering as well as t-SNEs gated for T cells, B cells, neutrophils/macrophages/littoral

cells, and endothelial cells. All analyses were performed with the Akoya CODEXMav plugin to

NIH Fiji using native algorithm parameters.

Extended Data Figure 8. CD31 in SPL highlight endothelium and associated elements

entering the *tunica media*. a, Photographs of human LN prior to (left) and after (right) tissue

clearing using the CLARITY protocol. **b,** Max intensity projection (*left*; 100 μm) and single cross-

section (*right*, 50 μm) of a CD31+ blood vessel in spleen. **c**, Max intensity projection of CD31+

splenic blood vessel with DAPI nuclear stain (20 μm). **d**, Cartoon depicting neuronal elements

(tyrosine hydroxylase [TH], ChAT, β3-tubulin, and PGP9.5) observed upon the arterial

architecture localized to the *tunica externa,* while GAP43 is limited within either the *tunica media*

or *tunica intima* (created with BioRender.com; adapted from[56]). **e**, Max intensity projection (*left,*

center left, and center right, 50 µm) and cross section (*far right*, 20 µm) of autofluorescence in

splenic blood vessels ensheathed by neuronal processes expressing TH demonstrating

localization to the *tunica externa*.

Extended Data Figure 9. LN Lymphatic vessel and nerve fiber networks. a, LSFM max

intensity projections of the extensive β3-tubulin+GAP43+ vessel network (10 µm). **b**, LSFM 3D

renderings of LN stained for β3-tubulin and GAP43 (*upper*), β3-tubulin+ vessel with a single

GAP43+ nerve fiber overlay (*three lower left panels*, 150 µm), as well as a max intensity

projection (75 µm) with inset image of cross-section (50 µm) demonstrating a GAP43+ filament

nested within the β3-tubulin+ vessel (*lower right*).

Extended Data Figure 10. 3D quantification of neuronal markers in vascular segments

associated with differential expression and lymphocyte accumulation in human LN and

SPL. a, 3D rendering of blood vessel segments from human LN demonstrating differential

expression of GAP43 and β3-tubulin, with the control segment demonstrating GAP43 volume at

55% of total β3-tubulin volume, whereas a target segment demonstrates reduced GAP43

volume at 34% of total β3-tubulin volume. **b**, 3D rendering of blood vessel segment in LN found

to have no appreciable β3-tubulin among a GAP43+ vessel. **c**, 3D rendering of blood vessel

segments from human SPL demonstrating differential expression of GAP43 and β3-tubulin, with

the control segments A and C illustrating GAP43 volume at 41% of total β3-tubulin volume in

regions without appreciable Coronin-1A$^+$ cells, whereas target segment B demonstrates GAP43

volume at 3% of total β3-tubulin volume in a region displaying significant Coronin-1A$^+$

lymphocyte accumulation. **d**, An additional vascular segment showing reduced GAP43$^+$ volume

at 4% of β3-tubulin volume in a region containing Coronin-1A$^+$ cell accumulation. Donor IDs are

defined to the right of each organ dataset. * Indicates coordination with primary figures.

Extended Data Figure 11. CODEX data demonstrating lymphocyte (CD20+ B cells and

CD45RO+ memory T cells) accumulation around vascular segments in human SPL.

Representative CODEX images of CD31+PDPN+ vascular segments within six CD20+ follicles

in a single donor SPL (750 μm). Numbered insets (90 μm) correspond to follicles identified in

top left image. Similar findings were observed across donors with bottom row showing follicles

from two more donor SPLs (90 μm). Note the distribution of CD45RO+ memory T cells with

regard to the vessels and follicle. Donor IDs and tissue CCF region are defined within each

representative image.

Lymph Node

Spleen

(a) CD21 LYVE1 PDPN CD31 CD34 SMA — 20-06-n1-reg02

Capsule, SCS, Sinus, Fo

(b) LYVE1 PDPN CD31 CD34 | LYVE1 PDPN SMA — 20-06-n3-reg03
Sinus

(c) LYVE1 Vimentin CD4 | LYVE1 Vimentin CD8 — 20-08-n7-reg03

(d) LYVE1 CD107a CD8 | LYVE1 CD163 CD8 — 20-09-n1-reg03

(e) CD20 CD163 PDPN LYVE1 CD8 — 20-08-n10-reg01
Fo

(f) CD20 SMA PDPN Vimentin CD34 — 21-014-n2.5-reg01
Fo

(g) CD20 SMA CD31 Vimentin CD163 — 20-09-n2-reg02
Fo

(h) CD21 LYVE1 PDPN CD31 CD34 SMA — 20-07-cc2-b
Sinus, Fo

(i) LYVE1 PDPN CD31 CD34 | LYVE1 PDPN SMA — 20-06-cc2-c
Sinus, Trabecula

(j) LYVE1 Vimentin CD4 | LYVE1 Vimentin CD8 — 20-07-cc2-b

(k) LYVE1 CD107a CD8 | LYVE1 CD163 CD8 — 20-08-cc2-b

(l) CD20 CD163 PDPN LYVE1 CD8 — 20-08-cc3-a
Fo

(m) CD20 SMA PDPN Vimentin CD34 — 20-06-cc2-c
Trabecula

(n) CD20 SMA CD31 Vimentin CD163 — 20-08-cc2-b
Fo, PALS

Lymph Node   **20-05-n2-reg01**   **Spleen**   **19-01-cc2-a**

Lymph Node

a β3-Tubulin | GAP43 | Overlay

b β3-Tubulin | GAP43 | Overlay **c** β3-Tubulin | GAP43

d β3-Tubulin | GAP43 | Overlay **e** β3-Tubulin | GAP43 | Overlay

f β3-Tubulin | GAP43 | Coronin1a | Overlay

Spleen

g β3-Tubulin | GAP43 | Coronin1a | Overlay

h CD20 CD31 CD8a | CD20 CD31 CD4 | CD20 CD31 CD45RO | CD45RO CD4 Colocalization threshold

20-06-n9-reg01

19-04-cc2-f

20-07-cc2-b