In vitro analysis of the effects of acetaminophen, ibuprofen and aspirin on

motility of female adult *Onchocerca volvulus* worms

Kenneth Bentum Otabil [1,2], Rhoda Kwakyewaa Antwi[1], Prince Nyarko[1], Rexford Kyei[1],

Joseph Ameyaw[4], Joseph Gyekye Bamfo[5] and Henk D F H Schallig[3]

[1]Department of Basic and Applied Biology, School of Sciences, University of Energy and

Natural Resources, Sunyani, Bono Region, Ghana; [2] The NeTroDis Research Group at the

Centre for Research in Applied Biology, School of Sciences, University of Energy and

Natural Resources, Sunyani, Bono Region, Ghana;[3]Amsterdam University Medical Centres,

Academic Medical Centre, University of Amsterdam, Department of Medical Microbiology,

Experimental Parasitology Unit, Amsterdam, The Netherlands; [4] Office of the Medical

Superintendent, Happy Family Hospital, Nkoranza, Bono East Region, Ghana and [5]Tain

District Hospital, Nsawkaw, Bono East Region, Ghana

* **Author for correspondence:** Kenneth Bentum Otabil, E-mail:

k.b.otabil@amsterdamumc.nl or   Kenneth.otabil@uenr.edu.gh

Abstract

In view of the very ambitious global timelines for elimination of onchocerciasis in 2030, the

search for alternative antifilarials cannot depend on drug development from scratch, and

repurposed drugs offer cheaper and faster alternatives. Previous studies had demonstrated the

presence and potential expression of amidase, an enzyme that can be targeted by repurposed

analgesics, *in Onchocerca volvulus worms*.   The aim of this study was to determine the

effects of acetaminophen, ibuprofen and aspirin on the motility of adult *O. volvulus* worms.

In total, thirty (30) female *O. volvulus* worms were exposed to acetaminophen, ibuprofen and

aspirin in concentrations of either 5mg/ml, 2.5mg/ml, 1.25mg/ml, 0.63mg/ml and control in

duplicates. Worm motility was observed and recorded using the WormAssay software and a

darkfield imaging apparatus starting on day 2 of incubation and ending on day 8.

Acetaminophen, ibuprofen and aspirin inhibited *O. volvulus* motility by 2-fold compared to

control in the first 24 hours of drug exposure. However, an extended exposure of the worms

to these test drugs rather improved the motility of the worms. The study has demonstrated

that a 24-hour exposure of *O. volvulus* worms to the analgesic drugs studied, results in

significant inhibition in worm motility when compared with the control group, but extended

duration of exposure led to an enhancement in motility of the previously immobile worms.

This finding supports the idea that aspirin and ibuprofen may have some longevity enhancing

properties. Further research on the utility of these analgesics as possible antifilarial drugs is

thus warranted.

Key words: onchocerciasis, amidase, wormassay, analgesics, repurposed drugs,
acetaminophen, ibuprofen, aspirin, antifilarials

Introduction

Onchocerciasis (river blindness) is a neglected tropical disease (NTD), caused by the parasite *Onchocerca volvulus* and transmitted by blackflies (WHO, 2019). Currently, the disease is still a major public health concern even after years of both vector control and Mass Drug Administration (MDA) with ivermectin (Otabil *et al.*, 2019). It is a chronic cutaneous and ocular disease characterized by skin nodules, severe itching, and ocular lesions that can progress to partial or total blindness (Noormahomed, Akrami and Mascaró-lazcano, 2016). Globally, about 21 million people are infected and a further 200 million people at risk of contracting the disease according to the 2017 Global Burden of Disease Study (WHO, 2020). Sub-Saharan Africa (SSA) bears approximately 99% of the disease burden.(Gebrezgabiher *et al.*, 2019) In Ghana, recent updates demonstrated that 15 of its 16 regions (120 districts) are endemic for onchocerciasis, with more than 7 million people at risk of contracting the disease (Marfo, 2018).

The key strategy for onchocerciasis control has been for decades the use of ivermectin, an effective microfilaricidal drug with no significant macrofilaricidal effects. However, some worrying reports from studies in West Africa have provided evidence of the development of sub-optimal response of *O. volvulus* parasites to ivermectin (Awadzi *et al.*, 2004; Osei-Atweneboana *et al.*, 2012; Eisenbarth, Achukwi and Renz, 2016). Aside this challenge, as ivermectin is purely microfilaricidal and does not kill adult worms (Abegunde, Ahuja and Okafor, 2016), control programmes have had to administer the drug for many years in order to interrupt the transmission of the disease in endemic communities (Debrah *et al.*, 2015). A major breakthrough in anti-filarial therapy was the discovery of anti-*Wolbachia* antibiotics such as doxycycline, which demonstrably had adulticidal effects (Hoerauf, 2008). However, logistical challenges associated with mass administration of antibiotics prevents the deployment of these antibiotics at the community level (Hoerauf, 2008; Hugues, Nana-

Djeunga *et al.*, 2014; Debrah *et al.*, 2015). As a result of these difficulties, coupled with the

suspension of the World Health Organization (WHO) sponsored Onchocerciasis Control

Programmes in endemic communities since 2002, there seems to be a resurgence of the

disease in some tropical foci (WHO, 2015).

It is against this background that the global health community, led by the WHO through

a recently published NTD roadmap for 2021–2030 (World Health Organization, 2020) has set

elimination of transmission (EOT) targets for onchocerciasis, with 12 countries verified for

EOT by 2030. Nevertheless, in some highly endemic foci, it might be impossible to achieve

elimination of the disease with a sole reliance on ivermectin (Basáñez *et al.*, 2006).

Therefore, the search for a microfilaricide and alternative drugs against onchocerciasis

remains a top research priority (Seidu *et al.*, 2016).

The discovery and development of new drugs requires a lot of time (10–15 years),

complex and expensive processes, and the success rate is low, only 2.01% (Ochoa-Maganda

et al., 2020). This makes it less advantageous for current onchocerciasis control efforts to

depend on development of new drugs from scratch in view of the very ambitious EOT

timelines of 2030. In this regard, drug repurposing may be the best strategy to discover

effective adulticidal drugs from existing approved drugs since these have already passed

preclinical and clinical stages (DiMasi, 2001; DiMasi, Hansen and Grabowski, 2003) and this

stands out as an attractive option to find alternative antifilarial options for onchocerciasis.

Meanwhile, a major discovery was made by Cotton *et al* (Cotton *et al.*, 2016) when they

sequenced the entire genome of *O. volvulus* parasites and its *Wolbachia* endosymbiont. They

provided a list of gene families that had key functions, which could potentially be exploited

as targets for repurposed drugs against the parasite (Cotton *et al.*, 2016). The investigators

also identified enzymes (and their gene targets) that are potentially chokepoint enzymes and

may thus be essential for *O. volvulus* survival (Cotton *et al.*, 2016). One of the top 16 *O.*

volvulus gene targets identified was the OVOC4110 gene which potentially codes for

amidase, which may be critical in metabolic pathways of *O. volvulus* (Cotton *et al.*, 2016).

The authors opined that analgesics such as acetaminophen may block the action of amidase

and consequently disrupt critical metabolic pathways in the worm (Cotton *et al.*, 2016). As

yet, no study has confirmed the expression of amidase in *O. volvulus* worms or the effects of

these analgesics on adult *O. volvulus* worms. For the present study we have investigated the

effects of acetaminophen, ibuprofen and aspirin on adult *O. volvulus* worms.

Acetaminophen (paracetamol), an analgesic and antipyretic, is the most widely used

over-the-counter medication in the world (Benista and Nowak, 2014). The mechanism of

action of acetaminophen is complex and includes the effects of both the peripheral (Cyclo-

oxygenase inhibition), and central (Cyclo-oxygenase, serotonergic descending neuronal

pathway, L-arginine/Nitric Oxide pathway, cannabinoid system) antinociception processes

and Redox mechanisms (Benista and Nowak, 2014). Though, some studies have reported

some antimicrobial effects of acetaminophen (AL-Janabi, 2010), the direct activity of

acetaminophen against helminths in general and *O. volvulus* in particular has not been

demonstrated by any study till date. Cotton *et al.* (2016) however speculated that

acetaminophen might bind to amidases in *O. volvulus* and potentially cause the death of the

worms, hence its choice in the present study.

Ibuprofen is a medication in the nonsteroidal anti-inflammatory drug (NSAID) class that

is used for treating pain, fever, and inflammation (Davies, 1998). Like other NSAIDs, it

works by inhibiting the production of prostaglandins by decreasing the activity of the enzyme

cyclooxygenase (Gunaydin and Bilge, 2018) though it might be a weaker anti-inflammatory

drug than other NSAIDs (Gunaydin and Bilge, 2018). The drug was chosen for the present

study because aside being an analgesic, some antimicrobial activity has been reported by

previous authors (AL-Janabi, 2010; Zimmermann and Curtis, 2017). However, till date no

study has investigated the direct effects of ibuprofen on *O. volvulus* survival.

Aspirin (acetylsalicylic acid, ASA), an analgesic, antipyretic and anti-inflammatory drug,

has emerged as an interesting option for repurposing drugs against parasites (Ochoa-Maganda

et al., 2020). ASA may also be used for secondary prevention of stroke and acute cardiac

events. ASA acts on the two cyclooxygenase isoforms (Cyclo-oxygenase 1and 2) via

acetylation of serine 532 in the active site; inhibits the action of both enzymes and prevents

the formation of prostaglandins from arachidonic acid (Lei *et al.*, 2015). Epidemiological and

clinical studies showed that ASA also reduces the incidence of epithelial tumours by the

acetylation of multiple proteins including transcription factors, cytoskeleton proteins, stress

response proteins (including the heat shock proteins, HSPs), membrane proteins, among

others (Flossmann and Rothwell, 2007; Chan and Giovannucci, 2010), suggesting other

mechanisms and molecular targets. Some studies have evidenced antiparasitic activity. For

instance, Singh and Rathaur (Singh and Rathaur, 2010) demonstrated that ASA improved the

antifilarial activity of diethylcarbamazine (Singh and Rathaur, 2010). Aspirin also

demonstrably affects dynamics of the actin cytoskeleton and decreases amebic movement in

Entamoeba histolytica (López-Contreras *et al.*, 2013).

The aim of the study therefore was to determine the effects of acetaminophen, ibuprofen

and aspirin on the survivability of adult *O. volvulus* worms. This was important because there

is the need to quickly discover and develop effective alternatives and complement to

ivermectin if global elimination targets of onchocerciasis is to be achieved.

Materials and methods

Patient recruitment and nodulectomies

Onchocerca nodules were obtained from 11 participants who had consented to participate in

this study by signing/thumbprinting a written, informed consent form. The study was

approved by the Institutional Ethics Committee of the Kintampo Health Research Centre in

Ghana (Approval number: KHRCIEC/2018-18). The participants were recruited from seven

onchocerciasis endemic communities in the Bono Region of Ghana. Participants were

certified as eligible for nodulectomies by the Study Clinician and Principal Investigator.

Using local anaesthesia, palpable nodules were surgically removed from the patients by the

Study Clinician and the medical team. Participants were transported to their homes and

wound care provided at the nearest health care facility until the wound was healed under the

supervision of the medical team. The excised nodules were preserved in 0.85% normal saline

in 20 ml plastic vials, placed on ice, transported to the laboratory and stored at room

temperature on the laboratory bench.

Collagenase digestion of excised nodule

In order to isolate the adult worms contained in the nodules, the latter were digested using the

collagenase technique (Nana-Djeunga *et al.*, 2014). Briefly, each nodule was incubated for

12–48 hours at 37°C (time dependant on the nodule's weight) in 5ml of Gibco's high glucose

Dulbecco's Modified Eagle's Medium (DMEM, Gibco, 12100061) containing type I

collagenase (Sigma Aldrich, SCR103) at a final concentration of 2.25 mg/ml. The product of

digestion (the worm mass and digested human tissues constituting the nodule) was placed in a

Petri dish containing 15 ml of DMEM, sodium bicarbonate (Dainess, Ghana), and

supplemented with Penicillin-Streptomycin (Gibco, 15140122) at a final concentration of 2

mg/ml. Individual worms were isolated under a dissecting microscope using entomological

forceps and needles. Each entire and living worm (dead or calcified and incomplete or broken

worms were counted but discarded from the further process) was then examined for sex.

Entire male worms were individually frozen for further studies. Thirty (30) female worms

were examined for activity in order to include them in the live worm assays after 24 hours of

separation from nodules. Males were excluded from this study because initial culture of the

male worms indicated an unusually fast decay in DMEM.

Drug preparation

Stock solutions of acetaminophen, ibuprofen, aspirin were prepared by grinding 500mg of the

commercially available drugs [Paracetamol (M&G Pharmaceuticals, Ghana); Ibuprofen

(Letap Pharmaceuticals, Ghana); Aspirin (M & A Pharmaceuticals, UK)] in a crucible and

dissolved in 50ml of DMEM solution in a 250 ml beaker. Subsequently, concentrations of

5mg/ml, 2.5 mg/ml, 1.25mg/ml and 0.63mg/ml were prepared using two-fold serial dilution

following a protocol developed by AL-Janabi, 2010 .The concentration of acetaminophen,

ibuprofen and aspirin in micromolars (mM) were; Acetaminophen (5mg/ml=33.08mM, 2.5

mg/ml=16.54mM, 1.25mg/ml = 8.27mM and 0.63mg/ml = 4.17mM), Ibuprofen (5mg/ml =

24.24mM, 2.5mg/ml = 12.12mM, 1.25mg/ml = 6.06mM and 0.63mg/ml = 3.06mM), Aspirin

(5mg/ml = 27.75mM, 2.5 mg/ml = 13.88mM, 1.25mg/ml = 6.94mM and 0.63mg/ml =

3.50mM). The drug dilutions were stored at 4° C in a refrigerator for 24 hours prior to usage

in the O. volvulus assays.

O. volvulus worm culture nodule

The worms were cultured following established protocols (Townson *et al.*, no date) with a

few modifications. Briefly, the experiment was set up by adding a single female adult O.

volvulus worm to each well of 6-well plates containing 5ml of DMEM +10% fetal bovine

serum (Gibco, A3840002) + 100 u/ml penicillin + 100 ug/ml streptomycin + sodium

bicarbonate (2 g/1). The worms were incubated at 37°c in a gas phase of 5% CO_2 in air for 24

hours prior to the addition of the analgesics to adapt them to the culture medium. After 24

hours of incubation in the above medium, 1ml of each concentration of analgesics (5, 2.5,

1.25 and 0.63mg/ml) was pipetted into the respectively labelled wells containing the pre-

incubated O. volvulus worms in the medium. Two wells with DMEM but no analgesics

added were used as the control for this study. The worms were again incubated at 37°c in a

gas phase of 5% CO_2 in air for 8 days with motility reading performed daily starting after 24

hours. Close examination of worms was performed to identify signs of disintegration and the

medium changed every other day as colour changes indicated depletion of glucose in the

culture medium.

The WormAssay and the darkfield imaging apparatus

The WormAssay software previously developed by Marcellino *et al.* (2012) for antifilarial

screening of *Brugia malayi* was repurposed in the present study for *O. volvulus* (Bedi, 1996).

The WormAssay is an inexpensive system for quantifying parasite movement based on worm

motility. The apparatus uses a commodity video camera, computer and a newly developed

free and open source software application (*GitHub - chrismarcellino/wormassay: automated*

whole plate screening of macroscopic parasites, no date) to provide quantitative

measurements of parasite motility on entire plates. The application can process multiple wells

simultaneously without user interaction, and automatically identifies each well in the plate

and labels the output data accordingly. This system can be used to assay large parasites such

as filarial nematodes as well as other macroparasites. WormAssay's automation of the video

capture step and lack of need for any interaction with the computer software during scoring

differentiates it from all existing motion-based schemes. For the video capture system, a dark

field macroscopic imaging apparatus based on the design of Marcellino *et al.* (2012) was

developed with slight modifications.

Measurement of motility using the WormAssay software and darkfield imaging system

Before the addition of the test drugs, the motility of the female worms to be used were

observed by technicians and judged to be of sufficient motility to include in the assays. Using

the WormAssay and the capture device, each 6-well microplate was observed and scored for

motility. One minute video recordings using the Lucas-Kande Optical Flow algorithm were

taken of each plate and mean optical flow movement units for each worm were converted to

percent motility by normalization in Graphpad Prism. The effective concentration to reduce

mean motility levels to 50% (EC50) of control values were also determined using Graphpad

Prism. The full profile of % mean motility scores (MMS) were illustrated graphically and

statistical comparisons made.

Statistical analyses

Statistical analysis was performed using Graphpad Prism (version 8.2.1). In plotting the dose

response curves, all movement units were normalized and the EC50 (the concentration of a

drug that gives half-maximal response was calculated for acetaminophen, ibuprofen and

aspirin) were determined. For the graphical plots, the mean of the normalized movement

units was plotted against the concentration or the day depending on the type of plot.

Differences in mean motility scores between groups were calculated using the Mann-Whitney

test with differences considered to be statistically significant at the 95% confidence interval if

$P < 0.05$. The standard deviations (SD) of the mean motility scores were determined and

subsequently converted to % SD for consistency.

Results

The results of the effects of acetaminophen, ibuprofen and aspirin on the survival of adult

female *O. volvulus* worms are presented in Figures 1-5. The indicator of survival of *O.*

volvulus female worms in this study was the percentage mean motility scores (% MMS).

Effects of acetaminophen, ibuprofen and aspirin on motility of adult worms

The results of the % MMS of the *O. volvulus* after a 24-hour exposure to the various

concentrations of analgesics are presented in Figure 1. It can be seen in Figure 1 that at the

0.63 mg/ml dose of acetaminophen, the % MMS was 41% (SD=11.6%). The value of %

MMS continued to increase as the concentration of acetaminophen increased [1.25 mg/ml

(63%, SD=100%), then 2.5 mg/ml (68%, SD=18%)]. The only concentration of

acetaminophen which performed best in terms of reducing motility of the worms was the

5mg/ml where the % MMS decreased to 5% (SD=0%). The EC50 (the concentration of a

drug that gives half-maximal response) value of acetaminophen in this study was 8.74mg/ml

(57.82mM).

The results show that at the lowest concentration of ibuprofen (0.63mg/ml), the % MMS

of worms in the ibuprofen group was 0% (SD=7%) compared to 100% (SD=100%) in the

control group. For ibuprofen concentrations of 1.25mg/ml, 2.5 mg/ml and 5mg/ml, the %

MMS of the female adult worms were respectively at 20 % (SD=0%), 23 % (SD=1%) and 24

% (SD=17%). The EC50 for ibuprofen was 0.65mg/ml (3.15 mM).

The findings on the effects of aspirin on the worms showed that at the startup aspirin

concentration (0.63mg/ml), 30% (SD=61%) of the worms were markedly motile within the

aspirin group compared to 100% in the control group. The lowest % MMS for worms in the

aspirin group (0%, SD=0%) was observed at an aspirin concentration of 1.25mg/ml.

However, at the 5mg/ml concentration of aspirin, 41% (SD=55%) of the female adult worms

in the group were still motile. The EC50 of aspirin in this study was 0.26mg/ml (1.44mM).

Figure 1: Dose response curve of the effects of Acetaminophen, Ibuprofen and Aspirin
261

on female adult O. volvulus worms after 24 hours of incubation
262

Response of O. volvulus parasites to varying concentrations of the test drugs during 8 days
264

of culture
265

The following figures (2-5) presents the results of the effects of varying concentrations of the
266

test drugs (control, 5 mg, 2.5mg, 1.25mg/ml and 0.63 mg/ml) on female adult worms in
267

culture for 8 days. Figure 2 summarizes the response of the of *O. volvulus* worms to 5mg/ml
268

concentrations of acetaminophen, ibuprofen, aspirin and control. The findings demonstrate
269

that at 24 hours of culture (day 2), all test drugs including control had no effects on the
270

motility of the worms as the % MMS of female adult worms in each group was 100 %. A
271

similar observation was made on day 3 as all test drugs again had similar effects on worm
272

motility. For day 4, the study observed that acetaminophen had the lowest effect on worm
273

motility with % MMS of 21% (SD = 1.6%), with this trend continuing even to the 8[th] day of
274

culture where 55% (SD=5.7%) of worms were still highly motile despite exposure to 5mg/ml
275

of acetaminophen. Generally, the differences in % MMS between the aspirin and control was

statistically significant (P = 0.026; Mann-Whitney test). There was however no statistically

significant difference in % MMS between acetaminophen and control (P = 0.052; Mann-

Whitney test) and ibuprofen and control (P = 0.127; Mann-Whitney test).

Figure 2: Comparison of 5mg concentration of Acetaminophen, Ibuprofen, Aspirin on

the motility of *O. volvulus* over the course of 8 days of culture

A summary of the effects of the 2.5mg/ml concentration of acetaminophen, ibuprofen and

aspirin on the motility of female adult *O. volvulus* are presented in Figure 3. After 24 hours

(day 2) of exposure of female adult *O. volvulus* to the test drugs, aspirin had the highest effect

of reducing the % MMS, but even then, still had a % MMS of about 59% (SD = 2.2%). For

the remaining days of culture (except on day 7), acetaminophen and ibuprofen outperformed

aspirin. For instance, on day 3 of culture, the % MMS of the acetaminophen and ibuprofen

group were only 5 % (SD = 56%) and 11% (SD = 16%) of worms in the, respectively, were

motile compared to 30% of the worms in the aspirin group. It is interesting to note that on

this day, the control group performed better than all the test drugs in terms of reduction in %

MMS. There was a statistically significant difference between ibuprofen and control (P =

0.037; Mann-Whitney test) as well as between aspirin and control (P = 0.026; Mann-Whitney

test). However, the difference between acetaminophen and control for this concentration

group was not statistically significant (P = 0.104; Mann-Whitney test).

Figure 3: Comparison of 2.5mg concentration of Acetaminophen, Ibuprofen, Aspirin on

the motility of *O. volvulus* over the course of 8 days of culture

For the 1.25mg /ml concentrations of the test drugs, aspirin performed best on day 2 (after 24

hours of exposure to the drug) though again, it was subsequently outperformed by all other

test drugs including control (Figure 4). The only exception was on day 6, when no motility

was observed for the aspirin group (0 % MMS, SD = 0 %) and day 8, when it performed

better than ibuprofen. Surprisingly however, the control group continued to demonstrate very

low motility throughout the culture period. For this drug concentration, there were

statistically significant differences between acetaminophen and control (P = 0.026; Mann-

Whitney test), ibuprofen and control (P = 0.037; Mann-Whitney test) but not between aspirin

and control (P = 0.071; Mann-Whitney test).

Figure 4: Comparison of 1.25 mg concentration of Acetaminophen, Ibuprofen, Aspirin

on the motility of *O. volvulus* over the course of 8 days of culture

The findings on the response of *O. volvulus* adult females to 0.63mg/ml concentration of the

various test agents as well as control is presented in Figure 5. On day 2, the best performing

drug in terms of reduction in % MMS was ibuprofen, but even then, the % MMS was 70%

(SD=7%) indicating a high worm motility in the group. Generally, however, acetaminophen

performed better at this concentration when compared with aspirin and ibuprofen. Again, the

control group continued to show relatively lower motility scores when compared with the

acetaminophen, ibuprofen and aspirin. These differences in % MMS was statistically

significant for ibuprofen versus control (P = 0.026; Mann-Whitney test) but not for aspirin

versus control (P = 0.052; Mann-Whitney test) and acetaminophen versus control (P = 0.104;

Mann-Whitney test).

Figure 5: Comparison of 0.63 mg/ml concentration of Acetaminophen, Ibuprofen,

Aspirin on the motility of *O. volvulus* over the course of 8 days of culture

Discussion

In response to the need to develop alternative drugs to augment ivermectin and based on

empirical evidence from the sequencing of the genome of *O. volvulus* by Cotton *et al.*(2016),

the present study tested the hypothesis that analgesics such as acetaminophen, ibuprofen and

aspirin has adulticidal effects on adult female *O. volvulus* worms.

     For acetaminophen, this study demonstrated that a the 24-hour exposure of the worms to

5mg/ml acetaminophen completely immobilized the worms suggesting a potential antifilarial

effect. However, the summary of results of the various concentrations of acetaminophen over

the course of the 8 days incubation revealed that all concentrations from this drug improved

the motility of the female adult worms and hence did not indicate any antifilarial effect. In

fact, the 1.25 mg/ml concentration of acetaminophen significantly improved worm motility.

This finding is contradictory to the suggestion by Cotton *et al.* (2016) that analgesics,

such as acetaminophen, could have adulticidal effects on *O. volvulus* by blocking amidase, a

potentially choke point enzyme. The ability of acetaminophen to enhance parasite motility

and survival demonstrated in this study may not be surprising as a previous study on

Plasmodium falciparum also demonstrated that the drug may prolong parasite clearance and

had no direct antiparasitic effects (Brandts *et al.*, 1997). Though the foregoing study was on

malaria parasites, the results from the current study suggest a similar worm enhancing effect.

It is therefore justified that prior to our study, acetaminophen had only been utilized as an

antipyretic in the treatment of helminthic infections, with no direct effects on parasite

intended. However, as the present study did not investigate the combination of

acetaminophen and antifilarials, it is still possible that such a combination, in the right

concentration, might be deleterious to *O. volvulus* parasite survival. This may not be far-

fetched as some investigators have demonstrated the antimicrobial ability of acetaminophen

on bacteria and fungi at different concentrations (AL-Janabi, 2010; Zimmermann and Curtis,

2017). Taken together, it is posited that there is a possible inhibitory effect of acetaminophen

which is both dose and time dependent as the 24-hour exposure of worms to 5mg/ml

acetaminophen completely immobilized them, but more research is needed in this area.

This study also investigated the effects of ibuprofen on the survivability of adult female

O. volvulus worms. The findings showed that ibuprofen reduced the motility of the parasites

at different concentrations within the first 24 hours of culture and this represented a two-fold

increase in inhibitory activity over that of control. This finding is novel as this study

represents the very first report of the inhibitory effects of ibuprofen on *O. volvulus* or on

helminths in general. The antimicrobial effects of ibuprofen have been demonstrated

previously for bacteria such as *Acinetobacter baumannii, Bacillus spp.*, *Helicobactor pylori*,

Klebsiella spp, etc as well as for fungi including *Candida albicans*, *Epidermophyton*

floccosum, *Penicillium expansum* etc (Zimmermann and Curtis, 2017). A rather surprising

finding was the fact that summary of results of long-term (8 days) of exposure of worms to

ibuprofen improved the motility of the worms. The improvement in worm motility by

ibuprofen was statistically significant at all concentrations with the exception of the 5 mg/ml

concentration when compared to the control group. This finding was particularly interesting

because based on a previous study by Cotton *et al.*(2016), the current study had hypothesized

that ibuprofen, being an analgesic, binds to amidase which is probably a chokepoint enzyme,

leading to worm death. Thus, the reason for this observed significant improvement in motility

of female adult worms following a relatively long-term exposure to ibuprofen is not known

though the idea that ibuprofen may improve the longevity of microbes is not new (He *et al.*,

2014). He *et al.* (2014) discovered that the common drug ibuprofen increases the lifespan of

yeast, worms and flies (specifically *Saccharomyces cerevisiae*, *Caenorhabditis elegans* and

Drosophila melanogaster respectively), indicative of conserved longevity effects. They

opined that in budding yeast, ibuprofen's pro-longevity action is independent of its known

anti-inflammatory role (He *et al.*, 2014). They also revealed that the critical function of

ibuprofen in longevity is to inhibit the uptake of aromatic amino acids, by destabilizing the

high-affinity tryptophan permease. Nevertheless, in consideration of the differences in effects

of ibuprofen in the short-term and after extended exposure in this study, it is suggested that

effect of ibuprofen just like acetaminophen is both dose and time dependent. The findings

from the present study therefore begs for more research in this area to unravel further details

on the effects of ibuprofen on *O. volvulus* worms and its potential role either as a treatment

option or as a supplement in culturing worms in the lab.

The study further investigated the effects of aspirin on female adult *O. volvulus* worms.

The findings showed that in the short-term, aspirin also demonstrated a 2-fold inhibition of

the motility of the worms compared to the control group. This finding was not surprising as

aspirin is a known nonsteroidal anti-inflammatory drug which can inhibit prostaglandin H

synthase and also induces apoptosis (Singh and Rathaur, 2010). In fact, Singh and Rathaur

(2010) working with *Setaria cervi,* a bovine filarial parasite, demonstrated that exposure of

the worms to a combination of diethylcarbamazine plus aspirin at a concentration of 100 mM

irreversibly paralyzed adult worms as well as microfilariae within 2 hours of exposure.

Evidence from their study also showed that the worms might have been killed due to

mitochondrial mediated apoptosis (Singh and Rathaur, 2010). In another study by Ochoa-

Maganda *et al.* (2020), aspirin showed a direct antigiardial activity and affected the adhesion

and growth of trophozoites in a time-dose-dependent manner (Ochoa-Maganda *et al.*, 2020).

However, for the present study, what was surprising in the aspirin group was the observation

that over the course of the entire period of incubation, aspirin seemed to enhance the motility

of the female adult worms. The reason for this is not known though aspirin has also long been

a subject of research on pro-longevity in humans. The results from our study thus begs for

further research in this area.

As observed in this study, the general trend is that at the onset of incubation,

acetaminophen, ibuprofen and aspirin, demonstrated some inhibitory effects on female adult

worms. However, after an extended period of exposure, the drugs improved the motility of

the worms indicating that the effects of these drugs are both dose and time dependent. It is

speculated, that at the beginning of the introduction of the drugs, the analgesics might bind to

amidase (Cotton *et al.*, 2016) and inhibit the worms within 24 hours duration of exposure. It

is however possible, that amidase might not be a chokepoint enzyme. Hence, the worms

could switch to alternative metabolic pathways (Seidu *et al.*, 2016) which does not require

amidase and might thus increase their survivability *in vitro*, thereby accounting for the

observed enhancement in worm motility scores in this study.

Conclusion

The study provides, for the first time, evidence on the direct effects of acetaminophen,

ibuprofen and aspirin on the motility of female adult *O. volvulus* worms. The study has

demonstrated that a 24-hour exposure of these parasitic worms to the study analgesic drugs

results in significant inhibition in worm motility when compared with the control group.

Longer duration of exposure of the worms to the different concentrations of the study drugs

however led to a rather surprising enhancement in motility of the previously immobile

worms. There is therefore the need for more research to determine the utility of these

analgesics as possible antifilarial drugs or supplements to improve *in vitro* culturing of *O.*

volvulus worms for other studies.

Supplementary material

Not applicable

Data

The datasets used and/or analyzed during the current study are available from the

corresponding author on reasonable request.

Acknowledgements

We thank all the staff and students of the Department of Basic and Applied Biology who

provided support in one way or the other to ensure the successful implementation of this

research. We also appreciate the efforts of the directors and staff of the Wenchi Municipal

Health Directorate, Nsawkaw District and the Subinso Health Center for their enormous

contributions towards the study. Our appreciation also goes to the chiefs and people of the

study communities who partnered with us in this study.

Author Contribution

Financial Support

This research received no specific grant from any funding agency, commercial or not-for-

profit sectors.

Conflicts of Interest

The authors declare there are no conflicts of interest.

Ethical Standards

This study received the ethical approval of Institutional Ethics Committee of the Kintampo

Health Research Centre, Kintampo, Bono Region, Ghana (Approval number:

KHRCIEC/2018-18).

References

Abegunde, AT, Ahuja, RM and Okafor, NJ (2016) 'Doxycycline plus ivermectin versus ivermectin alone for treatment of patients with onchocerciasis', *Cochrane Database of Systematic Reviews*. doi: 10.1002/14651858.CD011146.pub2.

AL-Janabi, H (2010) 'In Vitro antibacterial activity of ibuprofen and acetaminophen', *Journal of Global Infectious Diseases*, 2(2), p. 105. doi: 10.4103/0974-777x.62880.

Awadzi, K *et al.* (2004) 'An investigation of persistent microfilaridermias despite multiple treatments with ivermectin, in two onchocerciasis-endemic foci in Ghana', *Annals of Tropical Medicine & Parasitology*, 98(3), pp. 231–249. doi: 10.1179/000349804225003253.

Basáñez, MG *et al.* (2006) 'River blindness: A success story under threat?', *PLoS Medicine*, pp. 1454–1460. doi: 10.1371/journal.pmed.0030371.

Bedi, GS (1996) 'Comparative study of kallikrein-like serine proteinases from rat submandibular glands', *Preparative Biochemistry and Biotechnology*, 26(2), pp. 85–104. doi: 10.1080/10826069608000056.

Benista, MJ and Nowak, JZ (2014) 'Paracetamol□: mechanism of action , applications and safety concern', *Acta Pol Pharm*. Jan-Feb 2014;71(1):11-23.

Brandts, CH *et al.* (1997) 'Effect of paracetamol on parasite clearance time in *Plasmodium falciparum* malaria', *Lancet*, 350(9079), pp. 704–709. doi: 10.1016/S0140-6736(97)02255-1.

Chan, AT and Giovannucci, EL (2010) 'Primary Prevention of Colorectal Cancer', *Gastroenterology*, 138(6), p. 2029. doi: 10.1053/j.gastro.2010.01.057.

Cotton, JA *et al.* (2016) 'The genome of *Onchocerca volvulus*, agent of river blindness', *Nature Microbiology*, 2(November 2016), pp. 1–12. doi: 10.1038/nmicrobiol.2016.216.

Davies, NM (1998) 'Clinical Pharmacokinetics of Ibuprofen The First 30 Years', Clin

Pharmacokinet. 1998 Feb;34(2):101-54.

Debrah, AY *et al* (2015) 'Doxycycline Leads to Sterility and Enhanced Killing of Female *Onchocerca volvulus* Worms in an Area with Persistent Microfilaridermia after Repeated Ivermectin Treatment: A Randomized, Placebo-Controlled, Double-Blind Trial', *Clinical Infectious Diseases*, 61(4), pp. 517–526. doi: 10.1093/cid/civ363.

DiMasi, JA (2001) 'Risks in new drug development: Approval success rates for investigational drugs', *Clin Pharmacol Ther*, pp. 297–307. doi: 10.1067/mcp.2001.115446.

DiMasi, JA, Hansen, RW and Grabowski, HG (2003) 'The price of innovation: New estimates of drug development costs', *Journal of Health Economics*, 22(2), pp. 151–185. doi: 10.1016/S0167-6296(02)00126-1.

Eisenbarth, A, Achukwi, MD and Renz, A (2016) 'Ongoing transmission of *Onchocera volvulus* after 25 years of annual mass ivermectin treatments in the Vina Du Nord River Valley, in North Cameroon', *PLoS Negl Trop Dis*, 10(2).

Flossmann, E and Rothwell, PM (2007) 'Effect of aspirin on long-term risk of colorectal cancer: consistent evidence from randomised and observational studies', *Lancet*, 369(9573), pp. 1603–1613. doi: 10.1016/S0140-6736(07)60747-8.

Gebrezgabiher, G *et al.* (2019) 'Reaching the last mile: Main challenges relating to and recommendations to accelerate onchocerciasis elimination in Africa', *Infectious Diseases of Poverty*. doi: 10.1186/s40249-019-0567-z.

GitHub - chrismarcellino/wormassay: automated whole plate screening of macroscopic parasites. Retrieved from the website: https://github.com/chrismarcellino/wormassay (Accessed: 3 February 2021).

Gunaydin, C and Bilge, SS (2018) 'Effects of nonsteroidal anti-inflammatory drugs at the molecular level', *Eurasian Journal of Medicine*, 50(2), pp. 116–121. doi:

10.5152/eurasianjmed.2018.0010.

He, C *et al.* (2014) 'Enhanced Longevity by Ibuprofen, Conserved in Multiple Species,

Occurs in Yeast through Inhibition of Tryptophan Import', *PLoS Genetics*. Edited by S.

K. Kim, 10(12), p. e1004860. doi: 10.1371/journal.pgen.1004860.

Hoerauf, A (2008) 'Filariasis: new drugs and new opportunities for lymphatic filariasis and

onchocerciasis'. *Current Opinion in Infectious Diseases*. 2008 Dec;21(6):673-81. doi:

10.1097/QCO.0b013e328315cde7.

López-Contreras, *et al* (2013) 'Exploring the possible role of lysine acetylation on

Entamoeba histolytica virulence: a focus on the dynamics of the actin cytoskeleton',

BioMed research international, 2013. doi: 10.1155/2013/757392.

Lei, J *et al* (2015) 'Mechanistic insights into a classic wonder drug-aspirin', *Journal of the*

American Chemical Society, 137(1), pp. 70–73. doi: 10.1021/ja5112964.

Marcellino, C, *et al.* (2012) 'WormAssay: A novel computer application for whole-plate

motion-based screening of macroscopic parasites', *PLoS Neglected Tropical Diseases*,

6(1). doi: 10.1371/journal.pntd.0001494.

Marfo, K (2018) *Update of Oncho Program Status*. Retrieved from: https://dndi.org/wp-

content/uploads/2018/05/Marfo_CountryProgressGhana_UgandaMeeting2018.pdf.(Acce

ssed 16 February 2021)

Nana-Djeunga, H *et al.* (2014) 'Reproductive Status of Onchocerca volvulus after Ivermectin

Treatment in an Ivermectin-Naïve and a Frequently Treated Population from Cameroon',

PLoS Neglected Tropical Diseases. Edited by A. Hoerauf, 8(4), p. e2824. doi:

10.1371/journal.pntd.0002824.

Noormahomed, EV, Akrami, K and Mascaró-lazcano, C (2016) 'Onchocerciasis , an

undiagnosed disease in Mozambique: identifying research opportunities', *Parasites &*

Vectors, pp. 1–8. doi: 10.1186/s13071-016-1468-7.

Ochoa-Maganda, VY *et al.* (2020) 'Antigiardial activity of acetylsalicylic acid is associated with overexpression of hsp70 and membrane transporters', *Pharmaceuticals*, 13(12), pp. 1–16. doi: 10.3390/ph13120440.

Osei-Atweneboana, MY *et al* (2012) 'Genotypic analysis of β-tubulin in Onchocerca volvulus from communities and individuals showing poor parasitological response to ivermectin treatment', *International Journal for Parasitology: Drugs and Drug Resistance*, 2, pp. 20–28. doi: 10.1016/j.ijpddr.2012.01.005.

Otabil, KB, *et al.* (2019) 'Prevalence of onchocerciasis and associated clinical manifestations in selected hypoendemic communities in Ghana following long-term administration of ivermectin', *BMC Infectious Diseases*, 19(1), p. 431. doi: 10.1186/s12879-019-4076-2.

Seidu, MA, *et al.* (2016) 'Carbohydrate Catabolism in Adult Onchocerca volvulus□: An Carbohydrate Catabolism in Adult *Onchocerca volvulus*□: An Immunohistochemical Study'. *American Journal of Biomedical and Life Sciences* (May). doi: 10.11648/j.ajbls.20160403.13.

Singh, A and Rathaur, S (2010) 'Combination of DEC plus aspirin induced mitochondrial mediated apoptosis in filarial parasite *Setaria cervi*', *Biochimie*, 92(7), pp. 894–900. doi: 10.1016/j.biochi.2010.03.018.

Townson, S *et al* (1990) 'The effects of ivermectin used in combination with other known antiparasitic drugs on adult Onchocerca gutturosa and *O. volvulus* in vitro.', *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 84(3), pp. 411–6. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2260177 (Accessed: 9 April 2019).

Elliott, M, Nava, S, & Schönhofer, B (Eds.). (2018). Non-Invasive Ventilation and Weaning: Principles and Practice. 2[nd] ed. CRC Press. https://doi.org/10.1201/9781315153643

WHO (2019) 'WHO | Onchocerciasis elimination mapping of endemic countries is key to defeating river blindness', *WHO*. Available at:

https://www.who.int/neglected_diseases/news/Onchocerciasis-elimination-mapping-of-

endemic-countries-is-key/en/ (Accessed: 14 April 2019).

WHO (2020) *Onchocerciasis Factsheet*. Retrieved from: https://www.who.int/news-

room/fact-sheets/detail/onchocerciasis (Accessed on 30 July 2020).

World Health Organization (2020) 'Ending the neglect to attain the Sustainable Developmetn

Goals - A road map for neglected tropical diseases 2021–2030', (February 2019), pp. 1–

13. Available at: http://apps.who.int/bookorders.

Zimmermann, P. and Curtis, N. (2017) 'Antimicrobial Effects of Antipyretics.',

Antimicrobial agents and chemotherapy, 61(4), pp. e02268-16. doi:

10.1128/AAC.02268-16.