1 **Engineering transcription factor BmoR mutants for constructing multifunctional**

2 **alcohol biosensors**

3 Tong Wu[#], Zhenya Chen[#], Shuyuan Guo, Cuiying Zhang, Yi-Xin Huo[*]

Abstract

Native transcription factor-based biosensors (TFBs) have the potential for in situ detection of value-added chemicals or byproducts. However, their industrial application is limited by their ligand promiscuity, low sensitivity, and narrow detection range. Alcohols exhibit similar structures, and no reported TFB can distinguish a specific alcohol from its analogs. Here, we engineered an alcohol-regulated transcription factor, BmoR, and obtained various mutants with remarkable properties. For example, the generated signal-molecule-specific BmoRs could distinguish the constitutional isomers n-butanol and isobutanol, with insensitivity up to an ethanol concentration of 800 mM (36.9 g/L). Linear detection of 0–60 mM of a specific higher alcohol could be achieved in the presence of up to 500 mM (23.0 g/L) ethanol as background noise. Notably, two mutants with raised outputs and over 1.0×10^7-fold higher sensitivity, and one mutant with an increased upper detection limit (14.8 g/L n-butanol or isobutanol) were screened out. Using BmoR as an example, this study systematically explored the ultimate detection limit of a TFB towards its small-molecule ligands, paving the way for in situ detection in the biofuel and wine industries.

Keywords: transcription factor, BmoR, biosensor, specificity, high sensitivity, wider detection

Introduction

Upon sensing specific signal molecules, transcription factors (TFs) can bind or unbind to the DNA-regulatory sequences of target genes to induce or repress gene transcription. TFs can be utilized as biosensors by coupling the transcriptional alteration with the expression of a reporter protein[1]. Transcription factor-based biosensors (TFB) have been designed and constructed to detect toxic metals [2], regulate metabolic flux [3, 4], and screen highly active enzymes [5] or chemical overproducers [6, 7]. However, several drawbacks limit the industrial applications of TFBs. First, TFB signaling can be saturated by the medium concentration of intracellular signal products and cannot distinguish chemical overproducers from producers [8]. Second, TFBs can be interfered by the byproducts, which are structurally similar to the products [9]. Third, in many cases, TFBs are not sensitive enough to detect low concentrations of value-added chemicals[10, 11]. Therefore, it is necessary to develop TFBs with low ligand promiscuity, high sensitivity, and a wide detection range.

n-Butanol and isobutanol are considered as promising substitutes for gasoline because of their special properties[12]. Butanols are also crucial precursors for the production of plastics and polymers[13]. Metabolic engineering has been used to produce n-butanol and isobutanol in various hosts [14-22]. However, an efficient TFB with low ligand promiscuity, high sensitivity, and a wide detection range has yet to be achieved for in situ detection

and high-throughput screening. *Pseudomonas butanovora* BmoR is an activated

transcription factor that belongs to the bacterial enhancer-binding proteins (bEBPs) and

can induce the transcription of σ^{54}-dependent P_{bmo} promoter by combining with C2-C5

n-alcohols [23, 24]. The activation mechanism of BmoR is shown in Fig. 1A. The N-terminal,

central domain, and C-terminal of BmoR are responsible for sensing and combining the

signal molecules, contacting the σ^{54} factor of holoenzyme and hydrolyzing ATP, and

binding to the upstream sequence of the σ^{54}-dependent promoter, respectively [24]. Native

BmoR has been shown to respond to C3-C5 alcohols, and a BmoR-based biosensor has

been utilized to screen n-butanol or isobutanol producers [10].

In this study, we aimed to obtain efficient BmoR mutants with specificity, high sensitivity,

and a wide detection range towards special higher alcohols for the purpose of in situ

detection and high-throughput screening. The development of a biosensor that was

sensitive to butanols but completely insensitive to 0–36 g/L ethanol was another

significant purpose of this study. This type of biosensor is urgently needed in the wine

industry for the in situ detection of byproducts (n-butanol or isobutanol) production

without interfering with ethanol production. Here, we utilized error-prone PCR to

construct a BmoR random mutagenesis library based on the understanding of the BmoR

N-terminal and a variety of BmoR mutants with desired properties, such as specificity,

sensitivity or a wide detection range were obtained and are shown in Table 1. These

mutants could be widely utilized in various fields, including metabolic regulation,

enzyme or production strain screening, and byproduct detection.

Results and discussion

Analysis of the wild-type BmoR structure and establishment of the mutagenesis library

We first analyzed the binding regions of different molecules and attempted to utilize

rational design to endow BmoR with signal-molecule specificity. In our previous study,

the biofilm regulator FleQ (PDB code: 5EXP) and the transcription factor PobR (PDB

code: 5W1E) were used as templates to simulate wild-type BmoR (Fig. 1B). In this study,

molecular docking of BmoR with different alcohols (n-butanol, isobutanol, ethanol, and

glycol) was conducted. The results in Fig. 1C show that the binding pockets of different

signal molecules were similar, and the signal molecules bound to BmoR mainly via the

formation of hydrogen bonds with Arg211, Gln212, and Asn259. We first performed

site-saturation mutagenesis at these sites. The generated mutants had decreased response

values when compared to the wild-type (Fig. S1A, B, and C) and did not display

signal-molecule specificity. These results suggest the complexity of the response

mechanisms of BmoR and indicate that obtaining BmoR with desirable properties cannot

be easily achieved through rational design. Therefore, we performed random mutagenesis

of BmoR. The N-terminus of BmoR is responsible for recognizing and binding signal

molecules and modulating the activity of bEBP [25, 26]. Modifying the region could directly

influence the response of BmoR to the signal molecule and further change its properties.

Therefore, error-prone PCR was conducted at the N-terminal of BmoR to endow BmoR

with desirable properties. The entire screening process is illustrated in Fig. 1D. After

screening the mutagenesis library, we acquired 400 colonies that had different GFP

values when compared with wild-type BmoR. The *bmoR* genes in these colonies were

then extracted and sequenced.

Besides, we assumed the initiation mechanism of BmoR. BmoR and NtrC could both

activate the σ^{54}-dependent transcription. The regulatory mechanism of NtrC was positive

regulation[27]. In our previous study we have proved that the regulatory mechanism of

BmoR was also positive regulation[24]. Therefore, we speculated the initiation mechanism

of BmoR might be similar to NtrC because of their structural and functional similarity. To

induce the transcription, BmoR might be firstly expressed and formed as dimers. The

dimers of BmoR would then capture the signal molecules and the σ^{54} of σ^{54}-RNAP

holoenzyme ($E\sigma^{54}$) would bind to the -24 (GG) and -12 (TGC) regions of σ^{54}-dependent

promoter (P_{bmo}). After that, the dimers with signal molecules would form as hexamer to

initiate the transcription of P_{bmo} with the energy which would be released by ATP

hydrolysis. The whole activation process was shown in Fig. S2. The true regulation

mechanism of BmoR needed to be exploited by crystallization of BmoR with signal

molecules and in-depth analysis of the generated structures.

Specific response of BmoR mutants towards n-butanol

We isolated two special mutants (W21R/E54V and I183T/D273N) that show a significant response towards 10 mM of n-butanol and a weak response towards 10 mM of isobutanol (Fig. 2A). The GFP/OD$_{600}$ values of W21R/E54V and I183T/D273N towards 10 mM of n-butanol were 92.4 ± 14 and 15.6 ± 0.8, respectively (Fig. 2B). A series of gradient addition experiments with different concentrations of n-butanol or isobutanol were carried out to measure the apparent K_m value and to verify the detection potential of these two mutants. Significantly, the GFP/OD$_{600}$ values of W21R/E54V and I183T/D273N increased with the increase of n-butanol concentration (Fig. 2C), and the response values of W21R/E54V and I183T/D273N reached 325 ± 17 and 147 ± 12 when 100 mM n-butanol was added to the culture, respectively, while the response values of these two mutants have no obvious change with the increase of isobutanol concentration (Fig. 2D). These results suggest that W21R/E54V and I183T/D273N could perceive the change of n-butanol concentration and further give the corresponding response, and exhibited insensitivity towards isobutanol. In addition, the apparent K_m value of W21R/E54V was 42.6 mM towards n-butanol. The response value of I183T/D273N was linear to the concentration of n-butanol. Based on the above results, we verified the mutants W21R/E54V and I183T/D273N as n-butanol-specific BmoR mutants.

In addition, simulation and molecular docking of W21R/E54V and I183T/D273N with

isobutanol or n-butanol were conducted. The model in Fig. 2E shows that a hydrogen

bond was formed between n-butanol and the 261st glutamate of W21R/E54V, and there

was no force between isobutanol and W21R/E54V, suggesting that the interaction

between n-butanol and W21R/E54V was stronger than that between isobutanol and

W21R/E54V. In addition, we mutated the 261st glutamate to the other 19 amino acids.

The generated 19 mutants have significantly lower response values when compared with

W21R/E54V, which demonstrates the significance of Glu261 in the response to n-butanol

(Fig. S3). Similar to W21R/E54V, two hydrogen bonds between n-butanol and

I183T/D273N were observed in the complex, and no force was formed between

isobutanol and I183T/D273N, demonstrating that n-butanol could bind I183T/D273N

tightly (Fig. 2F).

Specific response of BmoR mutants towards isobutanol

In addition to the n-butanol-specific BmoR mutants, two mutants (M94V/F272L and

S240P) that significantly responded to 10 mM isobutanol and hardly responded to 10 mM

n-butanol, were screened from the mutagenesis library (Fig. 3A and B). The results of the

gradient addition experiment in Fig. 3C and D show that the GFP/OD$_{600}$ values of

M94V/F272L and S240P exhibited an increasing trend with the increase of isobutanol

concentration and had no significant change with the increase of n-butanol concentration.

These distinct responses illustrated that M94V/F272L and S240P have the ability to

perceive the change of isobutanol concentration and then output the corresponding

response signal; on the contrary, they could not sense n-butanol. Hence, the mutants

M94V/F272L and S240P could be considered as isobutanol-specific BmoR mutants.

Furthermore, complexes containing any of the above mutants and signal molecules

(n-butanol or isobutanol) were modeled, and the interactions between mutated BmoR and

the signal molecule were analyzed to explain the isobutanol specificity of M94V/F272L

and S240P. As shown in Fig. 3E, hydrophobic interactions existed between the signal

molecule isobutanol and Phe276 of M94V/F272L; meanwhile, no interaction was

observed in the complex of M94V/F272L and n-butanol. For the complexes of mutant

S240P and signal molecules, isobutanol bound to S240P via the formation of a hydrogen

bond at Glu261 and n-butanol did not interact with S240P (Fig. 3F). The interaction

between the isobutanol-specific mutants and signal molecules proved the isobutanol

specificity of the mutants.

Further verification of the specificity via the addition of mixed signal molecules

To further confirm the specificity and test whether the two kinds of signal molecules

competed to combine the mutants, the response of n-butanol-specific and

isobutanol-specific BmoR mutants was measured by adding mixed signal molecules to

the culture. We used an n-butanol-specific mutant (W21R/E54V) and an

isobutanol-specific mutant (M94V/F272L) as examples to conduct the experiments. As

shown in Fig. 4A, the GFP/OD$_{600}$ value of the n-butanol-specific mutant W21R/E54V

increased with an increase in the n-butanol proportion, reaching 342 ± 7 when the ratio of

n-butanol to isobutanol in the culture was 10:1 (100 mM:10 mM). As a comparison, the

GFP/OD$_{600}$ value of W21R/E54V did not change with the increase of isobutanol

proportion. The response of W21R/E54V only varied with the change of n-butanol

proportion, proving that there is no competition between n-butanol and isobutanol in the

n-butanol-specific mutant and that W21R/E54V could maintain its specificity in the case

of mixed signal molecules. Similar to W21R/E54V, the GFP/OD$_{600}$ value of

M94V/F272L only varied with the change of the isobutanol proportion. In summary, we

screened n-butanol-specific BmoR mutants (W21R/E54V and I183T/D273N) and

isobutanol-specific BmoR mutants (M94V/F272L and S240P) from the mutagenesis

library, and these mutants showed satisfactory specificity.

BmoR mutants with higher output and sensitivity

In addition to the above signal molecule-specific mutants, two mutants with higher

outputs were isolated from the random mutagenesis library. The sequencing results show

that these two mutants have mutations of E54G and V311A, respectively. Notably,

mutant E54G have a 27.5-fold higher GFP/OD$_{600}$ value (1941 ± 17) towards 1 mM

n-butanol and a 12.1-fold higher GFP/OD$_{600}$ value (1020 ± 17) towards 1 mM isobutanol

when compared with wild-type BmoR. In comparison, the GFP/OD$_{600}$ values of V311A

towards 1 mM n-butanol and 1 mM isobutanol were 34.2- and 19.5-fold higher than

wild-type BmoR, respectively (Fig. 4B). Based on this, we assumed that these two

mutants with higher output might respond to the low concentration of butanols,

signifying that the mutants have higher sensitivity. To validate this assumption, we added

0–1 mM n-butanol or isobutanol to the culture to test the response of mutants E54G and

V311A. The lower detection limit of BmoR was defined as the concentration of butanols

that could achieve a maximum GFP/OD$_{600}$ value of 75%. Based on this, we estimated the

lower detection limits of mutants E54G and V311A based on the Michaelis-Menten

equation of Origin8.5. As shown in Fig. 4C and D, the lower detection limits of mutants

E54G and V311A towards n-butanol were 2.64×10^{-6} mM and 2.13×10^{-6} mM,

respectively, which demonstrated that the sensitivity of these two mutants towards

n-butanol was over 1.0×10^{7}-fold higher than that of wild-type BmoR (28 mM). In

comparison, mutants E54G and V311A have lower detection limits of 2.16×10^{-6} mM

and 1.94×10^{-6} mM towards isobutanol, respectively, which suggests that these two

mutants possess over 1.0×10^{7}-fold higher sensitivity to isobutanol when compared with

wild-type BmoR (26 mM). After that, we fed 1×10^{-6} mM of n-butanol or isobutanol to

the cultures of mutants E54G and V311A, and used fluorescence microscope to observe

the green fluorescence. As shown in Fig. 4E, mutants E54G and V311A have significant

green fluorescence towards 1×10^{-6} mM of n-butanol or isobutanol. As a comparison,

wild-type BmoR has no obvious green fluorescence. These results suggested mutants

E54G and V311A possess high sensitivity to n-butanol and isobutanol. In addition, we

attempted to introduce the mutations in the mutant with high output into the signal

molecule-specific mutants in order to enhance the response values of the signal

molecule-specific mutants. However, the mutations in the mutant with high output did

not significantly enhance the output of signal molecule-specific mutants (data not shown).

Future work might focus on enhancing the sensitivity and output of signal

molecule-specific mutants in order to sense the trace accumulation of butanols.

BmoR mutant with wider detection range

To utilize the BmoR-based biosensor to screen industrial high-level n-butanol or

isobutanol production strains, we needed to broaden the detection range of BmoR. In this

study, we screened out a mutant (T12N) with a wider detection range (0–200 mM) when

compared with wild-type BmoR. Fig. 5A and B showed mutant T12N outputted

significantly distinct GFP/OD$_{600}$ values when feeding the 150 mM and 200 mM butanol

solutions and the apparent K_m values of T12N were 37.3 mM towards n-butanol and 89.9

mM towards isobutanol. These results suggest that the upper detection limit of mutant

T12N could reach 200 mM.

We then introduced mutant T12N into an isobutanol-producing strain to confirm its wider

detection range for *in vivo* host-producing isobutanol. First, plasmids pYH10 and

pYH10-T12N were individually introduced into the isobutanol-producing strain TW,

resulting in strains TW1 and TW2, respectively. TW1 and TW2 were used for subsequent

fermentation and GFP/OD$_{600}$ value detection. As shown in Fig. 5C and D, the GFP/OD$_{600}$

value of strain TW2 with T12N BmoR expression displayed an upward trend as the

isobutanol titer during the 84-hour fermentation. The GFP/OD$_{600}$ values at 72 h and 84 h

were significantly different, and the value at 84 h reached 835 ± 25. Meanwhile, 215 mM

isobutanol was accumulated in the culture. In comparison, the GFP/OD$_{600}$ value of strain

TW1 with wild-type BmoR expression increased with the production of isobutanol in the

first 36 h, and the isobutanol titer was 67 mM at 36 h. In the next 48 h, isobutanol was

continuously produced. However, the GFP/OD$_{600}$ value of TW1 did not increase

significantly. These results illustrate that mutant T12N demonstrates a wider detection

range (0–200 mM) towards *in vivo* host-producing isobutanol, and this mutant could be

used to construct a sensor to screen overproducers.

BmoR mutants with ethanol insensitivity

In the ethanol fermentation industry, n-butanol and isobutanol are usually produced as

byproducts. We assumed that the BmoR-based biosensor could serve as a powerful tool

to detect the accumulation of higher alcohols in ethanol-producing strains. However,

native BmoR showed a significant response to ethanol, which could induce signal

interference and lead to inaccurate measurement of higher alcohol concentrations.

Therefore, a biosensor that is insensitive to ethanol and sensitive to higher alcohols needs

to be urgently explored. Fortunately, the abovementioned two n-butanol-specific mutants

and two isobutanol-specific mutants did not respond to 0–200 mM ethanol (Fig. S4). The

apparent K_m values of mutants W21R/E54V and I183T/D273N towards ethanol were 228

and 174 mM, respectively, which were 23.5- and 17.9-fold higher than that of the

wild-type. Similarly, mutants M94V/F272L and S240P possess 13.6- and 12.1-fold higher

apparent K_m towards ethanol when compared with the wild type, respectively. The

insensitivity of these mutants to ethanol was maintained at a concentration of 800 mM

(Fig. 5E). The complexes in Fig. S5 show that there was no interaction between any of

the abovementioned mutants and ethanol. Notably, W21R/E54V and M94V/F272L

maintained their respective specificities even with background interference of 500 mM

ethanol (Fig. 5F). In addition, we tested the response of special BmoR mutants to other

alcohols including methanol, glycol, isopropanol, isopentanol, and 2-methyl-1-pentanol.

Fig. S6 shows that mutant T12N could respond to the abovementioned alcohols as well as

wild-type BmoR, and mutant W21R/E54V has a narrower signal molecule range when

compared with wild-type BmoR, suggesting that mutations in the mutants influenced the

range of signal molecules.

The promiscuity of TFBs is a double-edged sword. On the one hand, promiscuity could

provide some benefits. For instance, TFBs with promiscuity can expand the signal

molecule spectrum. On the other hand, the signal molecule diversity of TFBs could result

in an inaccurate response. Many studies have engineered biosensors to realize specific

responses [28, 29]. Endowing a BmoR-based biosensor with n-butanol specificity or

isobutanol specificity is required to expand its application. In addition to the native

sensing elements, some *de novo* designed biosensors that assemble specific protein

functional domains were also developed [30]. Based on the analysis of BmoR in this work,

the specific domains of BmoR might be served as crucial units for establishing novel

biosensors to detect other vital compounds.

Conclusions

The non-ideal properties of wild-type BmoR limit its applications for the screening of

single-higher-alcohol industrial-producing strains or the detection of byproducts in the

fermentation process. In this study, we engineered a significant region of BmoR and

acquired two n-butanol-specific mutants, two isobutanol-specific mutants, and two

ultrasensitive mutants. In addition, a mutant with a wider detection range (0–200 mM)

was screened out, which could also be reflected in the isobutanol-producing strain.

Notably, we observed that the signal-molecule-specific mutants display insensitivity

towards ethanol, indicating that these specific mutants could be used to detect the

production of byproducts (butanols) in the ethanol fermentation industry. In summary,

this work supplied various desirable BmoR mutants that could be employed for

constructing biosensors to screen ideal strains or establish dynamic control systems in the

field of metabolic engineering.

Methods

Strains, media, and materials

E. coli XL10-Gold was used for plasmid construction, screening, and construction of the

BmoR N-terminal-based random mutagenesis library. LB medium (10 g/L tryptone, 5 g/L

yeast extract, and 10 g/L NaCl) was used for strain incubation and library screening. M9

medium (6 g/L Na_2HPO_4, 3 g/L KH_2PO_4, 1 g/L NH_4Cl, and 0.5 g/L NaCl, 1 mM $MgSO_4$,

0.1 mM $CaCl_2$, 10 mg/L VB1, 4 g/L yeast extract, and 40 g/L glucose) was used for the

fermentation experiment. Plasmids pSA65, pSA69, pYH1, and pYH10 were obtained

from our previous study [16, 24]. The details of the strains and plasmids used in this study

are listed in Table S1.

Homology modeling and molecular docking of BmoR mutants with signal molecules

AUTODOCK and Chimera 1.14 were used for homology modeling and molecular

docking of BmoR with signal molecules (n-butanol, isobutanol, ethanol, or glycol). The

tertiary structure of wild-type BmoR, which was modeled in our previous study [24], was

used as a template to simulate and dock BmoR mutants with signal molecules. All BmoR

mutant models were evaluated by PROCHECK, and all models have satisfactory quality,

with over 80% of the residues in the most favored region of the calculated z-scores. A

grid box ($10 \times 17 \times 8$) encompassing the binding pocket of BmoR was set as the search

space to explore suitable substrate-binding regions. The interactions between the mutants

and signal molecules were analyzed and are shown in the corresponding figures.

Establishment of the BmoR N-terminal-based mutagenesis library

Error-prone PCR was carried out on the N-terminus of BmoR to generate an

N-terminal-based mutagenesis library. The details of the error-prone PCR method are

provided in the Supplementary Material. To carry out site-directed mutagenesis of *bmoR*,

a pair of primers, including the desired mutations, was synthesized and used to amplify a

linearized fragment from pYH1. The PCR products were purified and digested with DpnI

and then transferred into *E. coli* XL10-Gold. The primers used in this study are listed in

Table S2.

High-throughput screening of the mutagenesis library

Single colonies harboring plasmid pYH1 containing wild-type or mutated BmoR were

pre-inoculated into 5 mL LB with 100 μg/mL ampicillin and then cultured at 37 °C

overnight. Next, 50 μL of the seed culture was transferred into 950 μL LB, which was

supplemented with appropriate antibiotics and 10 mM n-butanol or isobutanol in

96-deep-well plates. The cultures were then left at 30 °C for 16 h. To measure the

apparent K_m values of the special BmoR mutants towards the butanols or ethanol, the

butanol concentrations in 96-deep-well plates ranged between 0–100 mM, and the

ethanol concentrations in 96-deep-well plates ranged between 0–800 mM. A microplate

reader (BioTek Cytation 3) was used to detect the OD_{600} values and GFP fluorescence.

The excitation and emission wavelengths were set at 470 nm and 510 nm, respectively.

The *bmoR* in the colonies that have distinct GFP/OD$_{600}$ values compared to the wild type

were sequenced. Apparent K_m values were estimated using Origin8.5 through non-linear

regression of the Michaelis-Menten equation. Fully automatic inverted biological

microscope (DMi 8) was used to observe green fluorescence. The entire screening

process is illustrated in Fig. 1D. The ratio of n-butanol to isobutanol in a mixture of

butanol in the culture is described in the Supplementary Material.

Fermentation verification of BmoR with a wider detection range

Plasmids pYH10 containing wild-type *bmoR* and pYH10-T12N containing *T12N* were

individually introduced into strain TW (JCL260 with pSA65, which contained genes *kivd*

and *adhA*, and pSA69, which contained genes *alsS*, *ilvC*, and *ilvD*), resulting in strains

TW1 and TW2, respectively. For the fermentation experiment, the single colonies were

pre-inoculated into 5 mL LB medium with associated antibiotics at 37 □ for 12–14 h.

Then, 200 μL of culture was inoculated into 20 mL M9 with 0.1 mM IPTG in a 250 mL

screw cap conical flask and left at 30 °C in a shaker at 220 rpm. After 36 h, 40 g/L

glucose was added to the culture. Samples were taken every 12 h for GFP/OD$_{600}$

measurements and isobutanol detection. GC analysis of the samples was performed as

described in our previous study [31].

AUTHOR INFORMATION

Corresponding Author

Yi-Xin Huo – *Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology, Beijing 100081, P. R. China; Tobacco Research Institute, Chinese Academy of Agricultural Sciences, Qingdao 266101, P. R. China;* Phone: 86-10-68918158;

Email: huoyixin@bit.edu.cn

Authors

Tong Wu – *Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology, Beijing 100081, P. R. China;*

Zhenya Chen – *Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology, Beijing 100081, P. R. China;*

Shuyuan Guo – *Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology, Beijing 100081, P. R. China;*

Cuiying Zhang – *Key Laboratory of Industrial Fermentation Microbiology, Ministry of Education, Tianjin Industrial Microbiology Key Laboratory, College of Biotechnology, Tianjin University of Science and Technology, Tianjin 300457, P. R. China;*

Autor Contributions

Y.-X.H. generated the idea. Y.-X.H., Z.C. and T.W. designed the project. T.W. and Z.C.

carried out the experiments. T.W., Z.C., S.G., C.Z. and Y.-X.H. analyzed the data. T.W.,

Z.C. and Y.-X.H. wrote the manuscript.

Autor Contributions

[#]T.W. and Z.C. contributed equally to this work.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Research reported in this publication was supported by the National Key R&D Program

of China (grant no. 2019YFA0904104), National Natural Science Foundation of China

(grant nos.21908007 and 32000059), the Science and Technology Innovation Program of

Chinese Academy of Agricultural Sciences (ASTIP TRIC05), and the Fundamental

Research Funds for the Central Universities. We thank Biological & Medical Engineering

Core Facilities (Beijing Institute of Technology) for providing advanced equipment.

References

(1) Hossain, G. S.,Saini, M.,Miyake, R.,Ling, H.,Chang, M. W., (2020) Genetic Biosensor Design for Natural Product Biosynthesis in Microorganisms. *Trends Biotechnol. 38* (7).

(2) Wan, X.,Volpetti, F.,Petrova, E.,French, C.,Maerkl, S. J.,Wang, B., (2019) Cascaded amplifying circuits enable ultrasensitive cellular sensors for toxic metals. *Nat. Chem. Biol. 15* (5), 540-548.

(3) Zhang, F.,Carothers, J. M.,Keasling, J. D., (2012) Design of a dynamic sensor-regulator system for production of chemicals and fuels derived from fatty acids. *Nat. Biotechnol. 30* (4), 354-359.

(4) Gao, C.,Hou, J.,Xu, P.,Guo, L.,Chen, X.,Hu, G.,Ye, C.,Edwards, H.,Chen, J.,Chen, W., (2019) Programmable biomolecular switches for rewiring flux in *Escherichia coli*. *Nat. Commun. 10* (1), 1-12.

(5) Xiong, D.,Lu, S.,Wu, J.,Liang, C.,Wang, W.,Wang, W.,Jin, J.-M.,Tang, S.-Y., (2017) Improving key enzyme activity in phenylpropanoid pathway with a designed biosensor. *Metab. Eng. 40*, 115-123.

(6) Liu, Y.,Zhuang, Y.,Ding, D.,Xu, Y.,Sun, J.,Zhang, D., (2017) Biosensor-based evolution and elucidation of a biosynthetic pathway in *Escherichia coli*. *ACS Synth Biol. 6* (5), 837-848.

(7) Yu, H.,Wang, N.,Huo, W.,Zhang, Y.,Zhang, W.,Yang, Y.,Chen, Z.,Huo, Y. X., (2019) Establishment of BmoR-based biosensor to screen isobutanol overproducer. *Microb. Cell Fact. 18* (1), 30.

(8) Koch, M.,Pandi, A.,Borkowski, O.,Batista, A. C.,Faulon, J. L., (2019) Custom-made transcriptional biosensors for metabolic engineering. *Curr. Opin. Biotechnol. 59*, 78-84.

(9) Liu, D.,Evans, T.,Zhang, F., (2015) Applications and advances of metabolite biosensors for metabolic engineering. *Metab. Eng. 31*, 35-43.

(10) Dietrich, J. A.,Shis, D. L.,Alikhani, A.,Keasling, J. D., (2013) Transcription factor-based screens and synthetic selections for microbial small-molecule biosynthesis. *ACS Synth Biol. 2* (1), 47-58.

(11) Boada, Y.,Vignoni, A.,Picó, J.,Carbonell, P., (2020) Extended Metabolic Biosensor Design for Dynamic Pathway Regulation of Cell Factories. *iScience 23* (7), 101305.

(12) Connor, M. R.,Liao, J. C., (2009) Microbial production of advanced transportation fuels in non-natural hosts. *Curr. Opin. Biotechnol. 20* (3), 307-315.

(13) Mascal, M., (2012) Chemicals from biobutanol: technologies and markets. *Biofuel Bioprod Bior. 6* (4), 483-493.

(14) Jin, C.,Yao, M.,Liu, H.,Chia-fon, F. L.,Ji, J., (2011) Progress in the production and application of n-butanol as a biofuel. *Renew Sust Energ Rev. 15* (8), 4080-4106.

(15) Atsumi, S.,Cann, A. F.,Connor, M. R.,Shen, C. R.,Smith, K. M.,Brynildsen, M. P.,Chou, K. J.,Hanai, T.,Liao, J. C., (2008) Metabolic engineering of *Escherichia coli* for 1-butanol production. *Metab. Eng. 10* (6), 305-311.

(16) Atsumi, S.,Hanai, T.,Liao, J. C., (2008) Non-fermentative pathways for synthesis of branched-chain higher alcohols as biofuels. *Nature 451* (7174), 86-89.

(17) Atsumi, S.,Wu, T.-Y.,Eckl, E.-M.,Hawkins, S. D.,Buelter, T.,Liao, J. C., (2010) Engineering the isobutanol biosynthetic pathway in *Escherichia coli* by comparison of three aldehyde reductase/alcohol dehydrogenase genes. *Appl. Microbiol. Biotechnol. 85* (3), 651-657.

(18) Chen, X.,Nielsen, K. F.,Borodina, I.,Kielland-Brandt, M. C.,Karhumaa, K., (2011) Increased isobutanol production in *Saccharomyces cerevisiae* by overexpression of genes in valine metabolism. *Biotechnol. Biofuels 4* (1), 1-12.

(19) Huo, Y.-X.,Cho, K. M.,Rivera, J. G. L.,Monte, E.,Shen, C. R.,Yan, Y.,Liao, J. C., (2011) Conversion
of proteins into biofuels by engineering nitrogen flux. *Nat. Biotechnol. 29* (4), 346-351.

(20) Huo, Y.-X.,Guo, L.,Ma, X., (2018) Biofuel production with a stress-resistant and growth
phase-independent promoter: mechanism revealed by in vitro transcription assays. *Appl. Microbiol.*
Biotechnol. 102 (6), 2929-2940.

(21) Li, S.,Wen, J.,Jia, X., (2011) Engineering *Bacillus subtilis* for isobutanol production by heterologous
Ehrlich pathway construction and the biosynthetic 2-ketoisovalerate precursor pathway overexpression.
Appl. Microbiol. Biotechnol. 91 (3), 577-589.

(22) Smith, K. M.,Cho, K.-M.,Liao, J. C., (2010) Engineering *Corynebacterium glutamicum* for isobutanol
production. *Appl. Microbiol. Biotechnol. 87* (3), 1045-1055.

(23) Kurth, E. G.,Doughty, D. M.,Bottomley, P. J.,Arp, D. J.,Sayavedra-Soto, L. A., (2008) Involvement of
BmoR and BmoG in n-alkane metabolism in '*Pseudomonas butanovora*'. *Microbiology 154* (1), 139-147.

(24) Yu, H.,Chen, Z.,Wang, N.,Yu, S.,Yan, Y.,Huo, Y.-X., (2019) Engineering transcription factor BmoR
for screening butanol overproducers. *Metab. Eng. 56*, 28-38.

(25) Wikström, P.,O'Neill, E.,Ng, L. C.,Shingler, V., (2001) The regulatory N-terminal region of the
aromatic-responsive transcriptional activator DmpR constrains nucleotide-triggered multimerisation. *J. Mol.*
Biol. 314 (5), 971-984.

(26) Xu, H.,Gu, B.,Nixon, B. T.,Hoover, T. R., (2004) Purification and characterization of the AAA+
domain of *Sinorhizobium meliloti* DctD, a σ54-dependent transcriptional activator. *J. Bacteriol. 186* (11),
3499-3507.

(27) De Carlo, S.,Chen, B.,Hoover, T. R.,Kondrashkina, E.,Nogales, E.,Nixon, B. T., (2006) The structural
basis for regulated assembly and function of the transcriptional activator NtrC. *Genes Dev. 20* (11),
1485-1495.

(28) Sun, X.,Li, Q.,Wang, Y.,Zhou, W.,Guo, Y.,Chen, J.,Zheng, P.,Sun, J.,Ma, Y., (2020) Isoleucyl-tRNA
synthetase mutant based whole-cell biosensor for high-throughput selection of isoleucine overproducers.
Biosens. Bioelectron. 172, 112783.

(29) Unger, E. K.,Keller, J. P.,Altermatt, M.,Liang, R.,Matsui, A.,Dong, C.,Hon, O. J.,Yao, Z.,Sun,
J.,Banala, S.,Flanigan, M. E.,Jaffe, D. A.,Hartanto, S.,Carlen, J.,Mizuno, G. O.,Borden, P. M.,Shivange, A.
V.,Cameron, L. P.,Sinning, S.,Underhill, S. M.,Olson, D. E.,Amara, S. G.,Temple Lang, D.,Rudnick,
G.,Marvin, J. S.,Lavis, L. D.,Lester, H. A.,Alvarez, V. A.,Fisher, A. J.,Prescher, J. A.,Kash, T.
L.,Yarov-Yarovoy, V.,Gradinaru, V.,Looger, L. L.,Tian, L., (2020) Directed Evolution of a Selective and
Sensitive Serotonin Sensor via Machine Learning. *Cell 183* (7), 1986-2002.e26.

(30) Quijano-Rubio, A.,Yeh, H. W.,Park, J.,Lee, H.,Langan, R. A.,Boyken, S. E.,Lajoie, M. J.,Cao,
L.,Chow, C. M.,Miranda, M. C.,Wi, J.,Hong, H. J.,Stewart, L.,Oh, B. H.,Baker, D., (2020) De novo design
of modular and tunable allosteric biosensors. *Nature*.

(31) Ma, L.,Guo, L.,Yang, Y.,Guo, K.,Yan, Y.,Ma, X.,Huo, Y.-X. J. B. f. b., (2020) Protein-based
biorefining driven by nitrogen-responsive transcriptional machinery. *13* (1), 1-14.

Figure legends

Fig. 1. The activation mechanism and simulated structure of wild-type BmoR. (A)
The simulation mechanism of wild-type BmoR. **(B)** The simulated structure and the
binding pocket of wild-type BmoR. **(C)** The binding pocket and binding sites of
wild-type BmoR with different signal molecules (isobutanol, n-butanol, ethanol or
glycol). **(D)** The whole screening process of the N-terminal mutagenesis library.

Fig. 2. Specific response of BmoR mutants towards n-butanol. (A) The green
fluorescence of mutants W21R/E54V and I183T/D273N towards 10 mM n-butanol or
isobutanol. **(B)** The response values of BmoR mutants W21R/E54V and I183T/D273N
towards 10 mM n-butanol or isobutanol. **(C)** The response curves of BmoR mutants
towards n-butanol. **(D)** The response curves of BmoR mutants towards isobutanol. **(E)**
Simulation and molecule docking of mutant W21R/E54V with n-butanol or isobutanol,
and confirmation of the binding sites with signal molecules. **(F)** Simulation and molecule
docking of mutant I183T/D273N with n-butanol or isobutanol, and confirmation of the
binding sites with signal molecules. Values and error bars represent mean and s.d. (n = 3),
respectively.

Fig. 3. Specific response of BmoR mutants towards isobutanol. (A) The green
fluorescence of mutants M94V/F272L and S240P towards 10 mM n-butanol or
isobutanol. **(B)** The response values of BmoR mutants M94V/F272L and S240P towards

10 mM n-butanol or isobutanol. **(C)** The response curves of BmoR mutants towards

n-butanol. **(D)** The response curves of BmoR mutants towards isobutanol. **(E)** Simulation

and molecule docking of mutant M94V/F272L with n-butanol or isobutanol, and

confirmation of the binding sites with signal molecules. **(F)** Simulation molecule docking

of mutant S240P with n-butanol or isobutanol, and confirmation of the binding sites with

signal molecules. Values and error bars represent mean and s.d. (n = 3), respectively.

Fig. 4. Verification of the specificity via adding mixed signal molecules and BmoR

mutants with higher outputs and sensitivity. **(A)** Maintaining the concentration (10

mM) of one kind of butanols and increasing the concentration of the other kind of

butanols with a gradient (0-100 mM) to confirm the specificity of BmoR mutants

W21R/E54V and M94V/F272L. **(B)** The response values of BmoR mutants E54G and

V311A towards 1 mM n-butanol or isobutanol. **(C)** The response curves of BmoR

mutants E54G and V311A towards n-butanol. **(D)** The response curves of BmoR mutants

E54G and V311A towards isobutanol. **(E)** The green fluorescence of mutants E54G and

V311A towards 1×10^{-6} mM n-butanol or isobutanol. Values and error bars represent

mean and s.d. (n = 3), respectively.

Fig. 5. BmoR mutant with wider detection range or ethanol insensitivity. **(A)** The

response curves of wild-type BmoR and mutant T12N towards n-butanol. **(B)** The

response curves of wild-type BmoR and mutant T12N towards isobutanol. **(C)** The

response curves of wild-type BmoR and mutant T12N towards isobutanol which was

produced by the isobutanol-producing strain. **(D)** The response values of wild-type BmoR

and mutant T12N towards 67 mM, 114 mM and 215 mM isobutanol. **(E)** The response

values of specific mutants via adding 0-800 mM ethanol in culture. **(F)** The response

values of W21R/E54V and M94V/F272L via adding 0-60 mM n-butanol or isobutanol

and 500 mM ethanol as noise. Values and error bars represent mean and s.d. (n = 3),

respectively. $^{*}P < 0.1$, $^{**}P < 0.01$ as determined by two-tailed t-test.

Table 1. Representative BmoR mutants with remarkable properties in this study

BmoR Mutant	Property					
	n-Butanol specificity	Isobutanol specificity	Insensitivity to ethanol	Higher sensitivity	Wider detection range	Higher output
W21R/E54V	√		√			
I183T/D273N	√		√			
M94V/F272L		√	√			
S240P		√	√			
E54G				√		√
V311A				√		√
T12N					√	

A

-24 -12

P_{bmo} **Inactive**

σ⁵⁴ RNAP

● Butanols

BmoR hexamer

Complex of BmoR hexamer with butanols

-24 -12

P_{bmo} **Active**

σ⁵⁴ RNAP

B

Glu261
Leu260
Asn259
Gln212
Arg211

C

n-Butanol

Isobutanol

Ethanol

Glycol

Binding sites	n-Butanol	Isobutanol	Ethanol	Glycol
	Arg 211 Gln 212 Asn 259 Glu 261	Arg 211 Gln 212 Asn 259	Arg 211 Gln 212 Asn 259 Leu 260 Glu 261	Arg 211 Gln 212 Leu 260

D

P_{bmoR}

bmoR

N

C

Amp

P_{bmo}

GFP

colE1

Wild-type BmoR

Mg²⁺

Mn²⁺

C

N

Error PCR for N-terminal

Mut-bmoR

P_{bmoR}

Amp

P_{bmo}

GFP

colE1

Mutagenesis library

Adding alcohols

GFP detection

BmoR mutants with various properties