Cytotoxic CD4[+] T cells driven by T-cell intrinsic IL-18R/MyD88 signaling predominantly infiltrate *Trypanosoma cruzi*-infected hearts

Carlos-Henrique. D. Barbosa[1,2], Ariel Gomes[1,3]*, Fábio B. Canto[1,4]*, Layza M. Brandão[1], Jéssica R. Lima[1], Guilherme A. Melo[1], Alessandra Granato[1,5], Eula G.A. Neves[6], Walderez O. Dutra[6], Ana C. Oliveira[7], Alberto Nóbrega[1] and Maria Bellio[1]**.

[1]Department of Immunology, Instituto de Microbiologia Paulo de Góes, Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, RJ, 21941-902 Brazil; Ph: +55-21-987496559

[2]present address: Department of Microbiology & Immunology, Chicago Medical School, Rosalind Franklin University of Medicine and Science, Chicago, IL, USA.

[3]present address: Faculdade de Medicina da Universidade Estácio de Sá, Angra dos Reis, RJ Brazil.

[4]present address: Departamento de Imunobiologia, Instituto de Biologia, Universidade Federal Fluminense (UFF) Niterói, RJ 24210-350, Brazil.

[5]present address: The Hospital for Sick Children - Genetics and Genome Biology. Toronto, ON, Canada

[6]Laboratório de Biologia das Interações Celulares, ICB, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

[7]Instituto de Biofísica Carlos Chagas Filho, UFRJ, Rio de Janeiro, RJ, Brazil.

Keywords: Infectious disease**,** cytotoxicity, myocarditis, parasite, tissue immunity.

* These two authors contributed equally to this work

**Corresponding Author and Lead Contact*:

e-mail: belliom@micro.ufrj.br or mariabellioufrj@gmail.com

Abstract:

Increasing attention has been directed to cytotoxic CD4$^+$ T cells (CD4CTLs) in different pathologies, both in humans and mice. The impact of CD4CTLs in immunity and the mechanisms controlling their generation, however, remain poorly understood. Here, for the first time, we showed that CD4CTLs abundantly differentiate during mouse infection with an intracellular parasite. CD4CTLs appear in the spleen in parallel to Th1 cells, display pathogen-derived peptide-specific cytotoxicity against antigen-presenting cells and express immunoregulatory and/or exhaustion markers. We demonstrated that CD4CTL absolute numbers and activity are severely reduced in both *Myd88*$^{-/-}$ and *Il18ra*$^{-/-}$ mice. Of note, the infection of mixed-bone marrow chimeras revealed that WT, but not *Myd88*$^{-/-}$, cells transcribe the CD4CTL gene signature and that *Il18ra*$^{-/-}$CD4$^+$ phenocopy *Myd88*$^{-/-}$CD4$^+$ T cells. Moreover, the adoptive transfer of WT CD4$^+$GzB$^+$ T cells to susceptible *Il18ra*$^{-/-}$ mice increased their survival. Importantly, cells expressing the CD4CTL phenotype predominate among CD4$^+$ T cells infiltrating the infected cardiac tissue, are increased in the circulation of Chagas patients and their frequency correlates with severe cardiomyopathy. Our findings describe CD4CTLs as a major player in immune response to a relevant human pathogen and disclose T-cell intrinsic IL-18R/MyD88 signaling as a key pathway controlling the magnitude of the CD4CTL response.

Introduction:

CD4CTLs are present in viral infections, cancer and autoimmunity (Cheroutre & Husain, 2013; Juno et al., 2017; Takeuchi & Saito, 2017). Although more frequently found in chronic conditions, CD4CTLs have also been described during acute viral infection (Brien et al., 2008; Brown et al., 2012; Fang et al., 2012; Jellison, et al., 2005; Wilkinson et al., 2012). At present, however, there are few studies on the CD4CTL response to infection with non-viral intracellular pathogens (Burel et al., 2016; Canaday et al., 2001; Kotov et al., 2018; Krueger et al., 2021). The role of CD4CTLs in immunity against intracellular parasites and the mechanisms operating in the generation and maintenance of this cell subset are poorly understood.

Following the recognition of the cognate antigen-MHC complex, and under the influence of cytokines secreted by antigen presenting cells (APCs), naïve CD4$^+$ T cells proliferate and differentiate into diverse subsets, such as Tfh, Th1, Th2 or Th17, with varied effector functions. Different pathogen-derived molecular patterns (PAMPs) elicit distinct innate immune pathways, resulting in tailored Th responses against the infectious agent (Kara et al., 2014). Interestingly, increasing heterogeneity in the transcriptional profile of effector CD4$^+$ T cells, dictated in a microbe-specific manner, has been recently found in the gut (Kiner et al., 2021). Additionally, tissue-derived factors can also be involved in defining CD4$^+$ T cell phenotypes (Poholek, 2021). In this context, the mechanism regulating the differentiation of CD4CTLs remains an open and relevant question. Although some authors consider CD4CTLs functional variants of Th1 cells, there is also evidence that CD4CTLs might constitute a Th1-independent subset, directly differentiating from naïve CD4$^+$ T cell precursors, or even a heterogeneous population congregating cytolytic cells that originate from different T-

cell subsets, such as Treg and Th2 cells [reviewed in (Cheroutre & Husain, 2013) and (Takeuchi & Saito, 2017)].

Receptors members of the TLR and IL-1R families are upstream of the MyD88 adaptor molecule and are expressed both in myeloid and lymphoid cell lineages. MyD88-dependent signaling in innate cells can influence Th1 cell differentiation, by inducing the release of IL-12 or IFN-γ (Joffre et al., 2009; Takeda et al., 1998), The T-cell intrinsic MyD88 adaptor function was also shown to be essential for the Th1 response in different contexts. The identity of the receptor acting upstream MyD88 on T cells, however, was not always determined (Frazer et al., 2013; LaRosa et al., 2008) and remains a matter of debate. Early *in vitro* studies have shown that IL-18 synergizes with IL-12 in the induction of IFN-γ production and T-bet expression by CD4$^+$ T cells and, therefore, impacts on Th1 development (Robinson et al., 1997). Intriguingly, however, the IL-18 contribution for a robust Th1 response *in vivo* has not always been confirmed (Haring & Harty, 2009; Monteforte et al., 2000). ST2 (IL-33 receptor) and IL-1R have also been implicated in the establishment of Th1 responses to viral infection (Baumann et al., 2015) and immunization (Ben-Sasson et al., 2009; Schenten et al., 2014), respectively. Recently, our group has disclosed the crucial role of T cell-intrinsic IL-18R/MyD88 signaling for the reinforcement and maintenance of the Th1 program *in vivo,* in response to mouse infection with *Trypanosoma cruzi,* where Th1 cells have a key protective function (Oliveira et al., 2017; Rodrigues et al., 2012).

T. cruzi is an intracellular protozoan parasite and the etiologic agent of Chagas disease (*American trypanosomiasis*), one of the most important parasitic diseases in the Western Hemisphere. Chagas disease is an important cause of heart failure, stroke, arrhythmia, and sudden death, being the leading cause of cardiomyopathy among younger individuals in Latin America (Bern, 2015; Nunes et al., 2018). Here, we

showed that CD4$^+$ T lymphocytes with CD4CTL phenotype are abundantly present in

the spleen of mice infected with *T. cruzi*, displaying *in vitro* and *in vivo* Ag-specific

cytotoxicity against APCs. Of note, we found that cells with CD4CTL phenotype

predominate among CD4$^+$ T lymphocytes infiltrating the infected cardiac tissue. Using

mixed bone marrow (mix-BM) chimeras, we showed that T cell-intrinsic IL-

18R/MyD88 signaling plays a crucial role on the expansion/survival of CD4$^+$ T cells

expressing the CD4CTL gene signature *in vivo*. In accordance with results obtained in

mix-BM chimeras, we found that absolute numbers of CD4$^+$ T cells displaying

cytotoxic markers and activity were severely reduced in *Myd88$^{-/-}$* and *Il18ra$^{-/-}$* mouse

lineages. Moreover, the adoptive transfer of WT CD4$^+$GzB$^+$ T cells to *Il18ra$^{-/-}$* mice

increased the survival of this susceptible mouse strain to infection. Interestingly, the

majority of CD4$^+$GzB$^+$ T cells in WT infected mice express high levels of

immunoregulatory molecules such as Tim-3, Lag-3 and CD39. Finally, and importantly,

we detected the presence of CD4$^+$ T cells expressing perforin in the circulation of

patients with the severe cardiac form of Chagas disease and correlation analyses point

to a possible pathogenic role of these cells in the chronic phase of Chagas myocarditis.

To our knowledge, the present study is the first to extensively characterize CD4CTLs

generated in response to a protozoan pathogen and to elucidate a central mechanism that

accounts for their expansion/survival. It is expected that pharmacological targeting of

the IL-18R/MyD88 signaling pathway might contribute to the strategic development of

new therapeutic and vaccine approaches aiming the control of the CD4CTL response.

Results:

CD4$^+$ T cells with cytolytic phenotype are abundantly present in the spleen of mice

infected with T. cruzi

CD4CTLs express granzyme B (GzB) and perforin (PRF), which are major

constituents of CD8CTL and NK cell-granules that promote target cell death in a

number of ways (Afonina et al., 2010). Here, we revealed the presence of a

TCR$\alpha\beta^+$CD8α^-NK1.1$^-$CD4$^+$GzB$^+$ T cell population, which represents around 30% of

CD4$^+$ T cells in the spleen of C57BL/6 (B6) mice, in the acute phase of infection with

T. cruzi (**Figures 1A-1B**). Equivalent percentages of CD4$^+$PRF$^+$ T cells were found

(**Figure 1C**) and the majority of CD4$^+$GzB$^+$ T cells express PRF and the cytolytic

marker 2B4 (CD244) (**Figures 1D-1E** and Figure 1-figure supplement 1A). Of note,

most CD4$^+$GzB$^+$ T cells (> 80%) do not express IFN-γ at day 14 post-infection (pi), the

peak time-point of the T cell response, and although CD4$^+$GzB$^+$ and CD4$^+$IFN-γ^+ cells

develop similar kinetics in the spleen, the absolute number of CD4$^+$GzB$^+$ T cells was

superior to that of CD4$^+$IFN-γ^+ cells, at every time point of infection (**Figures 1F-1G**).

It is known that intracellular parasitism by *T. cruzi* Y strain in the spleen is at its highest

point between day 12 and 15 pi, concomitant with the histologically observed massive

destruction of parasitized splenic macrophages (Cordeiro et al., 1997). Therefore, there

is a coincidence between parasite titers in the spleen and the frequency and absolute

numbers of CD4$^+$GzB$^+$ T cells. We also observed an increase in the percentage of

degranulating CD4$^+$ T cells at day 14 pi, as inferred by staining with anti-CD107a

(LAMP-1) mAb, comparable to the observed on CD8$^+$ T cells (**Figure 1H**), although

CD8CTLs are present at higher percentage (Figure 1-figure supplements 1B-E and 1G)

and absolute number (Figure 1-figure supplements 1F and 1H). In summary, these data

show the remarkable induction of CD4$^+$ T cells with cytolytic potential during infection

with *T. cruzi*, an effector subpopulation whose appearance in the spleen parallels and

largely outnumbers that of classic GzB-negative IFN-γ^+ Th1 cells.

CD4+ T cells from infected spleen exert Ag-specific cytotoxicity

To ascertain the presence of $CD4^+$ T lymphocytes with cytolytic function, we

performed a redirected cytotoxicity assay, where anti-CD3 mAb–coated B cell blasts

were used as target cells and co-incubated with either highly purified effector ($CD44^{hi}$)

$CD4^+$ T cells sorted from the spleen of infected mice, or naive ($CD44^{lo}$) $CD4^+$ T cells

from non-infected controls. The CTL activity was observed only when effector $CD4^+$ T

cells were employed (**Figure 2A**). Antigen-specific cytotoxicity using highly purified

$CD4^+$ T cells was also assayed *in vitro* against an I-A^{b+} macrophage cell line (IC-21),

loaded or not with *T. cruzi* amastigote total protein extract. Results shown in **Figure 2B**

demonstrate that CD4CTLs with cognate cytotoxic activity are present in the spleens of

infected mice. The FasL molecule is expressed on $CD8^+$ and $CD4^+$ T cells at the

immunological synapse following Ag-triggered degranulation and participates in

CD4CTL-mediated lysis of target cells (Bossi & Griffiths, 1999; Brown et al., 2009;

Kotov et al., 2018). As shown in **Figure 2B,** addition of anti-FasL mAb to the assay

significantly inhibited the specific lysis of target cells, evidentiating the involvement of

the FasL/Fas pathway in the killing process.

Then, we tested the *in vivo* killing of target splenocytes loaded with class II-

restricted peptides (SA85-1.1c and d), derived from a protein of the *T. cruzi*

transialidase superfamily (Millar & Kahn, 2000) (**Figure 2C).** The detailed Ag

repertoire of CD4CTLs could not be further assessed because of the scarce definition of

parasite-derived CD4 epitopes in the literature. To circumvent this, we infected mice

with ovalbumin (OVA)-expressing Y strain trypomastigotes (Y-OVA) (Gomes-Neto et

al., 2018) and compared the *in vivo* cytotoxic response against known OVA-derived

CD4 and CD8 epitopes: $OVA_{323-339}$ and $OVA_{257-264}$, respectively. As shown in **Figure**

2D, the same level of specific killing was attained against targets loaded with class I-

or class II-restricted OVA peptides. Together, these experiments confirmed the presence

of a CD4$^+$ T cell subpopulation with Ag-specific cytolytic function against MHC class

II-restricted epitopes in the spleen of mice infected with *T. cruzi.*

CD4$^+$GzB$^+$ T cells express transcription factors typical of CTLs

We then investigated the expression of transcription factors (TFs) typical of

cytotoxic cell subsets, such as CD8$^+$ T cells and Tregs. T-bet and Eomesodermin

(Eomes) bind to the promoters of the *Gzmb* and *Prf1* genes (Glimcher et al., 2004;

Intlekofer et al., 2005; Pearce et al., 2003) and are expressed in CD4CTLs (Curran et al.,

2013; Eshima et al., 2012; Qui et al., 2011). **Figures 2E** and **2F** show that the vast

majority of CD4$^+$GzB$^+$ T cells express T-bet, but do not express FoxP3. In accordance

with the cytolytic potential of CD4$^+$GzB$^+$ T cells in our model, 60% of the CD4$^+$GzB$^+$ T

cells express Eomes, while only a small percentage of GzB-negative CD4$^+$ T cells are

Eomes$^+$ (**Figure 2G**). Of note, at day 14 pi, the vast majority of both GzB$^+$ and GzB-

negative CD4$^+$ T lymphocytes are effector/memory cells CD44^{hi}CD62L^{lo} (Figure 2-

figure supplement 2).

CD4CTLs migrating to the gut gradually stop expressing ThPOK, while acquiring

the expression of RUNX3 (Mucida et al., 2013; Reis et al., 2013), which is known to

drive the CTL gene program in CD8$^+$ CTLs (Cruz-Guilloty et al., 2009; Egawa et al.,

2007). The long-form transcript encoding RUNX3 (*Runx3d*) is also expressed in Th1

cells, but not in Th2 or naïve cells (Djuretic et al., 2007). As shown in **Figure 2H**,

besides *Runx3d* mRNA, we found that CD4$^+$GzB$^+$ T cells, sorted from the spleen of

tamoxifen-treated and infected GzmbCreER^{T2}/ROSA26EYFP reporter mice (Bannard et

al., 2009) (Figure 2-figure supplement 3A), express ThPOK *(Zbtb7b gene)*-coding RNA

comparable to naive CD4$^+$ T cells, at day 14 pi, when their number and frequency are

at the highest levels in this organ (**Figure 1G**). The absence of CD8α-coding transcripts

confirms the purity of sorted CD4$^+$GzB$^+$ T cells (Figure 2-figure supplement 3B). These

data show that CD4$^+$GzB$^+$ T cells generated in response to infection with *T. cruzi,*

besides expressing, cytolytic markers, also express T-bet, Eomes and RUNX3, typical

TFs found in cytotoxic T cells.

CD4$^+$GzB$^+$ T cells express cytotoxic markers and immunoregulatory molecules

Tr1 cells are a subset of IL-10-producing CD4$^+$FoxP3$^-$ T cells that express GzB and

PRF and exert a potent suppressive function both in humans and mice (Roncarolo et al.,

2018). Since a high percentage of CD4$^+$GzB$^+$ T cells in our system express Eomes,

which drives the development of Tr1 cells (Zhang et al., 2017), we investigated whether

T. cruzi-induced CD4$^+$GzB$^+$ T cells would share other phenotypic characteristics with

Tr1 cells. We detected a very low frequency of IL-10-producing TCRαβ$^+$CD4$^+$ T cells

in the spleens of mice infected with *T. cruzi*, in agreement with previous data (Jankovic

et al., 2007), although IL-10$^+$ cells were found to be more frequent among CD4$^+$GzB$^+$ T

cells (**Figures 3A** and Figure 3-figure supplement 1A-B). The co-expression of Lag-3

and CD49b was described to define both human and mouse Tr1 cells (Gagliani et al.,

2013). Although most splenic CD4$^+$GzB$^+$ T cells express Lag3, only a minor percentage

of these cells co-express CD49b (**Figures 3B-3C** and Figure 3-figure supplement 1C).

On the other hand, a substantial fraction of splenic CD4$^+$GzB$^+$ T cells express the Tim-3

immunoregulatory molecule and downregulate CD27 (**Figures 3D-3E**), indicating that

these might be terminally differentiated T cells, as suggested in other models (Anderson

et al., 2016; Takeuchi & Saito, 2017).

Natural killer group 2 (NKG2) members A, C, D and E are C-type lectin-like

receptors and cytotoxic markers expressed by NK, NKT, γδ and activated CD8$^+$ T

cells (Gunturi et al., 2004; Stojanovic et al., 2018). Recently, NKG2C/E and NKG2D

molecules were shown to identify a CD4CTL population, present in mouse lungs during

influenza A virus (IAV) infection (Marshall et al., 2017). As shown in **Figures 3F-3G**,

the fraction of cells expressing NKG2A/C/E and NKG2D was increased among splenic

$CD4^+GzB^+$ T cells, when compared either to GzB-negative $CD4^+$ T cells of infected

mice, or with naive $CD4^+$ T cells of non-infected controls (Figure 3-figure supplement

2A-B). The class I-restricted T cell associated molecule (CRTAM) was described as

being critical to instruct the differentiation of CD4CTLs (Takeuchi et al., 2016;

Takeuchi & Saito, 2017). **Figure 3H** shows that the frequency of CRTAM-expressing

cells was also increased among splenic $CD4^+GzB^+$ T cells of *T. cruzi*-infected mice,

when compared with GzB-negative $CD4^+$ T cells. The frequencies of $CRTAM^+$,

$NKG2A/C/E^+$ and $NKG2D^+$ $CD8^+GzB^+$ T cells in infected mice are shown in Figure 3-

figure supplement 3. Since at day 14 pi, both GzB^- and GzB^+ $CD4^+$ T cells are in their

majority effector/effector memory cells (Figure 2-figure supplement 2), equivalent

results were obtained gating on $CD44^{hi}$ cells or not, as shown in Figure 3-figure

supplement 4. Together, these results show that a significant fraction of the $CD4^+GzB^+$

T cell subpopulation found in this unicellular parasite infection model complies with the

general phenotype of CD4CTLs and CD4CTL-precursors reported in response to viral

infection.

CD4CTLs are present in lower numbers in Myd88$^{-/-}$ infected mice

To extend our understanding on the requirements for the differentiation of

CD4CTLs and their relationship to Th1 cells, we next investigated the role of IL-12,

IFN-γ and MyD88 in their generation. IL-12 and IFN-γ are the major cytokines driving

Th1 differentiation and T-bet expression, but their requirement for CD4CTL

generation is controversial (Brown et al., 2009; Cooper et al., 2004; Curran et al., 2013;

Hua et al., 2013; Kotov et al., 2018; Krueger et al., 2021; Tagawa et al., 2016; Xie et al.,

2010). Here, we observed that $Il12p40^{-/-}$ mice infected with *T. cruzi* displayed lower

percentages of T-bet$^+$CD4$^+$ and IFN-γ$^+$CD4$^+$ T cells and exhibited very low levels of

CD4$^+$GzB$^+$ T cells (**Figure 4A**). Interestingly, in infected $Ifng^{-/-}$ mice, despite the lower

frequency of T-bet$^+$CD4$^+$ T cells, CD4$^+$GzB$^+$ T cells attained the same frequencies as

observed in the spleens of WT animals (**Figure 4B**). However, absolute numbers of

both total CD4$^+$ and CD4$^+$GzB$^+$ T cells were lower in $Ifng^{-/-}$ mice when compared to

WT controls. Thus, both IL-12p40 and IFN-γ play important but distinct roles in the

differentiation and expansion/survival of CD4$^+$GzB$^+$ T cells.

The above results obtained in $Il12p40^{-/-}$ and $Ifng^{-/-}$ mice, together with the fact that

$Myd88^{-/-}$ mice have lower levels of Th1 cells and are highly susceptible to infection with

T. cruzi (Campos et al., 2004; Oliveira et al., 2010), prompted us to address the role of

MyD88-dependent pathways in the generation of CD4CTLs. Although infected $Myd88^{-/-}$

mice displayed equivalent absolute numbers and percentages of total splenic CD4$^+$ T

cells when compared to WT mice (Figure 4-figure supplement 1A), the frequencies and

absolute numbers of CD4$^+$GzB$^+$ T cells were significantly lower in this mouse strain

(**Figures 4C-4E**). Furthermore, absolute numbers of CRTAM$^+$, NKG2D$^+$ and

NKG2A/C/E$^+$ CD4$^+$GzB$^+$ T cells were severely diminished in $Myd88^{-/-}$ mice (**Figures**

4C-4E). This picture does not result from lower numbers of total activated CD4$^+$ T cells

in infected $Myd88^{-/-}$ mice, as shown by the percentages and absolute numbers of CD44^{hi}

cells, which are equivalent to WT animals (Figure 4-figure supplement 2). On the other

hand, absolute numbers of CRTAM$^+$, NKG2D$^+$ and NKG2A/C/E$^+$ CD8$^+$GzB$^+$ T cells

were not affected in $Myd88^{-/-}$ mice (Figure 4-figure supplement 1C-D). Concerning the

expression of perforin, both the percentages and absolute numbers of PRF$^+$CD4$^+$ T

cells were also severely decreased in $Myd88^{-/-}$ compared to WT mice (**Figure 4F**). The

same was observed when analyzing the frequency and absolute numbers of PRF$^+$CD4$^+$

T cells among GzB-expressing cells (**Figure 4G**). In order to establish to what extent

the absence of MyD88 expression would impact on cytotoxicity mediated by CD4$^+$ T

splenocytes, we sorted activated CD4$^+$CD44^{hi} T cells from both WT and $Myd88^{-/-}$

infected mice to a high purity level and performed redirected cytotoxicity assay. Lower

levels of cytotoxicity were observed when sorted $Myd88^{-/-}$ CD4$^+$CD44^{hi} T cells were

employed, as compared to assays using equal numbers of sorted WT CD4$^+$CD44^{hi} T

cells (**Figure 4H**). Together, these results demonstrate the important role of MyD88

signaling for the establishment of a robust cytotoxic response mediated by CD4CTLs.

T cell-intrinsic IL-18R/MyD88 signaling plays a crucial role in the establishment of a

CD4$^+$ T cell subpopulation bearing the CD4CTL gene signature

MyD88 deficiency results in lower IL-12 production by APCs (Campos et al., 2004)

and this could explain the diminished numbers of CD4CTLs found in infected $Myd88^{-/-}$

mice. However, we have recently shown that IL-18R/MyD88 signaling intrinsic to

CD4$^+$ T cells is critical for Th1 cell proliferation and resistance to apoptosis in response

to infection with *T.* cruzi (Oliveira et al., 2017). If CD4CTLs herein described share a

common early differentiation process with (or derive from) Th1 cells, it is reasonable to

expect that besides IL-12 and IFN-γ, T-cell-intrinsic MyD88-dependent signaling would

also be important for the generation of a robust CD4CTL response. In order to test this

hypothesis, we constructed mix-BM chimeras, in which WT (B6 x B6.SJL) F1 mice

were irradiated and reconstituted with a 1:1 mix of WT and $Myd88^{-/-}$ BM cells. After

reconstitution, mix-BM chimeras were infected with *T. cruzi* and, at day 14 pi, we

sorted WT (CD45.1) and $Myd88^{-/-}$ (CD45.2) CD44$^+$CD4$^+$ T cells to a high purity level

for transcriptome analysis. We have then investigated whether the transcriptional

signature of CD4CTLs (Donnarumma et al., 2016) was present in sorted CD4$^+$ T

splenocytes and to what extent the absence of T-cell intrinsic MyD88-dependent

signaling would impact on the acquirement of this transcriptional program. This

signature, previously described in the experimental model of Friend virus infection,

distinguishes CD4CTLs from other CD4$^+$ T cell subsets, including Th1 and Tfh cells

(Donnarumma et al., 2016). Importantly, gene set enrichment analysis (GSEA),

comparing the transcriptome of sorted WT and *Myd88*$^{-/-}$ CD4$^+$ T cells, revealed that 51

genes of the CD4CTL signature were upregulated in WT CD4$^+$ T cells, while 78 were

downregulated in *Myd88*$^{-/-}$ CD4$^+$ T cells (**Figures 5A-5C**). We confirmed, by flow

cytometry, the upregulation of Blimp-1 (*Prdm1* gene) TF and of the ecto-nucleotidase

CD39 (*Entpd1*) in WT CD4$^+$GzB$^+$ T cells (**Figures 5D-5E**). Of note, Blimp-1 is

required for the cytotoxic function of CD4CTLs (Hua et al., 2013) and CD39 expression

level was found to be higher in CD4$^+$GzB$^+$ T cells than in Tregs (**Figure 5E**).

In order to identify the receptor signaling upstream MyD88 responsible for the

establishment of a robust CD4CTL response, we further constructed three groups of

mix-BM chimeras, by mixing 1:1 WT and (i) *Il1r1*$^{-/-}$, (ii) *Il18ra*$^{-/-}$ or (iii) *Myd88*$^{-/-}$ BM

cells. A fourth group of mix-BM chimeras, 1:1 WT(B6):WT (B6.SJL) control was also

generated. After infection with *T. cruzi*, the percentages and absolute numbers of GzB$^+$

cells among CD4$^+$ T cells were analyzed in each chimeric group (gate strategy in Figure

5-figure supplement 1). **Figure 5F** shows that the lack of IL-18R signaling in T cells,

but not of IL-1R, phenocopied the absence of T cell-intrinsic MyD88 signaling,

revealing that IL-18R is a critical pathway upstream MyD88 necessary for the

establishment of the CD4$^+$GzB$^+$ T cell population.

In order to further consolidate the role of IL-18R signaling in the generation of a

strong CD4CTL response, we analyzed the expression of PRF as well as of CRTAM,

NKG2D and NKG2A/C/E cytotoxic markers on CD4$^+$GzB$^+$ T cells from the spleens of

Il18ra$^{-/-}$ mice. **Figure 6A** shows that both the frequency and absolute number of

PRF$^+$CD4$^+$ T cells were significantly diminished in the spleens of infected *Il18ra$^{-/-}$*

compared to WT mice. The same was true for PRF-expressing cells among GzB$^+$CD4$^+$

T lymphocytes (**Figure 6B**). As observed in infected *Myd88$^{-/-}$* mice, the frequency of

GzB$^+$CD4$^+$ T cells and their absolute number were also decreased in infected *Il18ra$^{-/-}$*

compared to WT mice (**Figures 6C-6E**). Absolute numbers of CRTAM$^+$, NKG2D$^+$ and

NKG2A/C/E$^+$ GzB$^+$CD4$^+$ T cells were significantly diminished in *Il18ra$^{-/-}$* mice, as

well, again phenocopying results obtained in *Myd88$^{-/-}$* mice (**Figures 6C-6E**). The

cytotoxic potential of CD4$^+$ T splenocytes from *Il18ra$^{-/-}$* infected animals was also

evaluated by performing redirected cytotoxicity assays with equal numbers of

CD4$^+$CD44^{hi} T cells purified from infected *Il18ra$^{-/-}$* or WT mice. Again, as shown for

*Myd88$^{-/-}$*CD4$^+$ T cells, lower levels of target killing were observed when employing

purified *Il18ra$^{-/-}$*CD4$^+$CD44^{hi} T cells as effectors (**Figure 6F**). Collectively, the present

data demonstrate that T-cell intrinsic IL-18R/MyD88 signaling support the generation

and/or survival of CD4$^+$ T cells with cytotoxic potential in response to parasite

infection, revealing a parallel between the requirements for the development of robust

Th1 and CD4CTL responses.

The adoptive transfer of CD4$^+$GzB$^+$ T cells to Il18ra$^{-/-}$ mice increased survival to

infection

In vivo studies addressing the function of CD4$^+$GzB$^+$ T cells have been hampered by

the absence of a stable and abundantly expressed surface-marker of cytotoxicity,

which would allow purifying these cells alive. To circumvent this, we employed

infected and tamoxifen-treated GzmbCreER^{T2}/ROSA26EYFP reporter mice, from

whose spleens CD4$^+$EYFP$^+$ (GzB$^+$) cells were highly purified by flow cytometry and

adoptively transferred to infected *Il18ra$^{-/-}$* mice, following the scheme shown in Figure

6-figure supplement 1. The *Il18ra$^{-/-}$* mouse lineage was previously shown to be very

susceptible to infection with *T. cruzi* (Oliveira et al., 2017), and display lower numbers

of CD4$^+$GzB$^+$ and CD4$^+$PRF$^+$ T cells with cytotoxic potential, as demonstrated above.

The transfer of highly purified CD4$^+$GzB$^+$(EYFP$^+$) T cells to *Il18ra$^{-/-}$* mice significantly

extended their survival rate without, however, decreasing the parasite load in the hearts

(Figure 6G). In a previous work, we have demonstrated that the transfer of total WT

CD4$^+$ T cells protects infected *Il18ra$^{-/-}$* mice, and this property was attributed to Th1

cells, which are also diminished in this mouse lineage (Oliveira et al., 2017). Since

infected *Ifng$^{-/-}$* mice have the same frequency of CD4$^+$GzB$^+$ T cells as WT mice in their

spleens (**Figure 4B**), we tested whether the adoptive transfer of CD4$^+$ T cells sorted

from *Ifng$^{-/-}$* mice would also be able to provide some degree of protection to *Il18ra$^{-/-}$*

mice. As shown in Figure 6-figure supplement 2, indeed, this was the case. Although

IFN-γ expression by CD4$^+$ T cells is necessary for decreasing parasitemia (Oliveira et

al., 2017), this result indicated that the secretion of IFN-γ by adoptively transferred

CD4$^+$ T cells is not absolutely required for increasing the survival time of infected

Il18ra$^{-/-}$ recipients. Together, these data demonstrate the protective function of the

CD4$^+$GzB$^+$ T cell subset against mortality caused by an intracellular parasite, which is

independent of decreasing heart parasitism.

The great majority of CD4$^+$ T cells in the infected myocardium express the CD4CTL

phenotype and the frequency of CD4$^+$PRF$^+$ T cells correlates with cardiomyopathy in

Chagas patients

Previous studies in viral infection models suggested that CD4CTLs are present

in tissues where pathogen replication occurs (Marshall et al., 2017; Takeuchi et al.,

2016). Since *T. cruzi* heavily infects the heart, we performed intravascular staining with

anti-CD4 mAb and demonstrated that tissue-infiltrating CD4$^+$ T cells are abundant in

the myocardium of infected mice (Figure 7-figure supplement 1A-B). As shown in

Figures 7A-7B, CD4$^+$GzB$^+$ T cells infiltrating the cardiac tissue constitute the vast

majority of CD4$^+$ T cells. At the peak time-point of parasite load in the heart,

CD4$^+$GzB$^+$ T cells in the myocardium express the LAMP-1 degranulation marker at

levels comparable to those of CD8$^+$ T cells (**Figures 7C-D** and Figure 7-figure

supplement 1C). Importantly, the great majority of heart-infiltrating CD4$^+$GzB$^+$ T cells

expresses PRF, as well as the CRTAM and NKG2D cytotoxic markers (**Figure 7E-7F)**.

On the other hand, only few CD4$^+$GzB$^+$ T cells express NKG2A/C/E in the

myocardium (Figure 7-figure supplement 1D, left), although high percentages of

NKG2A/C/E$^+$CD8$^+$GzB$^+$ T cells were found both in the heart (Figure 7-figure

supplement 1D, right) and in the spleen (Figure 3-figure supplement 3A) of infected

mice.

Finally, we investigated the presence of PRF$^+$CD4$^+$ T cells in the circulation of

patients infected with *T. cruzi* and suffering from chronic Chagas cardiomyopathy (CC).

In order to ascertain the specificity of CD4$^+$ T cells against *T. cruzi*, PBMC from CC

patients or from healthy donors (HD) were incubated (or not) with trypomastigote-

derived antigens and analyzed for the expression of perforin by flow cytometry (Figure

7-figure supplement 2). As shown in **Figures 7G and 7H,** after treatment, both the

expression level of PRF (MFI) and the percentage of PRF$^+$CD4$^+$ T cells increased

among CD4$^+$ T cells from CC patients, but not from HD controls. Of note, CD4$^+$ T cells

expressing PRF were found at higher frequency in CC patients and their level of PRF

expression was also higher than the observed in CD4$^+$ T cells of HD (**Figure 7I**). We

next performed correlative analyses between the frequency of PRF-expressing CD4$^+$ T

cells in PBMC from CC patients and two echocardiographic parameters, usually

employed to evaluate cardiac function in CC patients: left ventricular ejection fraction

(LVEF) and left atrium diameter. As shown in **Figure 7J,** a clear inverse correlation

between the frequency of CD4$^+$PRF$^+$ T cells and LVEF, as well as a direct correlation

between the frequency of these cells and the left atrium diameter was found. These

correlations suggest that CD4CTLs may play a pathogenic role in the chronic phase of

Chagas disease. Together, data obtained here in CC patients and in the mouse

experimental model point to the heart as an important target-organ for CD4$^+$ T cells

with cytolytic potential generated in response to infection with *T. cruzi.*

Discussion:

Here we show, for the first time, that CD4CTLs are abundantly present in the

spleen of mice infected with *T. cruzi* and that cells with CD4CTL phenotype

predominate among CD4$^+$ T cells infiltrating the infected cardiac tissue. The expression

of LAMP-1 degranulation marker by CD4$^+$GzB$^+$ T cells in the heart and in the spleen

indicates that these cells may play important effector functions in both organs. These

data suggest that CD4$^+$GzB$^+$ T cells with cytotoxic potential are attracted from the

spleen to the cardiac tissue and preferentially accumulate there. CD4$^+$GzB$^+$ T cells

present in the *T. cruzi*-infected hearts express PRF, CRTAM and NKG2D cytotoxicity

markers, but differently from CD4CTLs found in the lungs of mice infected with IAV

(Marshall et al., 2017), do not express NKG2A/C/E molecules. This interesting

dissimilarity requires further investigation and could be due to differential regulation

exerted by tissue-specific factors or, alternatively, by different innate responses

influencing CD4CTL differentiation, which would be distinctly triggered by viral and

parasite pathogens.

CD4$^+$ T cells purified from the spleen of infected mice killed APCs presenting

parasite-derived antigens *in vitro*. *In vivo* Ag-specific cytotoxicity against I-A^b-

restricted epitopes was also observed. Killing infected APCs in the spleen would lead to

a decrease in the activation of effector T cells, dampening the exuberant immune

response against the parasite. Therefore, the action of CD4CTLs may contribute to

immunoregulation, avoiding the development of exacerbated immunopathology, which

can cause mortality in the acute phase of the infection with *T. cruzi*. Favoring this

hypothesis, we showed that the adoptive transfer of CD4$^+$GzB$^+$ T cells to susceptible

Il18ra$^{-/-}$ mice, which have lower levels of these cells, significantly increased their

survival to infection, without affecting parasite loads. Together with our previous work

(Oliveira et al., 2017), the current results point to possible complementary roles of

CD4CTL and Th1 cells. Both cell types might be required for survival against infection

with *T. cruzi*, working together to establish a balance between the control of parasite

load on one side, and dampening of the immunopathology resulting from this process,

on the other. This equilibrium is associated with the establishment of the asymptomatic

form of chronic infection (Nunes et al., 2018), in which increased levels of CD4$^+$GzB$^+$

T cells have been found in humans (Keesen et al., 2012). In accordance with a putative

regulatory role played by CD4$^+$GzB$^+$ T cells, we showed here that these cells express

surface co-inhibitory receptors at particularly high levels, such as Lag-3 and Tim-3, as

well as the ecto-ATPase CD39. These immunoregulatory molecules, typically

expressed by Treg and Tr1 cells, have been implicated in suppressive function

(Anderson et al., 2016; Mascanfroni et al., 2015; Park et al., 2019).

Chronic Chagas cardiomyopathy usually develops in around 30% of infected

individuals after decades, probably reflecting an imbalance between parasite and host

immune response (Nunes et al., 2018). Although the exact mechanism remains to be

elucidated, the association of NK, CD8CTL and GzA-expressing $CD4^+$ T cells with the

development of chronic carditis has been indicated (Ferreira et al., 2017; Menezes et al.,

2012; Nunes et al., 2018). Since CD4CTLs recognize the cognate antigen in the context

of MHC class II molecules, the lysis of infected MHC II^+ endothelial cells might trigger

immunopathology (Reis et al., 1993). In the present study we found that the frequency

of parasite-specific $CD4^+PRF^+$ T cells is increased in the circulation of chronic cardiac

patients with Chagas disease (CC) and that PRF expression in $CD4^+$ T cells is higher in

the CC group when compared to healthy donors. A correlation between the frequency of

PRF^+CD4^+ T cells and two echocardiographic parameters employed for evaluating

myocarditis in CC patients was also encountered. It is tempting to speculate that the loss

of regulatory function exerted by different $CD4^+$ T cell subsets, including CD4CTLs,

might lead to the chronic pathology. Therefore, further studies are necessary for fully

defining the role of tissue infiltrating CD4CTLs in Chagas disease. Interestingly,

NKG2D, expressed by most CD4CTLs in the heart, can mediate both activating and

inhibitory signals (Wensveen et al., 2018). Importantly, a polymorphism in one of the

NKG2D ligands, the major histocompatibility complex class I–related chain A gene

(MICA) molecule, was associated with left ventricular systolic dysfunction in patients

with chronic Chagas myocarditis (Ayo et al., 2015).

We have shown here that $CD4^+GzB^+$ T cells originating in response to *T. cruzi*

infection largely outnumber IFN-γ-producing $CD4^+$ T cells and that their appearance

in the spleen follows the same kinetics of classic Th1 cells, indicating thereby that

differentiation of CD4CTLs is not a late event during infection. Of note, more than 60%

of CD4$^+$GzB$^+$ T cells have downregulated CD27 expression in the spleen at day 14 pi, a

characteristic of terminally differentiated T cells (Takeuchi & Saito, 2017). We have

also shown here that the majority of CD4$^+$GzB$^+$ T cells express T-bet, Eomes and

Blimp-1. Although Eomes seems not to be essential for CD4CTLs in certain studies

(Donnarumma et al., 2016; Kotov et al., 2018), it was proposed to be the master

regulator for CD4CTLs and Tr1 cells by others (Curran et al., 2013; Eshima et al.,

2012; Qui et al., 2011; Zhang et al., 2017). The CD4$^+$GzB$^+$ T cell population described

herein presents some characteristics of Tr1 cells. However, we only detected a very low

frequency of CD4$^+$GzB$^+$ T cells co-expressing Lag-3 and CD49b (Gagliani et al., 2013)

or IL-10. In this regard, data from *T. cruzi* infection differs from those observed during

infection with other intracellular parasites such as *T. gondii* and certain *L. major* strains,

in which high frequencies of CD4$^+$ T cells producing both IL-10 and IFN-γ were

described (Anderson et al., 2007; Jankovic et al., 2007). In contrast with ThPOK-

negative CD4CTLs that accumulate in the gut (Cheroutre & Husain, 2013; Mucida et al.,

2013; Reis et al., 2013), CD4$^+$GzB$^+$ T cells in the *T. cruzi*-infected spleen express

ThPOK at the peak of infection. Interestingly, the absence of ThPOK downregulation

was also observed in mouse (Mittal et al., 2018) and human CD4CTLs (Serroukh et al.,

2018). Thus, different subsets of CD4CTL possibly exist, in which the interplay

between ThPOK and RUNX3 expression might be differentially regulated.

Although many authors consider CD4CTLs merely functional variants of Th1 cells,

in some models CD4CTLs fail to fit Th1-defining criteria, such as the dependence on

IL-12, STAT4 or T-bet for differentiation (Brown et al., 2009; Curran et al., 2013; Hua

et al., 2013; Kotov et al., 2018). Nonetheless, in agreement with other studies (Cooper

et al., 2004; Krueger et al., 2021; Tagawa et al., 2016; Xie et al., 2010), we showed here

that CD4$^+$GzB$^+$ T cells are scarcely generated in the absence of IL-12p40. Importantly,

we demonstrated that the expansion/survival of CD4CTLs also depends on T-cell

intrinsic IL-18R/MyD88 signaling, which we have previously shown to induce cell

cycling, anti-apoptotic and Th1-signature transcripts (Oliveira et al., 2017). Notably,

several genes of the CD4CTL signature, which distinguishes CD4CTLs from Th1 and

Tfh cells (Donnarumma et al., 2016), were expressed by WT, but not by $Myd88^{-/-}$ CD4$^+$

T cells from infected mix-BM chimeras. Therefore, analogous to its action on Th1 cells,

we showed here that T-cell intrinsic IL-18R/MyD88 signaling contributes for the

reinforcement and/or stabilization of CD4$^+$ T cell commitment into the CD4CTL

phenotype. IFN-γ is another key cytokine involved in Th1 differentiation and $Ifng^{-/-}$

mice are extremely susceptible to infection with $T.$ $cruzi$ (Campos et al., 2004).

Interestingly, we found no difference in the frequency of CD4$^+$GzB$^+$ T cells between

WT and $Ifng^{-/-}$ infected mice, in accordance with a study showing that IFN-γ had no

impact on the ability of CD4 effectors to acquire cytolytic activity (Brown et al., 2012).

However, both GzB$^+$ and total CD4$^+$ T-cell numbers were significantly lower in the

infected $Ifng^{-/-}$ animals, indicating that IFN-γ might also play a role in the

expansion/survival of both Th1 and CD4CTLs in our system. It is known that IFN-γ and

IL-18R participate in a reciprocal positive feedback loop in Th1 differentiation process:

while IFN-γ is required for IL-12-driven upregulation of IL-18R expression (Nakahira

et al., 2001; Smeltz et al., 2001), IL-18 is a major IFN-γ-inducing factor. Therefore, the

absence of IL-18R upregulation in CD4$^+$ T cells of $Ifng^{-/-}$ mice would prevent the

expansion of both CD4$^+$ T cell subpopulations. Thus, in $T.$ $cruzi$ infection model, IL-12,

IFN-γ and IL-18R/MyD88 signaling act as key pathways at early check-points in the

generation of CD4$^+$GzB$^+$ T cells with cytotoxic potential, suggesting that these cells

might share common early differentiating steps with the Th1 subset. It is not excluded,

however, that different precursors of CD4CTL and Th1 cells might exist (Takeuchi &

Saito, 2017). A cartoon summarizing our results and cited works published by others is

shown in Figure 7-figure supplement 3. Of note, in humans, the higher production of

IL-18 due to gene polymorphisms has been correlated to protection against chronic

Chagas cardiomyopathy (Leon Rodriguez et al., 2016; Nogueira et al., 2015).

It is possible that differentiation into CD4CTLs could also be influenced by

pathogen-derived factors. Very recently, Krueger *et al*. have shown that a subset of IL-

12-dependent Th1-derived cytotoxic cells was present in the spleen of *Il18$^{-/-}$* mice at

later stages of infection with *Salmonella enterica* (Krueger et al., 2021). Whether that

result was due to compensatory mechanisms, such as the known exacerbated IL-12

production by infected *Il18$^{-/-}$* mice (Monteforte et al., 2000), or to distinct innate

responses, differentially induced by phagosomal bacteria and cytoplasmatic parasites,

remains to be determined. Of note, other discrepancies between results obtained with

Il18$^{-/-}$ versus *Il18ra$^{-/-}$* mouse lineages have been described in different models in the past

(Gutcher et al., 2006; Lewis & Dinarello, 2006). It is known that different classes of

pathogens can trigger diverse APC responses and, therefore, elicit pathogen-tailored Th

programs (Kara et al., 2014; Kiner et al., 2021). Previous works have reported the

immunoregulatory action of *T. cruzi*-derived PAMPs on CD4$^+$ T cell activation, either

by directly acting as co-stimulatory signals for T cells, or by modulating APC functions

(Bellio et al., 1999; DosReis et al., 2002; Medeiros et al., 2007). It would be interesting

to test *T. cruzi*-derived molecules on their capacity to favor the generation of CD4CTLs.

In summary, the present work is the first to described a CD4CTL subset abundantly

present in the *T. cruzi*-infected cardiac tissue, bringing novel pieces to the scenario of

the immune response against a major human intracellular parasite. Importantly, our data

disclose a mechanistic framework involved in the generation of a robust CD4CTL

response, whose control and optimization may provide new therapeutic and vaccination

strategies to combat infection with intracellular pathogens and chronic diseases

resulting thereof.

Materials and Methods

Mice. C57BL/6, B6.SJL and F1 (B6 x B6.SJL) mice were from the Universidade

Federal Fluminense. $Myd88^{-/-}$ mice were generated by Dr. S. Akira (Osaka University,

Japan); $Il12p40^{-/-}$ and $Ifng^{-/-}$ mice were purchased from USP; $Il18ra^{-/-}$ and $Il1r1^{-/-}$ mice

from JAX Mice, USA. GzB reporter mice, GzmbCreER^{T2}/ROSA26EYFP, previously

described (Bannard et al., 2009) were obtained from Dr. Ricardo Gazinelli (UFMG, BH,

Brazil). Mice were on the C57BL/6 background, except GzmbCreER^{T2}/ROSA26EYFP,

which were on mixed Sv129/B6 background and backcrossed to B6 for 5 generations.

In each experiment, only male mice of the same age (6- to 8-week-old) were used. Mice

were housed at a maximum of 5 animals per cage, in forced-air-ventilated micro-

isolation rack system, in a room at 21–23°C under a 12/12 h light/dark cycle and

provided sterilized water and chow ad libitum. All mouse experiments were conducted

in accordance with guidelines of the Animal Care and Use Committee of the Federal

University of Rio de Janeiro (Comitê de Ética do Centro de Ciências da Saúde CEUA -

CCS/UFRJ, license: IMPPG022).

Experimental infection. Bloodstream trypomastigotes of the Y strain of *T. cruzi* (2 x

10^3 cells/0.2 mL) were inoculated intraperitoneally (i.p.). Y-OVA strain trypomastigotes

(Gomes-Neto et al., 2018) were obtained from LLC-MK2 infected cultures and 2 x 10^6

trypomastigotes (in 0.2 mL PBS) were inoculated i.p. Rare outliers (mice in which

the infection was not properly established) were identified by the lower size of the

spleen (total cell number) and lower percentage of GzB$^+$CD8$^+$ T cells (inferior of 50%)

in the spleen. These outlier mice were excluded from all analyses.

Generation of BM chimeras. Acidic drinking water (pH 2.5-3.0) was given for 8 days

to recipient mice prior to irradiation with a lethal single dose of 800 rad (TH780C -

Theratronics, Canada). The day after irradiation, BM cells were transferred and mice

were treated with neomycin sulfate (2 mg/mL) in the drinking water for 15 days.

Marrow was harvested from the femur and tibia of B6.SJL (CD45.1$^+$) and B6, *Myd88$^{-/-}$*,

Il1r1$^{-/-}$ or *Il18ra$^{-/-}$* (CD45.2$^+$) mice. For mixed BM chimeras, CD45.1$^+$ and CD45.2$^+$ BM

cells were transferred in a 1:1 ratio (2 x 10^6 cells each) to irradiated mice. In all

experiments, mice were infected 6 to 8 weeks after cell transfer, when reconstitution

was achieved.

Human population. A total of thirteen volunteers were enrolled in this study. Nine

volunteer patients had positive specific serology for *T. cruzi* and displayed the chronic

Chagas cardiomyopathy (CC) with right and/or left ventricular dilation, global left

ventricular dysfunction, alterations in the cardiac electric impulse generation and

conduction upon electrocardiogram, chest x-rays and echocardiography; four

individuals displayed negative specific serological tests for Chagas disease and were

included as control group (healthy donors, HD). Patients were from Chagas disease

endemic regions within Minas Gerais, Brazil, and were evaluated at the outpatient clinic

of the Universidade Federal de Minas Gerais. Healthy donors were from Belo

Horizonte, which is a non-endemic region for Chagas disease. We excluded from our

study individuals with any other chronic inflammatory diseases, diabetes,

heart/circulatory illnesses or bacterial infections. All individuals included in this work

were volunteers and treatment and clinical care were offered to all patients, as

needed, despite their enrollment in this research project. All participants provided

written informed consent for participation in the study. This cross-sectional study is part

of an extended project evaluating biomarkers of cardiomyopathy development in

Chagas disease, which has the approval of the National Committee of Ethics in

Research (CONEP#195/2007) and are in accordance with the Declaration for Helsinki.

PBMC flow cytometry. Peripheral blood samples were collected in heparin tubes.

PBMCs were obtained by separating whole blood over Ficoll (Sigma-Aldrich, St.

Louis, MO), as previously described (Menezes et al., 2012). Cells were washed and

resuspended in RPMI 1640 medium (Gibco, Rockville, MD) supplemented with 5%

heat-inactivated AB human serum, antibiotics (penicillin, 200 U/ml; and streptomycin,

0.1 mg/mL) and L-glutamine (1 mM) (Sigma-Aldrich) at a concentration of 1×10^7

cells/mL. Cells from each volunteer were cultured on 96-well plates (Costar, Corning

Incorporated, Corning, NY) in 200 μL of medium alone (ctr) or containing 20 μg/mL of

trypomastigote antigen (trypo) for 18h. Trypo was obtained from trypomastigotes

purified from VERO cell cultures: parasites were washed, freeze-thaw for 3 cycles,

followed by 5 cycles of 5 minutes each of sonication using 30 seconds between them.

Samples were ultracentrifugated at 5,000g for 60 min, at 4^oC. Supernatants were

collected, dialyzed for 24h and protein quantification performed by Bradford method.

Cells (2×10^5) were incubated with PerCP-anti-CD4 (clone A161A1) antibody

(Biolegend) for 15 min at 4°C, washed in phosphate-buffered saline (PBS) containing

1% bovine serum albumin (BSA), and fixed by a 20 min incubation with a 2%

formaldehyde. After washing with PBS, cells were permeabilized with 0.5% saponin

for 15 min and intracellularly stained with PE-anti-perforin (clone B-D48) antibody

(Biolegend) for 20 minutes, washed twice with 0.5% saponin solution, resuspended in

PBS and analyzed. A minimum of 100,000 gated events from each sample were

collected and analyzed using FlowJo software.

Flow cytometry of mouse cells. Reagents and antibodies: PB-anti-CD45.1, APC-Cy7-

anti-CD45.2, FITC- PE- or BV-605-anti-CD4, PE-anti-CD8, PE-Cy7-anti-NK1.1,

BV421-anti-TCRβ, PE-anti-2B4, FITC-anti-CD107a, PE-Cy7-anti-CD39, PE-anti-

FasL, PE-Cy7-anti-CRTAM, PE-anti-NKG2D, PE-anti-NKG2A/C/E, FITC- or

AlexaFluor647-anti-GzB, PE-anti-Eomes, PE- or A647-anti-T-bet, PE-anti-FoxP3,

AlexaFluor647-anti-Blimp1 and PE-anti-PRF were used. All mAbs were purchased

from Biolegend, BD Biosciences or eBioscience/Thermo Fisher Scientific. Spleen cells

were stained after red blood cell lysis with ACK buffer. Hearts were perfused with 20

mL of cold PBS, minced into small fragments and treated with collagenase D (1.0

mg/mL) and DNAse I (0.1 mg/mL), for 1 hour at 37ºC. Collagenase D, DNAse I,

monensin and saponin were purchased from Sigma-Aldrich. Single cell suspensions

were incubated with anti-CD16/CD32 (FcR block) for 5 min and then surface stained

for 30 min on ice. For IL-10 and IFN-γ intracellular staining, 2×10^{6} spleen cells were

cultured in the presence of PMA (16 nM), ionomycin (0.5 μM), monensin (5 μM) and

brefeldin A (3.0 μg/mL) for 4 hours. After staining for surface markers, cells were fixed

with paraformaldehyde 1% for 1 h, permeabilized with saponin 0.2% for 20 min and

stained with PE-Cy7-anti-IFN-γ mAb and/or PE-anti-IL-10 mAbs (Biolegend). For TF

and GzB/PRF staining, the FoxP3 staining buffer kit (eBioscience, ThermoFisher

Scientific) was employed. At least 20,000 events gated on $CD4^{+}$ T cells were acquired.

Analytical flow cytometry was conducted with MoFlo (Beckman Coulter/Dako-

Cytomation), FacsCanto, FacsCalibur or BD LSRFortessa (BD Bioscience) and data

analyzed with Summit V4.3 software (Dako Colorado, Inc).

***In vitro* and *in vivo* cytotoxicity assays.** For redirected cytotoxic assay, sorted B cells

(B220[+]), previously activated by LPS (InvivoGen) for 48 h and then incubated for 1 h

with anti-CD3ε antibody (10 µg/mL), were employed as target cells. FACS-sorted naïve

(CD44[low]CD4[+]) and effector/memory (CD44[hi]CD4[+]) T cells were purified from the

spleen of non-infected and infected mice, respectively. Sorted T cells were coincubated

at different E:T ratios, with 10^4 target B cells in round-bottom microplates (Costar-

Corning) for 14 h, at 37ºC in a 5% CO_2 incubator and in triplicates. After incubation,

living target cells were quantified by flow cytometry, after staining with propidium

iodide at 0.5 µg/ml (Sigma-Aldrich). For *T. cruzi*-specific cytotoxicity assays, CD4[+] T

cells were obtained from infected B6 mice, at day 14 pi, and purified by FACS sorting

(MoFlo cell sorter, Beckman Coulter/Dako-Cytomation). IC-21 macrophage cell

lineage (ATCC) were used as targets *in vitro.* For this, IC-21 cells were counted and

split into two tubes: half of the target cells were incubated for 45 min at 37º C in 5%

CO_2, in RPMI containing *T. cruzi* amastigote protein extract (30 µg/mL) obtained as

previously described (Oliveira et al., 2010). Ag-loaded cells were then stained with 2.5

µM CFSE (CFSE[Hi]) (Invitrogen, Life Technologies), and control cells with 0.5 µM

CFSE (CFSE[Low]) in 5 mL of warm PBS for 10 min at 37º C. After two washes in RPMI

1640 (10% FCS), cells were resuspended at a concentration of 2 x 10^5 cells/mL each

and mixed at 1:1 ratio (CFSE[Hi]:CFSE[Low]). Purified CD4[+] T cells were added at different

E:T ratios. Cells were seeded in duplicate to round-bottom 96-well plate in a final

volume of 200 µL. After ON incubation, cells were harvested and incubated with 0.5

µg/mL of propidium iodide and PI[+] cells were excluded from FACS analysis. The

percentage of specific lysis was calculated using the formula: % of specific lysis = 100x

[(CFSE[Low] / CFSE[Hi]) ctr − (CFSE[Low] / CFSE[Hi]) effector] x (CFSE[Low] / CFSE[Hi]) ctr[−1].

For *in vivo* cytotoxicity assays, splenocytes from non-infected B6 mice were

employed as target cells after being divided in 3 subpopulations and stained with 10.0

(High), 2.5 (Int) and 0.625 (Low) μM of CFSE or of CTV (Invitrogen, Life

Technologies). $CFSE^{Int}$ and $CFSE^{Low}$ cells were loaded with one of the following

peptides: SA85c (SHNFTLVASVIIEEA), SA85d (LVASVIIEEAPSGNT), $OVA_{323-339}$,

$OVA_{257-263}$ or PA8 (VNHRFTLV) at 2 μM, all from GenScript, EUA. $CFSE^{Hi}$ cells

were not loaded with any peptide and used as internal control. For calculating the

percentage of specific lysis, $CFSE^{Hi}$ and $CFSE^{Int}$ (or $CFSE^{Low}$) were gated together in

order to constitute 100% of cells. $CFSE^{Hi}$ and $CFSE^{Int}$ (or $CFSE^{Low}$) populations were

then individually gated and specific lysis was calculated using the following formula: %

of specific lysis = $100x$ $[(CFSE^{Low \ or \ Int} / CFSE^{Hi})$ ctr $- (CFSE^{Low \ or \ Int} / CFSE^{Hi})$

effector] x $(CFSE^{Low \ or \ Int} / CFSE^{Hi})$ ctr^{-1}.

Real time RT-PCR. RNA was extracted using the RNeasykit (QIAGEN). cDNA was

synthesized using the High Capacity cDNA Reverse Transcription Kit (Invitrogen, Life

Technologies), following the manufacturer's instructions. Gene expression was

analyzed using primers in combination with SYBR Green® on a StepOneReal Time

PCR System (Applied Biosystems, Life Technologies). The expression level of the

genes of interest was determined relative to the expression of 18S rRNA. Primers:

Runx3 Fwd.: 5'- ACA GCA TCT TTG ACT CCT TCC -3'; *Runx3* Rev.: 5- GTT GGT

GAA CAC GGT GAT TG- 3'; *Thpok* Fwd: 5'- ATG GGA TTC AAA TCA GGT CA-

3'; *Thpok* Rev.5'- TTCTTC CTA CAC CCT GTG CC -3'; *CD8a* Fwd: 5' - ACT GCA

AGG AAG CAA GTG GT – 3'; *CD8a* Rev: 5'-CAC CGC TAA AGG CAG TTC TC-

3'; 18S-Fwd.: 5'- GGG CGT GGG GCG GAG ATA TGC - 3' and 18S-Rev.: 5'- CG

CGG ACA CGA AGG CCC CAA A -3'. For parasite load quantification, hearts of

infected mice were perfused, minced and immediately homogenized in 1.0 ml of 4.0 M

Guanidine thiocyanate (SIGMA-Aldrich) containing 8.0 μl/mL of β-mercaptoethanol

and processed for DNA extraction. Generation of PCR standards and detection of

parasite tissue load by real-time PCR was carried out as described (Oliveira et al.,

2017); briefly, primers amplify a repeated satellite sequence of *T. cruzi* DNA of 195

base-pairs: TCZ-Fwd.: (5'-GCTCTTGCCCACAAGGGTGC-3') and TCZ-Rev.: (5'-

CCAAGCAGCGGATAGTTCAGG-3'). Reactions with TNF-a-Fwd: (5'-

CCTGGAGGAGAAGAGGAAAGAGA-3') and TNF-a-Rev.: (5'-

TTGAGGACCTCTGTGTATTTGTCAA-3') primers for *Mus musculus* TNF-α gene

were used as loading controls.

Gene-expression analysis. Gene expression data were extracted with GeneSpring

software GX 11.0 software using default parameters. Data were then normalized by

subtracting the average from all genes from each gene intensity and dividing the result

by the standard deviation of all gene intensities (i.e. z-score normalization). Z ratios

were calculated as previously described (Cheadle et al., 2003) and used to compare the

difference in expression between WT samples and *Myd88$^{-/-}$* samples. Gene Set

Enrichment Analysis (GSEA) (http://www.broadinstitute.org/gsea/index.jsp) was

performed on the genes pre-ranked by Z ratios (1,000 permutations and Weighted

Enrichment Statistic) using as data set 82 differentially expressed genes of sorted WT

and *Myd88$^{-/-}$*CD4$^+$ T cells (GEO accession no. GSE57738) (Oliveira et al., 2017); and

using as gene set the CD4$^+$GzmB$^+$ (CD4CTL) transcriptional signature (159 upregulated

genes) previously described (https://www.ebi.ac.uk/ena/data/view/PRJEB14043).

Adoptive CD4$^+$ T cell transfer assay. Eight-week-old male

GzmbCreER^{T2}/ROSA26EYFP mice were infected with 2 x 10^3 trypomastigotes and

treated with 4 mg/animal/day of tamoxifen citrate (Sandoz-Novartis) from day 7 to 11

pi by gavage. Spleen cells were harvested and treated with ACK for red blood cells lysis

at day 13 pi. Alternatively, CD4$^+$ T cells were purified from the spleen of non-treated

and infected *Ifng^{-/-}* mice. In both cases, spleen cells were labeled with anti-B220, anti-

CD11b, anti-NK1.1, anti-γδTCR, anti-CD8 and biotinylated PE-conjugated anti-CD4

antibodies for 30 minutes on ice, followed by staining with PE-Cy7-conjugated

Streptavidin for 20 min. Residual PE-Cy7$^+$ cells were excluded and EYFP$^+$PE$^+$

(CD4$^+$GzB$^+$ T cells) or PE$^+$CD4$^+$ (*Ifng^{-/-}*CD4$^+$ T cells) were sorted using MoFlo cell

sorter (Beckman Coulter/Dako-Cytomation). CD4$^+$GzB$^+$ T cells (2.8 x 10^5 cells/animal)

or *Ifng^{-/-}* CD4$^+$ T cells (1.3 x10^6 cells/animal), both presenting purity higher than 98%,

were adoptively transferred by *iv* injection into *Il18ra^{-/-}* recipient mice, previously

infected with 1 x 10^3 trypomastigotes (day 7 pi). Control groups of *Il18ra^{-/-}* and *Myd88^{-/-}*

infected mice received PBS. Survival was accompanied daily.

Statistics. Statistical analyses were performed using GraphPad Prism version 6 for Mac

OS (GraphPad Software, San Diego California USA, www.graphpad.com). Sample

sizes were chosen to allow the detection of statistical differences with $p < 0.5$ using

Student's *t* test, or the Log-rank (Mantel-Cox) test for mouse survival curves.

Correlation analysis employed Pearson correlation test. When comparing two groups,

the minimum number was n=3 per group. In some experiments, the number was

increased to n= 9 to permit $p < 0.05$ even considering the possibility of a single outlier

in a population. Data are expressed as mean $\pm$ SEM and considered statistically

significant if *p* values were < 0.05.

Acknowledgments:

General: We would like to thank Dr. Ricardo T. Gazzinelli (Fiocruz, MG, Brazil) for

enriching discussion and providing GzmbCreER^{T2}/ROSA26EYFP mouse lineage and

Dr. Helder I. Nakaya (USP, SP, Brazil) for helping with Bioinformatics Analyses.

Funding. This study was financed in part by the Coordenação de Aperfeiçoamento de

Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001; Fundação Carlos

Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ); Conselho

Nacional de Pesquisas (CNPq) and by The National Institute for Vaccine Development

and Technology (INCTV-CNPq) and National Institutes of Health (NIH). CDB and

LMB received PhD fellowships from CNPq; AG and EGAN received PhD fellowships

from CAPES; FBC and AG received PDJ-CNPq fellowships. GM received PIBIC-

CNPq fellowship. AN, WOD and MB received PQ fellowship from CNPq; AN and MB

received CNE grant from FAPERJ.

The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

Author contributions: ACO, FBC, WOD, AN and MB conceived and designed the

experiments. CDB, AG, FBC, LMB, JRL, AG, EGAN ACO, AN and MB performed

the experiments. CDB, JFG, FBC, LMB, GAM, ACO, EGAN, WOD, AN and MB

analyzed the data. ACO, WOD, AN and MB contributed parasites/ reagents/

mice/materials/analysis tools. MB wrote the paper.

Competing interests: The authors declare no competing financial interests.

References:

Afonina, I. S., Cullen, S. P., & Martin, S. J. (2010). Cytotoxic and non-cytotoxic roles

of the CTL/NK protease granzyme B. *Immunol Rev, 235*(1), 105-116. doi:

10.1111/j.0105-2896.2010.00908.x

Anderson, A. C., Joller, N., & Kuchroo, V. K. (2016). Lag-3, Tim-3, and TIGIT: Co-

inhibitory Receptors with Specialized Functions in Immune Regulation. *Immunity,*

44(5), 989-1004. doi: 10.1016/j.immuni.2016.05.001

Anderson, C. F., Oukka, M., Kuchroo, V. J., & Sacks, D. (2007). CD4(+)CD25(-

)Foxp3(-) Th1 cells are the source of IL-10-mediated immune suppression in chronic

cutaneous leishmaniasis. *J Exp Med., 204*(2), 285-297. Epub 2007

doi:10.1084/jem.20061886.

Ayo, C. M., Oliveira, A. P., Camargo, A. V., Mattos, C. C., Bestetti, R. B., & Mattos, L.

C. (2015). Association of the Functional MICA-129 Polymorphism With the Severity

of Chronic Chagas Heart Disease. *Clin Infect Dis, 61*(8), 1310-1313. doi:

10.1093/cid/civ540

Bannard, O., Kraman, M., & Fearon, D. T. (2009). Secondary replicative function of

CD8+ T cells that had developed an effector phenotype. *Science, 323*(5913), 505-509.

Baumann, C., Bonilla, W. V., Fröhlich, A., Helmstetter, C., Peine, M., Hegazy, A. N., .

. . Löhning, M. (2015). T-bet- and STAT4-dependent IL-33 receptor expression

directly promotes antiviral Th1 cell responses. *Proc Natl Acad Sci U S A, 112*(13),

4056-4061. doi: 10.1073/pnas.1418549112

Bellio, M., Oliveira, A. C., Mermelstein, C. S., Capella, M. A., Viola, J. P., Levraud, J.

P., . . . Mendonca-Previato, L. (1999). Costimulatory action of glycoinositol-

phospholipids from Trypanosoma cruzi: increased interleukin 2 secretion and

induction of nuclear translocation of the nuclear factor of activated T cells 1. *FASEB*

J., 13(12), 1627-1636. doi: 10.1096/fasebj.13.12.1627.

Ben-Sasson, S. Z., Hu-Li, J., Quiel, J., Cauchetaux, S., Ratner, M., Shapira, I., . . . Paul,

W. E. (2009). IL-1 acts directly on CD4 T cells to enhance their antigen-driven

expansion and differentiation. *Proc Natl Acad Sci U S A, 106*(17), 7119-7124. doi:

10.1073/pnas.0902745106.

Bern, Caryn. (2015). Chagas' Disease. *New England Journal of Medicine, 373*(5), 456-

466. doi: 10.1056/NEJMra1410150

Bossi, G., & Griffiths, G. M. (1999). Degranulation plays an essential part in regulating

cell surface expression of Fas ligand in T cells and natural killer cells. *Nat Med, 5*(1),

90-96. doi: 10.1038/4779

Brien, J. D., Uhrlaub, J. L., & Nikolich-Zugich, J. (2008). West Nile virus-specific CD4

T cells exhibit direct antiviral cytokine secretion and cytotoxicity and are sufficient

for antiviral protection. *J Immunol, 181*(12), 8568-8575. doi:

10.4049/jimmunol.181.12.8568.

Brown, D. M., Kamperschroer, C., Dilzer, A. M., Roberts, D. M., & Swain, S. L.

(2009). IL-2 and antigen dose differentially regulate perforin- and FasL-mediated

cytolytic activity in antigen specific CD4+ T cells. *Cell Immunol, 257*(1-2), 69-79.

doi: 10.1016/j.cellimm.2009.03.002

Brown, D. M., Lee, S., Garcia-Hernandez Mde, L., & Swain, S. L. (2012).

Multifunctional CD4 cells expressing gamma interferon and perforin mediate

protection against lethal influenza virus infection. *J Virol, 86*(12), 6792-6803. doi:

10.1128/jvi.07172-11

Burel, J. G., Apte, S. H., Groves, P. L., Klein, K., McCarthy, J. S., & Doolan, D. L.

(2016). Reduced Plasmodium Parasite Burden Associates with CD38+ CD4+ T

Cells Displaying Cytolytic Potential and Impaired IFN-gamma Production. *PLoS*

Pathog, 12(9), e1005839. doi: 10.1371/journal.ppat.1005839

Campos, M. A., Closel, M., Valente, E. P., Cardoso, J. E., Akira, S., Alvarez-Leite, J. I.,

. . . Gazzinelli, R. T. (2004). Impaired production of proinflammatory cytokines and

host resistance to acute infection with Trypanosoma cruzi in mice lacking functional

myeloid differentiation factor 88. *J Immunol, 172*(3), 1711-1718. doi:

10.4049/jimmunol.172.3.1711.

Canaday, D. H., Wilkinson, R. J., Li, Q., Harding, C. V., Silver, R. F., & Boom, W. H.

(2001). CD4(+) and CD8(+) T cells kill intracellular Mycobacterium tuberculosis by a

perforin and Fas/Fas ligand-independent mechanism. *J Immunol, 167*(5), 2734-2742.

doi: 10.4049/jimmunol.167.5.2734.

Cheadle, C., Vawter, M. P., Freed, W. J., & Becker, K. G. (2003). Analysis of

microarray data using Z score transformation. *J Mol Diagn, 5*(2), 73-81. doi:

10.1016/S1525-1578(10)60455-2.

Cheroutre, H., & Husain, M. M. (2013). CD4 CTL: living up to the challenge. *Semin*

Immunol, 25(4), 273-281. doi: 10.1016/j.smim.2013.10.022.

Cooper, D., Pride, M. W., Guo, M., Cutler, M., Mester, J. C., Nasar, F., . . . Natuk, R. J.

(2004). Interleukin-12 redirects murine immune responses to soluble or aluminum

phosphate adsorbed HSV-2 glycoprotein D towards Th1 and CD4+ CTL responses.

Vaccine, 23(2), 236-246. doi: 10.1016/j.vaccine.2004.05.011

Cordeiro, Z. M., Dahia, A. C., & Andrade, Z. A. (1997). Kinetics of Trypanosoma cruzi

destruction in the mouse spleen. *Rev Soc Bras Med Trop, 30*(1), 3-9. doi:

10.1590/s0037-86821997000100002

Cruz-Guilloty, F., Pipkin, M. E., Djuretic, I. M., Levanon, D., Lotem, J., Lichtenheld,

M. G., . . . Rao, A. (2009). Runx3 and T-box proteins cooperate to establish the

transcriptional program of effector CTLs. *J Exp Med, 206*(1), 51-59. doi:

10.1084/jem.20081242

Curran, M. A., Geiger, T. L., Montalvo, W., Kim, M., Reiner, S. L., Al-Shamkhani, A.,

. . . Allison, J. P. (2013). Systemic 4-1BB activation induces a novel T cell phenotype

driven by high expression of Eomesodermin. *J Exp Med, 210*(4), 743-755. doi:

10.1084/jem.20121190

Djuretic, I. M., Levanon, D., Negreanu, V., Groner, Y., Rao, A., & Ansel, K. M. (2007).

Transcription factors T-bet and Runx3 cooperate to activate Ifng and silence Il4 in T

helper type 1 cells. *Nat Immunol, 8*(2), 145-153. doi: 10.1038/ni1424.

Donnarumma, T., Young, G. R., Merkenschlager, J., Eksmond, U., Bongard, N., Nutt,

S. L., . . . Kassiotis, G. (2016). Opposing Development of Cytotoxic and Follicular

Helper CD4 T Cells Controlled by the TCF-1-Bcl6 Nexus. *Cell Rep, 17*(6), 1571-

1583. doi: 10.1016/j.celrep.2016.10.013

DosReis, G. A., Pecanha, L. M., Bellio, M., Previato, J. O., & Mendonca-Previato, L.

(2002). Glycoinositol phospholipids from Trypanosoma cruzi transmit signals to the

cells of the host immune system through both ceramide and glycan chains. *Microbes*

Infect., 4(9), 1007-1013. doi: 10.1016/s1286-4579(02)01616-7.

Egawa, T., Tillman, R. E., Naoe, Y., Taniuchi, I., & Littman, D. R. (2007). The role of

the Runx transcription factors in thymocyte differentiation and in homeostasis of

naive T cells. *J Exp Med, 204*(8), 1945-1957. doi: 10.1084/jem.20070133.

Eshima, K., Chiba, S., Suzuki, H., Kokubo, K., Kobayashi, H., Iizuka, M., . . .

Shinohara, N. (2012). Ectopic expression of a T-box transcription factor,

eomesodermin, renders CD4(+) Th cells cytotoxic by activating both perforin- and

FasL-pathways. *Immunol Lett, 144*(1-2), 7-15. doi: 10.1016/j.imlet.2012.02.013

Fang, M., Siciliano, N. A., Hersperger, A. R., Roscoe, F., Hu, A., Ma, X., . . . Sigal,

L. J. (2012). Perforin-dependent CD4+ T-cell cytotoxicity contributes to control a

murine poxvirus infection. *Proc Natl Acad Sci U S A, 109*(25), 9983-9988. doi:

10.1073/pnas.1202143109

Ferreira, L. R., Ferreira, F. M., Nakaya, H. I., Deng, X., Cândido, D. D., de Oliveira, L.

C., . . . Cunha-Neto, E. (2017). Blood Gene Signatures of Chagas Cardiomyopathy

With or Without Ventricular Dysfunction. *J Infect Dis, 215*(3), 387-395. doi:

10.1093/infdis/jiw540

Frazer, L. C., Sullivan, J. E., Zurenski, M. A., Mintus, M., Tomasak, T. E., Prantner, D.,

. . . Darville, T. (2013). CD4+ T cell expression of MyD88 is essential for normal

resolution of Chlamydia muridarum genital tract infection. *J Immunol, 191*(8), 4269-

4279.

Gagliani, N., Magnani, C. F., Huber, S., Gianolini, M. E., Pala, M., Licona-Limon, P., .

. . Roncarolo, M. G. (2013). Coexpression of CD49b and LAG-3 identifies human and

mouse T regulatory type 1 cells. *Nat Med, 19*(6), 739-746. doi: 10.1038/nm.3179

Glimcher, L. H., Townsend, M. J., Sullivan, B. M., & Lord, G. M. (2004). Recent

developments in the transcriptional regulation of cytolytic effector cells. *Nat Rev*

Immunol, 4(11), 900-911. doi: 10.1038/nri1490

Gomes-Neto, J. F., Sartorius, R., Canto, F. B., Almeida, T. S., Dias, A. A., Barbosa, C.

D., . . . Bellio, M. (2018). Vaccination With Recombinant Filamentous fd Phages

Against Parasite Infection Requires TLR9 Expression. *Front Immunol, 9*, 1173. doi:

10.3389/fimmu.2018.01173

Gunturi, A., Berg, R. E., & Forman, J. (2004). The role of CD94/NKG2 in innate and

adaptive immunity. *Immunol Res, 30*(1), 29-34. doi: 10.1385/ir:30:1:029

Gutcher, I., Urich, E., Wolter, K., Prinz, M., & Becher, B. (2006). Interleukin 18-

independent engagement of interleukin 18 receptor-alpha is required for

autoimmune inflammation. *Nat Immunol, 7*(9), 946-953. doi: 10.1038/ni1377.

Haring, J. S., & Harty, J. T. (2009). Interleukin-18-related genes are induced during the

contraction phase but do not play major roles in regulating the dynamics or function

of the T-cell response to Listeria monocytogenes infection. *Infect Immun, 77*(5),

1894-1903. doi: 10.1128/IAI.01315-08.

Hua, L., Yao, S., Pham, D., Jiang, L., Wright, J., Sawant, D., . . . Sun, J. (2013).

Cytokine-dependent induction of CD4+ T cells with cytotoxic potential during

influenza virus infection. *J Virol, 87*(21), 11884-11893. doi: 10.1128/jvi.01461-13

Intlekofer, A. M., Takemoto, N., Wherry, E. J., Longworth, S. A., Northrup, J. T.,

Palanivel, V. R., . . . Reiner, S. L. (2005). Effector and memory CD8+ T cell fate

coupled by T-bet and eomesodermin. *Nat Immunol, 6*(12), 1236-1244. doi:

10.1038/ni1268

Jankovic, Dragana, Kullberg, Marika C., Feng, Carl G., Goldszmid, Romina S.,

Collazo, Carmen M., Wilson, Mark, . . . Sher, Alan. (2007). Conventional T-

bet+FoxP3- Th1 cells are the major source of host-protective regulatory IL-10 during

intracellular protozoan infection. *J Exp Med, 204*(2), 273. doi: 10.1084/jem.20062175

Jellison, E. R., Kim, S. K., & Welsh, R. M. (2005). Cutting edge: MHC class II-

restricted killing in vivo during viral infection. *J Immunol, 174*(2), 614-618. doi:

10.4049/jimmunol.174.2.614.

Joffre, O., Nolte, M. A., Spörri, R., & Reis e Sousa, C. (2009). Inflammatory signals in

dendritic cell activation and the induction of adaptive immunity. *Immunol Rev,*

227(1), 234-247. doi: 10.1111/j.1600-065X.2008.00718.x

Juno, J. A., van Bockel, D., Kent, S. J., Kelleher, A. D., Zaunders, J. J., & Munier, C.

M. (2017). Cytotoxic CD4 T Cells-Friend or Foe during Viral Infection? *Front*

Immunol, 8, 19. doi: 10.3389/fimmu.2017.00019

Kara, E. E., Comerford, I., Fenix, K. A., Bastow, C. R., Gregor, C. E., McKenzie, D. R.,

& McColl, S. R. (2014). Tailored immune responses: novel effector helper T cell

subsets in protective immunity. *PLoS Pathog, 10*(2), e1003905. doi:

10.1371/journal.ppat.1003905

Keesen, T.S.L., Gomes, J.A.S, Fares, R.C.G., de Araújo, F.F, Ferreira, K.S, Chaves,

A.T, Rocha, M.O.C, Correa-Oliveira, R. (2012). Characterization of CD4+ cytotoxic

lymphocytes and apoptosis markers induced by *Trypanossoma cruzi* infection. *Scand*

J Immunol, 76(3), 311-319. doi: 10.1111/j.1365-3083.2012.02730.x.

Kiner, E., Willie, E., Vijaykumar, B., Chowdhary, K., Schmutz, H., Chandler, J., . . .

Benoist, C. (2021). Gut CD4(+) T cell phenotypes are a continuum molded by

microbes, not by T(H) archetypes. *Nat Immunol, 22*(2), 216-228. doi:

10.1038/s41590-020-00836-7

Kotov, D. I., Kotov, J. A., Goldberg, M. F., & Jenkins, M. K. (2018). Many Th Cell

Subsets Have Fas Ligand-Dependent Cytotoxic Potential. *J Immunol, 200*(6), 2004-

2012. doi: 10.4049/jimmunol.1700420

Krueger, P. D., Goldberg, M. F., Hong, S. W., Osum, K. C., Langlois, R. A., Kotov, D.

I., . . . Jenkins, M. K. (2021). Two sequential activation modules control the

differentiation of protective T helper-1 (Th1) cells. *Immunity, 54*(4), 687-701 e684.

doi: 10.1016/j.immuni.2021.03.006

LaRosa, D. F., Stumhofer, J. S., Gelman, A. E., Rahman, A. H., Taylor, D. K., Hunter,

C. A., & Turka, L. A. (2008). T cell expression of MyD88 is required for resistance to

Toxoplasma gondii. *Proc Natl Acad Sci U S A, 105*(10), 3855-3860. doi:

10.1073/pnas.0706663105.

Leon Rodriguez, D.A., Carmona, F. D., Echeverría, L. E., González, C. I., & Martin, J.

(2016). IL18 Gene Variants Influence the Susceptibility to Chagas Disease. *PLOS*

Negl Trop Dis, 10(3), e0004583. doi: 10.1371/journal.pntd.0004583

Lewis, E. C., & Dinarello, C. A. (2006). Responses of IL-18- and IL-18 receptor-

deficient pancreatic islets with convergence of positive and negative signals for the

IL-18 receptor. *Proc Natl Acad Sci U S A, 103*(45), 16852-16857. doi:

10.1073/pnas.0607917103.

Marshall, N. B., Vong, A. M., Devarajan, P., Brauner, M. D., Kuang, Y., Nayar, R., . . .

Swain, S. L. (2017). NKG2C/E Marks the Unique Cytotoxic CD4 T Cell Subset,

ThCTL, Generated by Influenza Infection. *J Immunol, 198*(3), 1142-1155. doi:

10.4049/jimmunol.1601297

Mascanfroni, I. D., Takenaka, M. C., Yeste, A., Patel, B., Wu, Y., Kenison, J. E., . . .

Quintana, F. J. (2015). Metabolic control of type 1 regulatory T cell differentiation by

AHR and HIF1-alpha. *Nat Med, 21*(6), 638-646. doi: 10.1038/nm.3868

Medeiros, M. M., Peixoto, J. R., Oliveira, A. C., Cardilo-Reis, L., Koatz, V. L., Van

Kaer, L., . . . Bellio, M. (2007). Toll-like receptor 4 (TLR4)-dependent

proinflammatory and immunomodulatory properties of the glycoinositolphospholipid

(GIPL) from *Trypanosoma cruzi. J Leukoc Biol, 82*(3), 488-496. doi:

10.1189/jlb.0706478.

Menezes, C. A., Sullivan, A. K., Falta, M. T., Mack, D. G., Freed, B. M., Rocha, M. O.,

. . . Dutra, W. O. (2012). Highly conserved CDR3 region in circulating

CD4(+)Vbeta5(+) T cells may be associated with cytotoxic activity in Chagas disease.

Clin Exp Immunol, 169(2), 109-118. doi: 10.1111/j.1365-2249.2012.04608.x.

Millar, A. E., & Kahn, S. J. (2000). The SA85-1.1 protein of the Trypanosoma cruzi

trans-sialidase superfamily is a dominant T-cell antigen. *Infect Immun, 68*(6), 3574-

3580. doi: 10.1128/IAI.68.6.3574-3580.2000.

Mittal, P., Abblett, R., Ryan, J. M., Hagymasi, A. T., Agyekum-Yamoah, A.,

Svedova, J., . . . Adler, A. J. (2018). An Immunotherapeutic CD137 Agonist Releases

Eomesodermin from ThPOK Repression in CD4 T Cells. *J Immunol, 200*(4), 1513-

1526. doi: 10.4049/jimmunol.1701039

Monteforte, Gina M., Takeda, Kiyoshi, Rodriguez-Sosa, Miriam, Akira, Shizuo, David,

John R., & Satoskar, Abhay R. (2000). Genetically Resistant Mice Lacking IL-18

Gene Develop Th1 Response and Control Cutaneous Leishmania major Infection. *J*

Immunol 164(11), 5890-5893. doi: 10.4049/jimmunol.164.11.5890

Mucida, D., Husain, M. M., Muroi, S., van Wijk, F., Shinnakasu, R., Naoe, Y., . . .

Cheroutre, H. (2013). Transcriptional reprogramming of mature CD4(+) helper T

cells generates distinct MHC class II-restricted cytotoxic T lymphocytes. *Nat*

Immunol, 14(3), 281-289. doi: 10.1038/ni.2523.

Nakahira, Masakiyo, Tomura, Michio, Iwasaki, Masayuki, Ahn, Hyun-Jong, Bian,

Yang, Hamaoka, Toshiyuki, . . . Fujiwara, Hiromi. (2001). An Absolute Requirement

for STAT4 and a Role for IFN-γ as an Amplifying Factor in IL-12 Induction of the

Functional IL-18 Receptor Complex. *J Immunol, 167*(3), 1306. doi:

10.4049/jimmunol.167.3.1306

Nogueira, Luciana Gabriel, Frade, Amanda Farage, Ianni, Barbara Maria, Laugier,

Laurie, Pissetti, Cristina Wide, Cabantous, Sandrine, . . . Cunha-Neto, Edecio. (2015).

Functional IL18 polymorphism and susceptibility to Chronic Chagas Disease.

Cytokine, 73(1), 79-83. doi: https://doi.org/10.1016/j.cyto.2015.01.037

Nunes, M. C. P., Beaton, A., Acquatella, H., Bern, C., Bolger, A. F., Echeverría, L. E., .

. . Marin-Neto, J. A. (2018). Chagas Cardiomyopathy: An Update of Current Clinical

Knowledge and Management: A Scientific Statement From the American Heart

Association. *Circulation, 138*(12), e169-e209. doi: 10.1161/cir.0000000000000599

Oliveira, A. C., de Alencar, B. C., Tzelepis, F., Klezewsky, W., da Silva, R. N.,

Neves, F. S., . . . Bellio, M. (2010). Impaired innate immunity in Tlr4(-/-) mice but

preserved CD8+ T cell responses against *Trypanosoma cruzi* in Tlr4-, Tlr2-, Tlr9- or

Myd88-deficient mice. *PLoS Pathog, 6*(4), e1000870. doi:

10.1371/journal.ppat.1000870

Oliveira, A. C., Gomes-Neto, J. F., Barbosa, C. D., Granato, A., Reis, B. S., Santos, B.

M., . . . Bellio, M. (2017). Crucial role for T cell-intrinsic IL-18R-MyD88 signaling in

cognate immune response to intracellular parasite infection. *Elife, 6*. doi:

10.7554/eLife.30883

Park, Young-Jun, Ryu, Heeju, Choi, Garam, Kim, Byung-Seok, Hwang, Eun Sook,

Kim, Hun Sik, & Chung, Yeonseok. (2019). IL-27 confers a protumorigenic activity

of regulatory T cells via CD39. *Proc Natl Acad Sci U S A,* 116(8):3106-3111. doi:

10.1073/pnas.1810254116

Pearce, Erika L., Mullen, Alan C., Martins, Gislâine A., Krawczyk, Connie M.,

Hutchins, Anne S., Zediak, Valerie P., . . . Reiner, Steven L. (2003). Control of

Effector CD8+ T Cell Function by the Transcription Factor Eomesodermin. *Science,*

302(5647), 1041. doi: 10.1126/science.1090148

Poholek, A. C. (2021). Tissue-Specific Contributions to Control of T Cell Immunity.

Immunohorizons, 5(6), 410-423. doi: 10.4049/immunohorizons.2000103

Qui, H. Z., Hagymasi, A. T., Bandyopadhyay, S., St Rose, M. C., Ramanarasimhaiah,

R., Menoret, A., . . . Adler, A. J. (2011). CD134 plus CD137 dual costimulation

induces Eomesodermin in CD4 T cells to program cytotoxic Th1 differentiation. *J*

Immunol, 187(7), 3555-3564. doi: 10.4049/jimmunol.1101244

Reis, B. S., Rogoz, A., Costa-Pinto, F. A., Taniuchi, I., & Mucida, D. (2013). Mutual

expression of the transcription factors Runx3 and ThPOK regulates intestinal CD4(+)

T cell immunity. *Nat Immunol, 14*(3), 271-280. doi: 10.1038/ni.2518.

Reis, D. D., Jones, E. M., Tostes, S., Lopes, E. R., Chapadeiro, E., Gazzinelli, G., . . .

McCurley, T. L. (1993). Expression of major histocompatibility complex antigens and

adhesion molecules in hearts of patients with chronic Chagas' disease. *Am J Trop Med*

Hyg, 49(2), 192-200. doi: 10.4269/ajtmh.1993.49.192.

Robinson, D., Shibuya, K., Mui, A., Zonin, F., Murphy, E., Sana, T., . . . O'Garra, A.

(1997). IGIF does not drive Th1 development but synergizes with IL-12 for

interferon-gamma production and activates IRAK and NFkappaB. *Immunity, 7*(4),

571-581. doi: 10.1016/s1074-7613(00)80378-7

Rodrigues, M. M., Oliveira, A. C., & Bellio, M. (2012). The Immune Response to

Trypanosoma cruzi: Role of Toll-Like Receptors and Perspectives for Vaccine

Development. *J Parasitol Res, 2012*, 507874. doi: 10.1155/2012/507874

Roncarolo, M. G., Gregori, S., Bacchetta, R., Battaglia, M., & Gagliani, N. (2018). The

Biology of T Regulatory Type 1 Cells and Their Therapeutic Application in Immune-

Mediated Diseases. *Immunity, 49*(6), 1004-1019. doi: 10.1016/j.immuni.2018.12.001

Schenten, Dominik, Nish, Simone A, Yu, Shuang, Yan, Xiting, Lee, Heung Kyu,

Brodsky, Igor, . . . Medzhitov, Ruslan. (2014). Signaling through the Adaptor

Molecule MyD88 in CD4+ T Cells Is Required to Overcome Suppression by

Regulatory T Cells. *Immunity, 40*(1), 78-90. doi: 10.1016/j.immuni.2013.10.023.

Serroukh, Y., Gu-Trantien, C., Hooshiar Kashani, B., Defrance, M., Vu Manh, T. P.,

Azouz, A., . . . Marchant, A. (2018). The transcription factors Runx3 and ThPOK

cross-regulate acquisition of cytotoxic function by human Th1 lymphocytes. *Elife, 7*.

doi: 10.7554/eLife.30496

Smeltz, R. B., Chen, J., Hu-Li, J., & Shevach, E. M. (2001). Regulation of interleukin

(IL)-18 receptor alpha chain expression on CD4(+) T cells during T helper (Th)1/Th2

differentiation. Critical downregulatory role of IL-4. *J Exp Med, 194*(2), 143-153.

doi: 10.1084/jem.194.2.143

Stojanovic, A., Correia, M. P., & Cerwenka, A. (2018). The NKG2D/NKG2DL Axis in

the Crosstalk Between Lymphoid and Myeloid Cells in Health and Disease. *Front*

Immunol, 9, 827. doi: 10.3389/fimmu.2018.00827

Tagawa, T., Albanese, M., Bouvet, M., Moosmann, A., Mautner, J., Heissmeyer, V., . . .

Hammerschmidt, W. (2016). Epstein-Barr viral miRNAs inhibit antiviral CD4+ T cell

responses targeting IL-12 and peptide processing. *J Exp Med, 213*(10), 2065-2080.

doi: 10.1084/jem.20160248

Takeda, K., Tsutsui, H., Yoshimoto, T., Adachi, O., Yoshida, N., Kishimoto, T., . . .

Akira, S. (1998). Defective NK cell activity and Th1 response in IL-18-deficient

mice. *Immunity, 8*(3), 383-390. doi: 10.1016/s1074-7613(00)80543-9.

Takeuchi, A., Badr Mel, S., Miyauchi, K., Ishihara, C., Onishi, R., Guo, Z., . . . Saito, T.

(2016). CRTAM determines the CD4+ cytotoxic T lymphocyte lineage. *J Exp Med,*

213(1), 123-138. doi: 10.1084/jem.20150519

Takeuchi, A., & Saito, T. (2017). CD4 CTL, a Cytotoxic Subset of CD4(+) T Cells,

Their Differentiation and Function. *Front Immunol, 8*, 194. doi:

10.3389/fimmu.2017.00194

Wensveen, F. M., Jelencic, V., & Polic, B. (2018). NKG2D: A Master Regulator of

Immune Cell Responsiveness. *Front Immunol, 9*, 441. doi:

10.3389/fimmu.2018.00441

Wilkinson, T. M., Li, C. K., Chui, C. S., Huang, A. K., Perkins, M., Liebner, J. C., . . .

Xu, X. N. (2012). Preexisting influenza-specific CD4+ T cells correlate with disease

protection against influenza challenge in humans. *Nat Med, 18*(2), 274-280. doi:

10.1038/nm.2612

Xie, Y., Akpinarli, A., Maris, C., Hipkiss, E. L., Lane, M., Kwon, E. K., . . . Antony,

P. A. (2010). Naive tumor-specific CD4(+) T cells differentiated in vivo eradicate

established melanoma. *J Exp Med, 207*(3), 651-667. doi: 10.1084/jem.20091921

Zhang, Ping, Lee, Jason S., Gartlan, Kate H., Schuster, Iona S., Comerford, Iain,

Varelias, Antiopi, . . . Hill, Geoffrey R. (2017). Eomesodermin promotes the

development of type 1 regulatory T (T_{R1}) cells. *Sci Immunol, 2*(10), eaah7152. doi:

10.1126/sciimmunol.aah7152

1069

 Figure Legends:

Figure 1: CD4$^+$GzB$^+$PRF$^+$ T cells are present at high frequency in the spleen of

mice infected with *T. cruzi*. (A) CD4$^+$ T cell gating strategy **(B and C)** Representative

flow-cytometry dot plots of GzB **(B)** or PRF **(C)** staining, gated on CD4$^+$ T cells of

non-infected and infected B6 mice (left) and mean frequency of CD4$^+$GzB$^+$ (or

CD4$^+$PRF$^+$) T cells in the spleen (right). **(D)** Representative dot plots of PRF and GzB

expression in CD4$^+$ T cells gated as in (A) and mean frequencies of PFR$^+$ among GzB$^+$

CD4$^+$ T cells (isotype controls on Figure 1-figure supplement 1A). **(E)** Representative

histogram of 2B4 (CD244) expression on CD8$^+$ (blue), CD4$^+$GzB$^+$ (red) or CD4$^+$GzB$^-$

T cells (pink), and correspondent mean MFI values; **(F)** Representative dot plots of

GzB and IFN-γ staining in gated CD4$^+$ T splenocytes (left), mean frequencies of

GzB$^+$IFN-γ$^-$ (red), GzB$^+$IFN-γ$^+$ (purple) and GzB$^-$IFN-γ$^+$ (blue) subsets among total

CD4$^+$ (center) or gated on CD4$^+$GzB$^+$ T cells (right). **(G)** Kinetics of the mean

frequency of CD4$^+$IFN-γ$^+$ (purple) and CD4$^+$GzB$^+$ (red) (left), and corresponding

absolute numbers (right). **(H)** Frequency of LAMP-1$^+$(CD107a$^+$) cells among

CD4$^+$GzB$^+$ (red) or CD8$^+$GzB$^+$ (blue) T cells in the infected spleen. Data obtained from

individually analyzed mice (n=4), at indicated or 14 day pi. Error bars= SEM. ND=

non-detected. *ns*= non-significant; **= p≤ 0.01 (Student t test). All data are

representative of 3 or more independent experiments.

1089

Figure 2: CD4 T cell-mediated Ag-specific cytotoxicity and TF expression by

CD4$^+$GzB$^+$ T cells. (A) Mean frequencies of dead target cells (B blasts) in redirected

cytotoxicity assay employing sorted naive (blue) or *in vivo* activated (red) CD4$^+$ T cells

as effectors, at different E:T ratios, in triplicates. **(B)** Specific lysis of IC-21

macrophages (T) loaded with amastigote antigens, coincubated with highly purified

CD4$^+$ T cells (E) at the indicated E:T ratios, in the presence of anti-FasL mAb (red) or

isotype control mAb (blue). Mean values of triplicates ± SEM are shown in **(A)** and **(B)**;

*p≤ 0.05 (two-tailed Student t-test). Data are representative of 3 independent

experiments. Gating strategy on Figure 2-figure supplement 1A. (**C and D)** Percentages

of Ag-specific cytotoxicity *in vivo*; bars represent mean values + SEM (error bars) of

individually analyzed mice (n=3) infected with *T. cruzi* Y in **(C)** or Y-OVA strain in

(D); class I-restricted (PA8 or OVA$_{257-264}$) and class II-restricted (SA851.1c+d or

OVA$_{323-339}$) peptides were employed. Data are representative of 2 independent

experiments. Cell survival was measured by flow cytometry (Figure 2-figure

supplement 1B) and specific lysis calculated as described in Methods. (**E and F)**

Representative dot plots of CD4$^+$ T cells (gated as in Figure 1A) and corresponding

mean frequencies of gated subpopulations: **(E)** GzB and FoxP3 staining and **(F)** GzB

and T-bet staining. **(G)** Representative histogram of Eomes expression in CD4$^+$ T

splenocytes of non-infected mice (ctr, gray) and in GzB$^-$ (black) and GzB$^+$ (red) CD4$^+$ T

cells from infected mice, gated as in Figs. 1A and 1B (left) and mean frequencies of

Eomes$^+$ cells among these different subpopulations (right). All experiments were done

at day 14 pi. Bars represent mean values of non-infected ctr group n=3 or infected

group, n=3 to 5, individually analyzed mice. Error bars= SEM. **(H)** Runx3d and ThPOK

(*Zbtb7b*) mRNA expression estimated by qRT-PCR in CD4$^+$EYFP$^+$ T cells sorted from

infected and tamoxifen-treated GzmbCreER^{T2}/ROSA26EYFP mice, as described in

Methods and shown in Figure 2-figure supplement 3A; ND = not detected. Data are

representative of 3 independent experiments.

1117

Figure 3: Expression of IL-10, immunoregulatory molecules and cytotoxic

markers by CD4$^+$GzB$^+$ T cells. (A) Representative dot plots of IL-10 staining and

mean frequencies of IL-10$^+$ cells among non-infected (ctr, gray) or GzB$^-$ (black) and

GzB$^+$ (red) CD4$^+$ T cells from infected mice. **(B)** Representative histogram of Lag-3

staining and mean frequency of Lag-3$^+$ cells. **(C)** Representative dot plots of Lag-3 and

CD49b staining and mean frequency of different cells subsets. **(D)** Representative

histogram of Tim-3 staining and mean frequency of Tim-3$^+$ cells among different cells

subsets. **(E)** Representative histogram of CD27 staining and mean frequency of CD27$^+$

cells. **(A-E)** CD4$^+$GzB$^+$ T cells gated as in Figures 1A-1B; staining controls in Figure 3-

figure supplement 1. **(F-H)** Representative dot plots and mean frequencies of **(F)**

NKG2A/C/E, **(G)** NKG2D and **(H)** CRTAM expression among gated CD4$^+$ T

splenocytes. All experiments were done at day 14 pi. Non-infected ctr group, n=3;

infected group, n=3 to 5 mice, individually analyzed. Error bars= SEM; *p≤ 0.05;

***p≤ 0.001; ****p≤ 0.0001 (two-tailed Student t-test). Data are representative of 3

independent experiments. Staining controls are shown on Figure 3-figure supplement 2.

1133

Figure 4: CD4CTLs are severely reduced in *Myd88$^{-/-}$* mice. (A-B) Representative dot

plots of GzB and T-bet staining on gated CD4$^+$ T splenocytes (left panels) and mean

frequencies of T-bet$^+$, IFN-γ$^+$ and GzB$^+$ CD4$^+$ T cells and absolute numbers of

GzB$^+$CD4$^+$ T cells (on the right) from **(A)** WT (B6) and *Il12p40$^{-/-}$* mice on day 14 pi

and **(B)** WT and *Ifng$^{-/-}$* mice on day 11 pi. **(C)** Representative dot plots and gating

strategy, **(D)** mean frequencies and **(E)** absolute numbers of GzB$^+$ or NKG2A/C/E$^+$,

NKG2D$^+$ and CRTAM$^+$ cells among GzB$^+$CD4$^+$ T cells in the spleen of WT (blue) and

Myd88$^{-/-}$ (red) mice at day 13 pi. Bars represent mean values in each group; n= 5

individually analyzed mice. Error bars= SEM; ns= non-significant; *p≤ 0.05; **p≤

0.01; ****p≤ 0.0001 (Student t-test). Representative dot plots of GzB, NKG2A/C/E,

NKG2D and CRTAM staining on CD4$^+$ T cells from non-infected animals are shown

in Figure 4-figure supplement 1B. **(F-G)** Frequencies (left) and absolute numbers

(right) of PRF$^+$ cells among CD4$^+$ T cells **(F)** and of PRF$^+$ cells among GzB$^+$CD4$^+$ T

cells **(G)** in the spleen of WT (blue bars) and *Myd88$^{-/-}$* (red bars) mice. GzB$^+$CD4$^+$ T

cells were gated as shown in Figs. 1A, 1B and 1G. Data in **(F)** and **(G)** are compiled

from four independent experiments with n= 3 animals in each group, each symbol

represents an individual analyzed mice. **(H)** Frequency of dead target cells (B cell

blasts) in cytotoxic assay, after 14 h of co-culture with CD4$^+$CD44^{hi} T cells sorted from

WT (blue line) or *Myd88$^{-/-}$* (red line) infected mice at day 13 pi. B cells were coated

with anti-CD3 (left) or not (right). Mean of triplicates +/- SEM for each E:T ratio point

are shown. Data are representative of 3 independent experiments.

1155

Figure 5: The CD4CTL gene program is absent in MyD88-deficient CD4$^+$ T cells.

(A) GSEA performed using pre-ranked list of genes expressed in WT or *Myd88$^{-/-}$* CD4$^+$

T cells sorted from mix-BM chimeras using the CD4CTL gene signature previously

described as gene set (66). **(B)** Venn diagram of CD4CTL genes upregulated in WT and

downregulated in *Myd88$^{-/-}$* CD4$^+$ T cells sorted from chimeric mice. **(C)** Heat map of

normalized CD4CTL-signature gene expression in WT or *Myd88$^{-/-}$* CD4$^+$ T cells sorted

from chimeric mice: values indicate mean Z-score, upregulated genes in red and

downregulated genes in blue. **(D)** Representative histograms of Blimp-1 expression

(right) and mean frequency of Blimp-1$^+$ splenocytes (left) among total gated CD4$^+$ T

cells from non-infected mice (ctr, gray shaded area and bar) or among CD4$^+$GzB$^+$ (red

line and bar) or CD4$^+$GzB$^-$ (black line and bar) from WT infected mice at day 14 pi.

Individually analyzed mice, n=4. Error bars=SEM, data are representative of 2

independent experiments. **(E)** Representative histograms (left and central panel) and

mean MFI values of CD39 expression (right panel) on gated FoxP3$^-$ (gray shaded area

and bar) or FoxP3$^+$ (blue line and bar) CD4$^+$ T cells from non-infected (ctr) mice (left);

GzB$^-$FoxP3$^-$ (black line and bar), GzB$^-$FoxP3$^+$ (purple line and bar) or GzB$^+$FoxP3$^-$ (red

line and bar) CD4$^+$ T cells from infected WT mice (center), at day 14 pi; individually

analyzed mice (n=4), error bars =SEM; data are representative of 4 independent

experiments. **(F)** Mean frequency (left) and absolute numbers (right) of GzB-expressing

cells among CD45.1$^+$ (B6.SJL WT) or CD45.2$^+$ (B6 WT or KO) CD4$^+$ T splenocytes in

mix-BM chimeric mice, at day 14 pi; WT:WT control chimera group, n=3; other

chimera groups, n=6 or 7. Error bars= SEM, *ns*= non-significant, ***p≤ 0.001,****p≤

0.0001 (Student t-test). Gating strategy and representative dot plots are shown in Figure

5-figure supplement 1. Data are representative of 3 independent experiments.

1180

Figure 6: CD4CTLs are severely reduced in *Il18ra$^{-/-}$* mice. (A and B) Frequencies

and absolute numbers of PRF$^+$ cells among CD4$^+$ T cells **(A)** and of PRF$^+$ cells among

GzB$^+$CD4$^+$ T cells **(B)** in WT (blue bars) and *Il18ra$^{-/-}$* (red bars) spleens. Data are

compiled from two independent experiments with n= 4-5 animals in each group, each

symbol represents an individual mouse. **(C)** Representative dot plots of GzB, CRTAM,

NKG2D and NKG2A/C/E staining in WT and *Il18ra$^{-/-}$* mice. **(D)** Mean frequencies and

(E) absolute numbers of GzB$^+$ or CRTAM$^+$, NKG2D$^+$ and NKG2A/C/E$^+$ cells among

CD4$^+$ and CD4$^+$GzB$^+$ T cells, respectively. **(A-E)** CD4$^+$ T cells gated as in Figs. 1A-1D;

n=4 individually analyzed mice in each group; error bars= SEM, *p≤ 0.05, **p≤ 0.01,

***p≤ 0.001, *ns*= non-significant (Student t-test). Data are representative of 2

independent experiments. **(F)** Frequency of dead target cells (B cell blasts) in cytotoxic

assay, after 14 h of co-culture with CD4$^+$CD44^{hi} T cells sorted from WT (blue line) or

Il18ra$^{-/-}$ (red line) infected mice at day 13 pi. B cells were coated with anti-CD3 (left) or

not (right). Mean of triplicates +/- SEM for each E:T ratio point are shown. **(G)**

Survival curve (left) and parasite load in the heart (right) of infected *Il18ra*$^{-/-}$ mice,

adoptively transferred with CD4$^+$GzB$^+$ cells at day 7 pi (red curve), or not (black

curve); time-line in Figure 6-figure supplement 1; n=10 mice in each group; $p = 0.0277$;

survival data are compiled from two independent experiments.

1199

Figure 7: Cells with CD4CTL phenotype predominate in the cardiac tissue of

infected mice and are at higher frequencies in patients with Chagas

cardiomyopathy. (A) Representative dot plots (gated as in Figure 7-figure supplement

1A), and mean frequency of CD8$^+$ and CD4$^+$ T cells infiltrating the heart of B6 infected

mice at day 14 pi. **(B and C)** Representative dot plot and mean frequency of **(B)** GzB$^+$

and **(C)** LAMP-1$^+$ (CD107a$^+$) cells, among gated CD4$^+$ T cells infiltrating the infected

heart; ctr= non-infected, ND= non-detected. **(D)** Mean parasite load in the cardiac tissue

at different time points pi, obtained by real-time PCR, of individually analyzed mice as

described in Methods; Data are representative of at least 8 **(A and B)** or 2 **(C and D)**

independent experiments.   **(E)** Representative dot plots and mean frequency of

CRTAM$^+$ and NKG2D$^+$ cells among gated intracardiac CD4$^+$GzB$^+$ T cells. DP=

CRTAM$^+$NKG2D$^+$ double positive cells. Individually analyzed mice, n= 4; error bars=

SEM; data are representative of 4 independent experiments. **(F)** Representative dot plots

and mean frequency of PFR$^+$ and GzB$^+$PFR$^+$ cells among intracardiac CD4$^+$GzB$^+$ T

cells; data are compiled from two independent experiments with n= 4 or 5 animals in

each group, each symbol represents an individual analyzed mice. **(G)** Intensity of PRF

staining (MFI) in and **(H)** Frequency of PRF$^+$ cells among CD4$^+$ T cells in PBMCs

from patients with chronic Chagas disease cardiomyopathy (CC; n=10) and healthy

donors (HD; n=4) cultured overnight in medium alone (ctr) or in the presence of

trypomastigote antigens (+trypo); *p$\leq$ 0.05, **p$\leq$ 0.01, *ns*= non-significant; data

analyzed by paired Student t test. **(I)** Frequency of PRF$^+$ cells among CD4$^+$ T cells from

PBMCs of patients with chronic Chagas disease cardiomyopathy (CC; n=10) and

healthy donors (HD; n=4) cultured ON in the presence of trypomastigote antigens.

Gating strategy and representative dot plots are shown in Figure 7-figure supplement 2.

(J) Correlation analysis between the percentage of CD4$^+$PRF$^+$ T cells among CD4$^+$ T

cells in PBMCs from CC patients and left ventricular ejection fraction (LVEF) (left

panel) or left atrium diameter (right panel); indicated *p* and *r* values were calculated

using Pearson correlation test.

SUPPLEMENTAL MATERIAL

Cytotoxic CD4[+] T cells driven by T-cell intrinsic IL-18R/MyD88 signaling predominantly infiltrate *Trypanosoma cruzi*-infected hearts

Carlos-Henrique. D. Barbosa[1,2], Ariel Gomes[1,3], Fábio B. Canto[1,4], Layza M. Brandão[1], Jéssica R. Lima[1], Guilherme A. Melo[1], Alessandra Granato[1,5], Eula G.A. Neves[6], Walderez O. Dutra[6], Ana C. Oliveira[7], Alberto Nóbrega[1] and Maria Bellio[1]*.

*corresponding author.
e-mail: belliom@micro.ufrj.br or mariabellioufrj@gmail.com

Supplemental Figures.

Figure 1–figure supplement 1: Comparative percentages and absolute numbers of GzB$^+$ and PRF$^+$ T cell subsets. (A) Isotype control staining of CD4$^+$ T cells, gated as in Fig. 1A, from infected mice. **(B)** Gating strategy for **C** and **D** plots. **(C)** Representative dot plots of GzB staining and mean frequency of GzB$^+$CD8$^+$T cells (on the right). **(D)** Representative dot plots of PRF and GzB staining in CD8$^+$ T cells. **(E)** Mean frequency and **(F)** absolute numbers of CD4$^+$ and CD8$^+$ T cells in the spleen of non-infected (ctr) and infected mice. **(G)** Mean frequency and **(H)** absolute numbers of GzB$^+$CD4$^+$ and GzB$^+$CD8$^+$ T cells. Mice individually analyzed at day 14 pi (n=4); error bars= SEM. Data are representative of 6 independent experiments.

A

B

Figure 2-figure supplement 1: Gating strategy of cytotoxic assays. **(A)** Gating strategies and PI staining of IC-21 macrophages loaded with total amastigote protein extract (CFSE^{lo}) or not (CFSE^{hi}) and cocultured with 4×10^5 purified CD4$^+$ T cells (right), or not (left); specific cytotoxicity values are shown in **Figure 2B**; **(B)** Gating strategies for CTV-stained target splenocytes, loaded with the indicated antigenic peptides or not (ctr), 20 h after i.v. injection in non-infected or Y strain-infected mice. The *T. cruzi*-derived PA8 (class I-restricted) and SA85 (class II-restricted) peptides were employed. Specific cytotoxicity values for the *in vivo* assay are shown in **Figure 2C**.

Figure 2-figure supplement 2: The majority of GzB$^+$CD4$^+$ and GzB$^-$CD4$^+$ T cells is composed of non-naive (activated effectors and memory cells). Representative dot plots and gating strategy of staining for GzB, CD44 and CD62L (on the left) and mean frequency of naive (CD44^{lo}) and non-naive (CD44^{hi}) CD4$^+$ T cells (on the right) in: **(A)** non-infected mice and gated on total CD4$^+$TCRβ$^+$ and **(B)** infected mice and gated on GzB$^+$ (top) or GzB$^-$ (bottom) CD4$^+$TCRβ$^+$ cells, at day 14 pi. Bars represent mean values of cell frequency of individually analyzed mice; n=4; Error bars = SEM. Data are representative of 3 independent experiments.

A

B

Figure 2-figure supplement 3: Expression of *Tbx21* and *Cd8a* in sorted GzB⁺CD4⁺ T cells. (A) Sorting gate and frequency of CD4⁺GzB⁺ (EYFP)⁺ T cells obtained from infected and tamoxifen-treated GzmbCreER^*T2*/ROSA26EYFP mice. CD4⁺ cells were enriched by negative selection with magnetic beads, as described in Methods, and then submitted to FACS-sorting. Pre-sorting CD4⁺ T cell frequency is shown on the left panel and post-sorting on the right. **(B)** Sorted cells had RNA extracted and *Tbx21*, *Runx3d*, *Zbtb7b* and *Cd8a* expression tested by qRT-PCR, as described in Methods. Results for *Runx3d* and *Zbtb7b* expression are shown in **Fig. 2H.** ND = not detected.

Figure 3-figure supplement 1: IL-10, Lag-3 and CD49b expression. (A) Representative dot plots and gating strategy for IL-10 expression. Total spleen cells (left) and gated CD4$^+$ T cells (middle and right panels) of non-infected control mice. **(B)** Representative dot plot of IL-10 staining in total splenocytes of infected mice. Dot plot and mean frequencies of IL-10$^+$ cells among CD4$^+$GzB$^-$ and CD4$^+$GzB$^+$ T cells in infected mice are shown in **Fig. 3A. (C)** Representative dot plot of Lag-3 and CD49b expression on gated B220$^+$, NK1.1$^+$ or total CD4$^+$ T cells from the spleen of non-infected mice (left, middle and right panels, respectively). Mean frequencies of Lag-3$^+$CD49b$^+$ and Lag-3$^+$CD49b$^-$ cells among CD4$^+$GzB$^-$ and CD4$^+$GzB$^+$ T splenocytes of infected mice are shown in **Fig. 3C.** Data are representative of 2 independent experiments.

Figure 3-figure supplement 2: Representative dot plots and gating strategies for NKG2A/C/E, NKG2D and CRTAM staining on CD4$^+$ and CD8$^+$ GzB$^+$ T cells. Staining of spleen cells from non-infected controls (right panels) and FMO and isotype controls of spleen cells from B6 infected mice (left panels) are shown. Representative dot plots and mean frequencies of CD4$^+$GzB$^+$ and CD8$^+$GzB$^+$ T cells

expressing these cytotoxic markers are shown in **Figure 3F-H** and Figure 3-figure supplement 3, respectively.

Figure 3-figure supplement 3: Expression of NKG2A/C/E, NKG2D and CRTAM cytotoxic markers by CD8⁺GzB⁺ T cells. Representative dot plots and gating strategies of spleen cells from B6 infected mice (left panels). Graphs on the right: bars represent mean values of cell frequency of each indicated cell subpopulation in individually analyzed mice; n=4; Error bars = SEM. Data are representative of 3 independent experiments.

Figure 3-figure supplement 4: Frequency of CRTAM-expressing cells among splenic GzB⁺ and GzB⁻ CD4⁺ T cells, gating or not on CD44^hi cells. Bars represent mean values of the frequency of CRTAM⁺ cells among: CD4 naive cells of non-infected (ctr) mice (white bar) and CD4⁺GzB⁻ (gray bar), CD4⁺GzB⁺ (orange bar), CD4⁺CD44^hiGzB⁻ (black bar) or CD4⁺CD44^hiGzB⁺ (red bars) of infected mice at day 14 pi; Gating as shown in **Sup. Figs. 3 and 6C**. Mice were individually analyzed, n= 4; Error bars = SEM; *ns*= non-significant, Student t test.

Figure 4-figure supplement 1: Equivalent frequencies and absolute numbers of total CD4[+] T cells in the spleens of infected *Myd88[-/-]* and WT mice. (A) Mean frequencies and absolute numbers of total CD4[+] and CD8[+] T cells in the spleens of non-infected (ctr) and infected *Myd88[-/-]* and WT (B6) mice, at day 13 pi. **(B)**

Representative dot plots and gating strategies for GzB, CRTAM, NKG2D and NKG2A/C/E staining of spleen cells from non-infected (ctr) WT (B6) and *Myd88*[-/-] mice. Staining results obtained with infected mice are shown on **Figures 4C-4E**. **(C)** Mean frequencies and **(D)** absolute numbers of GzB^+, GzB^+CRTAM^+, GzB^+NKG2D^+ and $GzB^+NKG2A/C/E^+$ cells among $CD8^+$ T cells in the spleen of WT and *Myd88*[-/-] infected mice at day 13 pi, gated as in Sup. Fig. 7. Bars represent mean values in each group; error bars = SEM; n= 5 individually analyzed mice. Results are representative of 2 independent experiments.

**Figure 4-figure supplement 2: Equivalent frequency and absolute number of activated CD4$^+$ T cells in WT and *Myd88$^{-/-}$* mice infected with *T. cruzi.* Representative dot plots (top) and mean frequencies and absolute numbers (bottom) of CD4$^+$CD44^{hi} T cells in the spleens of infected WT (B6) and *Myd88$^{-/-}$* mice, at day 14 pi. Mice were individually analyzed; n=4; error bars = SEM. Representative of 2 independent experiments.

Figure 5-figure supplement 1: GzB expression in CD4$^+$ T cells of mixed-BM chimeric mice. Gating strategy and representative dot plots of: **(A)** CD45.1$^+$ and CD45.2$^+$ cells (on the right) gated CD4$^+$CD8$^-$ T lymphocytes (on the left) from the spleens of mixed-BM chimeric mice. **(B)** GzB expression in CD4$^+$CD45.1$^+$ (WT) and CD4$^+$CD45.2$^+$ (WT or KO) T cells from the spleens of non-infected and infected chimeric mice at day 14 pi, gated as in **(A).** Mean frequencies and absolute numbers of CD4$^+$GzB$^+$ in each group are shown in **Figure 5F**.

Figure 6-figure supplement 1: Time-line of the adoptive transfer experiment shown on Figure 6G.

Figure 6-figure supplement 2: The adoptive transfer of *Ifng*[-/-] CD4[+] T cells increased survival to infection. Survival curve of *Il18ra*[-/-] mice, infected with 1 x 10^3 trypomastigotes of the Y strain and adoptively transferred with sorted (>98% pure) *Ifng*[-/-] CD4[+] T cells (green curve), or not (PBS group, black curve); n=6 mice in each group; p= 0.0295 Log-rank (Mantel-Cox) test.

Figure 7-figure supplement 1: CD4⁺ and CD8⁺ T cells in the heart of mice infected with *T. cruzi*. (A) Representative FSC x SSC dot plots and gating strategy of cells from the hearts of non-infected (left panel) and infected (right panel) WT mice, employed for data analysis shown on this figure and Figure 7. **(B)** Representative dot plots of blood samples from infected mice injected iv with anti-CD4-FITC mAb, 3 min before euthanasia: with no other staining (left panel) or after staining with anti-

CD4-PECy7 mAb *in vitro* (right panel). Representative dot plots of cells from the heart of non-infected and infected mice, injected *iv* with anti-CD4-FITC mAb and stained *in vitro* with anti-CD4-PECy7 (two panels on the right), and mean frequencies of intravenously stained CD4$^+$ T cells in the hearts of non-infected and infected mice. Mice were individually analyzed; n=3; error bars = SEM. Representative of 2 independent experiments. **(C)** Representative dot plots and gating strategy of isotype and GzB (left panels) or LAMP1 (right panels) staining and corresponding mean frequencies of GzB$^+$CD8$^+$ and LAMP1$^+$CD8$^+$ cells among gated CD8$^+$ T cells infiltrating the infected myocardium at day 14 pi. **(D)** Representative dot plots of isotype and NKG2A/C/E staining on gated CD4$^+$GzB$^+$ (left panels) or GzB$^+$CD8$^+$ (right panels) T cells and corresponding mean frequencies of NKG2A/C/E$^+$ cells among GzB$^+$CD4$^+$ and GzB$^+$CD8$^+$ T cells infiltrating the infected myocardium at day 14 pi. Individually analyzed mice, n= 4; error bars= SEM; ND= not detected. Data are representative of 2 independent experiments.

Figure 7-figure supplement 2: CD4$^+$ T cells expressing PRF are found in the circulation of patients suffering from chronic Chagas cardiomyopathy (CC). (A) Dot plot showing the gating strategy for CD4$^+$ T cell staining on PBMC. **(B)** Gating strategy and representative dot plots of PRF staining in gated CD4$^+$ T cells from CC patient incubated with medium (on the left), or with trypomastigote antigens (trypo) (on the right). Frequencies of PRF$^+$CD4$^+$ T cells and MFI of PRF staining for CC patients and HD controls are shown on **Figures 7G-7I**.

Figure 7-figure supplement 3: Predicted model for the generation of CD4CTLs cells in response to *T. cruzi* infection. This cartoon summarizes our results and also contains information of previous published studies, cited in this work. IL-12 produced during infection acts on TCR-triggered CD4[+] T cells, leading to T-bet expression, which promotes IFN-γ upregulation and induces *Runx3d* and *Eomes* gene expression. In turn, IFN-γ- and IL-12-signaling induce IL-18R expression. T-bet, Eomes and Runx3d drive the expression of cytotoxic effector molecules, including GzB, PRF and FasL. IL-10 expression can be induced by Blimp-1 and Eomes, although few IL-10[+] CD4[+]GzB[+] T cells were found in our infection model. As opposed to intraepithelial CD4CTLs of the gut, ThPOK expression is not downmodulated in splenic CD4[+]GzB[+] T cells, until day 14 pi. IFN-γ is produced by around 18% of these cells at the peak of the CD4[+] T cell response (14 dpi), and is necessary for CD4[+]GzB[+] T cell proliferation and/or survival. Most CD4[+]GzB[+] T cells express Lag-3, Tim-3 and CD39 immunoregulatory molecules, while CD27 is downmodulated. Notably, the expansion and/or survival of CD4[+]GzB[+] T cells expressing CRTAM, NKG2D, and NKG2A/C/E cytotoxic markers depend on T-cell intrinsic IL-18R/MyD88 signaling.