Widespread anthropogenic translocation of the African Clawed Frog across its native range is revealed by its co-introduced monogenean parasite in a co-phylogeographic approach

Anneke L Schoeman[1,2,3], Louis H du Preez[1,3], Nikol Kmentová[4,5], and Maarten PM Vanhove[4,5]

[1] African Amphibian Conservation Research Group, Unit for Environmental Sciences and Management, North-West University, Potchefstroom Campus, 11 Hoffman Street, Potchefstroom 2531, South Africa

[2] DSI-NRF Centre of Excellence for Invasion Biology, Stellenbosch, South Africa

[3] South African Institute for Aquatic Biodiversity, Private Bag 1015, Grahamstown 6140, South Africa

[4] Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2, 611 37 Brno, Czech Republic

[5] Hasselt University, Centre for Environmental Sciences, Research Group Zoology: Biodiversity & Toxicology, Agoralaan Gebouw D, B-3590 Diepenbeek, Belgium

Correspondence Anneke Lincoln Schoeman; email: anneke.lincoln@gmail.com

ABSTRACT

1. The management of biological invasions relies upon the development of methods to trace their origin and expansion. Co-introduced parasites, especially monogenean flatworms, are ideal tags for the movement of their invasive hosts due to their short generations, direct life cycles and host specificity. However, they are yet to be applied to trace the intraspecific movement of species in their native ranges.

2. As proof of this concept, we conducted a co-phylogeographic analysis based upon two mitochondrial markers of a globally distributed frog *Xenopus laevis* and its monogenean flatworm parasite *Protopolystoma xenopodis* in both its native range in southern Africa and its invasive range in Europe.

3. Translocation of lineages was largely masked in the frog's phylogeography. However, incongruent links between host and parasite phylogeography indicated host switches from one host lineage to the other after these were brought into contact due to human-mediated translocation in the native range. Thus, past translocation of host lineages is revealed by the invasion success of its co-introduced parasite lineage.

4. This study demonstrates the concept that parasite data can serve as an independent line of evidence in invasion biology, also on the intraspecific level, shedding light on previously undetected invasion dynamics. Based upon the distribution of these invasive parasite lineages, we infer that the widespread translocation of hosts is mainly facilitated by the frog's use as live bait by the local angling communities and not via official export routes.

5. *Synthesis and applications.* Data from co-introduced, host-specific parasites can add value to investigations in invasion biology and conservation. A better understanding of the translocation history and resulting genetic mixing of animals in their native ranges prior to introduction into new environments can inform management strategies in the invasive range. Knowledge of the intraspecific movement of different lineages of animals in their native ranges also has conservation implications, since contact between divergent lineages of hosts and parasites can facilitate host switches and altered parasite dynamics in both native and invasive populations. Therefore, we recommend the inclusion of parasite data as a more holistic approach to the invasion ecology of animals on the intraspecific level.

Keywords: biological invasion, co-invasion, holistic specimen, intraspecific cryptic invasion, introduction pathway, *Protopolystoma xenopodis*, *Xenopus laevis*

1 INTRODUCTION

The Anthropocene is characterised by the unparalleled human-mediated passage of organisms from their natural ranges to ecosystems across the world (Capinha *et al.*, 2015; Ricciardi, 2007). If these non-native organisms manage to disperse from their initial introduction point and seed new populations in their non-native range, they are regarded as invasive (Blackburn

et al., 2011; Richardson *et al.*, 2011). This reshuffling of the world's biota is a decisive agent of global change, as evidenced

by the fact that invasive species are implicated as the cause of a third of recent animal extinctions (Blackburn *et al.*, 2019).

Clearly, the development of methods to trace both the origin and expansion of biological invaders is an urgent, but not

always straightforward, task. For one, organisms can be both native and invasive within the borders of their country of origin,

especially in large countries with high climatic and habitat diversity (van Wilgen *et al.*, 2020). These so-called domestic

exotics occur outside their native range, yet remain within one administrative region, a feature which hinders their detection

and management (Guo & Ricklefs, 2010). Along the same lines is what is known as intraspecific cryptic invasion. This

describes the process where one lineage of a species spreads into parts of the species' native range where another lineage is

already present, often with human help (Morais & Reichard, 2018; Saltonstall, 2002).

Genomic tools can be highly informative for the reconstruction of invasion histories, the prediction of potential range

expansion and invasion success and the development of control strategies (e.g. de Busschere *et al.*, 2016; Dufresnes *et al.*,

2017; Dlugosch & Parker, 2008; Hiller & Lessios, 2017; Lee & Gelembiuk, 2008; Prentis *et al.*, 2009; Stouthamer *et al.*,

2017; Zahiri *et al.*, 2019). Nonetheless, in many cases the available molecular data or chosen molecular markers are of

insufficient resolution by themselves to trace the movement of species (e.g. Ahyong *et al.*, 2017; Criscione *et al.*, 2006). In

fact, the use of molecular population genetics at the short timescales and coarse-grained genomic resolution of most biological

invasions seems to be applicable only in rare instances (Fitzpatrick *et al.*, 2012).

A more holistic approach to solving the origin and range shifts of invasive species involves molecular data from co-

introduced parasites. Such an integrative approach has been proposed to investigate invasions that are difficult to trace

(Clavero *et al.*, 2016). For example, the long-debated cryptogenic history of the European marine snail *Littorina littorea*

was solved with the help of an associated digenean parasite (Blakeslee *et al.*, 2008). Likewise, intraspecific morphometric

variation of monogenean flatworm parasites identified both the origin and introduced life stage of invasive clupeid fish in

Africa (Kmentová *et al.*, 2019), shed light on the history of an invasive population of odontobutid fish in Europe (Ondračková

et al., 2012) and provided information on the introduction route of an invasive goby in Europe (Huyse *et al.*, 2015). Therefore,

the use of parasites as tags for host connectivity is not new (Gagne *et al.*, 2021). Yet, the co-phylogeographic analysis of host

and parasite data has never been applied to trace intraspecific cryptic invasions within the native ranges of species. In a proof

of concept investigation, we employed a globally distributed frog-parasite system to uncover within-species movement.

The African Clawed Frog *Xenopus laevis* (Daudin, 1802) (Anura: Pipidae) is arguably the world's most widely distributed

and well-studied amphibian (Measey *et al.*, 2012; van Sittert & Measey, 2016). The ability to trace the native origins of the

invasive populations of *X. laevis* is brought about by the high levels of phylogeographic structuring of this frog in its native range of southern Africa (de Busschere *et al.*, 2016; Furman *et al.*, 2015). Previous studies on the morphology and genetics of *X. laevis* have identified several differentiated populations in South Africa that are segregated by large mountain ranges and different climatic regions (de Busschere *et al.*, 2016; du Preez *et al.*, 2009; Furman *et al.*, 2015; Grohovaz *et al.*, 1996; Measey & Channing, 2003).

Based upon the widespread occurrence of both natural and human-mediated translocation of *X. laevis* in the region, a consequence of the domestic export of the frog for teaching and learning and the widely adopted practice of translocating the frog for use as live bait in recreational angling, some authors advocate for its recognition as a domestic exotic (Measey & Davies, 2011; Measey *et al.*, 2017; van Sittert & Measey, 2016; Weldon *et al.*, 2007). These translocations should leave a genetic trace (van Sittert & Measey, 2016). Nonetheless, the cryptic invasion of *X. laevis* over long distances and across major geographic barriers in the native range has not been confirmed in previous phylogeographic studies of this frog (de Busschere *et al.*, 2016; du Preez *et al.*, 2009; Furman *et al.*, 2015; Measey & Channing, 2003).

Thus, we posit that the bladder polystomatid flatworm *Protopolystoma xenopodis* (Price, 1943) Bychowsky, 1957 (Monogenea: Polystomatidae) is the ideal tool to trace the hitherto unconfirmed contact between different lineages of its host *X. laevis* across its native range of southern Africa. In natural populations across sub-Saharan Africa, *P. xenopodis*, the most widely distributed of the species of *Protopolystoma*, is restricted to a single host species complex, *X. laevis sensu lato*, even in sympatry with other *Xenopus* species (Tinsley & Jackson, 1998a,b). Patterns of host-specificity and speciation within *Protopolystoma* reflect the phylogenetic relationships among their hosts, hinting at a very close co-evolutionary association (Tinsley & Jackson, 1998a). Furthermore, in *P. xenopodis*, notable intraspecific morphological variation exists between populations of the polystomatid in southern Africa and those infecting more northerly host subspecies (Tinsley & Jackson, 1998b). These factors suggest that the phylogeny of *P. xenopodis* might mirror that of *X. laevis* also on an intraspecific level, meaning that this host-parasite system is a good candidate for a co-phylogeographic study.

In general, monogenean parasites (Platyhelminthes: Monogenea) exhibit sufficient levels of phylogeographic structuring to assess intraspecific spatial patterns in their hosts (Baldwin *et al.*, 2012; Hahn *et al.*, 2015; Huyse *et al.*, 2017; Kmentová *et al.*, 2018). At the very least, the phylogeographic structuring of monogenean lineages can support evidence revealed by the host data set or historic records, such as the well-documented anthropogenic introduction of salmonid fish host stocks across Europe that is reflected in the phylogeography of their gyrodactylid monogenean lineages (Hahn *et al.*, 2015). In particular, monogenean parasites are regarded as ideal tags of host genealogy due to their short generations, direct life cycles

and generally high host-specificity (Nieberding & Olivieri, 2007; Pariselle *et al.*, 2011).

All things considered, the well-studied, widely distributed and intimately associated *X. laevis* and *P. xenopodis* system

may act as a proof of concept that intraspecific parasite phylogeography can provide information on the cryptic introductions

of their host lineages. Therefore, in the present study, we will assess the performance of this integrative approach in the

detection of recent translocation of *X. laevis* lineages in the native range. We hypothesise that the phylogeography and

morphological divergence of *P. xenopodis* will mirror the documented intraspecific divergence of its host *X. laevis*. Thus, we

propose that a co-phylogeographic analysis of this host and parasite can identify recent human-mediated contact between

distinct lineages of *X. laevis* and provide new insights on invasion pathways in the native range and beyond.

87 2 MATERIALS AND METHODS

88 2.1 Host and parasite sampling

In the native range of southern Africa, 46 adult *X. laevis* were captured from March 2017 to April 2019 in liver baited funnel

traps from 30 localities (Fig. 1; Appendix S1 for locality details). The localities were representative of the phylogeography of

the frog, based upon the lineage assignment of de Busschere *et al.* (2016) (Appendix S2 for locality classification). In the

invasive range, 13 adult *X. laevis* were collected from 12 localities in western France and one in Portugal during the summers

of 2017, 2018 and 2019 (Appendix S1 for locality details). In France, frogs were captured in baited traps by members of

the EU LIFE CROAA team as part of the regional eradication programme for invasive amphibians. Frogs in Portugal were

obtained from the locally captured laboratory stock of the University of Lisbon.

All frogs underwent double euthanasia with anaesthesia in 6% ethyl-3-aminobenzoate methanesulfonate (MS222) (Sigma-

Aldrich Co., USA) followed by pithing as required by the institutional ethics committee. During subsequent dissection, 59

P. xenopodis (one per frog) were retrieved from the bladders for molecular analyses. Polystomatids and liver tissue from their

corresponding frog hosts were preserved in 70% ethanol. A further 23 polystomatids were retrieved from the same frogs hosts

at eight of the 30 sampled localities in South Africa for the morphometrical comparison of polystomatids from the southwest

of South Africa versus the northeast of South Africa (see Appendix S3 for morphometrical methods and findings).

102 2.2 Molecular data collection

103 The DNA was extracted from host and parasite tissue with the PCRBIO Rapid Extract PCR kit (PCR Biosystems Ltd., UK) or

104 the *Quick*-DNA^{TM} Miniprep Plus kit (Zymo Research, USA). Subsequently, amplification reactions for the mitochondrial

105 *COX1* and *rrnS* genes of both frogs and polystomatids were prepared with 2 to 5 μL genomic DNA, 1.25 μL forward primer

FIGURE 1 Localities in southern Africa where 46 adult *Xenopus laevis* were collected. Localities are classified according to the geographic extent of the native phylogeographic lineages of *X. laevis* (de Busschere *et al.*, 2016). Key localities are named according to the closest town. Jonkershoek (☆) is the source locality of the majority of historic *X. laevis* exports. Mountain ranges are indicated with shaded relief and rainfall seasonality is represented by the wettest month of each 25 km^2 grid cell. Monthly mean precipitation data, obtained from the WorldClim 2.1 database at a spatial resolution of 2.5 arc-minutes (Fick & Hijmans, 2017), were used to identify the months with the highest historical mean precipitation from 1970 to 2000 for each grid cell. Elevation data were obtained from the Natural Earth database. The map was constructed in QGIS *version* 3.10.2-A Coruña (QGIS Development Team, 2018) with the Mercator projection.

[0.2 μM], 1.25 μL reverse primer [0.2 μM], 12.5 μL master mix and PCR-grade water to the final volume of 25 μL. The 2×

PCRBIO HS Taq Mix Red (PCR Biosystems Ltd.) or the OneTaq® 2× Master Mix with Standard Buffer (New England

Biolabs Inc., USA) were used with the temperature of the elongation steps as per supplier instructions (68 °C or 72 °C).

For the frog *COX1*, the 'LCO1490'/'HCO2198' (Folmer *et al.*, 1994) and 'VF1'/'VR1' (Ivanova *et al.*, 2006; Ward *et al.*,

2005) primer pairs were implemented with the thermocycling profile: initial denaturation at 95 °C for 2 min, followed by

35 cycles of denaturation at 95 °C for 30 s, annealing for 30 s at 50 °C for the first primer pair and 54 °C for the second

and elongation for 2 min, terminated by a final elongation step of 7 min. In the case of the '12SA-F'/'12SB-R' primer pair

(Kocher *et al.*, 1989) to obtain the frog *rrnS*, the profile of Amaro *et al.* (2013) was followed. For the amplification of the

polystomatid *COX1* and *rrnS*, the profile of Héritier *et al.* (2015) was implemented for the primer pairs 'L-CO1p'/'H-Cox1R',

'L-CO1p'/'H-Cox1p2', '12SpolF1'/'12SpolR1' and '12SpolF1'/'12SpolR9' (Héritier *et al.*, 2015; Littlewood *et al.*, 1997).

For purification and sequencing, PCR products were sent to a commercial company (Inqaba Biotec, RSA) that used

the ExoSAP protocol (New England Biolabs Ltd.) for purification and obtained the sequences with BigDye® Terminator

version 3.1 Cycle Sequencing, utilising the corresponding primer pairs used in the PCR reaction, on an ABI3500XL analyser.

Sequences were assembled and manually edited in Geneious Prime *version* 2021.0.3 (www.geneious.com). The 236 obtained

sequences were deposited in GenBank (accession numbers to be added after article acceptance).

121 2.3 Phylogeographic analysis

The phylogeography of the 59 *X. laevis* and the 59 *P. xenopodis* was visualised in two haplotype networks. All genes were

aligned using the Clustal Omega algorithm under default settings (Sievers & Higgins, 2017) implemented in Geneious Prime.

For the protein-coding *COX1* gene, translation alignment was implemented under the vertebrate mitochondrial genetic code

(table 2) for the frog and the echinoderm and flatworm mitochondrial genetic code (table 9) for the polystomatid. The gene

alignments were trimmed to the length of the shortest sequence since gaps are masked during the subsequent analysis (Leigh

& Bryant, 2015). The two aligned genes of each species were concatenated by individual in Geneious Prime. Minimum

spanning networks (MSN) of the frog and polystomatid data sets were inferred in PopART *version* 1.7.2 (Bandelt *et al.*, 1999;

Leigh & Bryant, 2015).

The genetic diversity of the two host and the two parasite genes was studied based upon these trimmed alignments prior

to concatenation. Number of haplotypes and polymorphic sites, haplotype diversity, nucleotide diversity and mean percentage

of pairwise differences were calculated in in Arlequin *version* 3.5 (Excoffier & Lischer, 2010) for the alignments as a whole

and for predefined groups. Three groups were defined based upon collection locality (see Appendix S2 for full details),

namely specimens collected in South Africa to the southwest of the Great Escarpment (SA1, SA2, SA3, SA4 and SA7), to the

northeast of the Great Escarpment (SA5 and SA6), or in France. Specimens from Portugal were not included in the analyses

because of small sample size.

137 2.4 Phylogenetic analyses

138 For the phylogenetic analyses, duplicate sequences were identified with the help of CD-HIT (Fu *et al.*, 2012; Li & Godzik,

139 2006) implemented online (Huang *et al.*, 2010). Specimens with unique *COX1* or *rrnS* gene haplotypes were retained for the

140 analysis. The *COX1* and *rrnS* gene sequences of the congeneric Müller's clawed frog *X. muelleri* (Peters, 1844) (GenBank

accession numbers EU599031, AY581657; Evans *et al.*, 2004, Evans *et al.*, 2008) and its polystomatid *P. occidentalis* Tinsley *et* Jackson, 1998 (GenBank accession numbers KR856179, KR856121; Héritier *et al.*, 2015) were added as outgroups. This polystomatid only infects *X. muelleri*-like species, even in sympatry with *X. laevis* (Tinsley & Jackson, 1998b). Furthermore, its host *X. muelleri* reportedly reproduces sterile offspring when hybridisation occurs between this species and its sympatric congener *X. laevis* (Fischer *et al.*, 2000). Therefore, the selected outgroups are applicable to an intraspecific analysis of the *X. laevis* and *P. xenopodis* host-parasite system. The resulting data sets, after removal of duplicates and addition of the outgroups, were aligned and concatenated as for the haplotype networks, but not trimmed. Descriptive statistics of the two alignments were derived in DnaSP *version* 6.12.03 (Rozas *et al.*, 2017).

Initially, the concatenated alignments comprised four partitions each, namely the three codon positions of the partial *COX1* gene and the *rrnS* gene. The ModelFinder selection routine (Kalyaanamoorthy *et al.*, 2017), in conjunction with the FreeRate heterogeneity model (Soubrier *et al.*, 2012), implemented in IQ-TREE *version* 2.1.2 (Minh *et al.*, 2020), was employed to select the optimal model of molecular evolution for each partition. To reveal the optimal partitioning scheme and most suitable evolutionary models for each alignment, the initial partitions were sequentially merged (Chernomor *et al.*, 2016) until model fit, as evaluated by the Bayesian Information Criterion (BIC), ceased to improve. For both the data sets, the optimal partitioning scheme was a single partition of all genes and codon positions.

The maximum likelihood (ML) phylogeny of the host was inferred under the TIM2 model (Posada, 2003) with edge-unlinked FreeRate heterogeneity models (Soubrier *et al.*, 2012), proportions of invariant sites (Gu *et al.*, 1995) and discrete gamma models (Yang, 1994). Similarly, the ML tree of the polystomatid was inferred under the TPM3 model (Kimura, 1981) with a FreeRate heterogeneity model (Soubrier *et al.*, 2012). The analyses were performed in IQ-TREE *version* 2.1.2 (Minh *et al.*, 2020), with the assessment of branch support through ultrafast bootstrapping (UFboot2; Hoang *et al.*, 2018) and the Shimodaira-Hasegawa approximate likelihood ratio test (SH-aLRT; Guindon *et al.*, 2010), each with 10 000 replicates. The two phylogenies were rooted with respect to the outgroups in TreeGraph2 *version* 2.15.0-877 *beta* (Stöver & Müller, 2010). Clades with UFboot2 support lower than 95% or SH-aLRT lower than 80% were collapsed to polytomies in the same programme.

2.5 Co-phylogenetic analyses

The resulting phylogenies formed the basis of a global-fit approach to the co-phylogenetic analysis of *X. laevis* and *P. xenopodis*. The global congruence of the two phylogenies was assessed by the implementation of both Procrustean Approach to Co-

phylogeny (PACo) (Balbuena *et al.*, 2013) and the fourth-corner statistical approach of ParaFit (Legendre *et al.*, 2002). Both

these methods also consider the contribution of individual species interactions to phylogenetic congruence (Balbuena *et al.*,

2013; Legendre *et al.*, 2002; Poisot & Stouffer, 2015). Moreover, both methods can accommodate trees that are not fully

resolved and support instances of uneven numbers of terminal nodes of hosts and parasites.

In PACo, we assessed the dependency of the parasite phylogeny upon that of the host through asymmetrical Procrustean

superimposition of host and parasite distance matrices, whilst ParaFit tested the dependence of both phylogenies upon

one another (Balbuena *et al.*, 2013; Legendre *et al.*, 2002). Prior to analyses, the outgroups were removed from the ML

phylogenies. The phylogenies were converted to matrices of patristic distances by the package *ape* (Paradis & Schliep, 2019)

in R *version* 4.0.2 (R Core Team, 2020). These matrices served as input for both analyses, along with the corresponding

host-parasite association matrix, where the polystomatids were assigned to their original host individuals, or to a host with an

identical gene sequence to the original host individual.

The principal coordinates for PACo were derived with the R package *paco* (Hutchinson *et al.*, 2017) *version* o.4.2 for the

Procrustes superimposition in conjunction with the Cailliez correction (Cailliez, 1983) to avoid the production of negative

eigenvalues of non-Euclidean phylogenetic distances. The PACo analysis was performed, using the asymmetric Procrustes

statistic to test for phylogenetic tracking by the parasite of its host, in *paco* with 10 000 permutations, with conserved

row sums and number of interactions. The dynamics of the interacting clades were further explored by identifying those

interactions that most contribute to the possible concordance between the two phylogenies through a jackknife procedure in

paco. Those interactions with the smallest residual distances were interpreted as stronger supporters of overall congruence.

The ParaFit analysis (Legendre *et al.*, 2002) was carried out with *ape* based upon the patristic distance matrices, using

10 000 permutations with the Cailliez correction (Cailliez, 1983). Both the ParaFitLink tests were employed to assess the

significance of the individual links (Legendre *et al.*, 2002).

A tanglegram of the host and parasite ML phylogenies with a weighted interaction network, based upon the residuals

yielded by the PACo jackknife procedure, was constructed with the R package *phytools* (Revell, 2012) with the function

`cophylo()`, which has an internal algorithm that rotates nodes to optimally match tips. Due to the removal of duplicate

sequences, certain host and parasite terminal nodes were involved in multiple interactions.

 3 RESULTS

 3.1 Sequence diversity

195 The trimmed and concatenated alignment of the two host genes used for the network analysis was 1010 bp in length. In

196 the case of the two parasite genes, a concatenated alignment of length 900 bp included seven sites with gaps that were not

197 included for inference. The genetic diversity of the two parasite genes were generally higher than the two orthologous host

198 genes, with greater genetic diversity among the samples collected in South Africa than in France in both host and parasite

199 (Table 1). Genetic diversity was generally higher in the hosts collected to the southwest of the Great Escarpment than in those

200 collected from the northeast, which was not the case for the parasites (Table 1).

201 In terms of the alignments used for the phylogenetic analyses, analysis of redundancy identified 30 unique sequence

202 combinations in the concatenated *X. laevis* data set with 23 unique haplotypes of *COX1* and 10 of *rrnS*. Including the outgroup

203 *X. muelleri*, 31 haplotypes were aligned for the ML analysis to a total of 1153 bp of 717 bp *COX1* and 436 bp *rrnS* containing

204 189 sites with missing data and 183 variable sites, of which 88 were parsimony informative. Similarly, after removal of

205 duplicates, the concatenated parasite alignment contained 44 unique *P. xenopodis* sequences, a combination of 34 unique

TABLE 1 Genetic diversity of the bladder flatworm *Protopolystoma xenopodis* and its frog host *Xenopus laevis*, grouped according to sampling locality – collected in South Africa to the southwest of the Great Escarpment or to the northeast of the Great Escarpment, or in France. Indices were inferred separately by gene for each group and for all samples combined from the aligned and trimmed partial mitochondrial *COX1* and *rrnS* genes used in the haplotype network analyses.

	n	H	L	S	Hd	π	d (%)
X. laevis COX1							
Southwestern SA	30	11	641	78	0.823 ± 0.061	0.026 ± 0.013	2.6
Northeastern SA	16	6	641	74	0.742 ± 0.084	0.025 ± 0.013	2.5
France	12	3	641	50	0.591 ± 0.108	0.014 ± 0.008	1.4
All samples	58	16	641	86	0.880 ± 0.023	0.048 ± 0.024	4.8
X. laevis rrnS							
Southwestern SA	30	7	369	13	0.662 ± 0.088	0.008 ± 0.005	0.8
Northeastern SA	16	3	369	2	0.342 ± 0.140	0.001 ± 0.001	0.1
France	12	2	369	5	0.303 ± 0.148	0.004 ± 0.003	0.4
All samples	58	9	369	14	0.681 ± 0.044	0.009 ± 0.005	0.9
P. xenopodis COX1							
Southwestern SA	30	20	416	94	0.959 ± 0.022	0.070 ± 0.035	7.0
Northeastern SA	16	15	416	89	0.992 ± 0.025	0.072 ± 0.037	7.2
France	12	3	416	3	0.591 ± 0.108	0.004 ± 0.003	0.4
All samples	58	34	416	110	0.958 ± 0.014	0.072 ± 0.035	7.2
P. xenopodis rrnS							
Southwestern SA	30	14	484	78	0.903 ± 0.036	0.050 ± 0.025	5.0
Northeastern SA	16	14	484	87	0.983 ± 0.028	0.055 ± 0.029	5.5
France	12	2	484	1	0.485 ± 0.106	0.001 ± 0.001	0.1
All samples	58	28	484	98	0.930 ± 0.020	0.048 ± 0.024	4.8

Abbreviations: SA, South Africa; n, sample size; H, number of haplotypes; L, length of sequence; S, number of polymorphic sites; Hd, haplotype diversity; π, nucleotide diversity; d, mean percentage of pairwise differences.

206 haplotypes of *COX1* and 35 of *rrnS*. Along with the outgroup *P. occidentalis*, the 45 aligned *COX1* sequences of 423 bp

207 and the *rrnS* sequences of 511 bp resulted in a concatenated alignment of 934 bp in length that was used to obtain the ML

208 phylogeny of the parasite. Of these sites, the 799 complete sites contained 225 sites that were polymorphic with a subset of

209 174 parsimony informative sites.

3.2 Host and parasite phylogeographic clustering

The MSN of the frog host showed three major haplotype clusters (Fig. 2a) that clearly corresponds to the three major lineages

revealed by the phylogenetic analysis of *X. lavis* (Fig. 3). The northeastern frog cluster, which corresponded to lineage NE

in the frog phylogeny, included frog specimens from SA4, SA5 and SA6 localities and France. This cluster exhibited a

star-like topology with two dominant native haplotypes and four satellite haplotypes that were separated from the dominant

haplotypes by one to seven mutations. The two southwestern frog clusters, which corresponds to SW_{coast} and SW_{inland} from

the frog phylogeny, included frogs from SA1, SA2, SA3, SA4, SA6 and SA7 localities and France and Portugal. One central

dominant haplotype in the southwestern clusters represented specimens from SA1 and SA3, as well as France and Portugal.

Satellite haplotypes from SA1, SA2, SA3, SA4 and SA7 fanned out from the dominant southwestern haplotype with a

star-like topology and, with the exception of a cluster of four highly divergent haplotypes that differed from the most frequent

haplotype by 36 to 50 mutations, the rest were separated from the central haplotype by one to seven mutations. The specimens

in the divergent southwestern cluster, which corresponds to the SW_{inland} lineage identified by the phylogenetic analysis, also

included a specimen collected at one of the SA6 northeastern localities (Wakefield). Furthernore, frog specimens from SA6

localities (Wakefield and Ugie) had similar haplotypes that fell between the northeastern and southeastern clusters, separated

from the nearest northeastern haplotype by 29 mutations and the nearest southwestern haplotype by 45 mutations. Frog

specimens from the SA4 localities were present in all three the clusters.

In the polystomatid MSN (Fig. 2b), a cluster of parasite specimens from northeastern hosts, that corresponds to **ne** in the

polystomatid phylogeny, represented nine of the sixteen specimens collected from the northeastern SA5 and SA6 localities.

The seven remaining parasite specimens from SA5 and SA6 localities (Adelaide, Dullstroom, Hopetown, Johannesburg, Ugie,

Verkykerskop and Wakefield) were spread across the remainder of the network, therefore clustering with polystomatids from

southwestern hosts. Moreover, parasites collected at the SA4 localities were spread across the entire network.

The polystomatids from France and Portugal clustered together as a peripheral dominant haplotype that included frogs

from the southwestern SA1 and SA3 localities with two satellite haplotypes that differed by two or four mutations, respectively.

FIGURE 2 Median spanning haplotype networks of the partial *COX1* and *rrnS* genes of **(a)** 59 *Xenopus laevis* specimens and **(b)** their corresponding 59 *Protopolystoma xenopodis* parasites. For frogs and their polystomatids collected in the native southern Africa (see inset map), colours refer to the phylogeographic lineages of the host (de Busschere *et al.*, 2016). Co-introduced *X. laevis* and polystomatids are from France (grey) and Portugal (pink). Highlighted haplotypes correspond to the clades revealed by the phylogenetic analyses (Fig. 3): northeastern host (NE) and parasite (ne) and southwestern host (SW) and parasite (sw) clades. Key localities with host lineage switches are named, as well as Jonkershoek, the source locality of most of the *X. laevis* exports. Sizes of circles are proportional to haplotype frequencies (see scale). The number of mutations is marked on the connecting branches.

233 This haplogroup of polystomatid specimens from the invasive localities, which corresponds to **sw4** of the polystomatid

234 phylogeny, was separated from the specimen from Jonkershoek (Fig. 1), the source of the majority of *X. laevis* exports, by

235 255 to 260 mutations.

3.3 Host and parasite phylogeny and congruence

Tree reconstruction showed that frogs from the northeastern SA5 and SA6 localities in the native range form a well-supported

clade (clade NE in Fig. 3 with UFboot2 = 95% / SH-aLRT = 95.8%). In addition, two clades with unresolved relationships

to one another and to NE could be identified among sampled frogs from the southwestern localities in the native range of

X. laevis (clades SW_{coast} with UFboot2 = 81% / SH-aLRT = 86.1% and SW_{inland} with UFboot2 = 98% / SH-aLRT = 99.7% in

Fig. 3). Members of the first clade originate from localities closer to the coastline (clade SW_{coast} in Fig. 3 from SA1, SA2,

SA3 and SA7 localities) than the second one (clade SW_{inland} in Fig. 3 from SA3, SA4 and SA6 localities).

The two SA4 localities (near Three Sisters in the Karoo) are suggested as a contact zone between NE, SW_{coast} and

SW_{inland}, with sampled individuals from this area present in all three host lineages. Likewise, frogs from an SA6 locality

(Wakefield near Lesotho) were also present in both NE and SW_{inland}.

The non-native frog from the invasive population in Portugal, along with one of the non-native frogs from western France,

grouped together with the native frogs in SW_{coast} in the host phylogeny, thus placing these invasive frogs in the same clade as

the frog from Jonkershoek (Fig. 1), the source of the majority of *X. laevis* exports. Nonetheless, the invasive population of

X. laevis in France also harboured several frogs that clustered with frogs from the native northeastern localities in the NE

clade.

In the same manner as the host phylogeny, the ML inference of the *P. xenopodis* tree revealed a single, well-supported

clade of polystomatids from hosts from the northeastern SA5 and SA6 localities and the SA4 contact zone (clade **ne** in Fig. 3

with UFboot2 = 97% / SH-aLRT = 87.6%). Several clades basal to the **ne** group emerged among the polystomatids from

hosts collected at the southwestern SA1, SA2, SA3 and SA7 localities and the SA4 contact zone (clades **sw1**, **sw2**, **sw3**

and **sw4** in Fig. 3). The marked phylogenetic divergence of the polystomatid was supported by significant morphological

differences between polystomatids from the southwest of South Africa and the northeast of South Africa (see Appendix S3

for morphometrical methods and findings).

The earliest diverging clade is a well-supported grouping of polystomatids collected from SA1, SA3 and SA7 (clade **sw1**

in Fig. 3 with UFboot2 = 94%/SH-alrt = 88.1%). Two other native southwestern clades included a clade of polystomatids

from the SA3 localities (clade **sw2** in Fig. 3 with UFboot2 = 53%/SH-alrt = 90.7%) and another clade from a combination

of SA2 and SA3 localities (clade **sw3** in Fig. 3 with UFboot2 = 61% / SH-aLRT = 86.6%). Another well-supported clade,

FIGURE 3 The interaction network and phylogenies of the frog *Xenopus laevis* (left) and its polystomatid *Protopolystoma xenopodis* (right) with closely related *X. muelleri* and *P. occidentalis* as outgroups. Topologies are based on the maximum likelihood analyses of the concatenated *COX1* and *rrnS* gene alignments of 31 hosts and 45 parasites with low support clades (UFboot2 < 95% or SH-alrt < 80%) collapsed. Links indicate the observed host-parasite associations, weighted according to the contribution of each interaction to the overall phylogenetic congruence with thicker lines corresponding to greater support. Colours refer to geographic origin, following the host phylogeography of de Busschere *et al.* (2016): SA1 (dark blue), SA2 (light blue), SA3 (red), SA4 (green), SA5 (yellow), SA6 (brown) and SA7 (purple) (see Fig. 1) with main lineages named and highlighted. Invasive *X. laevis* and co-introduced *P. xenopodis* from France (grey) and Portugal (pink) are also included. Key localities are Adelaide (A), Dullstroom (D), Hopetown (H), Johannesburg (J), Ugie (U), Verkykerskop (V) and Wakefield (W). Jonkershoek (☆) is the origin of the majority of *X. laevis* exports. Scale bars indicate 0.1 nucleotide substitutions per site for the two phylogenies.

the sister clade to **ne**, contained polystomatids from SA1, France and Portugal (clade **sw4** in Fig. 3 with UFboot2 = 100%

/ SH-aLRT = 100%). However, overall low support values of many internal nodes in the parasite ML tree hampered the

interpretation of phylogenetic clustering among some of the polystomatids collected from the southwestern SA1, SA2, SA3,

SA4 and SA7 localities.

A few links represented mismatches between the host and parasite phylogenies that could be explained by long distance

translocation of frogs from the SW host lineages to the northeastern localities followed by spill-over of **sw1**, **sw2** and **sw3**

polystomatids to NE hosts. These indications of host switching were identified in frog-polystomatid pairs from SA5 and SA6

localities (Dullstroom, Hopetown, Johannesburg, Verkykerskop, Adelaide, Ugie and Wakefield).

The phylogeny of *X. laevis* exhibited significant overall congruence with that of *P. xenopodis* (PACo m_{xy}^2 = 0.1034, *p* =

0.0018, n = 10000; ParaFitGlobal = 0.0202, *p* < 0.0001, n = 10000), with the PACo analysis suggesting that the phylogeny

of *P. xenopodis* tracked that of *X. laevis*. According to PACo, links between frogs from SW$_{coast}$ and their corresponding

polystomatids that fall within **sw1** in the parasite phylogeny, generally contributed the most to the overall phylogenetic

congruence. The PACo jackknife procedure further identified the interactions between frog–polystomatid pairs from the

invasive range in France and Portugal as links conveying low support to congruence. The ParaFitLink tests reported 30 out of

48 links as significant at a level of 0.05. Out of the 18 links that did not significantly support congruence, 11 signified host

switches by parasites to another host lineage that occurred in the invasive range of France and Portugal and at SA5 and SA6

localities in the native range (Johannesburg, Three Sisters, Ugie and Wakefield).

279 4 DISCUSSION

In this study, the widespread, yet previously hidden, domestic expansion of *X. laevis* lineages is brought to light by the

intraspecific phylogeography of its co-invasive parasite, *P. xenopodis*, a participant in the invasion history that is often

forgotten. The use of parasite data to reveal host connectivity is hardly novel (reviewed in Gagne *et al.*, 2021). Yet, the

present study is the first application of host-parasite co-phylogeographic analyses to the detection of intraspecific cryptic

invasions. The study is not only a proof of concept that parasite data can provide additional information to trace biological

invaders on the intraspecific level but it also demonstrates how parasite data can illuminate invasion pathways, thus informing

management strategies.

287 4.1 Parasite clustering mirrors the intraspecific phylogeography of the host

The phylogeographic analyses of the parasite *P. xenopodis*, together with morphometric comparisons, reveal marked

intraspecific geographic structuring across the native range of the species. The observed structuring in the parasite corroborates

and expounds upon the phylogeography of the host *X. laevis* which also shows notable phylogeographic structure across

its native range (de Busschere *et al.*, 2016; du Preez *et al.*, 2009; Furman *et al.*, 2015; Grohovaz *et al.*, 1996; Measey &

Channing, 2003). Yet, the parasite data set provides an independent line of evidence and not a mere duplication of the

phylogeographic structure of the frog, since each parasite represents a separate infection event from a pool of free-swimming

larvae. Furthermore, with the exception of a brief free-swimming period that allows host switches to occur, *P. xenopodis*

completes the whole of its life cycle in the excretory system of *X. laevis* (Theunissen *et al.*, 2014), which translates to a very

specific and intimate host-parasite association that can faithfully portray the phylogeographic structure of the frog. Therefore, based upon the phylogenetic congruence between the host and parasite, the phylogeography of the parasite can provide insight on the translocation of the host.

The genetically distinct host-parasite clusters display distributions that roughly follow the two climatic regimes in the region, namely the summer rainfall region to the northeast and the winter rainfall region to the southwest, with the plateau edge of the Great Escarpment and the many ridges of the Cape Fold Mountains as assumed geographical barriers to natural dispersal (Furman *et al.*, 2015). This divergence is further echoed by the significant morphological differences between polystomatids from the northernmost and the southernmost localities. The more pronounced clustering among hosts and parasites and the higher haplotype diversity in hosts that hail from the southwest of the Great Escarpment is likely brought about by the presence of the many ridges of the Cape Fold Mountains (de Busschere *et al.*, 2016; du Preez *et al.*, 2009; Furman *et al.*, 2015; Grohovaz *et al.*, 1996; Measey & Channing, 2003). On the other hand, well-documented human-mediated translocation of *X. laevis* to the southwest of the Great Escarpment (van Sittert & Measey, 2016; Weldon *et al.*, 2007), in conjunction with widespread overland migration of the frog with the help of the many interconnected artificial water bodies in the region (de Villiers & Measey, 2017; Measey, 2016), ensure some gene flow – and exchange of parasites – between the divergent frog populations in this area.

4.2 Parasite spillover is an indication of contact between divergent host lineages

Parasite phylogeography offers a tool to recover historical insights of host dispersal pathways, since host lineage switches reflect past interactions between host organisms at various time scales (e.g. Barbosa *et al.*, 2012; Criscione *et al.*, 2006; Galbreath & Hoberg, 2015; Galbreath *et al.*, 2020; Nieberding *et al.*, 2004). The usefulness of parasite species in reflecting the introduction pathways of their hosts has been demonstrated for fish hosts and their monogenean parasites on a few occasions (Huyse *et al.*, 2015; Kmentová *et al.*, 2019; Ondračková *et al.*, 2012; Reyda *et al.*, 2020). However, these studies focused on species-level invasions from the native to the non-native range, rather than intraspecific invasions of different lineages within the native range. Our study, on the other hand, demonstrates the use of parasites as tags to trace intraspecific cryptic invasions. Multiple instances of host lineage switches revealed by the phylogeographic analyses of *P. xenopodis* act as an indicator for contact between phylogeographically distinct *X. laevis* lineages. This provides us with the means of recognising contact between *X. laevis* lineages even in cases where the locally non-native host lineage may have gone extinct after the introduction event or where they are not detected at a certain sampling effort.

In fact, thus far, genetic evidence for the admixture of the lineages of *X. laevis* with native ranges on opposite sides of the

Great Escarpment has not been found, despite historical records pointing towards large scale exports from the southwest to

the rest of southern Africa (van Sittert & Measey, 2016; Weldon *et al.*, 2007). In our study, the host phylogeographic analyses

suggest the presence of contact zones between the northeastern (NE) and southwestern lineages (SW$_{inland}$ & SW$_{coast}$) at

two localities to the northeast of the Great Escarpment (Fig. 2). Since these zones are situated to the north of a predicted

geographical barrier, namely the plateau edge of the Great Escarpment (Furman *et al.*, 2015), one could surmise that

anthropogenic translocation had a hand in bringing about this sympatry of multiple clades of *X. laevis*, pointing towards more

recent translocation events.

Moreover, it is this sympatry of multiple host clades at localities that makes possible host lineage switching by parasite

lineages. In turn, host switches provides more evidence for the translocation of *X. laevis* across southern Africa. Indeed,

our study confirmed the presence of southwestern parasite lineages (**sw1**, **sw2** & **sw3**) at seven localities to the northeast of

the Great Escarpment, where one would expect the northeastern parasite lineage (**ne**) to occur (Fig. 3). In these instances,

southwestern parasite lineages that are translocated along with their respective host lineages, thus locally non-native at these

new localities, spill over to the locally native northwestern host lineage when these divergent, previously allopatric host

lineages come into contact (Galbreath & Hoberg, 2015).

Differential survival between locally native and non-native *X. laevis* lineages in southern Africa can partially explain the

fewer observed instances of establishment by non-native *X. laevis* lineages when compared to its co-introduced polystomatid.

In translocation experiments of different lineages in the native range, *X. laevis* tadpoles and metamorphs exhibited remarkable

phenotypic plasticity when translocated between rainfall regimes at the cost of longterm survival (Kruger, 2020). Under

these circumstances, the locally native *X. laevis* lineage would outcompete the non-native lineage, preventing successful local

invasion. Thus, the employment of host phylogeography alone to trace long range translocation of *X. laevis* in southern Africa

would mask a number of introduction events.

Upon arrival in the new area, the co-introduced *P. xenopodis* lineage may rapidly spread in a population of locally native

and therefore naive *X. laevis*. It has been shown that primary infections by *P. xenopodis* in previously uninfected *X. laevis*

hosts elicit strong long term immunological responses against secondary infections by the same parasite species (Jackson &

Tinsley, 2001). Correspondingly, a cross-infectivity experiment between sympatric and allopatric *X. laevis* and *P. xenopodis*

lineages further suggested that *X. laevis* lineages in their native ranges might be more susceptible to infection by the locally

non-native *P. xenopodis* lineage than to their corresponding locally native parasite lineage (Jackson & Tinsley, 2005). In turn,

this effect will boost the nascent spread of non-native *P. xenopodis* lineages in the locally native lineage of *X. laevis*.

The rural locations of the parasite spillover events revealed by our analyses of polystomatid phylogeography suggest that

smaller scale human-mediated dispersal events, as opposed to large official consignments from the southwestern region to

centers of higher learning across South Africa, have far-reaching effects. The role of recreational fishing across South Africa

in the dispersal of *X. laevis*, and by extension, *P. xenopodis*, has been recognised before (Weldon *et al.*, 2007). Previously

documented interviews with local anglers and suppliers exposed the use of *X. laevis*, frequently bought from pet shops and

local breeders, as live bait for predominantly catfish angling (Weldon *et al.*, 2007). Our own informal discussions with local

anglers at the localities where we collected *X. laevis* affirm the widespread use of *X. laevis* as live bait for the capture of both

black bass (*Micropterus* spp.) and catfish (*Clarias gariepinus*).

In terms of sheer numbers, the impact of these translocation events on the domestic dispersal of *X. laevis* has been deemed

negligible when compared to exports from Jonkershoek, the most prolific exporter, and other official suppliers to the domestic

market (van Sittert & Measey, 2016; Weldon *et al.*, 2007). Nonetheless, previous studies on other species have pointed out

that the use of live bait in angling is an important pathway for introductions, even if the market might be small in economic

terms (Kalous *et al.*, 2013; Kilian *et al.*, 2012). Therefore, the impact of anglers on the spread of locally non-native *X. laevis*

lineages to more remote areas, as opposed to the bigger centra that received the documented official consignments, should not

be underestimated. Our informal interviews identified a myriad of ways for *X. laevis* to be released in new environments by

anglers, from escapees *en route*, to the deliberate release of surplus bait, to the intentional stocking of water bodies for future

use.

4.3 Parasite phylogeography as an underutilised tool in invasion biology

As shown above, the present study acts as a proof of concept for the potential of the co-phylogeographic approach to the

clarification of patterns in intraspecific host divergence and dispersal. In turn, this independent line of evidence can provide

additional insight on invasion pathways and inform management strategies in profound ways. Here, we highlight some of the

potential implications to demonstrate the broader applicability of this integrative approach.

Our approach was able to provide a new perspective on the origin of the relatively high haplotype diversity of the invasive

population of *X. laevis* in France, recovered in our analysis and others, when compared to other invasive populations in

Portugal, Sicily and Chile (de Busschere *et al.*, 2016; Lillo *et al.*, 2013; Lobos *et al.*, 2014). The idea of multiple introduction

events from various source populations in South Africa has been offered before as the most parsimonious explanation for

this genetic diversity in the invasive population of *X. laevis* in western France (de Busschere *et al.*, 2016). Since we found

evidence for host switches by *P. xenopodis* in the native range, an event that necessarily involves the sympatry of divergent

host lineages, it is a certainty that multiple host lineages co-occur at certain localities in the native range. In this, our results

build upon the evidence provided by historical export records (van Sittert & Measey, 2016; Weldon *et al.*, 2007), dispersal

patterns (de Villiers & Measey, 2017; Measey, 2016) and anecdotal information (Weldon *et al.*, 2007) to show that the mixing

of lineages most likely already happened in the source populations in South Africa prior to the co-introduction of *X. laevis*

and *P. xenopodis* to western France (van Sittert & Measey, 2016; Weldon *et al.*, 2007).

The novel insights provided by the addition of the parasite data set has implications for both the conservation of *X. laevis*

and its host-specific parasites. The widespread anthropogenic translocation of this host-parasite system disrupts interactions

between *X. laevis* lineages and the *P. xenopodis* lineages that tracked their phylogenetic divergence, ancient interactions that

deserve to be targets of conservation (Gómez & Nichols, 2013). Subsequent host switches between closely related lineages

set the stage for new diversity to arise and for novel ecological interactions to be established (Galbreath *et al.*, 2020; Sures

et al., 2017). In the case of host-specific parasites, translocation of lineages within the native range of a species may actually

be of more concern than translocation to far-flung non-native ecosystems, since these ecosystems will probably not harbour

close relatives of the invasive host to facilitate host switches (Blackburn & Ewen, 2017). Host-parasite interactions are

also something to keep in mind when conducting conservation re-introduction programmes, where captive-bred or locally

non-native individuals of a species are used to re-stock locally extinct populations of endangered species (Britt *et al.*, 2004).

Our results highlight that the evaluation of the genetic and parasitological suitability of a source population for translocation

should also encompass the potential history of local introductions that may have influenced the intraspecific parasitism of the

host population (Jørgensen, 2015; Northover *et al.*, 2018).

This approach further holds great promise in the management of invasive species. Prior to invasion, anthropogenic

translocations in the native range may have already altered the source populations of invasive species, as is the case of

X. laevis. We need to take altered phylogeographic structure into account to adequately monitor the course of an invasion or

the novel interactions that may arise. Parasites as passengers or invaders themselves easily slip under the radar in invasion

biology (Blackburn & Ewen, 2017; Solarz & Najberek, 2017). Yet, the role of parasites on the move in emerging infectious

diseases and habitat transformation via indirect density-mediated effects cannot be overemphasised (Amundsen *et al.*, 2013;

Bobadilla-Suarez *et al.*, 2017; Daszak, 2000; Dunn *et al.*, 2012; Goedknegt *et al.*, 2016; Lymbery *et al.*, 2014; Roy &

Lawson Handley, 2012; Sures *et al.*, 2017). Furthermore, the identification of the local angling community as a major player

in the translocation of *X. laevis* in the native range allows policy makers to focus their intervention efforts in the management

of *X. laevis* as a domestic exotic.

4.4 Recommendations to make parasite data sets work for invasion biologists

All in all, our findings agree with the necessity of a more holistic approach to data collection in invasion biology (Clavero

et al., 2016), especially with regard to parasites. The idea of the "holistic specimen" has recently gained some traction (Cook

et al., 2017; Galbreath *et al.*, 2019). This approach to the collection and study of organisms advocates for the integrated

specimen-based studies of animals and their associated parasites and pathogens. The optimisation of sampling effort is crucial

to improve the allocation of resources, both in terms of funding and specimen use, especially when working with endangered

species or specimens that require considerable effort to obtain, or in view of other ethical concerns.

The inclusion of parasite data during the data collection phase also ensures that the host and parasite data sets are comprised

of directly comparable individuals which greatly improves the feasibility and value of comparative phylogeographic analyses.

For example, the non-overlap between host and parasite data sets was cited as a limitation in the co-phylogeographic analyses

conducted by both Barbosa *et al.* (2012) and Huyse *et al.* (2017). Even in cases where parasites are not the targets of the

current project, the deposition of voucher material of both host and parasite into curated institutional collections, in other

words overlapping data sets, can be vital for later studies (Galbreath *et al.*, 2019; Thompson *et al.*, 2021).

Another aspect to keep in mind is the choice of markers for co-phylogeographic analyses. In general, it is regarded as best

practice to employ orthologous molecular markers in co-phylogenetic analyses (de Vienne *et al.*, 2013). Specifically, the

use of orthologous markers for host and parasite in co-phylogeographic analyses to answer questions in invasion biology

may have the benefit of making the findings from the two lines of evidence more directly comparable. The independent

yet comparable DNA signature provided by the parasite data set may ameliorate inference based upon the phylogeographic

structure of the host data set that is often blurred by the recent mixing of previously isolated genotypes in invasive populations

(Rius & Turon, 2020).

Ultimately, this study serves as a proof of concept that the use of parasite data sets will add value to investigations in

invasion biology, especially in the case of intraspecific cryptic invasions. In fact, we show that the utility of parasite data

sets can go beyond the provision of supporting evidence and provide greater resolution that may lead to elucidation of

patterns that went undetected in the host data set. Our study echoes the sentiments of Hoberg *et al.* (2015) who argued for an

integrated approach to parasitology as a discipline that holds the keys to understanding and managing global change in the

Anthropocene.

DATA AVAILABILITY STATEMENT

All DNA sequences used for co-phylogeographic analyses will be made available on the public database GenBank, along with relevant metadata and accession numbers will be added upon article acceptance. Detailed locality data and data for additional morphometrical analyses are contained in the supplementary information that can be made available upon request.

ACKNOWLEDGMENTS

ALS received funding from the DST-NRF Centre of Excellence for Invasion Biology (South Africa). The project utilised the infrastructure of the Unit for Environmental Sciences and Management, North-West University (South Africa). This research was supported by EMBRC Belgium – FWO project GOH3817N. MPMV is financed by the Special Research Fund of Hasselt University (BOF20TT06). The authors would like to express their sincere thanks to a number of persons who assisted in the collection of the frogs. Guénolé le Peutrec, Jean Secondi, Natasha Kruger and Rodolphe Olivier helped with the collection in France. In South Africa, several farm owners graciously gave permission for collection to take place on their properties and provided lodging for the research team: Fanus and Olga Kritzinger, William and Christa van Zyl, Dave Schlebusch, Fanus and Carin Oberholzer, Danie and Annalise Marais, Johan Hamann, Tobie Bielt, Gert Bench, Stoffel Labuschagne, Jannie and Susan van Rensburg, Jan Meintjies, Marthinus Hartman, Douw and Louise de Jager and Ernest de Villiers. Lastly, Mathys Schoeman, Clarke Scholtz, Andrea Darvall, Willie Landman, Ferdi de Lange, John Measey and Roxanne Viviers assisted with the collection of frogs at the remainder of the localities in South Africa. The authors are also indebted to numerous local recreational fishermen who were willing to share details about their use of *Xenopus laevis* across South Africa. The utilisation of the frogs and the research protocols were approved by the Animal Care, Health and Safety in Research Ethics (AnimCare) Committee of the Faculty of Health Sciences of the North-West University (ethics number: NWU-0380-16-A5-01). Animals from the South African populations were sampled under the permits 0056-AAA007-00224 (CapeNature) and FAUNA 1343-2017 (Northern Cape) provided by the Department of Economic Development, Environmental Affairs and Tourism.

CONFLICT OF INTEREST

All authors declare no conflict of interest.

AUTHORS' CONTRIBUTIONS

ALS conceived the initial idea, collected and analysed the data and led the writing of the manuscript. All authors refined the initial concept, designed the methodology, contributed to interpretation of data, contributed critically to the drafts and gave final approval for publication.

REFERENCES

Ahyong, S. T., Kupriyanova, E., Burghardt, I., Sun, Y., Hutchings, P. A., Capa, M. & Cox, S. L. 2017. Phylogeography of the invasive Mediterranean fan worm, *Sabella spallanzanii* (Gmelin, 1791), in Australia and New Zealand. *Journal of the Marine Biological Association of the United Kingdom*, 97:985–991. https://doi.org/10.1017/S0025315417000261

Amaro, R. C., Nunes, I., Canedo, C., Napoli, M. F., Junca, F. A., Verdade, V. K., ... Rodrigues, M. T. 2013. A molecular phylogeny recovers *Strabomantis aramunha* Cassimiro, Verdade and Rodrigues, 2008 and *Haddadus binotatus* (Spix, 1824) (Anura: Terrarana) as sister taxa. *Zootaxa*, 3741:S569–S582. https://doi.org/10.11646/zootaxa.3741.4.7

Amundsen, P. A., Lafferty, K. D., Knudsen, R., Primicerio, R., Kristoffersen, R., Klemetsen, A. & Kuris, A. M. 2013. New parasites and predators follow the introduction of two fish species to a subarctic lake: implications for food-web structure and functioning. *Oecologia*, 171:993–1002. https://doi.org/10.1007/s00442-012-2461-2

Balbuena, J. A., Miguez-Lozano, R. & Blasco-Costa, I. 2013. PACo: a novel procrustes application to cophylogenetic analysis. *PLOS One*, 8, e61048. https://doi.org/10.1371/journal.pone.0061048

Baldwin, R. E., Banks, M. A. & Jacobson, K. C. 2012. Interating fish and parasite data as a holistic solution for identifying the elusive stock structure of Pacific sardines (*Sardinops sagax*). *Reviews in Fish Biology and Fisheries*, 22:137–156. https://doi.org/10.1007/s11160-011-9227-5

Bandelt, H.-J., Forster, P. & Röhl, A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution*, 16:37–48. https://doi.org/10.1093/oxfordjournals.molbev.a026036

Barbosa, A. M., Thode, G., Real, R., Feliu, C. & Vargas, J. M. 2012. Phylogeographic triangulation: using predator-prey-parasite interactions to infer population history from partial genetic information. *PLoS One*, 7, e50877. https://doi.org/10.1371/journal.pone.0050877

Blackburn, T. M., Bellard, C. & Ricciardi, A. 2019. Alien versus native species as drivers of recent extinctions. *Frontiers in Ecology and the Environment*, 17:203–207. https://doi.org/10.1002/fee.2020

Blackburn, T. M. & Ewen, J. G. 2017. Parasites as drivers and passengers of human-mediated biological invasions. *EcoHealth*, 14:61–73. https://doi.org/10.1007/s10393-015-1092-6

Blackburn, T. M., Pyšek, P., Bacher, S., Carlton, J. T., Duncan, R. P., Jarošík, V., ... Richardson, D. M. 2011. A proposed unified framework for biological invasions. *Trends in Ecology & Evolution*, 26:333–339. https://doi.org/10.1016/j.tree.2011.03.023

Blakeslee, A. M. H., Byers, J. E. & Lesser, M. P. 2008. Solving cryptogenic histories using host and parasite molecular genetics: the resolution of *Littorina littorea*'s North American origin. *Molecular Ecology*, 17:3684–3696. https://doi.org/10.1111/j.1365-294x.2008.03865.x

Bobadilla-Suarez, M., Ewen, J. G., Groombridge, J. J., Beckmann, K., Shotton, J., Masters, N., ... Sainsbury, A. W. 2017. Using qualitative disease risk analysis for herpetofauna conservation translocations transgressing ecological and geographical barriers. *EcoHealth*, 14:47–60. https://doi.org/10.1007/s10393-015-1086-4

Britt, A., Welch, C., Katz, A., Iambana, B., Porton, I., Junge, R., ... Haring, D. 2004. The re-stocking of captive-bred ruffed lemurs (*Varecia variegata variegata*) into the Betampona Reserve, Madagascar: methodology and recommendations. *Biodiversity and Conservation*, 13:635–657. https://doi.org/10.1023/b:bioc.0000009497.24917.ae

Cailliez, F. 1983. The analytical solution of the additive constant problem. *Psychometrika*, 48:305–310. https://doi.org/10.1007/bf02294026

Capinha, C., Essl, F., Seebens, H., Moser, D. & Pereira, H. M. 2015. The dispersal of alien species redefines biogeography in the Anthropocene. *Science*, 348:1248–1251. https://doi.org/10.1126/science.aaa8913

Chernomor, O., von Haeseler, A. & Minh, B. Q. 2016. Terrace aware data structure for phylogenomic inference from supermatrices. *Systematic Biology*, 65:997–1008. https://doi.org/10.1093/sysbio/syw037

Clavero, M., Nores, C., Kubersky-Piredda, S., & Centeno-Cuadros, A. 2016. Interdisciplinarity to reconstruct historical introductions: solving the status of cryptogenic crayfish. *Biological Reviews*, 91:1036–1049. https://doi.org/10.1111/brv.12205

Cook, J. A., Galbreath, K. E., Bell, K. C., Campbell, M. L., Carrière, S., Colella, J. P., ... Hoberg, E. P. 2017. The Beringian Coevolution Project: holistic collections of mammals and associated parasites reveal novel perspectives on evolutionary and environmental change in the North. *Arctic Science*, 3:585–617. https://doi.org/10.1139/as-2016-0042

Criscione, C. D., Cooper, B. & Blouin, M. S. 2006. Parasite genotypes identify source populations of migratory fish more accurately than fish genotypes. *Ecology*, 87:823–828. https://doi.org/10.1890/0012-9658(2006)87[823:pgispo]2.0.co;2

Daszak, P. 2000. Emerging infectious diseases of wildlife – threats to biodiversity and human health. *Science*, 287:443–449. https://doi.org/10.1126/science.287.5452.443

de Busschere, C., Courant, J., Herrel, A., Rebelo, R., Rödder, D., Measey, G. J. & Backeljau, T. 2016. Unequal contribution of native South African phylogeographic lineages to the invasion of the African clawed frog, *Xenopus laevis*, in Europe. *PeerJ*, 4, e1659. https://doi.org/10.7717/peerj.1659

de Vienne, D. M., Refregier, G., Lopez-Villavicencio, M., Tellier, A., Hood, M. E. & Giraud, T. 2013. Cospeciation vs host-shift speciation: methods for testing, evidence from natural associations and relation to coevolution. *New Phytologist*, 198:347–385. https://doi.org/10.1111/nph.12150

de Villiers, F. A. & Measey, G. J. 2017. Overland movement in African clawed frogs (*Xenopus laevis*): empirical dispersal data from within their native range. *PeerJ*, 5, e4039. https://doi.org/10.7717/peerj.4039

Dlugosch, K. M. & Parker, I. M. 2008. Founding events in species invasions: genetic variation, adaptive evolution, and the role of multiple introductions. *Molecular Ecology*, 17:431–449. https://doi.org/10.1111/j.1365-294X.2007.03538.x

du Preez, L. H., Kunene, N., Hanner, R., Giesy, J. P., Solomon, K. R., Hosmer, A. & van der Kraak, G. J. 2009. Population-specific incidence of testicular ovarian follicles in *Xenopus laevis* from South Africa: a potential issue in endocrine testing. *Aquatic Toxicology*, 95:10–16. https://doi.org/10.1016/j.aquatox.2009.07.018

Dufresnes, C., Santo, L. D., Leuenberger, J., Schuerch, J., Mazepa, G., Grandjean, N., ... Dubey, S. 2017. Cryptic invasion of Italian pool frogs (*Pelophylax bergeri*) across Western Europe unraveled by multilocus phylogeography. *Biological Invasions*, 19:1407–1420. https://doi.org/10.1007/s10530-016-1359-z

Dunn, A. M., Torchin, M. E., Hatcher, M. J., Kotanen, P. M., Blumenthal, D. M., Byers, J. E., ... Fox, C. 2012. Indirect effects of parasites in invasions. *Functional Ecology*, 26:1262–1274. https://doi.org/10.1111/j.1365-2435.2012.02041.x

Evans, B. J., Carter, T. F., Tobias, M. L., Kelley, D. B., Hanner, R. & Tinsley, R. C. 2008. A new species of clawed frog (genus

Xenopus) from the Itombwe Massif, Democratic Republic of the Congo: implications for DNA barcodes and biodiversity

conservation. *Zootaxa*, 1780:55–68. https://doi.org/10.11646/zootaxa.1780.1.2

Evans, B. J., Kelley, D. B., Tinsley, R. C., Melnick, D. J. & Cannatella, D. C. 2004. A mitochondrial DNA phylogeny of

African clawed frogs: phylogeography and implications for polyploid evolution. *Molecular Phylogenetics and Evolution*,

33:197–213. https://doi.org/10.1016/j.ympev.2004.04.018

Excoffier, L. & Lischer, H. E. L. 2010. ARLEQUIN suite ver 3.5: a new series of programs to perform population genetics

analyses under Linux and Windows. *Molecular Ecology Resources*, 10:564–567. https://doi.org/10.1111/j.1755-0998.2010.

02847.x

Fick, S. E. & Hijmans, R. J. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas.

International Journal of Climatology, 37:4302–4315. https://doi.org/10.1002/joc.5086

Fischer, W., Koch, W. & Elepfandt, A. 2000. Sympatry and hybridization between the clawed frogs *Xenopus laevis laevis* and

Xenopus muelleri (Pipidae). *Journal of Zoology*, 252:99–107. https://doi.org/10.1111/j.1469-7998.2000.tb00824.x

Fitzpatrick, B., Fordyce, J., Niemiller, M. & Reynolds, R. G. 2012. What can DNA tell us about biological invasions?

Biological Invasions, 14:245–253. https://doi.org/10.1007/s10530-011-0064-1

Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. 1994. DNA primers for amplification of mitochondrial

cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*,

3:294–299

Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. 2012. CD-HIT: accelerated for clustering the next generation sequencing data.

Bioinformatics, 28:3150–3152. https://doi.org/10.1093/bioinformatics/bts565

Furman, B. L., Bewick, A. J., Harrison, T. L., Greenbaum, E., Gvozdik, V., Kusamba, C. & Evans, B. J. 2015. Pan-

African phylogeography of a model organism, the African clawed frog *Xenopus laevis*. *Molecular Ecology*, 24:909–925.

https://doi.org/10.1111/mec.13076

Gagne, R. B., Crooks, K., Craft, M. E., Chiu, E. S., Fountain-Jones, N. M., Malmberg, J. L., ... VandeWoude, S. 2021.

Parasites as conservation tools. *Conservation Biology*. https://doi.org/10.1111/cobi.13719

Galbreath, K. E. & Hoberg, E. P. 2015. Host responses to cycles of climate change shape parasite diversity across North America's Intermountain West. *Folia Zoologica*, 64:218–232. https://doi.org/10.25225/fozo.v64.i3.a4.2015

Galbreath, K. E., Hoberg, E. P., Cook, J. A., Armién, B., Bell, K. C., Campbell, M. L., ... Hope, A. G. 2019. Building an integrated infrastructure for exploring biodiversity: field collections and archives of mammals and parasites. *Journal of Mammalogy*, 100:382–393. https://doi.org/10.1093/jmammal/gyz048

Galbreath, K. E., Toman, H. M., Li, C. & Hoberg, E. P. 2020. When parasites persist: tapeworms survive host extinction and reveal waves of dispersal across Beringia. *Proceedings of the Royal Society B: Biological Sciences*, 287, art. 20201825. https://doi.org/10.1098/rspb.2020.1825

Goedknegt, M. A., Feis, M. E., Wegner, K. M., Luttikhuizen, P. C., Buschbaum, C., Camphuysen, K. C. J., ... Thieltges, D. W. 2016. Parasites and marine invasions: ecological and evolutionary perspectives. *Journal of Sea Research*, 113:11–27. https://doi.org/10.1016/j.seares.2015.12.003

Gómez, A. & Nichols, E. 2013. Neglected wild life: parasitic biodiversity as a conservation target. *International Journal for Parasitology: Parasites and Wildlife*, 2:222–227. https://doi.org/10.1016/j.ijppaw.2013.07.002

Grohovaz, G. S., Harley, E. & Fabian, B. 1996. Significant mitochondrial DNA sequence divergence in natural populations of *Xenopus laevis* (Pipidae) from South Africa. *Herpetologica*, 52:247–253. https://doi.org/10.1080/11250003.2013.847502

Gu, X., Fu, Y. X. & Li, W. H. 1995. Maximum likelihood estimation of the heterogeneity of substitution rate among nucleotide sites. *Molecular Biology and Evolution*, 12:546–557. https://doi.org/10.1093/oxfordjournals.molbev.a040235

Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology*, 59:307–321. https://doi.org/10.1093/sysbio/syq010

Guo, Q. & Ricklefs, R. E. 2010. Domestic exotics and the perception of invasibility. *Diversity and Distributions*, 16:1034–1039. https://doi.org/10.1111/j.1472-4642.2010.00708.x

Hahn, C., Weiss, S. J., Stojanovski, S. & Bachmann, L. 2015. Co-speciation of the ectoparasite *Gyrodactylus teuchis* (Monogenea, Platyhelminthes) and its salmonid hosts. *PLOS One*, 10, e0127340. https://doi.org/10.1371/journal.pone.0127340

Héritier, L., Badets, M., du Preez, L. H., Aisien, M. S., Lixian, F., Combes, C. & Verneau, O. 2015. Evolutionary processes involved in the diversification of chelonian and mammal polystomatid parasites (Platyhelminthes, Monogenea, Polystomatidae) revealed by palaeoecology of their hosts. *Molecular Phylogenetics and Evolution*, 92:1–10. https://doi.org/10.1016/j.ympev.2015.05.026

Hiller, A. & Lessios, H. A. 2017. Phylogeography of *Petrolisthes armatus*, an invasive species with low dispersal ability. *Scientific Reports*, 7, 3359. https://doi.org/10.1038/s41598-017-03410-8

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. 2018. UFBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution*, 35:518–522. https://doi.org/10.1093/molbev/msx281

Hoberg, E. P., Agosta, S. J., Boeger, W. A. & Brooks, D. R. 2015. An integrated parasitology: revealing the elephant through tradition and invention. *Trends in Parasitology*, 31:128–133. https://doi.org/10.1016/j.pt.2014.11.005

Huang, Y., Niu, B., Gao, Y., Fu, L. & Li, W. 2010. CD-HIT Suite: a web server for clustering and comparing biological sequences. *Bioinformatics*, 26:680–682. https://doi.org/10.1093/bioinformatics/btq003

Hutchinson, M. C., Cagua, E. F., Balbuena, J. A., Stouffer, D. B., Poisot, T. & Fitzjohn, R. 2017. paco: implementing Procrustean approach to cophylogeny in R. *Methods in Ecology and Evolution*, 8:932–940. https://doi.org/10.1111/2041-210x.12736

Huyse, T., Oeyen, M., Larmuseau, M. H. D. & Volckaert, F. A. M. 2017. Co-phylogeographic study of the flatworm *Gyrodactylus gondae* and its goby host *Pomatoschistus minutus*. *Parasitology International*, 66:119–125. https://doi.org/10.1016/j.parint.2016.12.008

Huyse, T., Vanhove, M. P. M., Mombaerts, M., Volckaert, F. A. M. & Verreycken, H. 2015. Parasite introduction with an invasive goby in Belgium: double trouble? *Parasitology Research*, 114:2789–2793. https://doi.org/10.1007/s00436-015-4544-6

Ivanova, N. V., Dewaard, J. R. & Hebert, P. D. N. 2006. An inexpensive, automation-friendly protocol for recovering high-quality DNA. *Molecular Ecology Notes*, 6:998–1002. https://doi.org/10.1111/j.1471-8286.2006.01428.x

Jackson, J. A. & Tinsley, R. C. 2001. *Protopolystoma xenopodis* (Monogenea) primary and secondary infections in *Xenopus laevis*. *Parasitology*, 123:455–463. https://doi.org/10.1017/s0031182001008745

Jackson, J. A. & Tinsley, R. C. 2005. Geographic and within-population structure in variable resistance to parasite species and strains in a vertebrate host. *International Journal for Parasitology*, 25:29–37. https://doi.org/10.1016/j.ijpara.2004.10.017

Jørgensen, D. 2015. Conservation implications of parasite co-reintroduction. *Conservation Biology*, 29:602–604

Kalous, L., Musil, J., Petrtỳl, M., Vajglová, T., Romočuskỳ, Š. & Bušta, L. 2013. The danger in the anglers' bucket: qualitative and quantitative insight into bait fish market in Prague (Czech Republic). *Acta Societatis Zoologicae Bohemicae*, 77:27–35

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*, 14:587–589. https://doi.org/10.1038/nmeth.4285

Kilian, J. V., Klauda, R. J., Widman, S., Kashiwagi, M., Bourquin, R., Weglein, S. & Schuster, J. 2012. An assessment of a bait industry and angler behavior as a vector of invasive species. *Biological Invasions*, 14:1469–1481. https://doi.org/10.1007/s10530-012-0173-5

Kimura, M. 1981. Estimation of evolutionary distances between homologous nucleotide sequences. *Proceedings of the National Academy of Sciences USA*, 78:454–458. https://doi.org/10.1073/pnas.78.1.454

Kmentová, N., Van Steenberge, M., Raeymaekers, J. A. M., Kolbmüller, S., Hablützel, P., Bukinga, F. M., ... Vanhove, M. P. 2018. Monogenean parasites of sardines in Lake Tanganyika: diversity, origin and intraspecific variability. *Contributions to Zoology*, 87:105–132. https://doi.org/10.1163/18759866-08702004

Kmentová, N., Van Steenberge, M., van den Audenaerde, D. F. E. T., Nhiwatiwa, T., Bukinga, F. M., N'sibula, T. M., ... Vanhove, M. P. M. 2019. Co-introduction success of monogeneans infecting the fisheries target *Limnothrissa miodon* differs between two non-native areas: the potential of parasites as a tag for introduction pathway. *Biological Invasions*, 21:757–773. https://doi.org/10.1007/s10530-018-1856-3

Kocher, T. D., Thomas, W. K., Meyer, A., Edwards, S. V., Pääbo, S., Villablanca, F. X. & Wilson, A. C. 1989. Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences USA*, 86:6196–6200

Kruger, N. 2020. *Do vulnerable life stages of Xenopus laevis reduce its invasion potential?* Stellenbosch: Stellenbosch University and Université Claude Bernard Lyon 1. (Thesis – PhD).

Lee, C. E. & Gelembiuk, G. W. 2008. Evolutionary origins of invasive populations. *Evolutionary Applications*, 1:427–448.

https://doi.org/10.1111/j.1752-4571.2008.00039.x

Legendre, P., Desdevises, Y. & Bazin, E. 2002. A statistical test for host-parasite coevolution. *Systematic Biology*, 51:217–234.

https://doi.org/10.1080/10635150252899734

Leigh, J. W. & Bryant, D. 2015. PopART: full-feature software for haplotype network construction. *Methods in Ecology and*

Evolution, 6:1110–1116. https://doi.org/10.1111/2041-210x.12410

Li, W. & Godzik, A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences.

Bioinformatics, 22:1658–1659. https://doi.org/10.1093/bioinformatics/btl158

Lillo, F., Dufresnes, C., Faraone, F. P., Lo Valvo, M. & Stöck, M. 2013. Identification and potential origin of invasive

clawed frogs *Xenopus* (Anura: Pipidae) in Sicily based on mitochondrial and nuclear DNA. *Italian Journal of Zoology*,

80:566–573. https://doi.org/10.1080/11250003.2013.847502

Littlewood, D. T. J., Rohde, K. & Clough, K. A. 1997. Parasite speciation within or between host species? – Phylogenetic

evidence from site-specific polystome monogeneans. *International Journal for Parasitology*, 27:1289–1297. https:

//doi.org/10.1016/S0020-7519(97)00086-6

Lobos, G., Mendez, M. A., Cattan, P. & Jaksic, F. 2014. Low genetic diversity of the successful invasive African clawed frog

Xenopus laevis (Pipidae) in Chile. *Studies on Neotropical Fauna and Environment*, 49:50–60. https://doi.org/10.1080/

01650521.2014.912865

Lymbery, A. J., Morine, M., Kanani, H. G., Beatty, S. J. & Morgan, D. L. 2014. Co-invaders: the effects of alien parasites on

native hosts. *International Journal for Parasitology: Parasites and Wildlife*, 3:171–177. https://doi.org/10.1016/j.ijppaw.

2014.04.002

Measey, G. J. 2016. Overland movement in African clawed frogs (*Xenopus laevis*): a systematic review. *PeerJ*, 4, e2474.

https://doi.org/10.7717/peerj.2474

Measey, G. J. & Channing, A. 2003. Phylogeography of the genus *Xenopus* in southern Africa. *Amphibia-Reptilia*, 24:321–330.

https://doi.org/10.1163/156853803322440781

Measey, G. J. & Davies, S. J. 2011. Struggling against domestic exotics at the southern end of Africa. *Froglog*, 97:28–30

Measey, G. J., Davies, S. J., Vimercati, G., Rebelo, A., Schmidt, W. & Turner, A. 2017. Invasive amphibians in southern

Africa: a review of invasion pathways. *Bothalia*, 47, a2117. https://doi.org/10.4102/abc.v47i2.2117

Measey, G. J., Rödder, D., Green, S. L., Kobayashi, R., Lillo, F., Lobos, G., ... Thirion, J. M. 2012. Ongoing invasions of

the African clawed frog, *Xenopus laevis*: a global review. *Biological Invasions*, 14:2255–2270. https://doi.org/10.1007/

s10530-012-0227-8

Minh, B. Q. Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler A. & Lanfear, R. 2020. IQ-TREE

2: new models and efficient methods for phylogenetic inference in the genomic era. *Molocular Biology and Evolution*,

37:1530–1534. https://doi.org/10.1093/molbev/msaa015

Morais, P. & Reichard, M. 2018. Cryptic invasions: a review. *Science of the Total Environment*, 613–614:1438–1448.

https://doi.org/10.1016/j.scitotenv.2017.06.133

Nieberding, C., Morand, S., Libois, R. & Michaux, J. R. 2004. A parasite reveals cryptic phylogeographic history of its host.

Proceedings of the Royal Society B: Biological Sciences, 271:2559–2568. https://doi.org/10.1098/rspb.2004.2930

Nieberding, C. M. & Olivieri, I. 2007. Parasites: proxies for host genealogy and ecology? *Trends in Ecology and Evolution*,

22:156–165. https://doi.org/10.1016/j.tree.2006.11.012

Northover, A. S., Lymbery, A. J., Wayne, A. F., Godfrey, S. S. & Thompson, R. C. A. 2018. The hidden consequences of

altering host-parasite relationships during fauna translocations. *Biological Conservation*, 220:140–148. https://doi.org//10.

1016/j.biocon.2017.12.037

Ondračková, M., Matějusová, I. & Grabowska, J. 2012. Introduction of *Gyrodactylus perccotti* (Monogenea) into Europe

on its invasive fish host, Amur sleeper (*Perccottus glenii*, Dybowski 1877). *Helminthologia*, 49:21–26. https://doi.org/10.

2478/s11687-012-0004-3

Paradis, E. & Schliep, K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R.

Bioinformatics, 35:526–528. https://doi.org/10.1093/bioinformatics/bty633

Pariselle, A., Boeger, W. A., Snoeks, J., Bilong Bilong, C. F., Morand, S. & Vanhove, M. P. M. 2011. The monogenean

parasite fauna of cichlids: a potential tool for host biogeography. *International Journal of Evolutionary Biology*, 2011:1–15.

https://doi.org/10.4061/2011/471480

Poisot, T. & Stouffer, D. 2015. When is cophylogeny evidence of coevolution? In: Morand, B. R., Krasnov, S. & Littlewood,

D. T., eds. *Parasite Diversity and Diversification*. Cambridge, UK: Cambridge University Press. pp. 420–434.

Posada, D. 2003. Using MODELTEST and PAUP∗ to select a model of nucleotide substitution. *Current Protocols in*

Bioinformatics, 00:6.5.1–6.5.14. https://doi.org/10.1002/0471250953.bi0605s00

Prentis, P. J., Sigg, D. P., Raghu, S., Dhileepan, K., Pavasovic, A. & Lowe, A. J. 2009. Understanding invasion history: genetic

structure and diversity of two globally invasive plants and implications for their management. *Diversity and Distributions*,

15:822–830. https://doi.org/10.1111/j.1472-4642.2009.00592.x

QGIS Development Team. 2018. *QGIS Geographic Information System*. Oregon, United States of America: Open Source

Geospatial Foundation Project. http://qgis.osgeo.org/

R Core Team. 2020. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical

Computing. https://www.R-project.org/

Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and*

Evolution, 3:217–223. https://doi.org/10.1111/j.2041-210X.2011.00169.x

Reyda, F. B., Wells, S. M., Ermolenko, A. V., Zietara, M. S. & Lumme, J. I. 2020. Global parasite trafficking: Asian

Gyrodactylus (Monogenea) arrived to the U.S.A. via invasive fish *Misgurnus anguillicaudatus* as a threat to amphibians.

Biological Invasions, 22:391–402. https://doi.org/10.1007/s10530-019-02097-4

Ricciardi, A. 2007. Are modern biological invasions an unprecedented form of global change? *Conservation Biology*,

21:329–336. https://doi.org/10.1111/j.1523-1739.2006.00615.x

Richardson, D. M., Pyšek, P. & Carlton, J. T. 2011. A compendium of essential concepts and terminology in invasion ecology.

In: Richardson, D. M., ed. *Fifty years of invasion ecology: the legacy of Charles Elton*. Oxford: Wiley-Blackwell.

Rius, M. & Turon, X. 2020. Phylogeography and the description of geographic patterns in invasion genomics. *Frontiers in*

Ecology and Evolution, 8, art. 595711. https://doi.org/10.3389/fevo.2020.595711

Roy, H. E. & Lawson Handley, L.-J. 2012. Networking: a community approach to invaders and their parasites. *Functional*

Ecology, 26:1238–1248. https://doi.org/10.1111/j.1365-2435.2012.02032.x

Rozas, J., Ferrer-Mata, A., Sánchez-DelBarrio, J., Guirao-Rico, S., Librado, P., Ramos-Onsins, S. & Sánchez-Gracia, A.

2017. DnaSP 6: DNA sequence polymorphism analysis of large datasets. *Molecular Biology and Evolution*, 34:3299–3302.

https://doi.org/10.1093/molbev/msx248

Saltonstall, K. 2002. Cryptic invasion by a non-native genotype of the common reed, *Phragmites australis*, into North

America. *Proceedings of the National Academy of Sciences USA*, 99:2445–2449. https://doi.org/10.1073/pnas.032477999

Sievers, F. & Higgins, D. G. 2017. Clustal Omega for making accurate alignments of many protein sequences. *Tools for*

Protein Science, 27:135–145. https://doi.org/10.1002/pro.3290

Solarz, W. & Najberek, K. 2017. Alien parasites may survive even if their original hosts do not. *EcoHealth*, 14, art. 3.

https://doi.org/10.1007/s10393-015-1092-6

Soubrier, J., Steel, M., Lee, M. S., Der Sarkissian, C., Guindon, S., Ho, S. Y. & Cooper, A. 2012. The influence of rate

heterogeneity among sites on the time dependence of molecular rates. *Molecular Biology and Evolution*, 29:3345–3358.

https://doi.org/10.1093/molbev/mss140

Stouthamer, R., Rugman-Jones, P., Thu, P. Q., Eskalen, A., Thibault, T., Hulcr, J., ... Zong, S. 2017. Tracing the origin of a

cryptic invader: phylogeography of the *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae) species complex.

Agricultural and Forest Entomology, 19:366–375. https://doi.org/10.1111/afe.12215

Stöver, B. C. & Müller, K. F. 2010. TreeGraph 2: combining and visualizing evidence from different phylogenetic analyses.

BMC Bioinformatics, 11, art. 7. https://doi.org/10.1186/1471-2105-11-7

Sures, B., Nachev, M., Pahl, M., Grabner, D. & Selbach, C. 2017. Parasites as drivers of key processes in aquatic ecosystems:

facts and future directions. *Experimental Parasitology*, 180:141–147. https://doi.org/10.1016/j.exppara.2017.03.011

Theunissen, M., Tiedt, L. & du Preez, L. H. 2014. The morphology and attachment of *Protopolystoma xenopodis* (Monogenea:

Polystomatidae) infecting the African clawed frog *Xenopus laevis*. *Parasite*, 21, art. 20. https://doi.org/10.1051/parasite/

2014020

Thompson, C. W., Phelps, K. L., Allard, M. W., Cook, J. A., Dunnum, J. L., Ferguson, A. W., ... Kilpatrick, C. W. 2021.

Preserve a voucher specimen! The critical need for integrating natural history collections in infectious disease studies.

mBio, 12, e02698–20. https://doi.org/10.1128/mbio.02698-20

Tinsley, R. C. & Jackson, J. A. 1998a. Correlation of parasite speciation and specificity with host evolutionary relationships. *International Journal for Parasitology*, 28:1573–1582. https://doi.org/10.1016/s0020-7519(98)00085-x

Tinsley, R. C. & Jackson, J. A. 1998b. Speciation of *Protopolystoma* Bychowsky, 1957 (Monogenea: Polystomatidae) in hosts of the genus *Xenopus* (Anura: Pipidae). *Systematic Parasitology*, 40:93–141. https://doi.org/10.1023/B:SYPA.0000004047.41228.a6

van Sittert, L. & Measey, G. J. 2016. Historical perspectives on global exports and research of African clawed frogs (*Xenopus laevis*). *Transactions of the Royal Society of South Africa*, 71:157–166. https://doi.org/10.1080/0035919x.2016.1158747

van Wilgen, B. W., Measey, J., Richardson, D. M., Wilson, J. R. & Zengeya, T. A. 2020. Biological invasions in South Africa: an overview. In: van Wilgen, B., Measey, J., Richardson, D., Wilson, J. & Zengeya, T., eds. *Biological Invasions in South Africa*. Springer International Publishing. pp. 3–31.

Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R. & Hebert, P. D. 2005. DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society of London B*, 360:1847–1857. https://doi.org/10.1098/rstb.2005.1716

Weldon, C., de Villiers, A. L. & du Preez, L. H. 2007. Quantification of the trade in *Xenopus laevis* from South Africa, with implications for biodiversity conservation. *African Journal of Herpetology*, 56:77–83. https://doi.org/10.1080/21564574.2007.9635553

Yang, Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods. *Journal of Molecular Evolution*, 39:306–314. https://doi.org/10.1007/BF00160154

Zahiri, R., Christian Schmidt, B., Schintlmeister, A., Yakovlev, R. V. & Rindoš, M. 2019. Global phylogeography reveals the origin and the evolutionary history of the gypsy moth (Lepidoptera, Erebidae). *Molecular Phylogenetics and Evolution*, 137:1–13. https://doi.org/10.1016/j.ympev.2019.04.021