1 A monogenean parasite reveals the widespread
2 translocation of the African Clawed Frog in its native
3 range

Anneke L Schoeman[1,2,3]**, Louis H du Preez**[1,3]**, Nikol Kmentová**[4]**, and Maarten PM**

Vanhove[4]

[1]**African Amphibian Conservation Research Group, Unit for Environmental Sciences and Management,**

North-West University, Potchefstroom Campus, 11 Hoffman Street, Potchefstroom 2531, South Africa

[2]**DSI-NRF Centre of Excellence for Invasion Biology, Stellenbosch, South Africa**

[3]**South African Institute for Aquatic Biodiversity, Private Bag 1015, Grahamstown 6140, South Africa**

[4]**Hasselt University, Centre for Environmental Sciences, Research Group Zoology: Biodiversity &**

Toxicology, Agoralaan Gebouw D, B-3590 Diepenbeek, Belgium

Correspondence Anneke Lincoln Schoeman; email: anneke.lincoln@gmail.com

ABSTRACT

1. The management of bio-invasions relies upon the development of methods to trace their origin and expansion. Co-introduced parasites, especially monogenean flatworms, are ideal tags for the movement of their hosts due to their short generations, direct life cycles and host specificity. However, they are yet to be applied to trace the intraspecific movement of host lineages in their native ranges.

2. As proof of this concept, we conducted a comparative phylogeographic analysis based upon two mitochondrial markers of a globally invasive frog *Xenopus laevis* and its monogenean parasite *Protopolystoma xenopodis* in its native range in southern Africa and invasive range in Europe.

3. Translocation of lineages was largely masked in the frog's phylogeography. However, incongruent links between host and parasite phylogeography indicated host switches from one host lineage to another after these were brought into contact in the native range. Thus, past translocation of host lineages is revealed by the invasion success of its co-introduced parasite lineage.

4. This study demonstrates that parasite data can serve as an independent line of evidence in invasion biology, also on the intraspecific level, shedding light on previously undetected invasion dynamics. Based upon the distribution of these invasive parasite lineages, we infer that there is widespread anthropogenic translocation of this frog, not only via official export routes, but also facilitated by the frog's use as live bait by angling communities.

5. *Synthesis and applications.* Data from co-introduced, host-specific parasites, as tags for translocation, can add value to investigations in invasion biology and conservation. A better understanding of the translocation history and resulting genetic mixing of host and parasite lineages in the native range can shed light on the genetic make-up of parasite assemblages co-introduced to the invasive range. Knowledge of the intraspecific movement of different lineages of animals in their native ranges also has conservation implications, since contact between divergent lineages of hosts and parasites can facilitate host switches and altered parasite dynamics in both native and invasive populations. Therefore, we recommend the inclusion of parasite data as a more holistic approach to the invasion ecology of animals on the intraspecific level.

Keywords: biological invasion, comparative phylogeography, holistic specimen, intraspecific cryptic invasion, parasite tag, *Protopolystoma xenopodis*, *Xenopus laevis*

1 INTRODUCTION

The Anthropocene is characterised by the unparalleled human-mediated passage of organisms from their natural ranges to ecosystems across the world (Capinha *et al.*, 2015; Ricciardi, 2007). If these non-native organisms manage to disperse from their initial introduction point and seed new populations in their non-native range, they are regarded as invasive (Blackburn *et al.*, 2011; Richardson *et al.*, 2011). This reshuffling of the world's biota is a decisive agent of global change, as evidenced by the fact that invasive species are implicated as the cause of a third of recent animal extinctions (Blackburn *et al.*, 2019).

Clearly, the development of methods to trace both the origin and expansion of biological invaders is an urgent task. Genomic tools can be informative for the reconstruction of invasion histories, the prediction of potential range expansion and invasion success, and the development of control strategies (e.g. de Busschere *et al.*, 2016; Dufresnes *et al.*, 2017; Dlugosch & Parker, 2008; Hiller & Lessios, 2017; Hudson *et al.*, 2022; Lee & Gelembiuk, 2008; Prentis *et al.*, 2009; Stouthamer *et al.*, 2017; Zahiri *et al.*, 2019). Nonetheless, in many cases the available molecular data or chosen molecular markers are of insufficient resolution by themselves to trace the movement of species (e.g. Ahyong *et al.*, 2017; Criscione *et al.*, 2006). Thus, despite the accessibilty of genomic tools and the widespread use of phylogeographic analyses in invasion science, these methods certainly have their limitations (Fitzpatrick *et al.*, 2012; Rius & Turon, 2020).

A more holistic approach to solving the origin and range shifts of invasive species involves molecular data from co-introduced parasites (Gagne *et al.*, 2021). Such an integrative approach has been proposed to investigate invasions that are difficult to trace (Clavero *et al.*, 2016). For example, the long-debated

cryptogenic history of the European marine snail *Littorina littorea* (Linnaeus, 1758) was solved with the help of an associated digenean (Platyhelminthes: Digenea) parasite (Blakeslee *et al.*, 2008). Likewise, intraspecific morphometric variation of monogenean flatworm (Platyhelminthes: Monogenea) parasites identified both the origin and introduced life stage of invasive clupeid fish in Africa (Kmentová *et al.*, 2019), shed light on the history of an invasive population of odontobutid fish in Europe (Ondračková *et al.*, 2012) and provided information on the introduction route of invasive gobiid fish in Europe (Huyse *et al.*, 2015). Moreover, the genetic variation among monogenean gill parasites were used to illuminate the introduction history of non-native cichlid fish in Africa (Geraerts *et al.*, 2022; Jorissen *et al.*, 2022).

Matters are complicated even further when we wish to trace movement of conspecific organisms between different populations. This phenomenon, also known as intraspecific cryptic invasion, describes the process where one lineage of a species spreads into parts of the species' native range where another lineage is already present, often with human help (Morais & Reichard, 2018; Saltonstall, 2002). Once again, co-translocated parasites hold great promise to provide information on the movement of lineages of their hosts. Monogenean parasites in particular exhibit sufficient levels of phylogeographic structuring to assess intraspecific spatial patterns in their hosts (Baldwin *et al.*, 2012; Hahn *et al.*, 2015; Huyse *et al.*, 2017; Kmentová *et al.*, 2018). At the very least, the phylogeographic structuring of monogenean lineages can support evidence revealed by the host data set or historic records, such as the well-documented anthropogenic introduction of salmonid fish host stocks across Europe that is reflected in the phylogeography of their gyrodactylid monogenean lineages (Hahn *et al.*, 2015). Monogenean parasites are regarded as ideal tags of host genealogy due to their short generations, direct life cycles and generally high host-specificity (Nieberding & Olivieri, 2007; Pariselle *et al.*, 2011). Yet, the comparative phylogeographic analysis of host and parasite data has never been applied to trace intraspecific cryptic invasions within the native ranges of species.

The African Clawed Frog *Xenopus laevis* (Daudin, 1802) (Anura: Pipidae) is arguably the world's most widely distributed and well-studied amphibian (Measey *et al.*, 2012; van Sittert & Measey, 2016). Previous studies on the morphology and genetics of *X. laevis* have identified several differentiated populations in South Africa that are segregated by large mountain ranges and different climatic regions, with the most notable divergence between northern and southern populations (de Busschere *et al.*, 2016; du Preez *et al.*, 2009; Furman *et al.*, 2015; Grohovaz *et al.*, 1996; Measey & Channing, 2003). In opposition to the clear phylogeographic structuring stands the widespread occurrence of both natural and human-mediated translocation of *X. laevis* in the region. This movement is a consequence of naturally occurring overland migration, the domestic export of the frog for teaching and learning and the widely adopted practice of the translocation of the frog for use as live bait in recreational angling (de Villiers & Measey, 2017; Measey & Davies, 2011; Measey, 2016; Measey *et al.*, 2017; van Sittert & Measey, 2016; Weldon *et al.*, 2007). One would expect these translocations to leave a genetic trace (van Sittert & Measey, 2016). Yet, the cryptic invasion of *X. laevis* over long distances and across major geographic barriers in the native range has not been confirmed in previous phylogeographic studies (de Busschere *et al.*, 2016; du Preez *et al.*, 2009; Furman *et al.*, 2015; Measey & Channing, 2003).

We posit that the bladder polystomatid flatworm *Protopolystoma xenopodis* (Price, 1943) Bychowsky, 1957 (Monogenea: Polystomatidae) is the ideal tool to trace the hitherto unconfirmed contact between northern and southern lineages of its host *X. laevis* across its native range of southern Africa. In natural populations across sub-Saharan Africa, *P. xenopodis*, the most widely distributed of the species of *Protopolystoma*, is restricted to a single host species group, the *leavis* group, even in sympatry with other *Xenopus* species (Tinsley & Jackson, 1998a,b). Furthermore, in *P. xenopodis*, notable intraspecific morphological variation exists between populations of the polystomatid infecting *X. laevis* in southern Africa and those infecting more northerly host species (Tinsley & Jackson, 1998b). This pattern is

echoed even on a smaller regional scale in South Africa, where notable morphological and phylogenetic divergence exists between northern and southern populations of *P. xenopodis* that parasitise *X. laevis* (Schoeman *et al.*, 2022). The combination of high host specificity and intraspecific divergence suggests that *P. xenopodis* could be useful in a holobiont approach to a bio-invasion study, since any inferred movement of the parasite has to be mediated by movement of the host.

Therefore, in the present study, we use the widely distributed and intimately associated *X. laevis* and *P. xenopodis* system as a proof of concept that intraspecific parasite phylogeography can provide information on the cryptic introductions of their host lineages. We hypothesise that the phylogeography of *P. xenopodis* in South Africa will magnify that of its host *X. laevis*, which will allow us to employ a comparative phylogeographic analysis of this host and parasite in the context of their domestic translocation. We postulate that the parasite phylogeography will reveal contact between distinct lineages of *X. laevis* where evidence for translocation events might be lacking in the frog phylogeography and provide new insights on invasion pathways in the native range and beyond.

2 MATERIALS AND METHODS

2.1 Host and parasite sampling

In the native range of southern Africa, 46 adult *X. laevis* were captured from March 2017 to April 2019 in liver baited funnel traps from 30 localities (Fig. 1; see Appendix S1 for locality details). The localities were representative of the phylogeography of the frog, based upon the lineage assignment of de Busschere *et al.* (2016) (see Appendix S2 for locality classification). In the invasive range, 13 adult *X. laevis* were collected from 12 localities in western France and one in Portugal during the summers of 2017, 2018 and 2019 (see Appendix S1 for locality details). In France, frogs were captured in baited traps by members of the EU LIFE CROAA team as part of the regional eradication programme for

FIGURE 1 Localities in southern Africa where 46 adult *Xenopus laevis* were collected. Localities are classified according to the geographic extent of the native phylogeographic lineages of *X. laevis* (de Busschere *et al.*, 2016). Key localities are named according to the closest town. Jonkershoek (☆) is the source locality of the majority of historic *X. laevis* exports. Mountain ranges are indicated with shaded relief and rainfall seasonality is represented by the wettest month of each 25 km^2 grid cell. Monthly mean precipitation data, obtained from the WorldClim 2.1 database at a spatial resolution of 2.5 arc-minutes (Fick & Hijmans, 2017), were used to identify the months with the highest historical mean precipitation from 1970 to 2000 for each grid cell. Elevation data were obtained from the Natural Earth database. The map was constructed in QGIS *version* 3.10.2-A Coruña (QGIS Development Team, 2018) with the Mercator projection.

invasive amphibians. Frogs in Portugal were obtained from the locally captured laboratory stock of the

University of Lisbon.

All frogs underwent double euthanasia with anaesthesia in 6% ethyl-3-aminobenzoate methanesul-

fonate (MS222) (Sigma-Aldrich Co., USA) followed by pithing as required by the institutional ethics

committee. During subsequent dissection, 59 *P. xenopodis* (one per frog) were retrieved from the

bladders for molecular analyses. Polystomatids and liver tissue from their corresponding frog hosts were

preserved in 70% ethanol.

111 2.2 Molecular data collection

For the comparative phylogeographic analyses, we obtained the genetic sequences of the mitochondrial

barcoding genes for cytochrome *c* oxidase subunit I (*COX1*) and for the small 12S subunit of ribosomal

RNA (*rrnS*) of both the frogs and the polystomatids. The DNA was extracted from host and parasite

tissue with the PCRBIO Rapid Extract PCR kit (PCR Biosystems Ltd., UK) or the *Quick*-DNA^{TM}

Miniprep Plus kit (Zymo Research, USA). Subsequently, amplification reactions for the mitochondrial

COX1 and *rrnS* genes of both frogs and polystomatids were prepared with 2 to 5 µL genomic DNA,

1.25 µL forward primer [0.2 µM], 1.25 µL reverse primer [0.2 µM], 12.5 µL master mix and PCR-grade

water to the final volume of 25 µL. The 2× PCRBIO HS Taq Mix Red (PCR Biosystems Ltd., UK) or

the OneTaq® 2× Master Mix with Standard Buffer (New England Biolabs Inc., USA) were used with

the temperature of the elongation steps as per supplier instructions (68 °C or 72 °C).

For the frog *COX1*, the 'LCO1490'/'HCO2198' (Folmer *et al.*, 1994) and 'VF1'/'VR1' (Ivanova

et al., 2006; Ward *et al.*, 2005) primer pairs were implemented with the following thermocycling profile:

initial denaturation at 95 °C for 2 min, followed by 35 cycles of denaturation at 95 °C for 30 s, annealing

for 30 s at 50 °C for the first primer pair and 54 °C for the second and elongation for 2 min, terminated

by a final elongation step of 7 min. In the case of the '12SA-F'/'12SB-R' primer pair (Kocher *et al.*,

1989) to obtain the frog *rrnS*, the profile of Amaro *et al.* (2013) was followed. For the amplification of

the polystomatid *COX1* and *rrnS*, the profile of Héritier *et al.* (2015) was implemented for the primer

pairs 'L-CO1p'/'H-Cox1R', 'L-CO1p'/'H-Cox1p2', '12SpolF1'/'12SpolR1' and '12SpolF1'/'12SpolR9'

(Héritier *et al.*, 2015; Littlewood *et al.*, 1997).

For purification and sequencing, PCR products were sent to a commercial company (Inqaba Biotec,

RSA) that used the ExoSAP protocol (New England Biolabs Ltd.) for purification and obtained the

sequences with BigDye® Terminator *version* 3.1 Cycle Sequencing, utilising the corresponding primer

pairs used in the PCR reaction, on an ABI3500XL analyser. Sequences were assembled and manually

edited in Geneious Prime *version* 2021.0.3 (www.geneious.com). The 236 obtained sequences were

deposited in GenBank (accession numbers to be added after manuscript revision).

137 2.3 Calculation of sequence statistics

An investigation of sequence diversity is an exploratory analysis that can reveal the relative differentiation

within and between geographic clusters. It can also give an indication of the markers' applicability to

subsequent analyses. For these analyses, we obtained alignments of each of the four gene sets: *COX1*

or *rrnS* of the frog and *COX1* or *rrnS* of the polystomatid. The Clustal Omega algorithm under default

settings (Sievers & Higgins, 2017) implemented in Geneious Prime was used for the alignment of all four

data sets. For the protein-coding *COX1* gene, translation alignment was implemented under the vertebrate

mitochondrial genetic code (table 2) for the frog and the echinoderm and flatworm mitochondrial genetic

code (table 9) for the polystomatid.

Genetic diversity indices were calculated for all specimens combined and for geographic clusters

in the host and parasite samples. Number of haplotypes and polymorphic sites, haplotype diversity,

nucleotide diversity and mean percentage of pairwise differences were calculated in Arlequin *version* 3.5

(Excoffier & Lischer, 2010). Three groups were defined based upon collection locality (see Appendix S2

for full details), namely specimens collected in South Africa to the southwest of the Great Escarpment,

the major geographical barrier to dispersal in the region (SA1, SA2, SA3, SA4 and SA7; n = 30), to

the northeast of the Great Escarpment (SA5 and SA6; n = 16), or in their non-native range in France

(n = 12). The specimen from Portugal was not included as a separate group in the analyses because of

small sample size.

2.4 Phylogenetic analyses

The phylogenetic relationships between the different populations of both the host *X. laevis* and the

parasite *P. xenopodis*, represented in phylogenetic trees derived from the *COX1* and *rrnS* genes, formed

the basis of the subsequent comparative phylogeographic analysis.

The sequences were prepared for the phylogenetic analyses through removal of duplicates, addition

of outgroups for rooting, re-alignment and concatenation. Duplicate sequences were identified with the

help of CD-HIT (Fu *et al.*, 2012; Li & Godzik, 2006) implemented online (Huang *et al.*, 2010). Only

specimens with unique *COX1* or *rrnS* gene haplotypes were retained for the analysis, leading to 30

unique sequence combinations (23 unique haplotypes of *COX1* and 10 of *rrnS*) in the case of *X. laevis*

and 44 (34 of *COX1* and 35 of *rrnS*) in the case of *P. xenopodis*. In order to root the trees, the *COX1*

and *rrnS* gene sequences of the congeneric Müller's clawed frog *X. muelleri* (Peters, 1844) (GenBank

accession numbers EU599031, AY581657; Evans *et al.*, 2004, Evans *et al.*, 2008) and its polystomatid

P. occidentalis Tinsley *et* Jackson, 1998 (GenBank accession numbers KR856179, KR856121; Héritier

et al., 2015) were added as outgroups. This polystomatid only infects species from the *muelleri* group,

even in sympatry with *X. laevis* (Tinsley & Jackson, 1998b). Furthermore, its host *X. muelleri* reportedly

reproduces sterile offspring when hybridisation occurs between this species and its sympatric congener

X. laevis (Fischer *et al.*, 2000). Therefore, the selected outgroups are applicable to an intraspecific

analysis of the *X. laevis* and *P. xenopodis* host-parasite system. The resulting data sets, after removal of

duplicates and addition of the outgroups, were re-aligned by gene as detailed above. The two aligned

genes of each species were concatenated by individual in Geneious Prime. Ultimately, 31 sequences of a

total of 1153 bp (717 bp *COX1* and 436 bp *rrnS*), containing 189 sites with gaps and 183 variable sites,

of which 88 were parsimony informative, were used for the phylogenetic analysis of the frog data set.

Likewise, the 45 aligned sequences consisting of a total of 934 bp (423 bp *COX1* and 511 bp *rrnS*), with

799 complete sites, of which 225 polymorphic sites with a subset of 174 parsimony informative sites,

were use to infer the phylogeny of the parasite.

Prior to phylogenetic inference, the most optimal models of molecular evolution of each gene

partition were determined. Initially, the concatenated alignments comprised four partitions each, namely

the three codon positions of the partial *COX1* gene and the *rrnS* gene. The ModelFinder selection

routine (Kalyaanamoorthy *et al.*, 2017), in conjunction with the FreeRate heterogeneity model (Soubrier

et al., 2012), implemented in IQ-TREE *version* 2.1.2 (Minh *et al.*, 2020), was employed to select the

optimal model of molecular evolution for each partition. To reveal the optimal partitioning scheme and

most suitable evolutionary models for each alignment, the initial partitions were sequentially merged

(Chernomor *et al.*, 2016) until model fit, as evaluated by the Bayesian Information Criterion (BIC),

ceased to improve.

Finally, we obtained phylogenies of the host and the parasite using the maximum likelihood (ML)

approach. The ML phylogeny of the host was inferred under the TIM2 model (Posada, 2003) with

edge-unlinked FreeRate heterogeneity models (Soubrier *et al.*, 2012), proportions of invariant sites (Gu

et al., 1995) and discrete gamma models (Yang, 1994) for all partitions. Similarly, the ML tree of the

polystomatid was inferred under the TPM3 model (Kimura, 1981) with a FreeRate heterogeneity model

(Soubrier *et al.*, 2012) for all partitions. The analyses were performed in IQ-TREE *version* 2.1.2 (Minh

et al., 2020), with the assessment of branch support through ultrafast bootstrapping (UFboot2; Hoang

et al., 2018) and the Shimodaira-Hasegawa approximate likelihood ratio test (SH-aLRT; Guindon *et al.*,

2010), each with 10 000 replicates. The two phylogenies were rooted with respect to the outgroups in

TreeGraph2 *version* 2.15.0-877 *beta* (Stöver & Müller, 2010). Clades with UFboot2 support lower than

95% or SH-aLRT lower than 80% were collapsed to polytomies in the same programme.

2.5 Comparative phylogeographic analyses

The assessment of the global fit between the two phylogenies at the level of lineages provided a framework to interpret the mismatches or incongruencies represented by certain individual host-parasite links as indications of long distance movement and subsequent parasite switching to different host lineages. The global congruence of the two phylogenies was assessed by the implementation of both Procrustean Approach to Co-phylogeny (PACo) (Balbuena *et al.*, 2013) and the fourth-corner statistical approach of ParaFit (Legendre *et al.*, 2002). Both these methods consider the contribution of individual specimen interactions to phylogenetic congruence (Balbuena *et al.*, 2013; Legendre *et al.*, 2002; Poisot & Stouffer, 2015). Moreover, both methods can accommodate trees that are not fully resolved and support instances of uneven numbers of terminal nodes of hosts and parasites.

In PACo, we assessed the dependency of the parasite phylogeny upon that of the host through asymmetrical Procrustean superimposition of host and parasite distance matrices, whilst ParaFit tested the dependence of both phylogenies upon one another (Balbuena *et al.*, 2013; Legendre *et al.*, 2002). Prior to analyses, the outgroups were removed from the ML phylogenies. The phylogenies were converted to matrices of patristic distances by the package *ape* (Paradis & Schliep, 2019) in R *version* 4.0.2 (R Core Team, 2020). These matrices served as input for both analyses, along with the corresponding host-parasite association matrix, where the polystomatids were assigned to their original host individuals, or to a host with an identical gene sequence to the original host individual.

The principal coordinates for PACo were derived with the R package *paco* (Hutchinson *et al.*, 2017) *version* 0.4.2 for the Procrustes superimposition in conjunction with the Cailliez correction (Cailliez, 1983) to avoid the production of negative eigenvalues of non-Euclidean phylogenetic distances. The PACo analysis was performed, using the asymmetric Procrustes statistic to test for phylogenetic tracking by the parasite of its host, in *paco* with 10 000 permutations, with conserved row sums and number

of interactions. The dynamics of the interacting clades were further explored by identifying those

interactions that most contribute to the possible concordance between the two phylogenies through a

jackknife procedure in *paco*. Those interactions with the smallest residual distances were interpreted as

stronger supporters of overall congruence, whilst those with larger residual distances represented likely

mismatches between host and parasite phylogeny.

The ParaFit analysis (Legendre *et al.*, 2002) was carried out with *ape* based upon the patristic distance

matrices, using 10 000 permutations with the Cailliez correction (Cailliez, 1983). Both the ParaFitLink

tests were employed to assess the significance of the individual links (Legendre *et al.*, 2002), which also

pointed out host-parasite links that signify incongruencies.

Visualisation was achieved through a tanglegram of the host and parasite ML phylogenies with a

weighted interaction network, based upon the residuals yielded by the PACo jackknife procedure. It

was constructed with the R package *phytools* (Revell, 2012) with the function *cophylo()*, which has an

internal algorithm that rotates nodes to optimally match tips. Due to the removal of duplicate sequences,

certain host and parasite terminal nodes were involved in multiple interactions.

Haplotype genealogies based upon bifurcarting phylogenetic trees and Fitch (1970) distances between

sequences are an alternative way of visualising phylogenies whilst giving an indication of the number

of individual specimens represented by each unique sequence in the data set. For this reason, we also

visualised the 59 *X. laevis* and the 59 *P. xenopodis*, that is with the inclusion of duplicates, in two

haplotype genealogies. The sequences were aligned and concatenated as detailed above. Two ML

phylogenies, one for the frog and one for the parasites, based on all 59 sequences of each and no

outgroups were derived using the procedure described above. These trees, along with the re-aligned

sequences, served as input for Fitchi (Matschiner, 2015), which produced two haplotype genealogy

graphs.

2.6 Use of frogs by anglers

As an addition to the information provided by (Weldon *et al.*, 2007) on the use of *X. laevis* as live bait by anglers in South Africa, we obtained anecdotal information on fishing practices from several stakeholders whilst conducting our fieldwork across the region. We asked about the practices surrounding the use of *X. laevis* as live bait during fishing and the translocation and release of *X. laevis* during fishing activities. Nine recreational and subsistence anglers in the Eastern Cape, Limpopo, Northwest, KwaZulu-Natal and Gauteng Provinces provided information on the use of *X. laevis* as live bait by their communities in the vicinity of some of our sampling localities.

3 RESULTS

3.1 Sequence diversity

The genetic diversity of the two parasite genes was generally higher than the two orthologous host genes, with greater genetic diversity among the samples collected in South Africa than in France in both host and parasite (Table 1). Genetic diversity was generally higher in the hosts collected to the southwest of the Great Escarpment than in those collected from the northeast, which was not the case for the parasites (Table 1).

3.2 Phylogenetic relationships of host and parasite

The phylogenetic analyses confirmed marked phylogeographic structuring in both the host and parasite in their native range, which provided the necessary resolution to assess spatial patterns. Tree reconstruction showed that frogs from the northeastern SA5 and SA6 localities in the native range form a well-supported clade, along with some of the frogs from the SA4 locality (clade NE in Fig. 2 with UFboot2 = 95% / SH-aLRT = 95.8%). In addition, two clades with unresolved relationships to one another and to NE could be identified among sampled frogs from the southwestern localities in the native range of *X. laevis*

TABLE 1 Genetic diversity of the bladder flatworm *Protopolystoma xenopodis* and its frog host *Xenopus laevis*, grouped according to sampling locality – collected in South Africa to the southwest or to the northeast of the Great Escarpment, or in France. Indices were inferred separately by gene for each group and for all samples combined from the aligned partial mitochondrial *COX1* and *rrnS* genes.

	n	H	L	S	Hd	π	d (%)
X. laevis COX1							
Southwestern SA	30	11	641	78	0.823 ± 0.061	0.026 ± 0.013	2.6
Northeastern SA	16	6	641	74	0.742 ± 0.084	0.025 ± 0.013	2.5
France	12	3	641	50	0.591 ± 0.108	0.014 ± 0.008	1.4
All specimens	58	16	641	86	0.880 ± 0.023	0.048 ± 0.024	4.8
X. laevis rrnS							
Southwestern SA	30	7	369	13	0.662 ± 0.088	0.008 ± 0.005	0.8
Northeastern SA	16	3	369	2	0.342 ± 0.140	0.001 ± 0.001	0.1
France	12	2	369	5	0.303 ± 0.148	0.004 ± 0.003	0.4
All specimens	58	9	369	14	0.681 ± 0.044	0.009 ± 0.005	0.9
P. xenopodis COX1							
Southwestern SA	30	20	416	94	0.959 ± 0.022	0.070 ± 0.035	7.0
Northeastern SA	16	15	416	89	0.992 ± 0.025	0.072 ± 0.037	7.2
France	12	3	416	3	0.591 ± 0.108	0.004 ± 0.003	0.4
All specimens	58	34	416	110	0.958 ± 0.014	0.072 ± 0.035	7.2
P. xenopodis rrnS							
Southwestern SA	30	14	484	78	0.903 ± 0.036	0.050 ± 0.025	5.0
Northeastern SA	16	14	484	87	0.983 ± 0.028	0.055 ± 0.029	5.5
France	12	2	484	1	0.485 ± 0.106	0.001 ± 0.001	0.1
All specimens	58	28	484	98	0.930 ± 0.020	0.048 ± 0.024	4.8

Abbreviations: SA, South Africa; n, sample size; H, number of haplotypes; L, length of sequence; S, number of polymorphic sites; Hd, haplotype diversity; π, nucleotide diversity; d, mean percentage of pairwise differences.

(clades SW_{coast} with UFboot2 = 81% / SH-aLRT = 86.1% and SW_{inland} with UFboot2 = 98% / SH-aLRT

= 99.7% in Fig. 2). On the whole, members of the first clade originate from localities closer to the

coastline (clade SW_{coast} in Fig. 2 from SA1, SA2, SA3, SA4 and SA7 localities) than the second one

(clade SW_{inland} in Fig. 2 from SA3, SA4 and SA6 localities). The non-native frog from the invasive

population in Portugal, along with one of the non-native frogs from western France, grouped together

with the native frogs in SW_{coast} in the host phylogeny, thus placing these invasive frogs in the same clade

as the frog from Jonkershoek (Fig. 1), the source of the majority of *X. laevis* exports. Nonetheless, the

invasive population of *X. laevis* in France also harboured several frogs that clustered with frogs from the

FIGURE 2 The interaction network and phylogenies of the frog *Xenopus laevis* (left) and its polystomatid *Protopolystoma xenopodis* (right) with *X. muelleri* and *P. occidentalis* as outgroups. Topologies are based on maximum likelihood analyses of concatenated *COX1* and *rrnS* gene alignments of 31 frogs and 45 polystomatids with low support clades (UFboot2 < 95% or SH-alrt < 80%) collapsed. Links indicate observed host-parasite associations, weighted according to the contribution of each interaction to the overall phylogenetic congruence with thicker lines corresponding to greater support. Colours refer to geographic origin to the southwest (SW) and northeast (NE) of the Great Escarpment, following the classification of de Busschere *et al.* (2016): SA1 (dark blue), SA2 (light blue), SA3 (red), SA4 (green), SA5 (yellow), SA6 (brown) and SA7 (purple) (see Fig. 1) with main lineages named and highlighted. Some parasite specimens from southwestern SA1, SA2, SA3, SA4 and SA7 localities are not assigned to any of the main lineages due to their paraphyletic grouping basal to lineages **sw3**, **sw4** and **ne**. Invasive *X. laevis* and co-introduced *P. xenopodis* from France (grey) and Portugal (pink) are also included. Key localities are Adelaide (A), Dullstroom (D), Hopetown (H), Johannesburg (J), Ugie (U), Verkykerskop (V) and Wakefield (W). Jonkershoek (☆) is the origin of the majority of *X. laevis* exports. Scale bars indicate 0.1 nucleotide substitutions per site.

native northeastern localities in the NE clade.

In the same manner as the host phylogeny, the ML inference of the *P. xenopodis* tree revealed a single,

well-supported clade of polystomatids from hosts from the northeastern SA5 and SA6 localities, along

with some of the parasites from the SA4 locality (clade **ne** in Fig. 2 with UFboot2 = 97% / SH-aLRT =

87.6%). Several clades basal to the **ne** group emerged among the polystomatids from hosts collected

at the southwestern SA1, SA2, SA3, SA4 and SA7 localities (clades **sw1**, **sw2**, **sw3** and **sw4** in Fig. 2).

The earliest diverging clade is a well-supported grouping of polystomatids collected from SA1, SA3

and SA7, along with one polystomatid from SA5 (clade **sw1** in Fig. 2 with UFboot2 = 94%/SH-alrt =

88.1%). Two other native southwestern clades included a clade of polystomatids from the SA3 localities

and one from SA6 (clade **sw2** in Fig. 2 with UFboot2 = 53%/SH-alrt = 90.7%) and another clade of

polystomatids from southwesten SA2 and SA3 localities and some northeastern SA5 and SA6 localities

(clade **sw3** in Fig. 2 with UFboot2 = 61% / SH-aLRT = 86.6%). Another well-supported clade, the sister

clade to **ne**, contained polystomatids from SA1, France and Portugal (clade **sw4** in Fig. 2 with UFboot2

= 100% / SH-aLRT = 100%). However, overall low support values of many internal nodes in the parasite

ML tree, especially the nodes basal to the **sw3**, **sw4** and **ne** lineages, hampered the interpretation of

phylogenetic clustering among some of the polystomatids collected from the southwestern localities.

Consequently, some parasite specimens were not assigned to any clades.

3.3 Congruence of host and parasite phylogeographies

The phylogeny of *X. laevis* exhibited significant overall congruence with that of *P. xenopodis* (PACo

m_{xy}^2 = 0.1034, p = 0.0018, n = 10000; ParaFitGlobal = 0.0202, $p < 0.0001$, n = 10000), with the PACo

analysis suggesting that the phylogeny of *P. xenopodis* tracked that of *X. laevis*. According to PACo, links

between frogs from SW$_{coast}$ and their corresponding polystomatids that fall within **sw1** in the parasite

phylogeny, generally contributed the most to the overall phylogenetic congruence. The establishment of

overall congruence between these two phylogenies allowed us to hone in on mismatches between the

two phylogenies, which signified past contact between divergent clades.

Based upon visual inspection of the tanglegram (Fig. 2), a few links represented mismatches between

the host and parasite phylogenies that could be explained by long distance translocation of frogs from the SW host lineages to the northeastern localities followed by spill-over of **sw1**, **sw2** and **sw3** polystomatids to NE hosts. These indications of host switching were identified in frog-polystomatid pairs from SA5 and SA6 localities (Dullstroom, Hopetown, Johannesburg, Verkykerskop, Adelaide, Ugie and Wakefield), where NE hosts harboured parasites that clustered with their southwestern **sw1**, **sw2** and **sw4** counterparts. The PACo jackknife procedure further identified the interactions between frog–polystomatid pairs from the invasive range in France and Portugal as links conveying low support to congruence. Moreover, interactions between frogs and polystomatids from the contact zone near Three Sisters (SA4) and Wakefield (SA6) demonstrate low support to overall congruence. Finally, the ParaFitLink tests reported 30 out of 48 links as significant at a level of 0.05. Out of the 18 links that did not significantly support congruence, 11 signified host switches by parasites to another host lineage that occurred in the invasive range of France and Portugal and at SA4, SA5 and SA6 localities in the native range (Johannesburg, Three Sisters, Ugie and Wakefield).

The haplotype genealogies of the frog host and the polystomatid parasite (Fig. 3) unsurprisingly demonstrated the same phylogeographic structuring as those revealed by the phylogenetic analysis of *X. laevis* (Fig. 2). The frog haplotype genealogy clearly revealed that all northeastern specimens clustered together, except for frog specimens from Wakefield (SA5), which clustered with both the NE and SW$_{inland}$ clades, and the frogs collected at Three Sisters (SA4), which were present in all three host lineages. This situation was somewhat different for the polystomatids, where the haplotype genealogy illustrated that the cluster of parasite specimens from northeastern hosts (**ne**) contained only nine of the sixteen specimens collected from the northeastern SA5 and SA6 localities. The seven remaining parasite specimens from SA5 and SA6 localities (Adelaide, Dullstroom, Hopetown, Johannesburg, Ugie, Verkykerskop and Wakefield) were spread across the remainder of the network, therefore clustering with

FIGURE 3 Unrooted haplotype genealogy graphs of the partial *COX1* and *rrnS* genes of **(a)** 59 *Xenopus laevis* specimens and **(b)** their corresponding 59 *Protopolystoma xenopodis* parasites. For frogs and their polystomatids collected in the native southern Africa (see inset map), colours refer to the phylogeographic lineages of the host (de Busschere *et al.*, 2016). Non-native *X. laevis* and *P. xenopodis* are from France (grey) and Portugal (pink). Highlighted haplotype groups correspond to the clades revealed by the phylogenetic analyses (Fig. 2): northeastern host (NE) and parasite (ne) and southwestern host (SW) and parasite (sw) clades. Key localities with host lineage switches are named, as well as Jonkershoek, the source locality of most of the *X. laevis* exports. Sizes of circles are proportional to haplotype frequencies (see scale). The number of mutational steps is marked on the connecting branches with dots. Dashed lines are extensions of dotted sections for improved visualisation.

325 polystomatids from southwestern hosts (**sw1**, **sw2**, **sw3**). Moreover, parasites collected at the Three

326 Sisters (SA4) localities clustered with both northeastern (**ne**) and southwestern (**sw2**) parasites.

3.4 Translocation of frogs by anglers

Our informal discussions with anglers confirmed the widespread use of *X. laevis* as live bait in five out of the nine South African provinces, with all nine anglers confirming the existence of the practice in their communities and surrounding areas. The frog was usually employed in the capture of black bass *Micropterus* Lacepède, 1802 and African sharp-toothed catfish *Clarias gariepinus* (Burchell, 1822). Black bass are non-native inhabitants of water bodies across South Africa (Hargrove *et al.*, 2015; Khosa *et al.*, 2019) and African sharp-toothed catfish are non-native to all areas to the south of the Great Escarpment (Weyl *et al.*, 2016). Seven out of the nine anglers indicated that specimens of *X. laevis* are generally caught and used locally, with short range translocation of the captured specimens. However, two of the anglers admitted to buying *X. laevis* from breeders and pet shops in urban areas (Potchefstroom and Johannesburg) and transporting them to more rural areas for use as live bait. These fishermen were active near some of our sampling localities, namely Potchefstroom, Johannesburg, Dullstroom and Verkykerskop. All fishermen confirmed that surplus live *X. laevis* are routinely released on site and that escapees are common. Notably, a fisherman in Ugie described the practice of acquiring live *X. laevis* from water bodies in the vicinity and translocating specimens to stock local dams for future use.

4 DISCUSSION

In this study, the widespread, yet previously hidden, domestic expansion of *X. laevis* lineages in South Africa is brought to light through a comparison with the phylogeography of its co-invasive parasite, *P. xenopodis*. The use of parasite data to reveal host connectivity is hardly novel (reviewed in Gagne *et al.*, 2021). Yet, the present study is the first application of host-parasite phylogeographic analyses to the detection of intraspecific cryptic invasions. Furthermore, it demonstrates that even a comparatively modest data set of less than 60 host-parasite pairs can reveal compelling evidence for

349 historic translocation.

4.1 Parasite clustering mirrors the intraspecific phylogeography of the host

If the phylogeographic structuring of a parasite mirrors that of its host, a phenomenon which is not

necessarily a consequence of co-evolutionary processes (de Vienne *et al.*, 2013), the genealogy of the

parasite can provide insight on the translocation of the host (Nieberding & Olivieri, 2007). Based on

our findings, there is phylogenetic congruence between the African Clawed Frog and its monogenean

parasite. This means that the observed phylogeographic structuring in the parasite, which is supported by

both molecular and morphological evidence (Schoeman *et al.*, 2022), can corroborate and expound upon

the phylogeography of the host *X. laevis*, which also shows notable phylogeographic structuring across

its native range (de Busschere *et al.*, 2016; du Preez *et al.*, 2009; Furman *et al.*, 2015; Grohovaz *et al.*,

1996; Measey & Channing, 2003). After a brief free-swimming stage, *P. xenopodis* completes the rest of

its life cycle in the excretory system of *X. laevis* (Theunissen *et al.*, 2014), which translates to a very

specific and intimate host-parasite association that can faithfully portray the phylogeographic structure

of the frog. However, the parasite data set provides an independent line of evidence and not a mere echo

of the phylogeographic structuring of the frog, since each parasite represents a separate infection event

from a pool of free-swimming larvae.

The genetically distinct host-parasite clusters display distributions that roughly follow the two

climatic regimes in the region, namely the summer rainfall region to the northeast and the winter rainfall

region to the southwest, with the plateau edge of the Great Escarpment and the many ridges of the

Cape Fold Mountains as assumed geographical barriers to natural dispersal (Furman *et al.*, 2015). The

more pronounced clustering among hosts and parasites and the higher haplotype diversity in hosts

that hail from the southwest of the Great Escarpment is likely brought about by the presence of the

many ridges of the Cape Fold Mountains (de Busschere *et al.*, 2016; du Preez *et al.*, 2009; Furman

et al., 2015; Grohovaz *et al.*, 1996; Measey & Channing, 2003). On the other hand, well-documented

human-mediated translocation of *X. laevis* in the area to the southwest of the Great Escarpment (van

Sittert & Measey, 2016; Weldon *et al.*, 2007), in conjunction with widespread overland migration of the

frog with the help of the many interconnected artificial water bodies in the region (de Villiers & Measey,

2017; Measey, 2016), ensure some gene flow – and exchange of parasites – between the divergent frog

populations in this area.

4.2 Parasite spillover is an indication of contact between divergent host lineages

Parasite phylogeography offers a tool to recover historical insights of host dispersal pathways, since host

lineage switches reflect past interactions between host organisms at various time scales (e.g. Barbosa

et al., 2012; Criscione *et al.*, 2006; Galbreath & Hoberg, 2015; Galbreath *et al.*, 2020; Nieberding *et al.*,

2004). The usefulness of parasite species in reflecting the introduction pathways of their hosts has been

demonstrated for fish hosts and their monogenean parasites on a few occasions (Geraerts *et al.*, 2022;

Huyse *et al.*, 2015; Jorissen *et al.*, 2022; Kmentová *et al.*, 2019; Ondračková *et al.*, 2012; Reyda *et al.*,

2020). However, these studies focused on species-level invasions from the native to the non-native range,

rather than intraspecific invasions of different lineages within the native range. Our study, on the other

hand, demonstrates the use of parasites as tags to trace intraspecific cryptic invasions. Multiple instances

of host lineage switches revealed by the phylogeographic analyses of *P. xenopodis* act as an indicator for

historic contact between distinct *X. laevis* lineages. This provides us with the means of recognising such

contact even in cases where the locally non-native lineage of *X. laevis* may have gone extinct after the

introduction event or where they are not detected at a certain sampling effort.

In fact, thus far, genetic evidence for the admixture of the lineages of *X. laevis* with native ranges on

opposite sides of the Great Escarpment has not been found, despite historical records pointing towards

large scale exports from the southwest to the rest of southern Africa (van Sittert & Measey, 2016; Weldon

et al., 2007). In our study, the host phylogeographic analyses suggest the presence of contact zones

between the northeastern (NE) and southwestern lineages (SW_{inland} & SW_{coast}) at two localities to the

northeast of the Great Escarpment where these lineages co-occur.

It is this sympatry of multiple host clades that makes host lineage switching by parasite lineages a

possibility. In turn, the observation of host switches from one lineage to another provides more evidence

for the translocation of *X. laevis* across southern Africa. Indeed, our study confirmed the presence

of southwestern parasite lineages (**sw1**, **sw2** & **sw3**) at seven localities to the northeast of the Great

Escarpment, where one would expect the northeastern parasite lineage (**ne**) to occur. In these instances,

southwestern parasite lineages that are translocated along with their respective host lineages, thus locally

non-native at these new localities, spill over to the locally native northwestern host lineage when these

divergent, previously allopatric host lineages come into contact (Galbreath & Hoberg, 2015).

Differential survival between locally native and non-native *X. laevis* lineages in southern Africa

can partially explain the fewer observed instances of establishment by non-native *X. laevis* lineages

when compared to its co-introduced polystomatid. In translocation experiments of the northern and

southern lineages in southern Africa, *X. laevis* tadpoles and metamorphs exhibited remarkable phenotypic

plasticity when translocated between rainfall regimes at the cost of longterm survival (Kruger, 2020).

Under these circumstances, the locally native *X. laevis* lineage would outcompete the non-native lineage,

preventing successful local invasion. Thus, the employment of host phylogeography alone to trace long

range translocation of *X. laevis* in southern Africa would mask a number of introduction events, as our

findings clearly demonstrate.

Upon arrival in the new area, the co-introduced *P. xenopodis* lineage may rapidly spread in a

population of locally native and therefore naive *X. laevis*. It has been shown that primary infections by

P. xenopodis in previously uninfected *X. laevis* elicit strong long term immunological responses against

secondary infections by the same parasite species (Jackson & Tinsley, 2001). Correspondingly, a cross-infectivity experiment between northeastern and southwestern lineages of *X. laevis* and *P. xenopodis* further suggested that lineages of *X. laevis* in their native ranges might be more susceptible to infection by the locally non-native lineage of *P. xenopodis* than to their corresponding locally native parasite lineage (Jackson & Tinsley, 2005). In turn, this effect will boost the nascent spread of non-native *P. xenopodis* lineages in the locally native lineage of *X. laevis*.

There are two possible explanations for the observed contact between divergent lineages of this host-parasite system in its native range: natural geographic expansion over the evolutionary history of the host-parasite system in response to past environmental fluctuations, or more recent human-mediated translocation of the host along with its corresponding parasite lineage. At first glance, it is noticeable that the non-native *P. xenopodis* specimens collected at northeastern localities present with haplotypes that were generally highly differentiated from their counterparts from the southwest, despite belonging to the same **sw** lineages. In contrast, the haplotypes of the co-introduced French and Portuguese polystomatids are identical or nearly identical to the source South African haplotypes, indicating low divergence after the introduction event. The apparently deep divergence between native southwestern polystomatids and domestically translocated southwestern polystomatids suggests a scenario where parasites that underwent natural geographic expansion in deep time persist to the present day, yet with plenty of time to accrue genetic polymorphisms that differentiate them from their close relatives in the southwest. However, at our level of sampling effort, such comparisons of haplotype diversity are premature. It is clear from the long branch lengths between native southwestern polystomatids in the haplotype genealogy that there is considerable differentiation among the southwestern populations of *P. xenopodis*. It is reasonable to assume that greater sampling effort will yield more southwestern haplotypes that could be very similar to the translocated southwestern haplotypes now found at northeastern localities. Furthermore,

it is worth noting that the number of domestic exports from the southwestern region of South Africa greatly dwarfs the number of the international exports (van Sittert & Measey, 2016), which translates to more introduction events of southwestern *X. laevis* to areas across South Africa, thus greater haplotype diversity among the domestically introduced frogs, than to France.

However, there is another reason affecting the plausibility of the scenario of earlier natural expansion of this host-parasite system. Of course, hosts and parasites can experience shared histories of geographic expansion and contact between isolated populations in response to historical environmental shifts, such as climate change (Galbreath & Hoberg, 2015; Hahn *et al.*, 2015; Huyse *et al.*, 2017; Kmentová *et al.*, 2018). Likewise, post expansion, the non-native host lineages may be rapidly lost from the population due to genetic drift or differential survival, whilst the non-native parasite lineages may persist and become widely distributed across the geographic range of its new host lineage, as has been demonstrated before on the species level for tapeworm parasites in Beringia (Galbreath *et al.*, 2020). Regardless, the formation of the Great Escarpment around 130 million years ago (Burke & Gunnel, 2008) predates the evolution of *X. laevis* over the last 15 million years (Furman *et al.*, 2015). The Great Escarpment is a dominant barrier to dispersal that has played a defining role in shaping the distribution and evolution of southern African fauna (e.g. Barlow *et al.*, 2013; Makokha *et al.*, 2007; Mynhardt *et al.*, 2015; Nielsen *et al.*, 2018; Predel *et al.*, 2012), not least among which *X. laevis* (Furman *et al.*, 2015). When considering the evidence from the phylogeographic structuring of insects, frogs, lizards, snakes and small mammals in southern Africa, where the Great Escarpment is a definite barrier to natural dispersal (Barlow *et al.*, 2013; Furman *et al.*, 2015; Makokha *et al.*, 2007; Mynhardt *et al.*, 2015; Nielsen *et al.*, 2018; Predel *et al.*, 2012), it seems highly unlikely that climatic fluctuations could have given rise to natural expansion of *X. laevis* beyond the plateau edge of the escarpment that may shape the distribution of the southwestern lineages of *P. xenopodis* to the current day.

Rather, we consider recent human-mediated translocation a more plausible explanation. In fact, we know that the bulk of exports of *X. laevis* since the 1930s from the southwest of Southern Africa, which amounts to over 400 000 animals, was sent to domestic destinations, especially to urban centers inland where *X. laevis* was used in research and teaching (van Sittert & Measey, 2016; Weldon *et al.*, 2007). Subsequent escapees from these centers of higher learning would lead to the presence of locally non-native *X. laevis* and *P. xenopodis* near these urban areas. Moreover, these escapees can spread even further through the ability of *X. laevis* to move overland, a mode of dispersal which is complemented by expansion via networks of artifical waterways (de Villiers & Measey, 2017; Measey, 2016). The rural locations of the parasite spillover events revealed by our analyses of polystomatid phylogeography suggest that smaller scale human-mediated dispersal events also have a role to play in the dispersal of *X. laevis*. The role of recreational fishing across South Africa in the dispersal of *X. laevis*, and by extension, *P. xenopodis*, has been recognised before (Weldon *et al.*, 2007). Previously documented interviews with local anglers and suppliers exposed the use of *X. laevis*, frequently bought from pet shops and local breeders, as live bait for predominantly catfish angling (Weldon *et al.*, 2007). Our own informal discussions with local anglers at the localities where we collected *X. laevis* affirm the widespread use of this frog, sometimes obtained from pet shops, as live bait for the capture of both black bass (*Micropterus* spp.) and catfish (*Clarias gariepinus*).

In terms of sheer numbers, the impact of these translocation events on the domestic dispersal of *X. laevis* has been deemed negligible when compared to exports from Jonkershoek, the most prolific exporter, and other official suppliers to the domestic market (van Sittert & Measey, 2016; Weldon *et al.*, 2007). Nonetheless, previous studies on other aquatic species have pointed out that the use of live bait in angling is an important pathway for introductions, even if the market might be small in economic terms (Kalous *et al.*, 2013; Kilian *et al.*, 2012). Therefore, the impact of anglers on the spread of locally

non-native *X. laevis* lineages to more remote areas, as opposed to the bigger centra that received the

documented official consignments, should not be underestimated. Our informal interviews identified a

myriad of ways for *X. laevis* to be released in new environments by anglers, from escapees *en route*, to

the deliberate release of surplus bait, to the intentional stocking of water bodies for future use. Therefore,

the combination of overland movement, dispersal along natural and artificial waterways and translocation

by the angling community could form a widespread undocumented dispersal network of *X. laevis* in

South Africa.

4.3 Parasite phylogeography as an underutilised tool in invasion biology

As shown above, the present study acts as a proof of concept for the potential of the comparative

phylogeographic approach to the clarification of patterns in intraspecific host divergence and dispersal.

In turn, this independent line of evidence can provide additional insight on invasion pathways and inform

management strategies. Here, we highlight some of the potential implications to demonstrate the broader

applicability of this integrative approach.

Our approach was able to provide a new perspective on the origin of the relatively high haplotype

diversity of the invasive population of *X. laevis* in France, recovered in our analysis and others, when

compared to other invasive populations in Portugal, Sicily and Chile (de Busschere *et al.*, 2016; Lillo

et al., 2013; Lobos *et al.*, 2014). The idea of multiple introduction events from various source populations

in South Africa has been offered before as the most parsimonious explanation for this genetic diversity

in the invasive population of *X. laevis* in western France (de Busschere *et al.*, 2016). Since we found

evidence for host switches by *P. xenopodis* in the native range, an event that necessarily involves the

sympatry of divergent host lineages, it is a certainty that multiple host lineages co-occur at certain

localities in the native range. In this, our results build upon the evidence provided by historical export

records (van Sittert & Measey, 2016; Weldon *et al.*, 2007), dispersal patterns (de Villiers & Measey,

2017; Measey, 2016) and anecdotal information (Weldon *et al.*, 2007) to show that the mixing of lineages

most likely already happened in the source populations in South Africa prior to the co-introduction of

X. laevis and *P. xenopodis* to western France (van Sittert & Measey, 2016; Weldon *et al.*, 2007).

The novel insights provided by the addition of the parasite data set has implications for both the

conservation of *X. laevis* and its host-specific parasites. The widespread anthropogenic translocation of

this host-parasite system disrupts interactions between *X. laevis* lineages and the *P. xenopodis* lineages

that tracked their phylogenetic divergence, ancient interactions that deserve to be targets of conservation

(Gómez & Nichols, 2013). Subsequent host switches between closely related lineages set the stage for

new diversity to arise and for novel ecological interactions to be established (Galbreath *et al.*, 2020;

Sures *et al.*, 2017). In the case of host-specific parasites, translocation of lineages within the native

range of a species may actually be of more concern than translocation to far-flung non-native ecosystems,

since these ecosystems will probably not harbour close relatives of the invasive host to facilitate host

switches (Blackburn & Ewen, 2017). Host-parasite interactions are also something to keep in mind

when conducting conservation re-introduction programmes, where captive-bred or locally non-native

individuals of a species are used to re-stock locally extinct populations of endangered species (Britt *et al.*,

2004). Our results highlight that the evaluation of the genetic and parasitological suitability of a source

population for translocation should also encompass the potential history of local introductions that may

have influenced the intraspecific parasitism of the host population (Jørgensen, 2015; Northover *et al.*,

2018).

This approach further holds great promise in the management of invasive species. Prior to invasion,

translocations in the native range may have already altered the source populations of invasive species, as

is the case of *X. laevis*. We need to take altered phylogeographic structure into account to adequately

monitor the course of an invasion or the novel interactions that may arise. Parasites as passengers or

invaders themselves easily slip under the radar in invasion biology (Blackburn & Ewen, 2017; Solarz & Najberek, 2017). Yet, the role of parasites on the move in emerging infectious diseases and habitat transformation via indirect density-mediated effects cannot be overemphasised (Amundsen *et al.*, 2013; Bobadilla-Suarez *et al.*, 2017; Daszak, 2000; Dunn *et al.*, 2012; Goedknegt *et al.*, 2016; Lymbery *et al.*, 2014; Roy & Lawson Handley, 2012; Sures *et al.*, 2017). Furthermore, the identification of the local angling community as a player in the translocation of *X. laevis* in the native range allows policy makers to broaden their intervention efforts in the management of *X. laevis* as a domestic exotic.

4.4 Recommendations to make parasite data sets work for invasion biologists

All in all, our findings agree with the necessity of a more holistic approach to data collection in invasion biology (Clavero *et al.*, 2016), especially with regard to parasites. The idea of the "holistic specimen" has recently gained some traction (Cook *et al.*, 2017; Galbreath *et al.*, 2019). This approach to the collection and study of organisms advocates for the integrated specimen-based studies of animals and their associated parasites and pathogens. The optimisation of sampling effort is crucial to improve the allocation of resources, both in terms of funding and specimen use, especially when working with endangered species or specimens that require considerable effort to obtain, or in view of other ethical concerns regarding invasive sampling.

The inclusion of parasite data during the data collection phase also ensures that the host and parasite data sets are comprised of directly comparable individuals which can improve the feasibility and value of comparative phylogeographic analyses. For example, the non-overlap between host and parasite data sets was cited as a limitation in the co-phylogeographic analyses conducted by both Barbosa *et al.* (2012) and Huyse *et al.* (2017). Overlap between data sets can be established through the sampling of host-parasite pairs from the same localities and time frames, whilst sampling of the corresponding parasite of each host specimen will not contribute to data set overlap, especially in the case of parasites with a free-living

stage. Even in cases where parasites are not the targets of the current project, the deposition of voucher material of both host and parasite from the sampling event into curated institutional collections, in other words overlapping data sets, can be vital for later studies (Galbreath *et al.*, 2019; Thompson *et al.*, 2021). Later on, DNA sequences could be linked to material deposited in these museum collections.

Similarly, another aspect to keep in mind is the choice of markers for our co-phylogeographic analyses. More often than not, investigators rely on markers for which primers are already available and entire analyses may even be based on a single marker (de Vienne *et al.*, 2013; Gutiérrez-García & Vásquez-Domínguez, 2011). Yet, successful comparative phylogeographic analyses rely on a combination of multiple markers with a sufficient degree of genetic divergence evident among populations (de Vienne *et al.*, 2013; Gutiérrez-García & Vásquez-Domínguez, 2011), which is often higher in the host than the parasite (Nieberding & Olivieri, 2007). Thus, the independent yet comparable DNA signature provided by the parasite data set may provide higher resolution than the host data set, where spatial patterns may be blurred by the recent mixing of previously isolated genotypes in invasive populations (Rius & Turon, 2020). As a next step to improve confidence in phylogeographic inferences, we also propose the addition of morphological data from the parasite specimens, as demonstrated by several recent studies on monogenean parasites (Huyse *et al.*, 2015; Kmentová *et al.*, 2019; Ondračková *et al.*, 2012; Schoeman *et al.*, 2022).

Ultimately, this study serves as a proof of concept that the use of parasite data sets will add value to investigations in invasion biology, especially in the case of intraspecific cryptic invasions. We show that the utility of parasite data sets can go beyond the provision of supporting evidence and provide greater resolution that may lreveal patterns that went undetected in the host data set. Our study echoes the sentiments of Hoberg *et al.* (2015), who argued for an integrated approach to parasitology as a discipline that holds the keys to understanding and managing global change in the Anthropocene.

DATA AVAILABILITY STATEMENT

All DNA sequences used for co-phylogeographic analyses will be made available on the public database GenBank, along with relevant metadata and accession numbers will be added upon article acceptance. Detailed locality data are contained in the supplementary information.

ACKNOWLEDGMENTS

ALS received funding from the DST-NRF Centre of Excellence for Invasion Biology (South Africa). The project utilised the infrastructure of the Unit for Environmental Sciences and Management, North-West University (South Africa). This research was supported by EMBRC Belgium – FWO project GOH3817N. NK and MPMV are financed by the Special Research Fund of Hasselt University (BOF21PD01 and BOF20TT06, respectively). The authors would like to express their sincere thanks to a number of persons who assisted in the collection of the frogs. Guénolé le Peutrec, Jean Secondi, Natasha Kruger and Rodolphe Olivier helped with the collection in France. Rui Rebelo provided access to frogs from Portugal. In South Africa, several farm owners graciously gave permission for collection to take place on their properties and provided lodging for the research team: Fanus and Olga Kritzinger, William and Christa van Zyl, Dave Schlebusch, Fanus and Carin Oberholzer, Danie and Annalise Marais, Johan Hamann, Tobie Bielt, Gert Bench, Stoffel Labuschagne, Jannie and Susan van Rensburg, Jan Meintjies, Marthinus Hartman, Douw and Louise de Jager and Ernest de Villiers. Lastly, Mathys Schoeman, Clarke Scholtz, Andrea Darvall, Willie Landman, Ferdi de Lange, John Measey and Roxanne Viviers assisted with the collection of frogs at the remainder of the localities in South Africa. The authors are also indebted to numerous local recreational fishermen who were willing to share details about their use of *Xenopus laevis* across South Africa. The utilisation of the frogs and the research protocols were approved by the Animal Care, Health and Safety in Research Ethics (AnimCare) Committee of the Faculty of Health

601 Sciences of the North-West University (ethics number: NWU-0380-16-A5-01). Animals from the South

602 African populations were sampled under the permits 0056-AAA007-00224 (CapeNature) and FAUNA

603 1343-2017 (Northern Cape) provided by the Department of Economic Development, Environmental

604 Affairs and Tourism.

605 CONFLICT OF INTEREST

606 All authors declare no conflict of interest.

607 AUTHORS' CONTRIBUTIONS

608 ALS conceived the initial idea, collected and analysed the data and led the writing of the manuscript.

609 All authors refined the initial concept, designed the methodology, contributed to interpretation of data,

610 contributed critically to the drafts and gave final approval for publication.

611 REFERENCES

Ahyong, S. T., Kupriyanova, E., Burghardt, I., Sun, Y., Hutchings, P. A., Capa, M. & Cox, S. L. 2017.

Phylogeography of the invasive Mediterranean fan worm, *Sabella spallanzanii* (Gmelin, 1791), in

Australia and New Zealand. *Journal of the Marine Biological Association of the United Kingdom*,

97:985–991. https://doi.org/10.1017/S0025315417000261

Amaro, R. C., Nunes, I., Canedo, C., Napoli, M. F., Junca, F. A., Verdade, V. K., ... Rodrigues, M. T. 2013.

A molecular phylogeny recovers *Strabomantis aramunha* Cassimiro, Verdade and Rodrigues, 2008

and *Haddadus binotatus* (Spix, 1824) (Anura: Terrarana) as sister taxa. *Zootaxa*, 3741:S569–S582.

https://doi.org/10.11646/zootaxa.3741.4.7

Amundsen, P. A., Lafferty, K. D., Knudsen, R., Primicerio, R., Kristoffersen, R., Klemetsen, A. &

Kuris, A. M. 2013. New parasites and predators follow the introduction of two fish species to a subarctic lake: implications for food-web structure and functioning. *Oecologia*, 171:993–1002. https://doi.org/10.1007/s00442-012-2461-2

Balbuena, J. A., Miguez-Lozano, R. & Blasco-Costa, I. 2013. PACo: a novel procrustes application to cophylogenetic analysis. *PLOS ONE*, 8, e61048. https://doi.org/10.1371/journal.pone.0061048

Baldwin, R. E., Banks, M. A. & Jacobson, K. C. 2012. Interating fish and parasite data as a holistic solution for identifying the elusive stock structure of Pacific sardines (*Sardinops sagax*). *Reviews in Fish Biology and Fisheries*, 22:137–156. https://doi.org/10.1007/s11160-011-9227-5

Barbosa, A. M., Thode, G., Real, R., Feliu, C. & Vargas, J. M. 2012. Phylogeographic triangulation: using predator-prey-parasite interactions to infer population history from partial genetic information. *PLoS One*, 7, e50877. https://doi.org/10.1371/journal.pone.0050877

Barlow, A., Baker, K., Hendry, C. R., Peppin, L., Phelps, T., Tolley, K. A., ... Wüster, W. 2013. Phylogeography of the widespread African puff adder (*Bitis arietans*) reveals multiple Pleistocene refugia in southern Africa. *Molecular Ecology*, 22:1134–1157

Blackburn, T. M., Bellard, C. & Ricciardi, A. 2019. Alien versus native species as drivers of recent extinctions. *Frontiers in Ecology and the Environment*, 17:203–207. https://doi.org/10.1002/fee.2020

Blackburn, T. M. & Ewen, J. G. 2017. Parasites as drivers and passengers of human-mediated biological invasions. *EcoHealth*, 14:61–73. https://doi.org/10.1007/s10393-015-1092-6

Blackburn, T. M., Pyšek, P., Bacher, S., Carlton, J. T., Duncan, R. P., Jarošík, V., ... Richardson, D. M. 2011. A proposed unified framework for biological invasions. *Trends in Ecology & Evolution*, 26:333–339. https://doi.org/10.1016/j.tree.2011.03.023

Blakeslee, A. M. H., Byers, J. E. & Lesser, M. P. 2008. Solving cryptogenic histories using host and

parasite molecular genetics: the resolution of *Littorina littorea*'s North American origin. *Molecular*

Ecology, 17:3684–3696. https://doi.org/10.1111/j.1365-294x.2008.03865.x

Bobadilla-Suarez, M., Ewen, J. G., Groombridge, J. J., Beckmann, K., Shotton, J., Masters, N.,

... Sainsbury, A. W. 2017. Using qualitative disease risk analysis for herpetofauna conserva-

tion translocations transgressing ecological and geographical barriers. *EcoHealth*, 14:47–60.

https://doi.org/10.1007/s10393-015-1086-4

Britt, A., Welch, C., Katz, A., Iambana, B., Porton, I., Junge, R., ... Haring, D. 2004. The re-stocking of

captive-bred ruffed lemurs (*Varecia variegata variegata*) into the Betampona Reserve, Madagascar:

methodology and recommendations. *Biodiversity and Conservation*, 13:635–657. https://doi.org/10.

1023/b:bioc.0000009497.24917.ae

Burke, K. & Gunnel, Y. 2008. The African erosion surface: a continental-scale synthesis of geomorphol-

ogy, tectonics, and environmental change over the past 180 million years. *Memoir of the Geological*

Society of America, 201:1–66

Cailliez, F. 1983. The analytical solution of the additive constant problem. *Psychometrika*, 48:305–310.

https://doi.org/10.1007/bf02294026

Capinha, C., Essl, F., Seebens, H., Moser, D. & Pereira, H. M. 2015. The dispersal of alien species

redefines biogeography in the Anthropocene. *Science*, 348:1248–1251. https://doi.org/10.1126/science.

aaa8913

Chernomor, O., von Haeseler, A. & Minh, B. Q. 2016. Terrace aware data structure for phylogenomic

inference from supermatrices. *Systematic Biology*, 65:997–1008. https://doi.org/10.1093/sysbio/syw037

Clavero, M., Nores, C., Kubersky-Piredda, S., & Centeno-Cuadros, A. 2016. Interdisciplinarity to reconstruct historical introductions: solving the status of cryptogenic crayfish. *Biological Reviews*, 91:1036–1049. https://doi.org/10.1111/brv.12205

Cook, J. A., Galbreath, K. E., Bell, K. C., Campbell, M. L., Carrière, S., Colella, J. P., ... Hoberg, E. P. 2017. The Beringian Coevolution Project: holistic collections of mammals and associated parasites reveal novel perspectives on evolutionary and environmental change in the North. *Arctic Science*, 3:585–617. https://doi.org/10.1139/as-2016-0042

Criscione, C. D., Cooper, B. & Blouin, M. S. 2006. Parasite genotypes identify source populations of migratory fish more accurately than fish genotypes. *Ecology*, 87:823–828. https://doi.org/10.1890/0012-9658(2006)87[823:pgispo]2.0.co;2

Daszak, P. 2000. Emerging infectious diseases of wildlife – threats to biodiversity and human health. *Science*, 287:443–449. https://doi.org/10.1126/science.287.5452.443

de Busschere, C., Courant, J., Herrel, A., Rebelo, R., Rödder, D., Measey, G. J. & Backeljau, T. 2016. Unequal contribution of native South African phylogeographic lineages to the invasion of the African clawed frog, *Xenopus laevis*, in Europe. *PeerJ*, 4, e1659. https://doi.org/10.7717/peerj.1659

de Vienne, D. M., Refregier, G., Lopez-Villavicencio, M., Tellier, A., Hood, M. E. & Giraud, T. 2013. Cospeciation vs host-shift speciation: methods for testing, evidence from natural associations and relation to coevolution. *New Phytologist*, 198:347–385. https://doi.org/10.1111/nph.12150

de Villiers, F. A. & Measey, G. J. 2017. Overland movement in African clawed frogs (*Xenopus laevis*):

empirical dispersal data from within their native range. *PeerJ*, 5, e4039. https://doi.org/10.7717/peerj.

4039

Dlugosch, K. M. & Parker, I. M. 2008. Founding events in species invasions: genetic variation,

adaptive evolution, and the role of multiple introductions. *Molecular Ecology*, 17:431–449. https:

//doi.org/10.1111/j.1365-294X.2007.03538.x

du Preez, L. H., Kunene, N., Hanner, R., Giesy, J. P., Solomon, K. R., Hosmer, A. & van der Kraak,

G. J. 2009. Population-specific incidence of testicular ovarian follicles in *Xenopus laevis* from South

Africa: a potential issue in endocrine testing. *Aquatic Toxicology*, 95:10–16. https://doi.org/10.1016/j.

aquatox.2009.07.018

Dufresnes, C., Santo, L. D., Leuenberger, J., Schuerch, J., Mazepa, G., Grandjean, N., ... Dubey, S. 2017.

Cryptic invasion of Italian pool frogs (*Pelophylax bergeri*) across Western Europe unraveled by multilo-

cus phylogeography. *Biological Invasions*, 19:1407–1420. https://doi.org/10.1007/s10530-016-1359-z

Dunn, A. M., Torchin, M. E., Hatcher, M. J., Kotanen, P. M., Blumenthal, D. M., Byers, J. E., ...

Fox, C. 2012. Indirect effects of parasites in invasions. *Functional Ecology*, 26:1262–1274. https:

//doi.org/10.1111/j.1365-2435.2012.02041.x

Evans, B. J., Carter, T. F., Tobias, M. L., Kelley, D. B., Hanner, R. & Tinsley, R. C. 2008. A new

species of clawed frog (genus *Xenopus*) from the Itombwe Massif, Democratic Republic of the

Congo: implications for DNA barcodes and biodiversity conservation. *Zootaxa*, 1780:55–68. https:

//doi.org/10.11646/zootaxa.1780.1.2

Evans, B. J., Kelley, D. B., Tinsley, R. C., Melnick, D. J. & Cannatella, D. C. 2004. A mitochondrial

DNA phylogeny of African clawed frogs: phylogeography and implications for polyploid evolution.

Molecular Phylogenetics and Evolution, 33:197–213. https://doi.org/10.1016/j.ympev.2004.04.018

Excoffier, L. & Lischer, H. E. L. 2010. ARLEQUIN suite ver 3.5: a new series of programs to perform

population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, 10:564–567.

https://doi.org/10.1111/j.1755-0998.2010.02847.x

Fick, S. E. & Hijmans, R. J. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global

land areas. *International Journal of Climatology*, 37:4302–4315. https://doi.org/10.1002/joc.5086

Fischer, W., Koch, W. & Elepfandt, A. 2000. Sympatry and hybridization between the clawed frogs

Xenopus laevis laevis and *Xenopus muelleri* (Pipidae). *Journal of Zoology*, 252:99–107. https:

//doi.org/10.1111/j.1469-7998.2000.tb00824.x

Fitch, W. M. 1970. Distinguishing homologous from analogous proteins. *Systematic Zoology*, 19:99–113

Fitzpatrick, B., Fordyce, J., Niemiller, M. & Reynolds, R. G. 2012. What can DNA tell us about

biological invasions? *Biological Invasions*, 14:245–253. https://doi.org/10.1007/s10530-011-0064-1

Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. 1994. DNA primers for amplification of

mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine*

Biology and Biotechnology, 3:294–299

Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. 2012. CD-HIT: accelerated for clustering the next generation

sequencing data. *Bioinformatics*, 28:3150–3152. https://doi.org/10.1093/bioinformatics/bts565

Furman, B. L., Bewick, A. J., Harrison, T. L., Greenbaum, E., Gvozdik, V., Kusamba, C. & Evans, B. J.

2015. Pan-African phylogeography of a model organism, the African clawed frog *Xenopus laevis*.

Molecular Ecology, 24:909–925. https://doi.org/10.1111/mec.13076

Gagne, R. B., Crooks, K., Craft, M. E., Chiu, E. S., Fountain-Jones, N. M., Malmberg, J. L., ...

VandeWoude, S. 2021. Parasites as conservation tools. *Conservation Biology*. https://doi.org/10.1111/

cobi.13719

Galbreath, K. E. & Hoberg, E. P. 2015. Host responses to cycles of climate change shape parasite

diversity across North America's Intermountain West. *Folia Zoologica*, 64:218–232. https://doi.org/10.

25225/fozo.v64.i3.a4.2015

Galbreath, K. E., Hoberg, E. P., Cook, J. A., Armién, B., Bell, K. C., Campbell, M. L., ... Hope, A. G.

2019. Building an integrated infrastructure for exploring biodiversity: field collections and archives

of mammals and parasites. *Journal of Mammalogy*, 100:382–393. https://doi.org/10.1093/jmammal/

gyz048

Galbreath, K. E., Toman, H. M., Li, C. & Hoberg, E. P. 2020. When parasites persist: tapeworms survive

host extinction and reveal waves of dispersal across Beringia. *Proceedings of the Royal Society B:*

Biological Sciences, 287, art. 20201825. https://doi.org/10.1098/rspb.2020.1825

Geraerts, M., Huyse, T., Barson, M., Bassirou, H., Bilong Bilong, C. F., Bitja Nyom, Arnold, R.,

... Vanhove, M. P. M. 2022. Mosaic or melting pot: The use of monogeneans as a biological tag

and magnifying glass to discriminate introduced populations of Nile tilapia in sub-Saharan Africa.

Genomics, 114, art. 110328. https://doi.org/10.1016/j.ygeno.2022.110328

Goedknegt, M. A., Feis, M. E., Wegner, K. M., Luttikhuizen, P. C., Buschbaum, C., Camphuysen,

K. C. J., ... Thieltges, D. W. 2016. Parasites and marine invasions: ecological and evolutionary

perspectives. *Journal of Sea Research*, 113:11–27. https://doi.org/10.1016/j.seares.2015.12.003

Gómez, A. & Nichols, E. 2013. Neglected wild life: parasitic biodiversity as a conservation target.

International Journal for Parasitology: Parasites and Wildlife, 2:222–227. https://doi.org/10.1016/j.ijppaw.2013.07.002

Grohovaz, G. S., Harley, E. & Fabian, B. 1996. Significant mitochondrial DNA sequence divergence in natural populations of *Xenopus laevis* (Pipidae) from South Africa. *Herpetologica*, 52:247–253. https://doi.org/10.1080/11250003.2013.847502

Gu, X., Fu, Y. X. & Li, W. H. 1995. Maximum likelihood estimation of the heterogeneity of substitution rate among nucleotide sites. *Molecular Biology and Evolution*, 12:546–557. https://doi.org/10.1093/oxfordjournals.molbev.a040235

Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology*, 59:307–321. https://doi.org/10.1093/sysbio/syq010

Gutiérrez-García, T. A. & Vásquez-Domínguez, E. 2011. Comparative phylogeography: designing studies while surviving the process. *BioScience*, 61:857–868. https://doi.org/10.1525/bio.2011.61.11.5

Hahn, C., Weiss, S. J., Stojanovski, S. & Bachmann, L. 2015. Co-speciation of the ectoparasite *Gyrodactylus teuchis* (Monogenea, Platyhelminthes) and its salmonid hosts. *PLOS ONE*, 10, e0127340. https://doi.org/10.1371/journal.pone.0127340

Hargrove, J. S., Weyl, O. L. F., Allen, M. S. & Deacon, N. R. 2015. Using tournament angler data to rapidly assess the invasion status of alien sport fishes (*Micropterus* spp.) in Southern Africa. *PLOS ONE*, 10, e0130056. https://doi.org/10.1371/journal.pone.0130056

Héritier, L., Badets, M., du Preez, L. H., Aisien, M. S., Lixian, F., Combes, C. & Verneau, O. 2015. Evolutionary processes involved in the diversification of chelonian and mammal polystomatid parasites

(Platyhelminthes, Monogenea, Polystomatidae) revealed by palaeoecology of their hosts. *Molecular*

Phylogenetics and Evolution, 92:1–10. https://doi.org/10.1016/j.ympev.2015.05.026

Hiller, A. & Lessios, H. A. 2017. Phylogeography of *Petrolisthes armatus*, an invasive species with low

dispersal ability. *Scientific Reports*, 7, 3359. https://doi.org/10.1038/s41598-017-03410-8

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. 2018. UFBoot2: improving

the ultrafast bootstrap approximation. *Molecular Biology and Evolution*, 35:518–522. https://doi.org/

10.1093/molbev/msx281

Hoberg, E. P., Agosta, S. J., Boeger, W. A. & Brooks, D. R. 2015. An integrated parasitology: revealing

the elephant through tradition and invention. *Trends in Parasitology*, 31:128–133. https://doi.org/10.

1016/j.pt.2014.11.005

Huang, Y., Niu, B., Gao, Y., Fu, L. & Li, W. 2010. CD-HIT Suite: a web server for clustering and

comparing biological sequences. *Bioinformatics*, 26:680–682. https://doi.org/10.1093/bioinformatics/

btq003

Hudson, J., Bourne, S. D., Seebens, H., hapman, M. A. & Rius, M. 2022. The reconstruction of invasion

histories with genomic data in light of differing levels of anthropogenic transport. *Philosophical*

Transactions of the Royal Society of London B, 377:art. 20210023. https://doi.org/10.1098/rstb.2021.

0023

Hutchinson, M. C., Cagua, E. F., Balbuena, J. A., Stouffer, D. B., Poisot, T. & Fitzjohn, R. 2017.

paco: implementing Procrustean approach to cophylogeny in R. *Methods in Ecology and Evolution*,

8:932–940. https://doi.org/10.1111/2041-210x.12736

Huyse, T., Oeyen, M., Larmuseau, M. H. D. & Volckaert, F. A. M. 2017. Co-phylogeographic study of the

flatworm *Gyrodactylus gondae* and its goby host *Pomatoschistus minutus*. *Parasitology International*,

66:119–125. https://doi.org/10.1016/j.parint.2016.12.008

Huyse, T., Vanhove, M. P. M., Mombaerts, M., Volckaert, F. A. M. & Verreycken, H. 2015. Parasite

introduction with an invasive goby in Belgium: double trouble? *Parasitology Research*, 114:2789–

2793. https://doi.org/10.1007/s00436-015-4544-6

Ivanova, N. V., Dewaard, J. R. & Hebert, P. D. N. 2006. An inexpensive, automation-friendly protocol

for recovering high-quality DNA. *Molecular Ecology Notes*, 6:998–1002. https://doi.org/10.1111/j.

1471-8286.2006.01428.x

Jackson, J. A. & Tinsley, R. C. 2001. *Protopolystoma xenopodis* (Monogenea) primary and secondary

infections in *Xenopus laevis*. *Parasitology*, 123:455–463. https://doi.org/10.1017/s0031182001008745

Jackson, J. A. & Tinsley, R. C. 2005. Geographic and within-population structure in variable resistance

to parasite species and strains in a vertebrate host. *International Journal for Parasitology*, 25:29–37.

https://doi.org/10.1016/j.ijpara.2004.10.017

Jørgensen, D. 2015. Conservation implications of parasite co-reintroduction. *Conservation Biology*,

29:602–604

Jorissen, M. W. P., Vanhove, M. P. M., Pariselle, A., Snoeks, J., Vreven, E., Šimková, A., ... Huyse, T.

2022. Molecular footprint of parasite co-introduction with Nile tilapia in the Congo Basin. *Research*

Square. https://doi.org/10.21203/rs.3.rs-995291/v1

Kalous, L., Musil, J., Petrtýl, M., Vajglová, T., Romočuský, Š. & Bušta, L. 2013. The danger in the

anglers' bucket: qualitative and quantitative insight into bait fish market in Prague (Czech Republic).

Acta Societatis Zoologicae Bohemicae, 77:27–35

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. 2017. ModelFinder:

fast model selection for accurate phylogenetic estimates. *Nature Methods*, 14:587–589. https://doi.

org/10.1038/nmeth.4285

Khosa, D., Marr, S. M., Wasserman, R. J., Zengaya, T. A. & Weyl, O. L. F. 2019. An evaluation of the

current extent and potential spread of Black Bass invasions in South Africa. *Biological Invasions*,

21:1721–1736. https://doi.org/10.1007/s10530-019-01930-0

Kilian, J. V., Klauda, R. J., Widman, S., Kashiwagi, M., Bourquin, R., Weglein, S. & Schuster, J. 2012.

An assessment of a bait industry and angler behavior as a vector of invasive species. *Biological*

Invasions, 14:1469–1481. https://doi.org/10.1007/s10530-012-0173-5

Kimura, M. 1981. Estimation of evolutionary distances between homologous nucleotide sequences.

Proceedings of the National Academy of Sciences USA, 78:454–458. https://doi.org/10.1073/pnas.78.1.

454

Kmentová, N., van Steenberge, M., Raeymaekers, J. A. M., Kolbmüller, S., Hablützel, P., Bukinga,

F. M., ... Vanhove, M. P. 2018. Monogenean parasites of sardines in Lake Tanganyika: diversity,

origin and intraspecific variability. *Contributions to Zoology*, 87:105–132. https://doi.org/10.1163/

18759866-08702004

Kmentová, N., van Steenberge, M., van den Audenaerde, D. F. E. T., Nhiwatiwa, T., Muterezi Bukinga,

F., Mulimbwa N'Sibula, T., ... Vanhove, M. P. M. 2019. Co-introduction success of monogeneans

infecting the fisheries target *Limnothrissa miodon* differs between two non-native areas: the potential

of parasites as a tag for introduction pathway. *Biological Invasions*, 21:757–773. https://doi.org/10.

1007/s10530-018-1856-3

Kocher, T. D., Thomas, W. K., Meyer, A., Edwards, S. V., Pääbo, S., Villablanca, F. X. & Wilson, A. C.

1989. Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with

conserved primers. *Proceedings of the National Academy of Sciences USA*, 86:6196–6200

Kruger, N. 2020. *Do vulnerable life stages of Xenopus laevis reduce its invasion potential?* Stellenbosch:

Stellenbosch University and Université Claude Bernard Lyon 1. (Thesis – PhD).

Lee, C. E. & Gelembiuk, G. W. 2008. Evolutionary origins of invasive populations. *Evolutionary*

Applications, 1:427–448. https://doi.org/10.1111/j.1752-4571.2008.00039.x

Legendre, P., Desdevises, Y. & Bazin, E. 2002. A statistical test for host-parasite coevolution. *Systematic*

Biology, 51:217–234. https://doi.org/10.1080/10635150252899734

Li, W. & Godzik, A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or

nucleotide sequences. *Bioinformatics*, 22:1658–1659. https://doi.org/10.1093/bioinformatics/btl158

Lillo, F., Dufresnes, C., Faraone, F. P., Lo Valvo, M. & Stöck, M. 2013. Identification and potential

origin of invasive clawed frogs *Xenopus* (Anura: Pipidae) in Sicily based on mitochondrial and nuclear

DNA. *Italian Journal of Zoology*, 80:566–573. https://doi.org/10.1080/11250003.2013.847502

Littlewood, D. T. J., Rohde, K. & Clough, K. A. 1997. Parasite speciation within or between host

species? – Phylogenetic evidence from site-specific polystome monogeneans. *International Journal*

for Parasitology, 27:1289–1297. https://doi.org/10.1016/S0020-7519(97)00086-6

Lobos, G., Mendez, M. A., Cattan, P. & Jaksic, F. 2014. Low genetic diversity of the successful invasive

African clawed frog *Xenopus laevis* (Pipidae) in Chile. *Studies on Neotropical Fauna and Environment*,

49:50–60. https://doi.org/10.1080/01650521.2014.912865

Lymbery, A. J., Morine, M., Kanani, H. G., Beatty, S. J. & Morgan, D. L. 2014. Co-invaders: the effects

of alien parasites on native hosts. *International Journal for Parasitology: Parasites and Wildlife*,

3:171–177. https://doi.org/10.1016/j.ijppaw.2014.04.002

Makokha, J. S., Bauer, A. M., Mayer, W. & Matthee, C. A. 2007. Nuclear and mtDNA-based phylogeny

of southern African sand lizards, *Pedioplanis* (Sauria: Lacertidae). *Molecular Phylogenetics and*

Evolution, 44:622–633

Matschiner, M. 2015. Fitchi: haplotype gegealogy graphs based on the Fitch algorithm. *Bioinformatics*,

32:1250–1252. https://doi.org/10.1093/bioinformatics/btv717

Measey, G. J. 2016. Overland movement in African clawed frogs (*Xenopus laevis*): a systematic review.

PeerJ, 4, e2474. https://doi.org/10.7717/peerj.2474

Measey, G. J. & Channing, A. 2003. Phylogeography of the genus *Xenopus* in southern Africa. *Amphibia-*

Reptilia, 24:321–330. https://doi.org/10.1163/156853803322440781

Measey, G. J. & Davies, S. J. 2011. Struggling against domestic exotics at the southern end of Africa.

Froglog, 97:28–30

Measey, G. J., Davies, S. J., Vimercati, G., Rebelo, A., Schmidt, W. & Turner, A. 2017. Invasive

amphibians in southern Africa: a review of invasion pathways. *Bothalia*, 47, a2117. https://doi.org/10.

4102/abc.v47i2.2117

Measey, G. J., Rödder, D., Green, S. L., Kobayashi, R., Lillo, F., Lobos, G., ... Thirion, J. M. 2012.

Ongoing invasions of the African clawed frog, *Xenopus laevis*: a global review. *Biological Invasions*,

14:2255–2270. https://doi.org/10.1007/s10530-012-0227-8

Minh, B. Q. Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler A. & Lanfear, R. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Molocular Biology and Evolution*, 37:1530–1534. https://doi.org/10.1093/molbev/msaa015

Morais, P. & Reichard, M. 2018. Cryptic invasions: a review. *Science of the Total Environment*, 613–614:1438–1448. https://doi.org/10.1016/j.scitotenv.2017.06.133

Mynhardt, S., Maree, S., Pelser, I., Bennett, N. C., Bronner, G. N., Wilson, J. W. & Bloomer, P. 2015. Phylogeography of a morphologically cryptic golden mole assemblage from South-Eastern Africa. *PLOS ONE*, 10, e0144995

Nieberding, C., Morand, S., Libois, R. & Michaux, J. R. 2004. A parasite reveals cryptic phylogeographic history of its host. *Proceedings of the Royal Society B: Biological Sciences*, 271:2559–2568. https://doi.org/10.1098/rspb.2004.2930

Nieberding, C. M. & Olivieri, I. 2007. Parasites: proxies for host genealogy and ecology? *Trends in Ecology and Evolution*, 22:156–165. https://doi.org/10.1016/j.tree.2006.11.012

Nielsen, S. V., Daniels, S. R., Conradie, W., Heinicke, M. P. & Noonan, B. P. 2018. Multilocus phylogenetics in a widespread African anuran lineage (Brevicipitidae: *Breviceps*) reveals patterns of diversity reflecting geoclimatic change. *Journal of Biogeography*, 45:2067–2079

Northover, A. S., Lymbery, A. J., Wayne, A. F., Godfrey, S. S. & Thompson, R. C. A. 2018. The hidden consequences of altering host-parasite relationships during fauna translocations. *Biological Conservation*, 220:140–148. https://doi.org//10.1016/j.biocon.2017.12.037

Ondračková, M., Matějusová, I. & Grabowska, J. 2012. Introduction of *Gyrodactylus perccotti* (Mono-

genea) into Europe on its invasive fish host, Amur sleeper (*Perccottus glenii*, Dybowski 1877). *Helminthologia*, 49:21–26. https://doi.org/10.2478/s11687-012-0004-3

Paradis, E. & Schliep, K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, 35:526–528. https://doi.org/10.1093/bioinformatics/bty633

Pariselle, A., Boeger, W. A., Snoeks, J., Bilong Bilong, C. F., Morand, S. & Vanhove, M. P. M. 2011. The monogenean parasite fauna of cichlids: a potential tool for host biogeography. *International Journal of Evolutionary Biology*, 2011:1–15. https://doi.org/10.4061/2011/471480

Poisot, T. & Stouffer, D. 2015. When is cophylogeny evidence of coevolution? In: Morand, B. R., Krasnov, S. & Littlewood, D. T., eds. *Parasite Diversity and Diversification*. Cambridge, UK: Cambridge University Press. pp. 420–434.

Posada, D. 2003. Using MODELTEST and PAUP∗ to select a model of nucleotide substitution. *Current Protocols in Bioinformatics*, 00:6.5.1–6.5.14. https://doi.org/10.1002/0471250953.bi0605s00

Predel, R., Neupert, S., Huetteroth, W., Kahnt, J., Waidelich, D. & Roth, S. 2012. Peptidomics-based phylogeny and biogeography of Mantophasmatodea (Hexapoda). *Systematic Biology*, 61:609–629

Prentis, P. J., Sigg, D. P., Raghu, S., Dhileepan, K., Pavasovic, A. & Lowe, A. J. 2009. Understanding invasion history: genetic structure and diversity of two globally invasive plants and implications for their management. *Diversity and Distributions*, 15:822–830. https://doi.org/10.1111/j.1472-4642.2009.00592.x

QGIS Development Team. 2018. *QGIS Geographic Information System*. Oregon, United States of America: Open Source Geospatial Foundation Project. http://qgis.osgeo.org/

R Core Team. 2020. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. https://www.R-project.org/

Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution*, 3:217–223. https://doi.org/10.1111/j.2041-210X.2011.00169.x

Reyda, F. B., Wells, S. M., Ermolenko, A. V., Zietara, M. S. & Lumme, J. I. 2020. Global parasite trafficking: Asian *Gyrodactylus* (Monogenea) arrived to the U.S.A. via invasive fish *Misgurnus anguillicaudatus* as a threat to amphibians. *Biological Invasions*, 22:391–402. https://doi.org/10.1007/s10530-019-02097-4

Ricciardi, A. 2007. Are modern biological invasions an unprecedented form of global change? *Conservation Biology*, 21:329–336. https://doi.org/10.1111/j.1523-1739.2006.00615.x

Richardson, D. M., Pyšek, P. & Carlton, J. T. 2011. A compendium of essential concepts and terminology in invasion ecology. In: Richardson, D. M., ed. *Fifty years of invasion ecology: the legacy of Charles Elton*. Oxford: Wiley-Blackwell.

Rius, M. & Turon, X. 2020. Phylogeography and the description of geographic patterns in invasion genomics. *Frontiers in Ecology and Evolution*, 8, art. 595711. https://doi.org/10.3389/fevo.2020.595711

Roy, H. E. & Lawson Handley, L.-J. 2012. Networking: a community approach to invaders and their parasites. *Functional Ecology*, 26:1238–1248. https://doi.org/10.1111/j.1365-2435.2012.02032.x

Saltonstall, K. 2002. Cryptic invasion by a non-native genotype of the common reed, *Phragmites australis*, into North America. *Proceedings of the National Academy of Sciences USA*, 99:2445–2449. https://doi.org/10.1073/pnas.032477999

Schoeman, A. L., Kmentová, N., du Preez, L. H. & Vanhove, M. P. M. 2022. Like host, like parasite: intraspecific divergence in a polystomatid flatworm parasite across South Africa echoes that of its frog host. *bioRxiv*. https://doi.org/10.1101/2022.03.15.483565

Sievers, F. & Higgins, D. G. 2017. Clustal Omega for making accurate alignments of many protein sequences. *Tools for Protein Science*, 27:135–145. https://doi.org/10.1002/pro.3290

Solarz, W. & Najberek, K. 2017. Alien parasites may survive even if their original hosts do not. *EcoHealth*, 14, art. 3. https://doi.org/10.1007/s10393-015-1092-6

Soubrier, J., Steel, M., Lee, M. S., Der Sarkissian, C., Guindon, S., Ho, S. Y. & Cooper, A. 2012. The influence of rate heterogeneity among sites on the time dependence of molecular rates. *Molecular Biology and Evolution*, 29:3345–3358. https://doi.org/10.1093/molbev/mss140

Stouthamer, R., Rugman-Jones, P., Thu, P. Q., Eskalen, A., Thibault, T., Hulcr, J., ... Zong, S. 2017. Tracing the origin of a cryptic invader: phylogeography of the *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae) species complex. *Agricultural and Forest Entomology*, 19:366–375. https://doi.org/10.1111/afe.12215

Stöver, B. C. & Müller, K. F. 2010. TreeGraph 2: combining and visualizing evidence from different phylogenetic analyses. *BMC Bioinformatics*, 11, art. 7. https://doi.org/10.1186/1471-2105-11-7

Sures, B., Nachev, M., Pahl, M., Grabner, D. & Selbach, C. 2017. Parasites as drivers of key processes in aquatic ecosystems: facts and future directions. *Experimental Parasitology*, 180:141–147. https://doi.org/10.1016/j.exppara.2017.03.011

Theunissen, M., Tiedt, L. & du Preez, L. H. 2014. The morphology and attachment of *Protopolystoma*

xenopodis (Monogenea: Polystomatidae) infecting the African clawed frog *Xenopus laevis*. *Parasite*,

21, art. 20. https://doi.org/10.1051/parasite/2014020

Thompson, C. W., Phelps, K. L., Allard, M. W., Cook, J. A., Dunnum, J. L., Ferguson, A. W., ...

Kilpatrick, C. W. 2021. Preserve a voucher specimen! The critical need for integrating natural history

collections in infectious disease studies. *mBio*, 12, e02698–20. https://doi.org/10.1128/mbio.02698-20

Tinsley, R. C. & Jackson, J. A. 1998a. Correlation of parasite speciation and specificity with host

evolutionary relationships. *International Journal for Parasitology*, 28:1573–1582. https://doi.org/10.

1016/s0020-7519(98)00085-x

Tinsley, R. C. & Jackson, J. A. 1998b. Speciation of *Protopolystoma* Bychowsky, 1957 (Monogenea:

Polystomatidae) in hosts of the genus *Xenopus* (Anura: Pipidae). *Systematic Parasitology*, 40:93–141.

https://doi.org/10.1023/B:SYPA.0000004047.41228.a6

van Sittert, L. & Measey, G. J. 2016. Historical perspectives on global exports and research of African

clawed frogs (*Xenopus laevis*). *Transactions of the Royal Society of South Africa*, 71:157–166.

https://doi.org/10.1080/0035919x.2016.1158747

Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R. & Hebert, P. D. 2005. DNA barcoding Australia's

fish species. *Philosophical Transactions of the Royal Society of London B*, 360:1847–1857. https:

//doi.org/10.1098/rstb.2005.1716

Weldon, C., de Villiers, A. L. & du Preez, L. H. 2007. Quantification of the trade in *Xenopus laevis*

from South Africa, with implications for biodiversity conservation. *African Journal of Herpetology*,

56:77–83. https://doi.org/10.1080/21564574.2007.9635553

Weyl, O. L. F., Daga, V. S., Ellender, B. R. & Vitule, J. R. S. 2016. A review of *Clarias gariepinus*

invasions in Brazil and South Africa. *Journal of Fish Biology*, 89:386–402. https://doi.org/10.1111/jfb.

12958

Yang, Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates

over sites: approximate methods. *Journal of Molecular Evolution*, 39:306–314. https://doi.org/10.

1007/BF00160154

Zahiri, R., Christian Schmidt, B., Schintlmeister, A., Yakovlev, R. V. & Rindoš, M. 2019. Global

phylogeography reveals the origin and the evolutionary history of the gypsy moth (Lepidoptera,

Erebidae). *Molecular Phylogenetics and Evolution*, 137:1–13. https://doi.org/10.1016/j.ympev.2019.

04.021