Non-motile subpopulations of *Pseudomonas aeruginosa* repress flagellar motility

in motile cells through a type IV pili- and Pel-dependent mechanism

Kimberley A. Lewis[a], Danielle M. Vermilyea[a], Shanice S. Webster[a], Jaime de Anda[b,c,d],

Gerard C. L. Wong[b,c,d], George A. O'Toole[a] and Deborah A. Hogan[a,#]

[a]Department of Microbiology and Immunology, Geisel School of Medical at Dartmouth,

Hanover, New Hampshire, United States of America.

[b]Department of Bioengineering, University of California Los Angeles, CA 90095

[c]Department of Chemistry and Biochemistry, University of California Los Angeles, CA

90095

[d]California NanoSystems Institute, University of California Los Angeles, CA 90095

Running Head: Non-motile *P. aeruginosa* represses flagellar swarming motility

[#]To whom correspondence should be addressed

Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth

Vail Building HB7550

Hanover, NH 03755

E-mail: dhogan@dartmouth.edu

Key words: swarming, *Pseudomonas aeruginosa*, motility, type IV pili, Pel, microbe-

microbe interaction

Abstract

The downregulation of *P. aeruginosa* flagellar motility is a key event in biofilm formation, host-colonization, and the formation of microbial communities, but the external factors that repress motility are not well understood. Here, we report that under swarming conditions, motility can be repressed by cells that are non-motile due to the absence of a flagellum or flagellar rotation. Non-motile cells, due to mutations that prevent either flagellum biosynthesis or rotation, present at 5% of the total population suppressed swarming of wild-type cells under the conditions tested in this study. Non-motile cells required functional type IV pili and the ability to produce the Pel exopolysaccharide to suppress swarming by the motile wild type. In contrast, motile cells required only type IV pili, but not Pel production, in order for swarming to be repressed by non-motile cells. We hypothesize that interactions between motile and non-motile cells may enhance the formation of sessile communities including those involving multiple genotypes, phenotypically-diverse cells, and perhaps other species.

Significance

Our study shows that, under the conditions tested, a small population of non-swarming cells can impact the motility behavior of the larger population. The interactions that lead to the suppression of swarming motility require type IV pili and a secreted polysaccharide, two factors with known roles in biofilm formation. These data suggest that interactions between motile and non-motile cells may enhance the transition to sessile growth in populations and promote interactions between cells with different genotypes.

Introduction

Microbial localization through processes other than cell-division are critical for the formation of spatially structured populations and communities. Thus, motility and its regulation by diverse chemical and physical stimuli are major drivers of intraspecies and interspecies microbial interactions (1, 2). Previously, we found that exogenous ethanol, a common product secreted by many microbes, markedly reduced *Pseudomonas aeruginosa* swimming and swarming at the population level, but only decreased the motile fraction of cells in the population from an average of 38% to 22% (3). Prompted by this finding, we sought to determine the impact of a subpopulation of non-motile cells on the motility of the larger motile *P. aeruginosa* population.

During *P. aeruginosa* biofilm formation, motile cells are recruited to microcolonies of sessile cells via a combination of processes driving colocalization, and these processes may differ between strains with distinct early-biofilm forming strategies (4, 5) or in response to different experimental conditions. For *P. aeruginosa* strain PA14, retractile type IV pili (T4P) participate in surface sensing, which results in the upregulation of cAMP and the subsequent induction of cyclic-di-GMP (6-8); flagellar motility is then down-regulated in order to facilitate surface attachment (6, 7, 9, 10) followed by Pel exopolysaccharide matrix production (11), which can connect cells to one another (10, 12). Both matrix production and T4P function have been shown to participate in motility repression (10, 12-14) suggesting that matrix materials may mediate physical interactions between neighboring cells. T4P also mediate cell-cell interactions in swarms (15).

In this study, we show that non-motile cells when present at 5 to 75% of the total

inoculum resulted in repression of swarming by WT motile cells. To repress motility,

non-motile cells required the ability to produce Pel matrix, and both motile and non-

motile cells required functional T4P for this interaction. These data have implications for

factors that contribute to population-level behaviors and intra- and inter-species

interactions. Interactions that promote cell-cell interactions may be relevant in situations

where non-motile and motile cells can be found together such as in cystic fibrosis (CF)-

associated lung infections (16-18), wounds (19), and ear infections (17).

Results

The presence of non-motile cells represses swarming motility by *P. aeruginosa*

strain PA14. To explore the effects of a subpopulation of non-motile cells on swarming

motility, mixes containing different proportions of motile *P. aeruginosa* strain PA14 wild-

type (WT) cells and non-motile Δ*flgK* mutant cells (which lack the flagellar hook protein)

were plated on 0.5% M8 agar, which supports swarming motility by the WT strain. The

percent of Δ*flgK* mutant cells ranged from 5% to 75% in these mixes, and single strains

were included as controls. As expected, cultures with 100% WT swarmed readily while

cultures with 100% Δ*flgK* showed no swarming (**Fig. 1A**). In agreement with published

literature, the addition of non-motile Δ*flgK* blocked WT swarming (20), although, we

found that swarming motility was completely repressed by the addition of non-motile

cells at 5% to 75% of the total population under the conditions tested (**Fig. 1A** and **Fig.**

1C).

To determine if the lack of swarming in WT:Δ*flgK* co-cultures was due to faster

growth of the non-motile cells compared to the WT, we competed the WT or Δ*flgK*

mutant against a WT strain tagged with a *lacZ* reporter at different ratios atop a 0.22 µm

filter placed on 0.5% M8 agar. After 16 h, the colony was disrupted and colony forming

units (CFUs) for each strain were enumerated on blue/white screening plates that

contained X-Gal. At all ratios tested, neither untagged WT nor the Δ*flgK* mutant could

outgrow the *lacZ*-expressing WT strain (**Fig. 1B**), indicating that the lack of swarming

observed in WT:Δ*flgK* mixes was not due to increased growth of non-motile cells and

thus overgrowth of the population. Rather, our data suggest that a subpopulation of

non-motile cells inhibits motility of the larger flagellated WT population.

Using conditions in which the non-motile strain comprised 25%, 50%, or 75% of

the population, we also found that, like the Δ*flgK* mutant, the Δ*fliC* mutant, another non-

motile strain that lacks the flagellin protein used to make the flagellar filament, inhibited

WT motility at all ratios tested (**Fig. 1C**). Similarly, the Δ*motABCD* mutant, which

produces a flagellum that is unable to rotate due to the absence of stators, repressed

motility of the WT; although, in comparison to Δ*flgK* and Δ*fliC*, the Δ*motABCD* mutant

was slightly less effective at repressing WT swarming (**Fig. 1C**). Together, these data

suggest that the absence of flagellar motility in a subpopulation of cells is sufficient to

suppress WT swarming in co-culture.

T4P are required for non-motile cells to affect swarming by motile cells. Several

studies have shown that *P. aeruginosa* strains that are deficient in T4P tend to have a

hyper-swarming phenotype (21-23). Additionally, *P. aeruginosa* cells have been shown

to interact via their T4P to form close cell associations that facilitate directional

swarming (15). Consistent with published results (4, 24, 25), the ∆*flgK* mutant has

functional T4P as evidenced by the formation of a large twitching motility zone using the

agar-plastic interface assay (**Fig. S1A** and **Fig. S2**). While both the WT and ∆*flgK*

formed twitch zones that were significantly larger than those formed by the T4P-null

∆*pilA* mutant, the ∆*flgK* twitch zone was ~25% smaller than that formed by the WT

(p<0.05; **Fig. S2A**). Using ∆*flgK* and ∆*flgKpilA* strains, we found that *pilA*, which

encodes the major pilin component of T4P, was required for the ∆*flgK* mutant to

suppress WT swarming on 0.5% M8 agar (**Fig. 2A**). In a similar assay using 0.3% M63

agar, which supports both flagellum-mediated swimming and swarming, we found that

inoculated spots of ∆*flgK* cells (**Fig. 2B, red dots**) decreased the local expansion of the

motile WT population, while spots of the ∆*flgKpilA* mutant (**Fig. 2B, purple dots**) did

not. In contrast to the effects of non-motile ∆*flgK* cells on flagellum-mediated motility,

the ∆*flgK* mutant did not alter T4P-dependent twitching motility of the WT in a 50:50

ratio when compared to the WT alone (**Fig. S2B**). Therefore, the data show that T4P

are required for non-motile cells to inhibit flagellum-mediated motility of WT cells in co-

culture.

Functional T4P in motile cells are required for non-motile cells to repress

swarming in co-culture. With the observation that T4P are required for non-motile

cells to suppress flagellum-mediated swarming motility by the WT, we determined if this

interaction also required T4P function for the motile cells. We analyzed the ability of

∆*flgK* to repress swarming when co-cultured with the following mutants: ∆*pilA,* which

lacks T4P; Δ*pilMNOP*, which lacks the alignment complex required for T4P function (26)

and cyclic-di-GMP signaling (27); and both Δ*pilT* and Δ*pilU*, which lack either of the

ATPases that mediate T4P retraction (28). Strains lacking *pilA*, *pilMNOP*, *pilT*, and *pilU*

were all defective for twitching motility (**Fig. S1** and **Fig. S2**).

The swarming motility of strains lacking *pilA* or *pilMNOP* (both of which are

twitching defective) or *pilT* (no T4P retraction) was no longer inhibited by the Δ*flgK*

mutant in co-culture (**Fig. 3** and **Fig. S3**). Interestingly, the swarming motility of the

Δ*pilU* mutant was still inhibited by the Δ*flgK* mutant in co-culture (**Fig. 3**). While PilT is

the main retractile ATPase, PilU has an accessory role in retraction; the Δ*pilT* mutant

completely lacks pilus retraction (29, 30), while the Δ*pilU* mutant has reduced retraction

but still retains some function (30). These differences were confirmed by the fact that

the WT and Δ*pilU* were sensitive to phage DMS3 infection, but the Δ*pilA* and Δ*pilT*

mutants are resistant to infection (**Fig. S1B**). Together, these data indicate that motile

cells likely need to have functional T4P in order for the non-motile subpopulation to

inhibit the swarming motility of the WT strain when grown in co-culture.

P. aeruginosa T4P participate in a cAMP-dependent signaling pathway (see **Fig.**

S4A for pathway) that involves FimS, PilJ, CyaAB adenylate cyclases, and the cAMP-

binding transcription factor Vfr. We found that this pathway was not required for the

repression of WT motility by the non-motile Δ*flgK* mutant (**Fig. S4B**) using published

mutants lacking components of the cAMP signaling pathway (Δ*pilJ*, Δ*fimS*, Δ*cyaAB*, and

Δ*vfr*) that retain expression of functional T4P as evidenced by their sensitivity to phage

DMS3 (**Fig. S4C**). Taken together, the data show that motile WT cells require functional

T4P, but not the cAMP-dependent surface-sensing pathway, in order for non-motile

cells to be able to repress swarming in co-culture.

Non-motile *P. aeruginosa* requires Pel matrix production to repress swarming

motility in the motile population. We next explored whether Pel matrix production

played a role in swarming repression. We found that swarming by the Δ*pelA* mutant,

which is a mutant capable of robust swarming but lacks Pel production, was repressed

by Δ*flgK* in co-culture like the WT (**Fig. 4**). In contrast, Δ*flgKpelA* did not repress WT

swarming (**Fig. 4A**). Interestingly, both Δ*flgK* and Δ*fliC*, which repress WT swarming

(**Fig. 1C**), had increased Pel production (**Fig. 4B** and **Fig. S5**). Together, these data

show that the non-motile strain needs to produce Pel matrix to repress swarming

motility, but the motile strain does not. Additionally, while T4P were required by the

Δ*flgK* mutant to suppress swarm motility, T4P were not required for the observed

increase in Pel production in the Δ*flgK* mutant background as evidenced by Congo Red

binding (**Fig. 4B**).

DISCUSSION

Based on the data presented here we propose a model (**Fig. 5**) whereby a

subpopulation of non-motile cells in a population limit swarming motility of the entire

population. Our analyses revealed that non-motile cells required T4P and Pel

polysaccharide to inhibit the larger swarming population, and that motile cells required

functional T4P but not the Vfr/cAMP signaling system to respond to the non-motile

population. These findings in regards to T4P and Pel are consistent with the recruitment

of cells to microcolonies during biofilm development and aggregate formation, which

has been reported in numerous contexts (e.g. (4)).

The involvement of T4P in swarming cells was previously reported by Anyan et

al. (15) who reported that T4P impact cell-cell interactions in swarms in ways that limit

the movement of cells away from the population front. T4P mutants have been shown to

have increased swarming motility, suggesting that the intercellular T4P interactions that

we report between motile and non-motile genotypes may also be occurring in single-

strain swarms in which all cells have the potential for flagellar motility (22, 23). Future

studies will determine if T4P play direct roles in cell-cell interactions or indirect roles in

structuring the community.

The WT and motile Δ*pelA* mutant responded similarly to non-motile cells while

Δ*flgKpelA* was no longer able to repress motility suggesting that not all cells need to

produce exopolysaccharide matrix to repress swarming. Our findings that Pel

production is only necessary in the non-motile subpopulation is supported by published

data showing that ethanol not only decreases *P. aeruginosa* flagellar motility (3), but

increases Pel production (31). Pel has been suggested to repress motility via the steric

hinderance of flagella (3, 32), but further studies are needed to test this hypothesis. The

involvement of Pel in the interaction between motile and non-motile cells is particularly

interesting in light of recent work by Whitfield et al. (33) which found that the Pel

biosynthetic locus is widespread across Gram-positive and Gram-negative bacteria.

Future studies will determine if Pel is specifically detected by a T4P-dependent

mechanism, or if other matrix materials play a similar role.

Boyle et al. (20) previously reported that Δ*flgK* was capable of repressing

swarming by WT strain PA14 when Δ*flgK* was added 5:1 or at 80% of the population,

while we show that Δ*flgK* at 5% of the population represses WT swarming. Our

previously published findings (3) also showed that while strain PA14 was capable of

swarming, an average of 38% of flagellated cells were motile at any given time.

Furthermore, we showed that ethanol strongly repressed *P. aeruginosa* swarming

motility by reducing the motile fraction of the population from 38% to 22% (3). Overall,

the data show that motility of individual flagellated WT cells is heterogeneous within a

population, but the addition of non-flagellated cells represses population-wide motility.

These findings suggest that the number of non-flagellated cells necessary to repress

WT motility may vary due to experimental conditions.

In the studies here, we seed a low percentage of non-swarming cells into a

swarming population, which is reminiscent of the situations where there are genetically

heterogeneous populations observed in the CF airway or other chronic infections (18,

34, 35) as well as during normal environmental growth (35, 36). Additionally, clinical *P.*

aeruginosa isolates from the CF lung have diverse motility phenotypes (18). This

diversity in motility is even observed within a single patient (18). We hypothesize that

such a mechanism promotes interspecies and intraspecies interactions. The strong

implication of these findings is that in a mixed population with a subpopulation of non-

motile cells, the entire population can be brought to a halt. Similarly, within a population

in which a significant number of cells have stopped swimming or swarming, the further

addition of even just a few non-motile cells can suppress motility for the entire

population (i.e., see the ethanol studies mentioned above). As *P. aeruginosa*

populations inherently have some number of non-motile or inactive cells (3), future

studies are required to determine how strain background or mutant genotype affects

phenotypic heterogeneity among isogenic cells.

Materials and Methods

Strains and media. Strains used in this study are listed in **Table 1**. *P. aeruginosa* strain

PA14 and derivatives were routinely cultured on lysogeny broth (LB) solidified with 1.5%

agar or in LB broth at 37^{o}C with shaking. For *P. aeruginosa* phenotypic assays, either

M63 [22 mM KH_2PO_4, 40 mM K_2HPO_4, and 15 mM $(NH_4)_2SO_4$] or M8 (42 mM Na_2PO_4,

22 mM KH_2PO_4, and 8.5 mM NaCl) minimal salts medium supplemented with $MgSO_4$ (1

mM), glucose (0.2%), and casamino acids (CAA; 0.5%) were used as indicated.

Swarming motility assays. Swarm assays were performed as previously described

(37). Briefly, M8 medium with 0.5% agar (swarm agar) was poured into 60 x 15 mm

plates and allowed to dry at room temperature for ~4 h prior to inoculation. The

indicated strains were grown for 8 to 16 h then each was normalized to an OD_{600} of 1.

Indicated isolates were then mixed at the indicated ratios in a final volume of 100 µl.

Each plate was inoculated with 0.5 µl of the liquid culture mixture and the plates

incubated face-up at 37^{o}C in stacks of no more than four plates for 16 h. Each mixture

was inoculated in four replicates and was assessed on at least three separate days.

Images were captured using a Canon EOS Rebel T6i camera and images assessed for

swarm repression.

Swimming motility assays. Swim assays were performed as previously described

(37). Briefly, M63 medium solidified with 0.3% agar (soft agar) was poured into petri

plates and allowed to dry at room temperature ($\sim 25^{o}$C) for ~4 h prior to inoculation.

Sterile toothpicks were used to inoculate bacterial mixes into the center of the agar

without touching the bottom of the plate; liquid cultures grown for 8 to 16 h were used as

inoculum. The indicated strains were normalized to an OD_{600} of 1 then were mixed at

the indicated ratios in a final volume of 100 μl. No more than four bacterial mixtures

were assayed per plate. Plates were incubated upright at 37^{o}C in stacks of no more

than four plates per stack for 18 to 20 h.

Twitching motility assays. T-broth [1% tryptone (w/v), 0.5% NaCl (w/v)] medium

solidified with 1.5% agar was poured into petri plates and allowed to dry at room

temperature ($\sim 25^{o}$C). Overnight (16 h) cultures were washed once in 1X PBS,

resuspended in 1X PBS, and then normalized to an OD_{600} of 1 in 1X PBS. Plates were

inoculated by dipping a sterile toothpick into the washed and normalized cultures and

then inserting the toothpick into the agar until it touched the bottom of the plate. Plates

were then incubated at 37^{o}C for 46 h after which time the agar was removed and the

plate incubated in 0.1% crystal violet for 10 min. Plates were then washed three times in

water and dried overnight at room temperature. Images of the dried, stained twitch area

were taken and twitch diameter was measured.

Competition experiments. Competition assays were performed to determine relative

growth of selected *P. aeruginosa* strains. Strains were grown overnight and 1 ml of

culture was pelleted at 15,682 *x g* for 2 min, and washed once in 1 mL 1X PBS followed

by resuspension in 1mL 1X PBS. The OD_{600} of each culture was normalized to 1. The

strains to be competed were mixed at the indicated ratios in a final volume of 100 µL

and then 0.5 µL of the combined suspension was spotted onto a 0.22 µm polycarbonate

filter (Millipore) placed on the surface of a swarm plate in triplicates. Plates were

incubated at 37˚C. Filters were then transferred into a 1.5 mL tube and the filter-

associated cells were resuspended by adding 1 ml PBS + 0.05% Triton X-100 detergent

and agitating the tubes at high speed for 2 minutes using a Genie Disruptor (Zymo).

This suspension as well as the inoculum were diluted, spread on LB plates

supplemented with 150 µg/mL 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-

Gal) using glass beads, and incubated at 30˚C until blue colonies were visible (~24h).

The number of blue and white colonies per plate were counted and recorded to

determine the relative abundance of each strain.

Phage susceptibility assays. Phage susceptibility was analyzed by the cross-streak

method or by spotting phage directly onto a lawn of *P. aeruginosa*. For the cross-streak

method, P. aeruginosa strains were grown for 16 h in LB before diluting 1:100 in LB and

then growing to an OD_{600} of 0.5-0.7. Then, 4 µl of phage DMS3 vir strain was spotted on

the side of an LB agar plate and then dragged across the agar surface in a straight line

before allowing to absorb into the agar. Once absorbed, 4 µl of each P. aeruginosa

strain was spotted at the top of the agar placed and dragged downward in a straight line

through the phage line in a straight line. Plates were incubated at

Alternatively, one percent M8 agar plates (60 x15 mm) containing 0.2% glucose

(v/v), 0.5% casamino acids (v/v), and 1 mM MgSO$_4$ were prepared and cooled to room

temperature. Fifty microliters of *P. aeruginosa* overnight cultures were added to 1 mL of

0.5% warm top agar (M8 medium and supplements). The mixture was gently mixed and

quickly poured onto M8 agar plates. Plates were swirled to ensure even spreading of

top agar. Once cooled, 2 µl of phage DMS3 vir strain were spotted onto the center of

the plate and allowed to dry before incubating plates at 37°C overnight.

Congo red assay. Cultures were grown in 5 ml LB for 16 h (overnight) at 37°C with

rolling. Overnight cultures were washed once and resuspended in sterile dH2O.

Washed cells were spotted (3 µl) onto Congo Red plates (1% tryptone [w/v], 1.5% agar,

40 µg/ml Congo Red, 20 µg/ml Coomassie Blue). Plates were grown for 16 h at 37°C

and then moved to room temperature for 3 days to allow for color development.

Statistical analysis. One-way ANOVA with multiple comparisons were performed

pairwise between all isolates and mixtures using the GraphPad Prism 6 software

(GraphPad, La Jolla, CA).

Acknowledgements. This work was supported by CFF HOGAN19G0 to DAH and

R37AI83256 and R01AI43730 to GAO. This work was also supported by P30DK117469

for the Applied Bioinformatics and Biostatistics Core. Sequencing services and

specialized equipment was provided by the Genomics and Molecular Biology Shared

Resource Core at Dartmouth supported by NCI Cancer Center Support Grant

5P30CA023108-37. Equipment used was supported by the NIH IDeA award to

Dartmouth BioMT P20GM113132. We thank Amy Baker for constructing the Δ*fliC*

mutant and Christine Toutain for construction of the *motABCD* deletion strain.

 Table 1. Strains used in this study

Strains (laboratory strain #)	Relevant genotype, description, or sequence	Source or reference
P. aeruginosa PA14 strains		
PA14 (DH123)	*P. aeruginosa* WT	(38)
WT att::*lacZ* (DH22)	WT with constitutive expression of *lacZ*	(39)
Δ*flgK* (DH1075)	*flgK*::Tn5B30(Tc), non-motile	(4)
Δ*fliC* (DH3543)	Unmarked deletion of *fliC*	This study
Δ*motABCD* (SMC7230)	Unmarked deletion of *motABCD* with pMQ72 EV	This study
Δ*pilMNOP* (DH2705)	Unmarked deletion of *pilMNOP*	(6)
Δ*pilA* (DH2636)	Unmarked deletion of *pilA*	(22)
Δ*pelA* (DH97/SMC2893)	Unmarked deletion of *pelA*	(40)
Δ*flgKpelA* (DH3541)	Unmarked deletion of *flgK* and *pelA*	(11)
Δ*flgKpilA* (DH3539)	Unmarked deletion of *flgK* and *pilA*	(11)
Δ*flgKpilApelA* (DH3542)	Unmarked deletion of *flgK*, *pilA*, and *pelA*	(11)
Δ*pilT* (DH3591/SMC7302)	Unmarked deletion of *pilT*	(41)
Δ*pilU* (DH3592/SMC7304)	Unmarked deletion of *pilU*	(41)
Δ*pilJ* (DH2637/SMC2992)	Unmarked deletion of *pilJ*	(42)
Δ*fimS* (SMC6967)	Unmarked deletion of *fimS*	(6)
Δ*cyaAB* (DH2648/SMC6707)	Unmarked deletion of *cyaAB*	(6)

Δ*vfr* (DH2701;SMC6722)   Unmarked deletion of *vfr*   (6)

References

1. Hogan DA, Kolter R. 2002. *Pseudomonas-Candida* interactions: An ecological role for virulence factors. Science 296:2229-2232.

2. Limoli DH, Warren EA, Yarrington KD, Donegan NP, Cheung AL, O'Toole GA. 2019. Interspecies interactions induce exploratory motility in *Pseudomonas aeruginosa*. Elife 8:e47365.

3. Lewis KA, Baker AE, Chen AI, Harty CE, Kuchma SL, O'Toole GA, Hogan DA. 2019. Ethanol decreases *Pseudomonas aeruginosa* flagellar motility through the regulation of flagellar stators. J Bacteriol 201:e00285-17.

4. O'Toole GA, Kolter R. 1998. Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development. Mol Microbiol 30:295-304.

5. Zhao K, Tseng BS, Beckerman B, Jin F, Gibiansky ML, Harrison JJ, Luijten E, Parsek MR, Wong GCL. 2013. Psl trails guide exploration and microcolony formation in *Pseudomonas aeruginosa* biofilms. Nature 497:388-391.

6. Luo Y, Zhao K, Baker AE, Kuchma SL, Coggan KA, Wolfgang MC, Wong GC, O'Toole GA. 2015. A hierarchical cascade of second messengers regulates *Pseudomonas aeruginosa* surface behaviors. MBio 6:e02456-14.

7. O'Toole GA, Wong GC. 2016. Sensational biofilms: Surface sensing in bacteria. Curr Opin Microbiol 30:139-146.

8. Lee CK, Vachier J, de Anda J, Zhao K, Baker AE, Bennett RR, Armbruster CR, Lewis KA, Tarnopol RL, Lomba CJ, Hogan DA, Parsek MR, O'Toole GA, Golestanian R, Wong GCL. 2020. Social cooperativity of bacteria during

reversible surface attachment in young biofilms: A quantitative comparison of

Pseudomonas aeruginosa PA14 and PAO1. mBio 11:e02644-19.

9.   Boyd CD, Smith TJ, El-Kirat-Chatel S, Newell PD, Dufrene YF, O'Toole GA.

2014. Structural features of the *Pseudomonas fluorescens* biofilm adhesin LapA

required for LapG-dependent cleavage, biofilm formation, and cell surface

localization. J Bacteriol 196:2775-88.

10.   Parsek MR. 2016. Controlling the connections of cells to the biofilm matrix. J

Bacteriol 198:12-4.

11.   Ribbe J, Baker AE, Euler S, O'Toole GA, Maier B. 2017. Role of cyclic di-GMP

and exopolysaccharide in type IV pilus dynamics. J Bacteriol 199:e00859-16.

12.   Ma L, Conover M, Lu H, Parsek MR, Bayles K, Wozniak DJ. 2009. Assembly and

development of the *Pseudomonas aeruginosa* biofilm matrix. PLoS Pathog

5:e1000354.

13.   Colvin KM, Irie Y, Tart CS, Urbano R, Whitney JC, Ryder C, Howell PL, Wozniak

DJ, Parsek MR. 2012. The Pel and Psl polysaccharides provide *Pseudomonas*

aeruginosa structural redundancy within the biofilm matrix. Environ Microbiol

14:1913-28.

14.   Wei Q, Ma LZ. 2013. Biofilm matrix and its regulation in *Pseudomonas*

aeruginosa. Int J Mol Sci 14:20983-1005.

15.   Anyan ME, Amiri A, Harvey CW, Tierra G, Morales-Soto N, Driscoll CM, Alber

MS, Shrout JD. 2014. Type IV pili interactions promote intercellular association

and moderate swarming of *Pseudomonas aeruginosa*. Proc Natl Acad Sci U S A

111:18013-8.

16.   Lozano C, Azcona-Gutierrez JM, Van Bambeke F, Saenz Y. 2018. Great

phenotypic and genetic variation among successive chronic *Pseudomonas*

aeruginosa from a cystic fibrosis patient. PLoS One 13:e0204167.

17.   Mahenthiralingam E, Campbell ME, Speert DP. 1994. Nonmotility and phagocytic

resistance of *Pseudomonas aeruginosa* isolates from chronically colonized

patients with cystic fibrosis. Infect Immun 62:596-605.

18.   Workentine ML, Sibley CD, Glezerson B, Purighalla S, Norgaard-Gron JC,

Parkins MD, Rabin HR, Surette MG. 2013. Phenotypic heterogeneity of

Pseudomonas aeruginosa populations in a cystic fibrosis patient. PLoS One

8:e60225.

19.   Turner KH, Everett J, Trivedi U, Rumbaugh KP, Whiteley M. 2014. Requirements

for *Pseudomonas aeruginosa* acute burn and chronic surgical wound infection.

PLoS Genet 10:e1004518.

20.   Boyle KE, Monaco HT, Deforet M, Yan J, Wang Z, Rhee K, Xavier JB. 2017.

Metabolism and the evolution of social behavior. Mol Biol Evol 34:2367-2379.

21.   Shrout JD, Chopp DL, Just CL, Hentzer M, Givskov M, Parsek MR. 2006. The

impact of quorum sensing and swarming motility on *Pseudomonas aeruginosa*

biofilm formation is nutritionally conditional. Mol Microbiol 62:1264-77.

22.   Kuchma SL, Ballok AE, Merritt JH, Hammond JH, Lu W, Rabinowitz JD, O'Toole

GA. 2010. Cyclic-di-GMP-mediated repression of swarming motility by

Pseudomonas aeruginosa: The *pilY1* gene and its impact on surface-associated

behaviors. J Bacteriol 192:2950-64.

23.   Kuchma SL, Griffin EF, O'Toole GA. 2012. Minor pilins of the type IV pilus

392   system participate in the negative regulation of swarming motility. J Bacteriol

393   194:5388-403.

394   24.   Shen Y, Siryaporn A, Lecuyer S, Gitai Z, Stone HA. 2012. Flow directs surface-

395   attached bacteria to twitch upstream. Biophys J 103:146-51.

396   25.   Landry RM, An D, Hupp JT, Singh PK, Parsek MR. 2006. Mucin-*Pseudomonas*

397   *aeruginosa* interactions promote biofilm formation and antibiotic resistance. Mol

398   Microbiol 59:142-51.

399   26.   Tammam S, Sampaleanu LM, Koo J, Sundaram P, Ayers M, Chong PA, Forman-

400   Kay JD, Burrows LL, Howell PL. 2011. Characterization of the PilN, PilO and PilP

401   type IVa pilus subcomplex. Mol Microbiol 82:1496-514.

402   27.   Webster SS, Lee CK, Schmidt WC, Wong GCL, O'Toole GA. 2021. Interaction

403   between the type 4 pili machinery and a diguanylate cyclase fine-tune c-di-GMP

404   levels during early biofilm formation. Proc Natl Acad Sci U S A

405   118:e2105566118.

406   28.   Burrows LL. 2005. Weapons of mass retraction. Mol Microbiol 57:878-88.

407   29.   Chiang P, Burrows LL. 2003. Biofilm formation by hyperpiliated mutants of

408   *Pseudomonas aeruginosa*. J Bacteriol 185:2374-8.

409   30.   Whitchurch CB, Mattick JS. 1994. Characterization of a gene, *pilU*, required for

410   twitching motility but not phage sensitivity in *Pseudomonas aeruginosa*. Mol

411   Microbiol 13:1079-1091.

412   31.   Chen AI, Dolben EF, Okegbe C, Harty CE, Golub Y, Thao S, Ha DG, Willger SD,

413   O'Toole GA, Harwood CS, Dietrich LE, Hogan DA. 2014. *Candida albicans*

ethanol stimulates *Pseudomonas aeruginosa* WspR-controlled biofilm formation

as part of a cyclic relationship involving phenazines. PLoS Pathog 10:e1004480.

32.   Merritt JH, Brothers KM, Kuchma SL, O'Toole GA. 2007. SadC reciprocally

influences biofilm formation and swarming motility via modulation of

exopolysaccharide production and flagellar function. J Bacteriol 189:8154-64.

33.   Whitfield GB, Marmont LS, Bundalovic-Torma C, Razvi E, Roach EJ, Khursigara

CM, Parkinson J, Howell PL. 2020. Discovery and characterization of a Gram-

positive Pel polysaccharide biosynthetic gene cluster. PLoS Pathog

16:e1008281.

34.   Darch SE, McNally A, Harrison F, Corander J, Barr HL, Paszkiewicz K, Holden S,

Fogarty A, Crusz SA, Diggle SP. 2015. Recombination is a key driver of genomic

and phenotypic diversity in a *Pseudomonas aeruginosa* population during cystic

fibrosis infection. Sci Rep 5:7649.

35.   Head NE, Yu H. 2004. Cross-sectional analysis of clinical and environmental

isolates of *Pseudomonas aeruginosa*: Biofilm formation, virulence, and genome

diversity. Infect Immun 72:133-44.

36.   Grosso-Becerra MV, Santos-Medellin C, Gonzalez-Valdez A, Mendez JL,

Delgado G, Morales-Espinosa R, Servin-Gonzalez L, Alcaraz LD, Soberon-

Chavez G. 2014. *Pseudomonas aeruginosa* clinical and environmental isolates

constitute a single population with high phenotypic diversity. BMC Genomics

15:318.

37.   Ha DG, Richman ME, O'Toole GA. 2014. Deletion mutant library for investigation

of functional outputs of cyclic diguanylate metabolism in *Pseudomonas*

aeruginosa PA14. Appl Environ Microbiol 80:3384-93.

38.   Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM. 1995.

Common virulence factors for bacterial pathogenicity in plants and animals.

Science 268:1899-902.

39.   Clay ME, Hammond JH, Zhong F, Chen X, Kowalski CH, Lee AJ, Porter MS,

Hampton TH, Greene CS, Pletneva EV, Hogan DA. 2020. *Pseudomonas*

aeruginosa lasR mutant fitness in microoxia is supported by an Anr-regulated

oxygen-binding hemerythrin. Proc Natl Acad Sci U S A 117:3167-3173.

40.   Friedman L, Kolter R. 2004. Genes involved in matrix formation in *Pseudomonas*

aeruginosa PA14 biofilms. Mol Microbiol 51:675-90.

41.   Lee CK, de Anda J, Baker AE, Bennett RR, Luo Y, Lee EY, Keefe JA, Helali JS,

Ma J, Zhao K, Golestanian R, O'Toole GA, Wong GCL. 2018. Multigenerational

memory and adaptive adhesion in early bacterial biofilm communities. Proc Natl

Acad Sci U S A 115:4471-4476.

42.   Belete B, Lu H, Wozniak DJ. 2008. *Pseudomonas aeruginosa* AlgR regulates

type IV pilus biosynthesis by activating transcription of the *fimU-pilVWXY1Y2E*

operon. J Bacteriol 190:2023-30.

Figure Legends

Fig. 1. Motility heterogeneity represses swarming motility independent of competition. A. Representative images from swarm assay of WT *P. aeruginosa* strain PA14 (motile) mixed with Δ*flgK* (non-motile) at the indicated ratios on M8 agar. In this and all figure panels in the manuscript, WT and derivatives are labeled in black, while non-motile mutants are labeled in red. **B.** Competition growth assay between WT PA14 tagged with a *lacZ* reporter (gray) and either an untagged WT strain (black) or Δ*flgK* mutant (red). Error bars represent the standard deviation between the replicate values. Statistical analysis using one-way ANOVA showed no difference between input and output for each ratio analyzed. **C.** Representative images from swarm assay of WT PA14 (motile) mixed with Δ*flgK* (non-motile), Δ*fliC* (non-motile), or Δ*motABCD* (non-motile, paralyzed flagellum) in the indicated ratios on M8 agar.

Fig. 2. Non-motile *Pseudomonas* requires T4P to repress flagellum-mediated motility of motile strains on soft agar. A. Representative images of swarm assays for wild-type *P. aeruginosa* strain PA14 (motile, Pel+, T4P+) mixed with Δ*flgK* (non-motile, Pel+, T4P+) or Δ*flgKpilA* (non-motile, Pel+, T4P-) on M8 agar at the indicated ratios. **B.** Representative image showing the interaction between WT (black dot), Δ*flgK* (red dots), and Δ*flgKpilA* (purple dots) for swimming motility in soft agar (0.3% M63 agar) after 42 h incubation. The colored dots indicate the points of inoculation for the respective strains.

Fig. 3. Motile *Pseudomonas* requires functional T4P to be repressed by the non-motile subpopulation. Representative images from swarm assays of the Δ*flgK* mutant

(functional T4P) mixed with either WT, the Δ*pilA* mutant (lacking T4P function), the Δ*pilT*

mutant (no T4P retraction), or the Δ*pilU* mutant (reduced T4P retraction) at the indicated

ratios on M8 agar.

Fig. 4. Pel matrix is only required by the non-motile *Pseudomonas* subpopulation

in order to repress overall motility. A. Representative images from swarm assays of

Δ*flgK* (non-motile and Pel+) mixed with WT (motile and Pel+) or Δ*pelA* (motile and Pel-)

and of WT mixed with Δ*flgK*Δ*pelA* (non-motile and Pel-) at the indicated ratios on M8

agar. **B.** Representative images of WT, Δ*pilA,* Δ*pelA,* Δ*flgK,* Δ*flgKpilA*, Δ*flgKpelA,* and

Δ*flgKpilApelA* grown on Congo Red plates to assess Pel production. WT and derivatives

are labeled in black while Δ*flgK* and derivatives are labeled in red.

Fig. 5. Summary model describing the components needed for repression of

motility by non-motile cells in *P. aeruginosa* strain PA14. We show that a

subpopulation of non-motile *P. aeruginosa* cells (red) in co-culture with motile cells

(black) impedes flagellum-mediated swarming motility in *P. aeruginosa* cells in a manner

that is dependent on retractable T4P. We also show that Pel matrix, only from the non-

motile strain, is required. PilU, PilJ, FimS, CyaAB, and Vfr were not required for

repression of motility. These data may indicate ways that *P. aeruginosa* cells come

together during biofilm formation.

Fig. 1. Motility heterogeneity represses swarming motility independent of competition. A. Representative images from swarm assay of WT *P. aeruginosa* strain PA14 (motile) mixed with Δ*flgK* (non-motile) at the indicated ratios on M8 agar. In this and all figure panels in the manuscript, WT and derivatives are labeled in black, while non-motile mutants are labeled in red. **B.** Competition growth assay between WT PA14 tagged with a *lacZ* reporter (gray) and either an untagged WT strain (black) or Δ*flgK* mutant (red). Error bars represent the standard deviation between the replicate values. Statistical analysis using one-way ANOVA showed no difference between input and output for each ratio analyzed. **C.** Representative images from swarm assay of WT PA14 (motile) mixed with Δ*flgK* (non-motile), Δ*fliC* (non-motile), or Δ*motABCD* (non-motile, paralyzed flagellum) in the indicated ratios on M8 agar.

Fig. 2. Non-motile *Pseudomonas* requires T4P to repress flagellum-mediated motility of motile strains on soft agar. A. Representative images of swarm assays for wild-type *P. aeruginosa* strain PA14 (motile, Pel+, T4P+) mixed with Δ*flgK* (non-motile, Pel+, T4P+) or Δ*flgKpilA* (non-motile, Pel+, T4P-) on M8 agar at the indicated ratios. **B.** Representative image showing the interaction between WT (black dot), Δ*flgK* (red dots), and Δ*flgKpilA* (purple dots) for swimming motility in soft agar (0.3% M63 agar) after 42 h incubation. The colored dots indicate the points of inoculation for the respective strains.

Fig. 3. Motile *Pseudomonas* requires functional T4P to be repressed by the non-motile subpopulation. Representative images from swarm assays of △*flgK* mutant (functional T4P) mixed with either WT, the △*pilA* mutant (lacking T4P function), the △*pilT* mutant (no T4P retraction), or the △*pilU* mutant (reduced T4P retraction) at the indicated ratios on M8 agar.

Fig. 4. Pel matrix is only required by the non-motile *Pseudomonas* subpopulation in order to repress overall motility. A. Representative images from swarm assays of Δ*flgK* (non-motile and Pel+) mixed with WT (motile and Pel+) or Δ*pelA* (motile and Pel-) and of WT mixed with Δ*flgK*Δ*pelA* (non-motile and Pel-) at the indicated ratios on M8 agar. **B.** Representative images of WT, Δ*pilA,* Δ*pelA*, Δ*flgK,* Δ*flgKpilA*, Δ*flgKpelA,* and Δ*flgKpilApelA* grown on Congo Red plates to assess Pel production. WT and derivatives are labeled in black while Δ*flgK* and derivatives are labeled in red.

Fig. 5. Summary model describing the components needed for repression of motility by non-motile cells in *P. aeruginosa* strain PA14. We show that a subpopulation of non-motile *P. aeruginosa* cells (red) in co-culture with motile cells (black) impedes flagellum-mediated swarming motility in *P. aeruginosa* cells in a manner that is dependent on retractable T4P. We also show that Pel matrix, only from the non-motile strain, is required. PilU, PilJ, FimS, CyaAB, and Vfr were not required for repression of motility. These data may indicate ways that *P. aeruginosa* cells come together during biofilm formation.