Non-motile subpopulations of *Pseudomonas aeruginosa* repress flagellar motility

in motile cells through a type IV pili- and Pel-dependent mechanism

Kimberley A. Lewis[a,†], Danielle M. Vermilyea[a,†], Shanice S. Webster[a], Christopher J.

Geiger[a], Jaime de Anda[b,c,d], Gerard C. L. Wong[b,c,d], George A. O'Toole[a] and Deborah A.

Hogan[a,#]

[a]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth,

Hanover, New Hampshire, United States of America.

[b]Department of Bioengineering, University of California Los Angeles, CA 90095

[c]Department of Chemistry and Biochemistry, University of California Los Angeles, CA

90095

[d]California NanoSystems Institute, University of California Los Angeles, CA 90095

[†]Kimberley A. Lewis and Danielle M. Vermilyea contributed equally. K.A.L.

conceptualized the project, generated most data for the first submission, and

contributed to manuscript preparation for first submission. D.M.V. generated data and

contributed to manuscript preparation for the first submission and was primarily

responsible for data generation and manuscript preparation for the second submission.

Author order was determined by who conceptualized and started the project.

Running Head: Non-motile *P. aeruginosa* represses flagellar motility

#To whom correspondence should be addressed

Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth

Vail Building HB7550

Hanover, NH 03755

E-mail: dhogan@dartmouth.edu

Key words: swarming, *Pseudomonas aeruginosa*, motility, type IV pili, Pel, microbe-

microbe interaction

Abstract

The downregulation of *Pseudomonas aeruginosa* flagellar motility is a key event in

biofilm formation, host-colonization, and the formation of microbial communities, but the

external factors that repress motility are not well understood. Here, we report that under

swarming conditions, swarming motility can be repressed by cells that are non-motile

due to the absence of a flagellum or flagellar rotation. Non-swarming cells, due to

mutations that prevent either flagellum biosynthesis or rotation, present at 5% of the

total population suppressed swarming of wild-type cells under the conditions tested in

this study. Non-swarming cells required functional type IV pili and the ability to produce

Pel exopolysaccharide to suppress swarming by the flagellated wild type. In contrast,

flagellated cells required only type IV pili, but not Pel production, in order for swarming

to be repressed by non-flagellated cells. We hypothesize that interactions between

motile and non-motile cells may enhance the formation of sessile communities including

those involving multiple genotypes, phenotypically-diverse cells, and perhaps other

species.

Importance

Our study shows that, under the conditions tested, a small population of non-swarming

cells can impact the motility behavior of the larger population. The interactions that lead

to the suppression of swarming motility require type IV pili and a secreted

polysaccharide, two factors with known roles in biofilm formation. These data suggest

that interactions between motile and non-motile cells may enhance the transition to

sessile growth in populations and promote interactions between cells with different

genotypes.

Introduction

Microbial localization through processes other than cell-division are critical for the formation of spatially structured populations and communities. Thus, motility and its regulation by diverse chemical and physical stimuli are major drivers of intraspecies and interspecies microbial interactions (1, 2). Previously, we found that exogenous ethanol, a common product secreted by many microbes, markedly reduced *Pseudomonas aeruginosa* swimming and swarming motilities at the population level, but only decreased the motile fraction of cells in the population by 16% (3). Prompted by this finding, we sought to determine the impact of a subpopulation of non-motile cells on the motility of the larger motile *P. aeruginosa* population.

During *P. aeruginosa* biofilm formation, motile cells are recruited to microcolonies of sessile cells via a combination of processes that may differ between strains with distinct early-biofilm forming strategies (4, 5) or in response to different experimental conditions. For *P. aeruginosa* strain PA14, retractile type IV pili (T4P) participate in surface sensing, which results in the upregulation of cAMP and the subsequent induction of cyclic-di-GMP (6-8); flagellar motility is then down-regulated in order to facilitate surface attachment (6, 7, 9, 10) followed by Pel exopolysaccharide matrix production (11), which can connect cells to one another (10, 12). Both matrix production and T4P function have been shown to participate in motility repression (10, 12-14) suggesting that matrix materials may mediate physical interactions between neighboring cells. T4P also mediate cell-cell interactions in swarms (15).

Boyle et al. (16) previously showed that a Δ*flgK* mutant lacking a flagellum repressed swarming (i.e., decreased the swarm radius) when in excess (83%) of the

flagellated wild type (WT) (17%). In support of the observation that a non-motile

subpopulation can suppress population-wide swarming, our previously published

findings (3) showed that while WT strain PA14 was capable of swarming, an average of

only 38% of flagellated cells were motile at any given time. Furthermore, we showed

that ethanol strongly repressed *P. aeruginosa* swarming motility by reducing the motile

fraction of the population from 38% to 22%; in other words, a 16% reduction in the

motile fraction was sufficient to repress swarming by the entire population (3). Taken

together, these findings show that a non-motile subpopulation is capable of repressing

swarming in the entire population. However, how non-motile cells, such as those lacking

a flagellum, suppress swarming of flagellated strains has not been elucidated.

In this study, we show that non-motile cells lacking a flagellum, when added to a

population at 5 to 75% of the total inoculum, resulted in repression of swarming by WT

cells. To repress flagellar motility, non-motile cells required the ability to produce Pel

polysaccharide, and both motile and non-motile cells required functional T4P for this

interaction. These data have implications for factors that contribute to population-level

behaviors and intra- and inter-species interactions. Interactions that promote cell-cell

interactions may be relevant in situations where non-motile and motile cells can be

found together such as in cystic fibrosis (CF)-associated lung infections (17-19),

wounds (20), and ear infections (18).

Results

The presence of non-flagellated cells represses swarming motility by *P.*

***aeruginosa* strain PA14.** To explore the effects of a subpopulation of non-motile cells

on swarming motility, mixes containing different proportions of flagellated *P. aeruginosa*

strain PA14 (WT) cells and non-flagellated Δ*flgK* mutant cells that lack the flagellar hook

protein were plated on 0.5% M8 agar, which supports swarming motility by the WT

strain. The percent of Δ*flgK* mutant cells ranged from 5% to 75% in these mixes, and

single strains were included as controls. As expected, cultures with 100% WT swarmed

readily while cultures with 100% Δ*flgK* showed no swarming (**Fig. 1A**). We found that

swarming motility was completely repressed by the addition of 5-75% Δ*flgK* cells under

the conditions tested (**Fig. 1A** and **Fig. 1C**). The addition of 1-4% Δ*flgK* was insufficient

to repress population-wide swarming (**Fig. S1**). It is worth noting that a swarming

phenotype can be variable between both technical replicates and biological replicates

(21, 22). While Δ*flgK* added at 5% of the population regularly repressed WT swarming,

the lower limit was variable (**Fig. S1, Fig. S5**). However, work by Boyle et al. (16)

reported that Δ*flgK* repressed swarming only when it was in excess of the WT.

Specifically, the authors reported repression as measured by a reduction in swarm

radius when there was 17% WT and 83% Δ*flgK*. Images for 1:1 WT:Δ*flgK* mixes may

also suggest effects on the swarm area even though swarm radius was not yet affected.

Therefore, while the lower limit of our assay was variable, Δ*flgK* repressed WT

swarming at significantly lower concentrations than previously reported and did not

depend on being in excess of the WT.

To determine if the lack of swarming in WT:Δ*flgK* co-cultures was due to faster

growth of the non-flagellated cells compared to the WT, we competed the WT or the

Δ*flgK* mutant against a WT strain tagged with a *lacZ* reporter at different ratios atop a

0.22 μm filter placed on 0.5% M8 agar. After 16 h, the colony was disrupted and colony

forming units (CFUs) for each strain were enumerated on blue/white screening plates

that contained X-Gal. At all ratios tested, neither untagged WT nor the $\Delta flgK$ mutant

could outgrow the *lacZ*-expressing WT strain (**Fig. 1B**), indicating that the lack of

swarming observed in WT:$\Delta flgK$ mixes was not due to increased growth of non-

flagellated cells and thus overgrowth of the population. Rather, our data suggest that a

subpopulation of cells incapable of flagellar motility inhibits motility of the larger

flagellated WT population.

Using conditions in which the non-motile strain comprised 25%, 50%, or 75% of

the population, we also found that, like the $\Delta flgK$ mutant, the $\Delta fliC$ mutant, another strain

incapable of flagellar motility due to the lack of the flagellin protein used to make the

flagellar filament, inhibited WT swarming at all ratios tested (**Fig. 1C**). Similarly, the

$\Delta motABCD$ mutant, which produces a flagellum that is unable to rotate due to the

absence of stators, repressed swarming of the WT; although, in comparison to $\Delta flgK$

and $\Delta fliC$, the $\Delta motABCD$ mutant was slightly less effective at repressing WT swarming

(**Fig. 1C**). Together, these data suggest that the absence of flagellar motility in a

subpopulation of cells is sufficient to suppress WT swarming in co-culture.

Functional T4P in flagellated cells are required for non-flagellated cells to repress

swarming in co-culture. Several studies have shown that *P. aeruginosa* strains that

are deficient in T4P tend to have a hyper-swarming phenotype (23-25), while

hyperpiliated strains (those that overproduce elongated pili due to a defect in retraction)

have inhibited swarming (15, 26). Additionally, *P. aeruginosa* cells have been shown to

interact via their T4P to form close cell associations that facilitate directional swarming

(15). Therefore, we determined if T4P are required by flagellated cells to facilitate the

ability of Δ*flgK* to inhibit population-wide swarming motility. To do this, we analyzed the

ability of Δ*flgK* to repress swarming when co-cultured with the following mutants: Δ*pilA*,

which lacks T4P; Δ*pilMNOP*, which lacks the alignment complex required for T4P

function (27) and cyclic-di-GMP signaling (28); and both Δ*pilT* and Δ*pilU*, hyperpiliated

mutants that lack either of the ATPases that mediate T4P retraction (29). Strains lacking

pilA, *pilMNOP*, *pilT*, and *pilU* were all defective for twitching motility (**Fig. S2** and **Fig.**

S3). The swarming motility of strains lacking *pilA* or *pilMNOP* was no longer inhibited by

the Δ*flgK* mutant in co-culture (**Fig. 2** and **Fig. S4**). Interestingly, despite its reported

hyperpiliation and in contrast to previous reports (15), Δ*pilU* was capable of swarming

and swarming motility was inhibited by the Δ*flgK* mutant in co-culture like the WT (**Fig.**

2, Fig. S5). However, the hyperpiliated Δ*pilT* mutant was incapable of swarming (**Fig.**

2), in agreement with previous models (15). Complementation of the Δ*pilU* and Δ*pilT*

mutants restored twitching in both strains as well as swarming in Δ*pilT* (**Fig. S6**).

While PilT is the main retractile ATPase, PilU has an accessory role in retraction;

the Δ*pilT* mutant completely lacks pilus retraction (30, 31), while the Δ*pilU* mutant has

reduced retraction but still retains some function (31). This accessory role is supported

by previous observations that the Δ*pilU* mutant retains sensitivity to infection by phage

PO4, while the Δ*pilT* mutant became resistant to infection (26). We confirmed that the

WT and Δ*pilU* mutant were sensitive to phage DMS3 infection and that the Δ*pilA* and

Δ*pilT* mutants were resistant to infection under our experimental conditions (**Fig. S2B**).

Additionally, while the Δ*pilU* mutant can form dense biofilms under static conditions, it is

unable to form a biofilm under flow conditions likely due to Δ*pilU* T4P being more fragile

than WT or even Δ*pilT* T4P, allowing for Δ*pilU* T4P to readily detach from the cell

surface (26, 30). Overall, the data show that Δ*pilU* displays an intermediate phenotype

in that it is hyperpiliated due to a reduction in T4P retraction that still results in inhibition

of T4P-mediated twitching motility, but not swarming motility. Together, these data

indicate that motile cells require T4P, and likely functional T4P, for the non-flagellated

subpopulation to inhibit the swarming motility of the WT strain when grown in co-culture.

P. aeruginosa T4P participate in a cAMP-dependent signaling pathway (see **Fig.**

S7A for pathway) that involves FimS, PilJ, CyaAB adenylate cyclases, and the cAMP-

binding transcription factor Vfr (6). We found that this pathway was not required for the

repression of WT swarming motility by the Δ*flgK* mutant (**Fig. S7B**) using published

mutants lacking components of the cAMP signaling pathway (Δ*pilJ*, Δ*fimS*, Δ*cyaAB*, and

Δ*vfr*) that retain expression of functional T4P as evidenced by their sensitivity to phage

DMS3 (**Fig. S7C**). Taken together, the data show that flagellated WT cells require

functional T4P, but not the cAMP-dependent surface-sensing pathway, in order for non-

flagellated cells to be able to repress swarming in co-culture.`

Non-flagellated cells require functional T4P to repress swarming motility of

flagellated cells in co-culture. Consistent with published results (4, 32, 33), the Δ*flgK*

mutant has functional T4P as evidenced by the formation of a large twitching motility

zone using the agar-plastic interface assay (**Fig. S2A** and **Fig. S3**). While both the WT

and Δ*flgK* formed twitch zones that were significantly larger than those formed by the

T4P-null Δ*pilA* mutant, the Δ*flgK* twitch zone was ~25% smaller than that formed by the

WT (p<0.05; **Fig. S3A**). Using Δ*flgK* and Δ*flgK*Δ*pilA* strains, we found that *pilA*, which

encodes the major pilin component of T4P, was required for the Δ*flgK* mutant to

suppress WT swarming on 0.5% M8 agar (**Fig. 3A**). To assess whether Δ*flgK* cells

needed functional pili to repress the WT, Δ*flgK* lacking either *pilU* (reduced T4P

retraction) or *pilT* (no T4P retraction) were mixed with the WT. At all ratios tested,

Δ*flgK*Δ*pilU* and Δ*flgK*Δ*pilT* were no longer able to repress WT swarming, like Δ*flgK*Δ*pilA,*

indicating that the Δ*flgK* mutant required fully functional pili to repress population-wide

swarming (**Fig. 3A, Fig. S8**). Complementation of *pilU* and *pilT* in Δ*flgK*Δ*pilU* and

Δ*flgK*Δ*pilU*, respectively, was confirmed to restore twitching motility (**Fig. S9**); the effect

of complementation on swarming motility was not assessed given that Δ*flgK* does not

swarm. In a similar assay using 0.3% M63 agar, which supports both flagellum-

mediated swimming and swarming, we found that inoculated spots of Δ*flgK* cells (**Fig.**

3B, red dots) decreased the local expansion of the motile WT population, while spots of

the Δ*flgK*Δ*pilA* mutant (**Fig. 3B, purple dots**) did not. In contrast to the effects of Δ*flgK*

cells on flagellar motility, the Δ*flgK* mutant did not alter T4P-dependent twitching motility

of the WT in a 50:50 ratio when compared to the WT alone (**Fig. S3B**). Therefore, the

data show that functional T4P are required for non-flagellated cells to inhibit swarming

of WT cells in co-culture.

Non-flagellated *P. aeruginosa* requires Pel matrix production to repress swarming

motility in the flagellated population. We next explored whether Pel matrix production

played a role in swarming repression by using a Δ*pelA* mutant, which is a mutant

capable of robust swarming but lacks PelA-mediated deacetylase and hydrolase

activities and, subsequently, secretion of properly modified Pel polysaccharide (34). We

found that swarming by the ΔpelA mutant was repressed by ΔflgK in co-culture like the

WT (**Fig. 4A**). In contrast, ΔflgKΔpelA did not repress WT swarming (**Fig. 4A**). Flagellar

mutants, such as ΔfliC, have been reported to overexpress Pel and Psl polysaccharides

(35). Both ΔflgK and ΔfliC, which repress WT swarming (**Fig. 1**), had increased Congo

Red binding, which is consistent with increased Pel or exopolysaccharide production

(**Fig. 4B**, **Fig. S10**). Additionally, while ΔflgKΔpilA (no T4P), ΔflgKΔpilU (reduced T4P

retraction), and ΔflgKΔpilT (no T4P retraction) no longer repressed population-wide

swarming motility (**Fig. 2A**), this does not appear to be due to a change in Pel

production as all ΔflgK mutants displayed an increase in Congo Red binding (Pel

production) (**Fig. 4B**, **Fig. S10**). Together, these data show that the non-flagellated

strain needs to produce Pel matrix to repress swarming motility, but the flagellated

strain does not.

DISCUSSION

Based on the data presented here we propose a model (**Fig. 5**) whereby a

subpopulation of cells defective in flagellar motility limit swarming motility of the entire

population. Our analyses revealed that non-flagellated cells required T4P and Pel

polysaccharide to inhibit the larger swarming population, and that flagellated cells

required functional T4P but not the Vfr/cAMP signaling system to respond to the non-

flagellated population. These findings in regards to T4P and Pel are consistent with the

recruitment of cells to microcolonies during biofilm development and aggregate

formation, which has been reported in numerous contexts (e.g. (4)). While cAMP was

not required for repression of swarming, our previous findings showed that ethanol

decreased swarming motility by increasing c-di-GMP (3). Future studies will investigate

the role of c-di-GMP levels in the Δ*flgK*-mediated repression of WT swarming.

Furthermore, given that Δ*flgK* requires retractable T4P and Pel to repress WT

swarming, and overproduces Pel compared to the WT, future studies will test how

varying the amount of Pel and TFP in WT and mutant populations contribute to

repression of population-wide swarming.

The involvement of T4P in swarming cells was previously reported by Anyan et

al. (15) who reported that T4P impact cell-cell interactions in swarms in ways that limit

the movement of cells away from the population front. T4P mutants have been shown to

have increased swarming motility, suggesting that the intercellular T4P interactions that

we report between flagellated and non-flagellated genotypes may also be occurring in

single-strain swarms in which all cells have the potential for flagellar motility (24, 25).

Future studies will determine if T4P play direct roles in cell-cell interactions or indirect

roles in structuring the community.

The WT and Δ*pelA* mutant responded similarly to non-flagellated cells while

Δ*flgKpelA* was no longer able to repress motility suggesting that not all cells need to

produce exopolysaccharide matrix to repress swarming. Our findings that Pel

production is only necessary in the non-flagellated subpopulation is supported by

published data showing that ethanol not only decreases *P. aeruginosa* flagellar motility

(3), but increases Pel production (36). Pel has been suggested to repress motility via

the steric hinderance of flagella (3, 37), but further studies are needed to test this

hypothesis. The involvement of Pel in the interaction between flagellated and non-

flagellated cells is particularly interesting in light of recent work by Whitfield et al. (38)

which found that the Pel biosynthetic locus is widespread across Gram-positive and

Gram-negative bacteria. Future studies will determine if Pel is specifically detected by a

T4P-dependent mechanism or if other matrix materials play a similar role.

Boyle et al. (16) previously reported that Δ*flgK* was capable of repressing WT

swarming when Δ*flgK* was added 5:1 or at 83% of the population, while we show that

Δ*flgK* at as low as 5% of the population represses WT swarming. Our previously

published findings (3) also showed that while strain PA14 was capable of swarming, an

average of 38% of flagellated cells were motile at any given time. Furthermore, we

showed that ethanol strongly repressed *P. aeruginosa* swarming motility by reducing the

motile fraction of the population from 38% to 22% (3). Overall, the data show that

motility of individual flagellated WT cells is heterogeneous within a population, but the

addition of non-flagellated cells represses population-wide motility. These findings

suggest that the number of non-flagellated cells necessary to repress WT motility may

vary due to experimental conditions such as medium type and composition.

In the studies here, we seed a low percentage of non-swarming cells into a

swarming population, which is reminiscent of the situations where there are genetically

heterogeneous populations observed in the CF airway or other chronic infections (19,

39, 40) as well as during normal environmental growth (40, 41). Additionally, clinical *P.*

aeruginosa isolates from the CF lung have diverse motility phenotypes (19). This

diversity in motility is even observed within a single patient (19). We hypothesize that

such a mechanism promotes interspecies and intraspecies interactions. The strong

implication of these findings is that in a mixed population with a subpopulation of non-

flagellated cells, the entire population can be brought to a halt. Similarly, within a

population in which a significant number of cells have stopped swimming or swarming,

the further addition of even just a few non-flagellated cells can suppress motility for the

entire population (i.e., see the ethanol studies mentioned above). As *P. aeruginosa*

populations inherently have some number of non-motile or inactive cells (3), future

studies are required to determine how strain background or mutant genotype affects

phenotypic heterogeneity among isogenic cells.

Materials and Methods

Strains and media. Strains used in this study are listed in **Table 1**. *P. aeruginosa* strain

PA14 and derivatives were routinely cultured on lysogeny broth (LB) solidified with 1.5%

agar or in LB broth at 37°C with shaking. For *P. aeruginosa* phenotypic assays, either

M63 [22 mM KH_2PO_4, 40 mM K_2HPO_4, and 15 mM $(NH_4)_2SO_4$] or M8 (42 mM Na_2PO_4,

22 mM KH_2PO_4, and 8.5 mM NaCl) minimal salts medium supplemented with $MgSO_4$ (1

mM), glucose (0.2%), and casamino acids (CAA; 0.5%) were used as indicated.

Complemented strains containing genes regulated by an arabinose-inducible promoter

(P_{BAD}) were grown in the presence of 0.25% arabinose.

Construction and complementation of mutant strains. Plasmids used in this study

are listed in **Table 1**. Plasmid constructs for making in-frame deletions and arabinose-

inducible complementation strains were constructed using a *Saccharomyces cerevisiae*

recombination technique described previously (42) or Gibson Assembly. All plasmids

were sequenced at the Molecular Biology Core at the Geisel School of Medicine at

Dartmouth. Plasmids were introduced into *P. aeruginosa* by conjugation via S17/lambda

pir *E. coli*. Merodiploids were selected by drug resistance and double recombinants

were obtained using sucrose counter-selection and genotype screening by PCR.

Swarming motility assays. Swarm assays were performed as previously described

(43). Briefly, 9 ml of M8 medium with 0.5% agar (swarm agar) was poured into 60 x 15

mm plates and allowed to dry at room temperature for 3 h prior to inoculation. The

indicated strains were grown for 16 h then each was washed in 1X PBS and then

normalized to an OD_{600} of 1. Indicated isolates were then mixed at the indicated ratios in

a final volume of 100 µl. Each plate was inoculated with 0.5 µl of the liquid culture

mixture and the plates incubated face-up at $37^{o}C$ in stacks of no more than four plates

for up to 22 h. Each mixture was inoculated in three to four replicates and was assessed

on at least three separate days. Images were captured using a Canon EOS Rebel T6i

camera and images assessed for swarm repression.

Swimming motility assays. Swim assays were performed as previously described

(43). Briefly, M63 medium solidified with 0.3% agar (soft agar) was poured into 100 x 15

mm petri plates and allowed to dry at room temperature (~$25^{o}C$) for 3 h prior to

inoculation. Sterile toothpicks or P20 pipette tips were used to inoculate bacterial mixes

into the center of the agar without touching the bottom of the plate; liquid cultures grown

for 16 h were used as inoculum. The indicated strains were normalized to an OD_{600} of 1

then were mixed at the indicated ratios in a final volume of 100 µl. No more than four

bacterial mixtures were assayed per plate. Plates were incubated upright at $37^{o}C$ in

stacks of no more than four plates per stack for 18 to 20 h.

Twitching motility assays. T-broth [1% tryptone (w/v), 0.5% NaCl (w/v)] medium

solidified with 1.5% agar was poured into petri plates and allowed to dry at room

temperature (~25°C) for 3 h. Overnight (16 h) cultures were washed once in 1X PBS

and then normalized to an OD_{600} of 1. Plates were inoculated by dipping a sterile

toothpick or P20 pipette tip into the washed and normalized cultures and then inserting

the toothpick into the agar until it touched the bottom of the plate. Plates were then

incubated at 37°C for 46 h after which time the agar was removed and the plate

incubated in 0.1% crystal violet for 10 min. Plates were then washed three times in

water and dried at room temperature. Images of the dried, stained twitch area were

taken and twitch diameter was measured.

Competition experiments. Competition assays were performed to determine relative

growth of selected *P. aeruginosa* strains. Strains were grown for 16 h and then 1 ml of

culture was pelleted at 15,682 *x g* for 2 min, and washed once in 1 ml 1X PBS followed

by resuspension in 1ml 1X PBS. The OD_{600} of each culture was normalized to 1. The

strains to be competed were mixed at the indicated ratios in a final volume of 100 µL

and then 0.5 µL of the combined suspension was spotted onto a 0.22 µm polycarbonate

filter (Millipore) placed on the surface of a swarm plate in triplicate. Plates were

incubated at 37˚C. Filters were then transferred into a 1.5 ml tube and the filter-

associated cells were resuspended by adding 1 ml 1X PBS + 0.05% Triton X-100

detergent and agitating the tubes at high speed for 2 minutes using a Genie Disruptor

(Zymo). This suspension as well as the inoculum were diluted, spread on LB plates

supplemented with 150 µg/ml 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal)

using glass beads, and incubated at 30°C until blue colonies were visible (~24 h). The

number of blue and white colonies per plate were counted and recorded to determine

the relative abundance of each strain.

Phage susceptibility assays. Phage susceptibility was analyzed by the cross-streak

method or by spotting phage directly onto a lawn of *P. aeruginosa*. For the cross-streak

method, *P. aeruginosa* strains were grown for 16 h in LB before diluting 1:100 in LB and

then growing to an OD_{600} of 0.5-0.7. Then, 4 µl of phage DMS3 vir strain was spotted on

the side of an LB agar plate and then dragged across the agar surface in a straight line

before allowing to absorb into the agar. Once absorbed, 4 µl of each *P. aeruginosa*

strain was spotted at the top of the agar and dragged downward in a straight line

through the phage line. Plates were incubated at 37°C for 16 h.

Alternatively, 1% M8 agar plates (60 x15 mm) containing 0.2% glucose (v/v),

0.5% casamino acids (v/v), and 1 mM $MgSO_4$ were prepared and cooled to room

temperature. 50 µl of *P. aeruginosa* 16 h cultures were added to 1 ml of 0.5% warm top

agar (M8 medium and supplements). The mixture was gently mixed and quickly poured

onto M8 agar plates. Plates were swirled to ensure even spreading of top agar. Once

cooled, 2 µl of phage DMS3 vir strain were spotted onto the center of the plate and

allowed to dry before incubating plates at 37°C for 16 h.

Congo red assay. Cultures were grown in 5 ml LB for 16 h at 37°C with rolling. Culture

aliquots were washed once and resuspended in sterile deionized water. Washed cells

were spotted (3 µl) onto Congo Red plates (1% tryptone [w/v], 1.5% agar, 40 µg/ml

Congo Red, 20 µg/ml Coomassie Blue). Plates were grown for 16 h at 37°C and then

moved to room temperature for 3 days to allow for color development.

Statistical analysis. One-way ANOVA with multiple comparisons was performed

pairwise between all isolates and mixtures using the GraphPad Prism 6 software

(GraphPad, La Jolla, CA).

Acknowledgements. This work was supported by CFF HOGAN19G0 to DAH,

R37AI83256 and R01AI43730 to GAO, and CFF VERMIL21F0 to DMV. This work was

also supported by P30DK117469 for the Applied Bioinformatics and Biostatistics Core.

Sequencing services and specialized equipment was provided by the Genomics and

Molecular Biology Shared Resource Core at Dartmouth supported by NCI Cancer

Center Support Grant 5P30CA023108-37. Equipment used was supported by the NIH

IDeA award to Dartmouth BioMT P20GM113132. We thank Amy Baker for constructing

the Δ*fliC* mutant and Christine Toutain for construction of the *motABCD* deletion strain.

Table 1. Strains and plasmids used in this study

Strain or plasmid	Lab Strain	Description	Source
P. aeruginosa **strains**			
PA14	DH123	Wild type (WT)	(44)
PA14 *att*::*lacZ*	DH22	WT with constitutive expression of *lacZ*	(45)
Δ*flgK*	DH1075	*flgK*::Tn5B30(Tc), non-motile	(4)
Δ*fliC*	DH3543	Unmarked deletion of *fliC*	This study
Δ*motABCD*	SMC7230	Unmarked deletion of *motABCD*	This study
Δ*pilMNOP*	DH2705	Unmarked deletion of *pilMNOP*	(6)
Δ*pilA*	DH2636	Unmarked deletion of *pilA*	(24)
Δ*pelA*	DH97 or SMC2893	Unmarked deletion of *pelA*	(46)
Δ*flgK*Δ*pelA*	DH3541	Unmarked deletion of *flgK* and *pelA*	(11)
Δ*flgK*Δ*pilA*	DH3539	Unmarked deletion of *flgK* and *pilA*	(11)
Δ*flgK*Δ*pilA*Δ*pelA*	DH3542	Unmarked deletion of *flgK*, *pilA*, and *pelA*	(11)
Δ*pilT*	DH3591 or SMC7302	Unmarked deletion of *pilT*	(47)

$\Delta pilU$	DH3592 or SMC7304	Unmarked deletion of $pilU$	(47)
$\Delta pilJ$	DH2637 or SMC2992	Unmarked deletion of $pilJ$	(48)
$\Delta fimS$	SMC6967	Unmarked deletion of $fimS$	(6)
$\Delta cyaAB$	DH2648 or SMC6707	Unmarked deletion of $cyaAB$	(6)
Δvfr	DH2701 or SMC6722	Unmarked deletion of vfr	(6)
$\Delta pilU$ att::P$_{BAD}$-ara	SMC7826	Unmarked $\Delta pilU$ (DH3592/SMC7304) with arabinose-inducible promoter at the att::Tn7 site, control, Gm^R	This study
$\Delta pilU$ att::P$_{BAD}$-$pilU$	SMC7827	Unmarked $\Delta pilU$ (DH3592/SMC7304) complemented with arabinose-inducible $pilU$ at the att::Tn7 site, Gm^R	This study
$\Delta pilT$ att::P$_{BAD}$-ara	SMC8882	Unmarked $\Delta pilT$ (DH3591/SMC7302) with arabinose-inducible promoter at the att::Tn7 site, control, Gm^R	This study

Δ*pilT att*::P*BAD*-*pilT*	SMC8883	Unmarked Δ*pilT* (DH3591/SMC7302) complemented with arabinose-inducible *pilT* at the *att*::Tn7 site, Gm^R	This study
Δ*flgK*Δ*pilU*	DH3947	Unmarked deletion of *flgK* and *pilU*	This study
Δ*flgK*Δ*pilT*	DH3948	Unmarked deletion of *flgK* and *pilT*	This study
Δ*flgK*Δ*pilU att*::P*BAD*-ara	DH3949	Unmarked deletion of *flgK* and *pilU* with arabinose-inducible promoter at the *att*::Tn7 site, control, Gm^R	This study
Δ*flgK*Δ*pilU att*::P*BAD*-*pilU*	DH3950	Unmarked deletion of *flgK* and *pilU* complemented with arabinose-inducible *pilU* at the *att*::Tn7 site, control, Gm^R	This study
Δ*flgK*Δ*pilT att*::P*BAD*-ara	DH3951	Unmarked deletion of *flgK* and *pilT* with arabinose-inducible promoter at the *att*::Tn7 site, control, Gm^R	This study
Δ*flgK*Δ*pilT att*::P*BAD*-*pilT*	DH3952	Unmarked deletion of *flgK* and *pilT* complemented with arabinose-inducible *pilT* at the *att*::Tn7 site, control, Gm^R	This study

E. coli strains

S17 λpir	DH71	Used as a conjugation partner for introducing pMQ30- and GH121-based plasmids	
Plasmids			
pUX-BF13	SMC1852	Helper plasmid that provides the transposition functions for Tn7 in trans, Ap^R	(49)
pMQ56	SMC3242	pMQ56-miniTn7 for insertion at the *att*::Tn7 site, Ap^R	(42)
pBAD-ara	SMC7453	pMQ56-miniTn7 plasmid containing arabinose-inducible P$_{BAD}$ and *araC* for insertion at the *att*::Tn7 site, Ap^R, Gm^R	This study
pPilU-His	SMC7777	pMQ56-miniTn7 plasmid containing 6X His-tagged *pilU* regulated by arabinose-inducible P$_{BAD}$ and *araC* for insertion at the *att*::Tn7 site, Ap^R, Gm^R	This study
pPilT-His	SMC8873	pMQ56-miniTn7 plasmid containing 6X His-tagged *pilT* regulated by arabinose-inducible P$_{BAD}$ and *araC*	This study

		for insertion at the *att*::Tn7 site, Ap^R, Gm^R	
pMQ30	DH2620	Suicide vector for allelic replacement; Gm^R	(42)
pMQ30-del-pilT	SMC7298	Construct for in-frame deletion of *pilT*	(47)
pMQ30-del-pilU	SMC7299	Construct for in-frame deletion of *pilU*	(47)

References

1.   Hogan DA, Kolter R. 2002. *Pseudomonas-Candida* interactions: An ecological

role for virulence factors. Science 296:2229-2232.

2.   Limoli DH, Warren EA, Yarrington KD, Donegan NP, Cheung AL, O'Toole GA.

2019. Interspecies interactions induce exploratory motility in *Pseudomonas*

aeruginosa. Elife 8:e47365.

3.   Lewis KA, Baker AE, Chen AI, Harty CE, Kuchma SL, O'Toole GA, Hogan DA.

2019. Ethanol decreases *Pseudomonas aeruginosa* flagellar motility through the

regulation of flagellar stators. J Bacteriol 201:e00285-17.

4.   O'Toole GA, Kolter R. 1998. Flagellar and twitching motility are necessary for

Pseudomonas aeruginosa biofilm development. Mol Microbiol 30:295-304.

5.   Zhao K, Tseng BS, Beckerman B, Jin F, Gibiansky ML, Harrison JJ, Luijten E,

Parsek MR, Wong GCL. 2013. Psl trails guide exploration and microcolony

formation in *Pseudomonas aeruginosa* biofilms. Nature 497:388-391.

6.   Luo Y, Zhao K, Baker AE, Kuchma SL, Coggan KA, Wolfgang MC, Wong GC,

O'Toole GA. 2015. A hierarchical cascade of second messengers regulates

Pseudomonas aeruginosa surface behaviors. MBio 6:e02456-14.

7.   O'Toole GA, Wong GC. 2016. Sensational biofilms: Surface sensing in bacteria.

Curr Opin Microbiol 30:139-146.

8.   Lee CK, Vachier J, de Anda J, Zhao K, Baker AE, Bennett RR, Armbruster CR,

Lewis KA, Tarnopol RL, Lomba CJ, Hogan DA, Parsek MR, O'Toole GA,

Golestanian R, Wong GCL. 2020. Social cooperativity of bacteria during

reversible surface attachment in young biofilms: A quantitative comparison of

Pseudomonas aeruginosa PA14 and PAO1. mBio 11:e02644-19.

9.   Boyd CD, Smith TJ, El-Kirat-Chatel S, Newell PD, Dufrene YF, O'Toole GA.

2014. Structural features of the *Pseudomonas fluorescens* biofilm adhesin LapA

required for LapG-dependent cleavage, biofilm formation, and cell surface

localization. J Bacteriol 196:2775-88.

10.   Parsek MR. 2016. Controlling the connections of cells to the biofilm matrix. J

Bacteriol 198:12-4.

11.   Ribbe J, Baker AE, Euler S, O'Toole GA, Maier B. 2017. Role of cyclic di-GMP

and exopolysaccharide in type IV pilus dynamics. J Bacteriol 199:e00859-16.

12.   Ma L, Conover M, Lu H, Parsek MR, Bayles K, Wozniak DJ. 2009. Assembly and

development of the *Pseudomonas aeruginosa* biofilm matrix. PLoS Pathog

5:e1000354.

13.   Colvin KM, Irie Y, Tart CS, Urbano R, Whitney JC, Ryder C, Howell PL, Wozniak

DJ, Parsek MR. 2012. The Pel and Psl polysaccharides provide *Pseudomonas*

aeruginosa structural redundancy within the biofilm matrix. Environ Microbiol

14:1913-28.

14.   Wei Q, Ma LZ. 2013. Biofilm matrix and its regulation in *Pseudomonas*

aeruginosa. Int J Mol Sci 14:20983-1005.

15.   Anyan ME, Amiri A, Harvey CW, Tierra G, Morales-Soto N, Driscoll CM, Alber

MS, Shrout JD. 2014. Type IV pili interactions promote intercellular association

and moderate swarming of *Pseudomonas aeruginosa*. Proc Natl Acad Sci U S A

111:18013-8.

16.   Boyle KE, Monaco HT, Deforet M, Yan J, Wang Z, Rhee K, Xavier JB. 2017.

444   Metabolism and the evolution of social behavior. Mol Biol Evol 34:2367-2379.

445   17.   Lozano C, Azcona-Gutierrez JM, Van Bambeke F, Saenz Y. 2018. Great

446   phenotypic and genetic variation among successive chronic *Pseudomonas*

447   *aeruginosa* from a cystic fibrosis patient. PLoS One 13:e0204167.

448   18.   Mahenthiralingam E, Campbell ME, Speert DP. 1994. Nonmotility and phagocytic

449   resistance of *Pseudomonas aeruginosa* isolates from chronically colonized

450   patients with cystic fibrosis. Infect Immun 62:596-605.

451   19.   Workentine ML, Sibley CD, Glezerson B, Purighalla S, Norgaard-Gron JC,

452   Parkins MD, Rabin HR, Surette MG. 2013. Phenotypic heterogeneity of

453   *Pseudomonas aeruginosa* populations in a cystic fibrosis patient. PLoS One

454   8:e60225.

455   20.   Turner KH, Everett J, Trivedi U, Rumbaugh KP, Whiteley M. 2014. Requirements

456   for *Pseudomonas aeruginosa* acute burn and chronic surgical wound infection.

457   PLoS Genet 10:e1004518.

458   21.   Ha DG, Kuchma SL, O'Toole GA. 2014. Plate-based assay for swarming motility

459   in *Pseudomonas aeruginosa*. Methods Mol Biol 1149:67-72.

460   22.   Tremblay J, Deziel E. 2008. Improving the reproducibility of *Pseudomonas*

461   *aeruginosa* swarming motility assays. J Basic Microbiol 48:509-15.

462   23.   Shrout JD, Chopp DL, Just CL, Hentzer M, Givskov M, Parsek MR. 2006. The

463   impact of quorum sensing and swarming motility on *Pseudomonas aeruginosa*

464   biofilm formation is nutritionally conditional. Mol Microbiol 62:1264-77.

24.   Kuchma SL, Ballok AE, Merritt JH, Hammond JH, Lu W, Rabinowitz JD, O'Toole

GA. 2010. Cyclic-di-GMP-mediated repression of swarming motility by

Pseudomonas aeruginosa: The *pilY1* gene and its impact on surface-associated

behaviors. J Bacteriol 192:2950-64.

25.   Kuchma SL, Griffin EF, O'Toole GA. 2012. Minor pilins of the type IV pilus

system participate in the negative regulation of swarming motility. J Bacteriol

194:5388-403.

26.   Whitchurch CB, Mattick JS. 1994. Characterization of a gene, *pilU*, required for

twitching motility but not phage sensitivity in *Pseudomonas aeruginosa*. Mol

Microbiol 13:1079-91.

27.   Tammam S, Sampaleanu LM, Koo J, Sundaram P, Ayers M, Chong PA, Forman-

Kay JD, Burrows LL, Howell PL. 2011. Characterization of the PilN, PilO and PilP

type IVa pilus subcomplex. Mol Microbiol 82:1496-514.

28.   Webster SS, Lee CK, Schmidt WC, Wong GCL, O'Toole GA. 2021. Interaction

between the type 4 pili machinery and a diguanylate cyclase fine-tune c-di-GMP

levels during early biofilm formation. Proc Natl Acad Sci U S A

118:e2105566118.

29.   Burrows LL. 2005. Weapons of mass retraction. Mol Microbiol 57:878-88.

30.   Chiang P, Burrows LL. 2003. Biofilm formation by hyperpiliated mutants of

Pseudomonas aeruginosa. J Bacteriol 185:2374-8.

31.   Whitchurch CB, Mattick JS. 1994. Characterization of a gene, pilU, required for

twitching motility but not phage sensitivity in Pseudomonas aeruginosa. Mol

Microbiol 13:1079-1091.

32. Shen Y, Siryaporn A, Lecuyer S, Gitai Z, Stone HA. 2012. Flow directs surface-
attached bacteria to twitch upstream. Biophys J 103:146-51.

33. Landry RM, An D, Hupp JT, Singh PK, Parsek MR. 2006. Mucin-*Pseudomonas*
aeruginosa interactions promote biofilm formation and antibiotic resistance. Mol
Microbiol 59:142-51.

34. Marmont LS, Whitfield GB, Rich JD, Yip P, Giesbrecht LB, Stremick CA, Whitney
JC, Parsek MR, Harrison JJ, Howell PL. 2017. PelA and PelB proteins form a
modification and secretion complex essential for Pel polysaccharide-dependent
biofilm formation in *Pseudomonas aeruginosa*. J Biol Chem 292:19411-19422.

35. Harrison JJ, Almblad H, Irie Y, Wolter DJ, Eggleston HC, Randall TE, Kitzman
JO, Stackhouse B, Emerson JC, McNamara S, Larsen TJ, Shendure J, Hoffman
LR, Wozniak DJ, Parsek MR. 2020. Elevated exopolysaccharide levels in
Pseudomonas aeruginosa flagellar mutants have implications for biofilm growth
and chronic infections. PLoS Genet 16:e1008848.

36. Chen AI, Dolben EF, Okegbe C, Harty CE, Golub Y, Thao S, Ha DG, Willger SD,
O'Toole GA, Harwood CS, Dietrich LE, Hogan DA. 2014. *Candida albicans*
ethanol stimulates *Pseudomonas aeruginosa* WspR-controlled biofilm formation
as part of a cyclic relationship involving phenazines. PLoS Pathog 10:e1004480.

37. Merritt JH, Brothers KM, Kuchma SL, O'Toole GA. 2007. SadC reciprocally
influences biofilm formation and swarming motility via modulation of
exopolysaccharide production and flagellar function. J Bacteriol 189:8154-64.

38. Whitfield GB, Marmont LS, Bundalovic-Torma C, Razvi E, Roach EJ, Khursigara
CM, Parkinson J, Howell PL. 2020. Discovery and characterization of a Gram-

positive Pel polysaccharide biosynthetic gene cluster. PLoS Pathog

16:e1008281.

39. Darch SE, McNally A, Harrison F, Corander J, Barr HL, Paszkiewicz K, Holden S,

Fogarty A, Crusz SA, Diggle SP. 2015. Recombination is a key driver of genomic

and phenotypic diversity in a *Pseudomonas aeruginosa* population during cystic

fibrosis infection. Sci Rep 5:7649.

40. Head NE, Yu H. 2004. Cross-sectional analysis of clinical and environmental

isolates of *Pseudomonas aeruginosa*: Biofilm formation, virulence, and genome

diversity. Infect Immun 72:133-44.

41. Grosso-Becerra MV, Santos-Medellin C, Gonzalez-Valdez A, Mendez JL,

Delgado G, Morales-Espinosa R, Servin-Gonzalez L, Alcaraz LD, Soberon-

Chavez G. 2014. *Pseudomonas aeruginosa* clinical and environmental isolates

constitute a single population with high phenotypic diversity. BMC Genomics

15:318.

42. Shanks RM, Caiazza NC, Hinsa SM, Toutain CM, O'Toole GA. 2006.

Saccharomyces cerevisiae-based molecular tool kit for manipulation of genes

from gram-negative bacteria. Appl Environ Microbiol 72:5027-36.

43. Ha DG, Richman ME, O'Toole GA. 2014. Deletion mutant library for investigation

of functional outputs of cyclic diguanylate metabolism in *Pseudomonas*

aeruginosa PA14. Appl Environ Microbiol 80:3384-93.

44. Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM. 1995.

Common virulence factors for bacterial pathogenicity in plants and animals.

Science 268:1899-902.

45.   Clay ME, Hammond JH, Zhong F, Chen X, Kowalski CH, Lee AJ, Porter MS,

535   Hampton TH, Greene CS, Pletneva EV, Hogan DA. 2020. *Pseudomonas*

536   *aeruginosa* lasR mutant fitness in microoxia is supported by an Anr-regulated

537   oxygen-binding hemerythrin. Proc Natl Acad Sci U S A 117:3167-3173.

538   46.   Friedman L, Kolter R. 2004. Genes involved in matrix formation in *Pseudomonas*

539   *aeruginosa* PA14 biofilms. Mol Microbiol 51:675-90.

540   47.   Lee CK, de Anda J, Baker AE, Bennett RR, Luo Y, Lee EY, Keefe JA, Helali JS,

541   Ma J, Zhao K, Golestanian R, O'Toole GA, Wong GCL. 2018. Multigenerational

542   memory and adaptive adhesion in early bacterial biofilm communities. Proc Natl

543   Acad Sci U S A 115:4471-4476.

544   48.   Belete B, Lu H, Wozniak DJ. 2008. *Pseudomonas aeruginosa* AlgR regulates

545   type IV pilus biosynthesis by activating transcription of the fimU-pilVWXY1Y2E

546   operon. J Bacteriol 190:2023-30.

547   49.   Hojberg O, Schnider U, Winteler HV, Sorensen J, Haas D. 1999. Oxygen-

548   sensing reporter strain of *Pseudomonas fluorescens* for monitoring the

549   distribution of low-oxygen habitats in soil. Appl Environ Microbiol 65:4085-93.

Figure Legends

Fig. 1. Motility heterogeneity represses swarming motility independent of competition. A. Representative images from swarm assay of wild-type (WT) *P. aeruginosa* strain PA14 (flagellated) mixed with Δ*flgK* (non-flagellated) at the indicated ratios on M8 agar. In this and all figure panels in the manuscript, WT and derivatives are labeled in black, while non-motile mutants are labeled in red. Data are representative of three individual experiments. **B.** Competition growth assay between WT tagged with a *lacZ* reporter (gray) and either an untagged WT strain (black) or Δ*flgK* mutant (red). Data show the average of three individual experiments. Error bars represent the standard deviation between the replicate values. Statistical analysis using one-way ANOVA showed no difference between input and output for each ratio analyzed. **C.** Representative images from swarm assay of WT (flagellated) mixed with Δ*flgK* (non-flagellated), Δ*fliC* (non-flagellated), or Δ*motABCD* (paralyzed flagellum) in the indicated ratios on M8 agar. Data are representative of three individual experiments. Scale bar: 30 mm

Fig. 2. Flagellated *P. aeruginosa* requires functional T4P to be repressed by the non-motile subpopulation. Representative images from swarm assays of the non-flagellated Δ*flgK* mutant (functional T4P) mixed with either wild-type (WT) *P. aeruginosa* strain PA14 (functional T4P), the Δ*pilA* mutant (lacking T4P), the Δ*pilT* mutant (no T4P retraction), or the Δ*pilU* mutant (reduced T4P retraction) at the indicated ratios on M8 agar. Data are representative of three individual experiments. Scale bar: 30 mm

Fig. 3. Non-flagellated *P. aeruginosa* requires T4P to repress flagellum-mediated motility of motile strains on soft agar. A. Representative images of swarm assays for wild-type (WT) *P. aeruginosa* strain PA14 (swarmer, Pel+, T4P+) mixed with Δ*flgK* (non-swarmer, Pel+, T4P+), Δ*flgK*Δ*pilA* (non-swarmer, Pel+, T4P-), Δ*flgK*Δ*pilU* (non-swarmer, Pel+, reduced T4P retraction), or Δ*flgK*Δ*pilT* (non-swarmer, Pel+, no T4P retraction) on M8 agar at the indicated ratios. WT is labeled in black while Δ*flgK* and derivatives are labeled in red. **B.** Representative image showing the interaction between WT (black dot), Δ*flgK* (red dots), and Δ*flgK*Δ*pilA* (purple dots) for flagellum-mediated swimming motility in soft agar (0.3% M63 agar) after 42 h incubation. The colored dots indicate the points of inoculation for the respective strains. Data are representative of three individual experiments. Scale bar: 30 mm

Fig. 4. Pel matrix is only required by the non-flagellated *P. aeruginosa* subpopulation in order to repress overall swarming motility. A. Representative images from swarm assays of Δ*flgK* (non-flagellated and Pel+) mixed with wild-type (WT) *P. aeruginosa* strain PA14 (flagellated and Pel+) or Δ*pelA* (flagellated and Pel-) and of WT mixed with Δ*flgK*Δ*pelA* (non-flagellated and Pel-) at the indicated ratios on M8 agar. Scale bar: 30 mm **B.** Representative images of WT, Δ*pilA*, Δ*pelA*, Δ*flgK*, Δ*flgK*Δ*pilA*, Δ*flgK*Δ*pelA,* and Δ*flgK*Δ*pilA*Δ*pelA* grown on Congo Red plates to assess Pel production. WT and derivatives are labeled in black while Δ*flgK* and derivatives are labeled in red. Scale bar: 10 mm

Fig. 5. Summary model describing the genes and their products needed for repression of population-wide flagellum-mediated motility by non-flagellated cells in *P. aeruginosa* strain PA14. We show that a subpopulation of *P. aeruginosa* cells defective in flagellar motility (red) in co-culture with flagellated cells (black) impedes flagellum-mediated population-wide swarming motility in a manner that is dependent on retractable T4P (PilMNOP, PilA, PilU, PilT). We also show that Pel matrix (PelA), but only from the non-flagellated strain, is required. PilU, PilJ, FimS, CyaAB, and Vfr were not required for repression of swarming motility. These data may indicate ways that *P. aeruginosa* cells come together during biofilm formation.

A

100:0   95:5   90:10   85:15   80:20   0:100

WT:*ΔflgK*

B

WT-*lacZ*:WT

WT-*lacZ*:*ΔflgK*

Percentage (%)

Initial ratio: 75:25   50:50   25:75

75:25   50:50   25:75

C

100:0   75:25   50:50   25:75   0:100

WT:*ΔflgK*

WT:*ΔfliC*

WT:*ΔmotABCD*