Molecular switch between mammalian MLL complexes dictates response to Menin-MLL inhibition

Yadira M. Soto-Feliciano[1,11,12,†], Francisco J. Sánchez-Rivera[2,11,12,,†], Florian Perner[3,4,5,†], Douglas W. Barrows[1,6], Edward R. Kastenhuber[2], Yu-Jui Ho[2], Thomas Carroll[6], Yijun Xiong[3,4], Disha Anand[3,5], Alexey Soshnev[1], Leah Gates[1], Mary Clare Beytagh[1], David Cheon[1], Shengqing Gu[7,8], X. Shirley Liu[7,8], Andrei V. Krivtsov[3,4], Maximiliano Meneses[9], Elisa de Stanchina[9], Richard M. Stone[10], Scott A. Armstrong[3,4,*], Scott W. Lowe[2,*], C. David Allis[1,*]

[1]Laboratory of Chromatin Biology & Epigenetics, The Rockefeller University, New York, NY 10065, USA.

[2]Cancer Biology and Genetics, MSKCC New York, NY 10065, USA.

[3]Department of Pediatric Oncology, Dana-Farber Cancer Institute, Boston, MA., 02215, USA.

[4]Division of Hematology/Oncology, Boston Children's Hospital and Harvard Medical School, Boston, MA, 02215, USA.

[5]Internal Medicine C, Greifswald University Medical Center, Greifswald, 17475, Germany.

[6]Bioinformatics Resource Center, The Rockefeller University, New York, NY 10065, USA.

[7]Department of Biostatistics and Computational Biology, Dana-Farber Cancer Institute, Harvard School of Public Health, Boston, 02215, MA, USA.

[8]Center for Functional Cancer Epigenetics, Dana-Farber Cancer Institute, Boston, 02215, MA, USA.

[9]Department of Molecular Pharmacology, MSKCC New York, NY 10065, USA.

[10]Leukemia Division, Department of Medical Oncology, Dana-Farber Cancer Institute, Boston,

MA, 02215, USA.

[11]**Current affiliation/address:** David H. Koch Institute for Integrative Cancer Research,

Massachusetts Institute of Technology, Cambridge, 02142, Massachusetts, USA

[12]**Current affiliation/address:** Department of Biology, Massachusetts Institute of Technology,

Cambridge, 02142, Massachusetts, USA

† Co-first authors

* Co-corresponding authors

Correspondence:

C. David Allis: The Rockefeller University, Allis Lab, Box #78, 1230 York Avenue, New York, NY

10065, USA. Email: alliscd@rockefeller.edu

Scott W. Lowe: Memorial Sloan Kettering Cancer Center (MSKCC), Sloan Kettering Institute,

Cancer Biology & Genetics Program, New York, NY, 10065, USA. Email: lowes@mskcc.org

Scott A. Armstrong: Harvard Medical School, Dana-Farber Cancer Institute, Department of

Pediatric Oncology, Boston, MA, 02115, USA. Email: Scott_Armstrong@dfci.harvard.edu

Keywords: Menin, MLL, UTX, chromatin, epigenetics, leukemia, CRISPR

ABSTRACT

The chromatin adaptor Menin interacts with oncogenic fusion proteins encoded by *MLL1*-rearrangements (*MLL1*-r), and small molecules that disrupt these associations are currently in clinical trials for the treatment of leukemia. Here, by integrating chromatin-focused and genome-wide CRISPR screens with genetic, pharmacological, and biochemical approaches in mouse and human systems, we discovered a molecular switch between the MLL1-Menin and MLL3/4-UTX chromatin modifying complexes that dictates response to Menin-MLL inhibitors. We show that MLL1-Menin safeguards leukemia survival by impeding binding of the MLL3/4-UTX complex at a subset of target gene promoters. Disrupting the interaction between Menin and MLL1 leads to UTX-dependent transcriptional activation of tumor suppressive genes, including factors that mediate cell cycle arrest and cellular senescence. We show that this epigenetic mechanism is operative in murine and human models of AML, and clinical responses to Menin-MLL inhibition in primary human leukemia are also accompanied by induction of these tumor suppressive genes. We provide proof-of-concept *in vivo* evidence that this mechanism can be rationally exploited to design combination therapies using FDA-approved CDK4/6 inhibitors to mitigate treatment resistance associated with MLL3/4-UTX complex mutations. These findings shed light on the context-dependent and often antagonistic roles that chromatin regulators exhibit in development and disease and provide mechanistic insight for rational design of targeted epigenetic therapies.

INTRODUCTION

Menin is an evolutionarily conserved nuclear factor that associates with chromatin to recruit (adapt) interacting proteins[1], including the Trithorax (*Trx*)-related MLL1 (KMT2A) and MLL2 (KMT2B) histone methyltransferase complexes[2,3] and the MLL1 oncogenic fusion proteins[4], transcription factors (e.g., c-MYC[5], JUND[6,7], SMADs[8,9]), and other chromatin-bound proteins (e.g.,

LEDGF[10] (reviewed in ref. 11)). However, the detailed underlying molecular mechanisms and

functional interactions associated with Menin remain elusive.

Menin is a core subunit of the MLL1 (ref. 12) and MLL2 complexes[2] and is responsible for targeting

these to chromatin[3]. Menin is required for MLL1/MLL2-dependent H3K4 trimethylation of *HOX*

genes and their stable long-term expression during development[2,13]. Menin has context-specific

functions in human diseases, acting as a tumor suppressor in neuroendocrine malignancies[14,15]

and in certain skin[16], lung[17], and CNS tumors[18], and as an oncogenic cofactor in other cancers,

including hepatocellular carcinoma[19] and *MLL1*-rearranged (*MLL1*-r) leukemias[4,20]. Furthermore,

over 1000 germline and somatic *MEN1* variants have been identified, some of which are linked

to cancer predisposition[21]. Owing to its pro-oncogenic roles in mixed lineage leukemia and other

malignancies, small molecule inhibitors targeting the Menin-MLL1 and Menin-MLL2 protein-

protein interactions have shown great promise for intercepting and treating different types of

cancers[19,22-27]. Notably, three structurally different Menin-MLL inhibitors have recently entered

clinical trials (NCT04065399, NCT04067336, NCT04811560) and at least one has been granted

fast track designation by the FDA for treatment of relapsed/refractory acute leukemias[25,27,28].

Thus, an understanding of the molecular mechanisms of action of these drugs would facilitate the

development of biomarkers to predict therapeutic response and resistance, and lead to rational

design of more effective combination treatments.

RESULTS

Functional interplay between MLL1-Menin and MLL3/4-UTX chromatin modifying

complexes

We performed a series of CRISPR-based genetic screens to molecularly dissect the dependency

of *MLL1*-r leukemias to Menin and identify factors that dictate response and resistance to Menin-

MLL inhibitors. First, we screened a chromatin-focused CRISPR library in Cas9-expressing

Menin-dependent mouse leukemia cells driven by a human MLL-AF9 transgene (hereafter

referred to as MLL-AF9 cells)[29] (**Fig. 1A, Extended Data Fig. 1A-D, Supplementary Table 1**).

Library-transduced cells were cultured in media with DMSO (vehicle) or the Menin-MLL inhibitor

MI-503 (ref. 22) for 12 cell population doublings (CPDs), followed by screen deconvolution using

next-generation sequencing (**Fig. 1A**, **Extended Data Fig. 1E-F**). A gene score (mean $\log_2$ fold-

change) was calculated for each gene included in the library by assessing the change in

abundance of sgRNAs during the culture period (**Fig. 1B**). As expected, sgRNAs targeting known

MLL1-r leukemia dependencies, such as *Dot1l*[29-32]*, Brd4* (ref. 33), and *Myb*[34] were strongly

depleted in both treatments while control sgRNAs remained largely neutral, thereby validating our

screening platform (**Supplementary Table 2**).

Among the sgRNAs that were enriched or depleted in the absence and presence of drug, sgRNAs

targeting the histone H3 lysine 27 (H3K27) demethylase *Utx* (*Kdm6a*) and the histone H3 lysine

4 (H3K4) mono-methyltransferases *Mll3* (*Kmt2c*) and *Mll4* (*Kmt2d*) were the most significantly

enriched in the Menin-MLL inhibitor context (**Fig. 1B**, red circles). This result was unexpected,

since the canonical activity of the MLL1/2-Menin complex (disrupted by MI-503) is to catalyze the

di/tri-methylation of H3K4 (H3K4me2/3) at *promoters*, resulting in loading of the transcriptional

machinery at such sites[13,35,36], while the MLL3/4-UTX complex regulates *enhancer* states by

serving as the major H3K4 mono-methyltransferase (H3K4me1) (refs. 37-41).

To assess whether these results were idiosyncratic of the cell line, library, or inhibitor used, we

performed a genome-wide CRISPR screen in an independently derived MLL-AF9 mouse cell line

using VTP-50469, a more potent, selective, and orally bioavailable Menin-MLL inhibitor that led

to the development of SNDX-5613, which is currently under investigation in a phase 1/2 clinical

trial for treatment of relapsed/refractory acute leukemia (NCT04065399) (refs. 25,27,28).

Remarkably, sgRNAs targeting *Utx* (*Kdm6a*), *Mll3* (*Kmt2c*), and *Mll4* (*Kmt2d*) were also among

the most significantly enriched candidate genes identified in this genome-wide screen (**Fig. 1C,**

Extended Data Fig. 2A-C, Supplementary Table 2). Interestingly, shared subunits between the

two types of MLL complexes scored similarly in both vehicle and Menin-MLL inhibitor conditions

(**Fig. 1B-D, Extended Data Fig. 1G, Extended Data Fig. 2D**). This suggests that core subunits

of the MLL3/4-UTX complex[42] modulate therapeutic responses to Menin-MLL inhibition and have

context-specific tumor suppressive functions[43,44], a result that is consistent with their

haploinsufficient tumor suppressor activity in some cancers[45].

Utx or *Mll3* loss-of-function did not affect leukemia cell proliferation *in vitro*, while *Mll4* inactivation

decreased cell growth, consistent with previous work[43] (**Extended Data Fig. 3A-D**). However, in

agreement with our screening results, *Utx* disruption using three distinct sgRNAs significantly

increased the viability of MI-503-treated leukemia cells, confirming that inactivation of *Utx* confers

resistance to Menin-MLL inhibition (**Fig. 1E, Extended Data Fig. 3E**). In addition, *Mll3-* and *Mll4-*

deficient cells treated with MI-503 exhibited a proliferative advantage over wild-type cells only in

the treatment context (**Extended Data Fig. 3F-G**). These orthogonal results establish the MLL3/4-

UTX complex as a central modulator of therapy response to Menin-MLL inhibition.

UTX was among the most significantly enriched chromatin factors in our screen (**Fig. 1B-C,**

Extended Data Fig. 1G, Extended Data Fig. 2D). As this factor is shared by both MLL3 and

MLL4 complexes[41,46,47], we focused on UTX disruption to functionally probe the molecular

mechanisms linked to resistance to Menin-MLL inhibition. We first tested whether genetic *Utx*

inactivation could rescue the effects of Menin-specific ablation in MLL-AF9 leukemia cells. While

CRISPR-mediated deletion of *Men1* led to robust inhibition of proliferation (**Extended Data Fig.**

4A), co-deletion of *Utx* suppressed this phenotype, such that *Men1*-deficient MLL-AF9 cells were

able to robustly proliferate (**Fig. 1F, Extended Data Fig. 4B-C**). Importantly, deletion of *Utx*, *Mll3*,

or *Mll4* in isolation did not confer any significant fitness advantage to cells, further validating our

findings obtained with pharmacologic inhibition of Menin-MLL (**Extended Data Fig. 3**).

To determine if the genetic interaction between Menin and UTX depends on the MLL-fusion

protein (MLL-FP), we ablated the human *MLL1-AF9* transgene expressed in MLL-AF9 leukemia

cells using sgRNAs selectively targeting exons 3 or 5 of the human *MLL1* gene (preceding the

breakpoint cluster region (exons 9-11) in *MLL1*-translocations)[48,49] (**Extended Data Fig. 5A**).

Cells deficient for *MLL-AF9* and *Utx* exhibited fitness defects similar to cells lacking *MLL-AF9*,

validating the exquisite dependency of leukemia cells to the pleiotropic gene regulatory activity of

this oncogenic fusion[20] (**Extended Data Fig. 5B-C**). However, MLL-AF9 leukemia cells deficient

for *Men1* and *Utx*, or those deficient for *Utx* and treated with Menin-MLL inhibitor, were able to

proliferate without reactivation of canonical fusion targets (**Extended Data Fig. 6**), demonstrating

that this genetic interaction is not entirely dependent on the oncogenic activity of the MLL-FP.

Similarly, co-deletion of *UTX* and *MEN1* in a Menin-dependent non *MLL1*-r human leukemia cell

line driven by *DNMT3A* and *NPM1* mutations[50,51] also rescued the fitness deficiency associated

with loss of *MEN1* alone (**Extended Data Fig. 5D-E**), suggesting that the functional Menin-UTX

interaction observed in our models is conserved in human cells. Similar results were obtained in

non *MLL1*-r AML1-ETO9a mouse leukemia cells treated with MI-503 (**Extended Data Fig. 5F-G**).

These data suggest that the phenotype of Menin-MLL inhibition is evolutionarily conserved, acts

in part through a pathway that involves crosstalk between Menin and UTX, and is downstream or

independent of MLL-FPs.

Menin is needed for expression of canonical MLL-FP target genes, such as *Meis1*, whose function

is required for leukemia maintenance[25,35,36,52,53], and *Meis1* over-expression was recently shown

to partially rescue the leukemic stem cell transcriptional program suppressed by Menin-MLL

inhibition[27]. We confirmed that treatment of mouse MLL-AF9 leukemia cells with MI-503 led to

robust downregulation of *Meis1* (**Extended Data Fig. 6A**). However, to our surprise, cells

genetically deficient for both *Men1* and *Utx* showed a similar reduction in *Meis1* levels (**Extended

Data Fig. 6B**), yet were able to proliferate *in vitro* (**Fig. 1F, Extended Data Fig. 4B**). We further

confirmed these findings in an additional Menin-dependent mouse MLL-AF9 cell line (**Extended**

Data Fig. 6C-D) and a human MLL-AF4-driven leukemia line[54] (**Extended Data Fig. 6E-F**). These

experiments reveal an epistatic relationship between Menin and UTX in Menin-dependent

mammalian cells and suggest that factors beyond *Meis1* can sustain the proliferative potential of

MLL1-r and non *MLL1*-r leukemia cells.

MLL1-Menin complex and NF-Y restrict chromatin occupancy of MLL3/4-UTX at promoter

regions

Menin is generally associated with chromatin-modifying and transcriptional complexes at

promoter regions[6,55], while UTX canonically resides in complexes that regulate enhancers[41]. To

determine the genomic distribution of Menin and UTX in the absence or presence of the Menin-

MLL inhibitor, we performed chromatin immunoprecipitation followed by sequencing (ChIP-Seq).

Menin showed strong enrichment at promoter regions (here defined as transcription start sites

(TSSs) ± 2kb), which was substantially decreased when its canonical interactions with MLL1/2

were disrupted by MI-503 (**Fig. 2A**). Genome-wide enrichment of Menin was also decreased, as

evidenced by a >10-fold reduction in the number of ChIP peaks post MI-503 treatment (**Extended**

Data Fig. 7A). Conversely, while only a very minor fraction of UTX appeared to be enriched on

chromatin at basal conditions, the genome-wide enrichment of UTX dramatically increased upon

treatment with MI-503, as evidenced by a ~70-fold increase in the number of peaks (**Extended**

Data Fig. 7A). UTX canonically resides in complexes that regulate enhancers[37,41] yet, contrary to

expectation, more than 70% of UTX binding peaks in MI-503 treated cells were localized at

promoters and proximal regions (**Fig. 2B, Extended Data Fig. 7A**)[56]. Importantly, the enhanced

enrichment of UTX on chromatin was not merely the result of increased expression and/or protein

stability (**Fig. 2C-D**). These data show that disruption of the Menin-MLL1/2 interaction leads to

dynamic and robust recruitment of UTX to promoter regions (**Fig. 2B**), suggesting a previously

unrecognized role for the canonical MLL3/4-UTX complex in promoter regulation and gene

transcription[37].

Not only did Menin depletion lead to recruitment of UTX to promoter regions, the loci that

accumulated UTX significantly overlapped with those where Menin was lost (**Fig. 2E-F**, **Extended**

Data Fig. 7B). Specifically, correlation analysis suggested that reduction of Menin binding by MI-

503 coincided with increased UTX chromatin occupancy at the same genomic loci (*Pearson*

coefficient (r) = -0.49) (**Fig. 2G**). This molecular switch was specific to Menin-dependent cells, as

MI-503 treatment of Menin-independent mouse fibroblasts did not lead to UTX redistribution or

increased binding to promoters (**Extended Data Fig. 8A-E**). Therefore, Menin excludes UTX from

binding to promoters in Menin-dependent cell types, and this antagonism can be reversed by MI-

503-mediated displacement of MLL1-Menin complexes from chromatin.

Inhibition of Menin-MLL1 leads to displacement of this complex from gene promoters, allowing

UTX to bind and potentially regulate target gene expression (**Fig. 2**). To further understand the

molecular switch between Menin and UTX at promoters, we examined the chromatin-binding

profiles of their cognate H3K4 methyltransferases and their enzymatic histone modifications

(MLL1 and H3K4me3 for Menin, and MLL3/4 and H3K4me1 for UTX) (**Extended Data Fig. 9A-**

B). Disruption of the Menin-MLL1 interaction led to a substantial decrease in MLL1 chromatin

binding and a concomitant decrease of its enzymatic product, H3K4me3, (**Extended Data Fig.**

9C-E). In contrast, Menin-MLL1 inhibition led to increased binding of the MLL3/4 enzymes at

promoter regions co-occupied by UTX and a concomitant increase in H3K4me1 signal at these

same loci (**Extended Data Fig. 9C, 9F-G**). These data suggest that disruption of the MLL1-Menin

complex induces targeting of the enzymatic subunits of the MLL3/4-UTX complex to non-

canonical sites that are normally bound by the MLL1-Menin complex and are distinct from those

bound by the MLL-FP (**Extended Data Fig. 10A-C**). These results provide additional evidence

for the existence of a molecular switch between the MLL complexes that is induced by Menin-

MLL inhibition.

The above results are provocative, as promoter-associated H3K4me1 facilitates transcriptional

repression in other cellular settings[57], suggesting that deposition of H3K4me1 at gene promoters

could depend on context[58]. Nevertheless, our finding that Utx^{KO}, $Mll3^{KO}$, and $Mll4^{KO}$ cells are

resistant to Menin-MLL inhibition (**Fig. 1**, **Extended Data Fig. 3E-G**) – combined with the

colocalization of UTX, MLL3, MLL4, and H3K4me1 at the promoters of target genes – led us to

hypothesize that the MLL3/4-UTX complex co-regulates gene expression in the context of Menin-

MLL inhibition in leukemia. To test this hypothesis, we leveraged H3 lysine-4-to-methionine (K4M)

'oncohistone' mutants as orthogonal molecular tools to destabilize the MLL3/4-UTX complex[59]

(**Extended Data Fig. 11A**). Supporting our model, expression of H3.1K4M in MLL-AF9 leukemia

cells led to a substantial proliferative advantage only in the context of Menin-MLL inhibition

(**Extended Data Fig. 11B-D**), demonstrating that destabilization of the MLL3/4-UTX complex

(**Extended Data Fig. 13E**) can phenocopy the intrinsic resistance of Utx-, $Mll3$-, and $Mll4$-deficient

cells to Menin-MLL1 inhibition (**Fig. 1E-F, Extended Data Fig. 3E-G**). These results further

support a model whereby the MLL3/4-UTX complex serves as a central modulator of therapy

response to Menin-MLL inhibition.

Given that Menin lacks a defined DNA binding domain[60], we hypothesized that a sequence-

specific DNA binding factor regulates the switch between MLL1-Menin and MLL3/4-UTX

occupancy at specific promoters. To test this hypothesis, we performed motif analysis on genomic

regions bound by Menin and UTX, uncovering NF-Y sequence motifs[61] as the most significantly

enriched (P=1e^{-130}) (**Extended Data Fig. 12A**). These motifs were specific to leukemia cells, as

motif enrichment analysis identified a different subset of sequences in mouse fibroblasts

(**Extended Data Fig. 12B-C**). ChIP-Seq of NF-YA (the DNA binding and transactivation subunit

of the NF-Y complex[61]) confirmed these observations, showing that NF-YA co-occupies sites

bound by Menin and UTX (**Extended Data Fig. 12D-E**). Consistent with our hypothesis, sites

bound by UTX in the context of Menin-MLL inhibition also had lower NF-YA occupancy (**Extended**

Data Fig. 12D-E), which was not due to decreased protein levels (**Extended Data Fig. 12F**).

Moreover, CRISPR-mediated disruption of *Nfya* in MLL-AF9 leukemia cells significantly

decreased the viability of MI-503-treated leukemia cells relative to control cells (**Extended Data**

Fig. 13A-C). This observation led us to hypothesize that NF-YA depletion could lead to a

concomitant decrease of Menin and increase of UTX occupancy at target genomic regions.

Supporting this model, treatment of *Nfya*[KO] cells with MI-503 led to a substantial increase in UTX

occupancy at genomic regions normally bound by Menin and NF-Y at steady-state (**Extended**

Data Fig. 10D-F). These results are also consistent with co-essentiality analyses in human cells[59]

showing that a subset of *MEN1*-dependent cells are codependent on members of the NF-Y

complex for survival (**Extended Data Fig. 13G-J**). Collectively, these results implicate NF-Y in

restricting the occupancy of UTX at promoter regions and further suggest that Menin binding to

these sites is mediated by the sequence-specific DNA binding activity of the NF-Y transcription

factor.

Transcriptional co-regulation of tumor suppressive pathways by Menin and UTX

The above results support a model whereby a molecular switch between the MLL1-Menin and

MLL3/4-UTX complexes at gene promoters dictates responses to Menin-MLL inhibition. We

hypothesized that this switch acts through transcriptional regulation of genes involved in pathways

that control cellular proliferation and survival. To test this, we performed transcriptional profiling

of MLL-AF9 leukemia cells treated with MI-503 and identified pathways that are reciprocally

regulated by Menin and UTX (**Extended Data Fig. 14A-B**). Although the catalytic activity of the

MLL1/2-Menin histone methyltransferase complex is canonically associated with actively

transcribed developmental genes[2,13], disruption of the complex by MI-503 resulted in both up- and

down-regulation of gene expression, with slight skewing towards gene activation (**Fig. 4A**).

Remarkably, the majority of significantly upregulated genes were reciprocally bound by Menin

and UTX, suggesting that their expression is tightly co-regulated by these chromatin factors and

gene repression mechanisms might be operative in this context (**Fig. 4B**). Consistent with this

observation, ChIP-Seq against acetylated H4K16 and H3K27 (histone marks associated with

gene activation;[62]) showed that levels of these marks increased at sites where the MLL3/4-UTX

complex is recruited to upon Menin-MLL inhibition (**Extended Data Fig. 15A-C**). Importantly, we

also observed increased levels of MOF, a histone H4K16 acetyltransferase[63-65], at the same loci

bound by UTX in the context of Menin-MLL inhibition (**Extended Data Fig. 15B**). These results

support a model whereby Menin-MLL inhibition leads to dynamic changes in gene expression due

to loss of MLL-dependent repressive activity and a concomitant increase in histone marks

associated with gene activation (including H4K16ac and H3K27ac)[66,67].

To determine if UTX was necessary for gene activation, we generated *Utx*-deficient (*Utx^{KO}*) MLL-

AF9 leukemia cells (**Extended Data Fig. 16A**), treated these with MI-503, and profiled by RNA-

Seq (**Extended Data Fig. 16B**). Genes bound by UTX and induced by MI-503 failed to get

activated in *Utx^{KO}* cells, suggesting that UTX function is necessary for their transcriptional

activation when Menin is displaced from chromatin (we refer to these genes as 'Menin-UTX

targets') (**Fig. 4C**, **Supplementary Table 4**). Importantly, *Utx^{KO}* cells treated with MI-503 exhibited

downregulation of canonical MLL-AF9 targets and induction of myeloid differentiation programs

(**Extended Data Fig. 17A-C**), as observed by others in *Utx* proficient cells[22,25,27]. As shown earlier,

Utx^{KO} leukemia cells treated with MI-503 were able to proliferate without re-expression of *Meis1*,

a critical Menin-MLL-FP target[25,27,52,53] (**Fig. 1E**, **Extended Data Fig. 6B, Extended Data Fig.**

17D). These data suggest a new paradigm whereby the effects of Menin-MLL inhibition on MLL-

FP target genes are independent of its effects on Menin-UTX transcriptional targets, and that

induction of a tumor suppressive gene expression program alongside repression of canonical

MLL-FP targets is required for the anti-leukemic activity of Menin-MLL1 inhibitors.

To gain insight into pathways directly regulated by Menin and UTX, we performed gene ontology

(GO) analysis[68,69] of Menin-UTX targets (**Supplementary Table 4**) and found significant evidence

for their association with transcriptional programs related to proliferation, differentiation, and

survival (**Extended Data Fig. 14B**). To further evaluate the relevance of these GO terms, we

performed Gene Set Enrichment Analysis (GSEA)[70] on transcriptional data from Utx^{WT} and Utx^{KO}

cells treated with MI-503 and observed a significant correlation between the presence of UTX and

the expression of senescence-associated genes following MI-503 treatment[71] (**Fig. 4D**, **Extended**

Data Fig. 18A-F, **Supplementary Table 4**). This correlation was independently confirmed using

a curated list of genes involved in cell cycle arrest and therapy-induced senescence[72] (**Fig. 4E**)

and is consistent with the fact that Menin-MLL inhibition induces a combination of cell cycle arrest,

apoptosis, and differentiation (**Extended Data Fig. 1E-G**)[22], as well as our observation that the

senescence-associated H4K16ac mark increases with MI-503 treatment[73] (**Extended Data Fig.**

15A-B). We also observed increased occupancy of RNA Polymerase II at the promoters of

senescence-associated genes that are up-regulated by MI-503 treatment, lending further support

to these findings (**Extended Data Fig. 18G**). These observations suggest that the interplay

between Menin and UTX in the context of Menin-MLL inhibition regulates the expression of genes

involved in cellular senescence.

The cellular senescence program is highly complex and is characterized in part by induction of

permanent cell cycle arrest and a senescence-associated secretory phenotype (SASP)[74].

Consistent with a direct role for the Menin-UTX switch in regulating cellular senescence, secretion

of prototypical SASP cytokines like IL-6, IFNβ-1, IL-3, and IL-15 was induced upon Menin-MLL

inhibition in a *Utx*-dependent manner (**Fig. 4F-G, Extended Data Fig. 19**). These results

demonstrate that Menin-MLL inhibition engages a potent tumor suppressive network that includes

therapy-induced cellular senescence and is transcriptionally regulated in large part by the MLL3/4-

UTX complex.

Combinatorial targeting of Menin and CDK4/6 overcomes resistance associated with loss of MLL3/4-UTX function

Given that senescence can be regulated at both transcriptional and post-transcriptional levels, and chromatin regulation has been functionally implicated in modulating this program[73,75,76], we hypothesized that the Menin-UTX molecular switch directly regulates the expression of cell cycle arrest- and senescence-associated genes by direct binding to their promoters[56,77]. Consistent with our hypothesis, we found that Menin is bound to the promoters of the cyclin-dependent kinase (CDK) inhibitors *Cdkn2c/Ink4c* and *Cdkn2d/Ink4d* at basal conditions[78], and that binding is abrogated upon Menin-MLL inhibition, leading to induction of their expression (**Fig. 4H**, **Extended Data Fig. 20A-B, Extended Data Fig. 21**). Conversely, we found that UTX binds to these promoters only in the context of Menin-MLL inhibition, leading to UTX-dependent upregulation of both CDK inhibitors (**Fig. 4H**, **Extended Data Fig. 20A-B**). These data show that the Menin-UTX molecular switch regulates the expression of these CDK inhibitors by direct chromatin regulation.

Since the proteins encoded by these two genes are natural inhibitors of the CDK4 and CDK6 kinases[78], we hypothesized that pharmacological inhibition of CDK4/6 would bypass the intrinsic resistance of *Utx^KO^* cells to MI-503 while retaining the therapeutic effects of Menin-MLL inhibition on MLL-FP targets (**Fig. 4I**). We therefore combined MI-503 with the FDA-approved CDK4/6 inhibitor Palbociclib[79], which has been shown to trigger durable cell cycle arrest and senescence[72,80,81]. *Utx^KO^* cells were more resistant to either MI-503 or Palbociclib alone when compared to *Utx^WT^* cells, likely due to higher basal levels of *Cdk6* transcripts (**Fig. 4J, Extended Data Fig. 20C-D**) [77,82]. However, combined inhibition of Menin-MLL and CDK4/6 resulted in a synergistic effect (CD=0.4) (ref. 83) on inhibiting cell proliferation to similar levels achieved by MI-503 treatment of *Utx^WT^* cells (**Fig. 4J**). These proof-of-concept data demonstrate that targeting pathways regulated by Menin and UTX can produce combinatorial therapeutic effects and further suggest that the anti-leukemic effects of MI-503[22] are primarily through reactivation of tumor

331 suppressor pathways and not solely through dampening transcription of MLL-FP targets like

332 *Meis1* (refs. 22,25,27,52). Thus, the combination of Palbociclib with Menin-MLL inhibitors may

333 represent a novel and more effective therapeutic strategy for Menin-dependent cancers[23,24,26,84,85].

Catalytic activity of UTX is not required to elicit UTX-dependent cellular and transcriptional

functions

The UTX protein contains several functional domains, including a JmjC demethylase domain that

catalyzes the removal of the H3K27me3 histone mark[86-89]. To determine the regions of UTX that

are necessary for UTX-dependent cellular and transcriptional phenotypes observed upon Menin-

MLL inhibition, we performed a series of structure-function-rescue experiments (**Fig. 4A**). We

generated lentiviral constructs encoding dual N-terminal HA- and C-terminal Flag-tagged

truncations of UTX spanning the full-length of the mouse protein and used them to stably

transduce and phenotype *Utx*-null leukemia cells treated with MI-503 (**Fig. 4A, Extended Data**

Fig. 22A-C). We found that full-length UTX or a UTX truncation harboring the first 500 amino

acids (1-500 aa) and lacking the JmjC demethylase domain[86-89], were sufficient to re-sensitize

cells to MI-503 treatment while truncations proximal to the C-terminus were unable to do so (**Fig.**

4B). Consistent with these cellular phenotypes, RNA-Seq of FACS-sorted *Utx* cDNA expressing

cells treated with MI-503 showed that full-length UTX and the 1-500 aa UTX truncation were

sufficient to rescue UTX-dependent transcriptional phenotypes associated with Menin-MLL

inhibition (**Fig. 4C**), including induction of Menin-UTX target genes like *Cdkn2c* (**Fig. 4D,**

Extended Data Fig. 23). These results demonstrate that an N-terminal truncation of UTX lacking

the JmjC demethylase domain[86-89] and a recently reported intrinsically disordered region[90] is both

necessary and sufficient to drive tumor suppressive responses in this context.

To further confirm that UTX employs non-catalytic mechanisms to regulate gene expression in

the context of Menin-MLL inhibition, we performed ChIP-Seq for H3K27me3, a modification

catalyzed by PRC2 (ref. 91) and removed by UTX[86-89]. We found that the genomic redistribution

of UTX induced upon treatment with MI-503 (**Fig. 2B, 2E**) did not affect global levels of H3K27me3

(**Extended Data Fig. 24A-B**), which led us to hypothesize that this histone mark does not play a

critical role in these phenotypes. To functionally test this hypothesis, we generated *Ezh2*^{KO} and

Ezh2^{KO};*Utx*^{KO} MLL-AF9 leukemia cells and confirmed by immunoblotting that these have no

detectable levels of H3K27me3 (**Extended Data Fig. 24C**). Remarkably, while *Ezh2*-null

leukemia cells remained exquisitely sensitive to MI-503, double *Ezh2*^{KO};*Utx*^{KO} cells were fully

resistant to Menin-MLL inhibition (**Extended Data Fig. 24D**). These results demonstrate that the

catalytic function of UTX is dispensable for UTX-dependent transcriptional phenotypes and

therapeutic responses to Menin-MLL inhibitors.

Response to Menin-MLL inhibitors in primary human AML is accompanied by

transcriptional induction of MLL3/4-UTX-dependent tumor suppressive programs

Small molecule inhibitors of the Menin-MLL interaction have shown promise in preclinical models

of acute lymphoid and myeloid leukemia[25,27,92] and are currently in Phase I/II clinical trials for

treatment of patients with acute leukemia (SNDX-5613 (NCT04067336), KO-539

(NCT04065399), and JNJ-75276617 (NCT04811560)). Notably, SNDX-5613 was recently

granted fast track designation by the FDA for treatment of relapsed/refractory acute leukemias.

We hypothesized that the Menin-UTX molecular switch described above would be operative in

leukemia patients treated with these Menin-MLL inhibitors. Consistent with our hypothesis,

longitudinal RNA-Seq analysis of AML cells derived from two patients with *NPM1c*-mutated

(Patient 1) and *MLL*-rearranged (Patient 2) leukemia treated with SNDX-5613 as part of the

AUGMENT-101 clinical trial (NCT04065399) showed concomitant suppression of canonical MLL-

FP target genes (e.g., *MEIS1*) and induction of CDK inhibitors (e.g., *CDKN2C*) (**Fig. 5A-E,**

Extended Data Fig. 25).

As an orthogonal approach, we analyzed a panel of genetically-annotated patient-derived

xenografts (PDXs) of *MLL*-rearranged acute leukemia for response and resistance to VTP-50469

(a close analog of SNDX-5613) *in vivo*[25] (**Fig. 5F, Extended Data Fig. 26A**). Mice transplanted

with an *MLL3* wild type PDX showed a potent response to VTP-50469 and significant induction

of Menin-UTX targets (**Fig. 5G-H, Extended Data Fig. 27A-B**), reflecting the activation of the

UTX-dependent tumor suppressive gene expression program defined in mouse models of AML

(**Fig. 3**). Conversely, an *MLL3* mutant PDX exhibited primary resistance to VTP-50469 and failed

to induce this gene expression program (**Fig. 5I-J, Extended Data Fig. 27A and 27C**), thereby

functionally linking the induction of Menin-UTX-targets to preclinical drug response. Targeted

DNA sequencing confirmed that the resistant PDX harbored a missense mutation in the *MLL3*

gene (**Extended Data Fig. 26B**). Thus, the gene expression programs regulated by the epistatic

relationship between the MLL1-Menin and MLL3/4-UTX complexes are at play in patients and

PDXs treated with orally bioavailable Menin-MLL inhibitors that are currently under clinical

investigation. Somatic mutations in members of the MLL3/4-UTX complex that impact catalytic

activity or protein complex composition may impact the induction of tumor suppressive gene

expression programs induced by Menin-MLL-inhibitors, thereby blunting clinical responses.

To test whether CDK4/6 inhibition can bypass the blunted induction of endogenous CDK inhibitors

in Menin inhibitor-resistant contexts, we treated mice harboring the *MLL3*-mutant AML PDX with

VTP-50469 (0.1% rodent diet) and Palbociclib (35mg/kg) (**Fig. 5I**). As predicted by our findings

(**Fig. 3I-J**), single treatment with Menin- or CDK4/6-inhibitors only led to a minor decrease in

leukemia burden in the bone marrow of recipient mice after 10 days of treatment (**Fig. 5K-L**).

However, combination treatment with VTP-50469 and Palbociclib induced significant regression

in the abundance of human leukemia cells (**Fig. 5K-L**). To determine whether treatment

responses were associated with transcriptional induction of UTX-dependent tumor suppressive

programs, we performed RNA-Seq from FACS-sorted human leukemia cells isolated from these

animals. Consistent with our findings, we observed significant induction of cell cycle arrest and

senescence-associated gene expression signatures in mice treated with both VTP-50469 and

Palbociclib (**Fig. 5M, Extended Data Fig. 27D**). Altogether, this data provides pre-clinical

evidence for the feasibility and efficacy of combined Menin- and CDK4/6-inhibition to overcome

the blunted induction of senescence-associated programs in patient-derived leukemias resistant

to Menin-inhibitor monotherapy.

Discussion

The Menin chromatin adaptor protein exhibits profound context-specific functions in different

tissues, acting as a tumor suppressor gene in neuroendocrine[14,15], lung[17,93], skin[16], and CNS

malignancies[18], and as an oncogenic cofactor in hepatocellular[19] and hematologic cancers[4,20].

Given that Menin can interact with similar cofactors in disparate settings, the basis for these

apparently paradoxical findings has remained largely unclear. For example, Menin functionally

cooperates with MLL proteins to activate transcription of the *Cdkn1b/p27^{Kip1}* and *Cdkn2d/Ink4d*

CDK inhibitors as a tumor suppressive mechanism in neuroendocrine tumors and lung

cancer[3,17,94,95], yet the same protein-protein interaction is critical for leukemia maintenance[4,10,20].

These contrasting observations suggest that Menin employs unique molecular mechanisms to

regulate chromatin and gene expression depending on the biological context.

Our study sheds light on these paradoxical observations by providing the first evidence for a

functional interaction between the mammalian histone methyltransferase complexes MLL1-Menin

and MLL3/4-UTX and challenges the paradigm that these complexes are restricted to certain

genomic compartments (**Fig. 6**). We show that the Menin-MLL and NF-Y complexes coordinately

co-repress the expression of a tumor suppressive network that involves the *Cdkn2c/Ink4c* and

Cdkn2d/Ink4d tumor suppressor genes in leukemia by direct binding to gene promoters.

Disruption of Menin-MLL or NF-Y complexes using genetic or pharmacological approaches

triggers a molecular switch that leads to recruitment of the MLL3/4-UTX complex to the promoters

of tumor suppressive genes, including the *Cdkn2c/Ink4c and Cdkn2d/Ink4d* CDK inhibitors[78]. This

molecular switch leads to UTX-dependent activation of gene expression and is associated with a

concomitant increase in the levels of the gene activation-associated histone marks H4K16ac and

H3K27ac, leading to robust induction of tumor suppressive programs like cellular senescence[73,75].

Thus, the molecular choreography between the MLL1-Menin and MLL3/4-UTX complexes

dictates cellular survival in leukemia.

Our study also shows that the first 500 amino acids of UTX (lacking the demethylase domain) are

sufficient to induce full length UTX-dependent cellular and transcriptional phenotypes. This finding

is particularly interesting given a recent study suggesting that UTX requires a much bigger protein

fragment to drive tumor suppressive activity and transcriptional regulation via phase separation

mechanisms[90]. Instead, our study decisively demonstrates that the first 500 amino acids of UTX

(lacking the core intrinsically disordered region defined by Shi *et al.*[90]) are necessary and sufficient

to drive tumor suppressive responses in the context of Menin-MLL inhibition. These results

establish a new paradigm by which UTX employs non-catalytic mechanisms to regulate gene

expression and cellular phenotypes that impact response to epigenetic therapies.

Based on our observation that UTX directly binds to and activates the expression of *Cdkn2c/Ink4c*

and Cdkn2d/Ink4d (which are natural inhibitors of the CDK4/6 kinases)[78], we combined Menin-

MLL inhibition with the FDA-approved CDK4/6 kinase inhibitor Palbociclib[79] and showed that this

combinatorial approach leads to a strong synergistic effect on inhibiting leukemia cell proliferation

in vitro and *in vivo*. As Palbociclib and other CDK4/6 inhibitors are already FDA-approved[81], our

proof-of-concept and pre-clinical studies suggest that this combinatorial approach could be a

viable therapeutic option to both boost the effectiveness of Menin inhibitors as monotherapies

while potentially overcoming the intrinsic resistance to Menin-MLL inhibition conferred by loss-of-

function of the MLL3/4-UTX complex.

Our study demonstrates — in murine and human systems — that the functional interplay between

MLL1-Menin and MLL3/4-UTX is critical for Menin-inhibitor responses due to induction of Menin-

UTX target genes. Notably, we were able to show that the induction of this gene expression

program is at play upon oral delivery of VTP-50469 to leukemia PDX models and SDNX5613 to

human AML subjects in the context of the AUGMENT-101 clinical trial (NCT04065399).

Importantly, the Menin-UTX signature was absent in a non-responsive PDX harboring a missense

mutation in *MLL3*, but could be re-engaged by additional treatment with an FDA-approved

CDK4/6 inhibitor, providing the first pre-clinical rationale to consider such combination therapy

approaches in clinical settings to induce cell cycle arrest and senescence programs in leukemia

cells when Menin-MLL inhibitor monotherapy fails to do so.

FIGURE LEGENDS

Figure 1. CRISPR screens uncover functional interplay between the mammalian

MLL1/2 and MLL3/4 chromatin modifying complexes. (A) CRISPR-Cas9-based

screening strategy to identify chromatin regulators that modulate response to Menin

inhibition. CPD, cell population doublings; sgRNA, single guide RNA; TF, final timepoint;

T0, initial timepoint. **(B)** Top 20 most significantly enriching (red) and depleting (blue)

genes in the MI-503-treated CRISPR screen. Gene scores are shown as the mean $\log_2$

fold-change in abundance of the 6 sgRNAs targeting each gene in each condition. **(C)**

Genome-wide screening data showing gene-level ranking based on differential

enrichment of sgRNAs under Menin-MLL inhibitor treatment (VTP-50469) relative to

vehicle (DMSO). Differential (Δ) beta-score between VTP-50469 and DMSO conditions

was calculated using MaGeCK. A positive Δ beta-score denotes enrichment of specific

gene-targeting sgRNAs. A negative Δ beta-score denotes depletion of specific gene-

targeting sgRNAs. Red circles denote MLL3/4-UTX complex subunits. Yellow circles

denote PRC1.1 complex subunits. **(D)** Schematic representation of the top scoring

chromatin regulators in the chromatin-focused MI-503 screen and their corresponding

protein complexes. Red denotes enriching subunits. Blue denotes depleting subunits.

Color scale represents the $\log_2$ fold-change in MI-503. **(E)** Cell viability assay from cells

treated with DMSO (vehicle, black) or MI-503 (red) after 96 hours (mean±SEM, n=3

infection replicates, *P*-value calculated by Student's t-test). sgCtrl, control sgRNA

targeting a non-genic region on chromosome 8. **(F)** Differential fitness is shown as the

relative fitness of double positive cells (sgMen1-RFP + sgUtx-BFP or sgMen1-RFP +

sgCtrl-BFP) to single positive cells (sgMen1-RFP) 16 days post-infection measured by

flow cytometry (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-

test).

Figure 2. MLL1-Menin complex and NF-Y restrict chromatin occupancy of MLL3/4-

UTX at promoter regions. (A) Average chromatin occupancy of Menin at transcription

start sites (TSS), and 2kb upstream and downstream of the TSS are also included.

Signals corresponding to cells treated with MI-503 (solid) compared to cells treated with

DMSO (dotted) after 96 hours are shown. RPM, reads per million. **(B)** Average chromatin

occupancy of UTX at transcription start sites (TSS), and 2kb upstream and downstream

of the TSS are also included. Signals corresponding to cells treated with MI-503 (solid)

after 96 hours compared to cells treated with DMSO (dotted) after 96 hours are shown.

RPM, reads per million. **(C)** Relative *Utx* mRNA levels determined by qPCR in MLL-AF9

cells treated with MI-503 (red) versus DMSO (black) after 96 hours (mean ± SEM, n=3

replicates, *P*-value calculated by Student's t-test). **(D)** Immunoblot analysis of Menin,

UTX, and HSP90 proteins (loading control) upon MI-503 treatment of MLL-AF9 cells after

96 hours. **(E)** Heatmaps displaying Menin (black) and UTX (purple) ChIP-Seq signals

mapping to a 4kb window around TSSs. Data is shown for DMSO and MI-503-treated

cells after 96 hours. Metaplot represents the average ChIP-Seq signal for Menin or UTX

at promoters that are UTX+ (green) or UTX- (black) post MI-503 treatment. **(F)** Genome

browser representation of ChIP-Seq normalized reads for Menin (black) and UTX (purple)

in MLL-AF9 leukemia cells treated with either vehicle or MI-503 after 96 hours. **(G)** ChIP-

Seq normalized reads per 10-kb bin for Menin (MI-503 vs DMSO, y-axis) and UTX (MI-

503 vs DMSO, x-axis). Pearson's correlation coefficient is indicated.

Figure 3. Transcriptional co-regulation of tumor suppressive pathways by Menin

and UTX. (A) Volcano plot of differentially expressed genes in mouse MLL-AF9 cells

treated with Menin inhibitor (MI-503) or vehicle (DMSO) after 96 hours. Significantly

($P<0.05$) downregulated (DN) genes are shown on the left (n=921 genes). Significantly

($P<0.05$) upregulated (UP) genes are shown on the right (n=1030). **(B)** Upset plot

showing significant overlap (red) between genes that undergo replacement of Menin by

UTX at their respective promoters and MI-503-induced genes. *P*-value for overlap is

shown. **(C)** Boxplot showing expression levels of genes that are induced by MI-503

treatment and are bound by UTX at their promoters in this condition and by Menin at

steady-state. Expression levels are shown for Utx^{WT} and Utx^{KO} leukemia cells. **(D)** Gene

set enrichment analysis (GSEA) showing that Menin-UTX target genes induced by MI-

503 are significantly enriched for genes and pathways regulating cellular senescence.

FDR, false discovery rate; NES, normalized enrichment score. **(E)** Heatmap denoting

relative gene expression levels of senescence-associated cell cycle and SASP genes

measured by RNA-Seq from Utx^{WT} and Utx^{KO} MLL-AF9 leukemia cells treated with Menin

inhibitor (MI-503) or vehicle (DMSO) after 96 hours. Senescence-associated cell cycle

and SASP gene list was manually curated[72]. **(F) - (G)** Secreted levels of IL-6 and IFNβ-1

in Utx^{WT} (black) and Utx^{KO} (red) MLL-AF9 leukemia cells treated with Menin inhibitor (MI-

503) or vehicle (DMSO) after four or six days. Data are quantified as pg/mL of secreted

cytokine per million cells and represent the mean of three biological replicates. *P*-value

calculated by Student's t-test. **(H)** Genome browser representation of ChIP-Seq (top) and

RNA-Seq (bottom) normalized reads (average RPKM) for *Cdkn2c* and *Cdkn2d* loci from

Utx^{WT} or Utx^{KO} MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours. **(I)**

Proposed model and rationale for combination therapies based on Menin-MLL and

CDK4/6 inhibitors. Our data support a model whereby **(1)** Menin restricts UTX-mediated

transcriptional activation of tumor suppressor genes, including *Cdkn2c* and *Cdkn2d*,

which are natural inhibitors of the CDK4 and CDK6 kinases, which in turn inhibit cell cycle

arrest and senescence. Our model predicts that CDK4/6 inhibition using Palbociclib

should boost the anti-cancer activity of Menin-MLL inhibitors, which we show induces a

MLL3/4-UTX tumor suppressive axis, by more potently inhibiting these downstream

kinases. On the other hand, Menin is known to be required for activation of MLL-FP

targets like *Meis1* and *Cdk6* itself to sustain leukemia **(2)**. Thus, our model predicts that

combination therapies based on Menin-MLL and CDK4/6 inhibitors should act

synergistically to suppress leukemia proliferation by potently engaging two parallel

pathways that converge on regulation of cell cycle progression. **(J)** Relative viability of

Utx^{WT} and Utx^{KO} MLL-AF9 leukemia cells treated with either DMSO, MI-503, Palbociclib,

or a combination of both inhibitors for 6 days (mean±SEM, n=3 replicates, *P*-value

calculated by Student's t-test).

Figure 4. Catalytic activity of UTX is not required to elicit UTX-dependent cellular

and transcriptional functions. (A) Schematic of UTX protein. Highlighted are the 8

tetratricopeptide repeats (TPR) (93-385aa) and the histone demethylase (JmjC) domain

(1095-1258aa). Three 500 amino acid long truncations are also represented. **(B)** Growth

competition assay in mouse *Utx*^{KO} MLL-AF9 leukemia cells expressing different RFP-

tagged *Utx* cDNAs and treated with MI-503 for 2 and 6 days. Graph shows the relative

growth of cells infected with RFP-tagged *Utx* cDNAs measured by flow cytometry

(mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test). **(C)**

Principal component analysis (PCA) using RNA-Seq gene expression data from *Utx*^{KO}

MLL-AF9 leukemia cells expressing different RFP-tagged *Utx* cDNAs and treated with

DMSO or MI-503 for 96 hours. **(D)** Mean normalized read counts of the cyclin-dependent

kinase inhibitor, *Cdkn2c*, across three RNA-Seq replicate samples from different *Utx*

truncations in mouse MLL-AF9 leukemia and treated with DMSO or MI-503 for 96 hours.

Figure 5. *In vivo* response to Menin-MLL inhibition is accompanied by induction of

MLL3/4-UTX-dependent tumor suppressive programs. (A) Longitudinal flow

cytometry analysis measuring the fraction of CD45lo, cKIT+ leukemia cells in the

peripheral blood of an *NPM1c* mutant patient (Patient 1) during treatment cycle 1 of

SDNX-5613 as part of the AUGMENT-101 clinical trial (NCT04065399). **(B)** Heatmap

denoting temporal gene expression changes among MLL-FP targets in FACS-sorted AML

blasts cells from Patient 1. Plotted are all MLL-FP targets that are differentially expressed

at day 14 vs. day 0 of treatment cycle 1. **(C)** Temporal expression levels of genes involved

in cell cycle arrest and senescence (*CDKN1A*, *CDKN2B*, and *CDKN2C*) and MLL-FP

targets (*HOXA9*, *CDK6*, *PBX3*, and *MEIS1*) in AML cells from Patient 1 treated with

SDNX-5613. **(D)** Heatmap denoting temporal gene expression level changes among

MLL-FP targets in FACS-sorted AML blasts cells from Patient 2 treated with SDNX-5613

as part of the AUGMENT-101 clinical trial (NCT04065399). Plotted are all MLL-FP targets

that are differentially expressed at day 10 vs. day 0 of treatment cycle 1. **(E)** Temporal

expression levels of genes involved in cell cycle arrest and senescence (*CDKN1A*,

CDKN1B, and *CDKN3*) and MLL-FP targets (*HOXA7*, *CDK6*, *MEF2C*, and *MEIS1*) in AML

cells from Patient 2 treated with SDNX-5613. **(F)** Schematic of *in vivo* treatment

experiments using genetically-defined AML patient-derived xenografts (PDXs). NSG mice

were transplanted with either *MLL3*-WT or *MLL3*-Mutant AML PDXs and, upon disease

engraftment, were randomized into VTP-50469 (0.1% in rodent diet) or normal chow for

a duration of four weeks. Disease progression was monitored weekly by bleeding and

AML cells were sorted at 7 days post initiation of treatment using magnetic mouse cell

depletion from the bone marrow of animals to perform RNA-Seq. **(G)** Disease progression

as measured by the percentage of human CD45+ cells in the peripheral blood of mice

harboring *MLL3*-WT leukemia treated with vehicle (grey) or VTP-50469 (blue). **(H)**

Boxplot denoting gene expression levels of Menin-UTX targets in AML cells isolated from

mice harboring *MLL3*-WT leukemia treated with vehicle (grey) or VTP-50469 (blue). **(I)**

Disease progression as measured by the percentage of human CD45+ cells in the

peripheral blood of mice harboring *MLL3*-mutant leukemia treated with vehicle (grey) or

VTP-50469 (blue). **(J)** Boxplot denoting gene expression levels of Menin-UTX targets in

AML cells isolated from mice harboring *MLL3*-mutant leukemia treated with vehicle (grey)

or VTP-50469 (blue). **(K)** Leukemia burden in the bone marrow of recipient mice

transplanted with the *MLL3*-mutant PDX and treated with VTP-50469, Palbociclib or the

combination of these two inhibitors (measured by % human CD45+ cells). VTP-50469

was administered via drug-supplemented rodent chow (0.1%) for 10 days, Palbociclib

was given once daily via intraperitoneal injections (35mg/kg) for 7 days. **(L)** FACS plots

showing the abundance of human leukemia cells in representative recipient mice from

each treatment group. **(M)** Heatmap denoting changes in a cell cycle-associated gene

expression signature in FACS-sorted human leukemia cells isolated from recipient mice

transplanted with the *MLL3*-mutant PDX and treated with VTP-50469, Palbociclib, or the

combination of these two inhibitors.

Figure 6. A molecular switch between mammalian MLL complexes dictates

response to Menin-MLL inhibition.

ACKNOWLEDGEMENTS

We thank all members of the Allis, Lowe, and Armstrong laboratories for their help and support;

Richard Phillips, Robert G. Roeder, Tom W. Muir, Benjamin A. Garcia, and Charles J. Sherr for

scientific discussions; David Chen (Chun-Wei Chen) for mouse MLL-AF9 cells, Zhaohui Feng for

human leukemia cells, and Laura Whitman (Agilent) for OLS support. C.D.A. was supported by

US National Institutes of Health (NIH) grants (P01CA196539 and 5R01CA204639-03), the

Leukemia and Lymphoma Society (LLS-SCOR 7006-13), and the Rockefeller University and St.

Jude Children's Research Hospital Collaborative on Chromatin Regulation in Pediatric Cancer.

S.W.L was supported by a NIH/NCI grant (R01 CA190261) and an Agilent Thought Leader Award,

as well as the MSKCC cancer center support grant (P30 CA008748). S.W.L. is the Geoffrey

Beene Chair of Cancer Biology and a Howard Hughes Medical Institute Investigator. S.A.A was

supported by NIH grants CA176745, CA206963, CA204639 and CA066996. Y.M.S.F was supported by the Damon Runyon-Sohn Pediatric Cancer Fellowship (DRSG-21-17) and NIGMS-MOSAIC K99/R00 Career Development Award (1K99GM140265-01). F.J.S.R. was partially supported by the MSKCC TROT program (5T32CA160001), a MSKCC GMTEC Postdoctoral Researcher Innovation Grant, and is a HHMI Hanna Gray Fellow. F.P. was supported by the German Research Foundation (DFG, PE 3217/1-1) and a Momentum Fellowship award by the Mark Foundation for Cancer Research. D.W.B. was supported by a Ruth L. Kirschstein National Research Service Award (5F32CA217068). E.R.K. was supported by an F31 NRSA predoctoral fellowship from the NIH/NCI (F31CA192835).

AUTHOR CONTRIBUTIONS

Y.M.S.F., F.J.S.R., F.P, S.A.A, S.W.L., and C.D.A. conceived the study and wrote the manuscript with input from all the authors; Y.M.S.F., F.J.S.R., F.P., Y.X., L.G., M.C.B., and D.C. conducted experiments; D.W.B., T.C., and Y.J.H. analyzed RNA-Seq and ChIP-Seq data; Y.J.H. and E.R.K. analyzed CRISPR-screen data. S.G. and X.S.L. provided genome-wide CRISPR library. A.V.K., M.M., and E.dS. provided patient-derived xenograft samples. R.M.S. provided primary AML samples from patients in the Syndax trial A.S. provided conceptual advice. S.A.A, S.W.L., and C.D.A. supervised the study and secured funding.

DECLARATION OF INTERESTS

C.D.A. is a co-founder of Chroma Therapeutics and Constellation Pharmaceuticals and a Scientific Advisory Board member of EpiCypher. S.W.L. is an advisor for and has equity in the following biotechnology companies: ORIC Pharmaceuticals, Faeth Therapeutics, Blueprint Medicines, Geras Bio, Mirimus Inc., and PMV Pharmaceuticals. S.W.L. also acknowledges receiving funding and research support from Agilent Technologies for the purposes of massively

637   parallel oligo synthesis. S.A.A. has been a consultant and/or shareholder for Vitae/Allergan

638   Pharmaceuticals, Epizyme Inc., Imago Biosciences, Cyteir Therapeutics, C4 Therapeutics, Syros

639   Pharmaceuticals, OxStem Oncology, Accent Therapeutics, and Mana Therapeutics. S.A.A. has

640   received research support from Janssen, Novartis, and AstraZeneca. The remaining authors

641   declare no competing interests.

MATERIALS AND METHODS

<u>Plasmids and sgRNA cloning</u>

To generate stable Cas9-expressing cell lines, we used lentiCas9-Blast (Addgene, 52962).

Human wild-type or mutant (K4M) histone H3.1 were cloned into pCDH-EF1-MCS-IRES-RFP

(System Biosciences, CD531A-2). To express sgRNAs, we generated the pUSEPR (U6-sgRNA-

EFS-Puro-P2A-TurboRFP) and pUSEPB (U6-sgRNA-EFS-Puro-P2A-TagBFP) lentiviral vectors

by Gibson assembly of the following DNA fragments: (i) PCR-amplified U6-sgRNA (improved

scaffold)[96] cassette, (ii) PCR-amplified EF1s promoter, (iii) PCR-amplified Puro-P2A-TurboRFP

or -TagBFP gene fragment (IDT), and (iv) BsrGI/PmeI-digested pLL3-based lentiviral backbone[97].

For sgRNA cloning, pUSEPR and pUSEPB vectors were linearized with BsmBI (NEB) and ligated

with BsmBI-compatible annealed and phosphorylated oligos encoding sgRNAs using high

concentration T4 DNA ligase (NEB). All sgRNA sequences used are listed in Supplementary

Table 1.

<u>Cell culture</u>

Mouse MLL-AF9 leukemia cells were kindly shared by David Chen (Chun-Wei Chen) and were

originally generated by transformation of female mouse bone marrow Lin-Sca1+cKit+ (LSK) cells

with an MSCV-IRES-GFP (pMIG) retrovirus expressing the human MLL-AF9 fusion protein and

transplanted into sub-lethally irradiated recipient mice as described previously[29,98]. Leukemic

blasts were harvested from moribund mice and cultured in vitro in IMDM (Gibco) supplemented

with 15% FBS (Gibco), mouse IL-6 (10ng/μL, PeproTech), mouse IL-3 (10ng/μL, PeproTech),

mouse SCF (20ng/μL, PeproTech), penicillin (100U/mL, Gibco), streptomycin (100μg/uL, Gibco),

L-glutamine (2mM, Gibco), and plasmocin (5μg/mL, InvivoGen). Human leukemia cell lines

MV4;11 and OCI-AML3 were kindly shared by Zhaohui Feng and were cultured in RPMI 1640

(Corning) supplemented with 10% FBS (Gibco), penicillin (100U/mL, Gibco), streptomycin

(100μg/uL, Gibco), L-glutamine (2mM, Gibco), and plasmocin (5μg/mL, InvivoGen). Mouse NIH-

3T3 cells were maintained in DMEM (Corning) supplemented with 10% FCS (ATCC), penicillin

(100U/mL, Gibco), streptomycin (100μg/uL, Gibco), and plasmocin (5μg/mL, InvivoGen). Human

HEK293 cells were maintained in DMEM (Corning) supplemented with 10% FBS (Gibco),

penicillin (100U/mL, Gibco), streptomycin (100μg/uL, Gibco), and plasmocin (5μg/mL,

InvivoGen). Cas9-expressing cells were generated by lentiviral transduction of lentiCas9-Blast

followed by Blasticidin (InvivoGen) selection and validation of Cas9 expression and activity. All

cells were confirmed to be free of Mycoplasma contamination and cultured at 37°C and 5% CO_2.

Virus production

Lentiviruses were produced by co-transfection of HEK293T (ATCC) cells with pUSEPR-EpiV2.0

sgRNA library, individual sgRNA plasmids, or lentiCas9-Blast, and packaging vectors psPax2

(Addgene, 12260) and pMD2.G (Addgene, 12259) using Lipofectamine 2000 (Invitrogen). Viral

supernatants were collected at 48 and 72 hours post transfection and stored at -80°C.

Transduction of cell lines

Leukemia cells were seeded at a density of 2.5 x 10^5 cells/well of a non-TC-treated 12-well plate

in complete medium containing polybrene (10μg/mL, EMD Millipore), and then transduced with

lentivirus by centrifugation at 2,500rpm for 90 minutes at 37°C. After a 24-hour incubation, cells

were transferred to a new plate containing fresh culture medium. Antibiotic selection or cell sorting

was done 48 hours post transduction.

<u>Drug treatments</u>

For MI-503 (Active Biochem) treatments, leukemia cells were seeded at a density of 4×10^5

cells/mL, treated with limiting dilutions of the inhibitor as indicated or 0.25% DMSO (vehicle). Cells

were re-plated every 4 days to the initial density and re-treated. Viability was assessed at various

time points by using the CellTiter-Glo Luminescent Cell Viability assay (Promega) following the

manufacturer's guidelines. Ratio of luminescence signal from metabolically active cells in MI-503

versus DMSO were plotted to calculate IC50 values (Prism 8, GraphPad). For MI-503 (Active

Biochem) and Palbociclib HCl (Selleckchem) combination treatments, 25,000 leukemia cells in

250µL of drug-containing medium were seeded in a 48-well plate and viability was assessed every

4 days by using the CellTiter-Glo Luminescent Cell Viability assay (Promega). For RNA-Seq and

ChIP-Seq experiments, cells were cultured at 4×10^5 cells/mL, treated with MI-503

(concentrations as indicated in figure legends) or 0.25% DMSO for 4 days. Cells were collected,

washed with PBS, pelleted, and flash-frozen before RNA or chromatin isolation. For *in vivo* VTP-

50649 treatment, mice were randomly assigned to either normal or 0.1% VTP-50469 rodent

special diet[26]. Mice were bled weekly to monitor leukemia burden and euthanized when showing

clinical signs of disease (experimental endpoint).

<u>Flow cytometric analyses</u>

Immunophenotyping of leukemia cells treated with MI-503 (or vehicle), was done by collecting

cells post treatment and staining using the indicated conjugated primary antibodies. Stained

samples were analyzed on an LSRFortessa (BD Biosciences) flow cytometer. Data analysis was

performed using FlowJo (BD Biosciences) software. Intracellular antigens detection was done by

using the Foxp3/Transcription Factor Staining Buffer Set (eBioscience) following the

manufacturer's guidelines. Conjugated primary antibodies used were: Pacific Blue anti-CD11b

(Biolegend, 101224), Alexa Fluor 647 anti-Cas9 (CST, 48796).

<u>Xenograft models of AML</u>

All animal experiments were performed with the approval of Dana-Farber Cancer Institute's

Institutional Animal Care and Use Committee (IACUC). NOD.Cg-*Prkdc^{scid} Il2rg^{tm1Sug}*/JicTac (NOG)

mice were obtained from Taconic Biosciences (Rensselaer, NY, USA). Non-irradiated 8-12 weeks

old adult mice were transplanted with previously established patient-derived xenografts

(PDXs)[25,99] via tail vein injection (250,000 cells/mouse). Engraftment of human cells (hCD45$^+$)

was analyzed and monitored longitudinally by weekly bleeding to quantify hCD45$^+$ cells in the

peripheral blood by flow cytometry with human CD45-PE and anti-mouse CD45-APC-Cy7

antibodies (Biolegend, San Diego, CA, USA). Mice were monitored closely to detect disease

onset and treatment started when hCD45$^+$ cells were detectable in the peripheral blood. Mice

were randomly assigned to either normal or 0.1% VTP-50469 rodent special diet[25]. Mice were

bled weekly to monitor leukemia burden as described above and euthanized when showing

clinical signs of disease (experimental endpoint). Leukemia cells from a subset of these animals

were harvested after seven days of treatment to perform RNA-Seq.

<u>Longitudinal analysis of AML patient treated with SNDX-5613</u>

Peripheral blood of patients was taken under informed consent according to the Declaration of

Helsinki during routine blood draws at screening and different timepoints during the first cycle of

treatment with SDNX-5613 within the AUGMENT-101 clinical trial (NCT04065399). Peripheral

blood mononuclear cells (PBMCs) were subsequently isolated using Ficoll (BD Bioscience,

Franklin Lakes, NJ, USA) gradient centrifugation, viably frozen, and banked at the Dana-Farber

Cancer Institute, Boston, MA (approved institutional protocol, IRB: #01-206). For longitudinal

analysis, samples were thawed, washed twice in PBS, and stained with anti-human CD45 (PE)

and anti-human CD117 (APC) (Biolegend, San Diego, CA, USA). CD45-low/CD117+ leukemia

cells were FACS sorted (MA900 sorter, Sony Biotechnology, San Jose, CA, USA) and

subsequently processed for RNA-Seq (see methods section on RNA-Seq).

<u>Immunoblotting</u>

Whole cell lysates were separated by SDS-PAGE, transferred to a PVDF membrane (EMD

Millipore), blocked in 5% non-fat milk in TBS plus 0.5% Tween-20 (Sigma-Aldrich), probed with

primary antibodies, and detected with HRP-conjugated anti-rabbit or anti-mouse secondary

antibodies (GE Healthcare). Primary antibodies used included: anti-Cas9 (CST, 14697), anti-UTX

(CST, 33510), anti-Menin (Bethyl, A300-105A), anti-NF-YA (Santa Cruz Biotechnology, sc-

17753), anti-Actin (abcam, ab8224), anti-HSP90 (BD Biosciences, 610418), anti-HA (Biolegend,

901501), anti-H3K4me1 (abcam, ab8895), anti-H3K4me3 (CST, 9751), anti-H3 (abcam, ab1791).

<u>Locus-specific DNA sequencing</u>

To determine the mutational status of the *Men1* and *Utx* loci in cells targeted by CRISPR-Cas9,

we performed next generation sequencing of PCR-amplified target regions. Genomic DNA

(gDNA) was isolated using the DNeasy Blood & Tissue Kit (Qiagen) following the manufacturer's

guidelines. Amplification of target regions was performed from 500ng of gDNA using Q5 High-

Fidelity 2X Master Mix (NEB) and primers listed in Supplementary Table 1. PCR products were

purified using the QIAquick PCR Purification Kit (Qiagen) and sequenced on Illumina instruments

at GENEWIZ (Amplicon-EZ service).

<u>RNA isolation, qRT-PCR analyses, and RNA sequencing</u>

Total RNA was isolated from cells using the RNeasy kit (Qiagen, Hilden, Germany). RNA was

reverse transcribed with High-Capacity cDNA Reverse Transcription kits (Applied Biosystems)

following the manufacturer's instructions. Quantitative PCRs (qRT-PCRs) were performed using

the TaqMan Gene Expression Master Mix (Applied Biosystems) with the StepOne Real-Time

System (Applied Biosystems). TaqMan gene expression assays were used. ActB was used as

the endogenous control for normalization and relative gene expression was calculated by using

the comparative CT method. The mouse gene probes used were: *ActB* (Mm02619580_g1),

Hoxa9 (Mm00439364_m1), *Meis1* (Mm00487664_m1), *Utx* (*Kdm6a*) (Mm00801998_m1), *Men1*

(Mm00484957_m1). Quality of extracted RNA for sequencing was assessed by RIN using a

Bioanalyzer (Agilent) and quantified by TapeStation (Agilent). Poly(A) mRNA enrichment and

library preparation was performed using the NEBNext Poly(A) mRNA Magnetic Isolation Module

and NEBNext Ultra II RNA Library Prep kit (NEB). Sequencing was done using the Illumina

NextSeq500 to obtain >20 million 75bp single-end or 37bp paired-end reads per sample or at

Genewiz (HiSeq, 150bp, paired end) (Illumina, South Plainfield, NJ, USA).

<u>Cloning of sgRNA library targeting mouse chromatin regulators</u>

sgRNA sequences (six per gene) targeting 616 mouse chromatin regulators (for a total of 3,696

sgRNAs) (Supplementary Table 1) were designed using the Broad Institute sgRNA Designer

tool[100]. We also included 36 non-targeting control sgRNAs obtained from the GeCKOv2 Mouse

CRISPR library[101] (for a total of 3,732 sgRNAs). This library was divided into 6 pools (each

composed of 616 targeting and 6 non-targeting sgRNAs), synthesized by Agilent Technologies,

and cloned into the pUSEPR lentiviral vector[102] using a modified version of the protocol published

by Doench et al[100] to ensure a library representation of >10,000X. Briefly, each sub-pool was

selectively amplified using barcoded forward and reverse primers that append cloning adapters

at the 5' and 3' ends of the sgRNA insert (Supplementary Table 1), purified using the QIAquick

PCR Purification Kit (Qiagen), and ligated into BsmBI-digested and dephosphorylated pUSEPR

vector using high-concentration T4 DNA ligase (NEB). A minimum of 1.2 ug of ligated pUSEPR

plasmid DNA per sub-pool was electroporated into Endura electrocompetent cells (Lucigen),

recovered for one hour at 37°C, plated across four 15cm LB-Carbenicillin plates (Teknova), and

incubated at 37°C for 16 hours. The total number of bacterial colonies per sub-pool was quantified

using serial dilution plates to ensure a library representation of >10,000X (>6.2 million colonies

per sub-pool). The next morning, bacterial colonies were scraped and briefly expanded for 4 hours

at 37°C in 500mL of LB-Carbenicillin. Plasmid DNA was isolated using the Plasmid Plus Maxi Kit

(Qiagen).

To assess sgRNA distribution in each of the sub-pools, as well as the master pool (composed of

equimolar amounts of plasmid DNA from each individual sub-pool), we amplified the sgRNA target

region using primers that append Illumina sequencing adapters on the 5' and 3' ends of the

amplicon, as well as a random nucleotide stagger and unique demultiplexing barcode on the 5'

end (Supplementary Table 1). Library amplicons were size-selected on a 2.5% agarose gel,

purified using the QIAquick Gel Extraction Kit (Qiagen), and sequenced on an Illumina NextSeq

instrument (75nt single end reads).

Chromatin-focused CRISPR-Cas9 genetic screening

To ensure that most cells harbor a single sgRNA integration event, we determined the volume of

viral supernatant that would achieve an MOI of ~0.3 upon spinfection of a population of Cas9-

expressing leukemia cells. Briefly, cells were plated at a concentration of $2.5x10^5$ per well in 12-

well plates along with increasing volumes of master pool viral supernatant (0, 25, 100, 200, 500,

1000, and 2000µL) and polybrene (10 µg/mL, EMD Millipore). Cells were then centrifuged at

1,500rpm for 2 hours at 37°C and incubated at 37°C overnight. Viral infection efficiency was

determined by the percentage of tRFP+ cells assessed by flow cytometry on an LSRFortessa (BD

Biosciences) instrument 72 hours post infection.

Each step of the screen - from infection to sequencing - was optimized to achieve a minimum

representation of 1000X. To ensure a representation of 1000X at the transduction step, we

spinfected a total of 20 million cells across seven 12-well plates in triplicate (for a total of twenty

one 12-well plates) using the volume of viral supernatant that would achieve a 30% infection rate

(6 million transduced cells per technical replicate).

24 hours after infection, cells were pooled into 2 x T225 flasks (Corning) per infection replicate

and selected with 2.5µg/mL puromycin (Gibco) for 4 days. Subsequently, 6 million puromycin-

selected cells were pelleted and stored at -20°C (T0/Input population) while the rest were plated

into either DMSO- or MI-503-containing media (at an IC50 concentration) and cultured until the

population reached 12 cumulative population doublings (TF/Final). At least 6 million cells were

harvested and pelleted for this final time point. Genomic DNA from MLL-AF9 cells was isolated

using the DNeasy Blood & Tissue Kit (Qiagen) following the manufacturer's guidelines.

As previously published[103], we assumed that each cell contains approximately 6.6 pg of genomic

DNA (gDNA). Therefore, deconvolution of the screen at 1000X required sampling ~4 million x 6.6

pg of gDNA, or ~26.4 ug. We employed a modified 2-step PCR version of the protocol published

by Doench et al[100]. Briefly, we perform an initial "enrichment" PCR, whereby the integrated sgRNA

cassettes are amplified from gDNA, followed by a second PCR to append Illumina sequencing

adapters on the 5' and 3' ends of the amplicon, as well as a random nucleotide stagger and unique

demultiplexing barcode on the 5' end. Each "PCR1" reaction contains 25µL of Q5 High-Fidelity

2X Master Mix (NEB), 2.5µL of Nuc PCR#1 Fwd Primer (10µM), 2.5µL of Nuc PCR#1 Rev Primer

(10µM), and 5µg of gDNA in 20µL of water (for a total volume of 50µL per reaction). The number

of PCR1 reactions were scaled accordingly; therefore, we performed six PCR1 reactions per

technical replicate, per time point (T0 or TF), and per condition (DMSO or MI-503). PCR1

amplicons were purified using the QIAquick PCR Purification Kit (Qiagen) and used as template

for "PCR2" reactions. Each PCR2 reaction contains 25µL of Q5 High-Fidelity 2X Master Mix

(NEB), 2.5µL of a unique Nuc PCR#2 Fwd Primer (10µM), 2.5µL of Nuc PCR#2 Rev Primer

(10µM), and 300ng of PCR1 product in 20µL of water (for a total volume of 50µL per reaction).

We performed two PCR2 reactions per PCR1 product. Library amplicons were size-selected on

828 a 2.5% agarose gel, purified using the QIAquick Gel Extraction Kit (Qiagen), and sequenced on

829 an Illumina NextSeq500 instrument (75nt single end reads). All primer sequences are available

830 in Supplementary Table 1. PCR Program for PCR1 and PCR2: 1) 98°C x 30s; 2) 98°C x 10s; 3)

831 65°C x 30s; 4) 72°C x 30s; 5) Go to step 2 x 24 cycles; 6) 72°C x 2 min; 7) 4°C forever.

832 <u>Genome-wide CRISPR-Cas9 genetic screening</u>

833 Paired mouse genome-scale CRISPR-Cas9 screening libraries (M1/M2) were provided by

834 Shengqing Gu and Xiaole Shirley Liu (Addgene Pooled Library #1000000173). The M1 and M2

835 libraries cover protein coding genes of the genome with a total of 10 guide RNAs per gene.

836 Lentivirus was produced using each separate library pool and used to transduce each 5×10^8 MLL-

837 AF9 cells at low MOI. 48 hours after library transduction, cells were selected with blasticidin

838 (5ug/ml). After 5 days of antibiotic selection, a baseline (T0) sample was collected, and cells were

839 cultured in duplicate before harvest of terminal samples after 12 days (TF). Subsequently,

840 genomic DNA was isolated using phenol-chloroform extraction and sgRNA libraries were

841 deconvoluted using next-generation sequencing essentially as described above.

842 <u>Analysis of CRISPR-Cas9 genetic screen data</u>

843 FASTQ files were processed and trimmed to retrieve sgRNA target sequences followed by

844 alignment to the reference sgRNA library file. Sequencing read counts were summarized at gene

845 level per sample and used as input to run differential analysis using DESeq2 package. The log2

846 fold change values were used as 'Gene Score' for the final visualization. Genome-wide screening

847 data was analyzed using MAGeCK MLE essentially as described in the original publication[104].

848 See Supplementary Table 2 for all raw screening data.

849 <u>Growth competition assays</u>

Cas9-expressing cells were virally transduced with the designated constructs (pUSEPR-sgRNA,

pUSEPB-sgRNA, pCDH-cDNA) in 12-well plates at 30-40% infection rate (three infection

replicates). Cells were monitored by flow cytometry over time using an LSRFortessa (BD

Biosciences) flow cytometer and relative growth of sgRNA-containing cells was assesed. Flow

cytometry data was analyzed with FlowJo software (BD Biosciences). The percentage of single

positive (SP) (tRFP+ or BFP+) or double positive (DP) (tRFP+/BFP+) cells was normalized to

their respective "T0" time-point values (assessed on day 2 or 3 post transduction, as indicated in

the figure legend). Normalized values were log2-transformed, and the differential fitness of cells

was calculated as follow:

Differential Fitness = log2(Normalized DP) - log2(Normalized SP)

<u>Chromatin immunoprecipitation (ChIP)</u>

Cross-linking ChIP in mouse leukemia and NIH-3T3 cells was performed with $10\text{-}20\times10^7$ cells per

immunoprecipitation. After drug (or vehicle) treatment, cells were collected, washed once with

ice-cold PBS, and flash-frozen. Cells were resuspended in ice-cold PBS and cross-linked using

1% paraformaldehyde (PFA) (Electron Microscopy Sciences) for 5 minutes at room temperature

with gentle rotation. Unreacted PFA was quenched with glycine (final concentration 125mM) for

5 minutes at room temperature with gentle rotation. Cells were washed once with ice-cold PBS

and pelleted by centrifugation (800g for 5 minutes). To obtain a soluble chromatin extract, cells

were resuspended in 1mL of LB1 (50mM HEPES pH 7.5, 140mM NaCl, 1mM EDTA, 10%

glycerol, 0.5% NP-40, 0.25% Triton X-100, 1X complete protease inhibitor cocktail) and incubated

at 4°C for 10 minutes while rotating. Samples were centrifuged (1400g for 5 minutes),

resuspended in 1mL of LB2 (10mM Tris-HCl pH 8.0, 200mM NaCl, 1mM EDTA, 0.5mM EGTA,

1X complete protease inhibitor cocktail), and incubated at 4°C for 10 minutes while rotating.

Finally, samples were centrifuged (1400g for 5 minutes) and resuspended in 1mL of LB3 (10mM

Tris-HCl pH 8.0, 100mM NaCl, 1mM EDTA, 0.5mM EGTA, 0.1% sodium deoxycholate, 0.5% N-

Lauroylsarcosine, 1X complete protease inhibitor cocktail). Samples were homogenized by

passing 7-8 times through a 28-gauge needle and Triton X-100 was added to a final concentration

of 1%. Chromatin extracts were sonicated for 14 minutes using a Covaris E220 focused

ultrasonicator. Lysates were centrifuged at maximum speed for 10 minutes at 4°C and 5% of

supernatant was saved as input DNA. Beads were prepared by incubating in 0.5% BSA in PBS

and antibodies overnight (100µL of Dynabeads Protein A or Protein G (Invitrogen) plus 20µL of

antibody). Antibodies used were: anti-Menin (Bethyl, A300-105A), anti-UTX (Bethyl, A302-374A),

anti-MLL1 (N-term-specific, Bethyl, A300-086A), anti-MLL3/4 (kindly provided by the Wysocka

laboratory[41], anti-NF-YA (Santa Cruz Biotechnology, sc-17753), anti-H3K4me1 (abcam, ab8895),

anti-H3K4me3 (Active Motif, 39159), and anti-H4K16ac (Active Motif, 39167). Antibody-Beads

mixes were washed with 0.5% BSA in PBS and then added to the lysates overnight while rotating

at 4°C. Beads were then washed six times with RIPA buffer (50mM HEPES pH 7.5, 500mM LiCl,

1mM EDTA, 0.7% sodium-deoxycholate, 1% NP-40) and once with TE-NaCl Buffer (10mM Tris-

HCl pH 8.0, 50mM NaCl, 1mM EDTA). Chromatin was eluted from beads in Elution buffer (50mM

Tris-HCl pH 8.0, 10mM EDTA, 1% SDS) by incubating at 65°C for 30 minutes while shaking,

supernatant was removed by centrifugation, and crosslinking was reversed by further incubating

chromatin overnight at 65°C. The eluted chromatin was then treated with RNase A (10mg/mL) for

1 hour at 37°C and with Proteinase K (Roche) for 2 hours at 55°C. DNA was purified by using

phenol-chloroform extraction followed with ethanol precipitation. The NEBNext Ultra II DNA

Library Prep kit was used to prepare samples for sequencing on an Illumina NextSeq500 (75bp

read length, single-end, or 37bp read length, paired-end).

ChIP-Seq analysis

ChIP-sequencing samples were sequenced using the Illumina NextSeq500. ChIP-seq reads were

aligned using Rsubread's align method and predicted fragment lengths calculated by the ChIPQC

R Bioconductor package[105,106]. Normalized, fragment extended signal bigWigs were created using

the rtracklayer R Bioconductor package. Peak calls were made in MACS2 software[107]. Read

counts in peaks were calculated using the featureCounts method in the Rsubread library[106].

Differential ChIP-seq signal were identified using the binomTest from the edgeR R Bioconductor

package[108]. Annotation of genomic regions to genes, biological functions, and pathways were

performed using the ChIPseeker R Bioconductor package[109]. Meta-peak plots were produced

using the soGGi package and ChIP-seq signal heatmaps generated using the Deeptools and

profileplyr software[110]. Plots showing ChIP-Seq read signal over transcription start sites (TSSs)

were made with the ngs.plot software package (v2.61) (ref. 111). Overlaps between peak sets

were determined using the ChIPpeakAnno R Bioconductor package with a maximum gap

between peaks set to 1kb (ref. 112). Peaks were annotated with both genes and the various types

of genomic regions using the ChIPseeker R Bioconductor package[109]. Range-based heatmaps

showing signal over genomic regions were generated using the soGGi and profileplyr R

Bioconductor package to quantify read signal and group the peak ranges and the deepTools

software package (v3.3.1) to generate the heatmaps[110]. Any regions included in the ENCODE

blacklisted regions of the genome were excluded from all region-specific analyses[113]. For some

ChIP-seq experiments, raw Illumina NextSeq BCL files were converted to FASTQs using Illumina

bcl2fastq v02.14.01.07 and reads trimmed using Trimmomatic v0.36 (phred quality threshold 33)

and uploaded to the to the Basepair-server (basepairtech.com). Alignment and ChIP-seq QC was

performed on the basepair platform (Bowtie2). Peak calling was performed using MACS (v.1.4)

within the basepair platform utilizing the default parameters.

<u>RNA-Seq analysis</u>

RNA-Seq samples were sequenced using the Illumina NextSeq500. Transcript abundance was

computed from FASTQ files using Salmon and the GENCODE reference transcript sequences,

transcript counts were imported into R with the tximport R Bioconductor package, and differential

gene expression was determined with the DESeq2 R Bioconductor package[114-116]. The data was

visualized using the ggplot2 R package. Normalized counts were extracted from the DESeq2

results and z-scores for the indicated gene sets were visualized using both heatmaps and

boxplots. Heatmaps showing gene expression changes across samples were generated using

the pheatmap R package and boxplots were made with the ggplot2 R package. Gene ontology

analysis using the KEGG 2019 database was performed using the Enrichr tool[56,68].

<u>Statistical analyses</u>

Statistical tests were used as indicated in Figure legends. Generation of plots and statistical

analyses were performed using Prism 8 (GraphPad). Error bars represent standard deviation,

unless otherwise noted. We used Student's t-test (unpaired, two-tailed) to assess significance

between treatment and control groups, and to calculate P values. $P<0.05$ was considered

statistically significant.

<u>Source data availability</u>

Data supporting the findings of this study are reported in Extended Data Figures 1-27 and

Supplementary Tables 1-4. All raw data corresponding to high-throughput approaches (CRISPR

screens, RNA-Seq, and ChIP-Seq) is available through NCBI GEO (Accession: GSE186711. Go

to   https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE186711   and   enter   token

wpkrckicxnsvxit into the box to get access to datasets). All reagents and materials generated in

this study will be available to the scientific community through Addgene and/or MTAs. Further

information and requests for resources and reagents should be directed to and will be fulfilled by

the Lead Contacts: C. David Allis (alliscd@rockefeller.edu), Scott W. Lowe (lowes@mskcc.org),

and Scott. A. Armstrong (Scott_Armstrong@dfci.harvard.edu).

REFERENCES

1   Guru, S. C. *et al.* Menin, the product of the MEN1 gene, is a nuclear protein.
948   *Proceedings of the National Academy of Sciences* **95**, 1630-1634,
949   doi:papers3://publication/doi/10.1073/pnas.95.4.1630 (1998).
950   2   Hughes, C. M. *et al.* Menin associates with a trithorax family histone
951   methyltransferase complex and with the Hoxc8 locus. *Mol Cell* **13**, 587-597,
952   doi:papers3://publication/doi/10.1016/s1097-2765(04)00081-4 (2004).
953   3   Milne, T. A. *et al.* Menin and MLL cooperatively regulate expression of cyclin-
954   dependent kinase inhibitors. *Proceedings of the National Academy of Sciences*
955   **102**, 749-754, doi:papers3://publication/doi/10.1073/pnas.0408836102 (2005).
956   4   Yokoyama, A. *et al.* The menin tumor suppressor protein is an essential
957   oncogenic cofactor for MLL-associated leukemogenesis. *Cell* **123**, 207-218,
958   doi:papers3://publication/doi/10.1016/j.cell.2005.09.025 (2005).
959   5   Bres, V., Yoshida, T., Pickle, L. & Jones, K. A. SKIP interacts with c-Myc and
960   Menin to promote HIV-1 Tat transactivation. *Mol Cell* **36**, 75-87,
961   doi:papers3://publication/doi/10.1016/j.molcel.2009.08.015 (2009).
962   6   Agarwal, S. K. & Jothi, R. Genome-wide characterization of menin-dependent
963   H3K4me3 reveals a specific role for menin in the regulation of genes implicated
964   in MEN1-like tumors. *PLoS ONE* **7**, e37952,
965   doi:papers3://publication/doi/10.1371/journal.pone.0037952 (2012).
966   7   Huang, J. *et al.* The same pocket in menin binds both MLL and JUND but has
967   opposite effects on transcription. *Nature 2018* **482**, 542-546,
968   doi:papers3://publication/doi/10.1038/nature10806 (2012).
969   8   Kaji, H., Canaff, L., Lebrun, J. J., Goltzman, D. & Hendy, G. N. Inactivation of
970   menin, a Smad3-interacting protein, blocks transforming growth factor type beta
971   signaling. *Proceedings of the National Academy of Sciences* **98**, 3837-3842,
972   doi:papers3://publication/doi/10.1073/pnas.061358098 (2001).
973   9   Sowa, H. *et al.* Inactivation of menin, the product of the multiple endocrine
974   neoplasia type 1 gene, inhibits the commitment of multipotential mesenchymal
975   stem cells into the osteoblast lineage. *J Biol Chem* **278**, 21058-21069,
976   doi:papers3://publication/doi/10.1074/jbc.M302044200 (2003).
977   10   Yokoyama, A. & Cleary, M. L. Menin critically links MLL proteins with LEDGF on
978   cancer-associated target genes. *Cancer Cell* **14**, 36-46,
979   doi:papers3://publication/doi/10.1016/j.ccr.2008.05.003 (2008).
980   11   Balogh, K., Rácz, K., Patócs, A. & Hunyady, L. Menin and its interacting proteins:
981   elucidation of menin function. *Trends Endocrinol. Metab.* **17**, 357-364,
982   doi:papers3://publication/doi/10.1016/j.tem.2006.09.004 (2006).
983   12   Yokoyama, A. *et al.* Leukemia proto-oncoprotein MLL forms a SET1-like histone
984   methyltransferase complex with menin to regulate Hox gene expression. *Mol Cell*
985   *Biol* **24**, 5639-5649, doi:papers3://publication/doi/10.1128/MCB.24.13.5639-
986   5649.2004 (2004).
987   13   Wang, P. *et al.* Global analysis of H3K4 methylation defines MLL family member
988   targets and points to a role for MLL1-mediated H3K4 methylation in the
989   regulation of transcriptional initiation by RNA polymerase II. *Mol Cell Biol* **29**,
990   6074-6085, doi:papers3://publication/doi/10.1128/MCB.00924-09 (2009).

991   14   Chandrasekharappa, S. C. *et al.* Positional cloning of the gene for multiple
992   endocrine neoplasia-type 1. *Science* **276**, 404-407,
993   doi:papers3://publication/doi/10.1126/science.276.5311.404 (1997).
994   15   Jiao, Y. *et al.* DAXX/ATRX, MEN1, and mTOR pathway genes are frequently
995   altered in pancreatic neuroendocrine tumors. *Science* **331**, 1199-1203,
996   doi:papers3://publication/doi/10.1126/science.1200609 (2011).
997   16   Asgharian, B. *et al.* Cutaneous tumors in patients with multiple endocrine
998   neoplasm type 1 (MEN1) and gastrinomas: prospective study of frequency and
999   development of criteria with high sensitivity and specificity for MEN1. *J. Clin.*
1000   *Endocrinol. Metab.* **89**, 5328-5336, doi:papers3://publication/doi/10.1210/jc.2004-
1001   0218 (2004).
1002   17   Pei, X.-H., Bai, F., Smith, M. D. & Xiong, Y. p18Ink4c collaborates with Men1 to
1003   constrain lung stem cell expansion and suppress non-small-cell lung cancers.
1004   *Cancer Res* **67**, 3162-3170, doi:papers3://publication/doi/10.1158/0008-
1005   5472.CAN-06-4517 (2007).
1006   18   Falchetti, A. *et al.* Multiple endocrine neoplasia type 1 (MEN1): not only inherited
1007   endocrine tumors. *Genet. Med.* **11**, 825-835,
1008   doi:papers3://publication/doi/10.1097/GIM.0b013e3181be5c97 (2009).
1009   19   Xu, B. *et al.* Menin promotes hepatocellular carcinogenesis and epigenetically
1010   up-regulates Yap1 transcription. *Proc Natl Acad Sci USA* **110**, 17480-17485,
1011   doi:papers3://publication/doi/10.1073/pnas.1312022110 (2013).
1012   20   Krivtsov, A. V. & Armstrong, S. A. MLL translocations, histone modifications and
1013   leukaemia stem-cell development. *Nat Rev Cancer* **7**, 823-833,
1014   doi:papers3://publication/doi/10.1038/nrc2253 (2007).
1015   21   Lemos, M. C. & Thakker, R. V. Multiple endocrine neoplasia type 1 (MEN1):
1016   analysis of 1336 mutations reported in the first decade following identification of
1017   the gene. *Hum. Mutat.* **29**, 22-32,
1018   doi:papers3://publication/doi/10.1002/humu.20605 (2008).
1019   22   Borkin, D. *et al.* Pharmacologic inhibition of the Menin-MLL interaction blocks
1020   progression of MLL leukemia in vivo. *Cancer Cell* **27**, 589-602,
1021   doi:papers3://publication/doi/10.1016/j.ccell.2015.02.016 (2015).
1022   23   Dreijerink, K. M. A. *et al.* Enhancer-Mediated Oncogenic Function of the Menin
1023   Tumor Suppressor in Breast Cancer. *Cell Rep* **18**, 2359-2372,
1024   doi:papers3://publication/doi/10.1016/j.celrep.2017.02.025 (2017).
1025   24   Grembecka, J. *et al.* Menin-MLL inhibitors reverse oncogenic activity of MLL
1026   fusion proteins in leukemia. *Nat Meth* **8**, 277-284,
1027   doi:papers3://publication/doi/10.1038/nchembio.773 (2012).
1028   25   Krivtsov, A. V. *et al.* A Menin-MLL Inhibitor Induces Specific Chromatin Changes
1029   and Eradicates Disease in Models of MLL-Rearranged Leukemia. *Cancer Cell*
1030   **36**, 660-673.e611, doi:papers3://publication/doi/10.1016/j.ccell.2019.11.001
1031   (2019).
1032   26   Malik, R. *et al.* Targeting the MLL complex in castration-resistant prostate cancer.
1033   *Nature Medicine* **21**, 344-352, doi:papers3://publication/doi/10.1038/nm.3830
1034   (2015).

27   Uckelmann, H. J. *et al.* Therapeutic targeting of preleukemia cells in a mouse
model of NPM1 mutant acute myeloid leukemia. *Science* **367**, 586-590,
doi:papers3://publication/doi/10.1126/science.aax5863 (2020).
28   Kurmasheva, R. T. *et al.* Evaluation of VTP-50469, a menin-MLL1 inhibitor,
against Ewing sarcoma xenograft models by the pediatric preclinical testing
consortium. *Pediatr Blood Cancer* **67**, e28284, doi:10.1002/pbc.28284 (2020).
29   Chen, C.-W. *et al.* DOT1L inhibits SIRT1-mediated epigenetic silencing to
maintain leukemic gene expression in MLL-rearranged leukemia. *Nature*
Medicine **21**, 335-343, doi:papers3://publication/doi/10.1038/nm.3832 (2015).
30   Bernt, K. M. *et al.* MLL-rearranged leukemia is dependent on aberrant H3K79
methylation by DOT1L. *Cancer Cell* **20**, 66-78,
doi:papers3://publication/doi/10.1016/j.ccr.2011.06.010 (2011).
31   Chang, M.-J. *et al.* Histone H3 lysine 79 methyltransferase Dot1 is required for
immortalization by MLL oncogenes. *Cancer Res* **70**, 10234-10242,
doi:papers3://publication/doi/10.1158/0008-5472.CAN-10-3294 (2010).
32   Mueller, D. *et al.* Misguided transcriptional elongation causes mixed lineage
leukemia. *PLoS Biol* **7**, e1000249,
doi:papers3://publication/doi/10.1371/journal.pbio.1000249 (2009).
33   Zuber, J. *et al.* RNAi screen identifies Brd4 as a therapeutic target in acute
myeloid leukaemia. *Nature 2018* **478**, 524-528,
doi:papers3://publication/doi/10.1038/nature10334 (2011).
34   Zuber, J. *et al.* An integrated approach to dissecting oncogene addiction
implicates a Myb-coordinated self-renewal program as essential for leukemia
maintenance. *Genes & Development* **25**, 1628-1640,
doi:papers3://publication/doi/10.1101/gad.17269211 (2011).
35   Milne, T. A. *et al.* MLL targets SET domain methyltransferase activity to Hox
gene promoters. *Mol Cell* **10**, 1107-1117,
doi:papers3://publication/doi/10.1016/s1097-2765(02)00741-4 (2002).
36   Nakamura, T. *et al.* ALL-1 is a histone methyltransferase that assembles a
supercomplex of proteins involved in transcriptional regulation. *Mol Cell* **10**,
1119-1128, doi:Doi 10.1016/S1097-2765(02)00740-2 (2002).
37   Dorighi, K. M. *et al.* Mll3 and Mll4 Facilitate Enhancer RNA Synthesis and
Transcription from Promoters Independently of H3K4 Monomethylation. *Mol Cell*
66, 568-576.e564, doi:papers3://publication/doi/10.1016/j.molcel.2017.04.018
(2017).
38   Herz, H.-M. *et al.* Enhancer-associated H3K4 monomethylation by Trithorax-
related, the Drosophila homolog of mammalian Mll3/Mll4. *Genes &*
Development **26**, 2604-2620,
doi:papers3://publication/doi/10.1101/gad.201327.112 (2012).
39   Hu, D. *et al.* The MLL3/MLL4 branches of the COMPASS family function as
major histone H3K4 monomethylases at enhancers. *Mol Cell Biol* **33**, 4745-4754,
doi:papers3://publication/doi/10.1128/MCB.01181-13 (2013).
40   Lee, J. *et al.* A tumor suppressive coactivator complex of p53 containing ASC-2
and histone H3-lysine-4 methyltransferase MLL3 or its paralogue MLL4. *Proc*
Natl Acad Sci USA **106**, 8513-8518,
doi:papers3://publication/doi/10.1073/pnas.0902873106 (2009).

41   Wang, S.-P. *et al.* A UTX-MLL4-p300 Transcriptional Regulatory Network
Coordinately Shapes Active Enhancer Landscapes for Eliciting Transcription. *Mol*
Cell **67**, 308-321.e306,
doi:papers3://publication/doi/10.1016/j.molcel.2017.06.028 (2017).
42   Issaeva, I. *et al.* Knockdown of ALR (MLL2) reveals ALR target genes and leads
to alterations in cell adhesion and growth. *Mol Cell Biol* **27**, 1889-1903,
doi:papers3://publication/doi/10.1128/MCB.01506-06 (2007).
43   Santos, M. A. *et al.* DNA-damage-induced differentiation of leukaemic cells as an
anti-cancer barrier. *Nature 2018* **514**, 107-111,
doi:papers3://publication/doi/10.1038/nature13483 (2014).
44   Sun, Y. *et al.* HOXA9 Reprograms the Enhancer Landscape to Promote
Leukemogenesis. *Cancer Cell* **34**, 643-658.e645,
doi:papers3://publication/doi/10.1016/j.ccell.2018.08.018 (2018).
45   Chen, C. *et al.* MLL3 is a haploinsufficient 7q tumor suppressor in acute myeloid
leukemia. *Cancer Cell* **25**, 652-665,
doi:papers3://publication/doi/10.1016/j.ccr.2014.03.016 (2014).
46   Gažová, I., Lengeling, A. & Summers, K. M. Lysine demethylases KDM6A and
UTY: The X and Y of histone demethylation. *Molecular Genetics and Metabolism*
127, 31-44, doi:papers3://publication/doi/10.1016/j.ymgme.2019.04.012 (2019).
47   Kooistra, S. M. & Helin, K. Molecular mechanisms and potential functions of
histone demethylases. *Nat Rev Mol Cell Biol* **13**, 297-311,
doi:papers3://publication/doi/10.1038/nrm3327 (2012).
48   Meyer, C. *et al.* The MLL recombinome of acute leukemias in 2017. *Leukemia*
32, 273-284, doi:papers3://publication/doi/10.1038/leu.2017.213 (2018).
49   Thirman, M. J. *et al.* Rearrangement of the MLL gene in acute lymphoblastic and
acute myeloid leukemias with 11q23 chromosomal translocations. *N Engl J Med*
329, 909-914, doi:10.1056/NEJM199309233291302 (1993).
50   Kuhn, M. W. *et al.* Targeting Chromatin Regulators Inhibits Leukemogenic Gene
Expression in NPM1 Mutant Leukemia. *Cancer Discov* **6**, 1166-1181,
doi:10.1158/2159-8290.CD-16-0237 (2016).
51   Tiacci, E. *et al.* The NPM1 wild-type OCI-AML2 and the NPM1-mutated OCI-
AML3 cell lines carry DNMT3A mutations. *Leukemia* **26**, 554-557,
doi:papers3://publication/doi/10.1038/leu.2011.238 (2012).
52   Chen, Y. *et al.* Distinct pathways affected by menin versus MLL1/MLL2 in MLL-
rearranged acute myeloid leukemia. *Exp Hematol* **69**, 37-42,
doi:10.1016/j.exphem.2018.10.001 (2019).
53   Kumar, A. R. *et al.* A role for MEIS1 in MLL-fusion gene leukemia. *Blood* **113**,
1756-1758, doi:papers3://publication/doi/10.1182/blood-2008-06-163287 (2009).
54   Lange, B. *et al.* Growth-Factor Requirements of Childhood Acute-Leukemia -
Establishment of Gm-Csf Dependent Cell-Lines. *Blood* **70**, 192-199,
doi:papers3://publication/doi/10.1182/blood.V70.1.192.192 (1987).
55   Scacheri, P. C. *et al.* Genome-wide analysis of menin binding provides insights
into MEN1 tumorigenesis. *PLoS Genet* **2**, e51,
doi:papers3://publication/doi/10.1371/journal.pgen.0020051 (2006).

1125 56 Wang, J. K. *et al.* The histone demethylase UTX enables RB-dependent cell fate
1126   control. *Genes & Development* **24**, 327-332,
1127   doi:papers3://publication/doi/10.1101/gad.1882610 (2010).
1128 57 Cheng, J. *et al.* A role for H3K4 monomethylation in gene repression and
1129   partitioning of chromatin readers. *Mol Cell* **53**, 979-992,
1130   doi:papers3://publication/doi/10.1016/j.molcel.2014.02.032 (2014).
1131 58 Morgan, M. A. J. & Shilatifard, A. Reevaluating the roles of histone-modifying
1132   enzymes and their associated chromatin modifications in transcriptional
1133   regulation. *Nat Genet* **52**, 1271-1281, doi:10.1038/s41588-020-00736-4 (2020).
1134 59 Jang, Y. *et al.* H3.3K4M destabilizes enhancer H3K4 methyltransferases
1135   MLL3/MLL4 and impairs adipose tissue development. *Nucleic Acids Res* **47**, 607-
1136   620, doi:papers3://publication/doi/10.1093/nar/gky982 (2019).
1137 60 La, P. *et al.* Direct binding of DNA by tumor suppressor menin. *J Biol Chem* **279**,
1138   49045-49054, doi:10.1074/jbc.M409358200 (2004).
1139 61 Oldfield, A. J. *et al.* NF-Y controls fidelity of transcription initiation at gene
1140   promoters through maintenance of the nucleosome-depleted region. *Nat*
1141   *Commun* **10**, 3072, doi:10.1038/s41467-019-10905-7 (2019).
1142 62 Jenuwein, T. & Allis, C. D. Translating the Histone Code. *Science* **293**, 1074-
1143   1080, doi:doi:10.1126/science.1063127 (2001).
1144 63 Akhtar, A. & Becker, P. B. Activation of Transcription through Histone H4
1145   Acetylation by MOF, an Acetyltransferase Essential for Dosage Compensation in
1146   Drosophila. *Mol Cell* **5**, 367-375, doi:10.1016/S1097-2765(00)80431-
1147   1 (2000).
1148 64 Bone, J. R. *et al.* Acetylated histone H4 on the male X chromosome is associated
1149   with dosage compensation in Drosophila. *Genes & Development* **8**, 96-104,
1150   doi:10.1101/gad.8.1.96 (1994).
1151 65 Smith, E. R. *et al.* The drosophila MSL complex acetylates histone H4 at lysine
1152   16, a chromatin modification linked to dosage compensation. *Mol Cell Biol* **20**,
1153   312-318, doi:10.1128/mcb.20.1.312-318.2000 (2000).
1154 66 Shogren-Knaak, M. *et al.* Histone H4-K16 acetylation controls chromatin
1155   structure and protein interactions. *Science* **311**, 844-847,
1156   doi:10.1126/science.1124000 (2006).
1157 67 Taylor, G. C., Eskeland, R., Hekimoglu-Balkan, B., Pradeepa, M. M. & Bickmore,
1158   W. A. H4K16 acetylation marks active genes and enhancers of embryonic stem
1159   cells, but does not alter chromatin compaction. *Genome Res* **23**, 2053-2065,
1160   doi:10.1101/gr.155028.113 (2013).
1161 68 Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list
1162   enrichment analysis tool. *BMC Bioinformatics* **14**, 128-114,
1163   doi:papers3://publication/doi/10.1186/1471-2105-14-128 (2013).
1164 69 Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis
1165   web server 2016 update. *Nucleic Acids Res* **44**, W90-97,
1166   doi:papers3://publication/doi/10.1093/nar/gkw377 (2016).
1167 70 Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based
1168   approach for interpreting genome-wide expression profiles. *Proceedings of the*
1169   *National Academy of Sciences* **102**, 15545-15550,
1170   doi:papers3://publication/doi/10.1073/pnas.0506580102 (2005).

71   Fridman, A. L. & Tainsky, M. A. Critical pathways in cellular senescence and
immortalization revealed by gene expression profiling. *Oncogene* **27**, 5975-5987,
doi:papers3://publication/doi/10.1038/onc.2008.213 (2008).
72   Ruscetti, M. *et al.* NK cell-mediated cytotoxicity contributes to tumor control by a
cytostatic drug combination. *Science* **362**, 1416-1422,
doi:papers3://publication/doi/10.1126/science.aas9090 (2018).
73   Dang, W. *et al.* Histone H4 lysine 16 acetylation regulates cellular lifespan.
Nature 2018 **459**, 802-807, doi:10.1038/nature08085 (2009).
74   Faget, D. V., Ren, Q. & Stewart, S. A. Unmasking senescence: context-
dependent effects of SASP in cancer. *Nat Rev Cancer* **19**, 439-453,
doi:papers3://publication/doi/10.1038/s41568-019-0156-2 (2019).
75   Kuilman, T., Michaloglou, C., Mooi, W. J. & Peeper, D. S. The essence of
senescence. *Genes & Development* **24**, 2463-2479,
doi:papers3://publication/doi/10.1101/gad.1971610 (2010).
76   Tasdemir, N. *et al.* BRD4 Connects Enhancer Remodeling to Senescence
Immune Surveillance. *Cancer Discov* **6**, 612-629, doi:10.1158/2159-8290.CD-16-
0217 (2016).
77   Placke, T. *et al.* Requirement for CDK6 in MLL-rearranged acute myeloid
leukemia. *Blood* **124**, 13-23, doi:papers3://publication/doi/10.1182/blood-2014-
02-558114 (2014).
78   Hirai, H., Roussel, M. F., Kato, J. Y., Ashmun, R. A. & Sherr, C. J. Novel INK4
proteins, p19 and p18, are specific inhibitors of the cyclin D-dependent kinases
CDK4 and CDK6. *Mol Cell Biol* **15**, 2672-2681,
doi:papers3://publication/doi/10.1128/mcb.15.5.2672 (1995).
79   Fry, D. W. *et al.* Specific inhibition of cyclin-dependent kinase 4/6 by PD 0332991
and associated antitumor activity in human tumor xenografts. *Molecular Cancer*
Therapeutics **3**, 1427-1437, doi:papers3://publication/uuid/E8D6C4BF-0FC6-
4134-969C-C9812E9A7B32 (2004).
80   Ruscetti, M. *et al.* Senescence-Induced Vascular Remodeling Creates
Therapeutic Vulnerabilities in Pancreas Cancer. *Cell* **181**, 424-441.e421,
doi:papers3://publication/doi/10.1016/j.cell.2020.03.008 (2020).
81   Sherr, C. J., Beach, D. & Shapiro, G. I. Targeting CDK4 and CDK6: From
Discovery to Therapy. *Cancer Discovery* **6**, 353-367,
doi:papers3://publication/doi/10.1158/2159-8290.CD-15-0894 (2016).
82   Yang, C. *et al.* Acquired CDK6 amplification promotes breast cancer resistance
to CDK4/6 inhibitors and loss of ER signaling and dependence. *Oncogene* **36**,
2255-2264, doi:10.1038/onc.2016.379 (2017).
83   Chen, L. *et al.* Autophagy inhibition contributes to the synergistic interaction
between EGCG and doxorubicin to kill the hepatoma Hep3B cells. *PLoS ONE* **9**,
e85771, doi:papers3://publication/doi/10.1371/journal.pone.0085771 (2014).
84   Kempinska, K. *et al.* Pharmacologic Inhibition of the Menin-MLL Interaction
Leads to Transcriptional Repression of PEG10 and Blocks Hepatocellular
Carcinoma. *Molecular Cancer Therapeutics* **17**, 26-38,
doi:papers3://publication/doi/10.1158/1535-7163.MCT-17-0580 (2018).

85 Svoboda, L. K. *et al.* Menin regulates the serine biosynthetic pathway in Ewing
sarcoma. *The Journal of pathology* **245**, 324-336,
doi:papers3://publication/doi/10.1002/path.5085 (2018).
86 Agger, K. *et al.* UTX and JMJD3 are histone H3K27 demethylases involved in
HOX gene regulation and development. *Nature 2018* **449**, 731-734,
doi:10.1038/nature06145 (2007).
87 Lan, F. *et al.* A histone H3 lysine 27 demethylase regulates animal posterior
development. *Nature 2018* **449**, 689-694, doi:10.1038/nature06192 (2007).
88 Lee, M. G. *et al.* Demethylation of H3K27 Regulates Polycomb Recruitment and
H2A Ubiquitination. *Science* **318**, 447-450, doi:doi:10.1126/science.1149042
(2007).
89 Hong, S. *et al.* Identification of JmjC domain-containing UTX and JMJD3 as
histone H3 lysine 27 demethylases. *Proceedings of the National Academy of*
Sciences **104**, 18439-18444, doi:10.1073/pnas.0707292104 (2007).
90 Shi, B. *et al.* UTX condensation underlies its tumour-suppressive activity. *Nature*
2018 **597**, 726-731, doi:10.1038/s41586-021-03903-7 (2021).
91 Cao, R. *et al.* Role of Histone H3 Lysine 27 Methylation in Polycomb-Group
Silencing. *Science* **298**, 1039-1043, doi:doi:10.1126/science.1076997 (2002).
92 Klossowski, S. *et al.* Menin inhibitor MI-3454 induces remission in MLL1-
rearranged and NPM1-mutated models of leukemia. *J Clin Invest* **130**, 981-997,
doi:papers3://publication/doi/10.1172/JCI129126 (2020).
93 Qiu, H. *et al.* MEN1 deficiency leads to neuroendocrine differentiation of lung
cancer and disrupts the DNA damage response. *Nat Commun* **11**, 1009,
doi:10.1038/s41467-020-14614-4 (2020).
94 Karnik, S. K. *et al.* Menin regulates pancreatic islet growth by promoting histone
methylation and expression of genes encoding p27Kip1 and p18INK4c.
Proceedings of the National Academy of Sciences **102**, 14659-14664,
doi:papers3://publication/doi/10.1073/pnas.0503484102 (2005).
95 Schnepp, R. W. *et al.* Menin induces apoptosis in murine embryonic fibroblasts. *J*
Biol Chem **279**, 10685-10691,
doi:papers3://publication/doi/10.1074/jbc.M308073200 (2004).
96 Chen, B. *et al.* Dynamic imaging of genomic loci in living human cells by an
optimized CRISPR/Cas system. *Cell* **155**, 1479-1491,
doi:10.1016/j.cell.2013.12.001 (2013).
97 Akama-Garren, E. H. *et al.* A Modular Assembly Platform for Rapid Generation of
DNA Constructs. *Sci. Rep.* **6**, 16836-16839,
doi:papers3://publication/doi/10.1038/srep16836 (2016).
98 Krivtsov, A. V. *et al.* Transformation from committed progenitor to leukaemia
stem cell initiated by MLL-AF9. *Nature 2018* **442**, 818-822,
doi:10.1038/nature04980 (2006).
99 Krivtsov, A. V. *et al.* Patient Derived Xenograft (PDX) Models Faithfully
Recapitulate The Genetic Composition Of Primary AML. *Blood* **122**, 1328-1328,
doi:10.1182/blood.V122.21.1328.1328 (2013).
100 Doench, J. G. *et al.* Optimized sgRNA design to maximize activity and minimize
off-target effects of CRISPR-Cas9. *Nat Biotechnol* **34**, 184-191,
doi:papers3://publication/doi/10.1038/nbt.3437 (2016).

101 Sanjana, N. E., Shalem, O. & Zhang, F. Improved vectors and genome-wide
libraries for CRISPR screening. *Nat Meth* **11**, 783-784,
doi:papers3://publication/doi/10.1038/nmeth.3047 (2014).
102 Phillips, R. E. *et al.* Target identification reveals lanosterol synthase as a
vulnerability in glioma. *Proc Natl Acad Sci USA* **116**, 7957-7962,
doi:papers3://publication/doi/10.1073/pnas.1820989116 (2019).
103 Strezoska, Ž. *et al.* Optimized PCR conditions and increased shRNA fold
representation improve reproducibility of pooled shRNA screens. *PLoS ONE* **7**,
e42341, doi:papers3://publication/doi/10.1371/journal.pone.0042341 (2012).
104 Li, W. *et al.* Quality control, modeling, and visualization of CRISPR screens with
MAGeCK-VISPR. *Genome Biol* **16**, 281, doi:10.1186/s13059-015-0843-6 (2015).
105 Carroll, T. S., Liang, Z., Salama, R., Stark, R. & de Santiago, I. Impact of artifact
removal on ChIP quality metrics in ChIP-seq and ChIP-exo data. *Front. Genet.* **5**,
doi:papers3://publication/doi/10.3389/fgene.2014.00075 (2014).
106 Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster,
cheaper and better for alignment and quantification of RNA sequencing reads.
Nucleic Acids Res **47**, e47, doi:papers3://publication/doi/10.1093/nar/gkz114
(2019).
107 Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**,
R137-139, doi:papers3://publication/doi/10.1186/gb-2008-9-9-r137 (2008).
108 Nikolayeva, O. & Robinson, M. D. edgeR for differential RNA-seq and ChIP-seq
analysis: an application to stem cell biology. *Methods Mol. Biol.* **1150**, 45-79,
doi:papers3://publication/doi/10.1007/978-1-4939-0512-6_3 (2014).
109 Yu, G., Wang, L.-G. & He, Q.-Y. ChIPseeker: an R/Bioconductor package for
ChIP peak annotation, comparison and visualization. *Bioinformatics* **31**, 2382-
2383, doi:papers3://publication/doi/10.1093/bioinformatics/btv145 (2015).
110 Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a
flexible platform for exploring deep-sequencing data. *Nucleic Acids Res* **42**,
W187-191, doi:papers3://publication/doi/10.1093/nar/gku365 (2014).
111 Shen, L., Shao, N., Liu, X. & Nestler, E. ngs.plot: Quick mining and visualization
of next-generation sequencing data by integrating genomic databases. *BMC*
Genomics **15**, 284-214, doi:papers3://publication/doi/10.1186/1471-2164-15-284
(2014).
112 Zhu, L. J. *et al.* ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq
and ChIP-chip data. *BMC Bioinformatics* **11**, 10,
doi:papers3://publication/doi/10.1186/1471-2105-11-237 (2010).
113 Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification
of Problematic Regions of the Genome. *Sci. Rep.* **9**, 9354-9355,
doi:papers3://publication/doi/10.1038/s41598-019-45839-z (2019).
114 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550,
doi:papers3://publication/doi/10.1186/s13059-014-0550-8 (2014).
115 Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides
fast and bias-aware quantification of transcript expression. *Nat Meth* **14**, 417-
419, doi:papers3://publication/doi/10.1038/nmeth.4197 (2017).

116   Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq:
1307   transcript-level estimates improve gene-level inferences. *F1000Research* **4**,
1308   1521, doi:papers3://publication/doi/10.12688/f1000research.7563.2 (2015).

FIGURE 1 - CRISPR screens uncover functional interplay between the mammalian MLL1/2 and MLL3/4 chromatin modifying complexes

FIGURE 2 - MLL1-Menin complex and NF-Y restrict chromatin occupancy of MLL3/4-UTX at promoter regions

FIGURE 3 - Transcriptional co-regulation of tumor suppressive pathways by Menin and UTX

FIGURE 4 - Catalytic activity of UTX is not required to elicit UTX-dependent cellular and transcriptional functions

FIGURE 5 - *In vivo* response to Menin-MLL inhibition is accompanied by induction of MLL3/4-UTX-dependent tumor suppressive programs

FIGURE 6 - Molecular switch between the mammalian MLL complexes dictates cellular response to Menin-MLL inhibition

Extended Data Figure 1. Generation and characterization of Cas9-expressing mouse MLL-AF9 leukemia cells. (A) Dose response curve analysis of mouse MLL-AF9 leukemia (WT-MA9) cells treated with DMSO (vehicle) or MI-503 (Menin inhibitor) for 96 hours (mean±SEM, n=3 replicates). **(B)** Immunoblot analysis of Cas9-expressing mouse MLL-AF9 leukemia cells. **(C)** Distribution of Cas9-expressing MLL-AF9 leukemia cells determined by intracellular staining of Cas9 and flow cytometry. **(D)** Growth competition assay to determine enzymatic activity of Cas9 in mouse MLL-AF9 leukemia cells (mean±SEM, n=3 replicates). **(E)** CD11b cell surface expression measured by flow cytometry after DMSO (grey) or MI-503 (blue) treatment of MLL-AF9 leukemia cells for 96 hours. **(F)** Quantification of %CD11b positive cells after DMSO or MI-503 treatment for 96 hours (mean±SEM, n=3 replicates, P-value calculated by Student's t-test). **(G)** Quantification of %Annexin V positive cells after DMSO or MI-503 treatment at different time points (mean±SEM, n=3 replicates, *P*-value calculated by Student's t-test). **(H)** log$_2$-fold change of individual sgRNAs from MI-503-treated vs DMSO-treated CRISPR screens in mouse MLL-AF9 leukemia cells.

Extended Data Figure 2. Genome-wide CRISPR screening identifies the MLL3/4-UTX complex as a determinant of response to Menin-MLL inhibition. (A) Genome-wide screening data showing gene-level ranking based on differential enrichment of sgRNAs under Menin-MLL inhibitor treatment (VTP-50469) relative to vehicle (DMSO). Differential (Δ) beta-score between VTP-50469 and DMSO conditions was calculated using MaGeCK. A positive Δ beta-score denotes enrichment of specific gene-targeting sgRNAs. A negative Δ beta-score denotes depletion of specific gene-targeting sgRNAs. Red circles denote genes represented by enriched sgRNAs (genes whose inactivation promotes resistance to Menin-MLL inhibition). Blue circles denote genes represented by depleted sgRNAs (genes whose inactivation promotes sensitivity to Menin-MLL inhibition). The top 100 candidate mediators of resistance were selected for gene ontology (GO) analysis. **(B)** Top-scoring GO categories based on *P*-values obtained from GO analysis. **(C)** Percentage of genes within a given gene set that scored among the top 100 candidate mediators of resistance. **(D)** Fold-change of individual sgRNAs relative to T0 from either VTP-50469-treated or DMSO-treated cells (*P*-value calculated by Student's t-test).

Extended Data Figure 3. MLL3/4-UTX complex dictates therapeutic outcome of Menin-MLL inhibition. (A) Layout of growth competition assay to assess the effect of genetic loss of *Utx*, *Mll3*, or *Mll4* in MLL-AF9 leukemia cells in response to Menin-MLL inhibition. **(B)** Growth competition assay in mouse MLL-AF9 (WT-MA9) cells. Graph shows the relative growth of cells infected with RFP-tagged *Utx* (*Kdm6a*) sgRNAs measured by flow cytometry (mean±SEM, n=3 infection replicates). sgControl targets a non-genic region in chromosome 8 as negative control. **(C)** Growth competition assay in mouse MLL-AF9 (WT-MA9) cells. Graph shows the relative growth of cells infected with RFP-tagged *Mll3* (*Kmt2c*) sgRNAs measured by flow cytometry (mean±SEM, n=3 infection replicates). **(D)** Growth competition assay in mouse MLL-AF9 (WT-MA9) cells. Graph shows the relative growth of cells infected with RFP-tagged *Mll4* (*Kmt2d*) sgRNAs measured by flow cytometry (mean±SEM, n=3 infection replicates). **(E)** Relative percentage of sgUtx-expressing (RFP+) cells over time after transduction of MLL-AF9 cells treated with MI-503 or DMSO. **(F)** Relative percentage of sgMll3-expressing (RFP+) cells over time after transduction of MLL-AF9 cells treated with MI-503 or DMSO. **(G)** Relative percentage of sgMll4-expressing (RFP+) cells over time after transduction of MLL-AF9 cells treated with MI-503 or DMSO.

Extended Data Figure 4. *Utx* deletion rescues the lethality associated with *Men1* deletion. (A) Growth competition assay in mouse MLL-AF9 (WT-MA9) cells. Graph shows the relative growth of cells infected with RFP-tagged *Men1* sgRNAs measured by flow cytometry (mean±SEM, n=3 infection replicates). sgRenilla Luciferase (RenLuc) and sgRpa3 were negative and positive controls, respectively. **(B)** Plots from flow cytometry analysis of leukemia cells co-expressing sgMen1-RFP and sgUtx-BFP (*e12* and *e23*) 2 and 16 days post-infection. **(C)** Amplicon sequencing results of the mouse *Men1* and *Utx* loci from leukemia cells co-expressing sgMen1 and sgUtx (*e12* and *e23*). R1 and R2 represent two infection replicates for each genotype.

Extended Data Figure 5. Men1 and Utx epistasis is independent of the MLL-fusion. (A) Growth competition assay in mouse MLL-AF9 (WT-MA9) cells. Graph shows the relative growth of cells infected with RFP-tagged human (hu) *MLL1* (*e3* = exon 3, *e5* = exon 5) sgRNAs measured by flow cytometry (mean±SEM, n=3 infection replicates). **(B)** Differential fitness of mouse WT-MA9 cells is shown as the relative fitness of double positive cells (sgMLL1(hu)_*e3*-RFP + sgUtx-BFP or sgMLL1(hu)_*e3*-RFP + sgControl-BFP) to single positive cells (sgMLL1(hu)_*e3*-RFP) 16 days post-infection measured by flow cytometry (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test). **(C)** Differential fitness of mouse WT-MA9 cells is shown as the relative fitness of double positive cells (sgMLL1(hu)_*e5*-RFP + sgUtx-BFP or sgMLL1(hu)_*e5*-RFP + sgControl-BFP) to single positive cells (sgMLL1(hu)_e5-RFP) 16 days post-infection measured by flow cytometry (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test). **(D)** Growth competition assay in human non *MLL1*-r (OCI-AML3) cells. Graph shows the relative growth of cells infected with RFP-tagged sgRNAs measured by flow cytometry (mean±SEM, n=3 infection replicates). **(E)** Differential fitness of human OCI-AML3 cells is shown as the relative fitness of double positive cells to single positive cells 20 days post-infection measured by flow cytometry (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test). **(F)** Viability of mouse AML1-ETO9a leukemia cells treated with MI-503 for 96 hours. **(G)** Differential fitness of mouse AML1-ETO9a leukemia cells is shown as the relative fitness of double positive cells to single positive cells 15 days post-infection measured by flow cytometry (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test).

Extended Data Figure 6. *Meis1* expression is dispensable for proliferation of *Men1*[KO];*Utx*[KO] double-knockout MLL-AF9 leukemia cells. (A) qPCR analysis for *Meis1* expression in mouse MLL-AF9 leukemia cells treated with DMSO (vehicle) or MI-503 for 96 hours (mean±SEM, n=3 replicates, *P*-value calculated by Student's t-test). **(B)** qPCR analysis for *Meis1* expression in mouse MLL-AF9 leukemia cells co-expressing sgMen1-RFP and sgUtx-BFP, or a control sgRNA (mean±SEM, n=3 replicates, *P*-value calculated by Student's t-test). **(C)** Growth competition assay in mouse MLL-AF9 (MC35) cells. Graph shows the relative growth of cells infected with RFP-tagged sgRNAs (sgControl and sgMen1_*e3*) measured by flow cytometry (mean±SEM, n=3 infection replicates). **(D)** Differential fitness of mouse MC35 cells is shown as the relative fitness of double positive cells (sgMen1-RFP + sgUtx-BFP or sgMen1-RFP + sgControl-BFP) to single positive cells (sgMen1-RFP) 9 days post-infection measured by flow cytometry (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test). **(E)** Growth competition assay in human MLL-AF4 (MV4;11) leukemia cells. Graph shows the relative growth of cells infected with RFP-tagged sgRNAs measured by flow cytometry (mean±SEM, n=3 infection replicates). **(F)** Differential fitness of human MV4;11 leukemia cells is shown as the relative fitness of double positive cells to single positive cells 20 days post-infection measured by flow cytometry (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test).

Extended Data Figure 7. Genomic distribution of Menin and UTX is different in the context of MI-503 treatment. (A) Genomic distribution of Menin and UTX ChIP-Seq peaks from mouse leukemia cells treated with DMSO or MI-503 for 96 hours. **(B)** Upset plot showing a comparison between the genomic distribution of Menin ChIP-Seq peaks in DMSO (grey), genomic distribution of UTX ChIP-Seq peaks in MI-503 (grey), and the overlap between UTX ChIP-Seq peaks in MI-503 and Menin ChIP-Seq peaks in DMSO (blue).

Extended Data Figure 8. Menin-independent mammalian cells do not exhibit chromatin changes associated with the Menin-UTX molecular switch. (A) Immunoblot analysis of Cas9-expressing mouse fibroblast (NIH-3T3) cells expressing different controls and *Men1* sgRNAs. **(B)** Growth competition assay in mouse fibroblast cells. Graph shows the relative growth of cells infected with RFP-tagged sgRNAs measured by flow cytometry (mean±SEM, n=3 infection replicates). **(C)** Genomic distribution of Menin ChIP-Seq peaks from mouse fibroblasts (top) or mouse MLL-AF9 leukemia cells (bottom). **(D)** Venn diagram comparing the number of Menin peaks detected in mouse MLL-AF9 leukemia cells (grey) versus NIH-3T3 fibroblasts (yellow). **(E)** Heatmaps displaying UTX ChIP-Seq signals mapping to a 4-kb window around TSSs. Data is shown for mouse MLL-AF9 cells (left) and NIH-3T3 mouse fibroblasts (right) treated with DMSO or MI-503 for 96 hours. Metaplot represents the average ChIP-Seq signal for UTX at promoters that are bound by UTX (green) or not (black) in MLL-AF9 cells treated with MI-503.

Extended Data Figure 9. Disruption of Menin-MLL1 leads to targeting of MLL3/4-UTX to sites normally bound by the MLL1-Menin complex. (A) Schematic representation of the mammalian MLL1 and MLL2 histone methyltransferase complexes. Highlighted in red is the enzymatic subunit of the complex (MLL1 and MLL2 are mutually exclusive). Yellow denotes shared subunits with the MLL3/4 complex. **(B)** Schematic representation of the mammalian MLL3 and MLL4 histone methyltransferase complexes. Highlighted in pink and purple are the enzymatic subunits of the complex (MLL3 and MLL4 are mutually exclusive). Yellow denotes shared subunits with the MLL1/2 complex. Green denotes non-enzymatic subunits of the MLL3/4 complex. **(C)** Heatmaps displaying MLL1 (red) and H3K4me3 (blue) ChIP-Seq signals mapping to a 4-kb window around TSSs. Data is shown for DMSO and MI-503-treated cells for 96 hours. Metaplot represents the average ChIP-Seq signal for each protein at promoters that are UTX+ (green) or UTX- (black). Heatmaps displaying MLL3/4 (pink) and H3K4me1 (grey) ChIP-Seq signals mapping to a 4-kb window around TSSs. Data is shown for DMSO and MI-503-treated cells for 96 hours. Metaplot represents the average ChIP-Seq signal for each protein at promoters that are UTX+ (green) or UTX- (black). **(D)** Density plot showing correlation between H3K4me3 and MLL1 ChIP-Seq signals (MI-503 vs DMSO). Signals correspond to summed signal ±2kb around TSSs that overlap with UTX peaks in MI-503 condition. **(E)** Density plot showing correlation between H3K4me3 and MLL1 ChIP-Seq signals (MI-503 vs DMSO). Signals correspond to summed signal ±2kb around TSSs that do not overlap with UTX peaks in MI-503 condition. **(F)** Density plot showing correlation between H3K4me1 and MLL3/4 ChIP-Seq signals (MI-503 vs DMSO). Signals correspond to summed signal ±2kb around TSSs that overlap with UTX peaks in MI-503 condition. **(G)** Density plot showing correlation between H3K4me1 and MLL3/4 ChIP-Seq signals (MI-503 vs DMSO). Signals correspond to summed signal ±2kb around TSSs that do not overlap with UTX peaks in MI-503 condition.

Extended Data Figure 10. MLL1-Menin targets are distinct genomic loci from MLL-FP targets.
(A) Heatmaps displaying Menin (black), MLL1 (red), H3K4me3 (blue), and H3K79me2 (green) ChIP-Seq signals mapping to MLL-AF9 target loci (139 sites) from MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours. Metaplot represents the average ChIP-Seq signal for each protein at promoters. **(B)** Schematic representation of ectopic chromatin complexes that are formed by MLL-FPs. **(C)** Heatmaps displaying H3K79me2 ChIP-Seq signals mapping to a 4-kb window around TSSs in MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours. Metaplot represents the average ChIP-Seq signal for H3K79me2 at promoters that are bound by UTX in MLL-AF9 cells treated with MI-503.

Extended Data Figure 11. Destabilization of the MLL3/4-UTX complex is sufficient to blunt cellular responses to Menin-MLL inhibition. (A) Layout of growth competition assay to assess the effects of expressing a H3.1K4M mutation in MLL-AF9 leukemia cells in response to Menin-MLL inhibition. **(B)** Relative percentage of RFP+ transgene (H3.1WT or H3.1K4M) cells over time after transduction of MLL-AF9 cells treated with MI-503 or DMSO (mean±SEM, n=3 replicates, *P*-value calculated by Student's t-test, P < 0.001 = ***). **(C)** Expression of H3.1WT-RFP or H3.1K4M-RFP in mouse MLL-AF9 leukemia cells treated with DMSO or MI-503 was monitored by flow cytometry over time. **(D)** Immunoblot analysis of HA-tag from MLL-AF9 leukemia cells expressing H3.1WT-HA or H3.1K4M-HA and treated with DMSO or MI-503 for 6 days. **(E)** Immunoblot analysis of MLL4, UTX, HSP90 (loading control), H3K4me1, and total H3 (loading control) from MLL-AF9 leukemia cells expressing H3.1WT or H3.1K4M and treated with DMSO or MI-503 for 6 days.

A

Overlap Menin (DMSO) and UTX (MI-503) in MLL-AF9 leukemia

Rank	Motif	Name	P-value	% of Target	% of Background
1		**NFY**	1e⁻¹³⁰	44.97%	9.13%
2		Sp5	1e⁻⁶³	54.14%	23.85%
3		Sp1	1e⁻⁵⁰	32.69%	11.10%
4		KLF5	1e⁻⁴⁵	50.74%	25.05%
5		KLF14	1e⁻⁴¹	59.47%	33.75%

B

Menin ChIP in NIH-3T3 fibroblasts

Rank	Motif	Name	P-value	% of Target	% of Background
1		CBF1	1e⁻⁶⁹	41.25%	1.86%
2		ARG80	1e⁻⁶⁹	40.62%	1.76%
3		TBX5	1e⁻⁶⁷	43.12%	2.35%
4		TEIL	1e⁻⁶⁴	42.50%	2.45%
5		HLH-2	1e⁻⁶²	40.00%	2.12%

C

UTX ChIP in NIH-3T3 fibroblasts

Rank	Motif	Name	P-value	% of Target	% of Background
1		VND	1e⁻¹⁹	0.48%	0.00%
2		GFI1	1e⁻¹⁶	0.43%	0.00%
3		RLR1	1e⁻¹⁶	0.43%	0.00%
4		HOX2A	1e⁻¹⁶	0.43%	0.00%
5		ARID5A	1e⁻¹⁶	0.48%	0.00%

D

E

F

Extended Data Figure 12. NF-Y restricts chromatin occupancy of UTX at promoter regions. (A) Motif analysis of overlapping ChIP-Seq peaks between Menin (in DMSO) and UTX (in MI-503) in mouse MLL-AF9 leukemia cells. **(B)** Motif analysis of Menin ChIP-Seq peaks detected in NIH-3T3 mouse fibroblasts. **(C)** Motif analysis of UTX ChIP-Seq peaks detected in NIH-3T3 mouse fibroblasts. **(D)** Heatmaps displaying UTX (purple), Menin (black), and NF-YA (red) ChIP-Seq signals mapping to a 4-kb window around TSSs. Data is shown for DMSO and MI-503-treated cells for 96 hours. Metaplot represents the average ChIP-Seq signal for each protein at promoters. **(E)** Genome browser representation of ChIP-Seq normalized reads (average RPKM) for representative loci bound by Menin (black), UTX (purple), and NF-YA (red) in cells treated with DMSO or MI-503 for 96 hours. **(F)** Immunoblot analysis of NF-YA and HSP90 proteins (loading control) upon MI-503 treatment of mouse MLL-AF9 leukemia cells after 96 hours.

Extended Data Figure 13. NF-Y facilitates promoter regulation by Menin. (A) Immunoblot analysis of NF-YA and HSP90 proteins (loading control) in mouse MLL-AF9 leukemia cells transduced with a control sgRNA (sgCtrl) or three independent *Nfya*-targeting sgRNAs. Direct blue staining of histones (bottom) serves as an additional loading control. **(B)** Cell viability assay from cells treated with DMSO (vehicle, black) or MI-503 (red) after 4 days (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test). sgCtrl, control sgRNA targeting a non-genic region on chromosome 8. **(C)** Cell viability assay from cells treated with DMSO (vehicle, black) or MI-503 (red) after 6 days (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test). **(D)** Heatmaps displaying UTX ChIP-Seq signals mapping to a 4-kb window around TSSs in *Nfya*^WT (red) or *Nfya*^KO (black) MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours. Metaplot represents the average ChIP-Seq signal for each protein at promoters. **(E) - (F)** Genome browser representation of Menin ChIP-Seq normalized reads (average RPKM) for representative loci bound by Menin in *Nfya*^WT (black) or *Nfya*^KO (blue) MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours. **(G)** Co-essentiality scatter plot showing relationship between *MEN1* loss-of-function and dependency to *NFYA*, *MLL1* (*KMT2A*), *MLL3* (*KMT2C*), and *UTX* (*KDM6A*). **(H)** Co-essentiality scatter plot showing relationship between *MEN1* loss-of-function and dependency to *CDKN2C* and *CDK6*. **(I)** Co-essentiality scatter plot showing relationship between MEN1 loss-of-function and dependency to *NFYA* and *MLL1* (*KMT2A*) **(J)** Co-essentiality scatter plot showing relationship between MEN1 loss-of-function and dependency to *MLL3* (*KMT2C*) and *CDKN2C*.

Extended Data Figure 14. Transcriptional assessment of MLL-AF9 leukemia cells treated with MI-503. (A) Heatmap showing differential gene expression in mouse MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours, as measured by RNA-Seq. **(B)** Gene ontology (GO) analysis of Menin-UTX targets using RNA-Seq data from MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours. Top 10 GO categories based on statistical significance are shown.

Extended Data Figure 15. Changes in histone acetylation accompany the Menin-UTX molecular switch upon Menin-MLL1 inhibition in human leukemia cells. (A) Heatmaps displaying UTX (purple) or H4K16ac (black) ChIP-Seq signals mapping to a 4-kb window around TSSs in MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours. Metaplot represents the average ChIP-Seq signal for each protein at promoters that are bound by UTX in mouse MLL-AF9 leukemia cells treated with MI-503 for 96 hours. **(B)** Genome browser representation of ChIP-Seq normalized reads (average RPKM) for UTX (purple), H4K16ac (grey), and MOF (blue) at Cdkn2c and Cdkn2d loci from MLL-AF9 cells treated with DMSO or MI-503 for 96 hours. **(C)** Schematic representation of the mammalian NSL histone acetyltransferase complex. Highlighted in blue is the enzymatic subunit of the complex, MOF. Yellow denotes other non-enzymatic subunits within the complex. **(D)** Average chromatin occupancy of H3K27ac at transcription start sites (TSS) +/-2kb of the Menin-UTX target genes. Signals corresponding to genes that are upregulated (red) or down-regulated (black) when cells are treated with MI-503 after 96 hours. RPM, reads per million.

Extended Data Figure 16. Transcriptional assessment of Utx-deficient MLL-AF9 leukemia cells. (A) Immunoblot analysis of UTX and β-actin (loading control) from Utx^{WT} and Utx^{KO} mouse MLL-AF9 leukemia cells. **(B)** Principal component analysis (PCA) using RNA-Seq gene expression data from Utx^{WT} and Utx^{KO} cells treated with DMSO or MI-503 for 96 hours.

Extended Data Figure 17. MLL-AF9 target genes are not regulated by the Menin-UTX molecular switch. (A) Boxplot showing expression levels of MLL-AF9 target genes, which are suppressed by the MI-503 treatment after 96 hours. Expression levels are shown for Utx^{WT} (left panel) and Utx^{KO} (right panel) mouse leukemia cells. **(B)** Heatmap of Z-scores for expression of MLL-AF9 target genes from Utx^{WT} and Utx^{KO} leukemia cells treated with DMSO or MI-503 for 96 hours. **(C)** GSEA plots showing changes in regulation of myeloid cell differentiation in genes induced by MI-503 after 96 hours. FDR, false discovery rate; NES, normalized enrichment score. **(D)** Mean normalized read counts for MLL-AF9 target genes *Hoxa9* (left) and *Meis1* (right) across three RNA-Seq replicate samples from Utx^{WT} and Utx^{KO} leukemia cells treated with DMSO or MI-503 for 96 hours (mean±SEM, n=3 replicates, *P*-value calculated by Student's t-test).

Extended Data Figure 18. Gene set enrichment analysis of Menin-UTX target genes. (A) GSEA plots showing changes in regulation of autophagy genes induced by MI-503 treatment after 96 hours. FDR, false discovery rate; NES, normalized enrichment score. **(B)** GSEA plots showing changes in insulin signaling pathway genes induced by MI-503 treatment after 96 hours. **(C)** GSEA plots showing changes in neurotrophin signaling pathway genes induced by MI-503 treatment after 96 hours. **(D)** GSEA plots showing changes in endocytosis genes induced by MI-503 treatment after 96 hours. **(E)** GSEA plots showing changes in MAPK signaling pathway genes induced by MI-503 treatment after 96 hours. **(F)** GSEA plots showing changes in mitophagy genes induced by MI-503 treatment after 96 hours. **(G)** RNA Pol II ChIP-Seq signal determined for senescence and cell-cycle arrest-associated genes that are Menin-UTX targets.

Extended Data Figure 19. MLL3/4-UTX-dependent induction of SASP cytokines upon Menin-MLL inhibition. (A) Secreted levels of IL-3 in Utx^{WT} and Utx^{KO} MLL-AF9 leukemia cells treated with Menin inhibitor (MI-503) or vehicle (DMSO) after four or six days. Data are quantified as pg/mL of secreted cytokine per million cells (mean±SEM, n=3 replicates, P-value calculated by Student's t-test). **(B)** Secreted levels of IL-15 in Utx^{WT} and Utx^{KO} MLL-AF9 leukemia cells treated with Menin inhibitor (MI-503) or vehicle (DMSO) after four or six days. Data are quantified as pg/mL of secreted cytokine per million cells (mean±SEM, n=3 replicates, P-value calculated by Student's t-test). **(C)** Heatmap of cytokine array results from Utx^{WT} and Utx^{KO} MLL-AF9 leukemia cells treated with Menin inhibitor (MI-503) or vehicle (DMSO) after four or six days. Data are quantified as pg/mL of secreted cytokine per million cells and represent the mean of three biological replicates. Note that the very high levels of IFNβ-1 secretion relative to the rest of the cytokines measured affect the dynamic range of data visualization.

Extended Data Figure 20. Induction of the Menin-UTX molecular switch regulates expression of the CDK inhibitors *Cdkn2c/d*. (A) - (B) Mean normalized read counts of the cyclin-dependent kinases inhibitors, *Cdkn2c* and *Cdkn2d*, across three RNA-Seq replicate samples from *Utx*^{WT} (black) and *Utx*^{KO} (red) leukemia cells treated with DMSO or MI-503 for 96 hours. **(C) - (D)** Mean normalized read counts of the cyclin-dependent kinases, *Cdk4* and *Cdk6*, across three RNA-Seq replicate samples from *Utx*^{WT} (black) and *Utx*^{KO} (red) leukemia cells treated with DMSO or MI-503 for 96 hours.

Extended Data Figure 21. Induction of the Menin-UTX molecular switch upon Menin-MLL1 inhibition in human leukemia cells. (A) Boxplot showing expression levels of SASP genes in MOLM13 human leukemia cells treated with DMSO or MI-503 for 96 hours. **(B)** Boxplot showing expression levels of SASP genes in OCI-AML3 human leukemia cells treated with DMSO or MI-503 for 96 hours. **(C)** Heatmap of Z-scores for expression of MLL-FP target genes from MOLM13 human leukemia cells treated with DMSO or MI-503 for 96 hours. **(D)** Heatmap of Z-scores for expression of MLL-FP target genes from OCI-AML3 human leukemia cells treated with DMSO or MI-503 for 96 hours. **(E)** Genome browser representation of ChIP-Seq normalized reads (average RPKM) for Menin (black) or MLL1 (red) at *CDKN2C* and *CDKN1A* loci from MOLM13 human leukemia cells treated with DMSO or VTP-50469 for 7 days.

A

pCDH-3XHA-*Utx* cDNA-3XFlag

B

C

Extended Data Figure 22. Expression of UTX truncations in MLL-AF9 *Utx*-null leukemia cells.
(A) Representation of the lentiviral vector used to stably co-express N-terminal HA-/C-terminal-Flag-tagged *Utx* cDNAs and the fluorescent protein RFP. **(B)** Flow cytometry plots showing RFP expression of the different mouse MLL-AF9 leukemia cell lines generated with the *Utx* truncations. **(C)** Immunoblotting validation of expression of the different *Utx* truncations in mouse MLL-AF9 leukemia cells.

Extended Data Figure 23. Transcriptional characterization of the different Utx truncations in mouse MLL-AF9 leukemia cells. Boxplot showing expression levels of Menin-UTX target genes in mouse MLL-AF9 leukemia cells treated with DMSO or MI-503 for 96 hours.

Extended Data Figure 24. UTX-dependent phenotypes are not associated with global changes in H3K27 methylation. (A) Genomic distribution of H3K27me3 ChIP-Seq peaks from mouse leukemia cells treated with DMSO or MI-503 for 96 hours. **(B)** Genome browser representation of ChIP-Seq normalized reads (average RPKM) for H3K27me3 from MLL-AF9 cells treated with DMSO or MI-503 for 96 hours. **(C)** Immunoblot analysis of EZH2, β-actin (loading control), H3K27me3, and total H3 (loading control) from *Ezh2*WT and *Ezh2*KO mouse MLL-AF9 leukemia cells. **(D)** Cell viability assay from mouse *Ezh2*WT (left) and *Ezh2*KO (right) MLL-AF9 leukemia cells treated with DMSO (vehicle, black) or MI-503 (red) after 96 hours (mean±SEM, n=3 infection replicates, *P*-value calculated by Student's t-test). sgCtrl, control sgRNA targeting a non-genic region on chromosome 8.

Extended Data Figure 25. Primary human AML samples from patients participating in the Syndax Phase I Menin-MLL inhibitor trial. (A) Immunophenotyping of primary human AML cells isolated from Patient 1 (*NPM1c* mutant AML). **(B)** Heatmap representation of Z-scores from longitudinal gene expression analysis of AML cells derived from Patient 1 (*NPM1c* mutant AML) treated with SNDX-5613. **(C)** Heatmap representation of Z-scores from longitudinal gene expression analysis of AML cells derived from Patient 2 (*MLL*-rearranged AML) treated with SNDX-5613.

A

PDX Sample	*MLL*-r	*MLL3* status
DFAM68555	MLL-AF9	wild-type
MSK042D	MLL-AF4	mutant

B

PDX sample	DNA sequencing	*MLL3* (*KMT2C*) alleles present

Extended Data Figure 26. Modeling of Menin-MLL1 inhibitor response in patient-derived xenografts. (A) Characteristics of PDX samples analyzed in this study. **(B)** Targeted sequencing validation of *MLL3* status in PDX samples.

Extended Data Figure 27. Transcriptional characterization of Menin-MLL1 inhibitor response in patient-derived xenografts. (A) PCA using RNA-Seq gene expression data obtained from *MLL3*-WT (DFAM68555) and *MLL3*-Mutant (MSK042D) PDX samples treated with DMSO or VTP-50469. PCA was performed using the top 500 differentially-expressed genes upon treatment of MLL3-WT PDXs with VTP-50469. Note that *MLL3*-Mutant PDX samples are virtually overlapping independent of the condition, suggesting that the induction of the Menin-UTX molecular switch is impaired in the context of *MLL3* mutation. **(B)** GSEA showing that Menin-MLL inhibition using VTP-50469 leads to induction of cellular senescence in *MLL3*-WT PDX. **(C)** GSEA showing that Menin-MLL inhibition using VTP-50469 fails to induce cellular senescence in *MLL3*-Mutant PDX. **(D)** Distance matrix for expression of all genes from RNA-Seq data obtained from *MLL3*-Mutant (MSK042D) PDX samples treated with DMSO, VTP-50469, Palbociclib, or a combination of the two inhibitors.