1   **Title of the article**:

2   Two structurally different oomycete MAMPs induce distinctive plant immune responses

3
4   **Authors:** Mohammad Shahjahan Monjil[1,2], Hiroaki Kato[3], Kentaro Matsuda[1], Natsumi

5   Suzuki[1], Shiho Tenhiro[1], Maurizio Camagna[1], Takamasa Suzuki[4], Aiko Tanaka[1], Ryohei

6   Terauchi[3], Ikuo Sato[1], Sotaro Chiba[1], Kazuhito Kawakita[1], Makoto Ojika[1] and Daigo

7   Takemoto[1*]

8
Affiliation:

[1] Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya, Aichi, 464-

8601, Japan

[2] Department of Plant Pathology, Bangladesh Agricultural University, Mymensingh-

2202, Bangladesh

[3] Graduate School of Agriculture, Kyoto University, Muko, Kyoto, 617-0001, Japan.

[4] College of Bioscience and Biotechnology, Chubu University, Kasugai, Aichi, 478-

8501 Japan

17
18   **Corresponding author**: Daigo Takemoto

19   E-mail: dtakemo@agr.nagoya-u.ac.jp, Telephone: +81 52 789 4029

20   Fax: +81 52 789 5525

21   Graduate School of Bioagricultural Sciences, Nagoya University, Chikusa, Nagoya,

22   464-8601 Japan

23
24   **Number of figures**: 6 Figures and 27 Supplemental Figures.

25
26

Abstract

Plants recognize a variety of external signals and induce appropriate mechanisms to

increase their tolerance to biotic and abiotic stresses. Precise recognition of attacking

pathogens and induction of effective resistance mechanisms are critical functions for

plant survival. Some molecular patterns unique to a certain group of microbes (MAMPs,

microbe-associated molecular patterns) are sensed by plant cells as non-self molecules

via pattern recognition receptors. While a variety of MAMPs of bacterial and fungal

origin have been identified, reports on MAMPs of oomycete pathogens are relatively

limited. This study aimed to identify unique MAMP elicitors from the oomycete

pathogen *Phytophthora infestans*, the causal agent of potato late blight. Using reactive

oxygen species (ROS) production and phytoalexin production in potato as markers for

the purification of oomycete elicitors, we identified two structurally different groups of

elicitors, namely ceramides and diacylglycerols. *P. infestans* ceramide (Pi-Cer) elicitors

induced ROS production, while diacylglycerol (Pi-DAG) elicitors, containing

eicosapentaenoic acid (EPA) as a substructure, induced the formation of phytoalexins in

potato. Pi-Cer and Pi-DAG are also contained in the mycelia of another oomycete

pathogen *Pythium aphanidermatum,* indicating that they are MAMPs of oomycetes.

When Arabidopsis was treated with Pi-Cer and EPA, partially overlapping but different

sets of genes were induced. Furthermore, simultaneous treatment with Pi-Cer and EPA

did not have a cumulative effect on induced genes, but rather the expression of some

genes induced by EPA was attenuated by the co-treatment with Pi-Cer. These results

indicate that plants may combine the signals from simultaneously recognized MAMP

elicitors to specifically adapt the defense response to a particular pathogen.

Keywords: Ceramide, EPA, MAMPs, Oomycete, Phytoalexins, *Phytophthora infestans*,

ROS.

Introduction

During the course of their evolution, plants were always faced with the challenges of environmental microorganisms but have survived by developing physical barriers or inducible resistance strategies against pathogens. The first step of inducible plant defense against potential pathogens is the recognition of molecular patterns of microbes, referred to as MAMPs or PAMPs (microbe- or pathogen-associated molecular patterns). Plant cells recognize molecular patterns unique to a certain group of microbes as non-self molecules via pattern recognition receptors (PRRs) (Ranf, 2017). In Arabidopsis, for instance, chitin, a major component of the fungal cell wall, is recognized by a PRR complex containing LysM-RLK and co-receptor CERK1, while a conserved sequence of bacterial flagellin (flg22) and an elongation factor EF-Tu (elf18) induce plant defense via PRRs, FLS2 and EFR, respectively, in conjuncture with co-receptor BAK1 (Ranf, 2017). While a variety of MAMPs of bacterial and fungal origin, as well as the mechanisms for their recognition in plants have been studied, reports for MAMPs of oomycete pathogens are relatively limited.

Oomycete plant pathogens resemble fungal pathogens in various aspects. Besides morphological similarities, such as filamentous hyphae, they are also able to form appressoria and haustoria during infection. These similarities are however merely the result of convergent evolution, since oomycetes are phylogenetically closely related to diatoms and brown algae (Stramenopiles), and only distantly related to fungi (Opisthokonta). Therefore, the natures of oomycetes and true fungi are rather different. For example, the somatic thallus of oomycetes is diploid, while it is haploid (or dikaryotic) for true fungi, and oomycetes have mitochondria with tubular cristae

whereas true fungi have flattened ones. More importantly in terms of interaction with

plants, the composition of the cell wall and cell membrane of oomycete and fungi is

distinctive. Although it varies from species to species, the fungal cell wall mainly

contains β-1,3-glucan and chitin (β-1,4-N-acetylglucosamine) (Free, 2013), whereas the

oomycete cell wall consists mainly of cellulose (β-1,4-glucan) and β-1,3-glucan

(Mélida et al., 2013). The plasma membrane of oomycetes contains fucosterol as the

end sterol, but ergosterol is the major end sterol for fungi (Gaulin et al., 2010).

Consequently, plants may have distinct mechanisms to recognize oomycete and fungal

pathogens to activate appropriate defense reactions.

Of the various oomycete genera, *Phytophthora*, *Pythium* and *Peronospora*

/*Hyaloperonospora* (downy mildew) are particularly noteworthy, as they include many

species that are problematic plant pathogens (Kamoun et al., 2015). These plant

pathogenic oomycete genera are predicted to have evolved from a common

phytopathogenic ancestor species (Thines and Kamoun, 2010). Undoubtedly the most

infamous representative of pathogenic oomycetes is *Phytophthora infestans*, the causal

agent of potato late blight, which is responsible for the Great Famine in Ireland of the

1840s. This pathogen is an ongoing problem even today, and the total costs for control

efforts and yield losses by *P. infestans* are estimated in the range of multi-billion dollars

annually (Garelik, 2002; Fry, 2008). Problems caused by *P. infestans* are particularly

serious in developing countries where fungicides are used as the primary solution for

disease management. Although potato cultivars with introduced resistance (*R*) gene(s)

have been employed, due to selection pressure on these effectors in the pathogen

population, *R*-gene dependent resistances have been lost in many varieties (Forbes,

2012).

In this study, we purified two structurally distinctive elicitors from *P. infestans*. Using

induction of reactive oxygen species (ROS) production or phytoalexin production in

potato as markers for the purification of elicitors, we identified two structurally different

groups of elicitors, namely ceramides and diacylglycerols. *P. infestans* ceramide (Pi-

Cer) elicitors induced ROS production, while diacylglycerol (Pi-DAG) elicitors,

containing eicosapentaenoic acid (EPA) as a substructure, induced the formation of

phytoalexins in potato tubers. RNAseq analysis was performed in Arabidopsis treated

with Pi-Cer, EPA, or the mixture of both elicitors, to investigate the difference in the

activity of the two elicitors on the induction of defense genes, and the effect of

simultaneous recognition of two MAMP elicitors.

RESULTS AND DISCUSSION

Purification of *Phytophthora infestans* elicitors which induce ROS production of

potato suspension-cultured cells

Previously, we have reported that crude elicitor derived from the mycelia of *P.*

infestans extracted by methanol (Pi-MEM) can induce the resistance reactions in potato

(Monjil et al., 2015). Treatment with Pi-MEM can induce the ROS production in potato

suspension cells and leaves, and production of sesquiterpenoid phytoalexins (rishitin,

lubimin and oxylubimin) in potato tubers (Figure 1). To purify the elicitor molecules

from the mycelia of *P. infestans* and determine their structure, Pi-MEM was first

fractionated by its solubility in water and butanol. In this study, the lipophilic (butanol

soluble) fraction was further separated by column chromatography, and the elicitor

fractions were selected using their ROS inducing activity in cultured potato cells

(Supplemental Figure S1). Only fractions with clear elicitor activity were used for

further purification processes. Six fractions with significant ROS-producing activity

were further purified (Supplemental Figure S1) and the chemical structures of these

elicitors were analyzed by NMR spectroscopy (Supplemental Figures S2, 3, 5, 6, 7 and

9). To compare the structural difference between elicitors and inactive related

substances, two fractions with no significant elicitor activity were also collected, and

the structures of substances in these fractions were analyzed (Supplemental Figures S1,

4 and 8). Further structural analyses (Supplemental Figures S10-16 and Supplemental

document) showed that these eight substances could be divided into two groups,

ceramide (Cer) and ceramide-phosphoethanolamine (CerPE). We designated the four

ceramide compounds Pi-Cer A-D (_P. infestans_ Ceramide) and the remaining four

compounds Pi-CerPE A-D respectively. Except for the additional phosphoethanolamine

in Pi-CerPE, compounds that share the same alphabetical suffix are otherwise

structurally identical (Figure 2 and Supplemental Figure S17A). Pi-Cer A, B and D

were able to induce ROS production, whereas Pi-Cer C had only a marginal ROS

inducing activity (Figure 2 and Supplemental Figure S1). Similarly, Pi-CerPE A, B and

D were active, whereas Pi-CerPE C was inactive (Supplemental Figures S1 and S17).

For both Pi-Cers and Pi-CerPEs, D is the most active, followed by B and A in that order

(Figure 2 and Supplemental Figure S17).

To investigate whether the substances corresponding to Pi-Cers are contained in

another phytopathogenic oomycetes, a modified purification process was applied to

partially purify the Pi-Cers from _Pythium aphanidermatum_. _Py. aphanidermatum_ is an

oomycete plant pathogen with a wide host range, causing seedling damping-off, root

and stem rots and blights of a wide range of important crops (e.g. tomato, soybean,

cucumber and cotton). Although the composition of Pi-Cers was different from that in *P.*

infestans, Pi-Cer A, B, C and D were detected in *Py. aphanidermatum*. On the other

hand, Pi-Cers were not detected in several tested fungal plant pathogens, such as

Botrytis cinerea, Fusarium oxysporum and *Colletotrichum orbiculare* (Supplemental

Figure S18). These results suggest that the Pi-Cers are molecules specific to oomycetes.

The most evident structural difference between active Pi-Cers and Pi-Cer C is the

8*E*,10*E*-double bonds and methyl-branching at C-9 in the sphingoid base, which are

found in Pi-Cer A, B and D, but not in Pi-Cer C (Figure 2). Kato et al. (2021) showed

that that 9-methyl branching and 4*E*-double bond structure of sphingoid base are

important for its elicitor activity in Arabidopsis. Previously, cerebrosides were identified

as elicitor molecules from rice blast pathogen, *Pyricularia oryzae* (Koga et al., 1998).

Cerebroside elicitors isolated from *P. oryzae* also contain 9-methyl-branching and a 4*E*-

double bond in the sphingoid base. Given that 9-methyl-branching in the sphingoid base

is essentially not found in plants (Sperling and Heinz, 2003), this structure appears to be

a molecular pattern that can be leveraged by the plant to recognize potentially

pathogenic microorganisms. By mutagenesis of the Arabidopsis pWRKY33-LUC

marker line and screening of mutants insensitive to Pi-Cer D, Kato et al. (2021)

revealed a lectin receptor kinase is required for the recognition of Pi-Cer D (9-methyl

sphingoid base) in Arabidopsis.

168 Pi-Cer D and Pi-CerPE D are elicitors of ROS production in dicot and monocot

169 plants

Because Pi-Cer D and Pi-CerPE D had the highest elicitor activity among Pi-Cers

and Pi-CerPEs, respectively, and were also more abundantly purified than the A and B

type substances (Supplemental Figure S1), Pi-Cer D and Pi-CerPE D were mainly used

in the following experiments. In potato suspension-cultured cells, ROS production

induced by Pi-MEM peaks approx. 3 h after treatment. The same pattern was observed

after Pi-Cer D and Pi-CerPE D treatment (Supplemental Figure S19), suggesting that

Pi-Cers and Pi-CerPEs could be the major ROS inducing substances in Pi-MEM. Potato

leaves were treated with Pi-Cer D, Pi-CerPE D or Pi-MEM by syringe infiltration and

induction of ROS production was detected. ROS production was induced by these

elicitors with a peak at approx. 12 h after the treatment. Similarly, ROS production was

detected in leaves of *A. thaliana* and rice treated with Pi-Cer D or Pi-CerPE D 12 h after

the treatment (Figure 3). These results indicated that Pi-Cer D and Pi-CerPE D are

recognized by both dicot (potato, Arabidopsis) and monocot (rice) plants as MAMPs.

184 Treatment of Pi-Cer D and Pi-CerPE D enhances the resistance of potato against *P.*

185 *infestans*

The effect of pre-treatment with ceramide elicitors on the resistance of potato against

P. infestans was investigated (Figure 4). Potato leaves were treated with 10 µg/ml Pi-

Cer D, Pi-CerPE D or 100 µg/ml Pi-MEM, and inoculated with a zoospore suspension

of *P. infestans* at 24 h after the elicitor treatment. Within 3 days after the inoculation,

leaves of the control plant showed water-soaked disease symptoms, and the lesions on

most leaves extended over the entire leaves within 7 days. In contrast, fewer and smaller

spots were developed in leaves treated with Pi-Cer D, Pi-CerPE D or Pi-MEM.

Evaluation of disease symptoms up to 7 days after the inoculation indicated that

pretreatment with Pi-Cer D and Pi-CerPE D can enhance the resistance of potato leaves

against *P. infestans* (Figure 4). Inoculated leaves were stained with aniline blue to

visualize the penetration of *P. infestans* as fluorescent spots of deposited callose beneath

the penetration sites at 24 h after the inoculation. Compared with control leaves, fewer

fluorescence spots were detected in leaves pretreated with Pi-Cer D, Pi-CerPE D or Pi-

MEM elicitors (Figure 4D), which indicates that Pi-Cer D and Pi-CerPE D can enhance

the penetration resistance of potato leaves against *P. infestans.*

Isolation of Pi-DAGs as elicitors for the induction of phytoalexins production in

potato tuber

In potato tubers, accumulation of phytoalexins (rishitin, lubimin and oxylubimin) was

induced by the treatment with the Pi-MEM (Figure 1). However, treatment of Pi-Cers or

Pi-CerPEs did not induce the production of phytoalexins in potato tubers (Supplemental

Figure S20), which suggested that Pi-MEM includes one or more additional elicitors

that can induce phytoalexin production. The initial step of our purification protocol

consisted of fractioning the butanol-soluble compounds via silica gel flash column

chromatography, yielding eight fractions. Of these eight fractions, the third (RM-I-130-

3) and sixth (RM-I-130-6) fraction had shown ROS inducing activity and were used for

the isolation of Pi-Cers and Pi-CerPEs (Supplemental Figures S1 and 21). The same

eight fractions were now used to evaluate their ability to induce phytoalexin production

214 in potato tubers. Substantial elicitor activity was only observed in the first fraction

215 (RM-I-130-1) and it was therefore further purified (Supplemental Figure S21). Among

216 the five fractions obtained from the following flash column chromatography, the third

217 (ST-I-8-3) and fourth (ST-I-8-4) fraction showed elicitor activity that resulted in

218 phytoalexin production. These fractions were later found to contain mainly 1,3-DAG

219 (diacylglycerol) and 1,2-DAG, respectively. Both ST-I-8-3 and -4 were further

220 separated by HPLC using an ODS column and at least 5 out of 9 resultant fractions

221 showed elicitor activity (Supplemental Figure S21). These results indicated that several

222 1,3- and 1,2-DAGs derived from *P. infestans* are elicitors that can induce the

223 phytoalexin production of potato tubers. Two fractions with relatively higher elicitor

224 activity, yield and purity, ST-I-14-4 and ST-I-23-4, were selected for further structural

225 analyses. Two additional fractions, which showed almost no elicitor activity (ST-I-14-8

226 and ST-I-23-8), were also subjected for structural analyses in order to narrow down the

227 molecular patterns that may be critical for the elicitor activity. Further structural

228 analyses (Supplemental Figures S22-25) revealed that active substances from ST-I-14-4

229 and ST-I-23-4 were 1,2- and 1,3-DAG which both contained an eicosapentaenoic acid

230 (EPA) and linoleic acid fatty acid chain each, and were designated as Pi-DAG A and B,

231 respectively (Figure 5 and Supplemental Figure S21). The inactive substances in ST-I-

232 14-8 and ST-I-23-8 were found to be 1,2- and 1,3-DAG containing palmitic acid and

233 linoleic acid as fatty acid chains, and were designated as Pi-DAG C and D, respectively.

234 Pi-DAG A and B, as well as Pi-DAG C and D, are also contained in *Py.*

235 *aphanidermatum*, suggesting that Pi-DAGs are molecules commonly found in oomycete

236 plant pathogens (Supplemental Figure S26).

Based on the structural difference of active and inactive Pi-DAGs (Figure 5A and B),

EPA was expected to be the essential structure for the elicitor activity of Pi-DAG A and

B. Consistently, EPA from *P. infestans* has previously been identified as an elicitor

molecule (Bostock et al., 1981). EPA can be considered as a MAMP of oomycete, as

EPA is not found in higher plants and higher fungi, but an essential component of the

plasma membrane of oomycete cells (Bostock et al., 2011). We confirmed that EPA, but

not Pi-Cer D, can induce the production of phytoalexins in potato tubers (Figure 5C and

D). On the contrary, Pi-Cer D, but not EPA, induced ROS production in potato leaves

(Supplemental Figure S27), thus indicating that the structurally different oomycete

MAMPs, EPA and Pi-Cer D, induce distinctive plant immune responses.

RNAseq analysis of Arabidopsis treated with EPA, Pi-Cer D or a mixture of EPA

and Pi-Cer D

To compare the elicitor activity of EPA and Pi-Cer D in Arabidopsis, an Arabidopsis

transformant containing a *LUC* transgene under the control of *WRKY33* (AT2G38470)

promoter (pWRKY33-LUC) (Kato et al., 2021) was treated with EPA and Pi-Cer D.

While Pi-Cer D can induce the activation of the *WRKY33* promoter, only marginal

induction of *WRKY33* promoter was detected by the treatment with EPA (Figure 6A).

RNAseq analysis was performed for Arabidopsis seedlings 24 h after the treatment with

20 µg/ml EPA, Pi-Cer D or a mixture of 20 µg/ml EPA and 20 µg/ml Pi-Cer D.

Clustering analysis of significantly up- or down-regulated genes revealed that treatment

with EPA or Pi-Cer D leads to distinct gene expression patterns. While some genes

respond similarly to either treatment, others genes are induced exclusively by only one

of the two treatments (Figure 6B). Furthermore, simultaneous treatment with Pi-Cer D

and EPA did not result in an expression pattern that would be expected by the mere

combination of either treatment. Instead, the expression of some genes induced by EPA

was attenuated by the co-treatment with Pi-Cer D, resulting in an expression pattern

different from either single treatment (Figure 6B). These results indicate that plants may

combine the signals from simultaneously recognized MAMP elicitors to specifically

adapt the defense response to a particular pathogen. Insight into how these elicitor

signals are processed, as well as how the response pathways of both elicitors affect one

another, will require further analysis. RNAseq analyses of potato and Arabidopsis

treated either with single, or simultaneous treatments of EPA and Pi-Cer D at different

time points should allow a better understanding of how these elicitors function.

MATERIALS AND METHODS

Biological materials, growth conditions and inoculation

Potato suspension-cultured cells were prepared from calluses induced from potato

tuber discs (cv. Sayaka carrying the *R1* and *R3* genes) in callus-inducing medium [1 ml/l

B5-vitamin (100 g/l Myo-inositol, 2.5 g/l glycine, 500 mg/l thiamine-HCl, 500 mg/l

pyridoxine-HCl, 5 g/l nicotinic acid, 50 mg/l biotin, 50 mg/l folic acid), 2 mg/l NAA,

0.5 mg/l kinetin, 3% sucrose, 0.2% phytagel and Murashige-Skoog basal medium],

under dark condition. Suspension-cultured potato cells were grown by agitation at 130

rpm, 23°C in 95 ml of MS medium supplemented with 30 mg/ml sucrose, 1 mg/l

thiamine, 100 mg/l myo-inositol, 200 mg/l KH_2PO_4, and 0.2 mg/l 2,4-

dichlorophenoxyacetic acid. Cells were sub-cultured every week and used for

experiments 4-5 days after the subculturing. Potato, rice and Arabidopsis (Col-0) plants

were grown at 23-25˚C under a 16 h photoperiod and an 8 h dark period in an

environmentally controlled growth room. Tubers of potato (cv. Rishiri) carrying the *R1*

gene were stored at 4˚C until use. *P. infestans* isolate 08YD1 and PI1234-1 (Shibata et

al., 2011) was maintained on rye-media at 10˚C. Inoculation of potato leaves with a

zoospore suspension of *P. infestans* was performed as described previously (Shibata et

al., 2010). *Pythium aphanidermatum* strain Pyaph isolated from cucumber, *Botrytis*

cinerea strain AI18 isolated from strawberry leaves in Mie Prefecture in 2018,

Fusarium oxysporum f. sp. *melonis* strain Mel02010 (Namiki et al., 1998) and

Colletotrichum orbiculare strain 104-T (Ishida and Akai 1969) were grown on potato

dextrose agar (PDA) at 25˚C and maintained at -80˚C in 20% glycerol.

Preparation of methanol extract of *Phytophthora infestans* mycelia (Pi-MEM)

The *Phytophthora infestans* isolates PI1234-1 were grown on rye-seed extract agar

medium in test tubes at 18°C in dark condition for 2 weeks. Parts of the growing

mycelia were placed in 100 ml flasks containing 20 ml of rye liquid nutrient medium of

rye seed-extract (60 g rye seed), 20 g of sucrose and 2 g of yeast extract per 1 liter and

incubated in the dark for 2 weeks at 18°C to allow the growth of the mycelia. The

mycelial mats grown in the liquid medium were washed thoroughly with water. To

remove excess water, water was filtered through filter paper (Toyoroshi No. 2) under

reduced pressure and the tissue then frozen at -20°C. The collected mycelia of *P.*

infestans were ground in liquid nitrogen by mortar and pestle. The ground mycelia were

transferred to a 50 ml tube containing methanol (10 ml/g mycelia). The mycelia

suspension was finely grounded using a polytron type homogenizer (HG30, Hitachi

Koki, Japan) for 2 min. After centrifugation at 4°C, 3000 × g for 30 min, the supernatant

was collected and dried using an evaporator and used as Pi-MEM elicitor. Further

procedures for the purification of Pi-Cer and Pi-DAG from Pi-MEM are described in

the supplemental document. Mycelia of *Pythium aphanidermatum*, *Botrytis cinerea*,

Fusarium oxysporum f. sp. *melonis* and *Colletotrichum orbiculare* were grown in 50 ml

of potato dextrose broth (PDB) in 100 ml flasks at 25°C with gentle shaking.

Measurement of ROS production in potato suspension-cultured cells

The relative intensity of ROS generation in potato suspension-cultured cells was

measured by counting photons from L-012-mediated chemiluminescence. The potato

suspension-cultured cells (50 mg/ml) were washed with the assay buffer (175 mM

mannitol, 50 mM MES-KOH, 0.5 mM $CaCl_2$ and 0.5 mM K_2SO_4, pH 5.7) twice for

removal of the liquid culture medium. For the detection of ROS produced in cultured

cells, the cells were resuspended in assay buffer, equilibrated for 1 h at 100 rpm at 23°C.

Cells were then treated with elicitors and incubated under the same condition for 3 h.

After the incubation, chemiluminescence was measured using 20 mM L-012 (Wako

Pure Chemical, Osaka, Japan) in a multimode microplate reader Mithras LB940

(Berthold Technologies, Bad Wildbad, Germany).

Measurement of ROS production in plant leaves

The relative intensity of ROS production was determined by counting photons from

L-012-mediated chemiluminescence. To detect the ROS production in potato,

Arabidopsis and rice leaves, 0.5 mM L-012 in 10 mM MOPS-KOH (pH 7.4) was

allowed to infiltrate to the intercellular space of leaves. For potato and Arabidopsis, a

syringe without a needle was used for leaf infiltration, whereas vacuum infiltration was

used for rice. Chemiluminescence was monitored using a photon image processor

equipped with a sensitive CCD camera in the dark chamber at 20°C (Aquacosmos 2.5;

Hamamatsu Photonics, Shizuoka, Japan) and quantified using the U7501 program

(Hamamatsu Photonics). For the data shown in Supplemental Fig. S27,

chemiluminescence was monitored using Lumino Graph II EM (ATTO, Tokyo, Japan).

Detection of potato phytoalexins by thin-layer chromatography (TLC)

Potato phytoalexins, exuded from potato tissue, were extracted with ethyl acetate as

described previously (Noritake et al., 1996). The extract was separated on TLC plates

(TLC aluminum sheet of silica gel 60, Merck, Whitehouse Station, NJ, USA), which

were developed with cyclohexane:ethyl acetate (1:1, v/v) and visualized by spraying

with sulfuric acid containing 0.5% vanillin followed by heating at 120°C.

Detection of potato phytoalexins by liquid chromatography/mass spectrometry

(LC/MS)

Potato tuber discs (2 cm in diameter and approx. 4 mm thick) were prepared and

incubated in a humidified chamber in dark at 23°C for 24 h before treatment with

elicitors. The upper side of incubated potato tubers was then treated with 100 µl of

elicitor solution and further incubated at 23°C in the dark for 48 h before the extraction

of phytoalexins. Four tuber disks/sample were immersed in 5 ml of ethyl acetate and

shaken for 1 h, then the organic solvent was collected and evaporated. The residual

material was redissolved in 100 µl of 50% (v/v) acetonitrile and 2 µl was injected for

the analysis with LC/MS (Accurate-Mass Q-TOF LC/MS 6520, Agilent Technologies,

Santa Clara, CA, USA).

Measuring the activation of *AtWRKY33* promoter and RNA-seq analysis

Seeds of Arabidopsis containing *Luciferase* marker gene under the control of

AtWRKY33 (AT2G38470) promoter (Kato et al., 2021) were surface sterilized with 3%

hydrogen peroxide and 50 % ethanol for 1 min with gentle shaking, washed with

sterilized water, and then individual sterilized seeds were placed in separate wells of a

96 microwell plate containing 150 µl of Murashige and Skoog (MS) liquid medium (1/2

MS salts, 0.05% [w/v] MES, 0.5% [w/v] Sucrose, adjusted to pH 5.8 with NaOH) with

50 µM D-Luciferin potassium salt (Biosynth Carbosynth, Compton, UK), covered with

a clear plastic cover. Plates containing Arabidopsis seeds were then incubated in a

growth chamber for approx. 12 days at 23ºC with 24 h light. After the treatment of

elicitors, chemiluminescence intensity derived from expressed luciferase was measured

using Mithras LB940 (Berthold) for 12 h. Total RNA was extracted from 6

seedlings/sample of 10-days old Arabidopsis 24 h after the treatment with elicitors using

the RNeasy Plant Mini Kit (QIAGEN, Hilden, Germany). Libraries were constructed

using KAPA mRNA Capture Kit (Roche, Basel, Switzerland) and MGIEasy RNA

Directional Library Prep Set (MGI, Shenzhen, China), and sequenced on DNBSEQ-

G400RS (MGI) with 150 bp paired-end protocol. The RNA-seq reads were filtered using

trim-galore v.0.6.6 (Martin, 2011, bioinformatics.babraham.ac.uk) and mapped to the

Arabidopsis genome (TAIR 10.1; RefSeq: GCF_000001735.4) using HISAT2 v.2.2.1

(Kim et al., 2019) and assembled via StringTie v.2.1.7 (Kovaka et al., 2019). Significant

differential expression was determined using DESeq2 v.1.32.0 (Love et al., 2014). All

software used during RNA-seq analysis was run with default settings. The expression

profile was calculated from the log2-fold expressions using the clustermap function

from seaborn v. 0.11.1 (Waskom, 2021).

ACKNOWLEDGMENTS

We thank Ms. Kayo Shirai (Hokkaido Central Agricultural Experiment Station, Japan)

and Dr. Seishi Akino (Hokkaido University, Japan) for providing *P. infestans* isolate, Dr.

Kenji Asano and Mr. Seiji Tamiya (National Agricultural Research Center for Hokkaido

Region, Japan) and Mr. Yasuki Tahara (Nagoya University, Japan) for providing tubers

of potato cultivars. We also thank Dr. Masaharu Kubota (National Agriculture and Food

Research Organization, Japan) for providing *Pythium aphanidermatum* strain, Prof.

Takashi Tsuge (Chubu University, Japan) for providing *Fusarium oxysporum* strain and

Prof. Yoshitaka Takano (Kyoto University, Japan) for providing *Colletotrichum*

orbiculare strain. This work was supported by a Grant-in-Aid for Scientific Research

(B) (17H03771 and 20H02985) to DT and (17H03963) to KK from the Japan Society

for the Promotion of Science.

SUPPLEMENTAL DATA

Supplemental Figure S1. Purification procedure of *Phytophthora infestans* ceramide

(Pi-Cer) and ceramide phosphoethanolamine (Pi-CerPE) elicitors.

Supplemental Figure S2. NMR spectra of Pi-Cer A (CDCl$_3$, 400 MHz for ^{1}H, 100
MHz for ^{13}C).

Supplemental Figure S3. NMR spectra of Pi-Cer B (CDCl$_3$, 600 MHz for ^{1}H, 150
MHz for ^{13}C).

Supplemental Figure S4. NMR spectra of Pi-Cer C (CDCl$_3$, 400 MHz for ^{1}H, 100
MHz for ^{13}C).

Supplemental Figure S5. NMR spectra of Pi-Cer D (CDCl$_3$, 600 MHz for ^{1}H, 150
MHz for ^{13}C).

Supplemental Figure S6. NMR spectra of Pi-CerPE A (CDCl$_3$-CD$_3$OD 4:1, 400 MHz
for ^{1}H, 100 MHz for ^{13}C).

Supplemental Figure S7. NMR spectra of Pi-CerPE B (CDCl$_3$-CD$_3$OD 4:1, 600 MHz
for ^{1}H, 150 MHz for ^{13}C).

Supplemental Figure S8. NMR spectra of Pi-CerPE D (CDCl$_3$-CD$_3$OD 4:1, 600 MHz
for ^{1}H, 150 MHz for ^{13}C).

Supplemental Figure S9. NMR spectra of Pi-CerPE D (CDCl$_3$-CD$_3$OD 4:1, 600 MHz
for ^{1}H, 150 MHz for ^{13}C).

Supplemental Figure S10. MS/MS fragmentation of Pi-Cer A (precursor ion: [M +
Na]$^+$).

Supplemental Figure S11. Two-dimensional NMR correlations of Pi-Cer B (thick
bonds: DQF-COSY, curved arrows: HMBC, l + m =15).

Supplemental Figure S12. MS/MS fragmentation of Pi-Cer B (precursor ion: [M +
Na]$^+$).

Supplemental Figure S13. Two-dimensional NMR correlations of Pi-Cer B (thick
bonds: DQF-COSY, curved arrows: HMBC, l + m + n = 19).

Supplemental Figure S14. DQF-COSY (thick bonds), HMBC (curved arrows), and
NOESY (dashed curves) correlations of Pi-Cer D.

Supplemental Figure S15. Linked-scan FAB MS/MS (negative ion mode) of the fatty
acid derived from Pi-Cer D by acid hydrolysis.

Supplemental Figure S16. DQF-COSY (thick bonds) and HMBC (curved arrows)
correlations of Pi-CerPE D.

Supplemental Figure S17. Ceramide phosphoethanolamine (Pi-CerPE) elicitors
purified from methanol extract of *P. infestans* mycelia (Pi-MEM), which can induce the
production of reactive oxygen species (ROS) in potato suspension-cultured cells.

Supplemental Figure S18. Pi-Cer contents in mycelia of oomycete and fungal plant
pathogens.

Supplemental Figure S19. Induction of reactive oxygen species (ROS) production by
Pi-Cer D and Pi-CerPE D treatment in potato suspension-culture cells.

Supplemental Figure S20. Production of phytoalexins in potato tuber is not induced by
Pi-Cers and Pi-CerPEs.

Supplemental Figure S21. Purification procedure of *Phytophthora infestans*
diacylglycerol (Pi-DAG) elicitors.

Supplemental Figure S22. ESI-TOF MS of Pi-DAG A. NMR spectra of Pi-DAG A
(CDCl$_3$, 400 MHz for ^{1}H and 100 MHz for ^{13}C). Two-dimensional NMR of Pi-DAG A
(CDCl$_3$, 400 MHz). Determination of fatty acids of Pi-DAG A by MS/MS.

Supplemental Figure S23. ^{1}H NMR spectrum of Pi-DAG B (CDCl$_3$, 400 MHz).
Determination of fatty acids in Pi-DAG B by negative ion FAB MS

Supplemental Figure S24. NMR spectra of Pi-DAG C (CDCl$_3$, 400 MHz for ^{1}H, 100
MHz for ^{13}C). Determination of fatty acids in Pi-DAG C by positive ion ESI MS/MS.
Confirmation of fatty acid linkage in Pi-DAG C.

Supplemental Figure S25. NMR spectra of Pi-DAG D (MeOD, 400 MHz for ^{1}H, 100
MHz for ^{13}C). Determination of fatty acids in Pi-DAG D by hydrolysis followed by
negative ion FAB MS and MS/MS analysis of two fatty acid products.

Supplemental Figure S26. Pi-DAG contents in mycelia of oomycete plant pathogens.

Supplemental Figure S27. Eicosapentaenoic acid (EPA) does not induce the
production of reactive oxygen species (ROS) in potato leaves.

LITERATURE CITED

Bostock RM, Kuć JA, Laine RA (1981). Eicosapentaenoic and arachidonic acids from

Phytophthora infestans elicit fungitoxic sesquiterpenes in the potato. Science **212:**

67–69

Bostock RM, Savchenko T, Lazarus C, Dehesh K (2011) Eicosapolyenoic acids:

novel MAMPs with reciprocal effect on oomycete-plant defense signaling networks.

Plant Signal Behav **6:** 531–533

Camagna M, Ojika M, Takemoto D (2020) Detoxification of the solanaceous

phytoalexins rishitin, lubimin, oxylubimin and solavetivone via a cytochrome P450

oxygenase. Plant Signal Behav **15:** 1707348

Forbes GA (2012) Using host resistance to manage potato late blight with particular

reference to developing countries. Potato Res **55:** 205–216

Free SJ (2013) Fungal cell wall organization and biosynthesis. Adv Genet **81:** 33–82

Fry W (2008) *Phytophthora infestans*: the plant (and *R* gene) destroyer. Mol Plant

Pathol **9:** 385–402

Garelik G (2002) Agriculture. Taking the bite out of potato blight. Science **298:**1702–

1704

Gaulin E, Bottin A, Dumas B (2010) Sterol biosynthesis in oomycete pathogens. Plant

Signal Behav 5:258–260

Ishida N, Akai S (1969) Relation of temperature to germination of conidia and

appressorium formation in *Colletotrichum lagenarium*. Mycologia **61:** 382–386

Kamoun S, Furzer O, Jones JD, Judelson HS, Ali GS, Dalio RJ, Roy SG, Schena L,

Zambounis A, Panabières F, Cahill D, Ruocco M, Figueiredo A, Chen XR,

Hulvey J, Stam R, Lamour K, Gijzen M, Tyler BM, Grünwald NJ, Mukhtar MS,

Tomé DF, Tör M, Van Den Ackerveken G, McDowell J, Daayf F, Fry WE,

Lindqvist-Kreuze H, Meijer HJ, Petre B, Ristaino J, Yoshida K, Birch PR,

Govers F (2015) The top 10 oomycete pathogens in molecular plant pathology. Mol

Plant Pathol **16:** 413–434

Kato H, Nemoto K, Shimizu M, Abe A, Asai S, Ishihama N, Daimon T, Ojika M,

Kawakita K, Onai K, Shirasu K, Ishiura M, Takemoto D, Takano Y, Terauchi R

(2021) Pathogen-derived 9-methyl sphingoid base is perceived by a lectin receptor

kinase in Arabidopsis. bioRxiv.

Koga J, Yamauchi T, Shimura M, Ogawa N, Oshima K, Umemura K, Kikuchi M,

Ogasawara N (1998) Cerebrosides A and C, sphingolipid elicitors of hypersensitive

cell death and phytoalexin accumulation in rice plants. J Biol Chem **273:** 31985–

31991

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol **37:**

907–915

Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M (2019)

Transcriptome assembly from long-read RNA-seq alignments with StringTie2.

Genome Biol **20:** 278

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol **15:** 550

Martin M (2011) Cutadapt removes adapter sequences from high-throughput

sequencing reads. EMBnet J **17:** 10–12

Mélida H, Sandoval-Sierra JV, Diéguez-Uribeondo J, Bulone V (2013) Analyses of

extracellular carbohydrates in oomycetes unveil the existence of three different cell

wall types. Eukaryot Cell **12:** 194–203

Monjil MS, Nozawa T, Shibata Y, Takemoto D, Ojika M, Kawakita K (2015)

Methanol extract of mycelia from *Phytophthora infestans*-induced resistance in

potato. C R Biol **338:** 185–196

Namiki F, Shiomi T, Nishi K, Kayamura T, Tsuge T (1998) Pathogenic and genetic

variation in the Japanese strains of *Fusarium oxysporum* f. sp. *melonis*.

Phytopathology **88:** 804–810

Noritake T, Kawakita K, Doke N (1996) Nitric oxide induces phytoalexin

accumulation in potato tuber tissues. Plant Cell Physiol **37:** 113–116

Ranf S (2017) Sensing of molecular patterns through cell surface immune receptors.

Curr Opin Plant Biol **38:** 68–77

Shibata Y, Kawakita K, Takemoto D (2010) Age-related resistance of *Nicotiana*

benthamiana against hemibiotrophic pathogen *Phytophthora infestans* requires both

ethylene- and salicylic acid-mediated signaling pathways. Mol Plant-Microbe

Interact **23:** 1130–1142

Shibata Y, Kawakita K, Takemoto D (2011) SGT1 and HSP90 are essential for age-

related non-host resistance of *Nicotiana benthamiana* against the oomycete pathogen

Phytophthora infestans. Physiol Mol Plant Pathol 75:120–128

Sperling P, Heinz E (2003) Plant sphingolipids: structural diversity, biosynthesis, first

genes and functions. Biochim Biophys Acta **1632:** 1–15

Thines M, Kamoun S (2010) Oomycete-plant coevolution: recent advances and future

prospects. Curr Opin Plant Biol **13:** 427–433

Waskom ML (2021) Seaborn: statistical data visualization. J. Open Res. Softw **6:** 3021

Figure 1. Elicitor activity of methanol extract of *Phytophthora infestans* mycelia (Pi-MEM) in potato.
(A) Potato suspension cultured cells were treated with 0.3% DMSO (Cont.) or 30 µg/ml Pi-MEM and production of reactive oxygen species (ROS) was detected as L-012-mediated chemiluminescence 3 h after the treatment. Data are means ± SE (n = 3). **(B)** Potato leaves were treated with 1 % DMSO (Cont.) or 100 µg/ml Pi-MEM and production of ROS was detected as L-012-mediated chemiluminescence. Data are means ± SE (n = 3). Data marked with asterisks are significantly different from control as assessed by two-tailed Student's *t* tests: **$P < 0.01$.
(C) Left, putative biosynthetic pathway of potato phytoalexins, lubimin, oxylubimin and rishitin. HPS, *Hyoscyamus muticus* premnaspirodiene synthase; HPO *H. muticus* premnaspirodiene oxygenase; SPH, sesquiterpenoid phytoalexins hydroxylase (Camagna et al., 2020). Potato tubers were treated with 3 % DMSO (Cont.) or 1 mg/ml Pi-MEM and phytoalexins were extracted 48 h after treatment. Produced phytoalexins were detected by thin-layer chromatography (middle) or LC/MS (right).

Figure 2. *Phytophthora infestans* ceramide elicitors (Pi-Cer) can induce the production of reactive oxygen species (ROS) in potato suspension cultured cells. **(A)** Potato suspension cultured cells were treated with 0.3% DMSO (Cont.), 30 µg/ml methanol extract of *P. infestans* mycelia (PiMEM), 3 µg/ml Pi-Cer A, B, C or D and production of ROS was detected as L-012 mediated chemiluminescence 3 h after the treatment. **(B)** Structures of Pi-Cer A, B, C and D. See Supplemental Figure S1 for the procedures of purification of elicitors and Supplemental Figures S2-5, S10-15 and supplemental document for details of their structural analysis.

Figure 3. Pi-Cer D and Pi-CerPE D can induce the production of reactive oxygen species (ROS) in different plant species.

(A) (Left) Potato leaves were treated with 1% DMSO (Cont.), 100 µg/ml methanol extract of *P. infestans* mycelia (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by syringe infiltration, and production of ROS was detected as L-012 mediated chemiluminescence 3-15 h after the treatment. Data are means ± SE (n = 3).

(B) (Left) *Arabidopsis thaliana* leaves were treated with 1% DMSO (Cont.), 100 µg/ml methanol extract of *P. infestans* (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by syringe infiltration, and production of ROS was detected as L-012 mediated chemiluminescence 12 h after the treatment. Data are means ± SE (n = 3).

(C) (Left) Rice leaves were treated with 1% DMSO (Cont.), 100 µg/ml methanol extract of *P. infestans* (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by vacuum infiltration, and production of ROS was detected as L-012 mediated chemiluminescence 12 h after the treatment. Data are means ± SE (n = 3). Data marked with asterisks are significantly different from control as assessed by two-tailed Student's *t* tests: **P < 0.01.

Figure 4. Pretreatment with Pi-Cer D and Pi-CerPE D enhances the resistance of potato leaves against *P. infestans*.

Potato leaves were treated with 0.5 % DMSO (control), 100 µg/ml Pi-MEM, 10 µg/ml Pi-Cer D or Pi-CerPE D and incubated for 24 hrs, and inoculated with a spore suspension of *P. infestans*. **(A)** Disease symptoms of *P. infestans* on potato leaves treated with DMSO, Pi-MEM, Pi-Cer D or Pi-CerPE D. Photographs were taken 6 days post inoculation. **(B)** Leaves representative of the disease severities used for classification. **(C)** Plots showing percentages of potato leaves with disease symptom severities using the classification depicted in (B). Leaves were pretreated with DMSO or elicitors, and disease severity of a subsequent *P. infestans* inoculation was observed from 1 - 7 days post inoculation (dpi). (n = 8). Data marked with asterisks are significantly different from control as assessed by one-tailed Mann–Whitney U-tests: *P < 0.05. **(D)** Left, Penetration sites of *P. infestans* in elicitor treated leaf-discs were detected as callose depositions by aniline blue staining 24 h after inoculation. Bars = 50 µm. Right, Number of fluorescent spots were counted in elicitor-treated leaf discs inoculated with *P. infestans* 24 hrs after inoculation. Data are means ± SE (n = 3). Data marked with asterisks are significantly different from control as assessed by two-tailed Student's *t* tests: p < 0.01.

Figure 5. *Phytophthora infestans* diacylglycerol (Pi-DAG) induce the production of phytoalexins in potato tubers. **(A)** Structures of Pi-DAG A and B. Structures equivalent to eicosapentaenoic acid (EPA) are shown in red, dotted boxes. **(B)** Structures of Pi-DAG C and D which have significantly weaker elicitor activity compared with Pi-DAG A and B. See Supplemental Figure S22 for the procedures of purification of Pi-DAGs and Supplemental Figures S23-26 for their structural analysis. **(C)** Potato tubers were treated with 0.3% DMSO (Cont.), 30 μg/ml methanol extract of *P. infestans* mycelium (Pi-MEM), or 100 μg/ml Pi-DAGs, and production of phytoalexins was detected by thin-layer chromatography. L, Lubimin; R, Rishitin; O, Oxylubimin. **(D)** EPA, but not Pi-Cer D, can induce the production of phytoalexins in potato tubers. Potato tubers were treated with 0.3% DMSO (Cont.), 100 μg/ml EPA or 100 μg/ml Pi-Cer D and produced phytoalexins were detected by LC/MS.

Figure. 6 Distinctive sets of Arabidopsis genes were upregulated by the treatment with EPA and Pi-Cer D. **(A)** Arabidopsis transformant pWRKY33-LUC containing *LUC* transgene under the control of *AtWRKY33* (AT2G38470.1) promoter was treated with 20 µg/ml EPA, Pi-Cer D or a mixture of 20 µg/ml EPA and Pi-Cer D. Chemiluminescence was monitored for 12 h after the treatment. (B) Expression profile of Arabidopsis genes treated with 20 µg/ml EPA, Pi-Cer D or a mixture of 20 µg/ml EPA and Pi-Cer D. RNAseq analysis was performed 24 h after the treatment.