Short title: Two oomycete MAMPs induce distinctive plant responses

Title of the article: Two structurally different oomycete lipophilic MAMPs induce

distinctive plant immune responses

Authors: Mohammad Shahjahan Monjil[1,2], Hiroaki Kato[3], Satomi Ota[1], Kentaro

Matsuda[1], Natsumi Suzuki[1], Shiho Tenhiro[1], Ayane Tatsumi[1], Sreynich Pring[1], Maurizio

Camagna[1], Takamasa Suzuki[4], Aiko Tanaka[1], Ryohei Terauchi[3], Ikuo Sato[1], Sotaro

Chiba[1], Kazuhito Kawakita[1], Makoto Ojika[1] and Daigo Takemoto[1*]

Affiliation:

[1] Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya, Aichi, 464-

8601, Japan

[2] Department of Plant Pathology, Bangladesh Agricultural University, Mymensingh-

2202, Bangladesh

[3] Graduate School of Agriculture, Kyoto University, Muko, Kyoto, 617-0001, Japan.

[4] College of Bioscience and Biotechnology, Chubu University, Kasugai, Aichi, 478-

8501 Japan

Corresponding author: Daigo Takemoto, E-mail:dtakemo@agr.nagoya-u.ac.jp,

Graduate School of Bioagricultural Sciences, Nagoya University, Chikusa, Nagoya,

464-8601 Japan

The author responsible for distribution of materials integral to the findings presented in

this article in accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plphys/pages/General-Instructions) is Daigo Takemoto

(dtakemo@agr.nagoya-u.ac.jp).

One-sentence summary: Two lipophilic MAMPs, Pi-Cer D and Pi-DAG, containing

microbe specific structures induce distinctive plant immune responses

Author Contribution: KK and DT designed the research. MSM, HK, SO, KM, NS, ST,

A Tatsumi, SP, TS, A Tanaka, MO and DT performed experiments. MSM, HK, KM, ST,

CM, TS, A Tanaka, MO and DT analyzed the data. MC, A Tanaka, RT, IS, SC, KK, MO

and DT supervised the experiments. MO and DT wrote the article, and MC, MO and DT

revised and edited the article.

Abstract

Plants recognize a variety of external signals and induce appropriate mechanisms to

increase their tolerance to biotic and abiotic stresses. Precise recognition of attacking

pathogens and induction of effective resistance mechanisms are critical functions for

plant survival. Some molecular patterns unique to a certain group of microbes, microbe-

associated molecular patterns (MAMPs), are sensed by plant cells as non-self molecules

via pattern recognition receptors. While MAMPs of bacterial and fungal origin have

been identified, reports on oomycete MAMPs are relatively limited. This study aimed to

identify MAMPs from an oomycete pathogen *Phytophthora infestans*, the causal agent

of potato late blight. Using reactive oxygen species (ROS) production and phytoalexin

production in potato as markers, two structurally different groups of elicitors, namely

ceramides and diacylglycerols were identified. *P. infestans* ceramides (Pi-Cer A, B and

D) induced ROS production, while diacylglycerol (Pi-DAG A and B), containing

eicosapentaenoic acid (EPA) as a substructure, induced phytoalexins production in

potato. The molecular patterns in Pi-Cers and Pi-DAGs essential for defense induction

were identified as 9-methyl-4,8-sphingadienine (9Me-Spd) and 5,8,11,14-tetraene-type

fatty acid (5,8,11,14-TEFA), respectively. These structures are not found in plants, but

in oomycetes and fungi, indicating that they are microbe molecular patterns recognized

by plants. When Arabidopsis was treated with Pi-Cer D and EPA, partially overlapping

but different sets of genes were induced. Furthermore, expression of some genes is

upregulated only after the simultaneous treatment with Pi-Cer D and EPA, indicating

that plants combine the signals from simultaneously recognized MAMPs to adapt the

defense response to pathogens.

Keywords: Ceramide, Diacylglycerol, MAMPs, Oomycete, Phytoalexins,

Phytophthora infestans, Reactive oxygen species.

Introduction

During the course of their evolution, plants were always faced with the challenges of

environmental microorganisms but have survived by developing physical barriers or

inducible resistance strategies against pathogens. The first step of inducible plant

defense against potential pathogens is the recognition of molecular patterns of microbes,

referred to as MAMPs or PAMPs (microbe- or pathogen-associated molecular patterns).

Plant cells recognize molecular patterns unique to a certain group of microbes as non-

self molecules via pattern recognition receptors (PRRs) (Ranf, 2017; Ngou et al., 2022).

In Arabidopsis, for instance, chitin, a major component of the fungal cell wall, is

recognized by a PRR complex containing LysM-RLK and co-receptor CERK1, while a

conserved sequence of bacterial flagellin (flg22) and an elongation factor EF-Tu (elf18)

induce plant defense via PRRs, FLS2 and EFR, respectively, in conjuncture with co-

receptor BAK1 (Ranf, 2017). While a variety of MAMPs of bacterial and fungal origin,

as well as the mechanisms for their recognition in plants have been studied, reports of

MAMPs from oomycete pathogens are relatively limited.

Oomycete plant pathogens resemble fungal pathogens in various aspects. Besides

morphological similarities, such as filamentous hyphae, they are also able to form

appressoria and haustoria during infection in host plants. These similarities are however

merely the result of convergent evolution, since oomycetes are phylogenetically closely

related to diatoms and brown algae (Stramenopiles), and only distantly related to fungi

(Opisthokonta). Therefore, the natures of oomycetes and true fungi are rather different.

For example, the somatic thallus of oomycetes is diploid, while it is haploid (or

dikaryotic) for true fungi, and oomycetes have mitochondria with tubular cristae

whereas true fungi have flattened ones. More importantly in terms of interaction with

plants, the composition of the cell wall and cell membrane of oomycete and fungi is

distinctive. Although varying from species to species, the fungal cell wall mainly

contains β-1,3-glucan and chitin (β-1,4-N-acetylglucosamine) (Free, 2013), whereas the

oomycete cell wall consists mainly of cellulose (β-1,4-glucan) and β-1,3-glucan

(Mélida et al., 2013). The plasma membrane of oomycetes contains fucosterol as the

end sterol, but ergosterol is the major end sterol for fungi (Gaulin et al., 2010).

Consequently, plants may have distinct mechanisms to recognize oomycete and fungal

pathogens to activate appropriate defense reactions.

Of the various oomycete genera, *Phytophthora*, *Pythium* and *Peronospora*

/*Hyaloperonospora* (downy mildew) are particularly noteworthy, as they include many

species that are problematic plant pathogens (Kamoun et al., 2015). These plant

pathogenic oomycete genera are predicted to have evolved from a common

phytopathogenic ancestor species (Thines and Kamoun, 2010). Undoubtedly the most

infamous representative of pathogenic oomycetes is *Phytophthora infestans*, the causal

agent of potato late blight, which is responsible for the Great Famine of the 1840s in

Ireland. This pathogen is an ongoing problem even today, and the total costs for control

efforts and yield losses by *P. infestans* are estimated in the range of multi-billion dollars

annually (Garelik, 2002; Fry, 2008). Problems caused by *P. infestans* are particularly

serious in developing countries where fungicides are used as the primary solution for

disease management. Although potato cultivars with introduced resistance (*R*) gene(s)

have been employed, due to selection pressure on these effectors in the pathogen

population, *R*-gene dependent resistances have been lost in many varieties (Forbes,

2012).

In this study, we purified two lipophilic elicitors from *P. infestans*. Using induction of

reactive oxygen species (ROS) production or phytoalexin production in potato as

markers for the purification of elicitors, we identified two structurally different groups

of elicitors, namely ceramides and diacylglycerols. *P. infestans* ceramide (Pi-Cer)

elicitors induced ROS production, while diacylglycerol (Pi-DAG) elicitors induced the

accumulation of phytoalexins in potato. RNAseq analysis was performed in Arabidopsis

treated with Pi-Cer, EPA (a substructure of Pi-DAG), or the mixture of both elicitors, to

investigate the difference in the activity of the two elicitors on the induction of defense

genes, and the effect of simultaneous recognition of both MAMP elicitors. We have

previously reported that a *P. infestans* ceramide Pi-Cer D is cleaved by an apoplastic

ceramidase NCER2 and resulting 9-methyl-branched sphingoid base is recognized by a

lectin receptor-like kinase RDA2 in Arabidopsis (Kato et al., 2022).

RESULTS

Purification of *Phytophthora infestans* elicitors which induce ROS production of

potato suspension-cultured cells

Previously, we have reported that crude elicitor derived from the mycelia of *P. infestans*

extracted by methanol (Pi-MEM) can induce the resistance reactions in potato (Monjil

et al., 2015). Treatment with Pi-MEM can induce the ROS production in potato

suspension cells and leaves, and production of sesquiterpenoid phytoalexins (rishitin,

lubimin and oxylubimin) in potato tubers (Fig. 1). To purify the elicitor molecules from

the mycelia of *P. infestans* and determine their structure, Pi-MEM was first fractionated

by its solubility in water and butanol. In this study, the lipophilic (butanol soluble)

fraction was further separated by column chromatography, and the elicitor fractions

were selected using their ROS inducing activity in cultured potato cells (Fig. S1). Only

fractions with clear elicitor activity were used for further purification processes. Six

fractions with significant ROS-producing activity were further purified (Fig. S1) and the

chemical structures of these elicitors were analyzed by NMR and mass spectroscopy

(Figs. S2, S3, S5-7 and S9). To compare the structural difference between elicitors and

inactive related substances, two fractions with no significant elicitor activity were also

structurally analyzed (Figs. S1, S4 and S8). Further structural analyses (Figures S10-

S16, and Supplemental document) showed that these eight substances could be divided

into two groups, ceramide (Cer) and ceramide-phosphoethanolamine (CerPE). We

designated the four ceramide compounds Pi-Cer A-D (*P. infestans* Ceramide) and the

remaining four compounds Pi-CerPE A-D respectively. Except for the

phosphoethanolamine in Pi-CerPE, compounds that share the same alphabetical suffix

are otherwise structurally identical (Fig. 2 and S17A). Pi-Cer A, B and D were able to

induce ROS production, whereas Pi-Cer C had only a marginal ROS inducing activity

(Figs. 2 and S1). Similarly, Pi-CerPE A, B and D were active elicitors for ROS

production (Figs. S1 and S17). For both Pi-Cers and Pi-CerPEs, the D type ones were

the most active, followed by B and A in that order (Figs. 2 and S17). An Arabidopsis

transformant containing a *LUC* transgene under the control of *WRKY33* (AT2G38470)

promoter (pWRKY33-LUC) (Kato et al., 2022) was employed for the detection of

elicitor activity of Pi-Cers and Pi-CerPEs in Arabidopsis. Treatment of Pi-Cers and Pi-

CerPEs can induce the transient activation of the promoter with a peak at approx. 90

min after elicitor treatment. Same as in the case of ROS production in potato suspension

cells (Figs. 2 and S17), Pi-Cer D and Pi-CerPE D showed the highest elicitor activity

and clearly lower elicitor activity was detected for Pi-Cer C and Pi-Cer PE C in

Arabidopsis (Fig. S18).

Pi-Cers are MAMPs of oomycete pathogens

To investigate whether the substances corresponding to Pi-Cers are contained in other

phytopathogenic oomycetes, a modified purification process was applied to partially

purify the Pi-Cers from *Pythium aphanidermatum*. *Py. aphanidermatum* is an oomycete

plant pathogen with a wide host range, causing seedling damping-off, root and stem rots

and blights of a wide range of important crops (e.g. tomato, soybean, cucumber and

cotton). Although the composition of Pi-Cers was different from that in *P. infestans*, Pi-

Cer A, B, C and D were detected in *Py. aphanidermatum*. On the other hand, Pi-Cers

were not detected in several tested fungal plant pathogens, such as *Botrytis cinerea,*

Fusarium oxysporum and *Colletotrichum orbiculare* (Fig. S19). These results suggest

that the Pi-Cers are molecules specific to oomycetes.

Pi-Cer D and Pi-CerPE D are elicitors of ROS production in dicot and monocot

plants

Because Pi-Cer D and Pi-CerPE D had the highest elicitor activity among Pi-Cers and

Pi-CerPEs, respectively, and were also more abundantly purified than the A and B type

substances (Fig. S1), Pi-Cer D and Pi-CerPE D were mainly used in the following

experiments. In potato suspension-cultured cells, Pi-MEM induced ROS production

peaks approx. 3 h after treatment (Fig. S20). The same pattern was observed after Pi-

Cer D and Pi-CerPE D treatment (Fig. S20), suggesting that Pi-Cers and Pi-CerPEs

could be the major ROS inducing substances in Pi-MEM. Potato leaves were treated

with Pi-Cer D, Pi-CerPE D or Pi-MEM by syringe infiltration and induction of ROS

production was detected. ROS production was induced by these elicitors with a peak at

approx. 12 h after the treatment. Similarly, ROS production was detected in leaves of *A.*

thaliana and rice treated with Pi-Cer D or Pi-CerPE D 12 h after the treatment (Fig. 3).

These results indicated that Pi-Cer D and Pi-CerPE D are recognized by both dicot

(potato, Arabidopsis) and monocot (rice) plants as MAMPs.

Treatment of Pi-Cer D and Pi-CerPE D enhances the resistance of potato against *P.*

infestans

The effect of pre-treatment with ceramide elicitors on the resistance of potato against *P.*

infestans was investigated (Fig. 4). Potato leaves were treated with 10 µg/ml Pi-Cer D,

Pi-CerPE D or 100 µg/ml Pi-MEM, and inoculated with a zoospore suspension of *P.*

infestans at 24 h after the elicitor treatment. Within 3 days after the inoculation, leaves

of the control plant showed water-soaked disease symptoms, and the lesions on most

leaves extended over the entire leaves within 7 days. In contrast, fewer and smaller

spots were developed in leaves treated with Pi-Cer D, Pi-CerPE D or Pi-MEM.

Evaluation of disease symptoms up to 7 days after the inoculation indicated that

pretreatment with Pi-Cer D and Pi-CerPE D can enhance the resistance of potato leaves

against *P. infestans* (Fig. 4). Inoculated leaves were stained with aniline blue to

visualize the penetration of *P. infestans* as fluorescent spots of deposited callose beneath

the penetration sites at 24 h after the inoculation. Compared with control leaves, fewer

fluorescence spots were detected in leaves pretreated with Pi-Cer D, Pi-CerPE D or Pi-

MEM elicitors (Fig. 4D), which indicates that Pi-Cer D and Pi-CerPE D can enhance

the pre-penetration resistance of potato leaves against *P. infestans*.

Isolation of Pi-DAGs as elicitors for the induction of phytoalexin production in

potato tuber

In potato tubers, accumulation of phytoalexins (rishitin, lubimin and oxylubimin) was

induced by the treatment with the Pi-MEM (Fig. 1). However, treatment of Pi-Cers or

Pi-CerPEs did not induce the production of phytoalexins in potato tubers (Fig. S21),
which suggested that Pi-MEM includes other elicitors that can induce phytoalexin
production. The initial step of our purification protocol consisted of fractioning the
butanol-soluble compounds via silica gel flash column chromatography, yielding eight
fractions. Of these eight fractions, the third (RM-I-130-3) and sixth (RM-I-130-6)
fraction had shown ROS inducing activity and were used for the isolation of Pi-Cers
and Pi-CerPEs (Figs. S1 and 22). The same eight fractions were now used to evaluate
their ability to induce phytoalexin production in potato tubers.   Substantial elicitor
activity was only observed in the first fraction (RM-I-130-1) and it was therefore further
purified (Fig. S22). Among the five fractions obtained from the following flash column
chromatography, the third (ST-I-8-3) and fourth (ST-I-8-4) fraction showed elicitor
activity that resulted in phytoalexin production. These fractions were later found to
contain mainly 1,3-DAG (diacylglycerol) and 1,2-DAG, respectively. Both ST-I-8-3 and
-4 were further separated by HPLC using an ODS column and at least 5 out of 9
resultant fractions showed elicitor activity (Fig. S22). These results indicated that
several 1,3- and 1,2-DAGs derived from *P. infestans* are elicitors that can induce the
phytoalexin production of potato tubers. Two fractions (ST-I-14-4 and ST-I-23-4) with
relatively higher elicitor activity, yield and purity were selected for further structural
analyses. Two additional fractions (ST-I-14-8 and ST-I-23-8) which showed almost no
elicitor activity were also subjected for structural analyses in order to narrow down the
molecular patterns that may be critical for the elicitor activity. Further structural
analyses (Figs. S23-26) revealed that the active substances from ST-I-14-4 and ST-I-23-
4 were 1,2- and 1,3-DAGs which both contained an eicosapentaenoic acid (EPA) and
linoleic acid as fatty acid chains each, and were designated as Pi-DAG A and B,
respectively (Figs. 5 and S22). The inactive substances (ST-I-14-8 and ST-I-23-8) were
found to be 1,2- and 1,3-DAGs containing palmitic acid and linoleic acid as fatty acid
chains, and were designated as Pi-DAGs C and D, respectively. A variety of Pi-DAGs
such as Pi-DAG A-D, are also contained in *Py. aphanidermatum*, suggesting that Pi-
DAGs are a group of molecules commonly found in oomycete plant pathogens (Fig.
S27).

Based on the structural difference of active and inactive Pi-DAGs (Fig. 5A and B), EPA
was predicted to be the essential structure for the elicitor activity of Pi-DAGs A and B.
Consistently, EPA from *P. infestans* has previously been identified as an elicitor
molecule (Bostock et al., 1981). EPA can be considered as a MAMP of oomycete
pathogens, as EPA is not found in higher plants, but an essential component of the
plasma membrane of oomycete cells (Bostock et al., 2011). We confirmed that EPA, but
not Pi-Cer D, can induce the production of sesquiterpenoid phytoalexins (rishitin and its
precursors, lubimin and oxylubimin) in potato tubers (Fig. 5C and D). On the contrary,
Pi-Cer D, but not EPA, induced ROS production in potato leaves (Fig. S28), thus
indicating that the structurally different oomycete MAMPs, EPA and Pi-Cer D, induce
distinctive plant immune responses.
257
Identification of the essential structure of unsaturated fatty acid to be recognized
as MAMPs in potato
EPA is an omega-3 polyunsaturated fatty acid with 20 carbon chain length and 5 double
bonds (20:5, Δ5,8,11,14,17, ω-3) (Fig. 6A). The primary unsaturated fatty acids
contained in plants are linoleic acid (18:2, Δ9,12, ω-6) and α-linolenic acid (18:3,
Δ9,12,15, ω-3) (Cahoon and Li-Beisson, 2020), thus it is presumed that plants recognize
the characteristic structure of unsaturated fatty acids not contained in plants as MAMPs.
To identify the essential molecular pattern of unsaturated fatty acid to be recognized as
a MAMP, structurally related unsaturated fatty acids were tested for their elicitor
activity on potato. Two days after the treatment of various fatty acids at 100 µg/ml on
potato tuber, accumulated rishitin was extracted and quantified by GC/MS. As predicted,
three unsaturated fatty acids derived from plants, linoleic acid (LA), α-linolenic acid
(ALA) and γ-linolenic acid (18:3, Δ6,9,12, ω-6, GLA) had no elicitor activity.
Treatment of arachidonic acid (20:4, Δ5,8,11,14, ω-6, ALA), which was previously
reported as an elicitor (Bostock et al., 1981), induced rishitin production, but other
structurally related unsaturated fatty acid, namely eicosadienoic acid (20:2, Δ11,14, ω-6,
EDA), eicosatrienoic acid (20:3, Δ8,11,14, ω-6, ETA), mead acid (20:3, Δ5,8,11, ω-9,
MA) and docosahexaenoic acid (22:6, Δ4,7,10,13,16,19, ω-3, DHA) did not show

elicitor activity on potato, even though DHA includes conjugated double bonds or MA

shares an identical structure to AA and EPA at the end of the carboxylic acid group (Fig.

6). Thus, 5,8,11,14-tetraene-type fatty acid (5,8,11,14-TEFA) is presumed to be the

essential structure recognized by potato as a MAMP.

Distinctive sets of genes were upregulated in Arabidopsis treated with EPA, Pi-Cer

D and their mixture

To analyze the differences in plant responses resulting from the recognition of EPA and

Pi-Cer D, an Arabidopsis pWRKY33-LUC transformant was treated with 100 µg/ml

EPA and Pi-Cer D. While Pi-Cer D can induce the transient activation of the *WRKY33*

promoter, only marginal induction of the *WRKY33* promoter was detected by the

treatment with EPA (Fig. 7A). Similarly, callose deposition, a form of penetration

resistance (Ellinger et al., 2013), was induced by either Pi-Cer D or EPA treatment on

leaves of Arabidopsis seedlings, but Pi-Cer D induced significantly greater number of

depositions (Fig. S29). RNAseq analysis was performed for Arabidopsis seedlings 12 h

after the treatment with 100 µg/ml EPA, Pi-Cer D or a mixture of EPA and Pi-Cer D.

When the number of upregulated genes was compared under the same criteria (Log2

fold change ≥2, p ≤0.05), Pi-Cer D had only 211 upregulated genes, compared to 1,422

genes for EPA treatment. Unexpectedly, simultaneous treatment with Pi-Cer D and EPA

did not result in an expression pattern that would be expected by the mere combination

of either treatment. Instead, a significantly a smaller number of genes (439 genes) was

upregulated by the mixture of EPA and Pi-Cer D compared with the single treatment of

EPA, and 76.8% of genes upregulated by the mixture of EPA and Pi-Cer D (337 genes)

were specifically induced by the co-treatment of both elicitors (Fig. 7B). Similarly,

clustering analysis of significantly up- or down-regulated genes further revealed that

treatment with EPA or Pi-Cer D leads to distinct gene expression patterns. While some

genes respond similarly to either treatment, other genes are induced exclusively by only

EPA or Pi-Cer D treatments (Fig. 7C). Expression of some genes was significantly

upregulated only by simultaneous treatment with both elicitors (Fig. S30A), whereas

other genes induced by EPA were attenuated by the co-treatment with Pi-Cer D (or vice

versa) (Fig. S30B and C), resulting in an expression pattern different from either single

treatment.

To interpret the overall influence of treatment with EPA, Pi-Cer D or their mixture,

upregulated genes were assigned to gene ontology (GO) terms using the analysis tool

PANTHER (Thomas et al., 2022) for GO enrichment analysis (Fig. 8). Biological

processes (BP) enhanced by Pi-Cer D treatment include categories related to plant

defense against microorganisms such as "Defense response to fungus (GO:0050832)",

"Defense response to bacterium (GO:0042742)". EPA treatment also enhanced

"Defense response to fungus (GO:0050832)", but also induced distinctive categories

such as "Oligopeptide transport (GO:0006857)", "Detoxification (GO:0098754)" and

"Inorganic ion homeostasis (GO:0098771)". It should be noted that although the

number of genes induced by EPA treatment (1422 genes) used for GO analysis was

approx. 7-fold compared to that of Pi-Cer D (211 genes), the number of enriched GO

terms hit was less than 1.5-fold (23 terms for Pi-Cer D and 34 terms for EPA, Fig. 8),

indicating that the genes induced by solo treatment of EPA are not focused on a specific

pathway or response. More importantly, GO analysis using 432 genes induced by

simultaneous treatment with EPA and Pi-Cer D detected the relatively larger enrichment

of much more GO terms (141 terms), including GOs involved in disease resistance,

such as "Camalexin biosynthetic process (GO:0010120)" (also see Fig. 7D), "Response

to chitin (GO:0010200)", "Innate immune response (GO:0045087)", "Defense response

to fungus (GO:0050832)", "Defense response to bacterium (GO:0042742)" and so on.

This result could indicate that the simultaneous recognition of multiple MAMPs by

plants better defines the appropriate choices of the stress responses in the plant cells.

DISCUSSION

As the major autotroph on land, terrestrial plants are suitable predation targets for many

organisms, including bacteria, fungi, oomycetes, insects, animals, or even some

parasitic plants, which consume the organic compounds produced by plants. Therefore,

since their ancestral phyla, defense against various types of foreign enemies has been of

utmost importance. While plants form surface physical barriers and superficially

accumulate antimicrobial substances on their surfaces as the first line of defense in
general, they also need to activate greater resistance when attacked by microorganisms,
and recognition of so-called MAMPs is the key event to activate the next line of plant
disease resistance.

Characteristic surface structures of microorganisms are recognized by plant cells as
MAMPs. The major components of fungal cell walls (such as chitin and β-glucan) and
the secretory proteins (NEP and elicitins), as well as the bacterial flagellin and
lipopolysaccharides are typical examples of MAMPs localized at the surface (Jones and
Takemoto, 2004; Ngou et al., 2022). Flagellin, lipopolysaccharides, chitin and β-glucan
are recognized as MAMPs for induction of innate immunity in animal cells by distinct
mechanisms from plants and are the results of convergent evolution (Takeuchi et al.,
1999; Hayashi et al., 2001, He et al., 2021). Whereas some MAMPs derived from
internal molecules, such as bacterial elongation factor EF-Tu or extracellular DNA and
RNA, are released after disruption of the microbial cells, and then recognized by the
plant cells (Kunze et al., 2004; Bhat and Ryu, 2016). When plant cell walls are damaged
by pathogen attack, fragments of oligosaccharides (such as cello-oligosaccharides and
xylo-oligosaccharides) are recognized by cell surface receptors as DAMPs (Martín-
Dacal et al., 2023; Pring et al., 2023).

In this study, lipophilic MAMPs derived from microbial cell membrane, Pi-Cers and Pi-
DAGs, were purified from a representative phytopathogenic oomycete, *P. infestans*, for
their activity in inducing ROS production and phytoalexin accumulation in potato.
These substances were also contained in polyxenous oomycete pathogen *Pythium*
aphanidermatum (Figs. S19, S27) and other oomycetes (Moreau et al., 1998, Fernandes
et al., 2019). Pi-Cers and Pi-DAGs (or EPA) activated resistance responses in potato,
Arabidopsis and other plant species, indicating that they are oomycete MAMPs
recognized by plants (Fig. 9, Bostock et al., 2011).

Two characteristic structures of microbial plasma membrane lipids are recognized
by plant cells

Ceramides and diacylglycerols are principal structural elements of cell membranes.

Ceramides are composed of a sphingoid base and a fatty acid joined via an amide bond.

The most evident structural difference between active Pi-Cers and inactive Pi-Cer C is

the 8,10-diene and methyl-branching at C-9 in the sphingoid base (Fig. 2). The same

structure was found in ceramide-phosphoethanolamines Pi-CerPE A, B and D purified

from other fractions (Figs. S1, S17). In a previous report, we indicated that 9-methyl

branching and the 4,8-double bond structure of the sphingoid base (9Me-Spd) represent

the epitope structure for its elicitor activity in Arabidopsis (Kato et al., 2022). Although

Pi-Cers are not detected in fungal pathogens (Fig. S19), cerebrosides (a group of

glycosphingolipids) have been identified as elicitor molecules from rice blast pathogen

Pyricularia oryzae (Koga et al., 1998, Fig. 9). Cerebroside elicitors isolated from *P.*

oryzae and other fungal pathogens also contain epitope structure 9Me-Spd (Umemura et

al., 2000). Given that 9Me-Spd is commonly found in (evolutionarily distant) oomycete

and fungal species but essentially not found in plants (Moreau et al., 1998; Sperling and

Heinz, 2003, Jiang et al., 2021), this structure appears to be a molecular pattern that can

be leveraged by the plant to recognize potentially pathogenic microorganisms. By

mutagenesis of the Arabidopsis pWRKY33-LUC marker line and screening of mutants

insensitive to Pi-Cer D, Kato et al. (2022) identified that a lectin receptor kinase RDA2

is the receptor of 9Me-Spd in Arabidopsis. Moreover, cleavage of Pi-Cer D by an

apoplastic ceramidase NCER2 is essential for perception of 9Me-Spd by RDA2 in

Arabidopsis, indicating that the excision of microbial components by the plant is an

important process for the activation of pattern-triggered immunity (PTI) of plants (Fig.

9).

Diacylglycerols consist of two fatty acid chains linked to a glycerol molecule through

covalent ester bonds. Pi-DAGs A and B, which have elicitor activity and induce potato

phytoalexin production, are 1,2- and 1,3-diacylglycerols, respectively, and share the

same fatty acids they consisted of were the same. Comparison of Pi-DAGs A and B with

Pi-DAGs C and D, which had no elicitor activity (Fig. 5), suggested that a structure

containing five unsaturated fatty acids is the site recognized by the plant. In most plants,

the predominant unsaturated fatty acids are oleic acid (18:1, ω-9), linoleic acid (18:2, ω-

6), and α-linolenic acid (18:3, ω-3) (Harwood, 1988), as well as γ-linolenate (18:3, ω-6),

which is found in a few seed oils (Nykiforuk et al., 2012). Natural unsaturated fatty

acids in plants contain one to three conjugated double bonds, and these unsaturated fatty

acids showed no elicitor activity on potato tuber (Fig. 6). In contrast, EPA, which is a

pentaene fatty acid (20:5, ω-3) contained in bacteria, fungi and oomycete (Adarme-

Vega et al., 2012), was previously reported as an elicitor of phytoalexin production in

potato by Bostock et al. 1981. In this study, we identified the essential and sufficient

structure of microbe-derived unsaturated fatty acids that act as MAMP. Although

Bostock et al. (1981) also reported that docosahexaenoic acid (DHA, 22:6, ω-3) had

elicitor activity, experiments conducted in this study with three different lots of DHA

showed that treatment with DHA has no elicitor activity for the induction of phytoalexin

production in potato (Fig. 6), suggesting that four or more conjugated double bonds

alone are not sufficient to be recognized as a MAMP by plants. Given that arachidonic

acid (20:4, ω-6), but not mead acid (20:3, ω-9), showed elicitor activity on potato,

5,8,11,14-tetraene-type fatty acid (5,8,11,14-TEFA) is presumed to be the essential

structure recognized by plants as a MAMP. Collectively, plants activate PTI by

recognizing characteristic microbial characteristic structures in the cell membrane.

Since plants secrete enzymes that partially degrade MAMPs such as Pi-Cers, chitin, and

β-glucans, such secreted enzyme (e.g. phospholipase or esterase) might also be involved

in the degradation of Pi-DAGs via the exposure of the 5,8,11,14-TEFA structure (Fig. 9).

Simultaneous recognition of two lipophilic MAMPs induces distinctive plant

immune responses

Over the course of infection by plant pathogens, plants are exposed to a series of

MAMPs, at different times of the infection process. While patterns on the microbial cell

wall and cell membrane are expected to be recognized first, intracellular components

and DAMPs (e.g. oligosaccharides derived from plant cell walls, Pring et al., 2023)

should be released at later stages of the infection. It is presumed that plants are able to

utilize these temporal cues to activate a response better suited to the disease progression.

Since we demonstrated that co-exposure to two MAMPs can result in a different

defense response, plants may leverage this mechanism to infer information of the
disease progression and fine-tune their response.

Two MAMPs derived from the oomycete membrane were isolated in this study, each
activating different resistance responses in potato, namely ROS generation and
phytoalexin accumulation respectively, indicating that distinct plant signaling is
activated by these two MAMPs. In Arabidopsis, Pi-Cer D treatment activates the
promoter of *AtWRKY33*, a central activator of disease resistance (Zheng et al., 2006),
while EPA hardly activates *AtWRKY33*. Consistently, distinctive sets of genes are
induced by the treatment with these two elicitors, though GO terms for up-regulated
genes after the treatment with Pi-Cer D or EPA are both includes terms categorized
under plant disease resistance. Importantly, simultaneous treatment with these two
MAMPs was shown to activate GO categories related to disease resistance that were not
triggered by the single treatment (Fig. 8). These results seem to indicate that plants may
combine the signals from simultaneously recognized MAMPs to specifically adapt the
defense response to a particular type of pathogen. Insight into how these elicitor signals
are processed, as well as how the response pathways of both elicitors affect one another,
will require further analysis. If additional MAMPs and DAMPs activating different
signaling systems were to be determined, we might be able to devise effective
combinations that effectively activate resistance. Such MAMP/DAMP recipes might be
tailored to act as efficient and stable resistance inducers for use in agricultural
production (Pring et al., 2023).

MATERIALS AND METHODS
Biological materials, growth conditions and inoculation
Potato suspension-cultured cells were prepared from calluses induced from potato tuber
discs (cv. Sayaka carrying the *R1* and *R3* genes) in callus-inducing medium [1 ml/l B5-
vitamin (100 g/l Myo-inositol, 2.5 g/l glycine, 500 mg/l thiamine-HCl, 500 mg/l
pyridoxine-HCl, 5 g/l nicotinic acid, 50 mg/l biotin, 50 mg/l folic acid), 2 mg/l 1-
naphthaleneacetic acid (NAA), 0.5 mg/l kinetin, 3% sucrose, 0.2% phytagel and
Murashige-Skoog basal medium], under dark condition. Suspension-cultured potato

cells were grown by agitation at 130 rpm, 23°C in 95 ml of Murashige-Skoog medium

supplemented with 30 mg/ml sucrose, 1 mg/l thiamine, 100 mg/l myo-inositol, 200 mg/l

KH_2PO_4, and 0.2 mg/l 2,4-dichlorophenoxyacetic acid. Cells were sub-cultured every

week and used for experiments 4-5 days after the subculturing. Potato, rice and

Arabidopsis (Col-0) plants were grown at 23-25°C under a 16 h photoperiod and an 8 h

dark period in an environmentally controlled growth room. Tubers of potato (cv.

Rishiri) carrying the *R1* gene were stored at 4°C until use. *P. infestans* isolate 08YD1

and PI1234-1 (Shibata et al., 2010, 2011) was maintained on rye-media at 10°C.

Inoculation of potato leaves with a zoospore suspension of *P. infestans* was performed

as described previously (Shibata et al., 2016). *Pythium aphanidermatum* strain Pyaph

isolated from cucumber, *Botrytis cinerea* strain AI18 isolated from strawberry leaves

(Kuroyanagi et al., 2022), *Fusarium oxysporum* f. sp. *melonis* strain Mel02010 (Namiki

et al., 1998) and *Colletotrichum orbiculare* strain 104-T (Ishida and Akai 1969) were

grown on potato dextrose agar (PDA) at 25°C and maintained at -80°C in 20% glycerol.

Preparation of methanol extract of *Phytophthora infestans* mycelia (Pi-MEM)

The *Phytophthora infestans* isolates PI1234-1 were grown on rye-seed extract agar

medium in test tubes at 18°C in dark condition for 2 weeks. Parts of the growing

mycelia were placed in 100 ml flasks containing 20 ml of rye liquid nutrient medium of

rye seed-extract (60 g rye seed), 20 g of sucrose and 2 g of yeast extract per 1 liter and

incubated in the dark for 2 weeks at 18°C to allow the growth of the mycelia. The

mycelial mats grown in the liquid medium were washed thoroughly with water. To

remove excess water, water was filtered through filter paper (Toyoroshi No. 2) under

reduced pressure and the tissue then frozen at -20°C. The collected mycelia of *P.*

infestans were ground in liquid nitrogen by mortar and pestle. The ground mycelia were

transferred to a 50 ml tube containing methanol (10 ml/g mycelia). The mycelia

suspension was finely grounded using a polytron type homogenizer (HG30, Hitachi

Koki, Japan) for 2 min. After centrifugation at 4°C, $3000 \times g$ for 30 min, the supernatant

was collected and dried using an evaporator and used as Pi-MEM elicitor. Further

procedures for the purification of Pi-Cers and Pi-DAGs from Pi-MEM and their

structural analyses with spectral data are described in the supplemental document.

Mycelia of *Pythium aphanidermatum*, *B. cinerea*, *F. oxysporum* f. sp. *melonis* and *C.*
orbiculare were grown in 50 ml of potato dextrose broth (PDB) in 100 ml flasks at 25˚C
with gentle shaking.

Measurement of ROS production in potato suspension-cultured cells
The relative intensity of ROS generation in potato suspension-cultured cells was
measured by counting photons from L-012-mediated chemiluminescence. The potato
suspension-cultured cells (50 mg/ml) were washed with the assay buffer (175 mM
mannitol, 50 mM MES-KOH, 0.5 mM $CaCl_2$ and 0.5 mM K_2SO_4, pH 5.7) twice for
removal of the liquid culture medium. For the detection of ROS produced in cultured
cells, the cells were resuspended in assay buffer, equilibrated for 1 h at 100 rpm at 23°C.
Cells were then treated with elicitors and incubated under the same condition for 3 h.
After the incubation, chemiluminescence was measured using 20 mM L-012 (Wako
Pure Chemical, Osaka, Japan) in a multimode microplate reader Mithras LB940
(Berthold Technologies, Bad Wildbad, Germany).

Measurement of ROS production in plant leaves
The relative intensity of ROS production was determined by counting photons from L-
012-mediated chemiluminescence. To detect the ROS production in potato, Arabidopsis
and rice leaves, 0.5 mM L-012 in 10 mM MOPS-KOH (pH 7.4) was allowed to
infiltrate to the intercellular space of leaves. For potato and Arabidopsis, a syringe
without a needle was used for leaf infiltration, whereas vacuum infiltration was used for
rice. Chemiluminescence was monitored using a photon image processor equipped with
a sensitive CCD camera in the dark chamber at 20°C (Aquacosmos 2.5; Hamamatsu
Photonics, Shizuoka, Japan) and quantified using the U7501 program (Hamamatsu
Photonics). For the data shown in Supplemental Fig. S28, chemiluminescence was
monitored using Lumino Graph II EM (ATTO, Tokyo, Japan).

Detection of potato phytoalexins by thin-layer chromatography (TLC)
Potato phytoalexins, exuded from potato tissue, were extracted with ethyl acetate as
described previously (Noritake et al., 1996). The extract was separated on TLC plates

(TLC aluminum sheet of silica gel 60, Merck, Whitehouse Station, NJ, USA), which

were developed with cyclohexane:ethyl acetate (1:1, v/v) and visualized by spraying

with sulfuric acid containing 0.5% vanillin followed by heating at 120°C.

Detection of potato phytoalexins by liquid chromatography/mass spectrometry

(LC/MS) and as chromatography/mass spectrometer (GC/MS)

Potato tuber discs (2 cm in diameter and approx. 4 mm thick) were prepared and

incubated in a humidified chamber in dark at 23°C for 24 h before treatment with

elicitors. The upper side of incubated potato tubers was then treated with 100 µl of

elicitor solution and further incubated at 23°C in the dark for 48 h before the extraction

of phytoalexins. Four tuber disks/sample were immersed in 5 ml of ethyl acetate and

shaken for 1 h, then the organic solvent was collected and evaporated. The residual

material was redissolved in 100 µl of 50% (v/v) acetonitrile and 2 µl was injected for

the analysis with LC/MS (6520 Accurate-Mass Q-TOF connected to Agilent 1100

Series HPLC, Agilent Technologies, Santa Clara, CA, USA) with a Cadenza CD-C18

column (i. d. 3 x 50 mm, Imtakt, Kyoto, Japan) as previously described (Imano et al.,

2022). Detection and quantification of rishitin produced in potato tuber by GC/MS

using an Agilent Technologies 7890A GC System with a DuraBond Ultra Inert column

(length 30 m; diameter 0.25 mm; film 0.25 µm, Agilent Technologies, Santa Clara, CA,

USA) as previously described (Camanga et al., 2020).

Measuring the activation of *AtWRKY33* promoter

Seeds of Arabidopsis containing *Luciferase* marker gene under the control of

AtWRKY33 (AT2G38470) promoter (Kato et al., 2022) were surface sterilized with 3%

hydrogen peroxide and 50 % ethanol for 1 min with gentle shaking, washed with

sterilized water, and then individual sterilized seeds were placed in separate wells of a

96 microwell plate containing 150 µl of Murashige and Skoog (MS) liquid medium (1/2

MS salts, 0.05% [w/v] MES, 0.5% [w/v] Sucrose, adjusted to pH 5.8 with NaOH) with

50 µM D-Luciferin potassium salt (Biosynth Carbosynth, Compton, UK), covered with

a clear plastic cover. Plates containing Arabidopsis seeds were then incubated in a

growth chamber for approx. 12 days at 23ºC with 24 h light. After the treatment of elicitors, chemiluminescence intensity derived from expressed luciferase was measured using Mithras LB940 (Berthold) for 12 h.

RNA-seq and gene ontology enrichment analysis

Total RNA was extracted from 6 seedlings/sample of 10-days old Arabidopsis 24 h after the treatment with elicitors using the RNeasy Plant Mini Kit (QIAGEN, Hilden, Germany). Libraries were constructed using KAPA mRNA Capture Kit (Roche, Basel, Switzerland) and MGIEasy RNA Directional Library Prep Set (MGI, Shenzhen, China), and sequenced on DNBSEQ-G400RS (MGI) with 150 bp paired-end protocol. The RNA-seq reads were filtered using trim-galore v.0.6.6 (Martin, 2011, bioinformatics.babraham.ac.uk) and mapped to the Arabidopsis genome (TAIR 10.1; RefSeq: GCF_000001735.4) using HISAT2 v.2.2.1 (Kim et al., 2019) and assembled via StringTie v.2.1.7 (Kovaka et al., 2019). Significant differential expression was determined using DESeq2 v.1.32.0 (Love et al., 2014). All software used during RNA-seq analysis was run with default settings. The expression profile was calculated from the log2-fold expressions using the clustermap function from seaborn v. 0.11.1 (Waskom, 2021). Gene ontology (GO) enrichment analysis was performed using the PANTHER statistical overrepresentation test (http://pantherdb.org; Version 17.0, Thomas et al., 2022) with default settings (Fisher's exact test, False discovery rate (FDR) < 0.05). The heatmap was plotted using SRplot (https://www.bioinformatics.com.cn/en). RNA-seq data reported in this work are available in GenBank under the accession numbers DRA017453.

ACKNOWLEDGMENTS

We thank Ms. Kayo Shirai (Hokkaido Central Agricultural Experiment Station, Japan) and Dr. Seishi Akino (Hokkaido University, Japan) for providing *P. infestans* isolate, Dr. Kenji Asano, Dr. Kotaro Akai and Mr. Seiji Tamiya (National Agricultural Research Center for Hokkaido Region, Japan) and Mr. Yasuki Tahara (Nagoya University, Japan) for providing tubers of potato cultivars. We also thank Dr. Masaharu Kubota (National Agriculture and Food Research Organization, Japan) for providing *Pythium*

aphanidermatum strain, Prof. Takashi Tsuge (Chubu University, Japan) for providing

Fusarium oxysporum strain and Prof. Yoshitaka Takano (Kyoto University, Japan) for

providing *Colletotrichum orbiculare* strain. This work was supported by a Grant-in-Aid

for Scientific Research (B) (17H03771, 20H02985, 23H02212) to DT and (17H03963)

to KK from the Japan Society for the Promotion of Science (JSPS), and by Japan

Science and Technology Agency (JST), PRESTO (JPMJPR22D2) to HK.

SUPPLEMENTAL DATA

Supplemental Figure S1. Purification procedure of *Phytophthora infestans* ceramide
(Pi-Cer) and ceramide phosphoethanolamine (Pi-CerPE) elicitors.

Supplemental Figure S2. NMR spectra of Pi-Cer A (CDCl$_3$, 400 MHz for ^{1}H, 100
MHz for ^{13}C).

Supplemental Figure S3. NMR spectra of Pi-Cer B (CDCl$_3$, 600 MHz for ^{1}H, 150
MHz for ^{13}C).

Supplemental Figure S4. NMR spectra of Pi-Cer C (CDCl$_3$, 400 MHz for ^{1}H, 100
MHz for ^{13}C).

Supplemental Figure S5. NMR spectra of Pi-Cer D (CDCl$_3$, 600 MHz for ^{1}H, 150
MHz for ^{13}C).

Supplemental Figure S6. NMR spectra of Pi-CerPE A (CDCl$_3$-CD$_3$OD 4:1, 400 MHz
for ^{1}H, 100 MHz for ^{13}C).

Supplemental Figure S7. NMR spectra of Pi-CerPE B (CDCl$_3$-CD$_3$OD 4:1, 600 MHz
for ^{1}H, 150 MHz for ^{13}C).

Supplemental Figure S8. NMR spectra of Pi-CerPE D (CDCl$_3$-CD$_3$OD 4:1, 600 MHz
for ^{1}H, 150 MHz for ^{13}C).

Supplemental Figure S9. NMR spectra of Pi-CerPE D (CDCl$_3$-CD$_3$OD 4:1, 600 MHz
for ^{1}H, 150 MHz for ^{13}C).

Supplemental Figure S10. MS/MS fragmentation of Pi-Cer A (precursor ion: [M +
Na]$^+$).

Supplemental Figure S11. Two-dimensional NMR correlations of Pi-Cer B (thick
bonds: DQF-COSY, curved arrows: HMBC, l + m =15).

Supplemental Figure S12. MS/MS fragmentation of Pi-Cer B (precursor ion: [M +
Na]$^+$).

Supplemental Figure S13. Two-dimensional NMR correlations of Pi-Cer B (thick
bonds: DQF-COSY, curved arrows: HMBC, l + m + n = 19).

Supplemental Figure S14. DQF-COSY (thick bonds), HMBC (curved arrows), and
NOESY (dashed curves) correlations of Pi-Cer D.

Supplemental Figure S15. Linked-scan FAB MS/MS (negative ion mode) of the fatty
acid derived from Pi-Cer D by acid hydrolysis.

Supplemental Figure S16. DQF-COSY (thick bonds) and HMBC (curved arrows)
correlations of Pi-CerPE D.

Supplemental Figure S17. Ceramide phosphoethanolamine (Pi-CerPE) elicitors
purified from methanol extract of *P. infestans* mycelia (Pi-MEM), which can induce the
production of reactive oxygen species (ROS) in potato suspension-cultured cells.

Supplemental Figure S18. Activation of *A. thaliana WRKY33* promoter by Pi-Cer and
Pi-CerPE elicitors.

Supplemental Figure S19. Pi-Cer contents in mycelia of oomycete and fungal plant
pathogens.

Supplemental Figure S20. Induction of reactive oxygen species (ROS) production by
Pi-Cer D and Pi-CerPE D treatment in potato suspension-culture cells.

Supplemental Figure S21. Production of phytoalexins in potato tuber is not induced by
Pi-Cers and Pi-CerPEs.

Supplemental Figure S22. Purification procedure of *Phytophthora infestans*
diacylglycerol (Pi-DAG) elicitors.

Supplemental Figure S23. ESI-TOF MS of Pi-DAG A. NMR spectra of Pi-DAG A
(CDCl$_3$, 400 MHz for ^{1}H and 100 MHz for ^{13}C). Two-dimensional NMR of Pi-DAG A
(CDCl$_3$, 400 MHz). Determination of fatty acids of Pi-DAG A by MS/MS.

Supplemental Figure S24. ^{1}H NMR spectrum of Pi-DAG B (CDCl$_3$, 400 MHz).
Determination of fatty acids in Pi-DAG B by negative ion FAB MS

Supplemental Figure S25. NMR spectra of Pi-DAG C (CDCl$_3$, 400 MHz for ^{1}H, 100
MHz for ^{13}C). Determination of fatty acids in Pi-DAG C by positive ion ESI MS/MS.
Confirmation of fatty acid linkage in Pi-DAG C.

Supplemental Figure S26. NMR spectra of Pi-DAG D (MeOD, 400 MHz for ^{1}H, 100
MHz for ^{13}C). Determination of fatty acids in Pi-DAG D by hydrolysis followed by
negative ion FAB MS and MS/MS analysis of two fatty acid products.

Supplemental Figure S27. Pi-DAG contents in mycelia of oomycete plant pathogens.

Supplemental Figure S28. Eicosapentaenoic acid (EPA) does not induce the
production of reactive oxygen species (ROS) in potato leaves.

Supplemental Figure S29. Callose deposition of Arabidopsis seedlings treated with
EPA or Pi-Cer D.

Supplemental Figure S30. Expression profiles of Arabidopsis genes upregulated by co-
treatment of EPA and Pi-Cer D, EPA or Pi-Cer D.

FIGURE LEGENDS
Figure 1 Elicitor activity of methanol extract of *Phytophthora infestans* mycelia (Pi-
MEM) in potato. (A) Potato suspension cultured cells were treated with 0.3% DMSO
(Cont.) or 30 µg/ml Pi-MEM and production of reactive oxygen species (ROS) was
detected as L-012-mediated chemiluminescence 3 h after the treatment. Data are means
± SE (n = 3). (B) Potato leaves were treated with 1 % DMSO (Cont.) or 100 µg/ml Pi-
MEM and production of ROS was detected as L-012-mediated chemiluminescence.
Data are means ± SE (n = 3). Data marked with asterisks are significantly different from
control as assessed by two-tailed Student's *t* tests: **$P < 0.01$. (C) Left, putative
biosynthetic pathway of potato phytoalexins, lubimin, oxylubimin and rishitin. HPS,
Hyoscyamus muticus premnaspirodiene synthase; HPO H. muticus premnaspirodiene
oxygenase; SPH, sesquiterpenoid phytoalexins hydroxylase (Camagna et al., 2020).
Potato tubers were treated with 3 % DMSO (Cont.) or 1 mg/ml Pi-MEM and
phytoalexins were extracted 48 h after treatment. Produced phytoalexins were detected
by thin-layer chromatography (middle) or LC/MS (right).

Figure 2. *Phytophthora infestans* ceramide elicitors (Pi-Cer) can induce the production
of reactive oxygen species (ROS) in potato suspension cultured cells. (A) Potato
suspension cultured cells were treated with 0.3% DMSO (Cont.), 30 µg/ml methanol
extract of P. infestans mycelia (Pi-MEM), 3 µg/ml Pi-Cer A, B, C or D and production
of ROS was detected as L-012 mediated chemiluminescence 3 h after the treatment. (B)
Structures of Pi-Cer A, B, C and D. See Supplemental Figure S1 for the procedures of
purification of elicitors and Supplemental Figures S2-5, S10-15 and supplemental
document for details of their structural analysis.

Figure 3. Pi-Cer D and Pi-CerPE D can induce the production of reactive oxygen
species (ROS) in different plant species.
(A) (Left) Potato leaves were treated with 1% DMSO (Cont.), 100 µg/ml methanol
extract of P. infestans mycelia (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by syringe
infiltration, and production of ROS was detected as L-012 mediated chemiluminescence
3-15 h after the treatment. Data are means ± SE (n = 3).
(B) (Left) Arabidopsis thaliana leaves were treated with 1% DMSO (Cont.), 100 µg/ml
methanol extract of P. infestans (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by
syringe infiltration, and production of ROS was detected as L-012 mediated
chemiluminescence 12 h after the treatment. Data are means ± SE (n = 3).
(C) (Left) Rice leaves were treated with 1% DMSO (Cont.), 100 µg/ml methanol extract
of P. infestans (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by vacuum infiltration, and
production of ROS was detected as L-012 mediated chemiluminescence 12 h after the
treatment. Data are means ± SE (n = 3). Data marked with asterisks are significantly
different from control as assessed by two-tailed Student's *t* tests: **P < 0.01.
695

Figure 4. Pretreatment with Pi-Cer D and Pi-CerPE D enhances the resistance of potato leaves against *P. infestans*. Potato leaves were treated with 0.5 % DMSO (Cont.), 100 µg/ml Pi-MEM, 10 µg/ml Pi-Cer D or Pi-CerPE D and incubated for 24 hrs, and inoculated with a spore suspension of *P. infestans*. (A) Disease symptoms of *P. infestans* on potato leaves treated with DMSO, Pi-MEM, Pi-Cer D or Pi-CerPE D. Photographs were taken 6 days post inoculation. (B) Leaves representative of the disease severities used for classification. (C) Plots showing percentages of potato leaves with disease symptom severities using the classification depicted in (B). Leaves were pretreated with DMSO or elicitors, and disease severity of a subsequent *P. infestans* inoculation was observed from 1 - 7 days post inoculation (dpi). (n = 8). Data marked with asterisks are significantly different from control as assessed by one-tailed Mann–Whitney U-tests: *$P < 0.05$. (D) Left, Penetration sites of *P. infestans* in elicitor treated leaf-discs were detected as callose depositions by aniline blue staining 24 h after inoculation. Bars = 50 µm. Right, Number of fluorescent spots were counted in elicitor-treated leaf discs inoculated with *P. infestans* 24 hrs after inoculation. Data are means ± SE (n = 3). Data marked with asterisks are significantly different from control as assessed by two-tailed Student's *t* tests: $p < 0.01$.

Figure 5. *Phytophthora infestans* diacylglycerol (Pi-DAG) induce the production of phytoalexins in potato tubers. (A) Structures of Pi-DAG A and B. Structures equivalent to eicosapentaenoic acid (EPA) are shown in red, dotted boxes. (B) Structures of Pi-DAG C and D which have significantly weaker elicitor activity compared with Pi-DAG A and B. See Supplemental Figure S22 for the procedures of purification of Pi-DAGs and Supplemental Figures S23-26 for their structural analysis. (C) Potato tubers were treated with 0.3% DMSO (Cont.), 30 µg/ml methanol extract of P. infestans mycelium (Pi-MEM), or 100 µg/ml Pi-DAGs, and production of phytoalexins was detected by thin-layer chromatography. L, Lubimin; R, Rishitin; O, Oxylubimin. (D) EPA, but not Pi-Cer D, can induce the production of phytoalexins in potato tubers. Potato tubers were treated with 0.3% DMSO (Cont.), 100 µg/ml EPA or 100 µg/ml Pi-Cer D and produced phytoalexins were detected by LC/MS.

Figure 6. Comparison of elicitor activity of polyunsaturated fatty acids for induction of rishitin production in potato tubers. (A) Structures of polyunsaturated fatty acids used in this study. Fatty acids in green letters indicate those derived from plant. (B) Potato tubers were treated with indicated polyunsaturated fatty acids (100 µg/ml) and rishitin was extracted 2 days after the treatment. Produced rishitin was quantified by GC/MS (n = 6 for AA and EPA, n = 3 for LA, ALA, GLA, EDA, ETA, MA. n = 12 for DHA).

Figure. 7 Distinctive sets of Arabidopsis genes were upregulated by the treatment with EPA and Pi-Cer D. (A) Arabidopsis transformant pWRKY33-LUC containing LUC transgene under the control of AtWRKY33 (AT2G38470.1) promoter was treated with 100 µg/ml EPA, Pi-Cer D or a mixture of 100 µg/ml EPA and Pi-Cer D. Chemiluminescence was monitored for 12 h after the treatment. (B) Venn diagram representing the up-regulated differentially expressed genes (DEGs) in Arabidopsis treated with 100 µg/ml EPA, 100 µg/ml Pi-Cer D, or a mixture of 100 µg/ml EPA and 100 µg/ml Pi-Cer D for 12 h, which are selected based on TPM ≥ 1, log2 ≥ 2 and p ≤

0.05. (C) Heatmap analysis of differentially expressed genes for the oligosaccharide treatments. The color bar represents the fold change value (log2) with p ≤ 0.05. (D) Expression profile of Arabidopsis genes for capsidiol biosynthesis after treatment with 100 µg/ml EPA, Pi-Cer D or a mixture of 100 µg/ml EPA and Pi-Cer D.

Figure 8. Gene ontology (GO) enrichment analysis of up-regulated genes in Arabidopsis treated with (A) 100 µg/ml eicosapentaenoic acid (EPA), (B) 100 µg/ml Pi-Cer D or (C) their mixculre (100 µg/ml EPA and 100 µg/ml Pi-Cer D) for 12 h. GO term enrichment is expressed as significantly different fold enrichment of mapped genes (FDR p<0.05). The dot size (and numbers beside the dots) indicates the number of significantly up-regulated genes (see Figure 7B) associated with the process and the dot color indicates the significance of the enrichment.

Figure 9. A model of Pattern-triggered immunity (PTI) activation by simultaneous recognition of multiple MAMPs (Microbe-associated molecular patterns) and DAMPs (Damage-associated molecular patterns). Plants secrete enzymes into the apoplast to release MAMPs from the pathogen. 9Me-Spd, 9-methyl-4,8-sphingadienine; 5,8,11,14-TEFA, 5,8,11,14-tetraene-type fatty acid; COS, Cellooligosaccharide; XOS, Xylooligosaccharide; CW, Cell wall; PM, Plasma membrane; CP, Cytoplasm.

REFERENCES

Adarme-Vega TC, Lim DK, Timmins M, Vernen F, Li Y, Schenk PM (2012) Microalgal biofactories: a promising approach towards sustainable omega-3 fatty acid production. Microb Cell Fact. **11: 96.**

Bhat A, Ryu CM. (2016) Plant perceptions of extracellular DNA and RNA. Mol Plant **9:** 956–958

Bostock RM, Kuć JA, Laine RA (1981) Eicosapentaenoic and arachidonic acids from *Phytophthora infestans* elicit fungitoxic sesquiterpenes in the potato. Science **212:** 67–69

Bostock RM, Savchenko T, Lazarus C, Dehesh K (2011) Eicosapolyenoic acids: novel MAMPs with reciprocal effect on oomycete-plant defense signaling networks. Plant Signal Behav **6:** 531–533

Cahoon EB, Li-Beisson Y (2020) Plant unusual fatty acids: learning from the less common. Curr Opin Plant Biol **55:** 66–73

Camagna M, Ojika M, Takemoto D (2020) Detoxification of the solanaceous phytoalexins rishitin, lubimin, oxylubimin and solavetivone via a cytochrome P450 oxygenase. Plant Signal Behav **15:** 1707348

Ellinger D, Naumann M, Falter C, Zwikowics C, Jamrow T, Manisseri C, Somerville SC, Voigt CA (2013) Elevated early callose deposition results in complete penetration resistance to powdery mildew in Arabidopsis. Plant Physiol **161:** 1433–1444

Fernandes BS, Dias O, Costa G, Kaupert Neto AA, Resende TFC, Oliveira JVC, Riaño-Pachón DM, Zaiat M, Pradella JGC, Rocha I (2019) Genome-wide

sequencing and metabolic annotation of *Pythium irregulare* CBS 494.86:
understanding Eicosapentaenoic acid production. BMC Biotechnol **19:** 41.

Forbes GA (2012) Using host resistance to manage potato late blight with particular
reference to developing countries. Potato Res **55:** 205–216

Free SJ (2013) Fungal cell wall organization and biosynthesis. Adv Genet **81:** 33–82

Fry W (2008) *Phytophthora infestans*: the plant (and *R* gene) destroyer. Mol Plant
Pathol **9:** 385–402

Garelik G (2002) Agriculture. Taking the bite out of potato blight. Science **298:**1702–
1704

Gaulin E, Bottin A, Dumas B (2010) Sterol biosynthesis in oomycete pathogens. Plant
Signal Behav 5: 258–260

Harwood JL (1988) Fatty acid metabolism. Annu Rev Plant Physiol Plant Mol Biol **39:**
101–138

Hayashi F, Smith KD, Ozinsky A, Hawn TR, Yi EC, Goodlett DR, Eng JK, Akira S,
Underhill DM, Aderem A. (2001) The innate immune response to bacterial flagellin
is mediated by Toll-like receptor 5. Nature **410:** 1099–1103

He X, Howard BA, Liu Y, Neumann AK, Li L, Menon N, Roach T, Kale SD,
Samuels DC, Li H, Kite T, Kita H, Hu TY, Luo M, Jones CN, Okaa UJ,
Squillace DL, Klein BS, Lawrence CB (2021) LYSMD3: A mammalian pattern
recognition receptor for chitin. Cell Rep **36:** 109392

Imano S, Fushimi M, Camagna M, Tsuyama-Koike A, Mori H, Ashida A, Tanaka A,
Sato I, Chiba S, Kawakita K, Ojika M, Takemoto D (2022) AP2/ERF
transcription factor NbERF-IX-33 is involved in the regulation of phytoalexin
production for the resistance of *Nicotiana benthamiana* to *Phytophthora infestans*.
Front Plant Sci **12:** 821574

Ishida N, Akai S (1969) Relation of temperature to germination of conidia and
appressorium formation in *Colletotrichum lagenarium*. Mycologia **61:** 382–386

Jiang C, Ge J, He B, Zeng B (2021) Glycosphingolipids in filamentous fungi:
Biological roles and potential applications in cosmetics and health foods. Front
Microbiol **12:** 690211

Jones DA, Takemoto D. (2004) Plant innate immunity - direct and indirect recognition
of general and specific pathogen-associated molecules. Curr Opin Immunol **16:** 48–
62

Kamoun S, Furzer O, Jones JD, Judelson HS, Ali GS, Dalio RJ, Roy SG, Schena L,
Zambounis A, Panabières F, Cahill D, Ruocco M, Figueiredo A, Chen XR,
Hulvey J, Stam R, Lamour K, Gijzen M, Tyler BM, Grünwald NJ, Mukhtar MS,
Tomé DF, Tör M, Van Den Ackerveken G, McDowell J, Daayf F, Fry WE,
Lindqvist-Kreuze H, Meijer HJ, Petre B, Ristaino J, Yoshida K, Birch PR,
Govers F (2015) The top 10 oomycete pathogens in molecular plant pathology. Mol
Plant Pathol **16:** 413–434

Kato H, Nemoto K, Shimizu M, Abe A, Asai S, Ishihama N, Matsuoka S, Daimon T,
Ojika M, Kawakita K, Onai K, Shirasu K, Yoshida M, Ishiura M, Takemoto D,
Takano Y, Terauchi R. (2022). Recognition of pathogen-derived sphingolipids in
Arabidopsis. Science **376:** 857–860

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol **37:**
907–915

Koga J, Yamauchi T, Shimura M, Ogawa N, Oshima K, Umemura K, Kikuchi M,
Ogasawara N (1998) Cerebrosides A and C, sphingolipid elicitors of hypersensitive
cell death and phytoalexin accumulation in rice plants. J Biol Chem **273:** 31985–
31991

Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M (2019)
Transcriptome assembly from long-read RNA-seq alignments with StringTie2.
Genome Biol **20:** 278

Kunze G, Zipfel C, Robatzek S, Niehaus K, Boller T, Felix G (2004) The N terminus
of bacterial elongation factor Tu elicits innate immunity in Arabidopsis plants. Plant
Cell **16:** 3496–3507

Kuroyanagi T, Bulasag AS, Fukushima K, Ashida A, Suzuki T, Tanaka A,
Camagna M, Sato I, Chiba S, Ojika M, Takemoto D (2022) *Botrytis cinerea*
identifies host plants via the recognition of antifungal capsidiol to induce expression
of a specific detoxification gene. PNAS Nexus **1:** pgac274

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol **15:** 550

Martin M (2011) Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet J **17:** 10–12

Martín-Dacal M, Fernández-Calvo P, Jiménez-Sandoval P, López G, Garrido-
Arandía M, Rebaque D, Del Hierro I, Berlanga DJ, Torres MÁ, Kumar V,
Mélida H, Pacios LF, Santiago J, Molina A (2023) Arabidopsis immune responses
triggered by cellulose- and mixed-linked glucan-derived oligosaccharides require a
group of leucine-rich repeat malectin receptor kinases. Plant J. **113:** 833–850

Mélida H, Sandoval-Sierra JV, Diéguez-Uribeondo J, Bulone V (2013) Analyses of
extracellular carbohydrates in oomycetes unveil the existence of three different cell
wall types. Eukaryot Cell **12:** 194–203

Monjil MS, Nozawa T, Shibata Y, Takemoto D, Ojika M, Kawakita K (2015)
Methanol extract of mycelia from *Phytophthora infestans*-induced resistance in
potato. C R Biol **338:** 185–196

Monjil MS, Kato H, Matsuda K, Suzuki N, Tenhiro S, Camagna M, Suzuki T,
Tanaka A, Terauchi R, Sato I, Chiba S, Kawakita K, Ojika M and Takemoto D
(2021) Two structurally different oomycete MAMPs induce distinctive plant immune
responses. bioRxiv 2021.10.22.465218 [Preprint]

Moreau RA, Young DH, Danis PO, Powell MJ, Quinn CJ, Beshah K, Slawecki RA,
Dilliplane RL (1998) Identification of ceramide-phosphorylethanolamine in
oomycete plant pathogens: *Pythium ultimum*, *Phytophthora infestans*, and
Phytophthora capsici. Lipids **33:** 307–317

Namiki F, Shiomi T, Nishi K, Kayamura T, Tsuge T (1998) Pathogenic and genetic
variation in the Japanese strains of *Fusarium oxysporum* f. sp. *melonis*.
Phytopathology **88:** 804–810

Ngou BPM, Ding P and Jones JDG (2022) Thirty years of resistance: Zig-zag through
the plant immune system. Plant Cell **34:** 1447–1478

Noritake T, Kawakita K, Doke N (1996) Nitric oxide induces phytoalexin accumulation in potato tuber tissues. Plant Cell Physiol **37:** 113–116

Nykiforuk CL, Shewmaker C, Harry I, Yurchenko OP, Zhang M, Reed C, Oinam GS, Zaplachinski S, Fidantsef A, Boothe JG, Moloney MM (2012) High level accumulation of gamma linolenic acid (C18:3Δ6.9,12 cis) in transgenic safflower (*Carthamus tinctorius*) seeds. Transgenic Res 21: 367–381

Pring S, Kato H, Imano S, Camagna M, Tanaka A, Kimoto H, Chen P, Shrotri A, Kobayashi H, Fukuoka A, Saito M, Suzuki T, Terauchi R, Sato I, Chiba S, Takemoto D (2023) Induction of plant disease resistance by mixed oligosaccharide elicitors prepared from plant cell wall and crustacean shells. Physiol Plant. **175:** e14052

Ranf S (2017) Sensing of molecular patterns through cell surface immune receptors. Curr Opin Plant Biol **38:** 68–77

Shibata Y, Kawakita K, Takemoto D (2010) Age-related resistance of *Nicotiana benthamiana* against hemibiotrophic pathogen *Phytophthora infestans* requires both ethylene- and salicylic acid-mediated signaling pathways. Mol Plant-Microbe Interact **23:** 1130–1142

Shibata Y, Kawakita K, Takemoto D (2011) SGT1 and HSP90 are essential for age-related non-host resistance of *Nicotiana benthamiana* against the oomycete pathogen *Phytophthora infestans*. Physiol Mol Plant Pathol 75:120–128

Shibata Y, Ojika M, Sugiyama A, Yazaki K, Jones DA, Kawakita K, Takemoto D (2016) The full-size ABCG transporters Nb-ABCG1 and Nb-ABCG2 function in pre- and postinvasion defense against *Phytophthora infestans* in *Nicotiana benthamiana*. Plant Cell **28:** 1163–1181

Sperling P, Heinz E (2003) Plant sphingolipids: structural diversity, biosynthesis, first genes and functions. Biochim Biophys Acta **1632:** 1–15

Pring S, Kato H, Imano S, Camagna M, Tanaka A, Kimoto H, Chen P, Shrotri A, Kobayashi H, Fukuoka A, Saito M, Suzuki T, Terauchi R, Sato I, Chiba S, Takemoto D. (2023) Induction of plant disease resistance by mixed oligosaccharide elicitors prepared from plant cell wall and crustacean shells. Physiol Plant **175:** e14052.

Takeuchi O, Hoshino K, Kawai T, Sanjo H, Takada H, Ogawa T, Takeda K, Akira S. (1999) Differential roles of TLR2 and TLR4 in recognition of gram-negative and gram-positive bacterial cell wall components. Immunity **11:** 443–451

Thines M, Kamoun S (2010) Oomycete-plant coevolution: recent advances and future prospects. Curr Opin Plant Biol **13:** 427–433

Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou LP, Mi H. (2022) PANTHER: Making genome-scale phylogenetics accessible to all. Protein Sci **31:** 8–22

Umemura K, Ogawa N, Yamauchi T, Iwata M, Shimura M, Koga J (2000) Cerebroside elicitors found in diverse phytopathogens activate defense responses in rice plants. Plant Cell Physiol **41:** 676–83

Waskom ML (2021) Seaborn: statistical data visualization. J. Open Res. Softw **6:** 3021

Zheng Z, Qamar SA, Chen Z, Mengiste T (2006) Arabidopsis WRKY33 transcription factor is required for resistance to necrotrophic fungal pathogens. Plant J **48:** 592–605

Figure 1. Elicitor activity of methanol extract of *Phytophthora infestans* mycelia (Pi-MEM) in potato.
(A) Potato suspension cultured cells were treated with 0.3% DMSO (Cont.) or 30 µg/ml Pi-MEM and production of reactive oxygen species (ROS) was detected as L-012-mediated chemiluminescence 3 h after the treatment. Data are means ± SE (n = 3). **(B)** Potato leaves were treated with 1 % DMSO (Cont.) or 100 µg/ml Pi-MEM and production of ROS was detected as L-012-mediated chemiluminescence. Data are means ± SE (n = 3). Data marked with asterisks are significantly different from control as assessed by two-tailed Student's *t* tests: **P < 0.01.
(C) Left, putative biosynthetic pathway of potato phytoalexins, lubimin, oxylubimin and rishitin. HPS, *Hyoscyamus muticus* premnaspirodiene synthase; HPO *H. muticus* premnaspirodiene oxygenase; SPH, sesquiterpenoid phytoalexins hydroxylase (Camagna et al., 2020). Potato tubers were treated with 3 % DMSO (Cont.) or 1 mg/ml Pi-MEM and phytoalexins were extracted 48 h after treatment. Produced phytoalexins were detected by thin-layer chromatography (middle) or LC/MS (right).

Figure 2. *Phytophthora infestans* ceramide elicitors (Pi-Cer) can induce the production of reactive oxygen species (ROS) in potato suspension cultured cells. **(A)** Potato suspension cultured cells were treated with 0.3% DMSO (Cont.), 30 µg/ml methanol extract of *P. infestans* mycelia (PiMEM), 3 µg/ml Pi-Cer A, B, C or D and production of ROS was detected as L-012 mediated chemiluminescence 3 h after the treatment. **(B)** Structures of Pi-Cer A, B, C and D. See Supplemental Figure S1 for the procedures of purification of elicitors and Supplemental Figures S2-5, S10-15 and supplemental document for details of their structural analysis.

Figure 3. Pi-Cer D and Pi-CerPE D can induce the production of reactive oxygen species (ROS) in different plant species.

(A) (Left) Potato leaves were treated with 1% DMSO (Cont.), 100 µg/ml methanol extract of *P. infestans* mycelia (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by syringe infiltration, and production of ROS was detected as L-012 mediated chemiluminescence 3-15 h after the treatment. Data are means ± SE (n = 3).

(B) (Left) *Arabidopsis thaliana* leaves were treated with 1% DMSO (Cont.), 100 µg/ml methanol extract of *P. infestans* (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by syringe infiltration, and production of ROS was detected as L-012 mediated chemiluminescence 12 h after the treatment. Data are means ± SE (n = 3).

(C) (Left) Rice leaves were treated with 1% DMSO (Cont.), 100 µg/ml methanol extract of *P. infestans* (Pi-MEM), 10 µg/ml Pi-Cer D or Pi-CerPE D by vacuum infiltration, and production of ROS was detected as L-012 mediated chemiluminescence 12 h after the treatment. Data are means ± SE (n = 3). Data marked with asterisks are significantly different from control as assessed by two-tailed Student's *t* tests: **P < 0.01.

Figure 4. Pretreatment with Pi-Cer D and Pi-CerPE D enhances the resistance of potato leaves against *P. infestans*.

Potato leaves were treated with 0.5 % DMSO (control), 100 µg/ml Pi-MEM, 10 µg/ml Pi-Cer D or Pi-CerPE D and incubated for 24 hrs, and inoculated with a spore suspension of *P. infestans*. **(A)** Disease symptoms of *P. infestans* on potato leaves treated with DMSO, Pi-MEM, Pi-Cer D or Pi-CerPE D. Photographs were taken 6 days post inoculation. **(B)** Leaves representative of the disease severities used for classification. **(C)** Plots showing percentages of potato leaves with disease symptom severities using the classification depicted in (B). Leaves were pretreated with DMSO or elicitors, and disease severity of a subsequent *P. infestans* inoculation was observed from 1 - 7 days post inoculation (dpi). (n = 8). Data marked with asterisks are significantly different from control as assessed by one-tailed Mann–Whitney U-tests: *P < 0.05. **(D)** Left, Penetration sites of *P. infestans* in elicitor treated leaf-discs were detected as callose depositions by aniline blue staining 24 h after inoculation. Bars = 50 µm. Right, Number of fluorescent spots were counted in elicitor-treated leaf discs inoculated with *P. infestans* 24 hrs after inoculation. Data are means ± SE (n = 3). Data marked with asterisks are significantly different from control as assessed by two-tailed Student's *t* tests: p < 0.01.

Figure 5. *Phytophthora infestans* diacylglycerol (Pi-DAG) induce the production of phytoalexins in potato tubers. **(A)** Structures of Pi-DAG A and B. Structures equivalent to eicosapentaenoic acid (EPA) are shown in red, dotted boxes. **(B)** Structures of Pi-DAG C and D which have significantly weaker elicitor activity compared with Pi-DAG A and B. See Supplemental Figure S22 for the procedures of purification of Pi-DAGs and Supplemental Figures S23-26 for their structural analysis. **(C)** Potato tubers were treated with 0.3% DMSO (Cont.), 30 µg/ml methanol extract of *P. infestans* mycelium (Pi-MEM), or 100 µg/ml Pi-DAGs, and production of phytoalexins was detected by thin-layer chromatography. L, Lubimin; R, Rishitin; O, Oxylubimin. **(D)** EPA, but not Pi-Cer D, can induce the production of phytoalexins in potato tubers. Potato tubers were treated with 0.3% DMSO (Cont.), 100 µg/ml EPA or 100 µg/ml Pi-Cer D and produced phytoalexins were detected by LC/MS.

Figure 6. Comparison of elicitor activity of polyunsaturated fatty acids for induction of rishitin production in potato tubers. **(A)** Structures of polyunsaturated fatty acids used in this study. Fatty acids in green letters indicate those derived from plant. **(B)** Potato tubers were treated with indicated polyunsaturated fatty acids (100 µg/ml) and rishitin was extracted 2 days after the treatment. Produced rishitin was quantified by GC/MS (n = 6 for AA and EPA, n = 3 for LA, ALA, GLA, EDA, ETA, MA. n=12 for DHA).

Figure. 7 Distinctive sets of Arabidopsis genes were upregulated by the treatment with EPA and Pi-Cer D. **(A)** Arabidopsis transformant pWRKY33-LUC containing *LUC* transgene under the control of *AtWRKY33* (AT2G38470.1) promoter was treated with 100 µg/ml EPA, Pi-Cer D or a mixture of 100 µg/ml EPA and Pi-Cer D. Chemiluminescence was monitored for 12 h after the treatment. (B) Venn diagram representing the upregulated differentially expressed genes (DEGs) in Arabidopsis treated with 100 µg/ml EPA, 100 µg/ml Pi-Cer D, or a mixture of 100 µg/ml EPA and 100 µg/ml Pi-Cer D for 12 h, which are selected based on TPM ≥ 1, log2 ≥ 2 and p ≤ 0.05. (C) Heatmap analysis of differentially expressed genes for the oligosaccharide treatments. The color bar represents the fold change value (log2) with p ≤ 0.05. (D) Expression profile of Arabidopsis genes for capsidiol biosynthesis after treatment with 100 µg/ml EPA, Pi-Cer D or a mixture of 100 µg/ml EPA and Pi-Cer D.

Figure 8. Gene ontology (GO) enrichment analysis of up-regulated genes in Arabidopsis treated with 100 μg/ml EPA, 100 μg/ml Pi-Cer D or their mixculre (100 μg/ml EPA and 100 μg/ml Pi-Cer D) for 12 h. GO term enrichment is expressed as significantly different fold enrichment of mapped genes (FDR p<0.05). The dot size (and numbers beside the dots) indicates the number of significantly up-regulated genes (see Figure 7B) associated with the process and the dot color indicates the significance of the enrichment.

Figure 9. A model of Pattern-triggered immunity (PTI) activation by simultaneous recognition of multiple MAMPs (Microbe-associated molecular patterns) and DAMPs (Damage-associated molecular patterns). Plant cells secrete enzymes into the apoplast to release MAMPs from the pathogen. 9Me-Spd, 9-methyl-4,8-sphingadienine; 5,8,11,14-TEFA, 5,8,11,14-tetraene-type fatty acid; COS, Cello-oligosaccharide; XOS, Xylo-oligosaccharide; CW, Cell wall; PM, Plasma membrane; CP, Cytoplasm.