Clostridioides difficile **infection increases circulating p-cresol levels and dysregulates brain**

dopamine metabolism: linking gut-brain axis to autism spectrum disorders?

Akhil A. Vinithakumari[1], Piyush Padhi[2], Belen Hernandez[1], Susanne Je-Han Lin[1], Aaron

Dunkerson-Kurzhumov[1], Lucas Showman[3], Matthew Breitzman[3], Caroline Stokes[1], Yousuf

Sulaiman[1], Chandra Tangudu[4], Deepa Ashwarya Kuttappan[5], Muhammed Shafeekh

Muyyarikkandy[5], Gregory Phillips[4], Vellareddy Anantharam[6], Ann Perera[3], Brett Sponseller[4],

Anumantha Kanthasamy[6], and Shankumar Mooyottu[1]*

[1]*Department of Veterinary Pathology, Iowa State University, Ames, IA, 50011*

[2]*Department of Veterinary Biomedical Sciences, Iowa State University, Ames, IA, 50011*

[3]*W.M. Keck Metabolomics Research Laboratory, Iowa State University, Ames, IA, 50011*

[4]*Department of Veterinary Microbiology and Preventive Medicine, Iowa State University, Ames,*

IA, 50011

[5]*Department of Animal Science, University of Connecticut, Storrs, CT, 06267*

[6]*Department of Biomedical Sciences, University of Georgia, Athens, GA, 30602*

*Corresponding author

Shankumar Mooyottu

shaan@iastate.edu

515-294-3877

Abstract

Gastrointestinal illnesses are one of the most common comorbidities reported in patients with neurodevelopmental diseases, including autism spectrum disorders (ASD). Gut dysbiosis, overgrowth of *C. difficile*, and gut microbiota-associated alterations in central neurotransmission have been implicated in ASD, where the dopaminergic axis plays an important role in the disease pathogenesis. Human *C. difficile* strains produce a significant amount of the toxic metabolite p-cresol, an inhibitor of dopamine beta-hydroxylase (DBH), which catalyzes the conversion of dopamine (DA) to norepinephrine (NE). p-Cresol is known to precipitate and exacerbate autistic behavior in rodents by increasing DA levels and altering DA receptor sensitivity in brain regions relevant to ASD. Therefore, we hypothesized that *C. difficile* infection dysregulates dopaminergic metabolism by increasing p-cresol levels in the gut and systemic circulation, and by inhibiting brain DBH, ultimately leading to elevated DA in different brain regions. For testing this hypothesis, we induced antibiotic-associated *C. difficile* infection in mice and determined the gut and serum p-cresol levels, serum DBH activity, and dopamine and its metabolite levels in different brain regions relevant to ASD. The results showed that *C. difficile* infection causes a significant increase in striatal DA, accompanied by significantly altered levels of DA metabolites and NE in different brain regions ($p < 0.05$). In addition, significantly increased circulating p-cresol levels and reduced DBH activity were observed in *C. difficile* infected mice ($p < 0.05$). Therefore, the results from this study suggest a potential link between *C. difficile* infection and alterations in the dopaminergic axis implicated in the precipitation and aggravation of ASD.

Keywords: *C. difficile*, gut-brain axis, p-cresol, gut dysbiosis

Introduction

Although the involvement of the gut-brain axis is increasingly implicated in the precipitation and manifestation of neurodevelopmental disorders such as autism, mechanistic evidence that directly connects gut bacteria, bacterial metabolites, and neuronal dysregulation is sparse.[1, 2] Disruption of the normal gut microbiota is known to cause an overgrowth of unfavorable bacterial communities that increase toxic metabolites in the gut.[3] Several of these metabolites are known to impact neurotransmission and brain functions, underscoring the importance of the gut-brain axis in health and disease.[4]

p-Cresol is a toxic phenolic compound produced by bacterial fermentation of protein in the human large intestine.[5] p-Cresol is readily absorbed into the bloodstream and secreted in urine.[6, 7] Moreover, it is already known that patients with autism spectrum disorders (ASD) often have a significantly higher urine p-cresol level, which is considered a biomarker for this disease complex.[8-10] A recent study demonstrated that intravenous p-cresol injection aggravates autistic behaviors in a murine genetic model of ASD by dysregulating the dopaminergic axis.[11] Another study revealed ASD-like behavior in rats intraperitoneally administered with p-cresol.[12] Similar results with dysregulation in dopaminergic neurotransmission were demonstrated following oral p-cresol administration in non-mutant mice.[13] Therefore, p-cresol mediated alterations in the dopaminergic pathway appear to be associated with the pathophysiology of ASD.

p-Cresol is a potent inhibitor of the enzyme dopamine beta-hydroxylase (DBH) that converts dopamine (DA) to norepinephrine (NE).[14-16] Thus, inhibition of DBH increases the concentration of DA in the central nervous system. Notably, a low maternal serum level of DBH and mutations in the DBH genes have been associated with autism in children.[17, 18] Additionally, increased urinary and cerebrospinal fluid levels of homovanillic acid, a DA metabolite, have

been reported in autistic patients.[19-21] Moreover, an elevated brain DA level, mutations in the DA

transporter protein, altered sensitivity of DA receptors, and impaired striatal dopaminergic

neurotransmission are observed in autistic patients and animal models of ASD.[22-25] In addition,

DA receptor antagonists are found to reduce autistic symptoms in patients.[25, 26]

Gut dysbiosis induced by prolonged antibiotic therapy is the most common cause of *C.*

difficile overgrowth and associated diarrhea in infants and adults.[27, 28] Human *C. difficile* strains

produce a significant amount of p-cresol, unlike most other gut bacteria.[29-31] *C. difficile* produces

10 to 1000 times more p-cresol than other known p-cresol producing bacteria in the gut.[5, 32]

Indeed, recent studies demonstrated p-cresol induced autistic behavior in mice by altering the

dopaminergic axis.[11, 13] What remains unknown specifically is the p-cresol mediated functional

consequence of *C. difficile* infection on the dopaminergic neurotransmission from a gut-brain

axis standpoint. Moreover, the involvement of possible mediators, such as DBH, that directly

connects gut microbial p-cresol to DA dysmetabolism needs to be demonstrated. Thus, the

current work investigates the effect of antibiotic-induced *C. difficile* infection and consequent *in*

vivo production of p-cresol on the DA metabolism in the brain.

Results

C. difficile UK1 produces a significant amount of p-cresol in vitro

To determine the p-cresol producing ability of the human hypervirulent *C. difficile* strain

(UK1), we performed a p-HPA-p-cresol conversion assay. For this experiment, we quantified

concentrations of total p-cresol and its substrate para-hydroxyphenyl acetic acid (p-HPA)

concentrations at different time points in *C. difficile* UK1 cultures by high-performance liquid

chromatography (HPLC) *in vitro*. Our results indicated that *C. difficile* UK1 completely

metabolized p-HPA to p-cresol in brain heart infusion (BHI) media by 24h of incubation in

anaerobic conditions **(Figure 1)**. This data confirm the p-cresol producing capability of human

hypervirulent *C. difficile* strain, suggesting that this bacterium could increase the gut p-cresol

concentrations in *C. difficile* infected host.

Antibiotic induced C. difficile infection increases gut p-cresol concentration

In order to quantitate *in vivo C. difficile* p-cresol production in the murine gut, we

performed a *C. difficile* challenge study. For this experiment, 3-week-old C57BL/6 mice were

administered with an oral antibiotic cocktail followed by an intraperitoneal clindamycin injection

in order to induce gut dysbiosis. Gut dysbiosis is required for *C. difficile* colonization in mice as

in humans.[33, 34] Mice were then challenged with 10^4 *C. difficile* UK1 spores **(Figure 2A)**. From

the previous experiments conducted by the authors and other investigators, it is known that 10^4

C. difficile UK1 spores reliably induces clinical CDI with less mortality in mice compared to

high dose (10^6 to 10^7 spores) challenge, simulating chronic persistent CDI in humans. Diarrhea

or wet tail were noticed in all *C. difficile* challenged mice on day-2 post-inoculation.

Histopathologic examination of the colonic sections revealed no significant lesions in the colon

of the control and antibiotic-control (Abx) groups **(Figure 2B)**. As expected, the colon of the

mice challenged with 10^4 *C. difficile* UK1 (C.Diff) exhibited significant pathologic changes in

the mucosal epithelium, as evidenced by necrosis, moderate neutrophilic inflammatory infiltrates

in the lamina propria and submucosa and marked submucosal edema, contributing to a

significantly increased cumulative colitis score **(Figure 2B & 2C)**. In addition, the total cecal p-

cresol level (free and bound) on day-2 post-challenge was determined using HPLC method

(Figure 2D).[35] A significant increase in total cecal p-cresol concentration was observed in C.

Diff group compared to controls ($p < 0.05$). An increasing trend in the cecal p-cresol, although

not statistically significant was observed in Abx group compared to control **(Figure 2D)**. Thus,

the data indicate that clinical *C. difficile* infection significantly increases gut p-cresol levels in

infected mice.

C. difficile infection increases serum p-cresol and reduces serum DBH activity

Since p-cresol produced in the intestine is readily absorbed into the blood stream, we

investigated the changes in the serum p-cresol levels in *C. difficile* challenged and control

groups. Previous studies demonstrated that p -cresol is a neurotoxin that can cross the blood

brain barrier, impair neurotransmission, and induce behavioral abnormalities.[12] In addition, p-

cresol is a potent inhibitor of DBH enzyme.[14-16] It is also known that the serum DBH is

predominantly contributed by the brain tissue, and hence the serum DBH activity meaningfully

reflects the brain DBH activity.[36, 37] Therefore, we quantitated serum p-cresol levels in *C.*

difficile challenged and control mice. Furthermore, we performed a DBH activity assay to

determine whether *C. difficile* induced changes in the circulating p-cresol inhibits the activity of

this enzyme, which may reduce DA to NE conversion in brain. As expected, a significant

increase in total serum p-cresol concentration was observed in C. Diff group compared to

controls ($p < 0.05$). Minor increases in the serum p-cresol (not significant) were observed in Abx

group compared to control **(Figure 3A).** Most importantly, a significant reduction in the serum

DBH activity ($p < 0.05$) was observed in *C. difficile* challenged mice as indicated by the lower

response in LC-MS detection of the reaction product (octopamine) compared to controls.

(Figure 3B). Thus, the results from this set of experiments demonstrates a significant increase in

serum p-cresol concentration and diminished DBH activity in *C. difficile* infected mice.

C. difficile infection increased DA and its metabolite levels in corpus striatum and prefrontal

cortex

Increased gut and serum concentrations of p-cresol have been demonstrated to alter

dopaminergic neurotransmission, which has been implicated in neurological disorders, including

autism.[11-13] Two major dopaminergic pathways in brain significant to ASD include the

mesocorticolimbic (MCL) circuit, and the nigrostriatal (NS) circuit.[38, 39] Substantia nigra and the

ventral tegmental area are the two major DA producing areas in the brain. Neurons from the

substantia nigra project to the dorsal striatum, forming the nigrostriatal (NS) circuit, which is

associated with the motor aspects of goal-directed behavior. The ventral tegmental area projects

to the prefrontal cortex and the ventral striatum forming the mesocorticolimbic (MCL) circuit,

which is involved in reward and motivation-related behavior. Dysregulation of the

mesocorticolimbic circuit leads to social deficits, and malfunction of the nigrostriatal circuit

leads to stereotyped behaviors seen in ASD.[39] Striatum is a critical area that receives projections

from both systems; and therefore, DA and its metabolite concentrations were examined in the

striatum in response to antibiotic treatment and *C. difficile* infection.

The results from the neurochemical analysis for striatal catecholamines **(Figure 4)**

showed a significantly increased DA concentration in the C.Diff group compared to the untreated

control (p < 0.05). However, concentrations of DA metabolic products 3,4-

dihydroxyphenylacetic acid (DOPAC) and homovanillic acid (HVA) did not significantly differ

from that of control animals. Similarly, no significant difference was observed in the NE

concentration in the striatum of *C. difficile* challenged mice compared to the negative control.

DA concentrations were unaltered or slightly reduced in the antibiotic control group (Abx)

compared to the untreated control. In contrast, the significantly lower DOPAC and HVA

concentrations (p < 0.05) were observed in Abx group compared to the negative control. No

statistically significant difference was observed in the NE levels in Abx compared to the negative control, although a decreasing trend was noted.

A comparable, similar trend was observed in the catecholamine concentrations in the prefrontal cortex **(Figure 5)**. An increasing trend in DA and DOPAC concentrations and a statistically significant ($p < 0.05$) increase in HVA in the *C. difficile* challenged mice were observed when compared to both negative and antibiotic control groups. Interestingly, although statistically not significant, a decreasing trend of NE concentration was observed in C.Diff group compared to both control groups.

Both antibiotic treatment and antibiotic induced* C. difficile *infection decreased NE levels in hippocampus

In contrast to the striatal and prefrontal cortex, the neurochemical profile of catecholamines in the hippocampus revealed significant reduction of the DA degradation product DOPAC ($p<0.05$) in C.Diff group, accompanied with a statistically nonsignificant, but decreasing trend in DA concentration compared to the control group **(Figure 6)**. More interestingly, the NE levels in both antibiotic control groups and *C. difficile* challenged group were significantly reduced in the hippocampus ($p < 0.05$ and $p < 0.001$ respectively). This result suggests hippocampal DA release in response to *C. difficile* infection is distinctively different compared to striatum and prefrontal cortex, where dopaminergic circuits predominate.

Antibiotic treatment, and antibiotic induced C. difficile infection differentially altered DA levels in substantia nigra

Substantia nigra is one of the major locations where neuronal DA is synthesized to be released to brain areas, such as striatum and prefrontal cortex through dopaminergic projections.

Although not statistically significant, a decreasing trend in concentrations of DA and its

degradation products were observed in the substantia nigra of *C. difficile* challenged mice

compared to negative control **(Figure 7)**. In contrast, the concentrations of DA and DOPAC

were significantly reduced in *C. difficile* challenged mice compared to the antibiotic control

group (Abx) ($p < 0.05$). The same trend was also observed in HVA concentration. Moreover, the

NE levels did not differ significantly across the treatment groups. The result suggests a reduced

DA synthesis in substantia nigra in *C. difficile* infected animals (C.Diff group) but not in

antibiotic (alone) treated animals (Abx group) compared to control, which will be further

discussed in the following section.

Discussion

Gastrointestinal disorders are one of the most common comorbidities reported in patients

with ASD.[40-42] Of particular note, gut dysbiosis, overgrowth of *C. difficile* in the gut, and gut

microbiota-associated alterations in central neurotransmission have been implicated in ASD,

where the dopaminergic axis plays an important role in the disease pathogenesis.[43] In addition, a

significant association has been demonstrated between antibiotic therapy and resultant alterations

in the gut microbiome (gut dysbiosis)-which predispose *C. difficile* overgrowth in gut in early

infancy contributing to ASD in children, although causation has not been proven.[44] At this point,

a specific underlying mechanism that connects gut dysbiosis to ASD-associated changes in the

brain has not been identified.

In children and adults with ASD, gastrointestinal symptoms are a common problem

frequently associated with dysbiosis and a higher incidence of *C. difficile* colonization[45, 46]. The

severity of ASD has been strongly correlated with the severity gastrointestinal symptoms in

autistic individuals[46]. Strikingly, fecal microbiome therapy (FMT), a radical treatment strategy

against *C. difficile* infection in humans, is also found to be effective in mitigating ASD

symptoms in both adults and children[47]. Additionally, transplantation of the fecal microbiome

from ASD patients induced autism-like behavior in mice[48]. Moreover, vancomycin, a primary

line of antibiotics used to treat *C. difficile* infection in humans, reduces autistic symptoms in

affected individuals[49]. In contrast, fluoroquinolones, a group of antibiotic that predispose to the

development of *C. difficile* infection in humans, aggravate ASD symptoms in autistic patients[49].

Thus, it appears that the factors associated with the predisposition, manifestation, and

suppression of autistic symptoms coincide with that of *C. difficile* infection in affected

individuals. Therefore, we hypothesized that gut-dysbiosis and dysbiosis-induced *C. difficile*

overgrowth and subsequent production of p-cresol dysregulate the dopaminergic axis in the

brain, which may have implications in neurodevelopmental disorders such as ASD.

As expected, the *in vitro* p-cresol production experiment showed that *C. difficile* UK1

strain efficiently converts the substrate p-HPA to p-cresol (**Figure 1**). Production of a very high

amount of p-cresol is an unusual property of *C. difficil*e compared to other gut bacteria.[50] *C.*

difficile ferments tyrosine and its metabolites such as p-HPA to synthesize p-cresol in the gut. p-

Cresol is a bacteriostatic compound that impairs the membrane integrity of Gram-negative

bacteria, which provides a competitive advantage to *C. difficile* over other bacteria, including

Escherichia and *Bacteroides sp.*[51] Similarly, the results from our infection trial indicated a

significant increase in p-cresol concentration in the cecal contents of *C. difficile* infected mice

compared to controls ($p < 0.05$) (**Figure 2D**). It is known that p-cresol is readily absorbed into

the bloodstream, as indicated by a significant increase in serum p-cresol levels in *C. difficile*

infected animals compared to controls ($p < 0.05$). Although not statistically significant, an

increased cecal p-cresol level was observed in the antibiotic control group (Abx). Previous

reports, including our studies have already established that the antibiotic treatment (oral

antibiotic cocktail and intraperitoneal clindamycin, in this case) causes significant dysbiosis in

the murine gut, causing overgrowth of unfavorable bacterial communities that produce toxic

metabolites, including p-cresol.[52, 53] As expected, a similar trend was also observed in serum p-

cresol levels of the antibiotic control mice (Abx group). These observations indicate that

antibiotic-induced *C. difficile* infection increases serum levels of p-cresol, which may have

significant extraintestinal, organ specific or systemic effects in the infected patient. Since p-

cresol can cross the blood-brain barrier, it is reasonable to expect an impact on the structural and

functional homeostasis of the central nervous system.

The enzymatic assay performed on the serum samples of *C. difficile* infected and control

mice **(Figure 3B)** demonstrated a significant reduction in the DBH activity coinciding with the

increased serum p-cresol levels. The brain is the primary source of the serum levels of DBH, and

either serum or plasma DBH activity has been considered as one of the reliable measures of brain

DBH activity.[36, 37] It has been previously established that p-cresol is a potent inhibitor of this

enzyme.[14-16] DBH catalyzes the conversion of DA to NE in dopaminergic and noradrenergic

neurons, and inhibition of DBH can increase the DA levels in the neurons and synapses with a

concurrent reduction in the NE levels.[14-16] The neurochemical profile of striatum and prefrontal

cortex of the *C. difficile* challenged mice showed increased DA levels and a lower NE levels in

these brain areas, suggesting that these alterations could be solely or partially a result of p-cresol

mediated inhibition of DBH enzyme. However, further investigations are required to rule out the

possibilities of reduced expression of DBH genes, and other pathways that affect

neurotransmitter release in response to *C. difficile* infection.

The most notable observation made in this study is the increased DA in the striatum and

pre-frontal cortex **(Figures 4 & 5)** with unchanged or decreased NE levels in mice with *C.*

difficile infection (increased DA to NE ratio). These observations corroborate a recently

published study which demonstrated increased DA levels in striatum and associated ASD like

behavior in mice intravenously administered with p-cresol.[11] DA release in striatum and

prefrontal cortex is generally associated with reward response, motivation, pleasure, and

addiction.[54] In contrast, an increase in DA levels in these brain regions in response to persistent

painful condition (dysbiosis and CDI associated typhlocolitis in this case) is unusual, as pain is a

significant factor that ought to be considered in a *C. difficile* infection study. Generally, DA is

considered as anti-nociceptive and persistent pain is often associated with a decreased DA levels

in brain as opposed to acute pain, such as a pin prick.[55-58] Therefore, a significantly increased DA

and its degradation products observed in the striatum and pre-frontal cortex could be potentially

attributed to a pathway that directly alters DA metabolism, such as p-cresol mediated inhibition

of DBH. Increased DA (or its degradation products) to norepinephrine ratio as seen in both

striatum and pre-frontal cortex of the *C. difficile* infected mice has been implicated in disorders

of DBH activity, such as low DBH levels due to mutations in DBH coding genes.[37] Therefore, *C.*

difficile cresol mediated inhibition of DBH could be a potential reason for this phenomenon.

In contrast, the hippocampal neurochemical profile revealed an unaltered or decreasing

trend of DA and DA degradation products in *C. difficile* infected animals **(Figure 6)**. Similar

results have also been reported in a recent study, where the mice were administered intravenous

p-cresol.[11] This may suggest the dominance of the pain and memory-associated non-

dopaminergic pathways in the hippocampus in response to *C. difficile* infection or high serum p-

cresol. Interestingly, a significant decrease in NE is noticed in hippocampus, suggesting a

reduced production of the NE in the nor-adrenergic neurons projecting to the hippocampus[59],

presumably due to p-cresol mediated inhibition of DBH. This enables exploration into β-

noradrenergic-mediated hippocampal-basolateral amygdalar control of social recognition

memory (SRM) consolidation, which is believed to disordered in individuals with ASD and

therefore fail to establish social relationship and social groups.[60] In addition, we examined the

catecholamine profile in the one of the two major dopamine producing areas in the brain that

projects to the striatum, i.e., substantia nigra (**Figure 7**). In addition to an unaltered NE level, the

concentrations of DA, DOPAC and HVA appears to follow a decreasing trend, suggesting a

potential feedback inhibition of DA production in response to accumulation of DA in striatum[61].

This is also in line with the hypothesis of altered DA to NE conversion in the brain in response

C. difficile p-cresol mediated inhibition of DBH.

One of the major results from this study shows that *C. difficile* infection induces an

increase in serum p-cresol, along with a concurrent increase in DA levels in the striatum, which

has direct implications in the pathogenesis and aggravation of ASD-like behavior in autistic

patients. A recent study by Pasucci *et al* demonstrated that intravenous p-cresol injection

aggravates autistic behaviors in BTBR mice, a murine genetic model of ASD.[11] Similar results

are also demonstrated following oral p-cresol administration in non-mutant mice by Bermudaz-

Martin *et al* [13]. In Pasucci's study, a single short-term intravenous p-cresol administration

aggravated autistic behavior in mice, and most notably, increased the striatal DA levels.

Interestingly, the serum p-cresol assay results and striatal neurochemical profile from *C. difficile*

infected mice corroborates Pasucci *et al*'s observation. This study also suggested potential

involvement of p-cresol mediated DBH enzyme inhibition to explain this observation. Similarly,

Bermudaz-Martin *et al*'s reported impaired excitability of dopaminergic neurons and

precipitation of autistic behavior in C57BL/6 mice continuously administered with oral p-cresol

in drinking water. In contrast, Bermudaz-Martin *et al* attributed p-cresol mediated behavioral and

neuronal alterations to remodeling of gut microbiota (gut dysbiosis) mediated by p-cresol, by

demonstrating similar changes in mice transplanted with fecal material from p-cresol

administered animals. However, Pasucci's results indirectly nullify this theory by demonstrating

similar behavioral and neurochemical alterations in short term, intravenous administration of p-

cresol in BTBR, which entirely bypasses the involvement of gut and leaves minimal room for

dysbiosis or microbial remodeling. Thus, an alternate but direct mechanistic pathway, such as

inhibition of DBH, as proposed by Pasucci *et al* and our current study could be a plausible

explanation for p-cresol induced alterations in dopaminergic neurotransmission. p-Cresol-

producing bacteria, such as *C. difficile,* are extremely resistant to high gut cresol levels.

Moreover, p-cresol strongly inhibits the growth of other bacteria in the gut providing a selective

advantage for p-cresol-producing bacteria to overgrow.[50, 51] This may result in further enhanced

p-cresol production in the gut, and p-cresol mediated neurologic alterations in mice transplanted

with such gut content, which more reasonably explains Bermudaz-Martin's observation.

Regardless, Bermudaz-Martin's experiment compliments Pasucci *et al*'s study in several ways,

as more prolonged nature of p-cresol administration in their study resulted in DA receptor

insensitivity. It has been previously reported that prolonged, high DA levels and increase

dopaminergic stimulation and impair DA receptor sensitivity in post synaptic neurons[62, 63],

supporting the results described in the current study and Pasucci *et al*'s report. Thus, the

published studies, together with our current observations, suggest a potential direct link between

C. difficile infection, gut and serum p-cresol levels, DBH activity, alterations in dopaminergic

activity in brain that have direct implications in the pathogenesis of ASD. Therefore, the results

from this study suggest a potential link between antibiotic induced *C. difficile* infection, bacterial

p-cresol production, reduced DBH activity, and an altered dopaminergic axis, supporting the

previous reports of p-cresol mediated ASD-like behavioral alterations in rodents.

Autism spectrum disorder is a group of heterogenous neurodevelopmental anomalies

where genetic and environmental factors play complex roles, most of which are poorly

understood. Our current study adds to the growing evidence of the 'dopamine hypothesis of

ASD', which is at the same time, can be considered as a limitation of this investigation.

Although dopaminergic system appears to be pivotal in the manifestation of classic autistic

behavioral anomalies i.e., social deficits and stereotypic behavior, several other neurotransmitters

and molecular pathways contribute to the pathogenesis of ASD. Another limitation of this study

is that p-cresol may not be the only neurotoxic compound produced during dysbiosis and *C.*

difficile infection in humans, and there could be several other unknown pathways implicated in

reduced DBH activity and altered catecholamine metabolism in the brain. However, this study

provides a logical association between gut dysbiosis, a known gut microbial toxin (p-cresol), and

a potential mechanistic pathway of deregulated neurotransmission relevant to ASD. However,

the scope of microbial mediated dopaminergic neurotransmission may not be restricted to ASD

alone, but also applicable to other neurologic disorders such as Parkinson's disease, where the

dopaminergic neuronal pathology is implicated. For example, *C. difficile* has been recently

shown to be associated with an increased short-term Parkinson's risk, re-enforcing this concept.[64]

In summary, the current study demonstrated that antibiotic induced *C. difficile* infection

causes significant alterations in the dopaminergic axis in mice. In addition, *C. difficile* infection

significantly increased circulating p-cresol levels and reduced DBH activity in infected animals.

Thus, the published studies, together with our current observations, suggest a potential direct link

between *C. difficile* infection, gut and serum p-cresol levels, DBH activity, and alterations in

dopaminergic activity in the brain that have direct implications in the pathogenesis of ASD.

However, additional mechanistic, and behavioral studies are required to definitively confirm this

hypothesis.

Materials and methods

Bacterial culture, media, and in vitro p-cresol production assay

The human epidemic *C. difficile* strain, UK1 (ribotype 027), was used in this study. *C.*

difficile was cultured on two pre-reduced media: 1) BHI broth supplemented with yeast, 2) BHI

broth supplemented with yeast and p-HPA, in anaerobic conditions (0% oxygen, 5% hydrogen,

5% CO2, and 90% nitrogen at 37°C) and incubated in an anaerobic workstation (AS-580,

Anaerobe Systems, Morgan Hill, CA, USA). Primary *C. difficile* cultures were grown on BHI

broth for 13 hours and then were diluted 1/100 into pre-equilibrated media with and without

0.3% p-HPA. After 24 hours, culture supernatant was collected after centrifugation at 5,000 x g

at 4°C for 15 minutes, followed by filter sterilization. Supernatants were immediately aliquoted

and saved at − 80 °C until analysis. All experiments were run in triplicates using pre-reduced

media in an anaerobic environment, with every experiment being repeated twice. p-Cresol and p-

HPA concentrations in the culture supernatant were estimated using HPLC method as per

standard protocol.[51]

Mouse model for antibiotic-induced **C. difficile** *infection*

All animal experiments were performed as per the protocols approved by ISU

Institutional Animal Care and Use Committee (IACUC-20-091). Fifty-four C57BL/6 mice (3 to

4 weeks old) with an equal male-female ratio were purchased and housed in groups of 2 per cage

with sterile food, water, and bedding. Mice were randomly assigned to each of the three groups:

Control, antibiotic control (Abx), and antibiotic induced *C. difficile* infection (C. Diff) groups.

After 4 days of acclimatization, a mixture of antibiotics and an intraperitoneal injection of

clindamycin was administered to Abx and C. Diff groups, and the C. Diff group mice were orally

gavaged with 1.4 x 10^4 *C. difficile* spores.[34] The animals were euthanized at approximately 48

hours post-inoculation, and the cecal contents and intestinal tissue samples, serum, and brain

sections were collected for further processing.

Histopathology

The cecum and colon tissues were fixed in 10% formalin and then embedded in paraffin.

A 5μm thickness of tissue sections was made and stained with hematoxylin and eosin. A

standardized scoring system was followed to assess CDI-associated histology injuries in

intestinal tissues, and scores were evaluated across the groups based on epithelial damage,

mucosal edema, and neutrophil infiltration.[65] A board-certified pathologist (sample ID-blinded)

performed microscopic analysis using an Olympus BX53 microscope (Olympus Optical

Company, Tokyo, Japan). Scores were combined into a cumulative histologic score for each

mouse ranging from 0 to 9.

HPLC-ECD for tissue catecholamine analysis

Mouse brain regions including striatum, pre-frontal cortex, substantia nigra and

hippocampus were micro-dissected with mouse brain matrix (ASI Instruments, RBM-1000S),

preserved immediately in an antioxidant mixture containing 0.2 M perchloric acid, 0.1%

$NA_2S_2O_5$, 0.05% Na-EDTA, and isoproterenol (internal standard) as described previously and

directly stored in dry ice.[66] The samples were homogenized, sonicated, and filtered in 0.2-micron

filter tubes. 20 µL samples were injected into High-Performance Liquid Chromatography –

Electrochemical Detector (HPLC-ECD) with pump (Thermo-Scientific ISO-3100SD-BM),

autosampler (Thermo-Scientific WPS-3000 TSL) and CoulArray 5600A-ECD detection system

operated at 23 °C connected to Agilent Eclipse Plus (3.5 µm, 100A) C18 HPLC column

(150x4.60 mm). Tissue neurochemical analytes were eluted in MD-TM MP mobile phase (cat.

no. 701332, Thermo Fisher Scientific) under isocratic conditions at 0.6 ml/min for 21 min. DA

(DA) and its major metabolites, Norepinephrine (N.E.), 3,4-dihydroxyphenylacetic acid

(DOPAC), and homovanillic acid (HVA), were analyzed using CoulArray Data Station (v. 3.10)

and quantified using a standard curve. Tissue neurochemical concentrations were normalized

using wet tissue weight (mg) and corrected using the internal standard extraction coefficient.

Quantification of p-cresol

Blood was collected by cardiac puncture. Serum was separated following clotting and

centrifugation at 2500 ×g for 15 min. The collected serum was aliquoted as 40 µL into multiple

microcentrifuge tubes and immediately temporarily stored in dry ice and further long-term

storage at − 80 °C until analysis. The gut contents were collected and snap-frozen using liquid

nitrogen and stored at − 80 °C until quantitative analysis via mass spectrometry.

Reagents: 4-ethyl phenol, p-cresol (pC), salicylic acid (S.A.) were purchased from Millipore-

Sigma, and p-cresol sulfate (pCS) and p-cresol glucuronide (pCG) were purchased from United

States Biological, (Salem, MA). All the solvents were LC-MS grade and purchased from Fisher

Chemical unless otherwise mentioned.

Instrumentation: Conjugated cresols were analyzed via an Agilent 6470 triple quadrupole (QQQ)

mass spectrometer with electrospray ionization (ESI) equipped with an Agilent 1290 Infinity II

UHPLC (Agilent Inc, MA, USA). Conjugated cresols were separated with the Agilent C18

column (2.1 x 100 mm, 1.8 µm) at 40°C. The mobile phase consisted of A: 5mM ammonium

acetate + 0.005% acetic acid and B: Acetonitrile with a 0.400 ml/min flow rate. The solvent

gradient was maintained at 0 min, 0% B; 0–0.25 min, 0–100% B; 0.5–15 min, 100-0% B; 15–20

min, with a 3-minute pre-equilibration.

Quantification: Analysis was performed in negative mode with the nozzle voltage set at 2000 V.

Nitrogen was used as both nebulizer, desolvation, and sheath gas. The nebulizer pressure was set

at 40 psi with a capillary voltage of 2500V. Desolvation gas was delivered at 12 L/min with

temperature was heated to 350°C. The sheath gas flow was 10 L/min at a temperature of 350°C.

High-purity nitrogen was used as a collision gas for collision induced dissociation. Multiple

reaction monitoring (MRM) was used to detect cresol conjugates, the precursor, product ions,

and collision energies are as follows: pCG precursor m/z 283 product m/z 107, 18 eV; pCS

precursor m/z 187 product m/z 107, 18 eV; S.A. precursor m/z 137 product m/z 93, 25 eV.

Free p-cresol was analyzed with an Agilent 5975C quadrupole mass spectrometer equipped with

7890A gas chromatograph. The capillary column used was HP-88 (60m x 0.25mm x 0.2µm). An

injection volume of 2 µL was used with the inlet operating in splitless mode. The oven GC

program used an initial temperature of 40°C for 1 min, then increased to 230°C at a 20°C/min

rate, with a final hold of 7.5 min. Inlet and MSD transfer line temperatures were held at 250°C.

Mass detection was conducted under standard settings with a mass detection range set to m/z 40–

600. Peak identification was performed using Qualitative Analysis (version 10.0) software and

the NIST Mass Spectral Search Program (version 2.3) with the NIST17 and Wiley 11 GC-MS

spectral library. Peak detection and integration were accomplished using Agilent MassHunter

Quantitative Analysis (version 10.0) software to monitor m/z 107 and m/z 108.

Sample preparation for conjugated p-cresol: The extraction protocol was adapted from Y.

Morinaga et al. and C. Shu et al. with modifications as described here. [6, 67] The analysis of p-

cresol conjugates was completed starting with 25-100µL sera samples or 150-500 mg gut

contents previously archived in microcentrifuge tubes. Samples were kept on ice throughout the

extraction. 4-ethyl phenol and salicylic acid were used as internal standards. These two internal

control samples were spiked into the samples as follows: 4-ethyl phenol, 10 µL of 0.25mg/mL in

ethyl acetate (2.5 µg); salicylic acid, 10µL of 0.05mg/mL in 50% methanol (0.5 µg). These

internal standards were also spiked into 200 µL of 0.9% NaCl spiked with internal standards was

used as a negative control. Three volumes (300µL/100mg-sample) of extraction solvent mixture

of acetonitrile and water (2:1 v/v) were mixed with the sample, and after a quick vortex, this was

incubated on ice for 10 minutes. The samples were sonicated for 10 min followed by a 10 min

vortex. The supernatant was recovered after 7 min centrifugation at 13,000g. Pellets were saved

in the freezer for free p-cresol extraction. 200 µL of the supernatant was filtered through a spin

filter (catalog # UFC30LG25, Merck Millipore). Samples were subjected to LC-MS/MS

analysis, 5 µL of which were directly injected into the LC-MS (QQQ).[67]

Sample preparation for free p-cresol: Analysis of free p-cresols was done with a continuation of

the above extract for conjugate cresols. The supernatants were recombined with their respective

pellets in the microcentrifuge tubes. Two volumes (200µL/100mg-sample) of 0.9% NaCl were

added to the mix, followed by the addition of a solvent mix of 1:1 hexane: ethyl acetate (200µL).

The mixture was sonicated for 10 min. After a 10 min vortexing, this was centrifuged for 7 min

to obtain the phase separation. The top layer, which contains free cresols, was carefully

transferred into a GCMS vial.

Preparation of calibration curves: Standards were made for conjugates in 2:1 acetonitrile:water

0.1-0.0005 mg/mL, and for p-cresol in ethyl acetate 0.75-0.0015mg/mL. 10µL of these standards

were added in to 200µL 0.9% NaCl. Internal standards were also added and extracted by

following the same extraction procedure for the samples.

DBH activity assay

Chemicals and reagents: All assay chemicals were purchased from Millipore-Sigma

unless otherwise stated. All solvents were LC-MS grade and purchased from Fisher Chemical.

Sample preparation and DBH assay reaction: Serum samples (40 µl) were prepared and assayed

as previously described by Punchaichira et al., 2018, with the addition of LC-MS quantification

as described below [68-70]. Briefly, 40 µl of the serum samples were added to a cocktail of 1.0 mM

tyramine HCl, 10 mM fumarate, 0.1 mg/ml catalase, 4 mM ascorbate, 2 µM copper sulphate, 30

mM NEM and 1.0 mM pargyline in 125 mM sodium acetate pH 5.2. The mixture was incubated

for 1 h at 37 °C followed by addition of 50 µl of 25 mM EDTA for stopping the reaction.

Instrumentation: Equipment included Agilent Technologies 1100 Series HPLC system coupled

to both a UV-Vis capable diode array detector (DAD) and an Agilent Technologies Mass

Selective Trap SL detector equipped with a SphereClone 5 µm ODS (2) 80 A, L.C. Column 250

x 4.6 mm (Phenomenex) held at 40°C, and an electrospray ion source operated in positive mode.

Running solvents consisted of A: 0.1% acetic acid in water and B: 0.1% acetic acid in

acetonitrile. The flow rate was 0.700 mL/min with isocratic conditions of 35% B. The ion source

drying gas was nitrogen which was set to flow at 12 mL/min at 350°C, the nebulizing gas was set

to 30psi. The mass scan range was m/z 75-300 with a max accumulation time of 300 ms.

470   *Quantification of enzyme activity:* To quantify DBH activity, 10μL of each prepared serum DBH

471   assay sample was injected for LC-MS analysis. The amount of octopamine converted from

472   tyramine was quantified by monitoring the MS/MS product ion m/z 119 fragmented from the

473   $[M+H]^+$ ion (m/z 154) of octopamine at 0.7 fragmentation amplitude. An octopamine standard

474   curve was generated in assay buffer ranging from 0-2 nmoles-octopamine/ sample. Quant

475   analysis for 6300 Series Ion Trap LC/MS Version 1.8 (Bruker Daltonik, Bremen, Germany)

476   software was used for the LC-MS/MS data analysis.

477   *Statistical analysis*

478     Statistical analysis was performed using GraphPad Prism 9 (GraphPad Software, San

479   Diego, CA) with $p < 0.05$ considered statistically significant. All results were expressed as means

480   ± standard errors of the means (SEM) unless otherwise indicated. The differences between the

481   experimental groups were compared using the analysis of variance (ANOVA). The differences

482   between the two groups were analyzed using an unpaired Student's t-test or Mann-Whitney test.

483   **Disclosure statement**

484   The authors declare no competing interest.

486   **Availability of data**

487   The data supporting the findings of this study are available from the corresponding author SM on

488   request.

References

492   1.   Kong X, Liu J, Liu K, Koh M, Tian R, Hobbie C, et al. Altered Autonomic Functions and Gut Microbiome in Individuals with Autism Spectrum Disorder (ASD): Implications for Assisting ASD Screening and Diagnosis. J Autism Dev Disord 2020.

2.   Lim JS, Lim MY, Choi Y, Ko G. Modeling environmental risk factors of autism in mice induces IBD-related gut microbial dysbiosis and hyperserotonemia. Mol Brain 2017; 10:14.

3.   Lin L, Zhang JQ. Role of intestinal microbiota and metabolites on gut homeostasis and human diseases. Bmc Immunol 2017; 18.

4.   Tran SM, Mohajeri MH. The Role of Gut Bacterial Metabolites in Brain Development, Aging and Disease. Nutrients 2021; 13.

5.   Russell WR, Duncan SH, Scobbie L, Duncan G, Cantlay L, Calder AG, et al. Major phenylpropanoid-derived metabolites in the human gut can arise from microbial fermentation of protein. Mol Nutr Food Res 2013; 57:523-35.

6.   Morinaga Y, Fuke C, Arao T, Miyazaki T. Quantitative analysis of cresol and its metabolites in biological materials and distribution in rats after oral administration. Leg Med (Tokyo) 2004; 6:32-40.

7.   Andersen A. Final report on the safety assessment of sodium p-chloro-m-cresol, p-chloro-m-cresol, chlorothymol, mixed cresols, m-cresol, o-cresol, p-cresol, isopropyl cresols, thymol, o-cymen-5-ol, and carvacrol. Int J Toxicol 2006; 25 Suppl 1:29-127.

8.   Persico AM, Napolioni V. Urinary p-cresol in autism spectrum disorder. Neurotoxicol Teratol 2013; 36:82-90.

9.   Gabriele S, Sacco R, Cerullo S, Neri C, Urbani A, Tripi G, et al. Urinary p-cresol is

elevated in young French children with autism spectrum disorder: a replication study.

Biomarkers 2014; 19:463-70.

10.   Shaw W. Increased urinary excretion of a 3-(3-hydroxyphenyl)-3-hydroxypropionic acid

(HPHPA), an abnormal phenylalanine metabolite of Clostridia spp. in the gastrointestinal tract,

in urine samples from patients with autism and schizophrenia. Nutr Neurosci 2010; 13:135-43.

11.   Pascucci T, Colamartino M, Fiori E, Sacco R, Coviello A, Ventura R, et al. P-cresol

Alters Brain Dopamine Metabolism and Exacerbates Autism-Like Behaviors in the BTBR

Mouse. Brain Sci 2020; 10.

12.   Tevzadze G, Oniani N, Zhuravliova E, Darchia N, Eliozishvili M, Gogichadze M, et al.

Effects of a Gut Microbiome Toxin, p-Cresol, on the Indices of Social Behavior in Rats.

Neurophysiology+ 2018; 50:372-7.

13.   Bermudez-Martin P, Becker JAJ, Caramello N, Fernandez SP, Costa-Campos R,

Canaguier J, et al. The microbial metabolite p-Cresol induces autistic-like behaviors in mice by

remodeling the gut microbiota. Microbiome 2021; 9:157.

14.   Southan C, DeWolf WE, Jr., Kruse LI. Inactivation of dopamine beta-hydroxylase by p-

cresol: evidence for a second, minor site of covalent modification at tyrosine 357. Biochim

Biophys Acta 1990; 1037:256-8.

15.   DeWolf WE, Jr., Carr SA, Varrichio A, Goodhart PJ, Mentzer MA, Roberts GD, et al.

Inactivation of dopamine beta-hydroxylase by p-cresol: isolation and characterization of

covalently modified active site peptides. Biochemistry 1988; 27:9093-101.

16.   Goodhart PJ, DeWolf WE, Jr., Kruse LI. Mechanism-based inactivation of dopamine

beta-hydroxylase by p-cresol and related alkylphenols. Biochemistry 1987; 26:2576-83.

17.   Nikolac Perkovic M, Nedic Erjavec G, Stefulj J, Muck-Seler D, Pivac N, Kocijan

Hercigonja D, et al. Association between the polymorphisms of the selected genes encoding

dopaminergic system with ADHD and autism. Psychiatry Res 2014; 215:260-1.

18.   Robinson PD, Schutz CK, Macciardi F, White BN, Holden JJ. Genetically determined

low maternal serum dopamine beta-hydroxylase levels and the etiology of autism spectrum

disorders. Am J Med Genet 2001; 100:30-6.

19.   Kaluzna-Czaplinska J, Socha E, Rynkowski J. Determination of homovanillic acid and

vanillylmandelic acid in urine of autistic children by gas chromatography/mass spectrometry.

Med Sci Monit 2010; 16:CR445-50.

20.   Garnier C, Comoy E, Barthelemy C, Leddet I, Garreau B, Muh JP, et al. Dopamine-beta-

hydroxylase (DBH) and homovanillic acid (HVA) in autistic children. J Autism Dev Disord

1986; 16:23-9.

21.   Lelord G, Callaway E, Muh JP, Arlot JC, Sauvage D, Garreau B, et al. [Modifications in

urinary homovanillic acid after ingestion of vitamin B6; functional study in autistic children

(author's transl)]. Rev Neurol (Paris) 1978; 134:797-801.

22.   DiCarlo GE, Aguilar JI, Matthies HJ, Harrison FE, Bundschuh KE, West A, et al.

Autism-linked dopamine transporter mutation alters striatal dopamine neurotransmission and

dopamine-dependent behaviors. J Clin Invest 2019; 129:3407-19.

23.   Squillace M, Dodero L, Federici M, Migliarini S, Errico F, Napolitano F, et al.

Dysfunctional dopaminergic neurotransmission in asocial BTBR mice. Transl Psychiatry 2014;

4:e427.

24.   Gangi DN, Messinger DS, Martin ER, Cuccaro ML. Dopaminergic variants in siblings at

high risk for autism: Associations with initiating joint attention. Autism Res 2016; 9:1142-50.

25.   Lee Y, Kim H, Kim JE, Park JY, Choi J, Lee JE, et al. Excessive D1 Dopamine Receptor

Activation in the Dorsal Striatum Promotes Autistic-Like Behaviors. Mol Neurobiol 2018;

55:5658-71.

26.   McDougle CJ, Scahill L, Aman MG, McCracken JT, Tierney E, Davies M, et al.

Risperidone for the core symptom domains of autism: results from the study by the autism

network of the research units on pediatric psychopharmacology. Am J Psychiatry 2005;

162:1142-8.

27.   Hookman P, Barkin JS. Clostridium difficile associated infection, diarrhea and colitis.

World JGastroenterol 2009; 15:1554.

28.   Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. Clostridium difficile infection.

Nat Rev Dis Primers 2016; 2:16020.

29.   Rees CA, Shen A, Hill JE. Characterization of the Clostridium difficile volatile

metabolome using comprehensive two-dimensional gas chromatography time-of-flight mass

spectrometry. J Chromatogr B Analyt Technol Biomed Life Sci 2016; 1039:8-16.

30.   Dawson LF, Donahue EH, Cartman ST, Barton RH, Bundy J, McNerney R, et al. The

analysis of para-cresol production and tolerance in Clostridium difficile 027 and 012 strains.

BMC Microbiol 2011; 11:86.

31.   Dawson LF, Stabler RA, Wren BW. Assessing the role of p-cresol tolerance in

Clostridium difficile. J Med Microbiol 2008; 57:745-9.

32.   Saito Y, Sato T, Nomoto K, Tsuji H. Identification of phenol- and p-cresol-producing

intestinal bacteria by using media supplemented with tyrosine and its metabolites. FEMS

MicrobiolEcol 2018; 94.

33.   Sun X, Wang H, Zhang Y, Chen K, Davis B, Feng H. Mouse relapse model of

Clostridium difficile infection. InfectImmun 2011; 79:2856.

34.   Chen X, Katchar K, Goldsmith JD, Nanthakumar N, Cheknis A, Gerding DN, et al. A

mouse model of Clostridium difficile-associated disease. Gastroenterology 2008; 135:1984.

35.   Koppe L, Alix PM, Croze ML, Chambert S, Vanholder R, Glorieux G, et al. p-Cresyl

glucuronide is a major metabolite of p-cresol in mouse: in contrast to p-cresyl sulphate, p-cresyl

glucuronide fails to promote insulin resistance. Nephrol Dial Transpl 2017; 32:2000-9.

36.   Rush RA, Geffen LB. Dopamine Beta-Hydroxylase in Health and Disease. Crc Cr Rev Cl

Lab Sc 1980; 12:241-77.

37.   Gonzalez-Lopez E, Vrana KE. Dopamine beta-hydroxylase and its genetic variants in

human health and disease. Journal of Neurochemistry 2020; 152:157-81.

38.   Paval D, Miclutia IV. The Dopamine Hypothesis of Autism Spectrum Disorder

Revisited: Current Status and Future Prospects. Dev Neurosci-Basel 2021; 43:73-83.

39.   Paval D. A Dopamine Hypothesis of Autism Spectrum Disorder. Dev Neurosci 2017;

39:355-60.

40.   Holingue C, Newill C, Lee LC, Pasricha PJ, Daniele Fallin M. Gastrointestinal symptoms

in autism spectrum disorder: A review of the literature on ascertainment and prevalence. Autism

Res 2018; 11:24-36.

41.   Chaidez V, Hansen RL, Hertz-Picciotto I. Gastrointestinal problems in children with

autism, developmental delays or typical development. J Autism Dev Disord 2014; 44:1117-27.

42.   Madra M, Ringel R, Margolis KG. Gastrointestinal Issues and Autism Spectrum

Disorder. Psychiatr Clin North Am 2021; 44:69-81.

43.   Finegold SM, Downes J, Summanen PH. Microbiology of regressive autism. Anaerobe

2012; 18:260-2.

44.   Wimberley T, Agerbo E, Pedersen CB, Dalsgaard S, Horsdal HT, Mortensen PB, et al.

Otitis media, antibiotics, and risk of autism spectrum disorder. Autism Res 2018; 11:1432-40.

45.   Cowan CSM, Dinan TG, Cryan JF. Annual Research Review: Critical windows - the

microbiota-gut-brain axis in neurocognitive development. J Child Psychol Psychiatry 2020;

61:353-71.

46.   Saurman V, Margolis KG, Luna RA. Autism Spectrum Disorder as a Brain-Gut-

Microbiome Axis Disorder. Dig Dis Sci 2020; 65:818-28.

47.   Adams JB, Borody TJ, Kang DW, Khoruts A, Krajmalnik-Brown R, Sadowsky MJ.

Microbiota transplant therapy and autism: lessons for the clinic. Expert Rev Gastroenterol

Hepatol 2019; 13:1033-7.

48.   Chen K, Fu Y, Wang Y, Liao L, Xu H, Zhang A, et al. Therapeutic Effects of the In Vitro

Cultured Human Gut Microbiota as Transplants on Altering Gut Microbiota and Improving

Symptoms Associated with Autism Spectrum Disorder. Microb Ecol 2020.

49.   Sandler RH, Finegold SM, Bolte ER, Buchanan CP, Maxwell AP, Vaisanen ML, et al.

Short-term benefit from oral vancomycin treatment of regressive-onset autism. J Child Neurol

2000; 15:429-35.

50.   Harrison MA, Faulds-Pain A, Kaur H, Dupuy B, Henriques AO, Martin-Verstraete I, et

al. Clostridioides difficile para-Cresol Production Is Induced by the Precursor para-

Hydroxyphenylacetate. JBacteriol 2020; 202.

51.   Passmore IJ, Letertre MPM, Preston MD, Bianconi I, Harrison MA, Nasher F, et al. Para-

cresol production by Clostridium difficile affects microbial diversity and membrane integrity of

Gram-negative bacteria. PLoS Pathog 2018; 14:e1007191.

52.   Theriot CM, Koenigsknecht MJ, Carlson PE, Hatton GE, Nelson AM, Li B, et al.

Antibiotic-induced shifts in the mouse gut microbiome and metabolome increase susceptibility to

Clostridium difficile infection. Nature Communications 2014; 5.

53.   Mooyottu S, Flock G, Upadhyay A, Upadhyaya I, Maas K, Venkitanarayanan K.

Protective Effect of Carvacrol against Gut Dysbiosis and Clostridium difficile Associated

Disease in a Mouse Model. Frontiers in microbiology 2017; 8.

54.   Klein MO, Battagello DS, Cardoso AR, Hauser DN, Bittencourt JC, Correa RG.

Dopamine: Functions, Signaling, and Association with Neurological Diseases. Cell Mol

Neurobiol 2019; 39:31-59.

55.   Sotres-Bayon F, Pellicer F. The role of the dopaminergic mesolimbic system in the

affective component of chronic pain. Salud Ment 2000; 23:23-9.

56.   Niikura K, Narita M, Butelman ER, Kreek MJ, Suzuki T. Neuropathic and chronic pain

stimuli downregulate central mu -opioid and dopaminergic transmission. Trends in

pharmacological sciences 2010; 31:299-305.

57.   Narita M, Niikura K, Yamashita A, Ikegami D, Kuzumaki N, Narita M, et al. Chronic

Pain Stimuli Downregulate Mesolimbic Dopaminergic Transmission: Possible Mechanism of the

Suppression of Opioid Reward. Contemp Clin Neurosc 2014; 17:49-58.

58.   Yang S, Boudier-Reveret M, Choo YJ, Chang MC. Association between Chronic Pain

and Alterations in the Mesolimbic Dopaminergic System. Brain Sciences 2020; 10.

59.   Lindvall O, Stenevi U. Dopamine and Noradrenaline Neurons Projecting to Septal Area

in Rat. Cell Tissue Res 1978; 190:383-407.

60.   Garrido Zinn C, Clairis N, Silva Cavalcante LE, Furini CR, de Carvalho Myskiw J,

Izquierdo I. Major neurotransmitter systems in dorsal hippocampus and basolateral amygdala

control social recognition memory. Proc Natl Acad Sci U S A 2016; 113:E4914-9.

61.   Best JA, Nijhout HF, Reed MC. Homeostatic mechanisms in dopamine synthesis and

release: a mathematical model. Theor Biol Med Model 2009; 6.

62.   Chen JC, Su HJ, Huang LI, Hsieh MM. Reductions in binding and functions of D2

dopamine receptors in the rat ventral striatum during amphetamine sensitization. Life Sci 1999;

64:343-54.

63.   Henry DJ, Hu XT, White FJ. Adaptations in the mesoaccumbens dopamine system

resulting from repeated administration of dopamine D1 and D2 receptor-selective agonists:

relevance to cocaine sensitization. Psychopharmacology (Berl) 1998; 140:233-42.

64.   Kang X, Ploner A, Ludvigsson J, Williams D, Pedersen N, Wirdefeldt K. Clostridium

Difficile Infection and Risk of Parkinson's Disease: A Swedish Population-Based Cohort Study.

Movement Disord 2019; 34:S747-S.

65.   Shelby RD, Tengberg N, Conces M, Olson JK, Navarro JB, Bailey MT, et al.

Development of a Standardized Scoring System to Assess a Murine Model of Clostridium

difficile Colitis. J Invest Surg 2020; 33:887-95.

66.   Ghosh A, Langley MR, Harischandra DS, Neal ML, Jin H, Anantharam V, et al.

Mitoapocynin   Treatment   Protects   Against   Neuroinflammation   and   Dopaminergic

Neurodegeneration in a Preclinical Animal Model of Parkinson's Disease. Journal of

Neuroimmune Pharmacology 2016; 11:259-78.

67.   Shu C, Chen XJ, Xia TY, Zhang F, Gao SH, Chen WS. LC-MS/MS method for

simultaneous determination of serum p-cresyl sulfate and indoxyl sulfate in patients undergoing

peritoneal dialysis. Biomed Chromatogr 2016; 30:1782-8.

68.   Nagatsu T, Department of Pharmacology SoM, Fujita Health University, Toyoake, Aichi,

Japan. Simple Photometric Assay of Dopamine-β-Hydroxylase Activity in Human Blood: Useful

in Clinical Chemistry. Clinical Chemistry 2021; 55:193-4.

69.   Nagatsu T, Roche Institute of Molecular Biology N, N.J. 07110, Udenfriend S, Roche

Institute of Molecular Biology N, N.J. 07110. Photometric Assay of Dopamine-β-Hydroxylase

Activity in Human Blood. Clinical Chemistry 2021; 18:980-3.

70.   Punchaichira TJ, Deshpande SN, Thelma BK. Determination of Dopamine-beta-

hydroxylase Activity in Human Serum Using UHPLC-PDA Detection. Neurochem Res 2018;

43:2324-32.

Figure 1. C. difficile *UK1 produces a significant amount of p-cresol in vitro*: *C. difficile* was

cultured anaerobically in supplemented brain heart infusion (BHIS) broth with the substrate 0.3

% p-HPA, and the concentrations of p-HPA and newly formed p-cresol in culture supernatant at

different time points were quantified using HPLC analysis. *C. difficile* metabolized the substrate

p-HPA within 24h.

Figure 2. *Antibiotic induced* **C. difficile** *infection increases gut p-cresol concentration*: **(A)**

Experimental outline: Three-four weeks old C57BL/6 mice are treated with an oral antibiotic

cocktail or PBS and an intraperitoneal clindamycin injection or PBS for inducing gut dysbiosis

and then challenged with 10^4 *C. difficile* spores or PBS. Animals were euthanized, and cecal

tissues were collected approximately two days post- *C. difficile* challenge; **(B)** *Histologic lesions*

in different treatment groups: mice colons from each treatment group were processed and stained

with hematoxylin and eosin. Minimal edema, minimal to absent neutrophilic infiltration, and no

necrosis were observed in the colonic epithelium of the Control and Abx groups. Significant

epithelial necrosis, mucosa, and submucosal neutrophilic infiltration and submucosal edema

were observed in the colons of the C. Diff group; **(C)** *Histopathology and colitis score:*

Histologic scores were assessed by a blinded pathologist across the groups based on epithelial

damage, mucosal edema, and neutrophil infiltration. The scores are combined into a cumulative

histologic score for each mouse ranging from 0 to 9; **(D)** *Gut p-cresol concentration:* Total cecal

703   p-cresol concentrations in control, antibiotic control (Abx), and *C. difficile* challenge (C. Diff)

704   groups at 2-day post-inoculation, quantitated by HPLC method**;** **p< 0.01, *p < 0.05.

Figure 3. C. difficile *infection increases serum p-cresol and reduces serum DBH enzyme activity***:** Three-four weeks old C57BL/6 mice are treated with an oral antibiotic cocktail or PBS and an intraperitoneal clindamycin injection or PBS for inducing gut dysbiosis and then challenged with 10^4 *C. difficile* spores or PBS. **(A)** *Serum p-cresol:* Sera were collected 2-day post-inoculation to quantify total serum p-cresol concentration in Control, antibiotic control (Abx), and *C. difficile* challenge (C. Diff) groups using the HPLC method**. (B)** *Relative serum DBH activity:* DBH enzyme activity in each treatment group was determined using an enzymatic assay coupled with LC-MS. The relative DBH activity was expressed as the peak area representing the tyramine octopamine transformation per hour in each treatment group. **p< 0.01, *p < 0.05.

Figure 4. *Antibiotic induced* **C. difficile** *infection increased striatal dopamine and its metabolite levels:* Three-four weeks old C57BL/6 mice are treated with an oral antibiotic cocktail or PBS and an intraperitoneal clindamycin injection or PBS for inducing gut dysbiosis and then challenged with 10^4 *C. difficile* spores or PBS. Striatal catecholamines- **(A)** dopamine (DA), **(B)** its metabolite 3, 4-dihydroxyphenylacetic acid (DOPAC), **(C)** homovanillic acid (HVA), and **(D)** norepinephrine (NE) levels in Control, antibiotic control (Abx), and *C. difficile*

727   challenge (C. Diff) groups at 2-day post-inoculation were quantitated by HPLC-ED method. *p <

728   0.05

Figure 5. *Antibiotic-induced* **C. difficile infection altered catecholamine levels in the prefrontal cortex:** Three-four weeks old C57BL/6 mice are treated with an oral antibiotic cocktail or PBS and an intraperitoneal clindamycin injection or PBS for inducing gut dysbiosis and then challenged with 10^4 *C. difficile* spores or PBS. Prefrontal Cortex (F.CTX) catecholamines- **(A)** dopamine (DA), **(B)** its metabolite 3,4-dihydroxyphenylacetic acid (DOPAC), **(C)** homovanillic acid (HVA), and **(D)** norepinephrine (NE) levels in Control,

738   antibiotic control (Abx), and *C. difficile* challenge (C. Diff) groups at 2-day post-inoculation

739   were quantitated by HPLC-ED method. *p < 0.05.

743   **Figure 6.** *Antibiotic treatment and antibiotic-induced C. difficile infection decreased NE levels*

744   *in the hippocampus:* Three-four weeks old C57BL/6 mice are treated with an oral antibiotic

745   cocktail or PBS and an intraperitoneal clindamycin injection or PBS for inducing gut dysbiosis

746   and then challenged with 10^4 *C. difficile* spores or PBS. Hippocampal catecholamines- **(A)**

747   dopamine (DA), **(B)** its metabolite 3, 4-dihydroxyphenylacetic acid (DOPAC), and **(C)**

748   norepinephrine (NE) levels in Control, antibiotic control (Abx), and *C. difficile* challenge (C.

749   Diff) groups at 2-day post-inoculation were quantitated by HPLC-ED method. (HVA levels were

750   below the detection limit in all groups) $**p< 0.01$, $*p < 0.05$.

751

Figure 7. *Antibiotic treatment and antibiotic-induced C. difficile infection differentially altered DA levels in substantia nigra*: Three-four weeks old C57BL/6 mice are treated with an oral antibiotic cocktail or PBS and an intraperitoneal clindamycin injection or PBS for inducing gut dysbiosis and then challenged with 10^4 *C. difficile* spores or PBS. Substantia Nigra **(SN)** catecholamines- **(A)** dopamine (DA), **(B)** its metabolite 3,4-dihydroxyphenylacetic acid

759 (DOPAC), **(C)** homovanillic acid (HVA), and **(D)** norepinephrine (NE) levels in Control,

760 antibiotic control (Abx), and *C. difficile* challenge (C. Diff) groups at 2-day post-inoculation

761 were quantitated by HPLC-ED method. *$p < 0.05$.