Modelling Eurasian lynx populations in Western Europe: What prospects for the next 50 years?

Authors

Bauduin Sarah, Germain Estelle, Zimmermann Fridolin, Idelberger Sylvia, Herdtfelder Micha, Heurich Marco, Kramer-Schadt Stephanie, Duchamp Christophe, Drouet-Hoguet Nolwenn, Morand Alain, Blanc Laetitia, Charbonnel Anaïs, and Gimenez Olivier

Bauduin Sarah, Office français de la biodiversité, 147 avenue de Lodève, 34990 Juvignac, sarah.bauduin@ofb.gouv.fr

Germain Estelle, Centre de Recherche et d'Observation sur les Carnivores, Lucy, estelle.pichenot@gmail.com

Zimmermann Fridolin, Carnivore Ecology and Wildlife Management. KORA, Thunstrasse 31, CH-3074 Muri bei Bern, Switzerland f.zimmermann@kora.ch; orcid.org/0000-0003-3861-246X

Idelberger Sylvia, Stiftung Natur und Umwelt Rheinland-Pfalz, Diether-von-Isenburg-Str. 7, 55116 Mainz, Sylvia.Idelberger@snu.rlp.de

Herdtfelder Micha, Forest Research Institute of Baden-Wuerttemberg, Wonnhalde 4, 79100 Freiburg, micha.herdtfelder@forst.bwl.de

Heurich Marco, Chair of Wildlife Ecology and Management, Faculty of Environment and Natural Resources, University of Freiburg, Tennenbacher Straße 4, 79106 Freiburg, Germany

Department of Visitor Management and National Park Monitoring, Bavarian Forest National Park, Freyunger Straße 2, 94481 Grafenau, Germany

Faculty of Forestry and Wildlife Management, Campus Evenstad, Inland Norway University of

Applied Sciences, 2480 Koppang, Norway

Marco.Heurich@npv-bw.bayern.de

Kramer-Schadt Stephanie, Department of Ecological Dynamics, Leibniz Institute for Zoo and

Wildlife Research, D-10315 Berlin, Germany; Institute of Ecology, Technische Universität

Berlin, Germany, kramer@izw-berlin.de; ORCID: 0000-0002-9269-4446

Duchamp Christophe, Office français de la biodiversité, Micropolis - La Bérardie F-05000 Gap,

christophe.duchamp@ofb.gouv.fr

Drouet-Hoguet Nolwenn, Office français de la biodiversité, 5 Allée de Bethléem, F-38610

Gières, nolwenn.drouet-hoguet@ofb.gouv.fr

Morand Alain, Centre d'études et d'expertise sur les Risques, l'Environnement, la Mobilité et

l'Aménagement, Direction territoriale Est, Ile du Saulcy, 57045 Metz, alain.morand@cerema.fr

Blanc Laetitia, CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

blanc.laetitia.esr34@gmail.com

Charbonnel Anaïs, Centre de Recherche et d'Observation sur les Carnivores, Lucy,

anais.charbonnel@laposte.net

Gimenez Olivier, CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

olivier.gimenez@cefe.cnrs.fr

Corresponding author

Bauduin Sarah, Office français de la biodiversité, 147 avenue de Lodève, 34990 Juvignac,

sarah.bauduin@ofb.gouv.fr

Abstract

Persistence of populations may be uncertain for large carnivore species, especially for those

established in human-dominated landscapes. Here, we studied the Eurasian lynx in Western

Europe established in the Upper Rhine meta-population (i.e., Jura, Vosges-Palatinian and Black

Forest populations) and in the Alpine population. These populations are currently considered as

endangered or critically endangered due to high anthropogenic mortality, small population size

and low genetic diversity, and isolation. We assessed lynx persistence over a 50-year time

horizon by implementing a spatially-explicit individual-based model, while accounting for road

mortality and habitat selection. Forecasts showed a steady growth rapidly reaching a more stable

phase for the Alpine and Jura populations, and a more heterogeneous positive growth with less

precision for the Vosges-Palatinian and Black Forest populations. Exchanges of individuals

between populations were limited, the Jura population playing the role of a crossroad. Finally,

persistence of lynx in Western Europe seems likely on a large scale over the next 50 years.

Indeed, simulations showed high female occupancy as well as average lynx density over the

whole core areas of the four studied populations. Nevertheless, these results should be interpreted

bearing in mind the model limitations concerning the absence of movement barriers and

inbreeding depression.

Keywords: *Lynx lynx,* population persistence, spatially-explicit individual-based model

Introduction

The Eurasian lynx (*Lynx lynx*) was eradicated in most of Europe between the 17[th] and 20[th] centuries. The main reasons for its disappearance were habitat degradation, human persecution and a decrease in prey availability (Breitenmoser et al., 2000). The species has recently recolonized parts of its historical range in Central and Western Europe thanks to different reintroduction programs which started in the 1970s. Nowadays there are eleven identified lynx populations in Europe (von Arx, 2020), and the species benefits from a conservation status across its whole range area. The species is considered as "least concerned" at the European level of the IUCN Red list. However, its status greatly differs from one population to another, notably because of their difference in population sizes, even though these populations share similar threats, mostly habitat destruction and fragmentation, illegal killings and collisions with vehicles (von Arx, 2020). Long-term persistence remains notably uncertain for the Alpine population (France and Switzerland) and for the Upper Rhine meta-population, which encompasses the Jura population (France and Switzerland), the Vosges-Palatinian population (France and Germany) and few individuals located in the Black Forest and its surroundings (e.g., the Swabian Alb, Germany) (Drouet-Hoguet et al., 2021; Germain & Schwoerer, 2021; Herdtfelder, Schraml, & Suchant, 2021; Idelberger et al., 2021; Molinari-Jobin et al., 2021). These populations are currently defined as endangered (Jura and Alpine) or critically endangered (Vosges-Palatinian) (von Arx, 2020). Indeed, the populations forming the Upper Rhine meta-population remain currently small and isolated. Only a few exchanges between them and with the Alpine population are documented, most likely because of habitat fragmentation and little functional connectivity (Zimmermann & Breitenmoser, 2007; Morand, 2016). Low female dispersal rates and movements most likely slow population expansion due to their conservative dispersal behavior

compared to male lynx (Port et al., 2021). This situation, added to the common origin from the

Carpathian population (Breitenmoser, Breitenmoser-Würsten, & Capt, 1998; Vandel et al., 2006)

of the few reintroduced individuals which did reproduced, resulting in a bottleneck and genetic

drift, may lead to high inbreeding within populations (Breitenmoser-Würsten & Obexer-Ruff,

2003; Bull et al., 2016; Premier et al., 2020). A national action plan has been defined in France

with several objectives in order to restore the lynx to a favorable conservation status in France

(Gatti, 2022).

In this context, wildlife conservationists, scientists and policy-makers face significant

challenges for lynx conservation when individuals inhabit human-dominated landscapes. To

inform decision-making processes for population conservation, several studies have used

individual-based models (IBMs; Railsback & Grimm, 2012). These bottom-up models flexibly

integrate species demography (dispersal, territory establishment, reproduction, mortality) and

how animals interact with their inhomogeneous environment (e.g., habitat selection, collisions,

illegal killings) and other individuals, while accounting for individual characteristics (e.g., sex,

age, dispersal status). Population-level results emerge from the individual-level simulations

(Railsback & Grimm, 2012). These models are used for the management and conservation of

large and small carnivores (e.g., Kramer-Schadt et al., 2004, 2011; Kramer-Schadt, Revilla, &

Wiegand, 2005; Marucco & McIntire, 2010; Hradsky et al., 2019). Recent applications of IBMs

for the lynx species allowed estimating illegal killing (Heurich et al., 2018), linking movements

and genetic diversity (Premier et al., 2020) and evaluating reintroduction scenarios (Ovenden et

al., 2019; Philips, 2020).

Lynx are large carnivores with extensive spatial requirements (Breitenmoser-Würsten et

al., 2007). They are territorial and live in vast connected forested areas (Vandel & Stahl, 2005;

Zimmermann & Breitenmoser, 2007), which sustain important prey populations (Basille et al.,

2009). Natural barriers (e.g., broad waterbodies) as well as anthropogenic barriers (e.g., urban

areas, road network) limit lynx movements (Kramer-Schadt et al., 2004). Roads are complex

infrastructures acting as semi-permeable movement barriers (Klar, Hermann, & Kramer-Schadt,

2006; Zimmermann & Breitenmoser, 2007) in addition to be an important source of mortality

(Schmidt-Posthaus et al., 2002; Ceia-Hasse et al., 2017; Bencin et al., 2019). Therefore, a major

modelling challenge when simulating lynx populations is to account for movements over large

areas and consequently, for the impact of landscape characteristics on different demographic

processes such as dispersal, territory establishment and survival rates. Simulation rules in IBMs

are defined at the individual level and result in different behavior of each simulated individual

due to their own characteristics. Therefore, these models appear as a great tool to integrate the

complex interactions between landscape characteristics and both lynx demographic and spatial

processes, to simulate population changes on a large scale.

Our study aimed to assess the long-term persistence of the Eurasian lynx in the large

range area delimited by the Upper Rhine meta-population and the Alpine population, using the

available population status in 2018. We build a spatially-explicit IBM that integrates up-to-date

information on lynx ecology while accounting for habitat preferences and the risk of collisions

with vehicles. We implement our model in the R programming language and share it publicly

with open-source code allowing full reproducibility and its adoption by ecologists and

conservationists. We use this model to forecast the fate of lynx populations over a 50-year time

horizon. Finally, we provide several population metrics that help in diagnosing the long-term

species persistence in its most western range area.

Material and Methods

Study area and populations

We conducted the study on the Eurasian lynx populations located in France, Germany and

Switzerland (Fig. 1). The three main populations inhabiting the vast mountainous and forested

areas of these countries are the Vosges-Palatinian (France-Germany), the Jura (France-

Switzerland) and the Alpine (France-Switzerland) populations (von Arx, 2020). Some individuals

were also observed in the Black Forest and its surroundings (e.g., Swabian Alb) in Germany (Fig.

1), but were not considered as a lynx population *per se* (von Arx, 2020). Therefore, we should

refer to "mountain ranges" when speaking of the Vosges-Palatinate, Jura, Alps and Black Forest

with its surroundings, but for simplicity and clarity, we use "populations" throughout our paper.

The Jura and Alpine populations originated from individuals reintroduced in Switzerland

in the 1970s (Breitenmoser, Breitenmoser-Würsten, & Capt, 1998; Vandel & Stahl, 2005),

followed by a natural recolonization of the territories. After the complete decline of the Vosges-

Palatinian population in the 18th century, a reintroduction of 21 lynx occurred in the southern part

of the Vosges Mountains (France) between 1983 and 1993 (Vandel et al., 2006). Only 10

individuals contributed to the local establishment of the species (Vandel et al., 2006), without

conclusive stabilization (Charbonnel & Germain, 2020). Since 2016, the lynx has been back in

the Palatinate Forest (Germany), thanks to a new reintroduction program finalized in 2020, with a

few individuals that already moved to the Vosges Mountains (Scheid, Germain, & Schwoerer,

2021; Schwoerer, 2021). Finally, only a few male individuals have been observed in the Black

Forest area since 2013, most of them coming from the Swiss Jura Mountains (Stiftung KORA,

2017; Ministerium für Ländlichen Raum und Verbraucherschutz, 2019; Wölfl et al., 2021).

Figure 1: Eurasian lynx presence as available in 2017-2018 in France (Fr.), Germany (Ger.) and

Switzerland (Sw.) over the study area (black rectangle). Data for France cover the period from

01/04/2013 to 31/03/2017 (OFB Réseau Loup-Lynx), for Germany from 01/05/2017 to

30/04/2018 (Bundesamt für Naturschutz) and for Switzerland from 01/01/2015 to 31/12/2017

(KORA). We used a standardized 10x10 km grid from the European Environment Agency for

France and Germany, and a grid derived from the 1:25,000 map for Switzerland). The four colors

are for the four different populations: the Vosges-Palatinian population (blue; 1,800 km^2) with

cells in France (Vosges Mountains) and Germany (Palatinate Forest), the Black Forest population

(purple; 900 km^2) in Germany, and the Jura (green; 12,057 km^2) and the Alpine (red; 11,190 km^2)

populations with cells in France and Switzerland. Top right corner inset: Europe with the location

of the study area.

Lynx population model

Based on previous works by Kramer-Schadt et al. (2004, 2005, 2011), we built a spatially-

explicit individual-based model (SE-IBM) to simulate lynx populations dynamics and dispersal,

while accounting for the risk of lynx-vehicle collisions and lynx habitat preferences. An SE-IBM

is an IBM where individual responses to behavioral rules are constrained by environmental

characteristics. Our lynx SE-IBM is made of four components (Fig. 2) for which the complete

description is available in Appendix A. The first component represents the impact of the road

network on lynx survival. Hereby, the "collision layer" (Fig. A.1) represents lynx-vehicle lethal

collision probabilities and is based on a collision model. The second component represents the

impact of land cover on lynx space use with a habitat model yielding a "habitat layer" (Fig. A.2)

and showing the distribution of the different lynx habitat types. These two spatial layers influence

the behavioral rules followed by the simulated lynx individuals. The third component represents

the initial lynx populations made of lynx individual's location (Fig. A.3) and characteristics (e.g.,

age, sex). The fourth component details all SE-IBM rules including lynx demography, dispersal

movement and territory establishment. The four components were combined to create an SE-IBM

and simulate lynx populations. The initial lynx populations are the starting individuals created at

the beginning of the simulation. Simulated individuals are either resident (i.e., established with a

territory) or dispersers (Fig. A.4). Dispersing lynx move through the landscape, driven by a

preference towards certain habitat types defined by the "habitat layer", a constant direction and

their memory (Fig. A.5). Dispersers can die with a fixed annual mortality probability or from

vehicle collisions (additive spatial mortality), defined using the "collision layer". When arriving

at a new location, dispersers look for a territory (Fig. A.6). Females need a large area of good

quality habitat to establish, whereas male establishment is driven by the presence of available

resident females. A female resident may reproduce if there is a resident male available nearby.

Residents may also die with a fixed mortality or by vehicle collision. This sequence represents

one year and is repeated with the updated individuals (e.g., newborns added, dead individuals

removed, individuals at different locations, new residents) as long as desired. A complete

description of the SE-IBM structure and processes following the Overview, Design concepts, and

Details (ODD) protocol (Grimm et al., 2006, 2010) is provided in Appendix B. The full list of

model parameters with their values is also provided in Appendix B (Table B.1). We calibrated

parameters to better fit the model to our studied populations. We also performed a sensitivity

analysis to evaluate the changes in the main results following the changes in parameter values.

Full details on model calibration, sensitivity analysis and validation are available in Appendix C.

Figure 2: The four components of the lynx SE-IBM. The first component represents the impact

of the road network on lynx survival through vehicle collision. A generalized linear model

predicts collision probabilities ("Collision layer", Fig. A.1). The second component represents the

impact of land cover on the lynx populations. A site-occupancy model predicts lynx habitats

influencing lynx movement ("Habitat layer", Fig. A.2). The third component represents lynx

populations with individuals' locations and characteristics (Fig. A.3). The fourth component

encompasses all ecological rules and SE-IBM parameters. All four components are included in

the SE-IBM to simulate lynx populations and assess their persistence with different metrics.

Population persistence

We ran 100 replicates of the lynx SE-IBM to forecast the populations over 50 years. We used a

different initial population (Appendix A) at each replicate to avoid bias due to initial locations of

simulated individuals. Initial populations were created based on real monitoring data (Appendix

A). We defined a burn-in phase of 2 years after the start of the simulation (i.e., only simulation

outputs from the 3rd year of simulation were included in the conclusions of the analyses) to let the

population settle down as all individuals from the initial population were defined as dispersers.

We did not simulate any mortality during the first year of simulation to allow all individuals to

find a territory, if they can, without dying while doing so. To evaluate lynx persistence, we

analyzed a) population growth rates, b) number of movements between populations looking at

establishments outside of the lynx native population, c) female territory occupancy, and d) lynx

density.

a) We extracted the number of individuals in each population, for every year and for each

replicate. Then, we calculated the growth rate for each replicate and each population over the

simulated years as the number of individuals at time t divided by the number of individuals at

time t-1. Finally, we calculated the mean growth rate per year and per population, and the 95%

confidence interval of the mean over the 100 replicates.

b) We extracted the number of times individuals established their territory in a population

area not corresponding to the one they were born in ("Population layer", Appendix A) for every

year and for each replicate. We did not account for individuals which moved to another

population area during their dispersal but finally came back to their native population area to

establish, nor those which died in another population area while dispersing. We calculated the

number of lynx establishing outside of their natal population per year for each replicate and for

each direction (e.g., from the Alpine to the Jura population and vice versa, from the Jura to the

Vosges-Palatinian population and vice versa). Finally, we calculated the mean and 95%

confidence interval of the mean across all replicates.

c) We extracted female territory locations at the last year of simulation for each replicate.

We focused on female territories as male territories are based on those of the females. We

assigned a "1" to each cell of the gridded study area (1 km^2) that was occupied by a female

territory and a "0" otherwise. Finally, we estimated territory occupancy by calculating the mean

value per cell from these rescaled maps over the 100 replicates. We also did a visual validation of

our model predictions using GPS and VHF tracks from several collared female residents. To this

end, we overlaid telemetry locations on the resulting territory occupancy map to see if they

mostly fall in cells often selected by simulated females, to establish their territory (details in

Appendix D).

d) We extracted the locations of all adult residents at the last year of simulation for each

replicate. As density measure, we computed the mean number of lynx on each 1 km^2 cell over all

replicates. Finally, we summed these means at the 100 km^2 resolution cells to obtain lynx density

values at this scale.

Results

Population growth rates

Simulations predicted similar growth rate patterns for all lynx populations, with growth rates above one (i.e., growing phase) slowly decreasing towards reaching one (i.e., stabilization phase at saturation) along the 50 years of simulation (Fig. 3). The Alpine and Jura populations had similar patterns with their maximum growth rates equal to 1.09 (sd = 0.04) and 1.05 (sd = 0.06) at the 10th and 13th year of simulation, respectively. The Jura population quickly reached a stabilization phase around the 20th year and fluctuated a little over the final time span of the simulation. The Alpine population growth rate similarly decreased rapidly, reaching a low value, but still significantly over 1. The Vosges-Palatinian and Black Forest populations had more fluctuating growth rates over the simulation. It reached a maximum of 1.18 (sd = 0.51) and 1.45 (sd = 0.93) at the 23rd and 14th year of simulation, respectively. Confidence intervals around mean growth rates were much larger than those of the Alpine and Jura populations. The Vosges-Palatinian and Black Forest populations also had their growth rates decreasing after reaching their maximum, until reaching values close to 1 but only towards the end of the simulation time period.

Figure 3: Annual rates of increase over the simulated years for each population. The grey area represents the 2-year burn-in phase at the beginning of the simulation. Points are means over 100 replicates and colored envelopes are 95% confidence intervals from 100 replicates. The dashed line represents a stabilization of the population (growth rate equals to 1).

Lynx establishments outside their native populations

The number of lynx establishments in a population area different from their native one showed that movements between populations were rare for the four populations. Indeed, the highest values were 2.5 lynx per year on average, concerning individuals that were born in the Alps and successfully established in the Jura when reaching 25 years of simulation (Fig. 4.a). The Jura population seemed at the center of lynx movements between populations in Western Europe (Fig. 4). However, the number of individuals born in Jura and establishing elsewhere decreased around

the 20[th] year of simulation, matching when the population started to stabilize (Fig. 3 and 4.b).

There was almost no exchange of individuals between the Alpine and Vosges-Palatinian

populations, in any direction (Fig. 4.a and 4.c); as these are the most distant populations, the other

ones must serve as stepping stones. Surprisingly, the Black Forest showed exchange of

individuals mainly with the Vosges-Palatinian population, even though the Black Forest seemed

evenly distant to all three other populations (Fig. 4).

a)   b)

c)   d)

Figure 4: Number of lynx establishments outside of their native populations per year when born

in the a) Alpine, b) Jura, c) Vosges-Palatinian or d) Black Forest population. Colors represent

different directions from the native populations (each figure) to those where individuals

established (different colored lines). The grey area represents the 2-year burn-in phase. Points are

mean over 100 replicates and colored envelopes are 95% confidence intervals around the mean

from 100 replicates.

Territory occupancy

At the end of the simulation period, female territories occupied most of the study area (Fig. 5). In

the core areas of the four populations, female occupancy was high, reaching values above 0.85 in

all of them. The overlaid VHF and GPS paths of collared resident females (Fig. 5 and Appendix

D) matched well the limits of the most frequently occupied cells. Most of the patches far from

these core areas were never occupied over the 100 replicate simulations (Fig. 5). Nevertheless,

there were distant patches, especially the bigger ones, occupied by female territories with low

probabilities (Fig. 5).

Figure 5: Occupancy by female territories over the study area at the last year of simulation.

Values, ranging from zero (excluded) to one represented by the color scale blue-yellow-red, are

mean occupancy probability per cell of 1 km² over 100 replicates (e.g., a cell equal to 1 means

the cell was included in a female territory, at the 50[th] year of simulation, in all 100 simulation

replicates). Grey cells are "breeding habitats" never included in a female territory across all

replicates (equal to zero). White cells are cells of a habitat type different than "breeding habitat"

and therefore could never be included in a female territory. GPS and VHF recorded paths for

female residents are overlaid as thin black lines. A zoom of the different areas with telemetry data

is presented in Appendix D. Axis (x and y) are in meters from the WGS 84 / Pseudo-Mercator

coordinate system.

Lynx density

Adult resident density ranged from 0 to 1.69 lynx per 100 km^2 over the study area, at the last year

of simulation (Fig. 6). The highest density value was in the Alpine population and this population

also had the highest mean density over the four populations (0.55, sd = 0.4). The Jura population

had the lowest density values (max = 0.97, mean = 0.26, sd = .024). The Vosges-Palatinian and

Black Forest had intermediate values (max = 1.13 and 1.16, mean = 0.33 and 0.47, sd = 0.29 and

0.34, respectively).

Figure 6: Adult resident lynx density per 100 km^2 at the last year of simulation. Lynx density

(color scale blue-yellow-red) are mean values over the 100 simulation replicates. White cells had

a density equals to 0. Axis (x and y) are in meters from the WGS 84 / Pseudo-Mercator

coordinate system.

Discussion

The well-established Alpine and Jura populations

The model predicted a steady growth for the Alpine and Jura populations, quickly reaching a

more stable phase indicating that they may soon be at saturation, especially for the Jura

population. However, saturation could be higher than what we found in our simulations,

depending on the strength of density-dependence processes (Zimmermann, Breitenmoser-

Würsten, & Breitenmoser, 2007). Although our model assumed that the number of individuals an

area can support is mainly defined by female territory size, it does not include a relationship

between lynx behavior and density. Yet, studies found that lynx density also influences home

range sizes (Pesenti & Zimmermann, 2013), and differently for females and males (Aronsson et

al., 2016). For instance, in the North-Western Swiss Alps, lynx territories were much larger in the

80s when lynx density was lower compared to the 90s (Breitenmoser-Würsten et al., 2001).

Simulated lynx density in 50 years were a little smaller compared to those estimated in the field

for both populations (Gimenez et al., 2019; Pesenti & Zimmermann, 2013). Our model did not

seem to allow a significant increase in lynx density. The Alpine and Jura areas could therefore

support more individuals if such density-dependent mechanisms would occur.

The Jura population as the crossroad of Western Europe lynx movements

The Jura population was found to be the only one connected by lynx exchanges with all the other

populations which is coherent with observed data, also in terms of number of individuals moving

per year (Drouet-Hoguet et al., 2021). Individual exchanges simulated were low, with only a

couple of individuals moving from their native population to establish their territory in another

one on average per year. Exchanges were predicted the highest for the individuals born in the

Alps and moving to settle down in the Jura Mountains, with two to three individuals per year

after 25 years of simulation. In Switzerland, only a few lynx movements between the Alpine and

Jura populations have been observed until now despite a large camera trapping effort over the

years. But even though camera trapping can detect movements between populations, it cannot

guarantee an exhaustive sampling nor ensuring that individuals successfully established in their

destination which contrast with the computed results. However, Alpine and Jura populations also

differ genetically, suggesting very few exchanges between them (Breitenmoser-Würsten &

Obexer-Ruff, 2003). Our model may slightly over-estimate these dispersal movements due to

absence of movement barriers in the "habitat layer".

When inspecting model output maps (Figs. 5 and 6), individuals are contained in the

Swiss Alps, restricted to an area of good habitat almost totally surrounded by less favorable

habitats (Fig. A.2). However, individuals have recently started to settle permanently on the Swiss

Plateau (between the Alps and the Jura Mountains) predicted with less favorable habitats. Some

have even reproduced successfully (F. Zimmerman, pers. comm.), indicating potential

connections between the two populations in the future. On the other hand, our model suggests

connectivity between the Jura and Alpine populations on the French side with a continuity of a

few forested corridors until the Chartreuse Mountains in the Alps (Zimmermann & Breitenmoser,

2007). Many observations were made via camera traps of lynx moving between the southern part

of the Jura population and the Chartreuse Mountains (Bailly, 2021). However, movement barriers

(e.g., highways) may prevent connections with the rest of the French Alps and the Alpine

populations in Switzerland. Illegal killings, notably in Switzerland in movement corridors as well

as a slow demographic dynamic of the Swiss population may also have prevented a larger

development of the lynx population in the French Alps (Arpin et al., 2024).

Lynx movements simulated from the Jura to establish within the Vosges-Palatinian

population were possible but rare. This is coherent with field monitoring that highlighted the

presence of one male in the south of the Vosges Mountains, coming from the Jura Mountains

during the winter 2014-2015 (Chenesseau & Briaudet, 2016; Hurstel & Laurent, 2016).

Connectivity between these two populations remains far from optimal because of multiple

barriers (e.g., highways, railroads and rivers) and restricted habitat that impede lynx dispersal

between the two mountain ranges and increase collision risk (Zimmermann & Breitenmoser,

2007; Morand, 2016).

Regarding the Black Forest (including the adjoint Swabian Alb area), six males are known

to have immigrated from the Swiss side of the Jura population since 2013 (unpublished data,

Forest Research Institute Baden-Wuerttemberg; Stiftung KORA, 2017; Drouet-Hoguet et al.,

2021), as well as two males from the Northeastern Alps (Herdtfelder, Schraml, & Suchant, 2021).

Simulated lynx movements towards the Black Forest seem therefore coherent with field data.

After 25 years of simulation, it is the Vosges-Palatinian population which seemed the most

connected to the Black Forest. This is concurrent with the growth of the Vosges-Palatinian

population, which could act as source population only after several years, as it started only with

few individuals. It lasted several years until the first lynx immigrated to the Black Forest from the

other populations in our simulation compared to the others. This is due to male establishment

which is driven by female presence in our model, and because the Black Forest population did

not have any female at first, new males arriving could not establish and were doomed to die while

searching for females. A more realistic rule for male establishment would be to let them search

for females but still allow them to settle after a defined period of time, even without females. It

happens quite regularly that males show territorial behavior even without females (M.

Herdtfelder, pers. Comm.).

The increasing Vosges-Palatinian and Black Forest populations

Projections for the Vosges-Palatinian and Black Forest populations showed a positive growth

over the next 50 years. However, there was a difference in their initial populations. The lynx

reintroduction program in the Palatinate Forest conducted since 2016 constituted a new larger

initial population for the Vosges-Palatinian one (Scheid, Germain, & Schwoerer, 2021;

Schwoerer, 2021) compared to the few individuals, only males (Drouet-Hoguet et al., 2021),

located in the Black Forest. Growth rates reached higher values for the Black Forest population

but growth rates were very heterogeneous along the simulated time period and confidence

intervals around the mean were very large. Demographic stochasticity impacted the Black Forest

population a lot more because of its small initial population. Precision, as well as robustness, to

interpret the results for this population were indeed lower than for the Vosges-Palatinian

population. In the field, one female was reintroduced in the Black Forest by the end of 2023 to

support this only-male population but it died in Summer 2024 (www.baden-wuerttemberg.de),

highlighting the high stochasticity of the recolonization of this habitat patch.

Exchanges of individuals between the Vosges-Palatinian population and the Black Forest

one increased only at the end of the simulation, probably when both populations had more

individuals. The two also seemed connected with the Jura population, highlighting the Upper

Rhine meta-population structure. However, functional connectivity within the meta-population

may not be at its best and may be altered by anthropogenic barriers. For example, in the Vosges-

Palatinian population, at the *Col de Saverne,* this forest bottleneck is fragmented by both a

highway and high-speed railway (Klar, Hermann, & Kramer-Schadt, 2006; Morand, 2016;

Scheid, Germain, & Schwoerer, 2021). However, movements are possible as lynx from the

Palatinate Forest are known to have crossed this pass, some even going back and forth

(Idelberger et al., 2021; Scheid, Germain, & Schwoerer, 2021). Moreover, no female lynx has

been observed until now dispersing from the Jura Mountains to the Black Forest crossing the

Rhine valley that separates the Black Forest population from the Jura one, probably due to their

conservative dispersal behavior compared to males (Port et al., 2021). Monitoring data from 2021

indicate that one female from the north-eastern Swiss population crossed this barrier for the first

time (M. Herdtfelder, pers. comm.).

Model limitations

Although predicted growth rates did not seem very sensitive to model calibration and model

predictions were well validated (Appendices C and D), several aspects of our IBM could still be

improved.

The "habitat layer" is defined in categories with strict "barriers" that the simulated lynx

cannot cross, whereas the species may be tolerant of human activities (Basille et al., 2009;

Bouyer et al., 2015). Lynx may live near urban areas and cross small lakes (F. Zimmermann,

pers. comm.), even sometimes large rivers. In that context, the "habitat layer" could be improved

by being defined as a continuous variable representing a degree of selection or permeability of

the landscape to avoid prohibiting the movement through certain landscape elements. The

"habitat layer" could also be improved by accounting for roads and their associated structures as

movement barriers (e.g., highways and other types of roads difficult to cross, over-pass

facilitating the movement) instead of only mortality sources (Klar, Hermann, & Kramer-Schadt,

2006; Marchand et al., 2017).

In our model, lynx movement is not impeded by roads. The inclusion of permeability of

these linear barriers to the lynx movements may help to refine the behavior rules (Marchand et

al., 2017). We could also redefine the movement behavior rules to be sex or age-dependent.

Indeed, females seem more conservative (i.e., disperse close to their natal range) compared to

males and some males are able to disperse over long distances (Port et al., 2021). Moreover, lynx

movement and therefore the impact of habitats and collision risk were modeled only for

dispersing individuals. Movements inside territories were not simulated for resident lynx and we

assumed an even use of the space. Model predictions could be different if resident individuals

would move across their territory in an uneven way, increasing or decreasing the averaged

collision risk of their home range. However, mortality risk for residents, thanks to their

knowledge about their territory to know where and how to cross linear infrastructures, would

probably be lower than for dispersing individuals.

Moreover, our model predictions were made in a static environment, as the landscape did

not change over the simulated period. A modification of the landscape (e.g., agricultural

intensification, forest cover increase) would give different lynx population predictions but a

landscape model would be necessary to predict a realistic future landscape, or different scenarios

could be defined (Bauduin et al., 2018). We used a medium-term simulation period of 50 years in

which a static landscape was a reasonable assumption.

Finally, we did not include any genetic aspect in the demographic rules of our SE-IBM

(Mueller et al., 2022). A collective expertise recently done in France on lynx viability showed

that inbreeding depression could greatly impact population size and extinction risk (Arpin et al.,

2024). We restricted our predictions over a 50-year period to limit the (non-included) effects of

inbreeding depression on population dynamics. However, even within this time frame, population

trends could shift from increasing when inbreeding depression is excluded, to potentially

decreasing when included. Therefore, our results need to be interpreted carefully and more in

lights of population relative differences than raw values for each population *per se*.

Perspectives for assessing lynx conservation strategies

Our model used to evaluate lynx population persistence could be applied to better understand

aspects of lynx conservation that we did not include yet, such as illegal killings (Heurich et al.,

2018). Moreover, thanks to its individual-based structure, a genetic component could easily be

included to track relatedness between individuals and allow studying inbreeding risk and Allee

effects (Arpin et al., 2024; Premier et al., 2020). Our model could also be used to test the effect of

different scenarios, either by modifying the lynx population sizes (e.g., due to illegal killings or

reintroduction programs) or the landscape at different scales (e.g., green bridges, roads

construction, habitat destruction or restoration) and assess the potential benefits or negative

effects on lynx population persistence. Modifications of road networks to improve connectivity,

such as the removal of road segments or the addition of overpasses or underpasses could be

tested. Then, their effect could be included through an updated layer of collision probabilities and

the population persistence calculated accordingly. Similarly, other modifications of land cover

(e.g., restoring forest areas) could also be tested. This model could be very useful for stakeholders

working on corridors and the reduction of lynx-vehicle collisions as well as reintroduction

programs and species acceptance. Helping reducing road mortality using model simulations is

one of the objectives of the French national action plan (Gatti, 2022).

Using IBMs, Herdtfelder (2012) showed that Black Forest population reinforcement with

lynx females might be one solution considering that habitat is of good quality in this area. The

collective expertise on lynx viability also used a model similar to the SE-IBM presented here to

make recommendations on lynx translocations in the Vosges area to support the population

viability (Arpin et al. 2024). IBMs are great tools to evaluate the impact of such management

actions. However, because our model does not include human attitudes towards the species, we

warn against its blind use to assess the effect of reinforcement on lynx long-term persistence

without accounting for the human dimension component. Illegal killings, as they occurred after

the 90's reintroduction program in the southern part of the Vosges Mountains (Vandel et al.,

2006), as well as more recent ones (Germain, 2020), may be an additional mortality that the

model does not account for. The extent of acceptance towards species reinforcement from some

local stakeholders is an important element (Charbonnel & Germain, 2020). In this case, we

recommend that our model be extended to include the dynamics of the whole socio-ecosystem

(Behr, Ozgul, & Cozzi, 2017; Guerrero et al., 2018).

Conclusion

In this paper, we built and analyzed a spatially-explicit individual-based model to forecast the

fate of lynx populations over the next 50 years. Our results suggest that exchanges of individuals

between Vosges-Palatinian, Black Forest, Jura and Alpine populations to establish new territories

were limited, and emphasize that the Jura population plays the role of a Western Europe

crossroad. Overall, lynx persistence in the Upper Rhine meta-population and the Alpine

population over the next 50 years seems likely on a large scale by considering road mortality and

habitat selection.

Data sharing

The lynx SE-IBM and all the necessary layers to run it (Appendix A), as well as the simulation

outputs and the code to analyze those are available on

https://github.com/SarahBauduin/appendix_lynxIBM.

Acknowledgements

We thank all the volunteers from the "Réseau Loup-Lynx" who collected data on the field. SB

and OG were funded by French National Research Agency (ANR☐16☐CE02☐0007). SB was

funded as well by OFB and OG was funded by CNRS and the "Mission pour l'Interdisciplinarité"

through the "Osez l'Interdisciplinarité" initiative. CEFE, Cerema and CROC were funded by

CILB, MTES (ITTECOP) and FRB through the research program ERC-Lynx. Cerema received

support from the " Direction générale des infrastructures de transports et de la mer (Ministère de

la transition écologique) ". CROC was funded in 2019/2020 by the European Union within the

framework of the Operational Program FEDER-FSE "Lorraine et Massif des Vosges 2014–

2020", the "Commissariat à l'Aménagement du Massif des Vosges" for the FNADT ("Fonds

National d'Aménagement et de Développement du Territoire"), the DREAL Grand Est

("Direction Régionale pour l'Environnement, l'Aménagement et le Logement"), the "Région

Grand Est" and the "Fondation d'entreprise UEM". This research is partly a product of the

DISCAR group funded by the synthesis centre CESAB of the French Foundation for Research on

Biodiversity. We thank OFB and Réseau Loup-Lynx for the lynx presence and lynx-vehicle

collision data to build the habitat and collision models; OFB, Réseau Loup-Lynx, KORA and

Bundesamt für Naturschutz for population data to create the initial populations; OFB, Réseau

Loup-Lynx and KORA for the collision data for model validation; and KORA and Stiftung Natur

und Umwelt Rheinland-Pfalz for the female telemetry data for model validation. We thank Elodie

Vercken, Hector Ruiz and Henrik Andren for their very constructive reviews.

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content

of this article. Olivier Gimenez is a PCIEcology recommender.

References

Arlettaz, R., Chapron, G., Kéry, M., Klaus, E., Mettaz, S., Roder, S., Vignali, S., Zimmermann,

F., & Braunisch, V. (2021). Poaching threatens the establishment of a lynx population,

highlighting the need for a centralized judiciary approach. *Front. Conserv. Sci.* **2**, 665000.

Aronsson, M., Low, M., López☐Bao, J. V., Persson, J., Odden, J., Linnell, J. D. C., & Andrén, H.

(2016). Intensity of space use reveals conditional sex specific effects of prey and

conspecific density on home range size. *Ecol. Evol.* **6**, 2957–2967.

Arpin, I., Sarrazin, F., Bal, G., Drouet-Hoguet, N., Dumez, R., Pichon, L., Régnier, A., Renault

G., Barnaud, C., Bauduin, S., Bessa-Gomes, C., Billet, P., Breitenmoser-Würsten, C.,

Duchamp, C., Gimenez, O., Mathevet, R., Mounet, C., Robert, A., Vandel, J.-M., Young, J.,

& Zimmermann, F. (2024). Expertise scientifique collective sur la viabilité des populations

de lynx boréal en France.

Bailly, J. (2021). Le Lynx en Chartreuse - Épine - Mont du Chat : bilan du suivi photographique

2017-2020 réalisé d'après les données collectées dans le cadre du réseau Loup-Lynx de

l'OFB.

Bauduin, S., McIntire, E. J. B., St-Laurent, M.-H., & Cumming, S. G. (2018). Compensatory

conservation measures for an endangered caribou population under climate change. *Sci. Rep.*

8, 1–10.

Basille, M., Herfindal, I., Santin-Janin, H., Linnell, J. D. C., Odden, J., Andersen, R., Arild

Høgda, K., & Gaillard, J. M. (2009). What shapes Eurasian lynx distribution in human

dominated landscapes: selecting prey or avoiding people? *Ecography (Cop.).* **32**, 683–691.

Behr, D. M., Ozgul, A., & Cozzi, G. (2017). Combining human acceptance and habitat suitability

in a unified socio-ecological suitability model: a case study of the wolf in Switzerland. *J.*

Appl. Ecol. **54**, 1919–1929.

Bencin, H. L., Prange, S., Rose, C., & Popescu, V. D. (2019). Roadkill and space use data predict

vehicle-strike hotspots and mortality rates in a recovering bobcat (*Lynx rufus*) population.

Sci. Rep. **9**, 15391.

Bouyer, Y., Gervasi, V., Poncin, P., Beudels-Jamar, R. C., Odden, J., & Linnell, J. D. C. (2015).

Tolerance to anthropogenic disturbance by a large carnivore: the case of Eurasian lynx in

586 south-eastern Norway. *Anim. Conserv.* **18**, 271–278.

587 Breitenmoser-Würsten, C., & Obexer-Ruff, G. (2003). Population and conservation genetics of

588  two re-introduced lynx (*Lynx lynx*) populations in Switzerland – a molecular evaluation 30

589  years after translocation. In *Proc. 2nd Conf. Status Conserv. Alp. Lynx Popul.* Vol. 2, pp.

590  28–31. Amden, Switzerland.

591 Breitenmoser-Würsten, C., Vandel, J.-M., Zimmermann, F., & Breitenmoser, U. (2007).

592  Demography of lynx (*Lynx lynx*)in the Jura Mountains. *Wildlife Biol.* **13**, 381–392.

593 Breitenmoser-Würsten, C., Zimmermann, F., Ryser, A., Capt, S., Laass, J., Siegenthaler, A., &

594  Breitenmoser, U. (2001). Untersuchungen zur Luchspopulation in den Nordwestalpen der

595  Schweiz 1997–2000. *KORA Bericht.*

596 Breitenmoser, U., Breitenmoser-Würsten, C., & Capt, S. (1998). Re-introduction and present

597  status of the lynx (*Lynx lynx*) in Switzerland. *Hystrix* **10**, 17–30.

598 Breitenmoser, U., Breitenmoser-Würsten, C., Okarma, H., Kaphegyi, T., Kaphygyi, U., &

599  Müller, U. M. (2000). Action Plan for the conservation of the Eurasian Lynx (*Lynx lynx*) in

600  Europe. *Nat. Environ.* Vol. 112.

601 Bull, J. K., Heurich, M., Saveljev, A. P., Schmidt, K., Fickel, J., & Förster, D. W. (2016). The

602  effect of reintroductions on the genetic variability in Eurasian lynx populations: the cases of

603  Bohemian–Bavarian and Vosges–Palatinian populations. *Conserv. Genet.* **17**, 1229–1234.

604 Ceia-Hasse, A., Borda-de-Água, L., Grilo, C., & Pereira, H. M. (2017). Global exposure of

605  carnivores to roads. *Glob. Ecol. Biogeogr.* **26**, 592–600.

606 Charbonnel, A., & Germain, E. (2020). Plan Régional d'Actions en faveur du Lynx boréal (*Lynx*

607  *lynx*) dans le Massif des Vosges : rétablir le Lynx dans un état de conservation favorable

608  dans le cadre d'une démarche participative, concertée et partagée avec les acteurs du

609  territoire. Lucy (57), France.

Chenesseau, D., & Briaudet, P.-E. (2016). Destin de lynx, trombinoscope insolite de lynx

identifiés par piégeage photographique□: Bingo□! Du massif jurassien aux Vosges, il n'y a

qu'un pas… de lynx. *Bull. Lynx du Réseau* **20**, 9–10.

Drouet-Hoguet, N., Chenesseau, D., Kunz, F., & Zimmermann, F. (2021). Situation of the lynx in

the Jura Mountains. *Cat News Spec. Issue* **14**, 29–34.

Gatti, S. (2022). Plan National d'Actions en faveur du Lynx boréal (*Lynx lynx*) : rétablir le Lynx

dans un état de conservation favorable en France (2022-2026), France.

Germain, E. (2020). Lettre d'information n°1 du PRA en faveur du Lynx dans le Massif des

Vosges. Janvier-Juin 2020. Lucy (57), France.

Germain, E., & Schwoerer, M.-L. (2021). Situation of the Eurasian lynx in the Vosges

Mountains. *Cat News Spec. Issue* **14**, 34–37.

Grimm, V., Berger, U., Bastiansen, F., Eliassen, S., Ginot, V., Giske, J., Goss-Custard, J., Grand,

T., Heinz, S. K., Huse, G., Huth, A., Jepsen, J. U., Jørgensen, C., Mooij, W. M., Müller, B.,

Pe'er, G., Piou, C., Railsback, S. F., Robbins, A. M., Robbins, M. M., Rossmanith, E.,

Rüger, N., Strand, E., Souissi, S., Stillman, R. A., Vabø, R., Visser, U., & DeAngelis, D. L.

(2006). A standard protocol for describing individual-based and agent-based models. *Ecol.*

Modell. **198**, 115–126.

Grimm, V., Berger, U., DeAngelis, D. L., Polhill, J. G., Giske, J., & Railsback, S. F. (2010). The

ODD protocol: A review and first update. *Ecol. Modell.* **221**, 2760–2768.

Guerrero, A. M., Bennett, N. J., Wilson, K. A., Carter, N., Gill, D., Mills, M., Ives, C. D.,

Selinske, M. J., Larrosa, C., Bekessy, S., Januchowski-Hartley, F. A., Travers, H., Wyborn,

C. A., & Nuno, A. (2018). Achieving the promise of integration in social-ecological

research: a review and prospectus. *Ecol. Soc.* **23**, 38.

Herdtfelder, M. (2012). Natur und sozialwissenschaftliche Analysen anthropogen bedingter

Mortalitätsfaktoren und deren Einfluss auf die Überlebenswahrscheinlichkeit des Luchses

(*Lynx lynx*). Albert-Ludwigs-Universität Freiburg.

Herdtfelder, M., Schraml, U., & Suchant, R. (2021). Steps towards a lynx population in the Black

Forest? *Cat News Spec. Issue* **14**, 45–46.

Heurich, M., Schultze-naumburg, J., Piacenza, N., Magg, N., Čeverny, J., Engleder, T.,

Herdtfelder, M., Sladova, M., & Kramer-Schadt, S. (2018). Illegal hunting as a major driver

of the source-sink dynamics of a reintroduced lynx population in Central Europe. *Biol.*

Conserv. **224**, 355–365.

Hradsky, B. A., Kelly, L. T., Robley, A., & Wintle, B. A. (2019). FoxNet: An individual-based

model framework to support management of an invasive predator, the red fox. *J. Appl. Ecol.*

56, 1460–1470.

Hurstel, A., & Laurent, A. (2016). Première preuve de dispersion du lynx d'Eurasie (*Lynx lynx*)

du Jura vers les Vosges. *Ciconia* **40**, 1–6.

Idelberger, S., Krebühl, J., Back, M., Ohm, J., Prüssing, A., Sandrini, J., & Huckschlag, D.

(2021). Reintroduction of Lynx in the Palatinate Forest Biosphere Reserve, Germany. *Cat*

News Spec. Issue **14**, 38–42.

Klar, N., Hermann, M., & Kramer-Schadt, S. (2006). Effects of roads on a founder population of

lynx in the biosphere reserve "Pfalzerwald-Vosges du Nord." *Naturschutz und*

Landschaftsplan. **38**, 330–337.

Kramer-Schadt, S., Kaiser, T. S., Frank, K., & Wiegand, T. (2011). Analyzing the effect of

stepping stones on target patch colonisation in structured landscapes for Eurasian lynx.

Landsc. Ecol. **26**, 501–513.

Kramer-Schadt, S., Revilla, E., & Wiegand, T. (2005). Lynx reintroductions in fragmented

landscapes of Germany: Projects with a future or misunderstood wildlife conservation? *Biol.*

Conserv. **125**, 169–182.

Kramer-Schadt, S., Revilla, E., Wiegand, T., & Breitenmoser, U. (2004). Fragmented landscapes,

road mortality and patch connectivity: modelling influences on the dispersal of Eurasian

lynx. *J. Appl. Ecol.* **41**, 711–723.

Marchand, P., Garel, M., Bourgoin, G., Duparc, A., Dubray, D., Maillard, D., & Loison, A.

(2017). Combining familiarity and landscape features helps break down the barriers between

movements and home ranges in a non-territorial large herbivore. *J. Anim. Ecol.* **86**, 371–383.

Marucco, F., & McIntire, E. J. B. (2010). Predicting spatio-temporal recolonization of large

carnivore populations and livestock depredation risk: wolves in the Italian Alps. *J. Appl.*

Ecol. **47**, 789–798.

Stiftung KORA. (2017). Jahresbericht, 24 Seiten. Muri.

Ministerium für Ländlichen Raum und Verbraucherschutz. (2019). Wildtierbericht für Baden-

Württemberg 2018. Stuttgart.

Molinari-Jobin, A., Breitenmoser, U., Breitenmoser-Würsten, C., Cerne, R., Drouet-Hoguet, N.,

Fuxjäger, C., Kos, I., Krofel, M., Marucco, F., Molinari, P., Nägele, O., Rauer, G., Sindicic,

M., Trbojevic, I., Wölfl, M., Wölfl, S., & Zimmermann, F. (2021). SCALP: Monitoring the

Eurasian lynx in the Alps and beyond. *Cat News Spec. Issue* **14**, 50–52.

Morand A. (2016). Le Lynx : risques routiers et mesures correctrices – état des lieux et

recommandations. Cerema pour la DGALN DEB / MTES. 82p + Annexes.

Mueller, S.A., Prost, S., Anders, O., Breitenmoser-Würsten, C., Kleven, O., Klinga, P., Konec,

M., Kopatz, A., Krojerová-Prokešová, J., Middelhoff, T.L., Obexer-Ruff, G., Reiners, T.E.,

Schmidt, K., Sindičič, M., Skrbinšek, T., Tám, B., Saveljev, A.P., Naranbaatar, G., &

Nowak, C. (2022). Genome-wide diversity loss in reintroduced Eurasian lynx populations

urges immediate conservation management. *Biol. Conserv.* **266**, 109442.

Ovenden, T. S., Palmer, S. C. F., Travis, J. M. J., & Healey, J. R. (2019). Improving

reintroduction success in large carnivores through individual-based modelling: How to

reintroduce Eurasian lynx (*Lynx lynx*) to Scotland. *Biol. Conserv.* **234**, 140–153.

Pesenti, E., & Zimmermann, F. (2013). Density estimations of the Eurasian lynx (*Lynx lynx*) in

the Swiss Alps. *J. Mammal.* **94**, 73–81.

Philips, I. (2020). An agent based model to estimate lynx dispersal if re-introduced to Scotland.

Appl. Spat. Anal. Policy **13**, 161–185.

Port, M., Henkelmann, A., Schröder, F., Waltert, M., Middelhoff, L., Anders, O., & Jokisch, S.

(2021). Rise and fall of a Eurasian lynx (*Lynx lynx*) stepping-stone population in central

Germany. *Mammal Res.* **66**, 45–55.

Premier, J., Fickel, J., Heurich, M., & Kramer-Schadt, S. (2020). The boon and bane of boldness:

movement syndrome as saviour and sink for population genetic diversity. *Mov. Ecol.* **8**, 1–

17.

Railsback, S. F., & Grimm, V. (2012). Agent-based and individual-based modeling: a practical

introduction. Princeton, NJ: Princeton University Press.

Scheid, C., Germain, E., & Schwoerer, M.-L. (2021). Les Lynx (*Lynx lynx*) du Pfälzerwald

s'installent progressivement dans le Massif des Vosges. *Ann. Sci. Rés. Bios. Trans. Vosges*

du Nord. **20**, 104–125.

Schmidt-Posthaus, H., Breitenmoser-Würsten, C., Posthaus, H., Bacciarini, L., & Breitenmoser,

U. (2002). Causes of mortality in reintroduced Eurasian lynx in Switzerland. *J. Wildl. Dis.*

38, 84–92.

Schwoerer, M.-L. (2021). Axe 3 : Suivi et conservation du Lynx, Action 10 / Bilan 2019-2020

du suivi conduit par le Réseau Loup-Lynx. *PRA Lynx Massif des Vosges, Lett. d'information*

n2 7–14.

Stiftung KORA. (2021). 50 Jahre Luchs in der Schweiz.

Vandel, J.-M., & Stahl, P. (2005). Distribution trend of the Eurasian lynx *Lynx lynx* populations

in France. *Mammalia* **69**, 145–158.

Vandel, J.-M., Stahl, P., Herrenschmidt, V., & Marboutin, E. (2006). Reintroduction of the lynx

into the Vosges mountain massif: From animal survival and movements to population

development. *Biol. Conserv.* **131**, 370–385.

von Arx, M. (2020). *Lynx lynx (*amended version of 2018 assessment). The IUCN Red List of

Threatened Species 2020: e.T12519A177350310

Wölfl, S., Anders, O., Middelhoff, T. L., Hohmann, U., Back, M., Idelberger, S., Krebühl, J.,

Ohm, J., Prüssing, A., Herdtfelder, M., Böcker, F., Erretkamps, J., Kopaniak, L., Wölfl, M.,

Jokisch, S., Hucht-Ciorga, I., Teubner, J., Trost, M., Zschille, J., Jeß, E., & Steinberg, C.

(2021). Status des Luchses in Deutschland. *Natur und Landschaft* **96**.

Zimmermann, F., Breitenmoser-Würsten, C., & Breitenmoser, U. (2007). Importance of dispersal

for the expansion of a Eurasian lynx *Lynx lynx* population in a fragmented landscape. *Oryx*

41, 358–368.

Zimmermann, F., & Breitenmoser, U. (2007). Potential distribution and population size of the

Eurasian lynx *Lynx lynx* in the jura Mountains and possible corridors to adjacent ranges.

Wildlife Biol. **13**, 406–416.

Supplementary Material

Appendix A – Detail of the lynx spatially-explicit individual-based model

A complete description of the model following the Overview, Design concepts, and Details

(ODD) protocol (Grimm et al., 2006, 2010) is provided in Appendix B.

Our lynx spatially-explicit individual-based model is made of four components (main text, Fig.

2). The first component represents the impact of road network on lynx survival via predicted

collision probabilities. The second component represents the impact of land cover on lynx space

use with the definition of different lynx habitat types. These first two components are spatial

layers influencing the behavioral rules followed by simulated lynx individuals. The third

component represents the initial lynx populations made of lynx individual's locations and

characteristics used to launch the SE-IBM. The fourth component details all SE-IBM rules

including lynx demography, dispersal movement and establishment.

Impact of road network (SE-IBM 1st component)

We first built a risk model to predict collision probabilities between lynx and vehicles within the

1 km² resolution gridded study area. We used lynx mortality events recorded by the wolf-lynx

monitoring framework implemented in France (Duchamp et al., 2012, Réseau Loup-lynx

https://www.loupfrance.fr/suivi-du-loup/reseau-loup-lynx/), which represents 84 collision events

recorded between 1982 and 2016, to train a logistic regression explaining lynx collisions using

lynx presence and both road and environmental characteristics (Visintin, van der Ree, &

Mccarthy, 2017; Visintin et al., 2018). We used the IGN route500 $^{©}$ for France (IGN ROUTE

500, 2018) and OpenStreetMap[©] for the other countries (Geofabrik OpenStreetMap, 2014) as

data sources to extract total road length per cell and the type of roads of the longest road segment

in each cell. We classified the road segments as "highways" (i.e., "Type autoroutier" in route500

[©] data, "motorway", "motorway_link", "trunk" and "trunk_link" in OpenStreeMap data), "main

road" (i.e., "Liaison principale" in route500 [©] data, "primary" and "primary_link" in

OpenStreeMap data), "secondary road" (i.e., "Liaison régionale" in route500 [©] data, and

"secondary" and "secondary_link" in OpenStreetMap data) and "local road" (i.e., "Liaison locale"

in route500 [©] data, "tertiary", "tertiary_link" and "unclassified" in OpenStreeMap data). For

environmental characteristics, we used Corine Land Cover[©] at 100 m of resolution for all Europe

(Copernicus, 2012) to calculate the proportion of urban area in each cell (i.e., human presence).

The best model identified included the total road length, the type of road of the longest road

segment and the proportion of urban area (Table A.1). Distance to highways and human density

were also tested to explain lynx-vehicles collisions but were not significant. We used this model

to predict a collision probability, ranging between zero and one in each cell of our study area, to

create the "collision layer" (Fig. A.1). Lynx presence was also included as explanatory variable in

the model. To predict collision probabilities, we defined the lynx as present everywhere on the

gridded study area because simulated individuals in the SE-IBM suffer from collision probability

on the cells they are located so the mortality value should already account for the lynx presence.

Study area grid cells without road intersecting them have a zero-collision probability.

Table A.1: Parameter estimates for the best model identified to explain lynx-vehicle collisions

(logistic regression parameters).

	Estimate	Standard error	Z value	P-value

Intercept	-6.29204	0.24903	-25.267	< 2e-16
Lynx presence	1.11454	0.15664	7.115	1.12e-12
Total road length	0.34289	0.10107	3.392	0.000693
Longest segment == highway	1.69306	0.53300	3.176	0.001491
Longest segment == main road	1.96672	0.27082	7.262	3.81e-13
Longest segment == secondary road	0.53213	0.31437	1.693	0.090517
Urban area	-0.41059	0.16709	-2.457	0.013996

Figure A.1: "Collision layer" with collision probabilities between lynx and vehicles estimated

between zero and one (yellow to red color scale). Collision probabilities are assigned to each cell

of the gridded landscape over the whole study area (black rectangle) used in the lynx SE-IBM

using the collision model. Limits of the countries (main text, Fig. 1) are overlaid in black over the

map.

Impact of land cover (SE-IBM 2nd component)

We built a habitat model to define a habitat type for each cell within the 1 km² resolution gridded

study area. We used a multi-year occupancy model (Isaac et al., 2014; Outhwaite et al., 2018) in

a Bayesian framework to explain regular lynx presence using land cover types important to lynx

(agricultural fields, forest and open land), the distance to highways (Basille et al., 2013), and

human density. The model was calibrated using French data for the lynx presence from 1994 to

2017 (data: Réseau Loup-Lynx https://www.loupfrance.fr/suivi-du-loup/reseau-loup-lynx/). The

model used both lynx detection and non-detection data over the surveyed area each year

(https://ecologicalstatistics.shinyapps.io/flexdashboard_effort/). We used Corine Land Cover[©] at

100 m of resolution for all Europe (Copernicus, 2012) to calculate the proportion of agricultural

and cultivated fields, forest, pasture and open land in each cell. We used the IGN route500[©] for

France (IGN ROUTE 500, 2018) and OpenStreetMap[©] for the other countries (Geofabrik

OpenStreetMap, 2014) as data sources to calculate the distance from each cell center to the

nearest highway. We also used IGN[©] data for France (IGN ADMIN-EXPRESS-COG, 2018) and

NASA[©] data for the other countries (CIESIN GPWv4, 2015) to calculate the mean human density

per cell. The best model identified to explain lynx presence included presence of agricultural

fields, forests, and open lands, distance to highways and human density (Table A.2). Shrub cover

and road length were also tested to explain lynx presence but were not significant. We used this

model to predict lynx occupancy probability in our entire gridded study area for the year 2017.

We then categorized the cells to obtain a "habitat layer" with the four habitat types defined by the

habitat categorization of Kramer-Schadt et al. (2004): "breeding", "dispersal", "matrix" and

"barrier" (Fig. A.2). All cells where lynx occupancy was predicted as non-null (above or equal a

threshold of 0.01) were considered "breeding" habitat. All forested areas not already "breeding"

habitat were considered as "dispersal" habitat. Cells with more than half of their surface covered

by water or urban area defined by Corine Land Cover[©] (Copernicus, 2012) were considered

"barrier" for lynx movements. The rest was considered "matrix" area (i.e., habitat not favorable

for lynx but that can be traversed by dispersers). The map of categorized cells for our study area

defined the "habitat layer" used in the SE-IBM (Fig. A.2).

Table A.2: Parameter estimates for the best model identified to explain lynx presence (Bayesian

occupancy model).

	Mean	Standard deviation
Arable lands	-1.06	0.23
Permanent cultures	-0.85	0.24
Pastures	0.68	0.15
Forests	1.67	0.16
Open lands	0.49	0.17
Distance to highways	-0.70	0.14
Human density	0.77	0.15

Figure A.2: "Habitat layer" with lynx habitat types as breeding habitat (dark green), dispersal

habitat (light green), matrix (yellow) and barrier (red). Habitat types are assigned to each cell of

the gridded landscape over the whole study area (black rectangle) used in the lynx SE-IBM using

the habitat model. Limits of the countries (main text, Fig. 1) are overlaid in black over the map.

Lynx initial populations (SE-IBM 3rd component)

The initial populations to launch the SE-IBM were built with generated locations and

characteristics for the lynx populations using the best data available at the time of the study (Fig.

A.3). We used cells of regular lynx presence for France (for the time period of 01/04/2013 to

31/03/2017 from the Réseau Loup-Lynx, main text, Fig. 1), Switzerland (for the time period of

01/01/2015 to 31/12/2017 from the KORA, main text, Fig. 1) and Germany (for the time period

of 01/05/2017 to 30/04/2018 from the Bundesamt für Naturschutz, main text, Fig. 1). In these

cells, we extracted "breeding" and "dispersal" areas and sampled random lynx locations in these

areas to locate individuals within the different populations. In France, we used the most reliable

lynx density estimate (1.14 lynx per 100 km^2; Gatti et al., 2014), regardless of differences in local

densities (Gimenez et al., 2019), to convert the area of regular presence into a number of

individuals to create. Ninety-two individuals were generated and dispatched over "breeding" and

"dispersing" areas in France: in the Vosges Mountains (500 km^2 of presence), in the French Jura

(7,700 km^2 of presence) and in the French Alps (500 km^2 of presence). A density of 1.14 lynx per

100 km^2 may be an over- or an under-estimate of the lynx density in certain areas but in the

absence of local density for each French population, we used this mean value. On average, this

method led to 4.5 individuals in the French Alps (sd = 2.1), 77.0 individuals in the French Jura

(sd = 3.4), and 8.4 individuals in the Vosges Mountains (sd = 2.7). In Switzerland, we generated

230 individuals distributed in the different areas of presence according to estimated local

population sizes in 2017 (data: F. Zimmermann, pers. comm.). In Germany, 11 lynx (i.e.,

reintroduced lynx still alive and their offsprings) were identified in the Palatinate at the end of

April 2018 (Scheid, Germain, & Schwoerer, 2021) and four male lynx were identified in the

Black Forest (Wölfl et al., 2021). Therefore, we generated 11 lynx in Palatinate Forest and four

lynx in the Black Forest area.

Except for reintroduced individuals for which we knew their characteristics, we randomly

assigned individuals' sex (male or female) according to the ratio 1:1 usually observed at birth

(Breitenmoser et al., 1993; Jedrzejewski et al., 1996). We also randomly assigned an age between

2 and 15 as lynx usually live until 15-17 years old (e.g., Breitenmoser-Würsten, Vandel, et al.,

2007). Age is defined as zero for the first year of life, one during the second year of life, etc. For

the Palatinate part of the Vosges-Palatinian population, we assigned known age and sex (five

females and six males). In the Vosges Mountains, only males were detected in 2017 thanks to the

camera trap survey design implemented (Charbonnel & Germain, 2020) and no case of

reproduction was reported during the years preceding our analyzes. We then only defined males

for the individuals located in the Vosges part of the Vosges-Palatinian population, with ages

randomly generated. We also only defined males for the Black Forest population. All generated

individuals were defined as "disperser" to avoid the bias of defining territories by ourselves. They

find their territories on their own as defined by the SE-IBM rules.

We also used lynx regular presence data to define the "population layer" (Fig. A.3). This

layer took the gridded study area where each cell was assigned one of the four lynx populations

(main text, Fig. 1), taking the population from the closest lynx regular presence cell. This

"population layer" delineates population areas. It is used to define in which population kitten

were born and identified individual exchanges between populations.

Figure A.3: "Population layer" with the example of an initial population over the whole study

area (black rectangle) used in the lynx SE-IBM using the habitat model. Limits of the countries

(main text, Fig. 1) are overlaid in black over the map. Areas for the Vosges-Palatinian (blue),

Black Forest (purple), Jura (green) and Alpine (red) populations were defined using the cells of

lynx regular presence as in 2017-2018. Each cell of the gridded study area was assigned the

population from its closest cell of lynx regular presence. White dots represent one initial

population (i.e., simulated lynx released at the beginning of one simulation replicate) generated

using the cells of lynx regular presence, the density or number of lynx in each population and the

"habitat layer" to place these theoretical individuals in breeding or dispersal habitats. A different

initial population was used at each simulation replicate; we used the same cells of lynx regular

presence, the same density or number of lynx in each population and the same "habitat layer",

only the generated locations, and chosen sex and age for unknown individuals, were different.

SE-IBM rules according to lynx ecology (SE-IBM 4th component)

Lynx individuals are simulated over a landscape represented as a grid of 1 km² resolution,

covering the whole study area and encompassing the four lynx populations (main text, Fig. 1).

The gridded study area resolution corresponds to lynx's perceptual range (Haller & Breitenmoser,

1986) as well as to the resolution of previous lynx IBMs (Kramer-Schadt et al., 2004, 2011;

Kramer-Schadt, Revilla, & Wiegand, 2005). Two variables characterize the gridded study area: a

probability of lynx-vehicle collision between zero and one ("Collision layer", Fig. A.1) and a

habitat type among "breeding", "dispersal", "matrix", and "barrier" ("Habitat layer", Fig. A.2).

Simulated individuals are characterized by their "disperser" (i.e., not established on a territory

and in search of one) or "resident" status (i.e., established on a defined territory), their age and

their sex.

Simulated resident individuals follow rules on a yearly time step (Fig. A.4). They do not

move (i.e., their movements inside their territory are not simulated), they hold a territory and they

may reproduce once a year. They suffer two types of annual mortality: a fixed baseline mortality

and a spatial one derived from the "collision layer". Simulated dispersing lynx follow rules on a

daily time step (Fig. A.4). They do not have a territory yet and move every day along the gridded

study area, searching for a place to establish themselves. Their dispersal movements (Fig. A.5)

and search for a territory (Fig. A.6) are driven by the "habitat layer". At each step, individuals

 can die from the spatial mortality derived from the "collision layer", and daily from a fixed

baseline mortality.

Figure A.4: Diagram of the main structure of the SE-IBM with the main events affecting both

resident and dispersing simulated lynx. Grey circles represent individuals. Yellow boxes

represent SE-IBM main events, detailed in model description. Blue boxes or blue writing

represent different mortality causes individuals suffer. Brown boxes represent environmental

layers used in the SE-IBM and brown arrows point where behavioral rules are constrained by

these layers. Black arrows indicate the flow of the model. Dispersers can go through dispersal

and territory search every day during a year, as long as they do not find a territory (Figs. A.5 and

A.6).

The modeling cycle starts with the dispersing individuals. Dispersers are individuals of

one year old and older that do not hold a territory yet. Both dispersing males and dispersing

females move on the gridded study area, one cell at a time, from one to 45 times per day (Fig.

A.5), following the same rules. The number of steps individuals move per day is sampled, each

day for each disperser, from a non-linear distribution (Kramer-Schadt et al., 2004). Dispersers

follow a correlated habitat-dependent walk in a two-step process: first, they favor their habitat

preferences and then, maintain their previous direction with a certain probability (i.e., correlation

factor; Kramer-Schadt et al., 2004). This movement process has been rigorously tested with

inverse fitting, and "pattern-oriented modelling" (Kramer-Schadt et al., 2007) using telemetry

data of six dispersing lynx (five females and one male) followed in the Swiss Jura Mountains

between 1988 and 1991. First, dispersers choose in which habitat type they will move next.

Dispersers favor "breeding" and "dispersal" habitats without distinction between the two when

moving and tend to avoid "matrix" habitats. By contrast, they never use "barrier" habitats. The

choice to move into the habitat "breeding/dispersal" or "matrix" depends on the types of their

nine available cells for their next step (i.e., their eight surrounding cells plus the one they are

currently on as they also can choose not to move). If the nine cells are all of one type, excluding

the "barrier", so they are either only "breeding/dispersal" (with or without "barrier") or only

"matrix" (with or without "barrier"), the only available habitat type is selected. If the available

cells for an individual are a mix between "breeding/dispersal" and "matrix" habitats (with or

without "barrier"), there is a probability of 0.03 times the number of "matrix" cells among the

nine ones, to choose the "matrix" habitat for the next step (Kramer-Schadt et al., 2004). For

example, if an individual has three "matrix" cells available, there is a probability of 0.09 that it

will choose a "matrix" cell for its following location. Second, once the habitat type is selected,

the choice of the particular cell to move on, among the ones of the selected habitat type, is given

by the correlation part of the movement. Individuals follow a correlated movement with a

probability equal to 0.5, except for the first step of the day where there is no correlated movement

(Kramer-Schadt et al., 2004). If the movement is not correlated, the choice of the next cell among

the ones of the selected habitat type is random. If the movement is correlated, the chosen cell is

the one maintaining the most of the individual's current direction among the ones of the selected

habitat type. The chosen cell is then where the simulated lynx is moving to. Dispersers try to

minimize their time spent in "matrix" habitat and they can use their memory to return to a

previously visited "breeding/dispersal" habitat when needed. Lynx do not move more than nine

cells inside "matrix" habitats, so if an individual already stepped nine consecutive times in

"matrix" cells and the chosen cell for its next step is again of "matrix" type, it will use its

memory and return to the last "breeding/dispersal" cell visited (Kramer-Schadt et al., 2004).

Finally, dispersers rotate towards their chosen cell and move on their center. Once dispersers

move to their next cell, they may die from the spatial mortality due to vehicle collision. This

spatial mortality is the collision probability from the "collision layer" for the cell of their new

location with a correction factor to transform the value into a per step collision mortality. If

dispersers survive, they search for a territory. If found, they stop moving, establish a territory,

and they will become resident the following year. Dispersers that do not find a territory to

establish on their new location keep moving, as many steps during the day as simulated for them

at the beginning of the day. At the end of the day, all individuals that dispersed during the current

year may die from a fixed daily mortality probability (i.e., baseline mortality, Table B.1).

Dispersers that have not established during the day can move and search for a territory every day

during the year. The daily spatial mortality is also applied to individuals which dispersed during

the year but already found a territory during previous days. Regarding spatial mortality, as

movements inside territories are not simulated and they were not considered as resident at the

beginning of the year, a per step spatial or annual mortality cannot be applied. A daily spatial

mortality is then calculated as their future spatial mortality with their resident status divided by

365.

Figure A.5: Diagram of simulated lynx dispersal movements in the lynx SE-IBM. The loop

represents one step. Individuals move one cell at the time per step, up to 45 a day or until they

find a territory to establish (yellow box). Individual movement is constrained by habitat types in

their surroundings defined by "habitat layer" (brown box), probabilities to step into different

habitat types and to keep a direction (i.e., correlated movement), and their memory. Blue boxes

represent mortalities an individual suffers, a spatial one each time reaching a new location,

derived from the "collision layer" (brown box) and a daily one.

Dispersing individuals arriving at a new location have a different strategy to search for a

new territory to establish regarding their sex (Fig. A.6). Females mainly look for good habitat

(which represent high prey availability) while males search for breeding opportunities and seek

female presence (Breitenmoser-Würsten, Zimmermann, et al., 2007b). Female dispersers need to

be on a cell of habitat type "breeding", defined by the "habitat layer", and with enough

unoccupied "breeding" habitats around their position to establish their territory. Territories can

have any shapes but they must only be composed of connected cells of "breeding" habitat (search

in 8-cell directions). The size of the territory females needs to define is the number of connected

"breeding" cells. It represents the female territory core area (Kramer-Schadt, Revilla, &

Wiegand, 2005) and depends on the population area where they are located in. We defined the

size of these territories to establish as the size of the 95% kernel ranges measured in the field as

kernel are computed based on the most used habitats (i.e., "breeding habitat"). When a female

builds a territory in the model, the cells used are no longer available for other females. Territory

size for females in the Alpine population is equal to 76 km^2 (Breitenmoser-Würsten et al., 2001)

and to 119 km^2 in the Jura population (Breitenmoser-Würsten, Zimmermann, et al., 2007b). As

we did not have a reliable estimate for the Vosges-Palatinian and the Black Forest populations,

we assigned them the value of the Jura population due to the similarity of the habitat structure

(Fig. A.2). Once a female has established a territory, she becomes resident. If a resident male

already established his territory nearby, the male includes the new resident female's territory into

his own, if he has less than three females already associated. Then, the two individuals may

potentially reproduce the following year. That means, the male's dispersal strategy to establish is

female-dependent. At their new location, dispersing males check if they arrived inside the

territory of a resident female available (i.e., with no male associated). If a male finds a female, he

pairs with her and his territory becomes the same as the one from the female je joined. He also

looks at nearby territories for up to three other available females to pair with. Males can pair with

females whose territories are within the size of their maximum territories. The maximum distance

between the male location and a female territory (i.e., a cell of her territory) is the radius of the

95% kernel ranges estimated for males. It is equal to 6.6 km for males in the Alpine population

(95% kernel range = 137 km^2; Breitenmoser-Würsten et al., 2001) and to 8.5 km for the other

populations (95% kernel range = 226 km^2; Breitenmoser-Würsten et al., 2007b). If there are

several females available for a particular male, the closest one(s) are chosen. The male can

potentially reproduce the following year with every three females. The male territory is defined

by the union of all the female territories he has paired with. If a disperser cannot find a territory

to create (for female) or to join (for male) at their location, it continues dispersing. If all the

females a male is paired with die, the male start to disperse again in search of new available

female.

1006

1007

Figure A.6: Diagram of the territory establishment in the lynx SE-IBM separated by sex. The

successive grey circles represent the successive status of the individual. Females are constrained

by their surrounding habitat defined by the "habitat layer" (brown box). Males are constrained by

female presence. The territory sizes females need to establish in "breeding" habitat and the

maximum distance to which males can look for available female residents are constrained by the

population in which the individuals are located ("population layer"; brown box).

1014

Lynx are solitary carnivores (i.e., each resident holds its own territory), except during

reproduction (i.e., mating between male and female) and when females are raising their kittens

(Breitenmoser et al., 1993; Stahl & Vandel, 1998). Sexually mature female residents reproduce if

a sexually mature resident male occurs on their territory. Females have 0.5 chance to be sexually

mature at one year old, and they are all sexually mature at two years old. Males have 0.5 chance

to be sexually mature at two years old, all being sexually mature at three years old (Kvam, 1991;

Kramer-Schadt, Revilla, & Wiegand, 2005; Breitenmoser-Würsten, Vandel, et al., 2007). In the

wild, lynx litter size averages two kittens and can be up to four (Breitenmoser-Würsten et al.,

2001; López-Bao et al., 2019). However, around 50% of lynx kittens die before reaching the age

of dispersal (Breitenmoser-Würsten, Vandel, et al., 2007). We simulated that resident females up

to 11 years old ("young" female in the SE-IBM model) produce one or two kittens, with a

probability of 0.5 for each litter size, that will survive until becoming dispersers the following

year (Henriksen et al., 2005; Kramer-Schadt, Revilla, & Wiegand, 2005). Senescence reduces

litter size, therefore "old" females in the model (12 years old and older) produce zero or one

kitten with an equal probability of 0.5 (Henriksen et al., 2005; Kramer-Schadt, Revilla, &

Wiegand, 2005). We defined that residents die from a fixed annual mortality (i.e., baseline

mortality, Table B.1) which does not include the mortality due to vehicle collisions (i.e., spatial

mortality added separately with the collision model) and illegal killing (i.e., unavailable

estimates). We did not define an increase of mortality due to senescence but we set an age

maximum of 20 years (Stahl & Vandel, 1998; Breitenmoser-Würsten, Vandel, et al., 2007; von

Arx et al., 2017). Residents can also die from a spatial annual mortality due to vehicle collisions

inside their territory. This spatial mortality is specific to each resident and corresponds to mean

collision probability inside their territory. A correction factor transforms the collision probability

from the "collision layer" into a yearly collision mortality.

At the end of the year, all individuals (i.e., individuals which were disperser or resident at

the beginning of the year) age by one year. Individuals still dispersing will do again this same

loop and the new residents will do the annual loop, as long as the simulation lasts or until they

die.

1043

The model is coded using R 4.1.0 (R Core Team, 2014). We used the package NetLogoR

(Bauduin, McIntire, & Chubaty, 2019) to facilitate the IBM structure implementation in R

language and the package SpaDES (Chubaty & McIntire, 2018) to schedule SE-IBM rules with

different time units. We also used the packages data.table (Dowle & Srinivasan, 2019),

randomcoloR (Ammar, 2019), raster (Hijmans & Van Etten, 2018), and testthat (Wickham,

RStudiom, & R Core Team, 2019).

1050

1051

Appendix B - Complete description of the lynx spatially-explicit individual-based model following the ODD protocol (Overview, Design concepts, and Details) developed by Grimm et al. (2006, 2010)

Overview

Purpose

The model simulates lynx population dynamics and dispersal, accounting for the impact of the road network via the risk of vehicle collisions and of the land cover to represent the lynx habitat preferences (Kramer-Schadt, Revilla, & Wiegand, 2005). Layers (i.e., maps) of collision probabilities, lynx habitats and lynx populations (Appendix A) are combined with SE-IBM rules simulating lynx dispersal (Kramer-Schadt et al., 2004, 2007) and demography (Kramer-Schadt, Revilla, & Wiegand, 2005; Kramer-Schadt et al., 2011).

Entities, state variables, and scales

The mobile entities of the model represent lynx individuals. Each simulated individual holds several characteristics:

- *id*: each lynx is unique and has a unique numerical identity (*id*);

- *heading*: direction of the lynx in degrees from 0 to 360 (0 is heading North) to one of its eight neighboring cells (i.e., 0º, 45º, 90º, 135º, 180º, 225º, 270º, 315º);

- *location*: coordinates of the lynx current position;

- *previous location*: lynx hold in memory the coordinates of their previous position;

- *population*: the population area where the individuals are born among Vosges-Palatinate, Black Forest, Jura or Alps;

- *sex*: male or female;

- *age*: numerical value to represent the lynx age. This number represents the age as for humans

with zero for the first year of life, one for the second year of life, etc.;

- *status*: resident (i.e., established with a territory) or disperser (i.e., without a territory);

- *steps*: number of steps dispersing lynx have to do during the current day;

- *last dispersing location*: dispersing lynx hold in memory the coordinates of the last cell of

habitat type "breeding" or "dispersal" they visited;

- *number matrix*: how many consecutive steps dispersing lynx move in "matrix" habitat;

- *male id*: for resident female only, the *id* of their male associated if they have one;

- *number females*: for resident male only, the number of resident females associated they have. It

cannot be more than 3;

- *road mortality territory*: mean collision probability inside its territory;

1087

Simulated lynx progress on a gridded study area of 1 km² resolution encompassing the four

populations of interest: Vosges-Palatinian, Black Forest, Jura and Alpine, over Germany, France

and Switzerland (main text, Fig. 1). The gridded study area resolution corresponds to the lynx

perceptual range (Haller & Breitenmoser, 1986) and original resolution for the lynx individual-

based models (Kramer-Schadt et al., 2004, 2011; Kramer-Schadt, Revilla, & Wiegand, 2005).

The gridded study area holds four different variables:

- *collision probability* (from the "collision layer"): cells crossed by roads have an estimated

probability of fatal collision with lynx, a value between zero and one. The collision probability is

equal to 0 for cells with no road in it;

- *habitat type* (from the "habitat layer"): each cell is of one of the following habitat types:

"breeding", "dispersal", "matrix" or "barrier";

- *individual territory*: this variable holds the *id* of the female residents for the cells included in

their territories.

1101

Process overview and scheduling

At the beginning of each simulated year, lynx individuals are differentiated by their resident or

disperser *status* (Fig. A.4). First, dispersing lynx move on the gridded study area (Fig. A.5) on a

daily time step. A number of *steps* to move per day is generated for each dispersing lynx at the

beginning of each day. All dispersers move simultaneously, one step (i.e., cell) at the time,

following a correlated habitat-dependent movement influenced by the *habitat type*. After each

step, dispersing lynx may suffer from a spatial mortality given by the *collision probability* at their

new location divided by a correction factor to transform the value into a per step collision

mortality. Then, the surviving ones search for a place where establish their territory (Fig. A.6).

Dispersing females are constrained by the *habitat type* in their surroundings to establish a new

territory while dispersing males are constrained by the presence of available resident females for

reproduction (Breitenmoser-Würsten, Zimmermann, et al., 2007b). Dispersing lynx that do not

establish a territory continue to disperse during their set number of *steps* per day, and every day

during the year until establish one. At the end of each day, lynx that dispersed suffer from a fixed

daily baseline dispersal mortality. On the other hand, residents progress on a yearly time step.

After the disperser movements, resident lynx may reproduce and kittens are born. Then, residents

may die from a fixed annual baseline mortality and from a spatial annual mortality, the latter

being defined by the mean *collision probability* inside their territories multiplied by a correction

factor to transform the value into a yearly collision mortality. All these happen once in the year.

At the end of a year, all lynx *age* are incremented by one and their *status* is updated if applicable.

Simulated lynx *id*, *population*, and *sex* variables do not change during a simulation. Their

heading, *location*, *previous location*, *last dispersing location*, *number matrix*, *male id*, *number*

females and *road mortality territory* may change at every dispersal step. The *steps* variable is

updated each day, and *age* and *status* are updated once a year. There is no modeling of the

landscape during the simulation, therefore the *collision probability* and *habitat type* are constant

during one simulation. However, the *territory id* may change at every step (i.e., each time there is

a new territory created).

1129

Design concepts

Basic principles

Our lynx SE-IBM is an assemblage of four components: three spatial layers and an existing

individual-based model (Kramer-Schadt et al., 2004, 2007, 2011; Kramer-Schadt, Revilla, &

Wiegand, 2005) parameterized with field data and inverse model fitting (main text, Fig. 2 and

Appendix C). The *collision probability* layer and the *habitat type* layer are output maps from a

collision model and a habitat model (Appendix A). The "population layer" was generated using

presence data and estimated number of lynx (Appendix A).

1138

Emergence

Simulated males and females follow the same rules, except for the territory establishment where

females are constrained by the habitat and males by the presence of females (Breitenmoser-

Würsten, Zimmermann, et al., 2007b). In an area with few individuals, females have many

possibilities of empty land to establish. On the other hand, males will have to move further to find

one of the few females available, inducing longer dispersal movement and higher chances of

dying. However, in a very dense area, this can be reverse. Females will have to move further to

find free land whereas males, if they are not too numerous, may move less as they may have

many females around them to pair with. If there are many males, all females may be already

taken, inducing again longer dispersal for the males.

To mate, residents are constrained by the presence of a partner on their territory and both

individuals need to be in age of reproduction. The presence of a suitable partner depends on their

past dispersal movement and territory search, and if they could establish a new territory or not.

Reproduction success is therefore stochastic, as well as the number of kittens. All this will

modify the number of new individuals in the population.

Residents and dispersers are subject to a fixed baseline and spatial mortality. Both

mortalities are stochastic and the spatial one depends on the location of the individuals (i.e.,

collision probability at their location). The location of the individuals depends on their current or

past dispersal movement that depends on the *habitat type*. All this will modify the number of

individuals dying each year.

1159

1160   *Adaptation*

1161   There is no adaptation in the SE-IBM (e.g., no density-dependent rules). All individuals follow

1162   the same behavioral rules dependent on their inner and environmental characteristics.

1163

1164   *Objectives*

1165   Simulated lynx do not have adaptive traits and they do not have a final goal to reach. They take

1166   decisions at each time step based only their current characteristics and their environment to fulfill

1167   an immediate goal (e.g., where to move next).

1168

Learning

Simulated lynx try to minimize their time spent in poor habitat quality. When they step into

"matrix" habitat, they store the position they just left (i.e., their last position in good habitat so

either in "breeding" or "dispersing" habitat) and start counting how many consecutive steps into

"matrix" they do. At the 10^{th} step, if the chosen cell is still located into "matrix", simulated lynx

will use their memory to find and move back to their last location of good habitat.

1175

Prediction

Simulated lynx cannot predict the future nor have a global view of the study area. Both residents

and dispersers sense the current state of their environment and population in their immediate

surroundings and react to this information only. For example, dispersing lynx move, making

decisions based only on their immediate surroundings. They cannot anticipate their path nor have

a global view of their environment to make advantageous decisions for the long term (e.g., they

cannot find a better territory and/or more quickly).

1183

1184   *Sensing*

1185   Simulated lynx can sense the *habitat types* and the *individual territories* of their surroundings.

1186   Dispersers establishing a territory and trying to find a partner can sense other individuals: it

1187   knows who hold nearby territories and the characteristics of these individuals, such as their *sex*

1188   and the availability to pair with.

1189

1190   *Interaction*

1191   There is a tacit competition between dispersing individuals during the territory search. Dispersers

1192   evaluate their surroundings before establishing one at the time (in a random order each time). The

1193   first female in the list looking for a territory has therefore more chance to find empty space to

establish than the last female. Similarly, the first male in the list looking for an available female

to join has more chance to find a partner than the last male.

Residents also interact with each other for reproduction. If a male and a female resident in

age of reproduction are on the same territory, they may reproduce and produce new individuals.

However, this interaction does not affect any of the individuals (i.e., a reproduction event does

not change the characteristics of the concerned individuals).

Stochasticity

The model involves several stochastic processes. Regarding demography, both reproduction (i.e.,

the reproduction event itself and the number of kittens) and mortality (both fixed baseline and

spatial for both residents and dispersers) are defined with probabilities and therefore outcomes

vary between individuals. During dispersal, the choice of the *habitat type* to select when there is a

mix of "breeding/dispersal" and "matrix" habitats in the lynx surroundings depends on the

probability to step into "matrix". Then, there is also stochasticity in the choice of the next

location within the selected *habitat type*, as the choice to follow a correlated movement or not are

defined by probabilities, as well as the choice of the next location among those available if the

movement is not correlated. Finally, the number of steps dispersing simulated lynx must move

during a day is sampled from a probability distribution and so, the outcomes are not the same for

all individuals all the time.

Collectives

When a resident male finds one or several available resident females, he may pair and reproduce

with them. Every year, the pairs persist unless one of the lynx dies and therefore frees the other

one to find another partner.

1218   Individuals belong to populations. The two rules different between populations are the

1219   size of the territory females have to reach and the maximum size of territory males can hold (i.e.,

1220   how far can they look for females). Other than that, no rule is defined at the population level and

1221   the population affiliation does not affect the individual progress. Individual progress according to

1222   their own characteristics.

1223

Observation

The state of all alive simulated individuals (i.e., their location and characteristics) and of the

individual territory map are available at the end of each yearly time step. All disperser

movements are also recorded on a single map over the whole simulation time. Events changing

the demography are saved each year: which individuals died and how (i.e., from the fixed

baseline mortality or from collision), which individuals reproduce, who are the kittens, and which

individuals became residents.

1231

Details

Initialization

At the beginning of the simulation, a grid covering the whole study area (main text, Fig. 1) with a

resolution of 1 km^2 is created. The values from the "habitat layer" and the "collision layer" are

transferred to the gridded study area variables *habitat type* and *collision probability*. The variable

individual territory is set to missing value (NA) everywhere. *Habitat type* on the borders of the

grid are set to "barrier" so that individuals "bounce" on the borders. We assumed that these

individuals would have had a hard time to go beyond the borders of the study area as we

encompassed the main populations and patches of good habitats and therefore would be unlikely

to leave the study area.

Using the "population layer", a lynx population is created with unique *id* for each

individual, *location*, *population*, as well with a *sex* and *age* (when known from field data) for

some individuals. A simulated individual cannot start in "matrix" or "barrier" habitats. Therefore,

if some individuals' *location* are located in these habitats, they are relocated to the closest cell of

"breeding" or "dispersal" habitat. At the start of the simulation, individuals cannot have a

previous location or a *last dispersing location*, so we used their current *location* for these two

variables. For individuals with unknown *sex* or *age* (no field data), a random *sex* (male or female

with ratio 1:1) and *age* (between 2 and 15) is randomly given. A random *heading* (i.e., initial

direction) is given to each individual, between 0° and 360°. At the start, all individuals have a

disperser *status*, 0 *steps* to do, 0 steps done in "matrix" (*number matrix*) and 0 for *road mortality*

territory. As all individuals are dispersers, no couples are made so *male id* for the females are set

to NA and the *number females* for males are set to 0 (Table B.1).

The parameter values used in the SE-IBM to represent the dynamics and dispersal of the

Eurasian lynx in our study area (main text, Fig. 1) are provided in Table B.1.

1256

1257   **Table B.1**: Parameter values used in the SE-IBM

Parameter	Explanation	Value	Reference
endTime	Number of years to simulate the lynx populations.	50	By default
pRepro	Probability of mating for a resident female which has a resident male associated on her territory when they are both in age	0.95	Model calibration (see Appendix C)

	of reproduction.		
nKittyYoungF	Number of kittens that survive until the age of dispersal when the female is considered "young" (maxAgeYoungF parameter).	1 or 2 (with equal probability for each value)	(Breitenmoser-Würsten et al., 2001; Henriksen et al., 2005; Kramer-Schadt, Revilla, & Wiegand, 2005; Breitenmoser-Würsten, Vandel, et al., 2007; López-Bao et al., 2019)
nKittyOldF	Number of kittens that survive until the age of dispersal when the female is considered "old" (maxAgeYoungF parameter).	0 or 1 (with equal probability for each value)	(Breitenmoser-Würsten et al., 2001; Henriksen et al., 2005; Kramer-Schadt, Revilla, & Wiegand, 2005; Breitenmoser-Würsten, Vandel, et al.,

			2007)
maxAgeYoungF	Age maximum for a female to be considered "young".	11	(Henriksen et al., 2005)
minAgeReproF	Age minimum at which all resident females can reproduce (i.e., are sexually mature). One year younger, only half of the females are mature.	2 (3rd year of life)	(Kvam, 1991; Henriksen et al., 2005; Breitenmoser-Würsten, Vandel, et al., 2007)
minAgeReproM	Age minimum at which all resident males can reproduce (i.e., are sexually mature). One year younger, only half of the males are mature.	3 (4th year of life)	(Kvam, 1991)
pMortResAlps	Fixed annual baseline probability of mortality for resident lynx (without the risk of collisions included) of the Alpine population.	0.15	Model calibration (see Appendix C)
pMortResJura	Fixed annual baseline probability of mortality for resident lynx (without the risk of collisions	0.15	Model calibration (see Appendix C)

	included) of the Jura population.		
pMortResVosgesPalatinate	Fixed annual baseline probability of mortality for resident lynx (without the risk of collisions included) of the Vosges-Palatinian population.	0.15	Same value as pMortResJura, see model calibration (Appendix C)
pMortResBlackForest	Fixed annual baseline probability of mortality for resident lynx (without the risk of collisions included) of the Black Forest population.	0.15	Same value as pMortResJura, see model calibration (Appendix C)
corrFactorRes	Correction factor for road mortality risk for residents. This factor multiplies the *road mortality territory* value to transform it into a yearly collision mortality.	3.9	Model calibration (see Appendix C)
corrFactorDisp	Correction factor for road mortality risk for dispersers. This factor divides the collision probability of from the "collision layer" to transform the value into a per step collision mortality.	170	Model calibration (see Appendix C)

ageMax	Age maximum a lynx can reach. When reaching this age, lynx die from natural causes.	20	(von Arx et al., 2017)
xPs	Exponent value of the power function to define the daily step distribution. This distribution is used to sample a number of steps per day to move for dispersing lynx.	11	(Kramer-Schadt et al., 2004)
sMaxPs	Maximum number of steps dispersing lynx can do in one day.	45	(Kramer-Schadt et al., 2004)
pMat	Probability of stepping into "matrix" cells. This probability is multiplied by the number of "matrix" cells available for a dispersing lynx to define its probability to choose this habitat type for its next location.	0.03	(Kramer-Schadt et al., 2004)
pCorr	Movement correlation probability for dispersal movement.	0.5	(Kramer-Schadt et al., 2004)
pMortDispAlps	Fixed daily baseline probability of mortality for dispersing lynx (without the risk of collisions	0.0007	Model calibration (see Appendix C)

	included) of the Alpine population.		
pMortDispJura	Fixed daily baseline probability of mortality for dispersing lynx (without the risk of collisions included) of the Jura population.	0.0009	Model calibration (see Appendix C)
pMortDispVosgesPalatinate	Fixed daily baseline probability of mortality for dispersing lynx (without the risk of collisions included) of the Vosges-Palatinian population.	0.0009	Same value as pMortDispJura, see model calibration (Appendix C)
pMortDispBlackForest	Fixed daily baseline probability of mortality for dispersing lynx (without the risk of collisions included) of the Black Forest population.	0.0009	Same value as pMortDispJura, see model calibration (Appendix C)
nMatMax	Maximum number of consecutive steps a dispersing lynx can do into "matrix" habitat. After this number of steps, disperser can use its memory to find and move in the last good habitat visited ("breeding" or "dispersal"	9	(Kramer-Schadt et al., 2004)

	habitat).		
coreTerrSizeFAlps	Territory size (in km^2) females located in the Alpine population need to reach.	76	(Breitenmoser-Würsten et al., 2001)
coreTerrSizeFJura	Territory size (in km^2) females located in the Jura population need to reach.	119	(Breitenmoser-Würsten, Zimmermann, et al., 2007b)
coreTerrSizeFBF	Territory size (in km^2) females located in the Black Forest population need to reach.	119	(Breitenmoser-Würsten, Zimmermann, et al., 2007b)
coreTerrSizeFVP	Territory size (in km^2) females located in the Vosges-Palatinian population need to reach.	119	(Breitenmoser-Würsten, Zimmermann, et al., 2007b)
terrSizeMAlps	Maximum territory size (in km^2) males located in the Alpine population can reach.	137	(Breitenmoser-Würsten et al., 2001)
terrSizeMJura	Maximum territory size (in km^2) males located in the Jura population can reach.	226	(Breitenmoser-Würsten, Zimmermann, et al., 2007b)

terrSizeMBF	Maximum territory size (in km^2) males located in the Black Forest population can reach.	226	(Breitenmoser-Würsten, Zimmermann, et al., 2007b)
terrSizeMVP	Maximum territory size (in km^2) males located in the Vosges-Palatinian population can reach.	226	(Breitenmoser-Würsten, Zimmermann, et al., 2007b)
nFem	Maximum number of resident females a resident male can pair with and potentially reproduce with.	3	(Kramer-Schadt, Revilla, & Wiegand, 2005)

1258

1259 *Input data*

1260   See Appendix A.

1261

Submodels

Dispersal: A number of steps to do during the day is sampled every day from the distribution

defined by Kramer-Schadt et al. (2004), one per dispersing individual. Dispersers move one cell

per step, into one of their eight neighboring cell or their current cell (i.e., so a total of nine

potential cells for their next location). First, there is a selection of the habitat type each disperser

chooses for its next location. Among those nine cells, all cells of "barrier" type are removed from

the potential choices. As individuals cannot be on "barrier" habitat, it cannot happen that all nine

cells are "barrier". For a dispersing individual, if all his cells are of "breeding" or "dispersal"

type, this type ("breeding/dispersal" without distinction between the two) is selected. If all his

cells are of "matrix" (i.e., habitat not favorable for lynx), this type is selected. If the cells are a

mix of "breeding/dispersal" and "matrix", there is a Bernoulli trial of mean 0.03 (Table B.1)

times the number of "matrix" cells among the nine to select the "matrix" habitat type.

Second, there is the choice of the cell for the next location, among all cells of the selected

habitat type. If there is only one cell of the selected type, the disperser goes on this one (for

example, if the selected cell type is "breeding/dispersal" and there is only one cell of

"breeding/dispersal" habitat, all other cells being "matrix" or "barrier", therefore the only cell of

"breeding/dispersal" is selected). Otherwise, there is a choice either random or governed by a

correlation movement to select the cell. The first step of the day is never correlated, otherwise

there is a Bernoulli trial of mean 0.5 (Table B.1) to determine if the movement is correlated for

each individual for each step. If there is no correlation of the movement, the choice for the cell is

done with an equal probability for all the cells of the selected type. If the movement is correlated,

the rotating angle between the individual's heading and each cell of the selected type is

calculated. There is a preference for the cell minimizing this rotating angle. If the cell that the

individual is on is of the selected habitat type (i.e., this cell can be selected for the individual's

next step which will make the lynx stay at its current location), the preference for this cell is

equal to the preference for the cells inducing a rotation of +/- 90 degrees, to favor a forward

movement. The selected cell is the one with the highest preference (i.e., smallest rotation

movement).

Third, if the selected cell is of "matrix" type, the individuals check how many consecutive

steps they have done in "matrix" habitat (*number matrix*). If a disperser has already step nine

consecutive times (Table B.1) in "matrix" and this 10th consecutive step is in "matrix" too, it used

its memory and select, for its next location, the last visited cell of good habitat ("breeding" or

"dispersal"), stored in his memory (*last dispersing location*) (Kramer-Schadt et al., 2004).

Finally, all dispersers rotate towards their selected cell and move to the center of it. They

update their *previous location* with the coordinates of the cell they were before moving. If their

new location is of type "breeding" or "dispersal", this location is stored in their memory for the

last dispersing location, and their *number matrix* is reset to zero. If their new location is of type

"matrix", their *number matrix* increments of one.

Now, disperser may be also subject to spatial mortality. There is a Bernoulli trial for each

disperser with mean equal to the *collision probability* value of their new location divided by a

correction factor (Table B.1). Dispersers that do not die from spatial mortality search for a new

home range to establish (*searchTerritory* sub-model).

The dispersal movement is done until all dispersing individuals have reached their number

of steps to do during the day, or they have established a territory and are now resident, or they

died from spatial mortality. At the end of the day, all individuals that were dispersers at the

beginning of the year (even the ones that established a territory during the day or previous day)

may die from a fixed daily baseline mortality using a Bernoulli trial for each individual. A daily

spatial mortality is also applied to individuals which dispersed during the year but already found

a territory during previous days. As movements inside territories are not simulated but they were

not considered as resident at the beginning of the year, a per step or an annual spatial mortality

cannot be applied. Their daily spatial mortality is then calculated as their spatial mortality with

their resident status divided by 365. If some of the individuals dying established a territory during

this day, there is an update of their characteristics similarly as the death of resident individuals

(*mortality* submodel).

At the beginning of the simulation, all initial individuals were created as dispersers to let

simulated individuals establish their initial territories. Individuals needs some time to do so,

therefore we did not apply any mortality (daily baseline nor spatial) during the first year of

simulation to allow dispersers to establish their territories.

1320

SearchTerritory: Dispersing individuals arriving at a new location search for a territory to

establish, one at the time, and females first (i.e., male establishment depending on female

territories). One at the time, in a random order each time, females evaluate their surroundings to

establish their territory. Females needs to be on a cell of "breeding" type that is not already

included in any territory to start creating their own. Then, they need to have enough empty

contiguous cells of "breeding" habitat (search in 8-cell directions) to create a big enough territory

(Table B.1). When enough empty contiguous cells of "breeding" habitat are present, females

establish their territory and becomes resident. Otherwise, they stay disperser and keep dispersing.

After a female establishes her territory, all the cells of this territory become occupied and are not

available anymore for the other females. The female *id* is given to the *individual territory* study

area variable for all cells in her territory. The mean *collision probability* of the territory is

calculated and given to the female for her *road mortality territory*. Then, the female checks if

there is a resident male around. If there is a male not further than the radius of the maximum

territory size for the male (Table B.1) and he has less than three females associated, she pairs

with him. Then, she obtains the male *id* for her *male id* and the *number females* of the male

increments of one.

After all dispersing females tried to establish, dispersing males try to disperse of their

own, one at the time in a random order each time. Each male checks if the cell they are currently

on is located inside the territory of an available female (i.e., without male associated yet). If there

is one, the male pairs with this female and becomes resident. His territory is the same one as the

female and its *number females* is set to one. The female obtains the male *id* for her *male id*

variables. Then, the male also looks if there are other nearby available females to pair with, as far

as the radius of its maximum territory size (Table B.1). He can pair with up to three available

females. The male territory represents the union of all the female territory with which he paired

with. His *road mortality territory* is the mean value of the *road mortality territory* of all the

females he paired with.

1347

Reproduction: A female resident in *age* of reproduction (one or two years old, see Table B.1)

with a resident male associated, also in *age* of reproduction (two or three years old, see Table

B.1), may reproduce with a Bernoulli trial (Table B.1) and have offspring. If the couple

reproduces, the female can produce one or two kittens with equal probability (Table B.1) if she is

"young" (11 years old or younger; Table B.1) or zero or one kitten with equal probability if she is

"old" (12 years old or older; Table B.1). Kittens obtain a unique *id*. The *sex* of the kittens is

randomly chosen between male and female with ratio 1:1 (Table B.1). Their *location*, *heading*

and *last dispersing location* are the ones from their mother. Their *previous location* is NA. Their

population is not the one of their parents but the one of the territory where they are born. Their

age is 0, their *status* is resident (i.e., they stay with their mother and cannot disperse at this age),

their *steps* and *number matrix* are zero, their *male id* is NA, and their *number females* and *road*

mortality territory are zero.

1360

Mortality: Resident individuals may die from a fixed baseline annual mortality (Table B.1) and a

spatial mortality using a Bernoulli trial. We did not simulate an effect of senescence but we set an

age maximum (20 years; Stahl & Vandel, 1998; Breitenmoser-Würsten, Vandel, et al., 2007; von

Arx et al., 2017) and all individuals reaching this age die from a non-collision source (i.e., old

lynx mortality is not included in the collision mortality). Resident individuals may also die from

spatial mortality, the mortality rate for female residents is the mean *collision probability* of the

cells inside of their territory (*road mortality territory*) multiplied by a correction factor (Table

B.1). For male residents, spatial mortality rate is the mean *road mortality territory* of all his

paired females multiplied by the same correction factor (Table B.1).

When a resident female dies, her territory disappears (i.e., her *id* is removed from the *individual*

territory study area variable) and the local male resident loses one female. If this male did not

have other females, he becomes disperser again to look for new available female resident as male

home ranges are adjusted to mate availability (Breitenmoser-Würsten, Zimmermann, et al.,

2007b). If the dead female had kittens during the year, they all die too. When a male resident

dies, its paired females become available to reproduce with other males.

1376

1377 Demography: All individuals' *age* increment of one. The *status* of the offspring of the year

1378 changes from resident to disperser.

1379

1380

1381

 Appendix C – Calibration of the parameters, sensitivity analysis and model validation

1383

Calibration of the parameters

We calibrated the lynx SE-IBM using a pattern-oriented modeling (Grimm & Railsback, 2012;

Gallagher et al., 2021) in a two-phase process. To better adapt the model to our study area and

simulated lynx populations, we (re)estimated the following 7 parameters:

- pMortResAlps / pMortDispAlps / pMortResJura / pMortDispJura: Fixed annual baseline

mortalities (i.e., probabilities) applied to resident / dispersing lynx, without the risk of collisions

included, for the Alpine and the Jura populations. These parameters do not exactly represent the

mortality rates for lynx when omitting collision mortality observed in the field as the mortality

due to old age in our model is additive to these estimated baseline probabilities. For the mortality

probabilities for the Vosges-Palatinian and the Black Forest populations, we used those of the

Jura population and they were not calibrated *per se*;

- corrFactorRes: Correction factor for road mortality risk for residents. This factor multiplies the

road mortality territory to transform the value into a yearly collision mortality;

- corrFactorDisp: Correction factor for road mortality risk for dispersers. This factor divides the

collision probability from the "collision layer" to transform the value into a per step collision

mortality;

- pRepro: Probability of mating for a resident female which has a resident male associated on the

same territory, when they are both in age of reproduction. This parameter is different than the

probability of reproducing observed in the field, detected by the presence of kitten, as, in our

model, old female can mate but produce 0 young because of her age.

For the first phase of calibration, we sampled one value for each parameter to calibrate

among wide ranges of values (Table C.1). We defined 50 calibration scenarios by sampling a

value within the range for each parameter for each scenario. We ran 15 replicates of each

calibration scenario. We ran the lynx SE-IBM forecasting the populations over 50 years and we

used a different initial population (Appendix A) at each replicate within each scenario to avoid

bias due to initial locations of simulated individuals. We did not simulate any mortality during

the first year of simulation to allow dispersing individuals to find a territory, if they can, without

dying while doing so. From simulation outputs, we computed 13 patterns for each scenario,

taking the mean value over the 15 replicates. Patterns were population emergent properties from

simulations that we could compare to reference values. We defined a burn-in phase of 2 years

after the start of the simulation to let the population settle down so that only outputs from the 3^{rd}

year of simulation were used to compute patterns. Reference values for the patterns were

extracted from literature or computed from field data. In order to calibrate the model, we looked

for the calibration scenarios (i.e., and their associated parameter values) which minimized the

differences between the pattern values extracted from their simulation outputs and the pattern

values from external sources. The patterns we tried to match with their values and 95%

confidence intervals from literature or data were:

- rAllRes_Alps = 0.24 [0.08;0.57] / rAllDisp_Alps = 0.24 [0.15;0.38]: Annual mortality rates

observed for resident / dispersing individuals from all causes of mortality (i.e., including

collisions) for the Alpine population (Premier et al., 2024).

- rCollRes_Alps = 0.02 [0;0.059] / rCollDisp_Alps = 0.077 [0;0.211]: Annual mortality rates

observed for resident / dispersing individuals from collisions with vehicles for the Alpine

population (Premier et al., 2024).

- nCollAlps = 2 [0;3]: Mean annual number of fatal collisions observed for the Alpine population

(data for 2017-2021: OFB Réseau Loup-Lynx and KORA).

- rAllRes_Jura = 0.17 [0.1;0.29] / rAllDisp_Jura = 0.35 [0.16;0.65]: Annual mortality rates for

resident / dispersing individuals from all causes of mortality (i.e., including collisions) observed

for the Jura population (Premier et al., 2024).

- rCollRes_Jura = 0.034 [0;0.08] / rCollDisp_Jura = 0.067 [0;0.185]: Annual mortality rates for

resident / dispersing individuals from collisions with vehicles observed for the Jura population

(Premier et al., 2024).

- nCollJura = 6 [3;9]: Mean annual number of fatal collisions observed for the Jura population

(data for 2017-2021; OFB, Réseau Loup-Lynx and KORA).

- rRepro = 0.81: Observed proportion of resident females reproducing (Breitenmoser-Würsten,

Vandel, et al., 2007a; López-Bao et al., 2019).

- distDispAlps = 26 [15;36] / distDispJura = 63 [39;87]: Mean dispersal distance in kilometers,

computed as a straight line, between the natal territory and the new established territory after

dispersal observed for the Alpine and the Jura population (Zimmermann, Breitenmoser-Würsten,

& Breitenmoser, 2005).

We removed from the calibration process the calibration of the parameter pRepro and its

associated pattern rRepro as quick tests showed that a pRepro of 0.95 produced a stable mean of

rRepro equal on average to 0.79 [0.78;0.80], independently from the other parameter values.

Among the 50 calibration scenarios, none produced patterns for which the mean values computed

from the simulation outputs fall into the 95% confidence intervals of the reference values for all

patterns simultaneously. We therefore omitted the patterns about dispersal distances for this step

(distDispAlps and distDispJura) as we chose to focus on the mortality process. Without these two

patterns about dispersal distances, 2 out 50 scenarios produced patterns within the confidence

intervals for the 10 remaining patterns (Table C.2).

For the second phase, we fined tuned the calibration of the 6 parameters to calibrate (i.e.,

still omitting rRepro). We explored a narrower range of values for each parameter, which

included the parameter values sampled in the two scenarios that resulted in the best calibrations

in the first phase (Table C.1). Similarly as in the first phase, we ran 50 new calibration scenarios

(with 15 replicates each time) in the same conditions, and computed the same patterns. Again,

none of the scenarios could satisfy the 13 patterns simultaneously. When removing the

reproduction (i.e., not relevant) and the dispersal distances (i.e., too constraining), 25 scenarios

produced patterns with their mean values falling within the confidence intervals of the reference

values for the 10 remaining patterns. We finally selected, among those 25, the scenario which

produced patterns with mean values the closest to the 13 pattern values simultaneously (Table

C.2). The parameter values sampled for this scenario (Table C.1) were the ones selected and used

to run the simulations to produce the results presented in the main text.

1464

Table C.1: Value tested or selected for the parameters to be calibrated. "Phase 1" and "Phase 2"

present the range of the values tested for each parameter for the two phases of the calibration.

One value was randomly sampled within each range for each calibration scenario. "Phase 1. Best

1" and "Phase 1. Best 2" are the values from the best two scenarios of the first phase of

calibration. "Selected" presents the selected values at the end of the phase 2 of the calibration

process.

	pMortResAlps	pMortDispAlps	pMortResJura	pMortDispJura	corrFactorRes	corrFactorDisp
Phase 1	[0.05;0.25]	[0.0001;0.002]	[0.05;0.25]	[0.0001;0.002]	[1;6]	[50;300]
Phase 1. Best 1	0.15	0.0007	0.19	0.0008	3.5	175

Phase 1. Best 2	0.15	0.001	0.12	0.0007	3.5	175
Phase 2	[0.13;0.17]	[0.0005;0.0012]	[0.10;0.21]	[0.0005;0.001]	[3;4]	[125;225]
Selected	0.15	0.0007	0.15	0.0009	3.9	170

1471

Table C.2: Patterns values. "Reference" are values (mean and 95% confidence intervals)

extracted from the literature or field data. "Phase 1. Best 1" and "Phase 1. Best 2" are the pattern

values (mean and 95% confidence intervals over the 15 replicates) computed with the simulation

outputs from the scenarios which were the best ones from the first phase of the calibration

process. "Selected" presents the pattern values (mean and 95% confidence intervals over the 15

replicates) computed with the simulation outputs from the best calibration scenario selected in the

second phase of the calibration process.

	rAllRes_Alps	rAllDisp_Alps	rCollRes_Alps	rCollDisp_Alps	nCollAlps	rAllRes_Jura	rAllDisp_Jura	rCollRes_Jura	rCollDisp_Jura	nCollJura	rRepro	distDispAlps	distDispJura
Reference	0.24 [0.08;0.57]	0.24 [0.15;0.38]	0.02 [0;0.059]	0.077 [0;0.211]	2 [0;3]	0.17 [0.1;0.29]	0.35 [0.16;0.65]	0.034 [0;0.08]	0.067 [0;0.185]	6 [3;9]	0.81 [NA;NA]	26 [15;36]	63 [39;87]
Phase 1. Best 1	0.187 [0.184;0.189]	0.228 [0.226;0.230]	0.0005 [0.0003;0.0006]	0.000003 [0;0.000008]	0.027 [0;0.064]	0.268 [0.264;0.272]	0.312 [0.309;0.315]	0.035 [0.033;0.037]	0.051 [0.050;0.053]	4.5 [4.0;5.0]	0.813 [0.809;0.816]	44.7 [44.1;45.3]	47.3 [46.2;48.4]
Phase 1. Best 2	0.188 [0.185;0.190]	0.309 [0.307;0.312]	0.0004 [0.0002;0.0005]	0.000004 [0;0.000013]	0.013 [0;0.040]	0.183 [0.180;0.187]	0.278 [0.274;0.280]	0.034 [0.032;0.035]	0.054 [0.053;0.056]	6.3 [5.7;6.9]	0.810 [0.806;0.812]	39.4 [38.8;40.0]	56.6 [55.2;57.9]
Selected	0.187 [0.186;0.190]	0.228 [0.226;0.230]	0.0004 [0.0003;0.0006]	0.00002 [0;0.00006]	0.053 [0.001;0.105]	0.222 [0.218;0.226]	0.336 [0.332;0.339]	0.039 [0.037;0.041]	0.051 [0.050;0.053]	4.9 [4.4;5.3]	0.814 [0.810;0.817]	44.2 [43.6;44.8]	50.1 [48.9;51.2]

1479

1480

Sensitivity analysis

The second phase of the calibration was used to perform the sensitivity analysis. Multiple values

were tested for each parameter, values smaller and larger than the ones selected. We compared

the percentage of variation of the parameter values used in the second phase of the calibration

(Fig. C.1) with the percentage of variation on the main result of the simulation, the population

growth rates (Fig. C.2).

1487

1488

1489   **Figure C.1**: Range of values tested during the second phase of calibration for each calibrated

1490   parameter. The value indicated within the range of each parameter was the selected one with the

1491   best calibration scenario. Percentages noted are the difference between the selected value and the

1492   minimum and maximum values of the range for each parameter.

1493

1494   a)

1495   c)

b)

d)

Figure C.2: Annual rates of increase over the simulated years for each population (a) Alpine, b)

Jura, c) Vosges-Palatinian, d) Black Forest). The grey area represents the 2-year burn-in phase at

the beginning of the simulation. Overlaid colored points are mean over the 15 replicates and

overlaid colored envelopes are 95% confidence intervals from the 15 replicates, for each of the

50 calibration scenarios from the second calibration phase. The black point and darker colored

envelope represent the mean and 95 % confidence interval from the 15 replicates for the best

calibration scenario selected during this second calibration phase. The dashed line represents a

stabilization of the population (growth rate equals to 1). The percentage noted above the lines is

the mean over all the years of the differences between the values for the selected scenario (means

over the 15 replicates) and the maximum values of each year (maximum means calculated over

the 15 replicates for the particular scenario this year). The percentage noted below is the same

value but calculated with the minimum values each year.

1508

The growth rates calculated for the Alpine and the Jura population were not sensitive to the

different parameter estimates as the variations in parameter values were much larger than those of

the result values. Parameters varied from -13.3% down to -44.4% and from +2.6% up to 40% in

values (Fig. C.1), whereas the growth rates for these two populations varied on average from -

2.9% up to +1.9% for the Alpine population and from -3.2% up to +3.5% for the Jura population

(Fig. C.2). The variation in the growth rates for the Vosges-Palatinian population were a little

larger (from -8% up to +13%) (Fig. C.2) but still smaller than most of the variations of the

parameters (Fig. C.1). The growth rates for the Black Forest population on the other hand seemed

sensitive to the model parameters with variations from -14.7% up to +30.2% (Fig. C.2). Results

concerning this population were therefore less robust to interpretation. However, this population

had very large confidence intervals regarding it growth rates compared to the other populations

which made the confidence in the population trend already less precise for interpretation (main

text, Fig. 3). The Vosges-Palatinian population and especially the Black Forest one had very few

starting individuals (i.e., more prone to stochasticity) and data from the literature, making the

model harder to calibrate regarding these populations and therefore less reliable to forecast their

future. In comparison, the Alpine and Jura populations had more starting individuals and data

from the literature to calibrate the model and therefore showed a stronger confidence and

precision in the results.

1527

1528

Model validation

Among the 13 patterns used for calibration, the selected set of parameters (i.e., calibration

scenario selected during the second phase) produced simulation results for which the pattern

values of 11 of them fell into the 95% confidence intervals of the reference values (Table C.2).

However, the confidence intervals for the pattern simulated values were extremely narrow

(Tables C.2 and C.3) compared to those of the reference values. Therefore, model stochasticity

did not seem very important. The reproduction rate reference value did not have a confidence

interval available but the simulated value was very close and the reference value was inside the

95% confidence interval of the simulations. We completed this validation phase looking at the

number of simulated collisions for the first few years with the concurrent field data. As the model

was parameterized with 2017 data, we compared the simulated collisions for the Alpine and Jura

populations for the first five years with the recorded collisions in these two populations for 2017

until 2021 (Table. C.3). From the calibration pattern results and this collision validation, we were

very confident in the selected parameter values regarding the demographic parameters (i.e.,

mortality and reproduction rates), and especially for the Jura population.

The simulated collisions data matched less the field data for the Alpine, although the

number of collisions were very low in this population (Table C.3). Also, the only discrepancy

between the reference values and the simulated results for the calibration patterns (Table C.2)

were for the dispersal distances for the Alpine population. The individuals seemed to have

dispersed further in the simulations than computed from the field data. The model may need more

attention on the interpretation for the Alpine population as the mortality may be slightly under-

estimated and dispersal distances slightly over-estimated. Both could be linked as individuals

could move further away as they were dying less doing so.

1552

Table C.3: Number of vehicle collisions with adult or subadult lynx. Collisions with juveniles

were not included in the recorded data as the SE-IBM did not simulate movement for the lynx in

their first year of life and therefore cannot reproduce juvenile collisions. "Recorded" were real

collisions recorded in the field. "Simulated" were collisions simulated with the 100 simulation

replicates using the parameter values selected (mean and 95% confidence interval over the 100

replicates).

	2017	2018	2019	2020	2021
Number of collisions recorded for the Jura population	6	10	4	5	5
Number of collisions simulated for the Jura population	5.7 [5.1;6.2]	5.1 [4.6;5.6]	5.5 [5.1;5.9]	5.3 [4.8;5.8]	4.8 [4.4;5.2]
Number of collisions recorded for the Alpine population	2	0	3	1	3
Number of collisions simulated for the Jura population	0.01 [0;0.03]	0.03 [0;0.06]	0.03 [0;0.06]	0.01 [0;0.03]	0.04 [0;0.08]

1559

1560

1561

1562 **Appendix D – Model validation using telemetry data**

1563

We performed a qualitative validation of the lynx model predictions using tracks from collared

female residents. The map below (Fig. D.1) shows the simulated territory occupancy with darker

red colors representing areas most often occupied by female territories across simulation

replicates and blue the least occupied (main text, Fig. 5). We overlaid GPS and VHF paths from

collared female residents on the map. Data in the Vosges-Palatinate consist in two females in the

Palatinate Forest (Germany) and one in the Vosges Mountains (France) (locations recorded in

2017-2018; Scheid et al., 2021). Data in the Jura represent four females in France and 13 in

Switzerland (locations recorded in 1988-1998; Breitenmoser-Würsten, Zimmermann, et al.,

2007b). Data in the Alps (Switzerland) represent 16 females (locations recorded in 1997-2000;

Breitenmoser-Würsten et al., 2001). We show the full study area and four zoomed areas on the

telemetry locations. Overall, the realized movements and territory emergence inferred from field

data were coherent with the frequencies across simulation replicates of the territories defined by

simulated lynx. Very few telemetry locations fall in white and grey cells (i.e., non-breeding

habitat and breeding habitat cells never selected as part of a female territory across the 100

simulation replicates respectively) and most of the recorded tracks were located in cells with a

high frequency of territory occupancy across the simulation replicates.

1580

1581

Figure D.1: Occupancy by female territories over the study area at the last year of simulation.

Values between above zero and one (color scale blue-yellow-red) are mean occupancy

probability per cell of 1 km^2 over 100 replicates (e.g., a cell equal to 1 means the cell was

included in a female territory, at the 50th year of simulation, in all 100 simulation replicates).

Grey cells are "breeding habitats" never included in a female territory across all replicates (equal

to zero). White cells are cells of a habitat type different than "breeding habitat" and therefore

could never be included in a female territory. GPS and VHF recorded paths for female residents

are overlaid as thin black lines. A zoom of different areas is presented on the sides of the central

map.

1591

1592

References (Appendices)

Ammar, R. (2019). randomcoloR: Generate Attractive Random Colors.

Basille, M., Van Moorter, B., Herfindal, I., Martin, J., Linnell, J. D. C., Odden, J., Andersen, R., & Gaillard, J. M. (2013). Selecting habitat to survive: the impact of road density on survival in a large carnivore. *PLoS One* **8**, e65493.

Bauduin, S., McIntire, E. J. B., & Chubaty, A. M. (2019). NetLogoR: a package to build and run spatially explicit agent□based models in R. *Ecography (Cop.).* **42**, 1841–1849.

Breitenmoser-Würsten, C., Vandel, J.-M., Zimmermann, F., & Breitenmoser, U. (2007a). Demography of lynx *Lynx lynx* in the Jura Mountains. *Wildlife Biol.* **13**, 381–392.

Breitenmoser-Würsten, C., Zimmermann, F., Ryser, A., Capt, S., Laass, J., Siegenthaler, A., & Breitenmoser, U. (2001). Untersuchungen zur Luchspopulation in den Nordwestalpen der Schweiz 1997–2000. *KORA Bericht.*

Breitenmoser-Würsten, C., Zimmermann, F., Stahl, P., Vandel, J.-M., Molinari-Jobin, A., Molinari, P., Capt, S., & Breitenmoser, U. (2007b). Spatial and social stability of a Eurasian lynx *Lynx lynx* population: an assessment of 10 years of observation in the Jura Mountains. *Wildlife Biol.* **13**, 365–380.

Breitenmoser, U., Kavczensky, P., Dötterer, M., Breitenmoser-Würsten, C., Capt, S., Bernhart, F., & Liberek, M. (1993). Spatial organization and recruitment of lynx (*Lynx lynx*) in a re-introduced population in the Swiss Jura Mountains. *J. Zool. London* **231**, 449–464.

Charbonnel, A., & Germain, E. (2020). Plan Régional d'Actions en faveur du Lynx boréal (*Lynx lynx)* dans le Massif des Vosges□ : rétablir le Lynx dans un état de conservation favorable dans le cadre d'une démarche participative, concertée et partagée avec les acteurs du territoire. Lucy (57), France.

Chubaty, A. M., & McIntire, E. J. B. (2018). Package "SpaDES": Develop and Run Spatially

Explicit Discrete Event Simulation Models. https://CRAN.R-project.org/package=SpaDES.

Dowle, M., & Srinivasan, A. (2019). data.table: Extension of "data.frame."

Duchamp, C., Boyer, J., Briaudet, P. E., Leonard, Y., Moris, P., Bataille, A., Dahier, T.,

Delacour, G., Millisher, G., Miquel, C., Poillot, C., & Marboutin, E. (2012). A dual frame

survey to assess time- and space-related changes of the colonizing wolf population in

France. *Hystrix* **23**, 1–12.

Gallagher, C.A., Chudzinska, M., Larsen-gray, A., Pollock, C.J., Sells, S.N., White, P.J.C., &

Berger, U. (2021). From theory to practice in pattern-oriented modelling: identifying and

using empirical patterns in predictive models. *Biol. Rev.* **96**, 1868–1888.

Gatti, S., Blanc, L., Gimenez, O., & Marboutin, E. (2014). Estimation des densités de lynx dans

le massif du Jura entre 2011 et 2014. *Faune Sauvag.* **304**, 4–8.

Gimenez, O., Gatti, S., Duchamp, C., Germain, E., Laurent, A., Zimmermann, F., & Marboutin,

E. (2019). Spatial density estimates of Eurasian lynx (*Lynx lynx*) in the French Jura and

Vosges Mountains. *Ecol. Evol.* **9**, 11707–11715.

Grimm, V., Berger, U., Bastiansen, F., Eliassen, S., Ginot, V., Giske, J., Goss-Custard, J., Grand,

T., Heinz, S. K., Huse, G., Huth, A., Jepsen, J. U., Jørgensen, C., Mooij, W. M., Müller, B.,

Pe'er, G., Piou, C., Railsback, S. F., Robbins, A. M., Robbins, M. M., Rossmanith, E.,

Rüger, N., Strand, E., Souissi, S., Stillman, R. A., Vabø, R., Visser, U., & DeAngelis, D. L.

(2006). A standard protocol for describing individual-based and agent-based models. *Ecol.*

Modell. **198**, 115–126.

Grimm, V., Berger, U., DeAngelis, D. L., Polhill, J. G., Giske, J., & Railsback, S. F. (2010). The

ODD protocol: A review and first update. *Ecol. Modell.* **221**, 2760–2768.

Grimm, V., & Railsback, S.F. (2012). Pattern-oriented modelling: a "multi-scope" for predictive

systems ecology. *Philos. Trans. R. Soc. London - Ser. B.* **367**, 298–310.

Haller, H., & Breitenmoser, U. (1986). Zur Raumorganisation der in den Schweizer Alpen

wiederangesiedelten Population des Luchses *Lynx lynx*. *Z. Saeugetierk* **51**, 289–311.

Henriksen, H. B., Andersen, R., Hewison, A. J. M., Gaillard, J.-M., Bronndal, M., Jonsson, S.,

Linnell, J. D. C., & Odden, J. (2005). Reproductive biology of captive female Eurasian lynx,

Lynx lynx. *Eur. J. Wildl. Res.* **51**, 151–156.

Heurich, M., Schultze-naumburg, J., Piacenza, N., Magg, N., Čeverny, J., Engleder, T.,

Herdtfelder, M., Sladova, M., & Kramer-Schadt, S. (2018). Illegal hunting as a major driver

of the source-sink dynamics of a reintroduced lynx population in Central Europe. *Biol.*

Conserv. **224**, 355–365.

Hijmans, R. J., & Van Etten, J. (2018). Package "raster": Geographic Data Analysis and

Modeling. http://CRAN.R-project.org/package=raster.

Isaac, N. J. B., van Strien, A. J., August, T. A., de Zeeuw, M. P., & Roy, D. B. (2014). Statistics

for citizen science: extracting signals of change from noisy ecological data. *Methods Ecol.*

Evol. **5**, 1052–1060.

Jedrzejewski, W., Jedrzejewska, B., Okarma, H., Schmidt, K., Bunevich, A. N., & Miłkowski, L.

(1996). Population dynamics (1869-1994), demography, and home ranges of the Lynx in

Białowieża Primeval Forest (Poland and Belarus). *Ecography (Cop.).* **19**, 122–138.

Kramer-Schadt, S., Kaiser, T. S., Frank, K., & Wiegand, T. (2011). Analyzing the effect of

stepping stones on target patch colonisation in structured landscapes for Eurasian lynx.

Landsc. Ecol. **26**, 501–513.

Kramer-Schadt, S., Revilla, E., & Wiegand, T. (2005). Lynx reintroductions in fragmented

landscapes of Germany: Projects with a future or misunderstood wildlife conservation? *Biol.*

Conserv. **125**, 169–182.

Kramer-Schadt, S., Revilla, E., Wiegand, T., & Breitenmoser, U. (2004). Fragmented landscapes,

road mortality and patch connectivity: modelling influences on the dispersal of Eurasian

lynx. *J. Appl. Ecol.* **41**, 711–723.

Kramer-Schadt, S., Revilla, E., Wiegand, T., & Grimm, V. (2007). Patterns for parameters in

simulation models. *Ecol. Modell.* **204**, 553–556.

Kvam, T. (1991). Reproduction in the European lynx, *Lynx lynx. Zeitschrift für Säugetierkd.* **56**,

146–158.

López-Bao, J. V., Aronsson, M., Linnell, J. D. C., Odden, J., Persson, J., & Andrén, H. (2019).

Eurasian lynx fitness shows little variation across Scandinavian human-dominated

landscapes. *Sci. Rep.* **9**, 8903.

Outhwaite, C. L., Chandler, R. E., Powney, G. D., Collen, B., Gregory, R. D., & Isaac, N. J. B.

(2018). Prior specification in Bayesian occupancy modelling improves analysis of species

occurrence data. *Ecol. Indic.* **93**, 333–343.

Premier, J., Bastianelli, M.L., Oeser, J., Anders, O., Andren, H., Aronsson, M., Bagrade, G.,

Belotti, E., Breitenmoser-Würsten, C., Bufka, L., Černe, R., Červený, J., Drouet-Hoguet, N.,

Ďuľa, M., Fuxjäger, C., Herdtfelder, M., Hočevar, L., Jędrzejewski, W., Kont, R., Koubek,

P., Kowalczyk, R., Krofel, M., Krojerová-Prokešová, J., Kubala, J., Kusak, J., Kutal, M.,

Linnell, J.D.C, Mattisson, J., Middelhoff, T.L., Melovski, D., Molinari-Jobin, A., Odden, J.,

Okarma, H., Ornicāns, A., Pagon, N., Persson, J., Schmidt, K., Sindičić, M., Slijepčević, V.,

Tám, B., Zimmermann, F., Kramer-Schadt, S., & Heurich, M. (2024). A continental scale

analysis of survival in a large carnivore occupying a human-dominated environment. *Cons.*

Biol.

R Core Team. (2014). R: A language and environment for statistical computing. Vienna, Austria:

R Foundation for Statistical Computing.

Scheid, C., Germain, E., & Schwoerer, M.-L. (2021). Les Lynx (*Lynx lynx*) du Pfälzerwald

s'installent progressivement dans le Massif des Vosges. *Ann. Sci. Rés. Bios. Trans. Vosges*

du Nord. **20**, 104–125.

Stahl, P., & Vandel, J.-M. (1998). Le Lynx boréal : *Lynx lynx* (Linné, 1758). In *Encycl. des*

Carniv. Fr. Société française pour l'Étude et la Protection des Mammifères, Bourges.

Visintin, C., Golding, N., van der Ree, R., & McCarthy, M. A. (2018). Managing the timing and

speed of vehicles reduces wildlife-transport collision risk. *Transp. Res. Part D* **59**, 86–95.

Visintin, C., van der Ree, R., & Mccarthy, M. A. (2017). Consistent patterns of vehicle collision

risk for six mammal species. *J. Environ. Manage.* **201**, 397–406.

von Arx, M., Breitenmoser-Würsten, C., Zimmermann, F., Kunz, F., Vogt, K., Ryser, A., Struch,

M., & Breitenmoser, U. (2017). Der Luchs im Jura - unter besonderer Berücksichtigung des

Solothurner Juras. *Naturforschende Gesellschaft des Kantons Solothurn* **43**, 177–234.

Wickham, H., RStudiom, & R Core Team. (2019). Package "testthat."

Wölfl, S., Anders, O., Middelhoff, T. L., Hohmann, U., Back, M., Idelberger, S., Krebühl, J.,

Ohm, J., Prüssing, A., Herdtfelder, M., Böcker, F., Erretkamps, J., Kopaniak, L., Wölfl, M.,

Jokisch, S., Hucht-Ciorga, I., Teubner, J., Trost, M., Zschille, J., Jeß, E., & Steinberg, C.

(2021). Status des Luchses in Deutschland. *Natur und Landschaft* **96**.

Zimmermann, F., Breitenmoser-Würsten, C., & Breitenmoser, U. (2005). Natal dispersal of

Eurasian lynx (*Lynx lynx*) in Switzerland. *J. Zool.* **267**, 381–395.

Zimmermann, F., Breitenmoser-Würsten, C., & Breitenmoser, U. (2007). Importance of dispersal

for the expansion of a Eurasian lynx *Lynx lynx* population in a fragmented landscape. *Oryx*

41, 358–368.

1710