Early epigenetic influence on EGR1, KLF2 and KLF4 transcriptional network associates with alteration of HSCs quiescence

Running title: Early epigenetic influence on HSPCs

Alexandre Pelletier[1,3,*], Arnaud Carrier[1,3,*], Yong Mei Zhao[2], Mickaël Canouil[1,3], Mehdi Derhourhi[1], Emmanuelle Durand[1], Lionel Berberian-Ferrato[1], John Greally[4], Francine Hughes[5], Philippe Froguel[1,3,6], Amélie Bonnefond[1,3,6], Fabien Delahaye[1,3].

1) Inserm U1283, CNRS UMR 8199, European Genomic Institute for Diabetes, Institut Pasteur de Lille, Lille, France.
2) Department of Obstetrics & Gynecology and Women's Health, Albert Einstein College of Medicine, 1300 Morris Park Ave, Bronx, NY, 10461, USA.
3) Lille University Hospital, University of Lille, Lille, France.
4) Department of Genetics, Albert Einstein College of Medicine, 1301 Morris Park Avenue, Price Building, Room 322, Bronx, New York 10461, USA.
5) Obstetrics & Gynecology and Women's Health, Division of Maternal-Fetal Medicine, Montefiore Medical Center/Albert Einstein College of Medicine, Bronx, NY 10461, USA.
6) Department of Metabolism, Digestion and Reproduction, Imperial College London, London, U.K.

* These authors contributed equally to this work

Corresponding authors:

Fabien Delahaye; Institut Pasteur de Lille-UMR1283, 1 rue du Prof Calmette, 59000 Lille, France ; fabien.delahaye@pasteur-lille.fr; (+33) 320877258

Philippe Froguel; EGID-UMR1283 1 place de Verdun 59045 Lille, France ; philippe.froguel@cnrs.fr; (+33) 374008101

Summary

Excessive fetal growth is associated with DNA methylation alterations in human hematopoietic stem and progenitor cells (HSPC), but their functional impact remains elusive. We implemented an integrative analysis combining single-cell epigenomics, single-cell transcriptomics, and *in vitro* analyses to functionally link DNA methylation changes to putative alterations of HSPC functions. We showed in hematopoietic stem cells (HSC) from large for gestational age neonates that both DNA hypermethylation and chromatin rearrangement target a specific network of transcription factors known to sustain stem cell quiescence. In parallel, we found a decrease expression of key genes regulating HSC differentiation including *EGR1*, *KLF2, SOCS3,* and *JUNB*. Our functional analyses showed that this epigenetic programming was associated with a decreased ability for HSCs to stay quiescent. Taken together, our multimodal approach using single-cell (epi)genomics showed that human fetal overgrowth affects hematopoietic stem cells quiescence maintenance via epigenetic programming.

Introduction

The hematopoietic stem cells (HSC) are involved in essential processes such as inflammation, cardiovascular repair, and immunity throughout the entire life.[1, 2] Thus, alterations in HSC ability to self-renew and to adequately produce differentiated progeny have been suggested to contribute to the onset and progression of age-related diseases such as cancer and cardiovascular diseases.[3, 4] Systemic alterations or the action of various stressors like aging[5, 6] can result in alteration of HSC destiny, and ultimately of hematopoietic functions. The early mechanisms that control their long-term functions in human are not well understood, in part due to the diversity of phenotypes and behaviors of HSCs.[7]

In mice, maternal high-fat diet during gestation limits fetal hematopoietic stem and progenitor cells (HSPC) expansion and repopulation ability while inducing myeloid-biased differentiation.[18] In humans, a limited number of studies have been conducted. Fetal growth was shown to alter the number of circulating CD34+ HSCs[16, 17]. We previously described a global increase of DNA methylation in cord blood derived CD34+ HSPCs from large for gestational age (LGA) infants compared to normal size neonates.[19] Still, the functional impacts of these early epigenetic alterations remain to be elucidated. Such an effort is essential to determine how these epigenetic modifications could mediate the association between early-life exposures and induce persistent life-long functional changes within the hematopoietic system.

We conducted a multimodal analysis combining single-cell epigenomics, single-cell transcriptomics, and *in vitro* analyses to link the DNA methylation alterations observed in LGA neonates with functional alterations in human cord blood derived HSPCs. We developed novel analytical approaches to improve integration of the single-cell

71 epigenomic and transcriptomic data. We found that the DNA methylation alterations

72 observed in LGA neonates are associated with an alteration of the expression of key

73 genes and transcription factors involved in the immediate early gene signalling

74 especially in hematopoietic stem cells as well as in the balance between quiescence

75 and proliferation in HSPCs.

Results

Optimized methylation gene set analysis reveals association between LGA

DNA hypermethylation and stem cell differentiation pathways

To confirm the LGA-associated DNA hypermethylation we previously observed we

significantly increased the power of our analysis. We expanded our original study

trough additional patient inclusions, thereby doubling the size of our cohort.[19] Using,

HELP-tagging assay (HpaII tiny fragment Enrichment by Ligation-mediated PCR), we

generated genome-wide DNA methylation data on 16 CTRL and 16 LGA cord-blood

derived human CD34+ HSPC samples. We independently retrieved in this new

dataset the global DNA hypermethylation initially found in LGA compared to controls[19]

(**Figure 1A**). Then, to increase our detection power, we pooled both dataset and

detected a total of 4815 differentially methylated CpGs (DMC) with 4787 CpGs

hypermethylated and 28 CpGs hypomethylated in LGA (*n*=36) compared to CTRL

(*n*=34, *p*-value<0.001 and |methylation difference|>25%; **Figure 1A, Supplemental**

Table 1). This new set of DMCs was then used throughout following analysis.

As the functional interpretation is performed at the gene level, each CpG (or DMC)

must be linked to a specific gene. Thus, our ability to adequately infer the regulatory

effect of a CpG and its target gene will affect our ability to identify relevant pathways.

Standard analytical approaches usually rely on distance between CpG and

transcription start site (TSS) of the targeted gene and often only consider the top

candidate DMC per gene, not taking into account the cell specific genomic context.

Therefore, we refined CpG-gene association to optimally assess the influence of DNA

methylation changes on gene expression and enhance functional interpretation. We

built a novel gene-methylation score considering 1) the distance between TSS and

CpG; 2) the CpG overlap with expression quantitative trait loci (eQTL) annotation, as

eQTL information allows us to identify tissue specific genomic region link to gene

expression changes; and 3) the regulatory annotation (e.g. Promoter, Enhancer)

based on cell-specific histone marks[25] and on the Ensembl Regulatory database, as

we know that the relationship between change in DNA methylation and change in gene

expression will depend on a cell specific genomic context (**Figure 1B**). We established

756470 CpG-gene associations including 34% of them found thanks to eQTL

annotation. We then summarized the CpG information at gene level, generating a

gene-methylation score for each gene (n=24857, **Supplemental Table 2**). We first

confirmed that the gene-methylation score properly recapitulates the influence of key

parameters in DMC analysis such as significance and effect size of the methylation

change, number of DMCs per gene, distance from TSS, as well as promoter and

enhancer localization (**Figure 1C**). We also confirmed that while preserving key

information from standard methylation metrics, the gene-methylation score presented

better association with DEGs than significance or methylation change alone. Thus, the

gene-methylation score appears as a better predictor of the methylation influence on

gene expression (**Figure 1C**). We then used our gene-methylation score to perform

pathway enrichment analysis and data integration especially considering integration

with gene expression data.

Using the gene ontology (GO) reference database, we performed methylation gene-

set enrichment analysis (GSEA) based on the gene-methylation score. We found that

change in DNA methylation in LGA HSPC samples targeted genes involved in

signaling regulating fetal development as well as in key stem cell pathways such as

Wnt signaling, cell fate specification, and cell fate commitment pathways (adjusted *p*-

value<0.01, **Figure 1D**) confirming previous findings[19].

Single-cell transcriptomic analysis confirms alteration of hyper-methylated

genes in pathways regulating stem cell differentiation among LGA HSCs

To identify genes altered in LGA and get further biological insight into the functional

consequences of the DNA methylation modifications observed in LGA, we performed

single-cell transcriptomic analysis comparing CTRL and LGA HSPCs.

To enable lineage specific transcriptomic analysis, we created a hematopoietic

reference map (*i.e.* hematomap) by integrating data generated from cord blood

derived CD34+ HSPC cells (*n*=18520) from 7 control neonates (**Figure 2A**). Based on

cluster specific gene expression, we identified 18 distinct clusters representative of

major lineages (Long-Term HSC, HSC, Multi-Potent progenitor, Lymphoid, Myeloid,

and Erythroid) of the hematopoietic compartment (**Figure 2B**, **Supplemental Figure**

2). Each cluster was annotated using cell type specific markers. Markers were then

ranked based on their expression fold change and the specificity for the cluster. Top

cluster-specific markers were compared with published cell type-specific genes[27-30]

(**Supplemental Table 3**). Candidate cell subpopulations were distributed as follows:

1% LT-HSC (*ID1*); 24% HSC (*AVP*); 45% MPP/LMPP (*CDK6*); Lymphoid (*CD99,*

LTB); 1% B cell (*IGHM*); 1% T cell (*CD7*); 14% Erythro-Mas (*GATA1*); <1%Mk/Er

(*PLEK, HBD*); 8% Myeloid (*MPO*); <1% DC (*CST3*, *CD83*).

To identify differentially expressed genes (DEG) between CTRL and LGA samples,

we implemented the Hash Tag Oligonucleotide (HTO) multiplexing strategy[31] allowing

simultaneous processing of CTRL and LGA samples. Multiplexing is a means to limit

the influence of technique-drive batch effects at every stage of the analysis to improve

biological relevance of the finding. We generated multiplexed single-cell transcriptomic

data from 6 LGA (*n*=6861 cells) and 7 CTRL (*n*=5823 cells) samples. We observed in

LGA a shift toward downregulated genes (**Supplemental Figure 3**) especially in the

HSC subpopulation (*n*=285 downregulated genes over 373 DEGs, adjusted *p*-

value<0.05 and log$_2$FC<(-0.5), **Figure 2C; Supplemental Table 4**). Notably, the well-

known *EGR1*, *JUNB,* and *KLF2* genes were among the top affected genes. Using GO

enrichment analysis, we found that downregulated genes were enriched in growth

related pathway (e.g. regulation of growth) as well as in stress related biological

process (e.g. response to temperature stimulus, cellular response to chemical stress;

Figure 2D, adjusted p-value<0.05).

To assess if these HSC specific transcriptomic changes may be associated with

epigenetic changes, we integrated bulk DNA methylation with single-cell gene

expression data using the gene-methylation score. We found that DEGs, and

particularly the down-regulated genes, mostly showed high gene-methylation score

(**Figure 3A, Figure 3B**). We then assessed the association between changes in DNA

methylation and gene expression at pathway level. We looked for enrichment for

differentially methylated genes considering pathways that were identified based on

DEGs. We found a significant overlap between GO terms enriched in LGA HSC

downregulated genes and GO terms enriched in hypermethylated genes (10 out of 46;

p-value <0.05, hypergeometric test). Among the overlapping terms, we found the

module "regulation of growth" including notably *SOCS3*, *SIRT1* and *SESN2* genes that

are both downregulated and within the top 10% of hypermethylated genes (**Figure**

3C).

DNA methylation changes occurs in HSCs and DEGs associated open

chromatin regions

We profiled chromatin accessibility at single-cell level (i.e. single-cell ATAC-seq) to

optimize the resolution of our epigenomic analyses. We generated open chromatin

data across 8733 cells in HSPCs from 6 CTRL and 5 LGA neonates. We first

annotated subpopulations using label transfer approach between ATAC-seq data and

the lineage labels from the Hematomap (**Figure 4A, Supplemental Figure 4A**). To

validate the relevance of our lineage annotation, we performed TF motif enrichment

and observed that lineage specific peaks were effectively associated with well-known

lineage specific TF (**Supplemental Figure 4B**).

We then integrated our bulk DNA methylation data with our single-cell ATAC-seq data

to assess DMCs distribution within open chromatin regions (OCRs). Overall, 31% of

the 211479 peaks contain CpGs queried by our genome wide methylation assay. We

first observed a strong enrichment for DMCs in OCRs with 74% of them located in

OCRs compare to only 34% of overall queried CpGs (pvalue < 0.001, hypergeometric

test). Such enrichment supports the putative regulatory influence of our DMCs. By

performing lineage specific analysis, we observed a DMCs enrichment in HSC-specific

open chromatin region with a total of 11% of HSC specific peaks containing DMCs

(adjusted *p*-value<0.001, **Figure 4B**), while no enrichment was observed for the others

lineages. This result corroborates the HSC specific transcriptional impact of the DNA

methylation change observed in LGA. Furthermore, we observed that DEGs in LGA

HSC and especially down regulated genes were enriched for OCRs containing DMCs

(**Figure 4C**).

Not limiting our analysis to the regulatory role of DMCs within open chromatin regions,

we then assessed the interaction between change in DNA methylation, change in

expression, and change in chromatin accessibility among LGA samples. We identified

278 open chromatin regions that significantly differ between LGA and CTRL in HSCs,

with 215 showing decreased and 63 showing increased accessibility in LGA versus

CTRL (adjusted *p*-value<0.001 and |log$_2$FC|>0.25, **Supplemental Figure 4C**). Motif

enrichment analysis identified EGR1, KLF2 and KLF4 among the top motif enriched

in regions with decreased accessibility (**Figure 4D, Supplemental Figure 4D**).

Regions with decreased accessibility were also strongly enriched in peaks including

DMCs and peaks associated with DEGs (**Figure 4E**), with 3-fold and a 2.5-fold

enrichment respectively. Finally, these regions were strongly enriched for peak

containing both DMCs and associated to DEGs (23-fold enrichment) illustrating that

early epigenetic programming is actually not limited to changes in DNA methylation

but also involves chromatin rearrangement targeting altered genes.

EGR1, KLF2 and KLF4 are key upstream regulators influenced by early

epigenetic programming in LGA

To further characterize the molecular mechanisms affected in LGA HSCs and

identified master regulators, we leveraged the single-cell resolution of our approaches

to perform a co-regulatory network analysis. This approach allowed us to model the

influence of upstream transcription factors (TF) on expression changes of downstream

target genes. We performed co-expression analysis to identify co-regulated genes, *i.e.*

regulons, and performed analysis of cis-regulatory motif to assign TF to each regulon

(SCENIC). We identified a total of 250 regulons but only considered for further

analyses the 106 regulons identified based on high confidence cis-regulatory motif.

These regulons only rely on associations for which the presence of TF motif was

validated. We scored the activity of regulons in each cell and identified lineage specific

regulons associated with key hematopoietic TFs such as GATA2, GATA3, MEIS1,

TAL1, TCF3, EGR1, CEBPB, HOXB4, SPI1 and STAT1/3 further supporting our

subpopulation annotation and the SCENIC approach (**Supplemental Figure 5A,**

Supplemental Table 5)[39].

To identify TF associated with the changes in gene expression observed in LGA HSC,

we compared the regulon activity between CTRL and LGA. We found seven regulons

with a significant decrease in activity in the LGA HSC population (adjusted *p*-

value<0.001 and |activity score fold change|>10%, **Supplemental Table 6**). No

regulons were upregulated. These regulons were associated to ARID5A, EGR1,

KLF2, KLF4, KLF10, FOSB, and JUN (**Figure 5A**). Among them, ARID5A, EGR1,

KLF2, FOSB and JUN were part of the 10 top active regulons in HSCs (**Supplemental**

Table 7). Based on functional enrichment analysis using as reference GO:BP gene

sets, and HSC signatures of quiescence or proliferative state[40], we showed that these

regulons were enriched in genes regulating stress response, proliferation, and HSC

differentiation (**Figure 5B**).

In parallel and to further support the association between change in DNA methylation

and change in gene expression previously identified at gene level, we performed

GSEA analysis to identify regulon enriched for both differentially methylated and

differentially expressed genes. We found 9 regulons enriched in both hypermethylated

and downregulated genes (adjusted *p*-value<0.01 and NES < -1.6), including the

differentially active regulons ARID5A, EGR1, FOSB, JUN, KLF2, and KLF4 (**Figure**

5C). We also found 9 regulons enriched in hypermethylated and upregulated genes

(adjusted *p*-value<0.01 and NES >1.6) with key HSPC specific regulons such as SPI1

promoting myeloid differentiation;[41] and HOX family (HOXA9, HOXA10, HOXB4)

promoting HSPC expansion (**Figure 5C**).[42-44]

To confirm the putative influence of methylation change on TF activity, we performed

TF motif analysis considering the proximal regions surrounding each DMCs (±20bp).

We found significant enrichment for 23 TF motifs (adjusted *p*-value<0.05, **Figure 5D**).

Among them, we found EGR1 and several members of the Kruppel-like factors (KLF)

family: KLF14, KLF5, KLF1, KLF6. Furthermore, by taking advantage of our single-cell

ATAC-seq data, we looked at the enrichment for TF motif in open chromatin region of

HSC containing DMCs. We found a strong enrichment in EGR1, KLF2, and KLF4

motifs indicating that DNA methylation change occurred in active regions of the

EGR1/KLF2/KLF4 TF network (**Figure 5E**).

Multimodal co-regulatory network recapitulating TF-gene interactions

influenced by early epigenetic programming in LGA

Based on the integration of the DNA methylation, single-cell ATAC-seq, and single-

cell RNA-seq data, we built a network recapitulating interaction between mains TFs

and downstream target genes within the principal regulons altered in LGA neonates:

EGR1, KLF2, and KLF4 (**Figure 6**). EGR1, KLF2, and KLF4 relies on highly

interconnected genes (**Figure 6A**). For each target gene, we confirmed the presence

of unique or shared upstream TF binding motif within the open chromatin regions. We

observed a high concordance between the regulons and open chromatin motif

analysis: 96%, 91% and 95% of genes included in EGR1, KLF2, and KLF4 regulons

respectively were associated to at least one peak containing the corresponding TF

motif supporting the association between genes and TFs. We then looked for evidence

of epigenetic modifications that may alter TF-target interactions. We annotated genes

with associated open chromatin regions containing at least one DMC (middle area) or

identified as differentially accessible between CTRL and LGA (inside area) (**Figure**

6B). Overall, 23% (n=27) of genes targeted by these TFs networks have epigenetic

alteration (DMCs or decrease accessibility) in open chromatin regions while 22%

(n=26 genes) appear downregulated in LGA. Finally, we highlight KLF2 as possible

master regulators influenced by early programming. Indeed, we identified KLF2 as

hypermethylated and down regulated gene that interacts directly with EGR1 and KLF4

suggesting downstream influence of KLF2 on these TFs. Conversely, KLF2 was not

identified as part of EGR1 and KLF4 regulons suggesting that KLF2 is not a target of

these TFs. This network also further validated *JUNB and SOCS3* being highly

epigenetically altered in cis-regulatory regions **(Figure 6C)***,* as well as *ID1, CDKN1A,*

IER2, IER3, and *IER5* as key downstream altered targets of KLF2, EGR1, and/or

KLF4, again highlighting how early programming alters signaling involved in the

regulation of cell proliferation and differentiation.

***In vitro* analysis confirms the alteration of HSPCs differentiation capacities in**

LGA

Our integrative analyses highlighted epigenetic and transcriptomic alterations

targeting signaling pathways involved in regulation of HSC differentiation and

proliferation. Thus, we decided to challenge HSPC differentiation and proliferation

potential *in vitro* using colony forming unit (CFU) assays. After 14 days of expansion,

colonies from 4 CTRL and 4 LGA samples were classified in 3 categories: those

derived from common myeloid progenitors (CFU_GEMM), erythroid progenitors (BFU-

E), and granulocyte-macrophage progenitors (CFU_GM) based on the morphology of

each colony. We observed a significant decrease in the number of common myeloid

progenitor colonies in LGA samples (*p*-value<0.05; **Figure 7A**) as well as striking

differences in shape and size of more differentiated colonies (**Figure 7B**).

CFU_GEMM colonies are the product of a non-committed hematopoietic progenitor

able to self-renew and differentiate in both erythroid and myeloid lineage. In our

samples, only HSC have these features, suggesting that the decreased CFU GEMM

proportion in LGA reflects either fewer HSC in starting cell subpopulation composition

or a decreased proliferation and differentiation capacity. To evaluate these two

possibilities, we monitored cell populations distribution across conditions at molecular

resolution using our single-cell expression dataset. We observed a decrease in HSC

cells (*p*-value=0.015) and a trend toward increased MPP cells (*p*-value=0.13, **Figure**

7C) in LGA compared to CTRL.

Another way to look at population shift is to use pseudotime, *i.e.* a measure that

reflects how far an individual cell is in a differentiation process. Indeed, cord-blood

derived CD34+ HSPCs represent a heterogeneous population of cells ranging from

progenitors to progressively restricted cells of the erythroid, myeloid, or lymphoid

lineages as confirmed by our single-cell transcriptomic analysis. To follow cell

distribution through these levels of differentiation and assess the influence of the LGA

environment we used pseudotime tool from Monocle.[32] Collecting the pseudotimes

across our different cell populations, we observed a positive correlation between

pseudotime and lineage differentiation as expected (*r*=0.99, Pearson correlation,

Figure 7D, **Supplemental Figure 6A**). We then compared the distribution of the

pseudotime between LGA and CTRL using the least differentiated cells as roots, i.e.

the long-term HSCs. At the population level, we observed an increase in pseudotime

in LGA (*p*-value<0.001, **Figure 7E**). Indeed, we observed a decrease in number of

cells presenting pseudotime associated with the HSC state in LGA samples (*p*-

value<0.05) and a shift toward cells presenting elevated pseudotime suggesting that

LGA HSCs exit quiescence and differentiate more quickly compared to CTRL HSCs

(**Figure 7E**). Altogether, our analysis supports the association between LGA exposure

and cell growth signaling targeted by DNA methylation and gene expression changes

with alteration of differentiation and proliferation capacities.

Discussion

Here, we interrogated three major layers of the regulatory landscape in cord-blood

derived CD34+ HPSCs, DNA methylation, chromatin conformation, and gene

expression. We characterized, in depth and at single-cell resolution, the functional

consequences associated with early DNA methylation changes observed in LGA

neonates. Through, the integration of multiple datasets and the development of novel

analytical approaches, we addressed a very challenging aspect of functional

(epi)genomics, the interpretation of DNA methylation changes. Focusing on HSPCs,

we believe that we contributed to a better understanding of how early environment

shapes the hematopoietic compartment development and long-term function.

We demonstrated in LGA neonates, a correlated increase of DNA methylation and

change in chromatin accessibility associated with decrease expression of downstream

target genes under the influence of key HSC transcription factors EGR1, KLF2, and

KLF4. EGR1, KLF2 and KLF4 are zinc-finger transcription factors involved in HSC

quiescence and differentiation. EGR1 has a known role in regulating cell growth,

development, and stress response in many tissues. In HSPC, EGR1 plays a role in

homeostasis of HSCs regulating proliferation[38]. EGR1 promotes quiescence and

decreases through differentiation. KLF family is implicated in key stem cell functions.

KLF4 is the most well-known factor of this family due to its role in reprogramming

somatic cells into induced pluripotent stem cells[47]. KLF4 has been identified as a target

for PU.1 transcription factor required for lineage commitment in HSPCs[48]. KLF2 and

KLF4 promote self-renewal in embryonic stem cells[34] but no study has looked

specifically at KLF2 and KLF2/KLF4 interactions in HSPCs. Our data suggests direct

and indirect (shared downstream target) interactions between these 3 transcriptions

factors in HSPCs. EGR1, KLF2, and KLF4 represent targets to be further explored in

order to challenge causality. Still, our findings lead to a better understanding of how

early exposure can affect long-term hematopoietic maintenance in human via

epigenetic programming of the EGR1, KLF2, and KLF4 signaling.

Interestingly, EGR1, KLF2, and KLF4 are not only involved in the regulation of

proliferation and differentiation *per se* but are key factors of the immediate early

response involved in stimulation-related cell activation. *EGR1* and *KLF2* expression

increase in response to extrinsic stimulation. Elevated *EGR1* and *KLF2* expression

promote self-renewal and quiescence in HSC[38,34]. Our transcriptomic data suggests

that such activation may be occurring in our samples with the activation of stress-

related signaling. The primary scope of our study was not to characterize the

environmental exposure that would trigger such response. However, one can

speculate that the activation could result from stress due to extended cold exposure

and handling time inherent to sample preparation. Still, the decrease activity observed

in LGA suggests that LGA HSCs capacities to respond to environmental challenges

are diminished. The alteration of HSCs capacity to respond to extrinsic stimulation fits

with the concept of early programming in which disease susceptibility relies not only

on early impairment of organ development but also on a decreased adaptability to

further environmental challenges to trigger disease. Indeed, fine-tuning of HSC

quiescence mechanisms is of crucial relevance for correct hematopoiesis. Not

responsive dormant HSC would lead to hematopoietic failure due to a lack of

differentiated blood cells. While highly responsive HSC would get to exhaustion of the

population and lack of long-term maintenance of the hematopoietic system.[46]

To validate findings from our integrative approach, we challenged HSPCs *in vitro* and

found significant decrease in common myeloid progenitors. These alterations can be

associated to difference in HSC population between condition, as the number of HSCs

seeded will influence the number of colonies generated, or to decreased differentiation

and proliferation capacities of HSC. Our data suggest that both are altered in LGA.

Indeed, the cell population analysis at transcriptomic level revealed a decrease in

HSCs in LGA neonates. While we also observed epigenomic and transcriptomic

alteration of signaling pathways and transcription factors involved in proliferation and

differentiation of HSCs. Yet, loss of stemness capacities in HSC is likely to drive the

decrease in HSC subpopulation observed in our data as supported by our *in silico*

differentiation analysis (pseudotimes). It showed that LGA HSCs appeared to

differentiate faster than their control counterpart, thus reducing the pool of HSCs in

the HSPCs population. These findings corroborate previous studies on developmental

programming of the hematopoietic system.[16,17] A reduction in self-renewal of HSPCs

and increase differentiation in both lymphoid and myeloid lineages have been

observed in a mouse model of maternal obesity.[18] These effects may drive long-term

consequences in human health as illustrated by the study performed by Kotowski et

al. in which the integrity of the hematopoietic system in neonates was associated with

susceptibility to onset of hematopoietic pathologies.[49]

Hematopoietic stem cell differentiation and self-renew rely on a synergic interplay

between genetically encoded signaling, cell-intrinsic and cell-extrinsic factors as well

as epigenetic modifiers.[45] This interplay appears altered in LGA neonates. We here

provide a comprehensive model recapitulating the functional influence of the

epigenetic early programming on HSPCs fitness to later environmental exposure

(**Figure 7F**). We also linked LGA associated epigenetic modifications to gene

expression and functional alterations through novel integrative approach. In this

regard, we identified targets to be further explored. We also brought a better

understanding of how early exposure can affect long-term tissue maintenance via

epigenetic programming of EGR1, KLF2, and KLF4 early immediate response

signaling associated TFs.

Methods

See the Supplemental Methods for additional information.

Clinical sample collection. Cord blood samples were obtained from CTRL and LGA

neonates. LGA were defined by birth weight and ponderal index values greater than

the 90th percentile adjusted for gestational age and sex. Control infants had normal

parameters (between 10th and 90th percentiles) for both birth weight and ponderal

index. Maternal and infant characteristics are shown in Supplemental Table 8.

Isolation of CD34+ HSPCs. Mononuclear cells were separated using PrepaCyte-

WBC following which CD34+ cells were obtained by positive immunomagnetic bead

selection, using the AutoMACS Separator (Miltenyi Biotech). Cells were

cryopreserved in 10% dimethyl sulfoxide using controlled rate freezing upon analysis.

Genome-Wide DNA methylation assay. DNA methylation levels for >1.7M CpGs

were obtained using the HELP-tagging assay as previously described.[50]

Single-cell RNA sequencing libraries preparation. After cell count and viability

check, cell suspension was loaded into the Chromium 10x Genomics controller and

library was generated using the chromium single-cell v3 chemistry following

manufacturer recommendations. Gene expression library was sequenced using

100bp paired-end reads on the Illumina NovaSeq 6000 system.

Single-cell ATAC sequencing libraries preparation. After cell count and viability

check, nuclei were isolated from cell suspension and incubated with transposase.

Transposed nuclei were then loaded into the Chromium 10x Genomics controller and

library was generated using the chromium single-cell ATAC v1.1 chemistry following

manufacturer recommendations. Gene expression library was sequenced using

150bp paired-end reads on the Illumina NovaSeq 6000 system.

HTO protocol. After cell counting and viability check and prior to cell suspension

loading on the Chromium 10x Genomics controller, cell hashtag (HTO) staining was

used following the cell-hashing protocol.[31]

Colony Forming Unit Assay. To assess clonogenic progenitor frequencies, 3×10^4

CD34+ HSPC cells were plated in methylcellulose containing SCF, GM-CSF, IL-3, and

EPO (H4434; STEMCELL Technologies). Colonies were scored 14 days later.

Data processing and statistical analysis. For DNA methylation analysis, low quality

CpGs were filtered out based on detection rate and confidence score. 754931 out of

1709224 CpGs were conserved for further analysis. Linear regression and statistical

modeling using LIMMA R package[51] were used to identify differentially methylated

CpGs (DMC). We assessed enrichment for biological pathways performing GSEA

using the ClusterProfiler package.[52] We performed transcription factor (TF) motif

enrichment analysis using HOMER tool[53] considering a 20bp region around the DMCs.

For single-cell RNAseq (scRNA-seq) analysis, data were preprocessed using the

CellRanger count pipeline. Data filtering, normalization and integration as well as

cluster identifications were performed using Seurat (v4) pipeline. Pseudo-bulk

differential expression analysis between LGA and CTRL cells within each

hematopoietic lineage was performed using DESeq2 R package.[54] Over

representation test was performed on differentially expressed genes (DEGs) using

enrichGO and enrichKEGG of the ClusterProfiler Package. The SCENIC workflow[55]

was used to identify co-regulated modules associated to a TF (regulons) and to

generate cell-specific activity score for each regulon. Differentiation trajectory analysis

and pseudotime attribution were conducted with Monocle.[32]

For single-cell ATAC-seq, data were preprocessed using the CellRanger ATAC

pipeline. Data filtering, normalization and integration as well as clustering were

performed using Signac pipeline. Cell type identification was based on scRNA-seq

annotation using a label transfer approach. Peaks calling at lineage level was

performed using MACS2 tool. Peaks specific to each lineage or differentially

accessible between LGA and Control were identified using FindMarkers function with

Logistic Regression (LR) models including cellular sequencing depth as latent

variable. TF motif enrichment on lineage or group specific peaks was performed using

FindMotifs function. All peaks enrichment analysis was performed using

hypergeometric tests. For final Gene Regulatory Network (GRN) construction, TF

target interactions inferred with SCENIC was filtered out based on presence of

corresponding TF motif in the peak associated to the target. Supplemental Table 9

contains information on number of cells per sample.

Gene-methylation score

To compute the gene-methylation score, 2 steps were needed: 1) to generate a CpG

score that reflects the association between CpG and gene, and 2) to concatenate

CpG-scores at gene level.

1) <u>CpG-score</u>

CpGScore = (-log10(p_{cpg}) × meth.change) × LinkWeight × RegWeight

Where p_{cpg} is the nominal *p*-value of the differential methylation analysis, and

meth.change is the difference between the percentage of methylation in LGA and the

percentage of methylation in CTRL. LinkWeight represents the confidence in CpG-

gene association and RegWeight represents the estimate regulatory influence of the

considered CpG based on CD34+ specific genomic annotation defined using CD34+

specific histone marks as previously described[4] and EnsRegScore refers to regulatory

regions defined based on the Ensembl Regulatory build hg19 genome annotation.[5]

<u>2) To concatenate CpG-Scores at gene level: gene-methylation score</u>

To summarize the CpG methylation change at the gene level, we aggregated the CpG-

Scores into a methylation gene-score by taking care of i) alleviate the arbitrary number

of CpGs per gene and ii) interpret differently CpG influences located on the promoter

of them in others genomic region.

The gene-methylation score is defined as:

Gene-methylation score $= (\sum CpG\ Score \times Weight\ _{n\_cpg})_{promoter} + (\sum |CpG\ Score| \times Weight\ _{n\_cpg})_{other\_regions}$

Where the Weight_ncpg was optimized to alleviate the influence of the number of CpGs

linked to a gene and defined as:

480
$$\text{Weight}_{nCpG} = \sqrt[3,8]{\frac{1}{\sum \frac{1}{|CpGScore+1|}}}$$

482 The code to perform the analyses in this manuscript is available at

483 https://github.com/umr1283/LGA_HSPC_PAPER.git.

Declarations

Ethics approval. This study was approved by the Institutional Review Board of the

Montefiore Medical Center and the Committee on Clinical Investigation at the Albert

Einstein College of Medicine and is in accordance with Health Insurance Portability

and Accountability Act regulations. Written informed consent was obtained from all

subjects before participation.

Acknowledgments. The authors thank the UMR 8199 LIGAN-PM Genomics platform

(Lille, France) which belongs to the 'Federation de Recherche' 3508 Labex EGID

(European Genomics Institute for Diabetes; ANR-10-LABX-46) and was supported by

the ANR Equipex 2010 session (ANR-10-EQPX-07-01; 'LIGAN-PM'). The LIGAN-PM

Genomics platform (Lille, France) is also supported by the FEDER and the Region

Nord-Pas-de-Calais-Picardie. This project is cofunded in the frame of CPER CTRL

program by the European Union - European Regional Development Fund (ERDF),

Hauts de France Region (contract n°17003781), Métropole Européenne de Lille

(contract n°2016_ESR_05), and French State (contract n°2017-R3-CTRL-Phase 1).

The present work was also supported by the National Center for Precision Diabetic

Medicine – PreciDIAB, which is jointly supported by the French National Agency for

Research (ANR-18-IBHU-0001), by the European Union (FEDER), by the Hauts-de-

France Regional Council and by the European Metropolis of Lille (MEL) and by the

European Research Council (ERC Reg-Seq – 715575).

Support for this project was provided by the Roadmap Epigenomics Program, R01

HD063791 (Einstein/Greally). Support was also provided by Einstein's Center for

Epigenomics, including the Epigenomics Shared Facility and Computational

Epigenomics Group.

Author contributions. AP, AC, YMZ, ED, LBF, and FD were responsible for

conducting research and analyzing data. MD and MC provided feedback on the data

analysis. AP, AC, FD, AB, and PF contributed to writing the manuscript. FD, FH, and

JG were responsible for designing the study.

Competing Interest. The authors declare no competing financial interests in relation

to the work described.

References

1. Doulatov S, Notta F, Laurenti E, Dick JE. Hematopoiesis: a human perspective*. Cell
Stem Cell*. 2012; **10**(2): p. 120-36.
2. Eaves CJ. Hematopoietic stem cells: concepts, definitions, and the new reality*. Blood*.
2015; **125**(17): p. 2605-13.
3. Balistreri CR, Garagnani P, Madonna R, Vaiserman A, Melino G. Developmental
programming of adult haematopoiesis system*. Ageing Res Rev*. 2019; **54**: p. 100918.
4. Pardali E, Dimmeler S, Zeiher AM, Rieger MA. Clonal hematopoiesis, aging, and
cardiovascular diseases*. Exp Hematol*. 2020; **83**: p. 95-104.
5. Lee CC, Fletcher MD, Tarantal AF. Effect of age on the frequency, cell cycle, and lineage
maturation of rhesus monkey (Macaca mulatta) CD34+ and hematopoietic progenitor
cells*. Pediatr Res*. 2005; **58**(2): p. 315-22.

6.   Chandel NS, Jasper H, Ho TT, Passegue E. Metabolic regulation of stem cell function in
tissue homeostasis and organismal ageing. *Nat Cell Biol*. 2016; **18**(8): p. 823-32.
7.   Guenechea G, Gan OI, Dorrell C, Dick JE. Distinct classes of human stem cells that differ
in proliferative and self-renewal potential. *Nat Immunol*. 2001; **2**(1): p. 75-82.
8.   Barker DJ, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart
disease in England and Wales. *Lancet*. 1986; **1**(8489): p. 1077-81.
9.   Barker DJ, Osmond C, Kajantie E, Eriksson JG. Growth and chronic disease: findings in
the Helsinki Birth Cohort. *Ann Hum Biol*. 2009; **36**(5): p. 445-58.
10. Cnattingius S, Villamor E, Lagerros YT, Wikstrom AK, Granath F. High birth weight and
obesity--a vicious circle across generations. *Int J Obes (Lond)*. 2012; **36**(10): p. 1320-4.
11. Wang Y, Gao E, Wu J, et al. Fetal macrosomia and adolescence obesity: results from a
longitudinal cohort study. *Int J Obes (Lond)*. 2009; **33**(8): p. 923-8.
12. Eriksson J, Forsen T, Tuomilehto J, Osmond C, Barker D. Size at birth, childhood growth
and obesity in adult life. *Int J Obes Relat Metab Disord*. 2001; **25**(5): p. 735-40.
13. Harder T, Rodekamp E, Schellong K, Dudenhausen JW, Plagemann A. Birth weight and
subsequent risk of type 2 diabetes: a meta-analysis. *Am J Epidemiol*. 2007; **165**(8): p.
849-57.
14. Pocobelli G, Dublin S, Enquobahrie DA, Mueller BA. Birth Weight and Birth Weight for
Gestational Age in Relation to Risk of Hospitalization with Primary Hypertension in
Children and Young Adults. *Matern Child Health J*. 2016; **20**(7): p. 1415-23.
15. Koyanagi A, Zhang J, Dagvadorj A, et al. Macrosomia in 23 developing countries: an
analysis of a multicountry, facility-based, cross-sectional survey. *Lancet*. 2013;
381(9865): p. 476-83.
16. Al-Sweedan SA, Musalam L, Obeidat B. Factors predicting the hematopoietic stem
cells content of the umbilical cord blood. *Transfus Apher Sci*. 2013; **48**(2): p. 247-52.
17. Chandra T, Afreen S, Kumar A, Singh U, Gupta A. Does umbilical cord blood-derived
CD34+ cell concentration depend on the weight and sex of a full-term infant? *J Pediatr
Hematol Oncol*. 2012; **34**(3): p. 184-7.
18. Kamimae-Lanning AN, Krasnow SM, Goloviznina NA, et al. Maternal high-fat diet and
obesity compromise fetal hematopoiesis. *Mol Metab*. 2015; **4**(1): p. 25-38.
19. Delahaye F, Wijetunga NA, Heo HJ, et al. Sexual dimorphism in epigenomic responses
of stem cells to extreme fetal growth. *Nat Commun*. 2014; **5**: p. 5187.
20. Stella CC, Cazzola M, De Fabritiis P, et al. CD34-positive cells: biology and clinical
relevance. *Haematologica*. 1995; **80**(4): p. 367-87.
21. Thurman RE, Rynes E, Humbert R, et al. The accessible chromatin landscape of the
human genome. *Nature*. 2012; **489**(7414): p. 75-82.
22. Delahaye F, Do C, Kong Y, et al. Genetic variants influence on the placenta regulatory
landscape. *PLoS Genet*. 2018; **14**(11): p. e1007785.
23. Dong D, Tian Y, Zheng SC, Teschendorff AE. ebGSEA: an improved Gene Set Enrichment
Analysis method for Epigenome-Wide-Association Studies. *Bioinformatics*. 2019;
35(18): p. 3514-3516.
24. Ren X, Kuan PF. methylGSA: a Bioconductor package and Shiny app for DNA
methylation data length bias adjustment in gene set testing. *Bioinformatics*. 2019;
35(11): p. 1958-1959.
25. Wijetunga NA, Delahaye F, Zhao YM, et al. The meta-epigenomic structure of purified
human stem cell populations is defined at cis-regulatory sequences. *Nat Commun*.
2014; **5**: p. 5195.

26. Buniello A, MacArthur JAL, Cerezo M, et al. The NHGRI-EBI GWAS Catalog of published
genome-wide association studies, targeted arrays and summary statistics 2019.
Nucleic Acids Res. 2019; **47**(D1): p. D1005-D1012.
27. Cabezas-Wallscheid N, Klimmeck D, Hansson J, et al. Identification of regulatory
networks in HSCs and their immediate progeny via integrated proteome,
transcriptome, and DNA methylome analysis. *Cell Stem Cell*. 2014; **15**(4): p. 507-522.
28. Karamitros D, Stoilova B, Aboukhalil Z, et al. Single-cell analysis reveals the continuum
of human lympho-myeloid progenitor cells. *Nat Immunol*. 2018; **19**(1): p. 85-97.
29. Zhang YH, Hu Y, Zhang Y, Hu LD, Kong X. Distinguishing three subtypes of
hematopoietic cells based on gene expression profiles using a support vector machine.
Biochim Biophys Acta Mol Basis Dis. 2018; **1864**(6 Pt B): p. 2255-2265.
30. Zheng S, Papalexi E, Butler A, Stephenson W, Satija R. Molecular transitions in early
progenitors during human cord blood hematopoiesis. *Mol Syst Biol*. 2018; **14**(3): p.
e8041.
31. Stoeckius M, Zheng S, Houck-Loomis B, et al. Cell Hashing with barcoded antibodies
enables multiplexing and doublet detection for single cell genomics. *Genome Biol*.
2018; **19**(1): p. 224.
32. Qiu X, Hill A, Packer J, Lin D, Ma YA, Trapnell C. Single-cell mRNA quantification and
differential analysis with Census. *Nat Methods*. 2017; **14**(3): p. 309-315.
33. Zhu H, Wang G, Qian J. Transcription factors as readers and effectors of DNA
methylation. *Nat Rev Genet*. 2016; **17**(9): p. 551-65.
34. Jiang J, Chan YS, Loh YH, et al. A core Klf circuitry regulates self-renewal of embryonic
stem cells. *Nat Cell Biol*. 2008; **10**(3): p. 353-60.
35. Zhao T, Liu C, Chen L. Roles of Klf5 Acetylation in the Self-Renewal and the
Differentiation of Mouse Embryonic Stem Cells. *PLoS One*. 2015; **10**(9): p. e0138168.
36. Lecuyer E, Herblot S, Saint-Denis M, et al. The SCL complex regulates c-kit expression
in hematopoietic cells through functional interaction with Sp1. *Blood*. 2002; **100**(7): p.
2430-40.
37. Gilmour J, O'Connor L, Middleton CP, et al. Robust hematopoietic specification
requires the ubiquitous Sp1 and Sp3 transcription factors. *Epigenetics Chromatin*.
2019; **12**(1): p. 33.
38. Min IM, Pietramaggiori G, Kim FS, Passegue E, Stevenson KE, Wagers AJ. The
transcription factor EGR1 controls both the proliferation and localization of
hematopoietic stem cells. *Cell Stem Cell*. 2008; **2**(4): p. 380-91.
39. Zhu J, Emerson SG. Hematopoietic cytokines, transcription factors and lineage
commitment. *Oncogene*. 2002; **21**(21): p. 3295-313.
40. Venezia TA, Merchant AA, Ramos CA, et al. Molecular signatures of proliferation and
quiescence in hematopoietic stem cells. *PLoS Biol*. 2004; **2**(10): p. e301.
41. Tenen DG, Hromas R, Licht JD, Zhang DE. Transcription factors, normal myeloid
development, and leukemia. *Blood*. 1997; **90**(2): p. 489-519.
42. Antonchuk J, Sauvageau G, Humphries RK. HOXB4-induced expansion of adult
hematopoietic stem cells ex vivo. *Cell*. 2002; **109**(1): p. 39-45.
43. Sun Y, Zhou B, Mao F, et al. HOXA9 Reprograms the Enhancer Landscape to Promote
Leukemogenesis. *Cancer Cell*. 2018; **34**(4): p. 643-658 e5.
44. Magnusson M, Brun AC, Miyake N, et al. HOXA10 is a critical regulator for
hematopoietic stem cells and erythroid/megakaryocyte development. *Blood*. 2007;
109(9): p. 3687-96.

45. Yu VWC, Yusuf RZ, Oki T, et al. Epigenetic Memory Underlies Cell-Autonomous
Heterogeneous Behavior of Hematopoietic Stem Cells. *Cell*. 2017; **168**(5): p. 944-945.
46. Wilson A, Laurenti E, Trumpp A. Balancing dormant and self-renewing hematopoietic
stem cells. *Curr Opin Genet Dev*. 2009; **19**(5): p. 461-8.
47. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic
and adult fibroblast cultures by defined factors. *Cell*. 2006; **126**(4): p. 663-76.
48. Feinberg MW, Wara AK, Cao Z, et al. The Kruppel-like factor KLF4 is a critical regulator
of monocyte differentiation. *EMBO J*. 2007; **26**(18): p. 4138-48.
49. Kotowski M, Safranow K, Kawa MP, et al. Circulating hematopoietic stem cell count is
a valuable predictor of prematurity complications in preterm newborns. *BMC Pediatr*.
2012; **12**: p. 148.
50. Suzuki M, Jing Q, Lia D, Pascual M, McLellan A, Greally JM. Optimized design and data
analysis of tag-based cytosine methylation assays. *Genome Biol*. 2010; **11**(4): p. R36.
51. Ritchie ME, Phipson B, Wu D, et al. limma powers differential expression analyses for
RNA-sequencing and microarray studies. *Nucleic Acids Res*. 2015; **43**(7): p. e47.
52. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological
themes among gene clusters. *OMICS*. 2012; **16**(5): p. 284-7.
53. Heinz S, Benner C, Spann N, et al. Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B
cell identities. *Mol Cell*. 2010; **38**(4): p. 576-89.
54. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol*. 2014; **15**(12): p. 550.
55. Aibar S, Gonzalez-Blas CB, Moerman T, et al. SCENIC: single-cell regulatory network
inference and clustering. *Nat Methods*. 2017; **14**(11): p. 1083-1086.
646

Figure Legend

Figure 1: LGA is associated with DNA hypermethylation targeting key stem cell

signaling pathways. (A) Volcano plot of DNA methylation score differences for LGA

compared to CTRL in cohort 1, cohort 2, and cohort 1 +2. Differentially methylated loci

with *p*-value <0.001 and |methylation difference| > 25 % are shown in red. (B)

Summary of calculation for the gene-methylation score. (C) Validation of the gene-

methylation score. Gene-methylation score distribution. Bar plot of the association

between gene-methylation score and genomic or methylation related features. Bar

plots representing the significance of the difference of gene-methylation score of

DEGs compared to non-DEGs considering different metrics. eQTR, region with

expression quantitative traits loci; DMC, differentially methylated CpGs. (D) Network

representation of GO Biological Process enriched in hypermethylated genes.

Significantly enriched GO terms were identify using GSEA based on the gene-

methylation score. Edges represent interactions (gene overlap) between pathways.

Figure 2: Lineage specific transcriptomic analysis. (A) Hematomap, UMAP

representation of distinct HSPC lineages. (B) Dot plot representing key markers used

to annotate cell populations. LT-HSC, long-term hematopoietic stem cell; HSC,

hematopoietic stem cell; MPP, multipotent progenitor; LMPP, lymphoid-primed

multipotent progenitors; Erythro-Mas, erythroid and mast precursor; Mk/Er,

megakaryocyte and erythrocyte; DC, dendritic cell. (C) MA plots representing gene

expression analysis in HSCs comparing LGA vs CTRL. Differentially expressed genes

with adjusted *p*-value <0.05 and |log$_2$FC| >0.5 are shown in red. (D) Network

representation of significantly enriched pathways identified through GO GSEA

analysis comparing LGA vs CTRL. Non redundant pathways annotations have been

used. Edges represent interactions between pathways.

Figure 3: Association between changes in DNA methylation and in gene

expression. (A) Dot plot representing the correlation between DNA methylation and

gene expression changes. Differentially expressed genes with adjusted *p*-value <0.05

and |log$_2$FC| >0.5 are shown in red. (B) Boxplots representing gene-methylation score

distribution associated to non DEG, up-regulated and down-regulated genes

(Wilcoxon test). (C) Network representation of significantly enriched pathways

identified through GO GSEA analysis based on DEG identified comparing LGA vs

CTRL. Nodes are color coded based on enrichment for differentially methylated genes

using the gene-methylation score. Edges represent interactions between pathways.

Figure 4: Chromatin accessibility analysis. (A) UMAP representing HSPCs lineage based on chromatin accessibility. Annotations is based on the Hematomap using transfer label approach. (B) Dot plot representing enrichment for DMC within lineage specific peaks. (C) Bar plots representing enrichment for peaks containing CpG or peaks containing DMC associated to DEGs, up-regulated and down-regulated genes (* p-value<0.05; *** p-value<0.001). (D) Dot plot representing enrichment for transcription factor motif within Down peaks identified comparing chromatin accessibility between LGA and CTRL. Dots are color coded based on percentage of peaks with motif and y-axis represent the significance of the enrichment. (E) Bar plots representing enrichment analysis considering accessible, down and up peaks. Enrichment is performed using peaks in expressed genes (expressed), peaks with CpGs (CpGs), peaks in DEG (DEGs), peaks with DMC (DMCs) and peaks in DEG with DMC (DMCs_DEGs) as reference gene sets.

Figure 5: Epigenetic programming of HSC specific regulons altered in LGA neonates. Regulons and TF target information were obtained through the SCENIC workflow. (A) Boxplots representing regulon activity score in CTRL and LGA HSC lineage. Barplot representing the change in regulon activity and significance comparing LGA vs CTRL. Only significantly affected regulons are represented (adjusted p-value <0.001 and |activity score fold change| >10%). (B) Heatmap representing association between altered regulons and selected gene sets annotation. (C) Volcano plot representing enrichment in change in expression and DNA methylation in regulons. Regulons enriched considering both expression and methylation (adjusted p-value<0.01 and NES>1.6) are in red. (D) Dot plot representing

enrichment for TF binding motifs using HOMER considering a ±20bp region around

DMCs. Dots are color coded based on significance of the enrichment and y-axis

represent the number of regions with binding motif among DMCs. (E) Dot plot

representing enrichment for TF binding motifs using HOMER considering peaks with

DMCs. Dots are color coded based on the fold-enrichment and y-axis represent the

significance of the enrichment.

Figure 6: Network recapitulating interaction between the epigenomic and

transcriptomic alterations in LGA. (A) Network representing interactions between

target genes and transcription factors considering our top affected regulons ARID5A,

EGR1, KLF2, KLF4, FOSB and JUN. Each dot represents a gene within the network,

triangle represents transcription factor, arrow represents interaction between

transcription factor and target genes, shapes are color coded to reflect the change in

gene-methylation score, and DEGs are labeled in red. Size of the shape represents

the number of interactions. Only genes with 2 or more interactions are represented.

(B) Tracks representing DNA methylation and chromatin accessibility for selected

representative regions. Histogram representing change in DNA methylation at CpG

level comparing LGA vs CTRL. Violin plot representing gene expression for selected

genes (C) Network based on integration of DNA methylation, gene expression and

chromatin accessibility representing transcription factors and downstream target

interactions within EGR1, KLF2 and KLF4 regulons. Only genes associated to peaks

with TF motifs of interest are annotated. Donuts represent different level of

interactions.

Figure 7: LGA is associated with decreased expansion capacity and a HSC shift

toward more differentiated cells. (A) Bar plot representing colonies distribution after

CFU assays. (B) Representative capture of colonies morphological differences found

in CTRL and LGA. (C) Boxplots representing the cell distribution across hematopoietic

main lineages in CTRL and LGA. (D) UMAPs representing pseudotimes across

lineages. (E) Box plots representing the cumulative percentage of cells per

pseudotime in CTRL and LGA. Boxplots in the vignette represent overall pseudotime

distribution in CTRL and LGA. Density plots correspond to cell populations distribution

across pseudotimes. (F) Model recapitulating the influence of LGA on the

hematopoietic compartment. (LT-HSC, long-term hematopoietic stem cell; HSC,

hematopoietic stem cell; MPP, multipotent progenitor; LMPP, lymphoid-primed

multipotent progenitors; Erythro-Mas, erythroid and mast precursor; Mk/Er,

megakaryocyte and erythrocyte; DC, dendritic cell; CFU-GEMM, common myeloid

progenitors; BFU-E, erythroid progenitors; CFU-GM, granulocyte-macrophage

progenitors.

A
DMC analysis (LGA vs CTRL)

Cohort 1

Cohort 2

Cohort 1 + 2

- −log(*p*-value)
- Methylation Change

• *p*-value < 0.001 & |Methylation Change| > 25
• All CpGs

B
Gene-Methylation Score

eQTL
Genomic Annotation — Promoter, Enhancer, Transcribed
TF Binding Site
CpG Regulatory Weight — Gene A, Gene B

$$Gene\text{-}Methylation\ Change\ Score_n = \sum \left(\begin{matrix} CpG_i\ Regulatory \\ Weight \end{matrix} \times \begin{matrix} CpG_i\ Methylation \\ Changes \end{matrix} \right)_{Promoter} + \sum \left(\begin{matrix} CpG_i\ Regulatory \\ Weight \end{matrix} \times \left| \begin{matrix} CpG_i\ Methylation \\ Changes \end{matrix} \right| \right)_{Enhancer}$$

C
Gene-Methylation Score Validation

Gene-Methylation Change Score association with covariates

Comparison DEGs vs no DEGs

D
GSEA analysis (GO Biological Process)

axon development

positive regulation of cell development

cell morphogenesis involved in neuron differentiation

positive regulation of neurogenesis

cell fate specification

embryonic organ development

cell fate commitment

Wnt signaling pathway

Adjusted *p*-value
genes

A Hematopmap

B Top Subpopulation Markers

C LGA vs CTRL in HSCs

DEG (adjusted *p*-value < 0.05 & |log

D GSEA analysis (GO Biological Process)

A

Correlation Methylation and Expression Changes at gene level

Gene Expression Change
$\log_2(\text{Fold Change})*-\log_{10}(\text{adjusted } p\text{-value})$

PLK3
KLF2
EGR1
ZFP36
DUSP2
JUNB
SOCS3
RGCC
SESN2

• DEG (adjusted p-value < 0.05 & |$\log_2$FC| > 0.5)

Gene-methylation Score (score percentiles)

B

p-value < 0.001

Gene-methylation score

non-regulated (n=8923)
upregulated (n=66)
downregulated (n=227)

Genes

C

Up-regulated | Down-regulated | GSEA network based on DEGs

regulation of protein serine/threonine kinase activity

regulation of endothelial cell apoptotic process

regulation of growth

positive regulation of hydrolase activity

signal transduction in response to DNA damage

regulation of DNA−binding transcription factor activity

negative regulation of synaptic transmission

blood coagulation

signal transduction by p53 class mediator

regulation of mitotic cell cycle

vasculature development

negative regulation of cellular component organization

regulation of DNA−templated transcription in response to stress

regulation of cAMP−dependent protein kinase activity

cellular response to chemical stress

response to temperature stimulus

regulation of ATPase activity

response to oxidative stress

Enrichment for differentially methylated genes
$-\log_{10}(\text{adjusted } p\text{-value})$

5   10

Genes overlap

20 40 60

A

ATAC cell subpopulations annotation based on the Hematomap

- Erythro−Mas
- HSC
- MPP/LMPP
- Myeloid
- DC
- Lymphoid
- T cell
- B cell

B

C

DMCs Enrichment Analysis

D

Peaks with decreased accessibility (LGA vs CTRL HSC)

E

Differentially Accessible Peak Enrichment Analysis

A

Regulon Activity Analysis

adjusted *p*-value

B

Regulon Annotation

adjusted *p*-value

C

Correlation Methylation and Expression at regulon level

D

Top 20 TF Motif enriched in DMC regions

E

Top 20 TF Motif enriched in HSC Peaks with DMCs

A

Co-regulatory network from enriched regulons

DEG

Gene with high
gene-methylation
score (> 500)

▲ Transcription Factor

● Target Gene

Number of interactions

● [2 to 3]
● [3 to 4]
● [4 to 126]

B

Multmodal co-regulatory network altered in LGA

Methylation

Gene-methylation
score >500

Expression
- ■ Downregulated FC>0.5
- ■ Downregulated 0<FC<0.5
- ■ Upregulated FC>0
- ■ No change

▲ Transcription Factors

● Target Gene

Genes co-regulated
- — EGR1 / KFL4
- — EGR1 / KFL2
- — KLF2 / KFL4

**TF motif ass
target**

C