General dimensions of human brain morphometry inferred from genome-wide

association data

Anna E. Fürtjes, MSc[1*], Ryan Arathimos, PhD[1,2], Jonathan R. I. Coleman, PhD[1,2], James H.

Cole, PhD[3,4,5], Simon R. Cox, PhD[6,7], Ian J. Deary, PhD[6,7], Javier de la Fuente, PhD[8,9],

James W. Madole, PhD[8], Elliot M. Tucker-Drob, PhD[8,9], Stuart J. Ritchie, PhD[1]

The names of the co-authors are listed in alphabetical order and grouped by affiliation.

[1] Social, Genetic and Developmental Psychiatry (SGDP) Centre, Institute of Psychiatry, Psychology

& Neuroscience, King's College London, SE5 8AF, UK

[2] National Institutes for Health Research Maudsley Biomedical Research Centre, South London and

Maudsley NHS Trust, London, SE5 8AF, UK

[3] Department of Neuroimaging, Institute of Psychiatry, Psychology & Neuroscience, King's College

London, London, SE5 8AF, UK

[4] Centre for Medical Image Computing, Department of Computer Science, University College

London, London, WC1V 6LJ, UK

[5] Dementia Research Centre, Institute of Neurology, University College London, London, WC1N

3BG, UK

[6] Department of Psychology, The University of Edinburgh, Edinburgh, EH8 9JZ, UK

[7] Lothian Birth Cohorts, University of Edinburgh, Edinburgh, EH8 9JZ, UK

[8] Department of Psychology, University of Texas at Austin, Austin, TX 78712-1043, USA

[9] Population Research Center and Center on Aging and Population Sciences, University of Texas at

Austin, Austin, TX 78712-1043, USA

24   **Word count:**

25   4,342 words

27   **Correspondence:**

28   Anna E. Fürtjes

29   anna.furtjes@kcl.ac.uk

Abstract

Background: Understanding the neurodegenerative mechanisms underlying cognitive

declines in the general population may facilitate early detection of adverse health outcomes in

late life. This study investigates biological pathways shared between brain morphometry,

ageing, and cognitive ability.

Methods: We develop Genomic Principal Components Analysis *(genomic PCA)* to

model general dimensions of variance in brain morphometry within brain networks at the

level of their underlying genetic architecture. With genomic PCA we extract genetic principal

components (PCs) that index global dimensions of genetic variance across phenotypes

(unlike ancestral PCs that index genetic similarity between participants). Genomic PCA is

applied to genome-wide association data for 83 brain regions which we calculated in 36,778

participants of the UK Biobank cohort. Using linkage disequilibrium score regression, we

estimate genetic overlap between brain networks and indices of cognitive ability and brain

ageing.

Results: A genomic principal component (PC) representing brain-wide dimensions of

shared genetic architecture accounted for 40% of the genetic variance across 83 individual

brain regions. Genomic PCs corresponding to canonical brain networks accounted for 47-

65% of the genetic variance in the corresponding brain regions. These genomic PCs were

negatively associated with brain age (r_g = -0.34). Loadings of individual brain regions on the

whole-brain genomic PC corresponded to sensitivity of a corresponding region to age (r = -

0.27). We identified positive genetic associations between genomic PCs of brain

morphometry and general cognitive ability (r_g = 0.17-0.21).

Conclusion: These results demonstrate substantial shared genetic etiology between

connectome-wide dimensions of brain morphometry, ageing, and cognitive ability, which

will help guide investigations into risk factors and potential interventions of ageing-related

cognitive decline.

1. Introduction

Progressive ageing-related neurodegenerative processes occurring across micro to

macro scales of the human brain are well-documented within otherwise healthy adults, and

are linked to ageing-related declines in multiple domains of cognitive function [1-3].

Understanding the biological processes underlying these links is paramount for identifying

mechanisms of cognitive ageing that can ultimately be targeted by intervention. The human

brain is a complex network of interconnected regions (the 'connectome') [4, 5], components

of which are interrelated with one another [6, 7], age unevenly over time [8], and may be

differentially relevant to adult cognitive ageing [1-3]. Whereas considerable attention has

been devoted separately to the genetic architecture of human brain morphometry [9-11] and

the genetic architecture of adult cognitive ability [12], relatively less work has been devoted

to scaffolding investigations of the genetic architecture of human brain morphometry onto the

well-established network organization of the brain (although see [13] for a recent exception),

or to investigating how genetic links between components of human brain networks relate to

ageing and cognition. Such investigations have the potential to provide insights into the

etiology of neurocognitive ageing.

Specifically, our genome-wide study builds on a previous study that investigated

dimensions of brain morphometric variation (i.e., principal components underlying brain

volumetric measures) across the human connectome in a large scale cohort ($N = 8,185$) [3].

This phenotypic study implicated well-studied macroscopic brain networks in cognitive

ageing, whereby connectome aging varied alongside those dimensions of morphometric

variation. Brain volumes in the central executive network tended to be most sensitive to age

(i.e., cross-sectionally correlated with age) and, albeit its small size, the central executive was

highlighted to play a disproportionate role in late-life cognitive ability.

Here, we hypothesise that morphometric network organisation, as described by

Madole et al. [3], corresponds to morphometric variation captured by genome-wide data (pre-

registration: https://osf.io/7n4qj). We suggest that a dissimilar organisation of phenotypic and

genetic brain architecture would contradict the neurobiological validity of canonical brain

networks (a similar organisation would be consistent with a measurable genetic foundation of

brain networks). This hypothesis relies on evidence that patterns of brain morphometry and

its organisation are highly heritable [10, 11, 14, 15]. Cheverud originally speculated that "If

genetically and environmentally based phenotypic variations are produced by similar

disruptions of developmental pathways, genetic and environmental correlations should be

similar." [16]. Strong correspondence between phenotypic and genetic correlations was

recently demonstrated for a wide range of morphometric human traits (for example, height

and body mass index) in the UK Biobank cohort [17], and we therefore expect that

phenotypic brain network structures should mirror the structure of genetic correlations within

the same networks.

We consider the same 'canonical' brain networks as Madole et al. [3], using common,

but not indisputable, definitions of the exact regions comprising them [5, 18, 19]. These brain

networks have been characterised embracing a whole-brain perspective, considering existing

literature describing synchronised (i.e., correlated) regional activity in functional MRI data

[3], in addition to converging evidence from other modalities (i.e., structural MRI and lesion-

based mapping [7, 20, 21]). Among the most reported networks are the central executive,

default mode, salience, and multiple demand networks. Specific network characterisations

considered in this study are displayed in Fig. 1 and listed in STable 2.

Brain networks are theorised to integrate information across the brain and,

collectively, to give rise to cognitive functions. The central executive network is thought to

underpin higher-level cognitive functions, including attention and working memory processes

[21, 22]; whereas the default mode network is associated with internally directed and abstract

thought [23]. The salience network is thought to detect salient sensory cues [24], helping to

integrate executive and default functions [22, 25]. Mental processes that organise multiple

cognitive requirements into a series of successive cognitive tasks are thought to be associated

with the multiple demand network [26].

Here, we use genome-wide association data summarising genetic correlates of grey

matter volumes to model morphometric network structures. We focus on brain volumes

because they are highly heritable [14], and are measured independent of mental processes

during MRI scanning (compared with functional MRI). Grey matter volume was

demonstrated to be a strong and robust predictor of general cognitive ability [27, 28], it

reflects atrophy; an important indicator of ageing and health outcomes [29], and, as discussed

above, regional brain volumes have been shown to capture dimensions of morphometric

variation implicated in aging and cognitive ability [3].

In order to model macroscopic brain networks using genome-wide association data, in

this pre-registered study (https://osf.io/7n4qj), we present our novel statistical genetics

method 'genomic PCA' (genomic Principal Component Analysis). With genomic PCA we

extract genetic principal components (PCs) that index global dimensions of genetic variance

across phenotypes (unlike ancestral PCs that index genetic similarity between participants),

the human structural connectome. We mirror previous phenotypic analyses to ensure

comparability with results in Madole et al. [3], that is, we estimate genetic associations

between genetic PCs across brain network structures and both cognitive ability and 'brain

age' [30]. Characterising genetic links between general dimensions of brain organisation,

aging and cognitive ability will help guide investigations into risk factors, biological

mechanisms, and potential interventions of ageing-related cognitive decline.

131   **2. Methods**

The considered UK Biobank sample consisted of 36,778 participants (54% females)

with available neuroimaging data and had an average age of 63.3 years at neuroimaging visit

(range from 40.0 to 81.8 years). Our methodological approach followed four major analysis

steps as displayed in Fig. 2. First, we calculated 83 genome-wide association study (GWAS)

summary statistics for 83 regional volumes that served as input data. Polygenic effects were

fitted in a linear mixed model using REGENIE [31]. Second, we calculated genetic

correlation matrices indicating genetic overlap between regional brain volumes using the

GenomicSEM software [32]. Genetic correlations formed the basis for subsequent analyses;

they are contrasted with phenotypic correlations in Section 3.1. Third, we extracted the first

principal component (PC) from genetic correlation matrices using principal components

analysis (PCA). The genetic PCs were calculated for the whole brain, as well as canonical

brain networks; they represent general dimensions of shared genetic morphometry between

regions included in either the whole brain, or the brain networks (Section 3.2). Based on

Pearson's correlations and Tucker congruence coefficient, we compared genetic correlation

structures with phenotypic correlation structures between the 83 regional volumes (Section

3.3). We also tested whether the relative ordering of phenotypic and genetic PC loadings

correlated with indices of a regions sensitivity to age (i.e. cross-sectional volume-age

correlations; Section 3.4). Finally, we presented a novel method to summarise shared

morphometric variance within brain networks on a genome-wide level in sets of univariate

summary statistics (i.e., genetic PCs underlying multiple brain volumes; see Supplementary

Methods). These univariate summary statistics can be viewed as a summary-based method of

computing GWAS summary statistics that would be obtained from a GWAS on individuals'

scores on the underlying genetic PCs. We used those summary statistics to represent general

dimensions of brain organisation across the whole brain and nine canonical networks at the

level of their underlying genetic architecture. We then quantified genetic correlations

between brain network structures and both general cognitive ability [32] (Section 3.5) and

brain age (Section 3.6). Detailed descriptions of the UK Biobank data used in this study and

the study design can be found in the Supplementary Methods. Our analysis code is displayed

at https://annafurtjes.github.io/Genetic_networks_project/.

162   **3.   Results**

3.1 Genetic correlations between brain-wide volumes recapitulated phenotypic

correlations

On a phenotypic level of analysis, correlations between the 83 brain volumes were

obtained using Pearson's correlations from volumetric phenotypes that were residualised for

age (mean = 63.3, range = 40.0-81.8 years) and sex (54% females). On a genetic level of

analysis, we calculated GWAS summary statistics to get genome-wide associations ($N =$

36,778) for 83 cortical and subcortical grey-matter volumes (Fig. 2.1). SNP-heritability

estimates ranged between 7% ($SE = 0.07$) for the frontal poles and 42% ($SE = 0.04$) for the

brain stem (mean = 0.23, $SD = 0.07$; Fig. 3A).

The GWAS summary statistics enabled the calculation of genetic correlations

between 83 volumes through linkage disequilibrium score regression (LDSC) [33] ($\frac{83(83-1)}{2} =$

3403 between-region correlations; Fig. 2.2). All bilateral regions were almost perfectly

correlated with the corresponding contralateral region. Between-region genetic correlations

ranged from $r_g = $ -0.08 ($SE = 0.09$) between right frontal pole and left pallidum, to $r_g = 0.87$

($SE = 0.08$) between left middle temporal and left inferior temporal (Fig. 3B, SFig. 1).

Corresponding standard errors ranged between 0.01 and 0.03 (mean = 0.014; $SD = 0.002$).

Genetic correlations within canonical networks are provided in SFig.s 2-10.

A positive and large association ($r = .84$; $b = 0.60$; SE = 0.007, $p < 2$ x10^{-16}, $R^2 =$

70%) was obtained between 3403 phenotypic and 3403 genetic correlations (Fig. 3&5A),

indicating that the same regions, that had strongly correlated phenotypic volumes, were also

genetically correlated. Phenotypic correlations were exclusively positive, as were 3,392 of

3,403 genetic correlations; the 11 (0.32%) negative genetic correlations were close to zero

(smallest r_g -0.083; Fig.3).

3.2 PCs of shared genetic variance across the whole-brain and canonical networks

Distributions of phenotypic PC loadings are in Fig. 4A (descriptive statistics for phenotypic shared morphometry in STable 3). On a genetic level of analysis, we extracted PCs from genetic correlation matrices. The first genetic whole-brain PC explained 40% of the genetic variance across 83 regional volumes - slightly larger than the 31% explained by the first phenotypic whole-brain PC. The second genetic whole-brain PC accounted for 6.7% of the total genetic variance; that is, 17% of the variance explained by the 1^{st} genetic PC (SFig. 20). We obtained loadings on this first genetic PC for each regional volume, quantifying how well an individual volume mapped onto the underlying dimension of shared morphometry across the whole brain. Their distribution ranged between 0.30 and 0.81 (mean = 0.62, SD = 0.13, median = 0.65; Fig. 4, STable 3a).

We used the same approach – extracting the first genetic PC and its genetic PC loadings – to examine nine predefined genetic brain-subnetworks (Fig. 1). Regions included in the networks are listed in STable 2. The percentage of genetic variance accounted for by the first network-specific PCs ranged between 65% for the central executive network and 47% for the temporo-amygdala-orbitofrontal network. While the central executive and the hippocampal-diencephalic networks had a narrow, unimodal distribution of PC loadings, the temporo-amygdala-orbitofrontal and cingulo-opercular networks had a wider, and bimodal distribution. That is, volumes included in the central executive network, for example, were more homogeneous and indexed more similar genetic variation, compared with the temporo-amygdala-orbitofrontal network (Fig. 4). Overall, percentages of explained variances were larger for networks including fewer volumes, potentially because larger networks tend to be more heterogeneous.

To test whether data-derived PCs explained more genetic variance than could be expected by chance, we present a version of Parallel Analysis which simulates PCs for

uncorrelated elements with matched genetic sampling variance (see Supplementary

Methods). Parallel Analysis confirmed that genetic PCs of the whole brain and the nine

canonical subnetworks explained substantially more variance than expected by chance (Scree

Plots SFig.s 11-20). Furthermore, we demonstrated that PCs extracted from 800 networks

with randomly included brain volumes explained substantially less averaged variance than

empirical canonical networks. Results from this simulation are presented in STable 5. In

summary, these results illustrate that genetic dimensions of shared morphometry are well

represented by the first underlying PC (i.e., accounts for the majority of genetic variance);

that the dimensions differ between networks, and that they explain similar magnitudes of

variance as their corresponding phenotypes.

3.3 General dimensions of phenotypic and genetic shared morphometry were similarly

organised

To quantify how closely patterns of shared variance between phenotypic and genetic

brain morphometry resemble each other, we calculated a linear regression between sets of 83

phenotypic and 83 genetic PC loadings. PC loadings indicate relative magnitudes of brain

regions' loadings on either phenotypic or genetic dimensions of shared morphometry, and

serve as an index of how well a volume represents trends across the brain (or the network).

The association between phenotypic PC loadings and genetic PC loadings was large and

significant ($b = 0.65$, SE $= 0.06$, $p = 5.07 \times 10^{-17}$, $R^2 = 58\%$), indicating that an increase in one

unit in the genetic PC loadings is associated with an increase of .65 units in the phenotypic

PC loadings (intercept $= 0.15$). This approach considers ordering relative to the mean.

The Tucker congruence coefficient was used to index the degree of similarity of

genetic and phenotypic PC loadings, taking into account both their relative ordering and their

absolute magnitudes [34]. The Tucker coefficient revealed very high congruence in the

deviation from zero between phenotypic and genetic PC loadings for the 83 volumes (Tucker

coefficient = 0.99). These results illustrate a close correspondence and an equivalent

organisation of phenotypic and genetic dimensions of shared morphometry; a finding that

aligns with Cheverud's Conjecture (Section 4.2).

3.4 Genetic dimensions of shared morphometry were associated with age sensitivity

240   Previous work demonstrated an association between phenotypic dimensions of shared

morphometry across the whole brain, represented by phenotypic PC loadings, and indices of

age sensitivity [3]. Age sensitivity is approximated by a correlation of a regional brain

volume with age across the sample, which is typically negative in adult populations. Here, we

replicated this association between phenotypic shared morphometry (i.e., phenotypic PC

loadings) and age sensitivity ($r = -0.43$, $p = 4.4 \times 10^{-5}$; Fig. 5c), and we found a significant,

but smaller association for genetic PC loadings ($r = -0.27$, $p = 0.012$; Fig. 5d). This

demonstrates that the more the genetic variation of a brain volume resembles general

morphometric trends across the brain (larger genetic PC loading), the stronger this volume is

negatively correlated with age. Note that these results emerged even though PC loadings were

extracted from brain volumes *residualised* for age and were nevertheless associated with age

sensitivity. In summary, these results show that phenotypic PC loadings and genetic PC

loadings both display associations with age sensitivity, as indexed by cross-sectional age-

volume correlations (Section 4.3).

3.5 General dimensions of shared morphometry were genetically correlated with

general cognitive ability

To quantify genetic correlations between general dimensions of network morphometry and

general cognitive ability, we indexed shared genetic variance across brain networks, by

extracting underlying genome-wide PCs. Genome-wide PCs were calculated by summarising

per-SNP effects from multiple brain volume GWAS summary statistics, weighted by volume-

and network-specific PC loadings (novel method presented in Fig. 2.4). Using GenomicSEM

software [32], we calculated genetic correlations between brain networks and seven cognitive

traits [32] (SFig. 21). The cognitive traits mostly had high loadings on a genetic general

cognitive ability factor (median = 0.81, range = 0.30-0.95); the Reaction Time task had the

lowest loading on the factor (SFig. 22). Strong genetic overlap between brain networks

indicated that they indexed very similar polygenic signal (r_g between networks = 0.63-0.97).

All networks were significantly genetically associated with the general cognitive ability

factor; correlation magnitudes across all networks ranged between r_g = 0.17-0.21 (Table 1).

According to commonly-used rules of thumb from Hu and Bentler [35](CFI > 0.95, RMSEA

< 0.08), all models showed good model fit (STable 4).

Based on phenotypic findings that have highlighted the importance of the central

executive network to general cognitive function [3], we hypothesised to find a stronger

genetic association between general cognitive ability and the central executive network

relative to other subnetworks (see pre-registered plan https://osf.io/7n4qj). There was no

evidence for significant differences in total correlation magnitudes between the central

executive network and general cognitive ability compared with other brain networks, even

after accounting for network sizes (see Methods; Fig. 23, STable 6). We found no significant

difference in model fit using GenomicSEM [32] comparing one model accounting for

network size, and another model not accounting for it ($\Delta \chi^2$ p-values ranged between .072 and

1.00; STable 5).

We also investigated whether genetic associations were driven by specific cognitive

traits. We obtained non-significant Q_{trait} heterogeneity indices [36] for all brain networks,

demonstrating that the general cognitive ability factor accounted well for the patterns of

association between specific cognitive abilities and the brain networks (SFig. 24). The fact

that the general cognitive ability factor accounted well for specific abilities, and that the

specific abilities were mostly significantly associated with the networks, confirms that the

genetic associations between specific cognitive abilities and brain networks are likely general

and act through a factor of general cognitive ability (Section 4.4).

3.6 General dimensions of shared morphometry were genetically correlated with brain

age

Finally, we calculated a genetic correlation between shared morphometry across the

whole brain and *brain age*. Brain age is based on individual-level predictions of how much

older (or younger) an individual's brain appears from structural MRI measures, relative to

their chronological age [37] (see Supplementary Methods). We found a moderate negative

genetic association (r_g = -0.34; SE = 0.06) between general dimensions of shared

morphometry across the whole-brain and brain age, suggesting that consistently larger

volumes across the whole brain indicate younger brain age (Section 4.3).

299   **4. Discussion**

This genetically-informed study provides fundamental insights into the complex

biology shared between brain organisation, ageing, and cognitive ability. Using genomic

PCA, we demonstrated that general morphometric dimensions underlying brain network

structures genetically overlapped with general cognitive ability, brain age, and sensitivity of a

corresponding region to age, albeit being distinctly measured demographic, psychological

and neuroimaging concepts. Our findings highlight measurable biological pathways giving

rise to genetic variation in brain morphometry which may influence pathways underlying

cognitive ability and vulnerability towards ageing. Discovery of shared genetic etiology and

its associated neurodegenerative mechanisms should inform efforts of detecting and

mitigating cognitive decline in ageing societies [3, 38, 39].

4.1 Characteristics of genetic brain network organisation

We demonstrated that genetic dimensions of shared morphometry underlying brain

networks (i.e., first genetic PC) accounted for substantial systematic variance shared between

brain volumes (e.g., 40% of genetic variance across the whole brain). These major genetic

dimensions even explained more variance than their phenotypic analogue (31% of phenotypic

variance across the whole brain). All genetic networks explained substantially more variance

than was expected by chance. These findings provide a new line of evidence characterising

and underpinning the existence of a genetic foundation for canonical brain networks that have

featured prominently in neuroscientific studies [e.g., 7].

4.2 Analogous organisation of phenotypic and genetic dimensions of shared

morphometry

We discovered a high degree of similarity between phenotypic and genetic features of

brain network organisation (e.g., $r_{\text{genetic vs. phenotypic correlations}} = 0.84$; Tucker congruence = 0.99;

$r_{phenotypic\ vs.\ genetic\ PC\ loadings} = 0.76$). According to Cheverud's Conjecture [16], this indicates that

brain organisation, indexed through both phenotypic and genetic variance, seems to be

underpinned by similar, overlapping developmental pathways.

4.3 Genetic PCs as indices of brain regions' age sensitivity and brain age

As previously demonstrated, phenotypic PC loadings onto an underlying brain-wide

dimension of shared morphometry resembled patterning of sensitivity of a corresponding

region towards age (i.e., cross-sectional age-volume correlations) [3]. Here, we replicated this

negative association, and showed that it also exists, albeit to a lesser degree, between age-

volume correlations and genetic instead of phenotypic PC loadings. This suggests that

dimensions along which brain regions share morphometric variance (i.e., generally larger

volumes across an individuals' brain) are structured similarly to patterns by which brain

regions display increased vulnerability to ageing. This finding needs to be triangulated by

either future longitudinal studies, or cross-sectional studies modelling within-person atrophy

by incorporating information on prior brain size (e.g., intracranial volume).

One potential explanation for this association is that brain regions that are genetically

predisposed to be large volumes, that share higher levels of morphometric variance with the

rest of the brain, and that are more central to heavily-demanding cognitive processes, might

come under more strenuous developmental and environmental pressure, perhaps through

increased metabolic burden, compared with other, less central regions. Thus, the embedding

of a brain volume within the whole brain's organisation, and the genetic foundation of its

positioning in the brain, could govern the functional stresses and other influences to which

certain areas are exposed. This might alter disproportionately the speed at which some

regions atrophy with advancing age.

That dimensions of shared morphometry resemble patterns of age sensitivity is of

interest because it emerged from shared variance among brain phenotypes that had been

residualised for age. Consequently, we suggest that patterns of brain structural ageing, a

construct labelled *brain age* [29], might not capture how quickly an individual's regional

volumes decline compared to their peers, but rather, general healthy morphometry across the

brain. Previous research showed that a younger-appearing brain, relative to the individual's

chronological age, predicted better physical fitness, better fluid intelligence, and longevity

[29]. Healthy brain morphometry could vary between people for many non-age-related

reasons, including genetic predisposition. Individuals that are genetically predisposed

towards consistently larger brain volumes might have generally healthier, better-integrated

brains, which could be more resilient towards harmful environmental factors.

In line with this theory, we found that younger brain age was genetically associated

with a major dimension of brain-wide shared morphometry as indexed by a genetic PC (r_g = -

0.34; SE = 0.06). Thus, consistently larger volumes across the brain indicate a younger

structural brain organisation, and this is the first study to quantify the degree to which these

two concepts overlap . It motivates further investigation into the possibility that they are

underpinned by the same general shared biological pathways.

4.4 Genetic PCs as indices of cognitive performance

This study demonstrated that cognitive ability is positively associated with genetic

morphometric variance shared across the whole brain, and across smaller canonical networks.

This was investigated by modelling a genetic factor of general cognitive ability using

GenomicSEM [32]. We calculated the genetic correlation between general cognitive ability

and genetic PCs across the whole brain, and nine canonical subnetworks. The whole brain

and all nine networks were significantly genetically correlated with general cognitive ability

at magnitudes between 0.17 and 0.21. This was the same level of genetic association with

general cognitive ability that was previously found for broad measures of total brain volume

[40]. There was no evidence to suggest that those magnitudes statistically differed between

the networks; probably because the polygenic signal indexed by the genetic PCs were highly

similar between brain networks (mean r_g between networks 0.83, SD = 0.09).

This indicates that the genetic association between brain morphometry and cognitive

ability was not driven by specific network configurations. Instead, genetic PCs indexed

genetic variance relevant to larger brain volumes and a brain organisation that is

advantageous for better cognitive performance. This was regardless of how many brain

regions and from which regions the measure of shared genetic morphometry was extracted.

This lack of differentiation between networks, in how strongly they correlate with cognitive

ability, is in line with the suggestion that the total number of neurons in the mammalian

cortex, which should at least partly correspond to its volume, is a major predictor of higher

cognitive ability [41]. These findings suggest that highly shared brain morphometry between

regions, and its genetic analogue, predict a generally bigger, and cognitively better-

functioning brain.

Unexpectedly, genetic correlations between networks and cognitive ability did not

suggest any prominent role of the central executive network (a previous *phenotypic* study [3]

demonstrated that the central executive network was disproportionately predictive of

cognitive abilities relative to its few included volumes). On a genetic level of analysis, we

also expected a stronger correlation with cognitive ability for the central executive network

compared with the other networks. The lack of differentiation between networks, taken

together with previous phenotypic evidence for a disproportionately large association

between the cognitive ability and the central executive, suggests nongenetic mechanisms to

play important roles, perhaps developmental and environmental influences, through which

the central executive network matures, and specialises for cognitive performance.

4.5 Limitations

Analyses in this study come with limitations. Genetic correlations are representative for genetic associations across the entire genome, but do not give direct insight into specific genomic regions of sharing. As genetic correlations were calculated using LDSC, the limitations that apply to LDSC methodology are relevant to our study (discussion in Supplementary Note). We conclude based on heritability estimates, indexing signal-to-noise ratios in GWAS, that there was sufficient polygenic signal to warrant LDSC analysis (heritability ranged 7-42%). LDSC intercepts were perfectly associated with phenotypic correlations ($R^2 = 0.99$), indicating that the analyses successfully separated confounding signal (including environmental factors) from the estimates of genetic correlations.

This study was conducted in the UK Biobank sample, which is not fully representative of the general population: its participants are more wealthy, healthy and educated than average [42]. Cohort effects may affect the degree to which differential cortical regional susceptibility to ageing can be inferred from cross-sectional data. It remains to be tested whether our results can be extrapolated to socio-economically poorer subpopulations, or outside European ancestry. Results were also dependent on the choice of brain parcellation to divide the cortex into separate regions.

4.6 Conclusion

This genetically-informed study delivered evidence for shared etiology between factors that may contribute to neurodegenerative mechanisms underlying ageing-related cognitive decline. Using genome-wide data, we quantified a substantial overlap of genetic variation between distinct measures of ageing, cognitive ability, and brain morphometry, all of which are variables of interest due to their potential social and economic consequences for ageing societies. These fundamental insights will help guide investigations into risk factors, biological mechanisms, and potential interventions of ageing-related cognitive decline.

More specifically, we demonstrated that younger brain age genetically captured

interindividual variation substantially related to brain network structures (i.e., consistently

enlarged volumes). Because the network structures were modelled based on variance

independent of age, this suggests that younger brain age could primarily be an index of brain

health. Contrary to previous phenotypic findings, our genetic analyses did not provide

evidence for a disproportionate role of the central executive network in cognitive

performance. This motivates future investigations into environmental influences on the

specialisation of brain networks. Altogether, our new genomic PCA methodology and the

resulting insights of this study provide a basis for future investigations that aim to interrogate

the genetic and environmental bases of ageing and cognitive decline.

Supplementary Methods

Study Design

UK Biobank data

Magnetic resonance imaging (MRI) data was collected by the UK Biobank study with identical hardware and software in Manchester, Newcastle, and Reading. Brain volumetric phenotypes were pre-processed by an imaging-pipeline developed and executed on behalf of UK Biobank [43]. More information on T1 processing can be found in the UK Biobank online documentation [44]. Briefly, cortical surfaces were modelled using FreeSurfer, and volumes were extracted based on Desikan-Killiany surface templates [45]; subcortical areas were derived using FreeSurfer aeseg tools [46]. Volumetric measures (mm^3) have been generated in each participant's native space. We used 83 available imaging-derived phenotypes (IDPs) of cortical and subcortical grey-matter volumes in regions of interest spanning the whole brain (UK Biobank category 192 & 190; STable 1). We assume the IDPs to be normally-distributed.

Phenotypic quality control

Excluding participants who withdrew consent, we considered 41,776 participants with non-missing T1-weighted IDPs that had been processed in conjunction with T2-weighted FLAIR (UK Biobank field ID 26500) where available. Using both T1 and T2 measures ensures more precise cortical segmentation [47]. Extreme outliers outside of 4 standard deviations from the mean were excluded, which resulted in between 41,686 to 41,769 available participants depending on the IDP. 381 participants were excluded as they self-reported non-European ethnicity. Across the 83 brain volumes variables and the covariates, this phenotypic quality control resulted in 39,947 complete cases, for whom the following genetic quality control steps were performed.

Genetic quality control

Out of the 39,947 UK Biobank participants, genetic data were available for 38,957

participants. Genetic data was quality controlled on by UK Biobank and were downloaded

from the full release [48]. We applied additional quality control as previously described in

Coleman et al. [49] using PLINK2 [50]. 38,038 participants were of European ancestry

according to 4-means clustering on the first two genetic principal components available

through UK Biobank [51]. Of those participants, we removed 72 due to quality assurance

provided by UK Biobank and 204 participants due to high rates of missingness (2%

missingness). To obtain a sample of unrelated individuals, 956 participants were removed

using the greedyRelated algorithm (KING $r < 0.044$ [52]). The algorithm is "greedy" because

it maximises sample size; for example, it removes the child in a parent-child-trio. Finally, 28

participants were removed because genetic sex did not align with self-reported sex, resulting

in a total of 36,778 participants (STable 10). Genetic sex was identified based on measures of

X-chromosome homozygosity (F_X ; removal of participants with $F_X < 0.9$ for phenotypic

males, $F_X > 0.5$ for phenotypic females). The final sample ($N = 36,778$) included 19,888

females (54 %) and had an average age of 63.3 years at the neuroimaging visit (range from

40.0 to 81.8 years).

Out of 805,426 available directly genotyped variants, 104,771 were removed for high

rates of missing genotype data (> 98%). 103,137 variants were removed due to a minimum

allele frequency of 0.01, and 9,935 variants were removed as they failed the Hardy-Weinberg

exact test (p-value $= 10^{-8}$). After excluding 16,326 variants on the sex chromosomes and

those with chromosome labels larger than 22, we obtained a final sample of 571,257 directly

genotyped SNPs. Imputed genotype data was obtained by UK Biobank with reference to the

Haplotype Reference Consortium [53], and we filtered them for a minor allele frequency of

above 0.01 and an IMPUTE INFO metric of above 0.4.

Measures of cognitive performance

UK Biobank collected cognitive performance data using assessment on a touchscreen

computer. The following seven tests were implemented: *Matrix Pattern Completion task* for

nonverbal reasoning, *Memory – Pairs Matching Test* for memory, *Reaction Time* for

perceptual motor speed, *Symbol Digit Substitution Task* for information processing speed,

Trail Making Test – B and *Tower Rearranging Task* for executive functioning, and *Verbal*

Numerical Reasoning Test for verbal and numeric problem solving, or fluid intelligence.

Despite the non-standard and unsupervised delivery of assessment, these cognitive tests

demonstrate strong concurrent validity compared with standard reference tests ($r = .83$) and

good test-retest reliability (Pearson r range for different cognitive tests = 0.4–0.78) [54].

In this study, we considered GWAS summary statistics of performance in these seven

cognitive tests by de la Fuente, Davies [12] that were calculated with between 11,263 and

331,679 participants for each test. We consider the HapMap 3 reference SNPs with the MHC

regions removed.

Statistical analysis

GWAS summary statistics calculation. GWAS summary statistics for the 83

regional brain volumes (continuous variables) were calculated using REGENIE [31], which

fits polygenic effects in a linear mixed model using Ridge regression. The REGENIE pipeline

is split into two steps: First, blocks of directly genotyped SNPs are used to fit a cross-

validated whole-genome regression model using Ridge regression, to determine the amount

of phenotypic variance explained by genetic effects. Second, the association between the

phenotype and imputed genetic variants is calculated conditional upon Ridge regression

predictions from the first step. Proximal contamination is circumvented by using a leave-one-

chromosome-out scheme.

507   Covariates included in the GWAS analyses were *age at neuroimaging visit*, *sex*,

genotyping batch, and *40 genetic principal components* as provided by UK Biobank. We also

derived the variables *time of year*, *head position*, and *acquisition site,* but excluded them

from our set of GWAS covariates because they were not associated with the brain volumes at

the pre-registered arbitrary cut-off of $r \leq .10$ (STable 9), and therefore explained less than 1%

of the phenotype variance. Note that, in contrast to other existing brain-volume GWAS in UK

Biobank [e.g., 55], our analyses were conducted *without* controlling for brain size (or any

other global brain measure such as total grey-matter volume or intracranial volume). Genetic

correlations calculated relative to such global measures are known to attenuate genetic

correlations among volumes, as well as with other traits such as cognitive abilities [15]. In the

context of this study, we aim to model general dimensions of variance shared between brain

volumes which will closely covary with brain size. Attenuated genetic correlations would

hide major dimensions of variance across genetic brain networks, because much of the

variance shared between volumes overlaps with variance indexed by brain size and would

therefore not tag general dimensions of shared genetic variance between brain volumes. This

variance is of interest because general intelligence yields global rather than a region-specific

associations with grey matter volume [28]. Equally, aging affects the whole brain rather than

individual regions [56].

525   **Genetic and phenotypic correlation matrices between brain volumes.** To derive

dimensions of shared morphometry across brain volumes, we calculated both a phenotypic

and a genetic correlation matrix from 83 grey-matter volume variables. Phenotypic regional

brain volumes were residualised for age at neuroimaging visit and sex, and then used to

estimate a phenotypic correlation matrix through Pearson's correlations with complete

pairwise observations. The genetic correlation matrix was inferred through LDSC, a

technique quantifying shared polygenic effects between traits using GWAS summary

statistics. Cross-trait LDSC regresses the product of effect sizes in two GWAS onto linkage

disequilibrium scores, indicating how correlated a genetic variant is with its neighbouring

variants [33]. The slope indexes the genetic correlation, while the intercept captures signal

uncorrelated with LD, such as population stratification, environmental confounding, and

sample overlap.

To quantify the relationship between phenotypic and genetic correlations, we

estimated the correlation between 3403 phenotypic and genetic between-region correlations

($\frac{83(83-1)}{2} = 3403$ correlations between 83 volumes). Additionally, we calculated genetic

correlation matrices for smaller canonical networks including fewer brain volumes than the

whole brain. For example, the central executive network included eight regional volumes

(STable 2 lists volumes included in the nine canonical networks). We reported SNP-

heritability estimates for each brain volume inferred through LDSC.

Principal component analysis (PCA) **of genetic and phenotypic correlation**

matrices. PCA was applied to the phenotypic and genetic correlation matrices indicating

genetic overlap between brain volumes described above to obtain their respective first

principal component (PC). The first PC represents an underlying dimension of common

structural sharing across regional volumes, which we refer to as general dimensions of shared

morphometry throughout this manuscript. PC loadings were calculated for all volumes in the

whole brain, as well as volumes in smaller canonical networks to quantify contributions of

regional volumes to this either brain-wide, or network-specific dimension of shared

morphometry.

Parallel analysis. We tested whether genetic PCs explained more variance than

expected by chance, that is, whether they explained more than 95% of their corresponding

PCs generated under a simulated null correlation matrix. We developed a version of parallel

analysis to generate null distributions of eigenvalues by simulating null correlation matrices

sampled from a diagonal population correlation matrix, where the multivariate sampling

distribution is specified to take the form of the sampling distribution of the standardised

empirical genetic correlation matrix (the V_{STD} matrix, as estimated using GenomicSEM [32]).

This sampling correlation matrix serves as an index of the precision of the elements in the

empirical genetic covariance matrix (i.e., heritabilities and co-heritabilities across traits) and

the sampling dependencies among these when generating the random null models. We

specified 1,000 replications to simulate the null correlation matrices and use a 95% threshold

for distinguishing true eigenvalues from noise.

Simulation of networks with randomly included brain volumes. We performed an

additional sensitivity analysis simulating networks with randomly included brain volumes, to

determine whether shared structural variance relied on network membership, or arose through

phenotypic properties common to all regional brain volumes. To compare explained

variances between canonical networks and random networks, we quantified the expected

explained variance in random networks by randomly sampling regions 800 times each, for

different numbers of included volumes (because networks including fewer volumes generally

tend to explain a larger percentage of variance, as larger networks are more heterogeneous).

That is, simulations were run for 8, 10, 12, 16, 30, and 36 included regions, to obtain a

distribution for each networks size to compare the corresponding network's explained

variance to. We reported the mean explained variance by PCs for networks with randomly

included volumes and a 95% confidence interval. Comparisons between explained variances

for random and empirical networks were done for the same number of included volumes.

Correlation between phenotypic and genetic PC loadings. To compare whether

genetic correlations structures of regional brain morphometry resembled the phenotypic

correlation structure of the same regions, we calculated an un-standardised linear regression

with a vector of 83 phenotypic whole-brain PC loadings as the dependent variable, and a

vector containing 83 genetic whole-brain PC loadings as the independent variable. We

calculated the Tucker congruence coefficient to quantify the relative similarity between the

two sets of PC loadings independent of their absolute magnitude. The coefficient is

insensitive to scalar multiplication [57].

Correlation between genetic PC loadings with age sensitivity. Pearson's

correlations between 83 phenotypic grey-matter volumes and age at neuroimaging visit were

calculated to quantify cross-sectional age-volume-correlations for each of the 83 brain

volumes. These age-volume correlations are referred to as *age sensitivity* throughout the rest

of the manuscript. We estimated the correlation between a vector containing indices of age

sensitivity and (1) a vector of *genetic* whole-brain PC loadings, and for comparison (2) a

vector of *phenotypic* whole-brain PC loadings.

Genome-wide shared genetic variance of morphometry across the whole brain

and canonical networks. To statistically represent genome-wide shared morphometric

variance across brain volumes (i.e., genetic PCs), we developed a novel method summarising

genome-wide by-variant effects contained in the grey-matter volume GWAS summary

statistics, which were weighted by their respective (region-specific) PC loadings obtained

through PCA. We derived GWAS summary statistics for a genetic principal component of

multiple GWAS phenotypes derived from samples of unknown degrees of overlap by

adapting existing software for genome-wide multivariate meta-analysis by Baselmans et al.

[58] and using GenomicSEM [32]. Fig. 2 illustrates this approach in a four-step procedure.

The input data for our approach are GWAS summary statistics for 83 cortical and subcortical

brain volumes (step 1). We have made them publicly available online. Using the

GenomicSEM software [32], we obtained a genetic correlation matrix indicating genetic

overlap between these 83 brain volumes (step 2). We extracted PC loadings on the underlying

general dimension of shared genetic variance for each of the 83 regions (step 3). Finally, we

modified the existing genome-wide multivariate meta-analysis software package [58], in

order to create summary statistics for an underlying genetic PC. Genome-wide SNP effects

were calculated as an average of all SNP effects contributed by the 83 GWAS phenotypes,

weighted by their respective PC loading, with standard errors computed using a method that

corrects for sample overlap, as estimated by LDSC (step 4). We used this approach to

calculate univariate summary statistics to represent general dimensions of shared

morphometry between regional volumes across the whole brain (83 GWAS phenotypes), as

well as nine smaller canonical networks.

We had tested and validated this novel approach in an independent set of GWAS

summary statistics of four risky behaviours [59]. In addition to the risky behaviour GWAS,

another set of summary statistics is available for a phenotypic PC underlying these risky

behaviour phenotypes that the authors had calculated phenotypically before running GWAS

analyses. We compared these phenotypic PC GWAS summary statistics by Linnér, Biroli

[59] with summary statistics for a *genetic* PC underlying the four risky behaviours GWAS

that we calculated using our novel method outlined above (Fig. 2). We found that they

correlated at a magnitude of $r_g = 0.99$ (*SE* = 0.037) confirming that our method captures the

same signal as can be obtained from phenotypic PCs, by simply relying on publicly available

GWAS data. For details of the analysis and code refer to:

https://annafurtjes.github.io/genomicPCA/ .

Genetic correlation between general dimensions of shared morphometry across

the whole-brain and brain age. Using LDSC [33], we calculated a genetic correlation

between genetic morphometric sharing across the whole brain and *brain age*. The summary

statistics indexing dimensions of shared morphometry across brain volumes were created

using the novel method presented above (Fig. 2). We downloaded the brain age GWAS

summary statistics online [37]. Brain age is a phenotype based on individual-level predictions

of how much older (or younger) an individual's brain appears, relative to their chronological

age. It is estimated using parameters characterising the relationship between age and

structural neuroimaging measures (volume, thickness, and surface area) that were tuned using

machine learning in an independent sample. The final brain age phenotype indexed in the

GWAS was calculated as the difference between participants chronological age and their age

as predicted based on structural brain characteristics.

Genetic correlations between brain networks and a factor of general cognitive

ability. We assessed genetic correlations between brain networks and general cognitive

ability using GenomicSEM [32]. Using univariate network-specific summary statistics (as

describe above; Fig. 2) and a genetic general cognitive ability factor modelled from seven

cognitive ability GWAS summary statistics , the GenomicSEM software [32] was used to

model general cognitive ability and perform multivariate LDSC using diagonally weighted

least squares. To quantify model fit, we reported default fit indices calculated by the

GenomicSEM package: χ_2 values, the Akaike Information Criterion (AIC), the Comparative

Fit Index (CFI) and the Standardised Root Mean Square Residuals (SRMR). The multiple

testing burden was addressed by correcting *p*-values from the genetic correlations for

multiple testing with a false-positive discovery rate of 5% [60].

We preregistered that we would test for significant differences in correlation

magnitudes between the networks that yielded a significant association with general cognitive

abilities. Because we hypothesised a particularly strong association for the central executive

network, we planned to perform this comparison between the central executive and all other

networks, to reduce the multiple testing burden. We fitted two GenomicSEM models in

which correlation magnitudes between general cognitive ability and both the central

executive and another network were either freely estimated, or they were forced to be the

same. A significant decrease in model fit between the freely estimated model and the

constrained model ($df = 1$) would indicate that there likely are differences in correlation

magnitudes between the networks in how strongly they correlate with general cognitive

ability (SFig. 23).

660   Additionally, we assessed whether the central executive network was

disproportionately genetically correlated with general cognitive ability considering its small

size (i.e., few included volumes). Similar to the approach described above, we fitted two

models: One, in which we freely estimate the correlation between the central executive and

general cognitive ability, and the correlation between another network and general cognitive

ability. We then divided the correlation magnitude by the number of regions included in the

network (i.e., magnitude was divided by 8 for the central executive network, it was divided

by 16 for the default mode, by 36 for the P-FIT etc.). The second model had the same set up,

but we forced the adjusted correlations for the two networks to be equal (e.g., $r_{central\ executive}$ / 8

$== r_{default}$ / 16). We assessed whether there was a significant difference in χ^2 model fit

between these two models. As above, a significant decrease in model fit between the freely

estimated model and the constrained model ($df = 1$) would indicate that there likely are

differences in relative correlation magnitudes (i.e., magnitudes adjusted for network sizes).

Based on previous findings, we expected the relative magnitude for the central executive

network to be significantly larger than the relative magnitude for any other network.

675   To probe whether any specific cognitive ability might have driven the genetic

associations between brain networks and general cognitive ability, we reported genetic

correlations between the significant networks and three specific cognitive abilities: (1) *Matrix*

Pattern Completion task to represent nonverbal reasoning, (2) *Memory – Pairs Matching Test*

to represent memory, and (3) *Symbol Digit Substitution Task* to represent information

processing speed. Reducing the analyses to only three consistent and representative cognitive

measures reduced the burden of multiple testing.

We calculated Q_{trait} heterogeneity indices [36] to evaluate whether the general

cognitive ability factor that we fit in the models above accounts well for the specific

cognitive abilities. To this end, we compared the fit of two models for each network as

displayed in SFig. 24. One model allows for independent associations between the seven

cognitive traits, and both general cognitive ability and the brain network. The second model

forces the association between the seven cognitive traits and the brain network to go through

the general cognitive ability factor. We obtained χ^2 fit statistics for both models and tested

their difference for statistical significance ($\Delta \chi^2 \neq 0$; $df = 6$). Non-significant results ($p >$

0.05/10) would suggest that genetic associations between cognitive abilities and brain

networks are likely general and act through a factor of general cognitive ability.

Data and code availability. Access to phenotypic and genetic UK Biobank data was

granted through the approved application 18177. We have made the 83 GWAS summary

statistics of regional volumes available at the GWAS catalogue

(https://www.ebi.ac.uk/gwas/). GWAS summary statistics for the seven cognitive traits by de

la Fuente, Davies [12] were downloaded at https://datashare.ed.ac.uk/handle/10283/3756.

The pre-registration for this analysis can be found online (https://osf.io/7n4qj). Full analysis

code including results for this study are available at

https://annafurtjes.github.io/Genetic_networks_project/index.html.

Acknowledgements

AEF is funded by the Social, Genetic and Developmental Psychiatry Centre, King's College London and the National Institute of Health (NIH) grant R01AG054628. SJR is funded by the Jacobs Foundation. JHC is funded by a UK Research & Innovation(UKRI) Innovation Fellowship (MR/R024790/1; MR/R024790/2). JF is funded by the National Institutes of Health (NIH) grant R01AG054628. JF and EMTD are members of the Population Research Center (PRC) and Center on Aging and Population Sciences (CAPS) at The University of Texas at Austin, which are supported by NIH grants P2CHD042849 and P30AG066614. JD, JWM, and EMTD were supported by NIH R01AG054628. IJD is with the Lothian Birth Cohorts group, which is funded by Age UK (Disconnected Mind grant), the Medical Research Council (grant no. MR/R024065/1) and the University of Edinburgh's School of Philosophy, Psychology and Language Sciences. The contribution by RA represents independent research part-funded by the National Institute for Health Research (NIHR) Maudsley Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. The contribution by JRIC represents independent research part-funded by the National Institute for Health Research (NIHR) Maudsley Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. SRC is supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (Grant Number 221890/Z/20/Z). This research was funded in part by the Wellcome Trust [221890/Z/20/Z]. For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.
The authors gratefully acknowledge the UK Biobank resource (https://www.ukbiobank.ac.uk/) and its research team, who have made this work possible (project number 18177). The authors acknowledge use of the research computing facility at King's College London, Rosalind (https://rosalind.kcl.ac.uk), which is delivered in partnership with the National Institute for Health Research (NIHR) Biomedical Research Centres at South London & Maudsley and Guy's & St. Thomas' NHS Foundation Trusts, and part-funded by capital equipment grants from the Maudsley Charity (award 980) and Guy's & St. Thomas' Charity (TR130505).

Author contributions

Conceptualisation and methodology: SJR, EMTD, JHC, AEF, SRC

Supervision: SJR, EMTD, JHC

Network characterisation: SRC

Idea to investigate genetic brain age – shared morphometry correlation: JWM

Script used to perform genetic parallel analysis: JF

Data access: CML

Genetic quality control: AEF, JRIC

GWAS calculation: AEF, RA

Data analysis: AEF

Writing: AEF

Visualisations: AEF

Reviewed draft: all authors

Disclosures

Ian Deary is a participant in UK Biobank. All other authors have no conflicts of interest to

declare.

Figures

Fig. 1. Nine canonical brain subnetworks. The networks were visualized with the BrainNet

Viewer (http://www.nitrc.org/projects/bnv/) [61]. Regions of interest were visualised using

scripts by Dr. Colin Buchanan (University of Edinburgh). Included brain regions and their

abbreviations are listed in STable 2.

1. Input data:

83 GWAS summary statistics for cortical and
subcortical brain volumes

2. Genetic correlation matrix:

inferred through linkage disequilibrium score regression for 83
brain volumes in GenomicSEM [32]

3. Principal Component Analysis:

extract principal component (PC) loadings on the first
PC of genetic correlation matrix with eigen() R function

First Principal Component

0.74   0.88   0.54   0.64

Volume *1*   Volume *2*   Volume *3*   ...   Volume *83*

4. Summary statistics representing underlying genetic PC:

Genome-wide SNP effects calculated as an average of all SNP effects
contributed by multiple volumes, weighted by respective PC
loadings. Computed using modified software package by Baselmans
et al. [58]

GWAMA software package by Baselmans[1], our modified GWAMA
function[2], and a step-by-step demonstration of Genomic PCA[3] available
on GitHub

Fig. 2. Four-step procedure to obtain statistical representation of genetic brain network

structure from GWAS summary statistics. (1) GWAS summary statistics for 83 grey-matter

volumes in UK Biobank from European ancestry were used as input data ($N = 36,778$). They

were calculated as described in Methods and are publicly available. (2) Linkage

disequilibrium score regression (LDSC) was used to infer genetic correlations between 83

brain volumes using GenomicSEM [32]. (3) Genetic correlations are analysed using PCA to

derive PC loadings on the first PC, representing an underlying dimension of shared

morphometry. (4) We developed a method to derive univariate summary statistics for a

genetic PC of multiple GWAS phenotypes (derived from samples of unknown degrees of

overlap). A genetic PC underlying several brain volumes is interpreted throughout the

manuscript to index general dimensions of regionally shared morphometry. Genome-wide

SNP effects are calculated as an average of all SNP effects contributed by multiple

phenotypes, weighted by their respective PC loadings. Standard errors are computed using a

method that corrects for sample overlap, as estimated by LDSC. We have validated this novel

approach in an independent set of GWAS summary statistics [59]. All software we used is

available on https://github.com/. [1] The software by Baselmans, Jansen [58], containing the

GWAMA function is available at https://github.com/baselmans/multivariate_GWAMA/. [2]

Our modified version of the GWAMA function is at

https://github.com/AnnaFurtjes/Genetic_networks_project/blob/main/my_GWAMA_260320

20.R and [3] a step-by-step demonstration of genomic PCA is at

https://annafurtjes.github.io/genomicPCA/.

Fig. 3. **(A)** Distribution of SNP-heritability estimates for 83 regional grey-matter volumes

inferred through univariate LDSC. **(B)** Distribution of genetic correlations among 83 regional

grey-matter volumes inferred through between-region LDSC. This figure only depicts

between-region correlations but not the very high genetic inter-region correlations between

regions and their homologous counterpart in the opposite hemisphere (excluding brain stem).

(C) Distribution of phenotypic correlations among 83 regional grey-matter volumes inferred

through Pearson's correlations. The raincloud plots were created based on code adapted from

Allen et al. [62].

Fig. 4. (A) Density distributions of principal component (PC) loadings on the first PC

underlying phenotypic and (B) genetic networks. Vertical lines indicate quantiles. (C)

Variance explained by phenotypic and genetic first PC in each network.

Fig. 5. **(A)** Association between phenotypic and genetic between-region correlations of 83 regional grey-matter volumes. The dashed red line is the line of identity, with a slope of 1 and an intercept of 0. The dashed grey line indicates $r_g = 0$. **(B)** Correlation between phenotypic and genetic PC loadings on the first PC underlying 83 regional volumes. The dashed red line is the line of identity. **(C)** Correlation between *phenotypic* PC loadings and age sensitivity as indexed by phenotypic cross-sectional age-volume correlations. **(D)** Correlation between *genetic* PC loadings and age sensitivity as indexed by phenotypic cross-sectional age-volume correlations.

Tables

Table 1. Genetic correlations between general cognitive ability and nine canonical brain networks

Network	Included	r_g	95% CI	*p*-value	FDR *q*-value

	volumes				
Whole brain	83	0.21	0.13-0.29	1.00×10^{-7}	3.00×10^{-7}
Central executive	8	0.20	0.12-0.27	1.00×10^{-7}	3.00×10^{-7}
Cingulo-opercular	10	0.20	0.13-0.27	1.00×10^{-7}	3.00×10^{-7}
Default Mode	16	0.19	0.12-0.26	2.00×10^{-7}	3.00×10^{-7}
Hippocampal-Diencephalic	12	0.17	0.09-0.24	2.66×10^{-5}	2.66×10^{-5}
Multiple Demand	12	0.19	0.12-0.27	7.00×10^{-7}	9.00×10^{-7}
P-FIT	36	0.20	0.12-0.27	2.00×10^{-7}	3.00×10^{-7}
Salience	10	0.19	0.12-0.26	3.00×10^{-7}	4.00×10^{-7}
Sensorimotor	12	0.19	0.11-0.27	1.20×10^{-7}	1.30×10^{-6}
Temporo-amygdala-orbitofrontal	30	0.20	0.12-0.27	2.00×10^{-7}	4.00×10^{-7}

r_g = genetic correlation between brain network and a factor of general cognitive ability modelled from seven

cognitive traits, SE = standard error, 95% CI = 95% confidence interval, *p*-value = original *p*-value as indicated

by the GenomicSEM model, false discovery rate (FDR) *q*-value = *p*-value corrected using 5% false discovery

rate.

Supplemental Information titles and legends

Supplementary Table 1. 83 cortical and subcortical grey-matter regions of interest

Supplementary Table 2. Network characterisation

Supplementary Table 3. Explained variance and descriptive statistics of PC loadings within

phenotypic canonical networks

Supplementary Table 4. Model fit for genetic correlations between genetic general cognitive

ability and each canonical network

Supplementary Table 5. Fit indices for the comparison between freely-varying or constrained

correlations with general cognitive ability between central executive and other

networks

Supplementary Table 6. Fit indices for the *adjusted* comparison between freely-varying or

constrained correlations with general cognitive ability between central executive and

other networks

Supplementary Table 7. Genetic correlations between three cognitive abilities and brain

networks

Supplementary Table 8. Canonical networks explain more variance than networks with

randomly included volumes

Supplementary Table 9. Associations between brain volumes and potential covariates

Supplementary Table 10. Genetic quality control exclusion criteria resulting in a total GWAS

sample of 36,778 out of 39,947 participants

Supplementary Fig. 1. Genetic correlation matrix inferred through LDSC across the whole

brain (83 volumes).

Supplementary Fig. 2. Genetic correlations inferred through LDSC among the central

executive network (8 volumes).

Supplementary Fig. 3. Genetic correlations inferred through LDSC among the cingulo-

opercular network (10 volumes).

Supplementary Fig. 4. Genetic correlations inferred through LDSC among the default mode

network (16 volumes).

Supplementary Fig. 5. Genetic correlations inferred through LDSC among the hippocampal-

diencephalic network (12 volumes).

Supplementary Fig. 6. Genetic correlations inferred through LDSC among the multiple

demand network (12 volumes).

Supplementary Fig. 7. Genetic correlations inferred through LDSC among the P-FIT network

(36 volumes).

Supplementary Fig. 8. Genetic correlations inferred through LDSC among the salience

network (10 volumes).

Supplementary Fig. 9. Genetic correlations inferred through LDSC among the sensorimotor

network (12 volumes).

Supplementary Fig. 10. Genetic correlations inferred through LDSC among the temporo-

amygdala-orbitofrontal network (30 volumes).

Supplementary Fig. 11. Parallel analysis in the central executive network

Supplementary Fig. 12. Parallel analysis in the cingulo-operular network

Supplementary Fig. 13. Parallel analysis in the default mode network

Supplementary Fig. 14. Parallel analysis in the hippocampal-diencephalic network

Supplementary Fig. 15. Parallel analysis in the multiple demand network

Supplementary Fig. 16. Parallel analysis in the P-FIT network

Supplementary Fig. 17. Parallel analysis in the salience network

Supplementary Fig. 18. Parallel analysis in the sensorimotor network

Supplementary Fig. 19. Parallel analysis in the temporo-amygdala-orbitofrontal network

Supplementary Fig. 20. Parallel analysis in the whole brain with 83 nodes

Supplementary Fig. 21. Genetic correlations between seven cognitive traits and brain

868   networks. Descriptively, performance in the Tower Rearranging Task has the largest

869   association with brain networks in comparison with other cognitive tasks.

870   Abbreviations: Matrix = Matrix Pattern Completion task; Memory = Memory – Pairs

871   Matching Test; RT = Reaction Time; Symbol Digit = Symbol Digit Substitution Task;

872   Trails-B = Trail Making Test – B; Tower = Tower Rearranging Task; VNR = Verbal

873   Numerical Reasoning Test; central exec = central executive; cingulo = cingulo-

874   opercular; hippocampal = hippocampal-diencephalic; multiple = multiple demand; p fit

875   = parieto-frontal integration theory; sensori = sensorimotor; temporo = temporo-

876   amygdala-orbitofrontal

877   *Supplementary Fig. 22.* Genetic correlation between the central executive network and factor *g*

878   modelled for correlation structure of seven cognitive traits. The seven cognitive traits and the

879   network are inferred through LDSC, and the factor through factor analysis. Matrix = Matrix

880   Pattern Completion task; Memory = Memory – Pairs Matching Test; RT = Reaction Time;

881   Symbol Digit = Symbol Digit Substitution Task; Trails-B = Trail Making Test – B; Tower =

882   Tower Rearranging Task; VNR = Verbal Numerical Reasoning Test. Model fit: χ^2 = 124.04,

883   df = 20, *p*-value = 2.1 x10^{-20}, AIC = 174.04, CFI = 0.97, SRMR = 0.079

Supplementary Fig. 23. Illustration of the genomic structural equation models used to test

whether correlation magnitudes with genetic general cognitive ability differ between

the central executive network and other significantly associated networks. The model

on the right freely estimates correlation parameters between two networks and genetic

g while allowing for correlations between the networks. In the left model, we force

the correlation magnitudes to be the same, and assess whether model fit deteriorates

significantly, to conclude whether correlation magnitudes between networks are likely

different from each other.

Supplementary Fig. 24. Structural equation models to calculate Q_{trait} heterogeneity indices

References

1. Fjell AM, Walhovd KB. (2010). Structural Brain Changes in Aging: Courses, Causes and Cognitive
Consequences. Reviews in the Neurosciences, *21*(3), 187-222.
doi:10.1515/REVNEURO.2010.21.3.187
2. Cox SR, Ritchie SJ, Tucker-Drob EM, Liewald DC, Hagenaars SP, Davies G, et al. (2016). Ageing and
brain white matter structure in 3,513 UK Biobank participants. Nature Communications,
7(1), 13629. doi:10.1038/ncomms13629
3. Madole JW, Ritchie SJ, Cox SR, Buchanan CR, Hernández MV, Maniega SM, et al. (2021). Aging-
Sensitive Networks Within the Human Structural Connectome Are Implicated in Late-Life
Cognitive Declines. Biological Psychiatry, *89*(8), 795-806. doi:10.1016/j.biopsych.2020.06.010
4. Sporns O. (2011). The human connectome: a complex network. Annals of the New York Academy
of Sciences, *1224*(1), 109-25. doi:10.1111/j.1749-6632.2010.05888.x
5. Power Jonathan D, Cohen Alexander L, Nelson Steven M, Wig Gagan S, Barnes Kelly A, Church
Jessica A, et al. (2011). Functional Network Organization of the Human Brain. Neuron, *72*(4),
665-78. doi:10.1016/j.neuron.2011.09.006
6. Yeo BTT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, et al. (2011). The
organization of the human cerebral cortex estimated by intrinsic functional connectivity.
Journal of Neurophysiology, *106*(3), 1125-65. doi:10.1152/jn.00338.2011
7. Bressler SL, Menon V. (2010). Large-scale brain networks in cognition: emerging methods and
principles. Trends in Cognitive Sciences, *14*(6), 277-90. doi:10.1016/j.tics.2010.04.004
8. Raz N, Ghisletta P, Rodrigue KM, Kennedy KM, Lindenberger U. (2010). Trajectories of brain aging
in middle-aged and older adults: Regional and individual differences. NeuroImage, *51*(2),
501-11. doi:https://doi.org/10.1016/j.neuroimage.2010.03.020
9. Zhao B, Luo T, Li T, Li Y, Zhang J, Shan Y, et al. (2019). Genome-wide association analysis of 19,629
individuals identifies variants influencing regional brain volumes and refines their genetic co-
architecture with cognitive and mental health traits. Nature Genetics, *51*(11), 1637-44.
doi:10.1038/s41588-019-0516-6
10. Anderson KM, Ge T, Kong R, Patrick LM, Spreng RN, Sabuncu MR, et al. (2021). Heritability of
individualized cortical network topography. Proceedings of the National Academy of
Sciences, *118*(9), e2016271118. doi:10.1073/pnas.2016271118
11. Meer Dvd, Kaufmann T, Shadrin AA, Makowski C, Frei O, Roelfs D, et al. (2021). The genetic
architecture of human cortical folding. Science Advances, *7*(51), eabj9446.
doi:doi:10.1126/sciadv.abj9446
12. de la Fuente J, Davies G, Grotzinger AD, Tucker-Drob EM, Deary IJ. (2021). A general dimension of
genetic sharing across diverse cognitive traits inferred from molecular data. Nature Human
Behaviour, *5*(1), 49-58. doi:10.1038/s41562-020-00936-2
13. Arnatkevičiūtė A, Fulcher BD, Oldham S, Tiego J, Paquola C, Gerring Z, et al. (2021). Genetic
influences on hub connectivity of the human connectome. Nature Communications, *12*(1),
4237. doi:10.1038/s41467-021-24306-2
14. Zhao B, Ibrahim JG, Li Y, Li T, Wang Y, Shan Y, et al. (2019). Heritability of Regional Brain Volumes
in Large-Scale Neuroimaging and Genetic Studies. Cerebral Cortex, *29*(7), 2904-14.
doi:10.1093/cercor/bhy157
15. de Vlaming R, Slob EAW, Jansen PR, Dagher A, Koellinger PD, Groenen PJF, et al. (2021).
Multivariate analysis reveals shared genetic architecture of brain morphology and human
behavior. bioRxiv, 2021.04.19.440478. doi:10.1101/2021.04.19.440478
16. Cheverud JM. (1988). A Comparison of Genetic and Phenotypic Correlations. Evolution, *42*(5),
958-68. doi:10.2307/2408911
17. Sodini SM, Kemper KE, Wray NR, Trzaskowski M. (2018). Comparison of Genotypic and
Phenotypic Correlations: Cheverud's Conjecture in Humans. Genetics, *209*(3), 941-8.
doi:10.1534/genetics.117.300630

18. Smith SM, Fox PT, Miller KL, Glahn DC, Fox PM, Mackay CE, et al. (2009). Correspondence of the
brain's functional architecture during activation and rest. Proceedings of the National
Academy of Sciences, *106*(31), 13040-5. doi:10.1073/pnas.0905267106
19. Uddin LQ, Yeo BTT, Spreng RN. (2019). Towards a Universal Taxonomy of Macro-scale Functional
Human Brain Networks. Brain Topography, *32*(6), 926-42. doi:10.1007/s10548-019-00744-6
20. Jung RE, Haier RJ. (2007). The Parieto-Frontal Integration Theory (P-FIT) of intelligence:
Converging neuroimaging evidence. Behavioral and Brain Sciences, *30*(2), 135-54.
doi:10.1017/S0140525X07001185
21. Menon V, Uddin LQ. (2010). Saliency, switching, attention and control: a network model of insula
function. Brain Structure and Function, *214*(5), 655-67. doi:10.1007/s00429-010-0262-0
22. Sridharan D, Levitin DJ, Menon V. (2008). A critical role for the right fronto-insular cortex in
switching between central-executive and default-mode networks. Proceedings of the
National Academy of Sciences, *105*(34), 12569-74. doi:10.1073/pnas.0800005105
23. Buckner RL, DiNicola LM. (2019). The brain's default network: updated anatomy, physiology and
evolving insights. Nature Reviews Neuroscience, *20*(10), 593-608. doi:10.1038/s41583-019-
0212-7
24. Downar J, Crawley AP, Mikulis DJ, Davis KD. (2002). A Cortical Network Sensitive to Stimulus
Salience in a Neutral Behavioral Context Across Multiple Sensory Modalities. Journal of
Neurophysiology, *87*(1), 615-20. doi:10.1152/jn.00636.2001
25. Li R, Zhang S, Yin S, Ren W, He R, Li J. (2018). The fronto-insular cortex causally mediates the
default-mode and central-executive networks to contribute to individual cognitive
performance in healthy elderly. Human Brain Mapping, *39*(11), 4302-11.
doi:10.1002/hbm.24247
26. Duncan J. (2010). The multiple-demand (MD) system of the primate brain: mental programs for
intelligent behaviour. Trends in Cognitive Sciences, *14*(4), 172-9.
doi:10.1016/j.tics.2010.01.004
27. Cox SR, Ritchie SJ, Fawns-Ritchie C, Tucker-Drob EM, Deary IJ. (2019). Structural brain imaging
correlates of general intelligence in UK Biobank. Intelligence, *76*, 101376.
doi:10.1016/j.intell.2019.101376
28. Hilger K, Winter NR, Leenings R, Sassenhagen J, Hahn T, Basten U, et al. (2020). Predicting
intelligence from brain gray matter volume. Brain Structure and Function, *225*(7), 2111-29.
doi:10.1007/s00429-020-02113-7
29. Cole JH, Ritchie SJ, Bastin ME, Valdés Hernández MC, Muñoz Maniega S, Royle N, et al. (2018).
Brain age predicts mortality. Molecular Psychiatry, *23*(5), 1385-92. doi:10.1038/mp.2017.62
30. Cole JH, Franke K. (2017). Predicting Age Using Neuroimaging: Innovative Brain Ageing
Biomarkers. Trends in Neurosciences, *40*(12), 681-90. doi:10.1016/j.tins.2017.10.001
31. Mbatchou J, Barnard L, Backman J, Marcketta A, Kosmicki JA, Ziyatdinov A, et al. (2021).
Computationally efficient whole-genome regression for quantitative and binary traits.
Nature Genetics. doi:10.1038/s41588-021-00870-7
32. Grotzinger AD, Rhemtulla M, de Vlaming R, Ritchie SJ, Mallard TT, Hill WD, et al. (2019). Genomic
structural equation modelling provides insights into the multivariate genetic architecture of
complex traits. Nature Human Behaviour, *3*(5), 513-25. doi:10.1038/s41562-019-0566-x
33. Bulik-Sullivan B, Finucane HK, Anttila V, Gusev A, Day FR, Loh P-R, et al. (2015). An atlas of
genetic correlations across human diseases and traits. Nature Genetics, *47*(11), 1236.
doi:10.1038/ng.3406
34. Lorenzo-Seva U, Berge JMFt. (2006). Tucker's Congruence Coefficient as a Meaningful Index of
Factor Similarity. Methodology, *2*(2), 57-64. doi:10.1027/1614-2241.2.2.57
35. Hu L-t, Bentler PM. (1998). Fit indices in covariance structure modeling: Sensitivity to
underparameterized model misspecification. Psychological Methods, *3*(4), 424.
doi:10.1037/1082-989X.3.4.424

36. Grotzinger AD, Mallard TT, Akingbuwa WA, Ip HF, Adams MJ, Lewis CM, et al. (2020). Genetic
Architecture of 11 Major Psychiatric Disorders at Biobehavioral, Functional Genomic, and
Molecular Genetic Levels of Analysis. medRxiv, 2020.09.22.20196089.
doi:10.1101/2020.09.22.20196089
37. Kaufmann T, van der Meer D, Doan NT, Schwarz E, Lund MJ, Agartz I, et al. (2019). Common brain
disorders are associated with heritable patterns of apparent aging of the brain. Nature
Neuroscience, 22(10), 1617-23. doi:10.1038/s41593-019-0471-7
38. Comas-Herrera A, Wittenberg R, Pickard L, Knapp M. (2007). Cognitive impairment in older
people: future demand for long-term care services and the associated costs. International
Journal of Geriatric Psychiatry, 22(10), 1037-45. doi:https://doi.org/10.1002/gps.1830
39. Tucker-Drob EM. (2019). Cognitive Aging and Dementia: A Life-Span Perspective. Annual Review
of Developmental Psychology, 1(1), 177-96. doi:10.1146/annurev-devpsych-121318-085204
40. Jansen PR, Nagel M, Watanabe K, Wei Y, Savage JE, de Leeuw CA, et al. (2020). Genome-wide
meta-analysis of brain volume identifies genomic loci and genes shared with intelligence.
Nature Communications, 11(1), 5606. doi:10.1038/s41467-020-19378-5
41. Herculano-Houzel S. (2017). Numbers of neurons as biological correlates of cognitive capability.
Current Opinion in Behavioral Sciences, 16, 1-7. doi:10.1016/j.cobeha.2017.02.004
42. Fry A, Littlejohns TJ, Sudlow C, Doherty N, Adamska L, Sprosen T, et al. (2017). Comparison of
Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those
of the General Population. American Journal of Epidemiology, 186(9), 1026-34.
doi:10.1093/aje/kwx246
43. Alfaro-Almagro F, Jenkinson M, Bangerter NK, Andersson JLR, Griffanti L, Douaud G, et al. (2018).
Image processing and Quality Control for the first 10,000 brain imaging datasets from UK
Biobank. NeuroImage, 166, 400-24. doi:10.1016/j.neuroimage.2017.10.034
44. Smith S, Alfaro-Almagro F, Miller K. UK Biobank Brain Imaging Documentation. 2020 [Available
from: https://biobank.ctsu.ox.ac.uk/crystal/crystal/docs/brain_mri.pdf.
45. Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. (2006). An automated
labeling system for subdividing the human cerebral cortex on MRI scans into gyral based
regions of interest. NeuroImage, 31(3), 968-80. doi:10.1016/j.neuroimage.2006.01.021
46. Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, et al. (2002). Whole Brain
Segmentation: Automated Labeling of Neuroanatomical Structures in the Human Brain.
Neuron, 33(3), 341-55. doi:10.1016/S0896-6273(02)00569-X
47. Lindroth H, Nair VA, Stanfield C, Casey C, Mohanty R, Wayer D, et al. (2019). Examining the
identification of age-related atrophy between T1 and T1+T2-FLAIR cortical thickness
measurements. Scientific Reports, 9(1), 11288. doi:10.1038/s41598-019-47294-2
48. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. (2018). The UK Biobank
resource with deep phenotyping and genomic data. Nature, 562(7726), 203-9.
doi:10.1038/s41586-018-0579-z
49. Coleman JRI, Peyrot WJ, Purves KL, Davis KAS, Rayner C, Choi SW, et al. (2020). Genome-wide
gene-environment analyses of major depressive disorder and reported lifetime traumatic
experiences in UK Biobank. Molecular Psychiatry, 25(7), 1430-46. doi:10.1038/s41380-019-
0546-6
50. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. (2015). Second-generation PLINK:
rising to the challenge of larger and richer datasets. GigaScience, 4(1). doi:10.1186/s13742-
015-0047-8
51. Warren HR, Evangelou E, Cabrera CP, Gao H, Ren M, Mifsud B, et al. (2017). Genome-wide
association analysis identifies novel blood pressure loci and offers biological insights into
cardiovascular risk. Nature Genetics, 49(3), 403-15. doi:10.1038/ng.3768
52. Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen W-M. (2010). Robust relationship
inference in genome-wide association studies. Bioinformatics, 26(22), 2867-73.
doi:10.1093/bioinformatics/btq559

53. McCarthy S, Das S, Kretzschmar W, Delaneau O, Wood AR, Teumer A, et al. (2016). A reference
panel of 64,976 haplotypes for genotype imputation. Nature Genetics, *48*(10), 1279-83.
doi:10.1038/ng.3643
54. Fawns-Ritchie C, Deary IJ. (2020). Reliability and validity of the UK Biobank cognitive tests. PLOS
ONE, *15*(4), e0231627. doi:10.1371/journal.pone.0231627
55. Smith SM, Douaud G, Chen W, Hanayik T, Alfaro-Almagro F, Sharp K, et al. (2020). Enhanced
Brain Imaging Genetics in UK Biobank. bioRxiv, 2020.07.27.223545.
doi:10.1101/2020.07.27.223545
56. Cole JH, Marioni RE, Harris SE, Deary IJ. (2019). Brain age and other bodily 'ages': implications for
neuropsychiatry. Molecular Psychiatry, *24*(2), 266-81. doi:10.1038/s41380-018-0098-1
57. Tucker LR. A method for synthesis of factor analysis studies. Educational Testing Service
Princeton Nj; 1951.
58. Baselmans BML, Jansen R, Ip HF, van Dongen J, Abdellaoui A, van de Weijer MP, et al. (2019).
Multivariate genome-wide analyses of the well-being spectrum. Nature Genetics, *51*(3), 445-
51. doi:10.1038/s41588-018-0320-8
59. Linnér RK, Biroli P, Kong E, Meddens SFW, Wedow R, Fontana MA, et al. (2019). Genome-wide
association analyses of risk tolerance and risky behaviors in over 1 million individuals identify
hundreds of loci and shared genetic influences. Nature Genetics, *51*(2), 245-57.
doi:10.1038/s41588-018-0309-3
60. Benjamini Y, Hochberg Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B
(Methodological), *57*(1), 289-300. doi:10.1111/j.2517-6161.1995.tb02031.x
61. Xia M, Wang J, He Y. (2013). BrainNet Viewer: A Network Visualization Tool for Human Brain
Connectomics. PLOS ONE, *8*(7), e68910. doi:10.1371/journal.pone.0068910
62. Allen M, Poggiali D, Whitaker K, Marshall T, Kievit R. (2019). Raincloud plots: a multi-platform
tool for robust data visualization [version 1; peer review: 2 approved]. Wellcome Open
Research, *4*(63). doi:10.12688/wellcomeopenres.15191.1

1074

1075

1076

1077

1078

1079

1080