Title: Pattern Recognition Receptor for Bacterial Lipopolysaccharide in the Cytosol of Human Macrophages

Authors: Maricarmen Rojas-Lopez[1†], Amanda S. Zajac[1†], Thomas E. Wood[1†], Kelly A. Miller[1†], María Luisa Gil-Marqués[1], Austin C. Hachey[1], Vritti Kharbanda[1], Keith T. Egger[1], Marcia B. Goldberg[1,2,3,4]*

Affiliations:

[1]Center for Bacterial Pathogenesis, Division of Infectious Diseases, Massachusetts General Hospital, Boston, MA, USA.

[2]Department of Microbiology, Blavatnik Institute, Harvard Medical School, Boston, MA, USA.

[3]Broad Institute of MIT and Harvard, Cambridge, MA, USA.

[4]Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Boston, MA, USA.

[†]These authors contributed equally to this work.

*Corresponding author. Email: marcia.goldberg@mgh.harvard.edu

Abstract:

Endotoxin, a component of bacterial lipopolysaccharide (LPS), is a driver of the lethal infection sepsis through activation of innate immune responses. When delivered to the cytosol of macrophages, LPS (cLPS) induces the assembly of an inflammasome that contains caspases-4/5 in humans or caspase-11 in mice. Whereas activation of all other inflammasomes is triggered by sensing of pathogen products by a specific host cytosolic pattern recognition receptor protein, whether pattern recognition receptors for cLPS exist has been doubted by many investigators, as caspases-4, -5, and -11 bind and activate LPS directly *in vitro*. Here we show that the primate-specific protein NLRP11 is a pattern recognition receptor for cLPS required for efficient activation of the caspase-4 inflammasome in human macrophages. *NLRP11* is present in humans and other primates, but absent in mice, likely explaining why it has been missed in screens looking for innate immune signaling molecules, most of which have been carried out in mice. NLRP11 is a previously missing link and a component of the human caspase-4 inflammasome activation pathway.

One Sentence Summary: Discovery that human macrophages contain a cytosolic sensor for endotoxin-containing bacterial lipopolysaccharide.

Main Text:

INTRODUCTION

Inflammasomes assemble in response to the presence in the cytosol of pathogen-associated molecules, including dsDNA, flagellin, type 3 secretion system structural proteins, pore-forming toxins, ion flux, ATP, and LPS (*1-4*). Canonical inflammasomes consist of a pattern recognition receptor, which senses a pathogen-associate molecule protein, and an inflammatory caspase, frequently linked by an adaptor protein. Inflammasome assembly activates inflammatory caspases, which proteolytically activate the pore-forming protein gasdermin D to form plasma membrane pores, leading to pyroptotic cell death (*5-9*). The presence of bacterial LPS in the cytosol of human macrophages activates a non-canonical inflammasome, which in humans, contains caspases-4 and -5, and in mice, caspase-11 (*10-12*), and which also leads to caspase-mediated processing of gasdermin D, pore formation, and pyroptotic cell death. To date, no pattern recognition receptor protein of cytosolic LPS (cLPS) has been identified, and because caspase-4, -5, and -11 bind LPS *in vitro* (*13*), the current paradigm is that the non-canonical inflammasome lacks a pattern recognition receptor.

Here we identify primate specific nucleotide-binding oligomerization domain, leucine rich repeat and pyrin domain containing protein-11 (NLRP11) as a cytosolic pattern recognition receptor of bacterial LPS in human macrophages. We show that efficient activation of the non-canonical pathway in human-derived macrophages depends on NLRP11, and that NLRP11-dependent cell death is specifically due to detection of cLPS. Our results demonstrate that NLRP11 senses cLPS and promotes LPS-dependent activation of caspase-4.

RESULTS

NLRP11 required for cell death due to intracellular gram-negative pathogens

In a forward genetic screen that enriched for genes required for death during infection with the gram-negative intracellular bacterial pathogen *Shigella flexneri*, we identified NLRP11 (*14*) (Fig. 1A, and fig. S1); the screen was conducted using a human-derived myeloid lineage cell line. The NLRP family includes proteins such as NLRP1 and NLRP3 that sense a range of pathogen-associated molecular patterns in the cell cytosol. In human-derived monocyte THP-1 cells containing either of two targeted deletions of *nlrp11* (fig. S2), cell death (release of lactate dehydrogenase, LDH) during *S. flexneri* infection was decreased by >50% (Fig. 1B). *S. flexneri* mutants unable to access the cytosol by virtue of inactivation of the type 3 secretion system induced minimal cell death, indicating that cell death depended on bacterial cytosolic localization and/or type 3 secretion activity (fig. S3A).

To test whether the dependence on NLRP11 for efficient cell death was generalizable to other intracellular gram-negative bacterial pathogens, we infected *NLRP11*$^{-/-}$ cells with the cytosolic pathogen *Burkholderia thailandensis* and the largely vacuolar pathogen *Salmonella enterica* serovar Typhimurium. As with *S. flexneri*, in *NLRP11*$^{-/-}$ macrophages, cell death during infection with *B. thailandensis* or *S.* Typhimurium was significantly reduced (Fig. 1, C and D). Upon viral infection and certain Toll-like receptor interactions, the NF-κB pathway is repressed by NLRP11-dependent targeted degradation of TRAF6 (*15-17*); however, *S. flexneri*-induced NLRP11-dependent cell death was independent of TRAF6, whether or not *Shigella* OspI, which suppresses TRAF6 signaling (*18*), was present (fig. S3, B, C, and D), indicating that the role of NLRP11 in bacterial infection-induced cell death is distinct from its role in NF-κB signaling.

Thus, NLRP11 is required for efficient cell death induced by multiple intracellular gram-negative pathogens.

Bacterial LPS recognized by NLRP11 in cytosol

We tested whether LPS, which is specific to gram-negative bacteria, might be required for NLRP11-dependent cell death. Infection of THP-1 macrophages with the cytosolic gram-positive pathogen *Listeria monocytogenes* induced minimal levels of cell death during the experimental timeframe, but when coated with *S. enterica* LPS, induced cell death that was largely dependent on NLRP11 (Fig. 2A). The observed cell death was dependent on cytosolic localization, as *L. monocytogenes* lacking listeriolysin O (LLO), which is required to lyse the vacuole, induced minimal cell death. Introduction of LPS alone directly into the cytosol by electroporation similarly induced cell death that was significantly dependent on NLRP11, both in *NLRP11*$^{-/-}$ and NLRP11 siRNA-treated THP-1 macrophages (Fig. 2, B and C), whereas cell death induced by extracellular LPS was independent of NLRP11 (fig. S4). Therefore, NLRP11 functions in a cell death pathway that is activated by LPS localized to the cytosol (cLPS).

For pathogens such as *Salmonella* that do not themselves efficiently lyse the vacuole, delivery of LPS to the cytosol occurs upon guanylate-binding protein (GBP) 2-mediated vacuolar lysis (*19*). This provides a potential explanation for the lack of differential NLRP11-dependent cell death induced by wild-type *S.* Typhimurium, which is largely vacuolar, and *sifA* mutant *S.* Typhimurium, which is predominantly cytosolic (*20*) (Fig. 1D).

NLRP11 required for efficient activation of non-canonical inflammasome

Activation of caspase-4 and gasdermin D were dependent on NLRP11; during *S. flexneri* infection, processing of each, which releases their active domains, was significantly reduced in *NLRP11*[-/-] macrophages (Fig. 3). Upon *CASP4* depletion, the absence of NLRP11 caused no significant additional decrease in cell death induced by LPS transfection (Fig. 4 and fig. S5A). Depletion of *CASP1* or *MyD88* induced no significant decrease in cell death in response to cLPS as compared to control depletion for either *NLRP11*[-/-] or WT THP-1 macrophages (Fig. 4 and fig. S5, B and C). Despite testing several antibodies, we could not detect native cellular NLRP11 by western blot, presumably due to very low levels of native protein. These data indicate that NLRP11 is required for efficient activation of the non-canonical inflammasome, induces caspase-4-dependent cell death in the presence of LPS, and that NLRP11 function is independent of caspase-1 and extracellular Toll-like receptor signaling. Consistent with NLRP11 functioning specifically in the non-canonical inflammasome pathway, NLRP11 was not required for activation of the canonical NLRP3 inflammasome (fig. S6).

NLRP11 interaction with LPS and caspase-4

In contrast to NLRP3 and NLR family CARD domain containing 4 (NLRC4) inflammasomes, activation of the non-canonical pathway is independent of the adaptor protein ASC (*10, 13, 15*). We tested whether NLRP11 interacts with other core components of the non-canonical pathway by ectopic expression of selected components in HEK293T cells, which lack most innate immune proteins, including ASC and inflammatory caspases. Biotin-conjugated LPS precipitated NLRP11 in the absence of caspase-4 (Fig. 5A), indicating that NLRP11 binds LPS independently of caspase-4. LPS specifically precipitated caspase-4 and its catalytically inactive derivative (CASP4 C258A), as described (*13*), including in the presence of NLRP11 (Fig. 5B), whereas the triacylated lipid Pam3CSK4 did not precipitate either NLRP11 or caspase-4 (Fig.

5A). In addition to binding to LPS independent of caspase-4, NLRP11 specifically precipitated caspase-4 (Fig. 5C). We developed a plate binding assay using FLAG-NLRP11 in HEK293T lysates and purified caspase-4 (fig S7). FLAG-NLRP11 binding depended on caspase-4, as in the absence of caspase-4, bound FLAG-NLRP11 was at background levels (Fig. 5D, diamonds). Binding of NLRP11 to caspase-4 increased as a function of increasing concentrations of both NLRP11 lysate and caspase-4, resembling the asymptotic phase of a sigmoidal curve, beginning to reach saturation at 8-10 µg/ml caspase-4 coating the wells (Fig. 5D).

Caspase specificity of NLRP11

In addition to caspase-4, NLRP11 co-precipitated with caspase-1; it did not co-precipitate with caspase-5, a paralog of caspase-4 that binds LPS and a component of some non-canonical inflammasomes (*13, 21, 22*), or the more divergent caspase-9 (Fig. 6 and fig. S8). Although caspase-1 co-precipitates with NLRP11 (Fig. 6), NLRP11 does not activate caspase-1 under conditions in which AIM2 does (*15*), and NLRP11 function in LPS-induced cell death of THP-1 macrophages is independent of caspase-1 (Fig. 4), indicating that the observed interaction of caspase-1 with NLRP11 contributes minimally to activation of the non-canonical inflammasome.

NLRP11 role independent of interferon-gamma and guanylate-binding protein-1

Host GBPs are cytosolic proteins that bind intracellular LPS, promote non-canonical inflammasome activation (*23-27*), and are encoded by genes that are strongly induced by interferon-gamma (IFNγ) (*28*). Human GBP1 binds LPS on the *S. flexneri* surface, whereupon it enhances caspase-4 recruitment (*29*). We tested whether IFNγ stimulation altered the role of NLRP11 in activation of the non-canonical inflammasome. IFNγ pre-treatment of THP-1 macrophages induced GBP1, as expected, yet NLRP11 was still required for efficient activation

of *S. flexneri*-induced cell death (Fig. S9, A and B). Moreover, during the timeframe of this infection, GBP1 was not induced by *S. flexneri* infection alone, depending exclusively on pre-treatment with IFNγ (Fig. S9C), indicating that GBP1 is not required for non-canonical inflammasome induced pyroptosis or for NLRP11 function in inducing non-canonical pyroptosis. These results also indicate during *S. flexneri* infection of human macrophages, priming is not required for activation of the non-canonical inflammasome. Moreover, IFNγ and GBP1 do not bypass the requirement for NLRP11 in this pathway.

Requirement for NLRP11 in primary human macrophages

As in THP-1 macrophages, in primary human monocyte-derived macrophages, siRNA depletion of NLRP11 was associated with significantly diminished cLPS-induced cell death (Fig. 7A). Cell death depended on caspase-4, but not MyD88 (Fig. 7, B and C, and fig. S10), indicating that the trigger was cLPS rather than extracellular LPS, and demonstrating that the requirement for NLRP11 in the non-canonical inflammasome pathway extends to primary human cells.

DISCUSSION

The current dogma for LPS recognition – that caspase-4 is the sensor for cytosolic LPS – is based on the simplest interpretation of data that show that *in vitro* caspase-4 binds, is induced to oligomerize, and is activated by LPS (*13*). LPS consists of lipid A, an acylated hydrophobic diglucosamine backbone that anchors LPS in the bacterial outer membrane, core oligosaccharide, and serotype-specific O-antigen. The moiety of LPS that binds caspase-4 is lipid A (*13*). We speculate that caspase-4 binding and activation by LPS under the described *in vitro* conditions (*13*) is due, at least in part, to the presence of a multiligand platform of lipid A presented by LPS in micelles or aggregates. Like other caspases, activation of caspase-4 is triggered by proximity-induced dimerization (*13, 30-37*); dimerization is sufficient to activate the caspase to undergo auto-processing, thereby releasing the small and large active subunits, which form an active heterotetrametric complex. As LPS species self-associate in micelles and aggregates at concentrations above critical micelle concentrations estimated to be 40 nM or lower (*38*), and caspase-4 binds preferentially to LPS in micelles or bacterial outer membrane vesicles (*39*), it seems likely that LPS micelles and aggregates provide a multiligand platform for caspase-4 dimerization. Supporting this possibility is that although at 100-250 nM, purified LPS binds, oligomerizes, and activates caspase-4 *in vitro*, in the presence of >0.25% Tween-20, LPS-mediated caspase-4 oligomerization and activation are disrupted (*13*).

The data presented here identify NLRP11 as a primate-specific pattern recognition receptor of bacterial LPS in the cytosol of human macrophages. As such, NLRP11 is a critical component activation of the caspase-4 dependent inflammasome and a critical mediator of caspase-4-dependent pyroptosis. Nevertheless, it is likely that delivery of high concentrations of LPS to the macrophage cytosol is sufficient to activate caspase-4 directly and independently of NLRP11,

perhaps principally when multiligand LPS platforms are present. Consistent with this, a proportion of caspase-4-dependent cell death appeared to be independent of NLRP11 (Fig. 4); in these experiments, LPS was transfected at relatively high concentration, and cells were infected with ten bacteria per cell. We speculate that NLRP11 may be particularly critical for non-canonical inflammasome activation when LPS is delivered in low amounts, such as during human infection with gram-negative pathogens, wherein the intracellular burden of organisms and intracellular LPS is likely very low.

Limitations of the investigations presented here are that, although performed with human cells, they were *ex vivo*, and that due to the restriction of NLRP11 to primates, no mouse model for testing the relevance of NLRP11 in animals currently exists. It remains uncertain whether the unique presence of NLRP11 in primates and the absence of identifiable homologs in mice contributes to the exquisite sensitivity of humans to endotoxic shock as compared with mice; humans display $\sim 10^4$-fold increased susceptibility to gram-negative bacterial sepsis and bloodstream LPS relative to mice (*40, 41*). Additional insights into these relationships would facilitate much-needed improved mouse models for the study of bacterial sepsis.

MATERIALS AND METHODS

Bacterial strains, plasmids, antibodies, and reagents. *Shigella flexneri* serotype 2a wildtype strain 2457T and the *S. flexneri* mutant strains *virB*::Tn5, BS103, and Δ*ospC3*, which are isogenic to 2457T, have been described previously (*42-45*). *Salmonella enterica* serovar Typhimurium and the isogenic Δ*sifA* mutant (*20*) were gifts from L. Knodler. *Burkholderia thailandensis* (*46*) was a gift from P. Cotter. *Listeria monocytogenes* 10403s and 10403 LLO⁻ (*47*) were gifts from M. Kielian and J. Miller. *S. flexneri* was grown overnight in tryptic soy broth at 37°C with aeration. *L. monocytogenes* was grown overnight in brain heart infusion broth at 30°C at a stationary slant. *B. thailandensis* was grown overnight in low salt Luria broth (LB) at 37°C with aeration. *S. enterica* Typhimurium was grown in LB broth at 37°C with aeration.

Caspase-1 (Abcam, ab207802) rabbit monoclonal antibody was used at 0.5 µg/mL (1:1000). Caspase-4 (Santa Cruz, sc-56056), and gasdermin D (Santa Cruz, sc-81868) mouse monoclonal antibodies were used at 0.5 µg/mL (1:200). FLAG M2 mouse monoclonal antibody (Sigma, F3165) was used at 76 ng/mL (1:5000). Myc rabbit monoclonal antibody (Cell Signaling, 2278) was used at 1:1000. GBP1 mouse monoclonal antibody (Abcam, ab119236) was used at 1:1000. For western blots of immunoprecipitations, the secondary antibody used was TrueBlot horseradish peroxidase-conjugated anti-mouse Ig (Rockland, 18-8817-31). For detecting β-actin, horseradish peroxidase-conjugated anti- β-actin antibody (Sigma, A3854) was used. For all other western blots, the secondary antibody used was horseradish peroxidase-conjugated goat anti-rabbit or goat anti-mouse IgG antibodies (Jackson ImmunoResearch, 115-035-003 or 111-035-144). All antibodies were diluted in 5% milk in TBST (0.1% Tween).

Strain and plasmid construction. The *S. flexneri ospI* deletion mutant was generated in the 2457T strain background using the lambda red method by introducing a chloramphenicol acetyltransferase (*cat*) cassette into the *orf169b* gene (which encodes OspI), as described (*48*).

pCMV-CASP4-MYC and pCMV-CASP5-MYC were made from pCMV-CASP4-MYC-FLAG and pCMV-CASP5-MYC-FLAG, gifts from S. Shin. The isoform of caspase-5 that was used was caspase-5/a, the most abundant isoform; this isoform contains nearly all sequences present in other isoforms. pCMV-CASP4-MYC C258A and pCMV-CASP4-MYC K19E C258A was generated by splicing by overlap extension PCR (SOE-PCR). pcDNA-NLRP11 (gift from H. Wu) was subcloned into pCMV-FLAG with an N-terminal FLAG tag for FLAG-NLRP11 and an N-terminal MYC tag for MYC-NLRP11. All FLAG tags are 3xFLAG. All N-terminal Myc tags are 6xMyc; all C-terminal Myc tags are 1xMyc. All DNA used for transfection was prepared LPS-free (Macherey-Nagel, 740410.50).

Generation of CRISPR knockouts. THP-1 cells were transduced with lentivirus packaged with pLentiCas9-Blast (Addgene, 52962) generated from HEK293T cells. THP-1 cell suspensions were treated with 8 µg/mL polybrene, combined with lentivirus and centrifuged in 12-well plates for 2 h at 1000 x g at 30°C. After centrifugation, cells were incubated for 2 h and washed once with fresh media. After 48 h, blasticidin (10 µg/mL) was added to the transduced cells. Guide RNA sequence ("sgRNA3", AGGCATGCAAAGCTGTCATG) was subcloned into pLentiGuide-Puro (Addgene, 59702). Guide RNA expressing plasmids were packaged into lentivirus using HEK293T cells. Polybrene treated THP-1 cells were transduced and selected with 1 µg/mL puromycin. Cells were limiting diluted and single cell clones were screened by SURVEYOR (IDT, 706020) and Sanger sequencing coupled with TIDE analysis (*49*).

Generation of *TRAF6* depletion cell lines. pLKO.1-shTRAF6 ((*50*), gift of A. Goldfeld), which encodes shRNA to *TRAF6*, was packaged into lentivirus in HEK293T cells. Lentivirus were introduced into NLRP11+/+ and NLRP11-/- THP-1 cell lines and cells containing pLKO.1-shTRAF6 were selected, expanded by maintenance in 0.8 µg/mL puromycin, as described (*50*), and stable clones were isolated by limiting dilution.

Mammalian cell culture. All cell lines were obtained from ATCC. HEK293T cells were maintained in Dulbecco's Modified Eagle's medium (Gibco, 11965118). THP1 cells were maintained in RPMI-1640 (Gibco, 21870092). All media was supplemented with 10% (vol/vol) fetal bovine serum (heat inactivated, sterile filtered, R&D Systems). For THP-1 cells, RPMI-1640 media was also supplemented with 10 mM HEPES (Gibco,15630080) and 2 mM L-glutamine (Gibco, 25030081). All cells were grown at 37°C in 5% CO_2. To differentiate monocyte cell lines into macrophage-like cells, 50 ng/mL phorbol-12-myristrate-13-acetate (PMA) was added 48 h before experiments. Where indicated, cells were pretreated with 10 ng/mL IFNγ 16-20 h prior to infection.

Precipitation experiments. HEK293T cells were transiently transfected using Fugene6 (Promega). A stock solution of 10x Buffer A (100 mM HEPES, 1500 mM NaCl, 10 mM EGTA, 1 mM $MgCl_2$, pH 7.4) was prepared. The cell culture medium was removed 24-48 h after transfection. Cells were washed with cold PBS and lysed in 1x Buffer A, 0.5% Triton-X. Cell lysates were incubated on ice for 15 min, vortexed for 15 s and centrifuged at 13,200 rpm for 10 min at 4°C. The pellet was discarded, and Bradford assay was used to determine protein concentration. 30 µL of FLAG M2 Magnetic Beads (Millipore-Sigma, M8823) were washed twice in 1x Buffer A, 0.1% Triton-X and equilibrated in the same buffer at 4°C for 2 h. 500 µg of

protein lysate was incubated with the beads for 2 h at 4°C. Beads were washed 3 times with 1x Buffer A, 0.1% Triton-X and eluted with 50 µL of 2x Laemmli buffer.

Streptavidin pulldown was largely as described (*13*), with minor modifications. Cell lysates were diluted to 4 mg/mL before a 2 h incubation with beads. For co-precipitation experiments, separate cells were transfected with each DNA construct, and pre-specified amounts of lysates were combined with each other or with untransfected lysate, such that the total amount of lysate were similar in each experimental condition; these lysates were then incubated at 37°C overnight. To reduce non-specific binding, lysates were pre-cleared with streptavidin beads and then were incubated for 2 h at 30°C with 4 µg biotinylated *E. coli* O111:B4 LPS loaded onto 8 µg of Streptavidin Mag sepharose (GE, 28985738, or Pierce, 88816).

Plate binding assays. Clear flat bottom Nunc Maxisorp ELISA plates (Thermo Fisher, 442404) were coated with various concentrations of purified recombinant caspase-4 (Origene, TP760359) overnight at 4°C. The plates were washed 3x with PBS then blocked for 2 h at RT with 1% bovine serum albumin (Fisher, BP9700100) in PBS. Excess blocking solution was removed. HEK 293T cells transiently expressing FLAG-NLRP11 and control untransfected cells were lysed into 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 5 mM $MgCl_2$, 100 µg/ml digitonin (Thermo Fisher Scientific, BN2006) and complete Mini EDTA-free protease inhibitor (Roche), as described (*26*). Total protein concentration of lysates was normalized to 2 mg/mL. Serial dilutions were in untransfected HEK293T cell lysate. Varying concentrations of cell lysates were incubated with pre-blocked ELISA plates for 1 h at RT. Excess lysate was removed by washing 3 times with PBS. For detection of FLAG-NLRP11, wells were incubated with 0.8 µg/mL anti-FLAG rabbit (Sigma, F7425-.2MG) was incubated for 1 h at RT, and after washing with PBS,

0.16 μg/mL goat-anti-rabbit conjugated to horseradish peroxidase (HRP) (Jackson, 111-035-144). To detect caspase-4 bound to the plate, wells were incubated with 0.5 μg/mL anti-caspase-4 (Santa Cruz, sc-56056) and, after washing with PBS, 0.16 μg/mL goat-anti-mouse conjugated to HRP (Jackson, 115-035-003), each for 1 hr. After 6 washes with PBS, 50 μL of One Step Ultra-TMB (Fisher Scientific, PI34029) was added to each well and incubated for 5 min before stopping with 4 N sulfuric acid and reading in a BioTek plate reader.

Bacterial infection. Overnight cultures were started from a single colony taken from a freshly streaked plate. Infections with *S. flexneri* were performed with exponential phase bacteria, after back-dilution from overnight cultures. Infections with *L. monocytogenes* were performed with stationary phase bacteria from overnight cultures. Infections were with a multiplicity of infection (MOI) of 10 for *S. flexneri* cultures and 1 for *L. monocytogenes*. Bacteria were centrifuged onto cells at 2000 rpm for 10 min, followed by incubation at 37°C. After 30 min, the cells were washed twice, and then 25 μg/mL gentamicin was added to the media to kill extracellular bacteria. Where appropriate *S. enterica* serovar Minnesota LPS was added to 10 μg/mL at the time of infection and MCC950 to 1 μM 30 min prior to infection and maintained during infection. For LDH assays of *S. flexneri* infected cells, 100 μL of cell culture supernatant was collected at 2 h and 30 min and at 5 h and/or 30 min of infection; for detection of caspases and/or gasdermin D by western blot, 100 μL of cell culture supernatant was collected at the same time points and was mixed with 35 μL 4x Laemmli buffer supplemented with cOmplete EDTA-free protease inhibitors (Roche, 5056489001). For parallel analyses of *L. monocytogenes* infected cells, gentamicin was used at 15 μg/mL, and samples were collected at 6 h of infection. For western blot analyses, whole cell lysates were collected in 2x Laemmli buffer.

Infections with *B. thailandensis* and *S.* Typhimurium were performed with exponential phase bacteria at a multiplicity of infection (MOI) of 10. Bacteria were centrifuged onto cells at 2000 rpm for 10 min, followed by incubation at 37°C. For *S.* Typhimurium, after 30 min incubation, the cells were washed twice, and then 25 μg/mL gentamicin was added to the media to kill extracellular bacteria. For *B. thailandensis*, after 2 h incubation, the cells were washed twice with cRPMI. For LDH assays with *S.* Typhimurium infection, 100 μL of cell culture supernatant was collected at 2 h and 30 min and at 5 h and 30 min of infection and, for LDH assays with *B. thailandensis*, 100 μL of cell culture supernatant was collected at 24 h.

Isolation of primary human peripheral blood mononuclear cells (PBMCs) and CD14 monocytes. Cells were isolated from leukopaks from individual healthy donors (Crimson Core Laboratories, Brigham and Women's Hospital) using density-gradient centrifugation. Briefly, blood was diluted 1:2 with RPMI-1640 (Gibco), layered on top of Ficoll-Paque Plus (GE Healthcare) using SepMate™-50 (IVD) (StemCell Technologies), and centrifuged at 1,200g for 20 min. The PBMC layer was collected and resuspended in 50 ml RPMI-1640 (Gibco) and centrifuged again at 300 g for 10 min. The cells were counted, resuspended in Cryostor CS10 (StemCell Technologies), and aliquoted in cryopreservation tubes at a concentration of 1×10^8 cells per mL. The tubes were placed at −80°C overnight, then transferred to liquid nitrogen for long-term storage. To isolate CD14+ cells, EasySep™ Human CD14 Positive Selection Kit II (StemCell Technologies) was used, following the manufacturer's instructions. Cells were cultured in complete RPMI, differentiated with 25 ng/mL m-CSF (StemCell Technologies), and maintained at 37°C in 5% CO_2.

LPS transfection. Electroporation of cells with LPS was performed using the Neon Transfection System (Invitrogen) according to the manufacturer's instructions using *S.* Minnesota LPS at indicated concentrations. THP-1 cells were seeded in low-attachment 10 cm dishes with 50 ng/mL PMA in RPMI. After 48 h, the cells were lifted in PBS (without calcium or magnesium). For electroporation, 1.5×10^6 THP-1 cells were pelleted, resuspended in Resuspension Buffer R with 2 μg/mL of LPS or water control. Using a 10 μL Neon tip, cells were pulsed once at 1400 V and 10 ms pulse width. For primary human macrophage cells, 7.5×10^5 CD14 macrophages were pelleted, then resuspended in Resuspension Buffer T with 2 μg/mL of LPS or water control. Using a 10 μL Neon tip, cells were pulsed 2 times at 2250 V and 20 ms pulse width. Electroporated cells were immediately resuspended in 2 mL of RPMI and placed into wells of a 24 well plate. Cells were incubated for 2.5 h at 37°C before measuring LDH release.

siRNA transient knockdowns. siRNAs used for NLRP11 knockdowns were Hs_NALP11_1 FlexiTube siRNA and Hs_NALP11_5 FlexiTube siRNA at 10 nM total (5 nM each siRNA); for caspase-4 knockdowns were 3941, 4036 and 4130 at 30 nM total (10 nM each siRNA) (ThermoFisher); for Caspase-1 Hs_CASP1_14 FlexiTube siRNA and Hs_CASP1_13 FlexiTube siRNA at 30 nM total (15 nM each siRNA); and for MyD88 Hs_MYD88_5 FlexiTube siRNA and Hs_MYD88_11 FlexiTube siRNA at 30 nM total (15 nM each siRNA) (Qiagen). AllStars Negative Control siRNAs (Qiagen) were used as negative controls at 10 nM or 30 nM, accordingly. siRNAs were transfected using the HiPerfect reagent (Qiagen) into primary human monocyte-derived macrophages 72 h before or into THP-1 cells 48 h before experimental assays, following the manufacturer's protocol for "Transfection of Differentiated Macrophage Cell Lines." After transfection, cells were maintained at 37ºC in 5% CO_2. Degree of knock-down was assessed by western blot of lysates of equivalent numbers of cells for each condition.

Nigericin. PMA differentiated THP-1 cells were washed once with fresh RPMI and primed with 1 µM O55:B5 *E. coli* LPS for 3 h, at which time 10 µM nigericin or vehicle control was added. LDH release was assayed at indicated times after treatment.

LDH assay. LDH was measured using an LDH Cytotoxicity Kit (Pierce, 88953) or Cytotoxicity Detection Kit (Sigma, 11644793001) per the manufacturer's instructions. Measurements at 490 nm and 680 nm were taken immediately after addition of Stop Solution using a BioTek Plate Reader. To determine LDH activity, the 680 nm absorbance value (background) was subtracted from the 490-nm absorbance value, and cytotoxicity was calculated for each sample as LDH activity as a fraction of the total LDH activity for cells treated with the Triton X-100 lysis buffer provided by the manufacturer. For experiments involving primary cells, a different donor was used for each biological replicate; due to inherent variability from donor to donor in the baseline levels of death in cell death assays, within-donor data were normalized to the negative control condition before compilation of the biological replicates.

Statistics and reproducibility. Unless otherwise indicated, all results are data from at least 3 independent biological replicates. On bar graphs, each individual symbol is data from an independent biological replicate. Statistical analysis was performed with Prism version 9.

Data availability. All relevant data are included in the manuscript and will be made available by request to the corresponding author, marcia.goldberg@mgh.harvard.edu.

List of Supplementary Materials:

Figs. S1 to S10

References and Notes:

1. S. M. Brewer, S. W. Brubaker, D. M. Monack, Host inflammasome defense mechanisms and bacterial pathogen evasion strategies. *Curr Opin Immunol* **60**, 63-70 (2019).
2. P. Broz, V. M. Dixit, Inflammasomes: mechanism of assembly, regulation and signalling. *Nat Rev Immunol* **16**, 407-420 (2016).
3. I. Jorgensen, M. Rayamajhi, E. A. Miao, Programmed cell death as a defence against infection. *Nat Rev Immunol* **17**, 151-164 (2017).
4. S. S. Wright, S. O. Vasudevan, V. A. Rathinam, Mechanisms and Consequences of Noncanonical Inflammasome-Mediated Pyroptosis. *J Mol Biol*, 167245 (2021).
5. N. Kayagaki, I. B. Stowe, B. L. Lee, K. O'Rourke, K. Anderson, S. Warming, T. Cuellar, B. Haley, M. Roose-Girma, Q. T. Phung, P. S. Liu, J. R. Lill, H. Li, J. Wu, S. Kummerfeld, J. Zhang, W. P. Lee, S. J. Snipas, G. S. Salvesen, L. X. Morris, L. Fitzgerald, Y. Zhang, E. M. Bertram, C. C. Goodnow, V. M. Dixit, Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. *Nature* **526**, 666-671 (2015).
6. J. Shi, Y. Zhao, K. Wang, X. Shi, Y. Wang, H. Huang, Y. Zhuang, T. Cai, F. Wang, F. Shao, Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature* **526**, 660-665 (2015).
7. J. Ding, K. Wang, W. Liu, Y. She, Q. Sun, J. Shi, H. Sun, D. C. Wang, F. Shao, Pore-forming activity and structural autoinhibition of the gasdermin family. *Nature* **535**, 111-116 (2016).
8. W. T. He, H. Wan, L. Hu, P. Chen, X. Wang, Z. Huang, Z. H. Yang, C. Q. Zhong, J. Han, Gasdermin D is an executor of pyroptosis and required for interleukin-1beta secretion. *Cell Res* **25**, 1285-1298 (2015).
9. X. Liu, Z. Zhang, J. Ruan, Y. Pan, V. G. Magupalli, H. Wu, J. Lieberman, Inflammasome-activated gasdermin D causes pyroptosis by forming membrane pores. *Nature* **535**, 153-158 (2016).
10. N. Kayagaki, S. Warming, M. Lamkanfi, L. Vande Walle, S. Louie, J. Dong, K. Newton, Y. Qu, J. Liu, S. Heldens, J. Zhang, W. P. Lee, M. Roose-Girma, V. M. Dixit, Non-canonical inflammasome activation targets caspase-11. *Nature* **479**, 117-121 (2011).
11. N. Kayagaki, M. T. Wong, I. B. Stowe, S. R. Ramani, L. C. Gonzalez, S. Akashi-Takamura, K. Miyake, J. Zhang, W. P. Lee, A. Muszynski, L. S. Forsberg, R. W. Carlson, V. M. Dixit, Noncanonical inflammasome activation by intracellular LPS independent of TLR4. *Science* **341**, 1246-1249 (2013).
12. J. A. Hagar, D. A. Powell, Y. Aachoui, R. K. Ernst, E. A. Miao, Cytoplasmic LPS activates caspase-11: implications in TLR4-independent endotoxic shock. *Science* **341**, 1250-1253 (2013).
13. J. Shi, Y. Zhao, Y. Wang, W. Gao, J. Ding, P. Li, L. Hu, F. Shao, Inflammatory caspases are innate immune receptors for intracellular LPS. *Nature* **514**, 187-192 (2014).
14. B. C. Russo, L. M. Stamm, M. Raaben, C. M. Kim, E. Kahoud, L. R. Robinson, S. Bose, A. L. Queiroz, B. B. Herrera, L. A. Baxt, N. Mor-Vaknin, Y. Fu, G. Molina, D. M. Markovitz, S. P. Whelan, M. B. Goldberg, Intermediate filaments enable pathogen docking to trigger type 3 effector translocation. *Nat Microbiol* **1**, 16025 (2016).
15. K. Ellwanger, E. Becker, I. Kienes, A. Sowa, Y. Postma, Y. Cardona Gloria, A. N. R. Weber, T. A. Kufer, The NLR family pyrin domain-containing 11 protein contributes to the regulation of inflammatory signaling. *The Journal of biological chemistry* **293**, 2701-2710 (2018).
16. C. Wu, Z. Su, M. Lin, J. Ou, W. Zhao, J. Cui, R. F. Wang, NLRP11 attenuates Toll-like receptor signalling by targeting TRAF6 for degradation via the ubiquitin ligase RNF19A. *Nat Commun* **8**, 1977 (2017).
17. Y. Qin, Z. Su, Y. Wu, C. Wu, S. Jin, W. Xie, W. Jiang, R. Zhou, J. Cui, NLRP11 disrupts MAVS signalosome to inhibit type I interferon signaling and virus-induced apoptosis. *EMBO Rep* **18**, 2160-2171 (2017).
18. T. Sanada, M. Kim, H. Mimuro, M. Suzuki, M. Ogawa, A. Oyama, H. Ashida, T. Kobayashi, T. Koyama, S. Nagai, Y. Shibata, J. Gohda, J. Inoue, T. Mizushima, C. Sasakawa, The Shigella flexneri effector OspI deamidates UBC13 to dampen the inflammatory response. *Nature* **483**, 623-626 (2012).
19. E. Meunier, M. S. Dick, R. F. Dreier, N. Schurmann, D. Kenzelmann Broz, S. Warming, M. Roose-Girma, D. Bumann, N. Kayagaki, K. Takeda, M. Yamamoto, P. Broz, Caspase-11 activation requires lysis of pathogen-containing vacuoles by IFN-induced GTPases. *Nature* **509**, 366-370 (2014).
20. C. R. Beuzon, S. Meresse, K. E. Unsworth, J. Ruiz-Albert, S. Garvis, S. R. Waterman, T. A. Ryder, E. Boucrot, D. W. Holden, Salmonella maintains the integrity of its intracellular vacuole through the action of SifA. *The EMBO journal* **19**, 3235-3249 (2000).

21.   P. J. Baker, D. Boucher, D. Bierschenk, C. Tebartz, P. G. Whitney, D. B. D'Silva, M. C. Tanzer, M. Monteleone, A. A. Robertson, M. A. Cooper, S. Alvarez-Diaz, M. J. Herold, S. Bedoui, K. Schroder, S. L. Masters, NLRP3 inflammasome activation downstream of cytoplasmic LPS recognition by both caspase-4 and caspase-5. *Eur J Immunol* **45**, 2918-2926 (2015).

22.   N. J. Bitto, P. J. Baker, J. K. Dowling, G. Wray-McCann, A. De Paoli, L. S. Tran, P. L. Leung, K. J. Stacey, A. Mansell, S. L. Masters, R. L. Ferrero, Membrane vesicles from Pseudomonas aeruginosa activate the noncanonical inflammasome through caspase-5 in human monocytes. *Immunol Cell Biol* **96**, 1120-1130 (2018).

23.   D. M. Pilla, J. A. Hagar, A. K. Haldar, A. K. Mason, D. Degrandi, K. Pfeffer, R. K. Ernst, M. Yamamoto, E. A. Miao, J. Coers, Guanylate binding proteins promote caspase-11-dependent pyroptosis in response to cytoplasmic LPS. *Proc Natl Acad Sci U S A* **111**, 6046-6051 (2014).

24.   D. Fisch, H. Bando, B. Clough, V. Hornung, M. Yamamoto, A. R. Shenoy, E. M. Frickel, Human GBP1 is a microbe-specific gatekeeper of macrophage apoptosis and pyroptosis. *The EMBO journal* **38**, e100926 (2019).

25.   J. C. Santos, M. S. Dick, B. Lagrange, D. Degrandi, K. Pfeffer, M. Yamamoto, E. Meunier, P. Pelczar, T. Henry, P. Broz, LPS targets host guanylate-binding proteins to the bacterial outer membrane for non-canonical inflammasome activation. *The EMBO journal* **37**, (2018).

26.   M. P. Wandel, B. H. Kim, E. S. Park, K. B. Boyle, K. Nayak, B. Lagrange, A. Herod, T. Henry, M. Zilbauer, J. Rohde, J. D. MacMicking, F. Randow, Guanylate-binding proteins convert cytosolic bacteria into caspase-4 signaling platforms. *Nat Immunol* **21**, 880-891 (2020).

27.   J. C. Santos, D. Boucher, L. K. Schneider, B. Demarco, M. Dilucca, K. Shkarina, R. Heilig, K. W. Chen, R. Y. H. Lim, P. Broz, Human GBP1 binds LPS to initiate assembly of a caspase-4 activating platform on cytosolic bacteria. *Nat Commun* **11**, 3276 (2020).

28.   B. H. Kim, A. R. Shenoy, P. Kumar, C. J. Bradfield, J. D. MacMicking, IFN-inducible GTPases in host cell defense. *Cell host & microbe* **12**, 432-444 (2012).

29.   M. Kutsch, L. Sistemich, C. F. Lesser, M. B. Goldberg, C. Herrmann, J. Coers, Direct binding of polymeric GBP1 to LPS disrupts bacterial cell envelope functions. *The EMBO journal* **39**, e104926 (2020).

30.   J. L. Poyet, S. M. Srinivasula, M. Tnani, M. Razmara, T. Fernandes-Alnemri, E. S. Alnemri, Identification of Ipaf, a human caspase-1-activating protein related to Apaf-1. *The Journal of biological chemistry* **276**, 28309-28313 (2001).

31.   C. Pop, J. Timmer, S. Sperandio, G. S. Salvesen, The apoptosome activates caspase-9 by dimerization. *Mol Cell* **22**, 269-275 (2006).

32.   R. A. MacCorkle, K. W. Freeman, D. M. Spencer, Synthetic activation of caspases: artificial death switches. *Proc Natl Acad Sci U S A* **95**, 3655-3660 (1998).

33.   B. Faustin, L. Lartigue, J. M. Bruey, F. Luciano, E. Sergienko, B. Bailly-Maitre, N. Volkmann, D. Hanein, I. Rouiller, J. C. Reed, Reconstituted NALP1 inflammasome reveals two-step mechanism of caspase-1 activation. *Mol Cell* **25**, 713-724 (2007).

34.   B. C. Baliga, S. H. Read, S. Kumar, The biochemical mechanism of caspase-2 activation. *Cell Death Differ* **11**, 1234-1241 (2004).

35.   P. Karki, G. R. Dahal, I. S. Park, Both dimerization and interdomain processing are essential for caspase-4 activation. *Biochemical and biophysical research communications* **356**, 1056-1061 (2007).

36.   K. Wang, Q. Sun, X. Zhong, M. Zeng, H. Zeng, X. Shi, Z. Li, Y. Wang, Q. Zhao, F. Shao, J. Ding, Structural Mechanism for GSDMD Targeting by Autoprocessed Caspases in Pyroptosis. *Cell* **180**, 941-955 e920 (2020).

37.   Q. Yin, H. H. Park, J. Y. Chung, S. C. Lin, Y. C. Lo, L. S. da Graca, X. Jiang, H. Wu, Caspase-9 holoenzyme is a specific and optimal procaspase-3 processing machine. *Mol Cell* **22**, 259-268 (2006).

38.   H. Sasaki, S. H. White, Aggregation behavior of an ultra-pure lipopolysaccharide that stimulates TLR-4 receptors. *Biophys J* **95**, 986-993 (2008).

39.   M. A. Wacker, A. Teghanemt, J. P. Weiss, J. H. Barker, High-affinity caspase-4 binding to LPS presented as high molecular mass aggregates or in outer membrane vesicles. *Innate Immun* **23**, 336-344 (2017).

40.   A. M. Taveira da Silva, H. C. Kaulbach, F. S. Chuidian, D. R. Lambert, A. F. Suffredini, R. L. Danner, Brief report: shock and multiple-organ dysfunction after self-administration of Salmonella endotoxin. *N Engl J Med* **328**, 1457-1460 (1993).

41.   R. W. Schaedler, R. J. Dubos, The susceptibility of mice to bacterial endotoxins. *J Exp Med* **113**, 559-570 (1961).

42. E. H. Labrec, H. Schneider, T. J. Magnani, S. B. Formal, Epithelial Cell Penetration as an Essential Step in the Pathogenesis of Bacillary Dysentery. *J Bacteriol* **88**, 1503-1518 (1964).

43. H. J. Wing, A. W. Yan, S. R. Goldman, M. B. Goldberg, Regulation of IcsP, the outer membrane protease of the Shigella actin tail assembly protein IcsA, by virulence plasmid regulators VirF and VirB. *J Bacteriol* **186**, 699-705 (2004).

44. A. T. Maurelli, B. Blackmon, R. Curtiss, 3rd, Loss of pigmentation in Shigella flexneri 2a is correlated with loss of virulence and virulence-associated plasmid. *Infect Immun* **43**, 397-401 (1984).

45. K. A. Miller, A. C. Garza-Mayers, Y. Leung, M. B. Goldberg, Identification of interactions among host and bacterial proteins and evaluation of their role early during Shigella flexneri infection. *Microbiology (Reading)* **164**, 540-550 (2018).

46. V. Novem, G. Shui, D. Wang, A. K. Bendt, S. H. Sim, Y. Liu, T. W. Thong, S. P. Sivalingam, E. E. Ooi, M. R. Wenk, G. Tan, Structural and biological diversity of lipopolysaccharides from Burkholderia pseudomallei and Burkholderia thailandensis. *Clin Vaccine Immunol* **16**, 1420-1428 (2009).

47. M. M. Gedde, D. E. Higgins, L. G. Tilney, D. A. Portnoy, Role of listeriolysin O in cell-to-cell spread of Listeria monocytogenes. *Infect Immun* **68**, 999-1003 (2000).

48. K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640-6645 (2000).

49. E. K. Brinkman, T. Chen, M. Amendola, B. van Steensel, Easy quantitative assessment of genome editing by sequence trace decomposition. *Nucleic Acids Res* **42**, e168 (2014).

50. S. Ranjbar, L. D. Jasenosky, N. Chow, A. E. Goldfeld, Regulation of Mycobacterium tuberculosis-dependent HIV-1 transcription reveals a new role for NFAT5 in the toll-like receptor pathway. *PLoS Pathog* **8**, e1002620 (2012).

51. T. Kobayashi, M. Ogawa, T. Sanada, H. Mimuro, M. Kim, H. Ashida, R. Akakura, M. Yoshida, M. Kawalec, J. M. Reichhart, T. Mizushima, C. Sasakawa, The Shigella OspC3 effector inhibits caspase-4, antagonizes inflammatory cell death, and promotes epithelial infection. *Cell host & microbe* **13**, 570-583 (2013).

52. Z. Li, W. Liu, J. Fu, S. Cheng, Y. Xu, Z. Wang, X. Liu, X. Shi, Y. Liu, X. Qi, X. Liu, J. Ding, F. Shao, Shigella evades pyroptosis by arginine ADP-riboxanation of caspase-11. *Nature* **599**, 290-295 (2021).

Acknowledgments: We thank Jonathan Kagan, Hao Wu, Kate Fitzgerald, Sunny Shin, Ann Goldfeld, Margaret Kielian, Peggy Cotter, and Jeffrey Miller for reagents.

Funding:

Real Colegio Complutense Fellowship, Real Colegio Complutense at Harvard (MLGM)

National Institutes of Health grant R01 AI081724 (MBG)

National Institutes of Health grant F32 AI145128 (ASZ)

National Institutes of Health grant F32 AI126765 (KAM)

National Institutes of Health grant T32 AI007061 (ASZ)

Author contributions

Conceptualization: MBG

Methodology: MRL, ASZ, TEW, MBG

Investigation: MRL, ASZ, TEW, KAM, MLGM, ACH, VK, KTE

Funding acquisition: ASZ, KAM, MLGM, MBG

Supervision: MRL, KAM, MBG

Writing – original draft: MBG

Writing – review & editing: MRL, ASZ, TEW, KAM, MLGM, MBG

Competing interests: The authors declare no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials. All materials are available by request to MBG, restricted by institutional MTAs.

Fig. 1. *NLRP11* **is required for efficient cell death induced by infection of human-derived macrophage-like cells with intracellular gram-negative bacteria.**

(**A**) Prevailing (blue/black arrows) and proposed (pink/black arrows) paradigms of non-canonical recognition of cLPS, high cLPS concentration activating both pathways. CASP4, caspase-4; GSDMD, gasdermin D; GSDMD-N, GSDMD N-terminal domain. (**B**-**C**) *NLRP11*-dependent death of THP-1 macrophages upon infection with cytosolic *S. flexneri* (**B**) or *B. thailandensis* (**C**), or predominantly vacuolar *S.* Typhimurium (**D**), in two independent (**B**) or one (**C**-**D**) *NLRP11*$^{-/-}$ THP-1 cell lines. $n = 5$ (**B**: WT, KO #1) or 3 experiments (**B**: KO #2, **C**-**D**). *S.* Typhimurium *sifA* is largely cytosolic. $n = 3$ experiments. Cell death, by LDH release, as percent of Triton X-100 cell lysis. Mean ± S.E.M. Two-way ANOVA with Dunnett *post hoc* test (**B**, **D**), or student T-test (**C**). *, $p<0.05$; ***; $p<0.001$; ****, $p<0.0001$.

Fig. 2. *NLRP11* potentiates macrophage cell death in response to cytosolic LPS.

(**A**) *NLRP11*-dependent death of THP-1 macrophage-like cells upon infection with cytosolic gram-positive pathogen *L. monocytogenes* coated with *S. enterica* serovar Minnesota LPS. Cell death depends on LPS and vacuolar escape, as *L. monocytogenes LLO⁻* is restricted to the phagosome. $n = 4$ experiments. (**B-C**) In *NLRP11⁻ᐟ⁻* THP-1 macrophages (**B**) or following NLRP11 siRNA (**C**), cell death in response to electroporated LPS is defective. $n = 3$ (**B**) or 4 (**C**) experiments. Cell death measured as in Fig 1. Tfxn, transfection; mock, carrier alone; scRNA, scrambled siRNA. Two-way ANOVA with Sidak *post hoc* test (**A**) or Tukey *post hoc* test (**B-C**). Mean ± S.E.M. **, $p<0.01$; ***; $p<0.001$; ****, $p<0.0001$.

Fig. 3. *NLRP11* **is required for processing of caspase-4 and gasdermin D during** *S. flexneri* **infection.**

S. flexneri infection of THP-1 macrophages. (**A**) Processing of caspase-4 (CASP4), with release of p32 polypeptide. (**B**) Processing of gasdermin D (GSDMD), with release of p30 polypeptide. Western blot of culture supernatants (top) and corresponding band densitometry (bottom); in **B**, densitometry is normalized to cell death upon wildtype (WT) *S. flexneri* infection of NLRP11[+/+] THP-1 macrophages. In the absence of OspC3, processing was similar to induced by WT *S. flexneri*; in mouse bone marrow-derived macrophages and human epithelial cell lines, OspC3 inhibits caspase-4 processing of gasdermin D (*51, 52*). $n = 3$ experiments. MW markers in kD. Mean ± S.E.M. Two-way ANOVA with Sidak *post hoc* test. *, $p < 0.05$; **, $p < 0.01$; ***; $p < 0.001$; ****, $p < 0.0001$.

Fig. 4. NLRP11-dependent cell death in response to cLPS is dependent on caspase-4 but independent of caspase-1 and MyD88.

Cell death of THP-1 macrophages in response to electroporated LPS following siRNA depletion of *CASP4*, *CASP1*, or *MyD88*. *n* = 5 experiments. Cell death measured as in Fig 1; normalized to mock transfection of *NLRP11*[+/+] cells following scRNA treatment. Tfxn, transfection; mock, carrier alone; scRNA, scrambled siRNA; CASP4, caspase-4; CASP1, caspase-1. Two-way ANOVA with Tukey *post hoc* test. Mean ± S.E.M. **, p<0.01; ****, p<0.0001; n.s., not significant.

Fig. 5. NLRP11 interactions with LPS and caspase-4

Precipitations from reconstituted HEK293T lysates expressing NLRP11 and/or caspase-4. (**A-B**)

Precipitation of NLRP11 and wildtype caspase-4 (CASP4) by biotin-LPS, separately (**A**) and

together (**B**), and not biotin-PAM3CSK4 (Pam3) (**A**). PD: SA, precipitation with streptavidin

beads. $n = 3$ experiments. (**C**) NLRP11 precipitates wildtype CASP4. $n = 3$ experiments. (**D**)

Increasing interaction between NLRP11 and CASP4 as a function of concentration of each

protein. FLAG-NLRP11 in transfected HEK293T cells bound to plates coated with purified

wildtype CASP4; detection of bound FLAG. NLRP11 lysate dilutions: circles, undiluted;

squares, 1:2 dilution; triangles, 1:4 dilution; inverted triangles, 1:8 dilution. Diamonds,

untransfected (untfxd) lysate. $n = 3$ experiments. a.u., arbitrary units.

Fig. 6. Specificity of NLRP11 for caspases

Precipitations from reconstituted HEK293T lysates expressing NLRP11 and indicated caspases.

FLAG-tagged inactive caspase (CASP1[C285A]-FLAG, CASP5a[C315A]-FLAG,

CASP9[C287S]-FLAG) precipitation of Myc-NLRP11. CASP*, catalytically inactive caspase.

IP: FLAG, precipitation with FLAG antibody-coated beads. Western blot detection with anti-

FLAG or anti-Myc antibody. $n = 3$ experiments.

Fig. 7. Requirement for NLRP11 for cell death in response to cLPS in primary human macrophages.

NLRP11-dependent cell death in response to electroporated LPS in isolated primary human monocyte-derived macrophages (CD14+), treated with siRNA to *NLRP11* (**A**), *CASP4* (**B**), or *MyD88* (**C**). Each experiment performed with cells from a different donor. Cell death measured as in Fig 1; normalized to mock transfection following scRNA treatment. $n = 4$ (**A-B**) or 3 (**C**) experiments. Tfxn, transfection; mock, carrier alone; scRNA, scrambled siRNA; CASP4, caspase-4. Two-way ANOVA with mixed-effects analyses. Mean ± S.E.M. ***; p<0.001; n.s., not significant.