Title:

Cytosolic Sensor of Bacterial Lipopolysaccharide in Human Macrophages

Authors:

Maricarmen Rojas-Lopez[1]*, Amanda S. Zajac[1]*, Thomas E. Wood[1]*, Kelly A. Miller[1]*,

María Luisa Gil-Marqués[1], Austin C. Hachey[1], Vritti Kharbanda[1], Marcia B.

Goldberg[1,2,3,4]†

Affiliations:

[1]Center for Bacterial Pathogenesis, Division of Infectious Diseases, Department of

Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA,

USA.

[2]Department of Microbiology, Blavatnik Institute, Harvard Medical School, Boston, MA,

USA.

[3]Broad Institute of MIT and Harvard, Cambridge, MA, USA.

[4]Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of

Public Health, Boston, MA, USA.

*These authors contributed equally to this work.

†Correspondence should be addressed to M.B.G. (marcia.goldberg@mgh.harvard.edu),

ORCID number: 0000-0003-4266-9733

Summary paragraph

Inflammasomes are cytosolic supramolecular organizing centers that, in response to

pathogen-derived molecules and endogenous danger signals, assemble and activate

innate immune responses. Bacterial lipopolysaccharide (LPS) is an inflammasome

trigger and a major mediator of inflammation during infection, including during the

potentially lethal condition sepsis. Activation of most inflammasomes is triggered by

sensing of pathogen products by a specific host cytosolic nucleotide-binding

oligomerization domain, leucine rich repeat and pyrin domain containing protein (NLRP)

or other sensor protein that in turn activates a pro-inflammatory caspase. LPS that

accesses the cell cytosol (cLPS) induces cell-autonomous activation of a non-canonical

inflammasome that contains caspases-4/5 in humans or caspase-11 in mice[1-3].

Whereas the NLRPs that sense most pathogen triggers have been identified, no NLRP

is known to sense cLPS, which together with the observation that caspases-4, -5, and -

11 bind LPS *in vitro*[4], has led to the postulate that inflammasome activation by cLPS

occurs independent of an NLRP. Here we show that primate-specific NLRP11 senses

cLPS and promotes the activation of caspase-4. We found that in response to infection

by each of several gram-negative intracellular bacterial pathogens or to LPS

transfection, efficient activation of the non-canonical pathway in human-derived

macrophages depends on NLRP11. Further, we found that in both immortalized human-

derived macrophages and primary human macrophages, the dependence of the non-

canonical pathway on NLRP11 is due to detection of cLPS. Moreover, in cell lysates,

NLRP11 binds LPS independently of caspase-4 and binds caspase-4 independently of

LPS. Our results demonstrate that NLRP11 senses cLPS and promotes LPS-dependent

activation of caspase-4. NLRP11 is a previously missing link in the human non-

canonical inflammasome activation pathway.

Inflammasomes assemble in response to the presence in the cytosol of pathogen-

associated molecules, including dsDNA, flagellin, type 3 secretion system structural

proteins, pore-forming toxins, ion flux, ATP, and LPS[5]. Most inflammasomes consist of a

sensor protein and an inflammatory caspase, frequently linked by an adaptor protein.

Inflammasome assembly activates inflammatory caspases, which proteolytically

process the pore-forming protein gasdermin D[6-8], including caspase-1-dependent

proteolytically processing of the pro-inflammatory cytokines IL-18 and IL-1β[9,10]. Pores

formed in the plasma membrane by the N-terminal domain of gasdermin D enable the

release of activated pro-inflammatory cytokines and trigger pyroptotic cell death[6-8,11].

The presence of bacterial LPS in the cytosol of human macrophages activates a non-

canonical inflammasome, which in humans, contains caspases-4 and -5, and in mice,

caspase-11[1-3]. To date, no sensor protein of cytosolic LPS has been identified, and

because caspase-4, -5, and -11 bind LPS *in vitro*[4], the current paradigm is that the non-

canonical inflammasome lacks a sensor protein.

NLRP11 required for cell death due to intracellular gram-negative pathogens

In a forward genetic screen that enriched for cellular genes required for efficient cell

death of a human-derived myeloid lineage cell line by the gram-negative intracellular

bacterial pathogen *Shigella flexneri*, we identified *NLRP11*[12]. We performed targeted

deletion of *nlrp11* in the human-derived macrophage cell line THP-1 and found that cell

death during *S. flexneri* infection (as measured by lactate dehydrogenase (LDH)

release) was significantly reduced for cell lines that lack NLRP11. In each of two THP-1-

derived cell lines in which *nlrp11* was disrupted by a CRISPR-Cas9 induced indel

(Extended Data Fig. 1), release of LDH upon infection with *S. flexneri* was reduced by

more than 50% (Fig. 1a). Consistent with previous reports, *S. flexneri*-induced

macrophage cell death was dependent on the bacterial type 3 secretion system, which

mediates both bacterial uptake[13] and vacuolar escape[14,15], as a mutant that lacked

either the virulence plasmid that encodes this secretion system or *virB*, the

transcriptional activator of this system, displayed minimal cell death (Extended Data Fig.

2a).

To test whether the dependence on NLRP11 for efficient cell death was generalizable to

other intracellular gram-negative bacterial pathogens, we infected *NLRP11*[-/-] cells with

the cytosolic pathogen *Burkholderia thailandensis* and the intracellular but largely

vacuolar pathogen *Salmonella enterica* serovar Typhimurium. As with *S. flexneri*

infection, in the absence of NLRP11, cell death during infection with *B. thailandensis* or

S. Typhimurium was significantly reduced (Fig. 1b-c). In response to viral infection and

certain Toll-like receptor interactions, the NF-κB pathway is repressed by NLRP11-

dependent targeted degradation of TRAF6[16-18]; however, *S. flexneri*-induced NLRP11-

dependent cell death was independent of TRAF6, whether or not the *Shigella* effector

OspI, which suppresses TRAF6 signaling[19], was present (Extended Data Fig. 2b-d),

indicating that the role of NLRP11 in bacterial infection-induced cell death is distinct

from its role in NF-κB signaling. Together, these results indicate that NLRP11 is

required for efficient cell death induced by multiple intracellular gram-negative

pathogens and raised the possibility that the implicated pathogen-associated trigger of

cell death is generally present among gram-negative bacteria and depends on

intracellular presentation.

NLRP11 recognizes cLPS

We tested whether LPS, which is specific to the cell envelope of gram-negative
bacteria, might be required for NLRP11-dependent cell death. Infection of THP-1
macrophages with the cytosolic gram-positive pathogen *Listeria monocytogenes*
induced minimal levels of cell death during the timeframe of the experiment, but when
coated with *S. enterica* LPS, induced cell death that was largely dependent on NLRP11
(Fig. 2a). The observed cell death was dependent on cytosolic localization, as a *L.*
monocytogenes strain lacking listeriolysin O (LLO), which is required to lyse the
vacuole[20], induced minimal cell death. Introduction of LPS alone directly into the cytosol
by electroporation similarly induced cell death that was significantly dependent on
NLRP11, both in *NLRP11*[-/-] and NLRP11 siRNA-treated THP-1 macrophages (Fig. 2b-
c), whereas cell death induced by extracellular LPS was independent of NLRP11
(Extended Data Fig. 3a). These data indicate that NLRP11 functions in a cell death
pathway that is activated by LPS localized to the cytosol (cLPS). For pathogens such as
Salmonella that do not themselves efficiently lyse the vacuole, delivery of LPS to the
cytosol occurs upon guanylate-binding protein-2 (GBP2)-mediated vacuolar lysis[21], a
likely explanation for the lack of differential NLRP11-dependent cell death induced by
wild-type *S.* Typhimurium, which are largely vacuolar, and *sifA* mutant *S.* Typhimurium,
which is predominantly cytosolic[22] (Fig. 1c).

NLRP11 required for efficient activation of non-canonical inflammasome

We tested whether NLRP11 was required for activation of the non-canonical
inflammasome, which in humans, contains caspases-4 and -5. Activation of caspase-4

and the downstream pore-forming protein gasdermin D were dependent on NLRP11;

during *S. flexneri* infection, processing of caspase-4 and gasdermin D, leading to the

release of active domains of each, was significantly reduced in the absence of NLRP11

(Fig. 1d-e). In THP-1 macrophages depleted of caspase-4, the absence of NLRP11

caused no significant additional decrease in cell death induced by LPS transfection (Fig.

2d and Extended Data Fig. 4a). Depletion of *CASP1* was associated with no significant

decrease in cell death in response to cLPS as compared to control depletion for either

THP-1 macrophages containing NLRP11 or THP-1 macrophages lacking NLRP11 (Fig.

2d and Extended Data Fig. 4b). Despite testing several antibodies, we were unable to

detect native cellular NLRP11 by western blot, presumably due to very low levels of

native protein. These data indicate that NLRP11 is required for efficient activation of the

non-canonical inflammasome, likely functioning in the caspase-4 pathway, and that

NLRP11 function is independent of caspase-1. Consistent with NLRP11 functioning

specifically in the non-canonical inflammasome pathway, NLRP11 was not required for

activation of the canonical NLRP3 inflammasome, as cell death induced by the NLRP3

agonist nigericin was independent of NLRP11 (Extended Data Fig. 3b-d).

NLRP11 interacts with LPS and caspase-4

In contrast to NLRP3 and NLR family CARD domain containing 4 (NLRC4)

inflammasomes, activation of the non-canonical pathway is independent of the adaptor

protein ASC, as signaling in response to cytosolic LPS is normal in human or mouse-

derived macrophages lacking ASC[2,4] and NLRP11 does not co-localize with ASC

specks[16,23]. We tested whether NLRP11 interacts with other core components of the

non-canonical pathway by ectopic expression of selected components in HEK293T

cells, which lack most innate immune proteins. Biotin-conjugated LPS precipitated

NLRP11 in the absence of caspase-4 (Fig. 3a), indicating that NLRP11 binds LPS

independently of caspase-4. As previously described[4], LPS also specifically precipitated

caspase-4 and its catalytically inactive derivative (CASP4 C258A), including in the

presence of NLRP11 (Fig. 3b), whereas the triacylated lipid Pam3CSK4 did not

precipitate either NLRP11 or, as previously described[4], caspase-4 (Fig. 3a). In addition

to binding to LPS independent of caspase-4, NLRP11 specifically precipitated caspase-

4 (Fig. 3c). To characterize the binding relationship of NLRP11 to caspase-4, we

developed a plate binding assay using FLAG-NLRP11 in HEK293T cell lysates and

purified caspase-4 (Methods and Extended Data Fig. 5a). In this assay, FLAG-NLRP11

binding depended on caspase-4, as binding of FLAG-NLRP11 in the absence of

caspase-4 was at background levels (Fig. 3d, diamonds). Moreover, binding of NLRP11

to caspase-4 increased as a function of increasing concentrations of both NLRP11

lysate and caspase-4, resembling the asymptotic phase of a sigmoidal curve, beginning

to reach saturation at 8-10 µg/ml caspase-4 coating the wells (Fig. 3d).

NLRP11 displays caspase specificity

In addition to caspase-4, NLRP11 co-precipitated with caspase-1; of note, NLRP11 did

not co-precipitate with caspase-5, a paralog of caspase-4 that binds LPS and is a

component of at least some non-canonical inflammasomes [4,10,24], or the more divergent

caspase-9 (Fig. 3e and Extended Data Fig. 6a). Although caspase-1 co-precipitates

with NLRP11 (Fig. 3e), in THP-1 macrophages, NLRP11 function in LPS-induced cell

death is independent of caspase-1 (Fig. 2d), indicating that the observed interaction of

caspase-1 with NLRP11 does not functionally contribute to activation of the non-

canonical inflammasome.

NLRP11 role in non-canonical inflammasome activation is independent of

interferon-gamma and guanylate-binding protein-1

Host guanylate-binding proteins (GBPs) bind intracellular LPS, promote non-canonical

inflammasome activation[25-29], and are encoded by genes that are strongly induced by

interferon-gamma (IFNγ)[30,31]. Human GBP1 binds LPS on the *S. flexneri* surface,

whereupon it enhances caspase-4 recruitment[32]. We tested whether IFNγ stimulation

altered the role of NLRP11 in activation of the non-canonical inflammasome. IFNγ pre-

treatment of THP-1 macrophages induced GBP1, as expected, yet NLRP11 was still

required for efficient activation of caspase-4 and cell death (Extended Data Fig. 7a-c),

indicating that IFNγ and GBP1 do not bypass the requirement for NLRP11 in non-

canonical inflammasome induced pyroptosis.

Requirement for NLRP11 in primary human macrophages

Following treatment of primary human monocyte-derived macrophages with siRNA to

NLRP11, although overall levels of cell death are lower than for immortalized THP-1

cells, cell death induced by cLPS was significantly diminished (Fig. 2e). As in THP-1

macrophages, in primary macrophages, the defect in cLPS-induced cell death

depended on caspase-4, but not MyD88 (Figs. 2f-g and Extended Data Fig. 4c-d),

indicating that in these assays, the trigger was cLPS rather than extracellular LPS, and

demonstrating that the requirement for NLRP11 in the non-canonical inflammasome

pathway extends to primary human cells.

In summary, the data reported here identify NLRP11 as a primate-specific cytosolic

sensor of bacterial LPS in the activation of the caspase-4 dependent inflammasome and

caspase-4 dependent pyroptosis. Galectin-3, one among a family of β-galactoside–

binding proteins, also senses cytosolic pathogens, promotes delivery of GBPs to

ruptured pathogen-containing vacuoles, and enhances assembly of LPS-triggered

caspase-4/11 oligomerization and activation[33-35]. Whether galectin-3 activity synergizes

with the interaction of LPS and NLRP11 is uncertain; however, the independence of

NLRP11 function from GBP1 (Extended Data Fig. 7) suggests that any contribution of

galectin-3 to this process would be minor. Additionally, it remains uncertain whether the

unique presence of NLRP11 in primates and the absence of it or identifiable homologs

in mice contributes to the observed $\sim 10^4$-fold increased susceptibility to gram-negative

bacterial sepsis and LPS of humans as compared to mice[36-42]; additional insights into

these relationships would potentially enable much-needed improved mouse models for

bacterial sepsis.

Methods

Bacterial strains, plasmids, antibodies, and reagents. *Shigella flexneri* serotype 2a

wildtype strain 2457T and the *S. flexneri* mutant strains *virB*::Tn5, BS103, and Δ*ospC3*,

which are isogenic to 2457T, have been described previously[43-46]. *Salmonella enterica*

serovar Typhimurium and the isogenic Δ*sifA* mutant[22] were gifts from L. Knodler.

Burkholderia thailandensis[47] was a gift from P. Cotter. *Listeria monocytogenes* 10403s

and 10403 LLO[-48] were gifts from M. Kielian and J. Miller. *S. flexneri* was grown

overnight in tryptic soy broth at 37°C with aeration. *L. monocytogenes* was grown

overnight in brain heart infusion broth at 30°C at a stationary slant. *B. thailandensis* was

grown overnight in low salt Luria broth (LB) at 37°C with aeration. *S. enterica*

Typhimurium was grown in LB broth at 37°C with aeration.

Caspase-1 (Abcam, ab207802) rabbit monoclonal antibody was used at 0.5 μg/mL

(1:1000). Caspase-4 (Santa Cruz, sc-56056), and gasdermin D (Santa Cruz,

GSDMDC1 64-Y) mouse monoclonal antibodies were used at 0.5 μg/mL (1:200). FLAG

M2 mouse monoclonal antibody (Sigma, F3165) was used at 76 ng/mL (1:5000). Myc

rabbit monoclonal antibody (Cell Signaling, 2278S) was used at 1:1000. GBP1 mouse

monoclonal antibody (Abcam, ab119236) was used at 1:1000. For western blots of

immunoprecipitations, the secondary antibody used was TrueBlot horseradish

peroxidase-conjugated anti-mouse Ig (Rockland, 18-8817-31). For detecting β-actin,

horseradish peroxidase-conjugated anti-β-actin antibody (Sigma, A3854) was used. For

all other western blots, the secondary antibody used was horseradish peroxidase-

conjugated goat anti-rabbit or goat anti-mouse IgG antibodies (Jackson

ImmunoResearch, 115-035-003 or 111-035-144). All antibodies were diluted in 5% milk

in TBST (0.1% Tween).

Strain and plasmid construction. The *S. flexneri ospI* deletion mutant was generated

in the 2457T strain background using the lambda red method by introducing a

chloramphenicol acetyltransferase (*cat*) cassette into the *orf169b* gene (which encodes

OspI), as described[49].

pCMV-CASP4-MYC and pCMV-CASP5-MYC were made from pCMV-CASP4-MYC-

FLAG and pCMV-CASP5-MYC-FLAG, gifts from S. Shin. pCMV-CASP4-MYC C258A

and pCMV-CASP4-MYC K19E C258A was generated by splicing by overlap extension

PCR (SOE-PCR). pcDNA-NLRP11 (gift from H. Wu) was subcloned into pCMV-FLAG

with an N-terminal FLAG tag for FLAG-NLRP11 and an N-terminal MYC tag for MYC-

NLRP11. All FLAG tags are 3xFLAG. All N-terminal Myc tags are 6xMyc; all C-terminal

Myc tags are 1xMyc. All DNA used for transfection was prepared LPS-free (Macherey-

Nagel, 740410.50).

Generation of CRISPR knockouts. THP-1 cells were transduced with lentivirus

packaged with pLentiCas9-Blast (Addgene, 52962) generated from HEK293T cells.

THP-1 cell suspensions were treated with 8 μg/mL polybrene, combined with lentivirus

and centrifuged in 12-well plates for 2 h at 1000 x g at 30°C. After centrifugation, cells

were incubated for 2 h and washed once with fresh media. After 48 h, blasticidin

(10 μg/mL) was added to the transduced cells. Guide RNA sequence ("sgRNA3",

AGGCATGCAAAGCTGTCATG) was subcloned into pLentiGuide-Puro (Addgene,

59702). Guide RNA expressing plasmids were packaged into lentivirus using HEK293T

cells. Polybrene treated THP-1 cells were transduced and selected with 1 μg/mL

puromycin. Cells were limiting diluted and single cell clones were screened by

SURVEYOR (IDT, 706020) and Sanger sequencing coupled with TIDE analysis

(Brinkman et. al 2014).

Generation of *TRAF6* depletion cell lines. pLKO.1-shTRAF6 ([50], gift of A. Goldfeld),

which encodes shRNA to *TRAF6*, was packaged into lentivirus in HEK293T cells.

Lentivirus were introduced into NLRP11+/+ and NLRP11-/- THP-1 cell lines and cells

containing pLKO.1-shTRAF6 were selected, expanded by maintenance in 0.8 µg/mL

puromycin, as described[50], and stable clones were isolated by limiting dilution.

Mammalian cell culture. All cell lines were obtained from ATCC. HEK293T cells were

maintained in Dulbecco's Modified Eagle's medium (Gibco, 11965118). THP1 cells were

maintained in RPMI-1640 (Gibco, 21870092). All media was supplemented with 10%

(vol/vol) fetal bovine serum (heat inactivated, sterile filtered, R&D Systems). For THP-1

cells, RPMI-1640 media was also supplemented with 10 mM HEPES (Gibco,15630080)

and 2 mM L-glutamine (Gibco, 25030081). All cells were grown at 37$\degree$C in 5% CO_2. To

differentiate monocyte cell lines into macrophage-like cells, 50 ng/mL phorbol-12-

myristrate-13-acetate (PMA) was added 48 h before experiments. Where indicated,

cells were pretreated with 10 ng/mL IFNγ 16-20 h prior to infection.

Precipitation experiments. HEK293T cells were transiently transfected using Fugene6

(Promega). Stock solutions of 10x Buffer A (100 mM HEPES, 1500 mM NaCl, 10 mM

EGTA, 1 mM $MgCl_2$, pH 7.4) and 20% Triton X-100 were prepared as described[51]. The

cell culture medium was removed 24-48 h after transfection. Cells were washed with

cold PBS and lysed in 1x Buffer A, 0.5% Triton-X. Cell lysates were incubated on ice for

15 min, vortexed for 15 s and centrifuged at 13,200 rpm for 10 min at 4°C. The pellet

was discarded, and Bradford assay was used to determine protein concentration. 30 μL

of FLAG M2 Magnetic Beads (Millipore-Sigma, M8823) were washed twice in 1x Buffer

A, 0.1% Triton-X and equilibrated in the same buffer at 4°C for 2 h. 500 μg of protein

lysate was incubated with the beads for 2 h at 4°C. Beads were washed 3 times with 1x

Buffer A, 0.1% Triton-X and eluted with 50 μL of 2x Laemmli buffer.

Streptavidin pulldown was largely as described[4], except that 4 μg biotinylated *E. coli*

O111:B4 LPS was loaded onto 8 μg of Streptavidin Mag sepharose (GE, 28985738, or

Pierce, 88816). Cell lysates were diluted to 4 mg/mL before a 2 h incubation with beads.

For co-precipitation experiments, separate cells were transfected with each DNA

construct, and equivalent amounts of lysates were combined with each other or with

untransfected lysate, such that the amount of target protein and the total amount of

lysate were similar in each experimental condition; these lysates were then incubated at

37°C overnight. Lysates were cleared for non-specific binding with streptavidin beads

and then incubated for 2 h at 30°C with 4 μg biotinylated *E. coli* O111:B4 LPS loaded

onto Streptavidin Mag sepharose.

Plate binding assays. Clear flat bottom Nunc Maxisorp ELISA plates (Thermo Fisher,

442404) were coated with various concentrations of purified recombinant caspase-4

(Origene, TP760359) overnight at 4°C. The plates were washed 3x with PBS then

blocked for 2 h at RT with 1% bovine serum albumin (Fisher, BP9700100) in PBS.

Excess blocking solution was removed. HEK 293T cells transiently expressing FLAG-

NLRP11 and control untransfected cells were lysed into 50 mM Tris-HCl pH 7.4, 150

mM NaCl, 5 mM $MgCl_2$, 100 µg/ml digitonin (Thermo Fisher Scientific, BN2006) and

complete Mini EDTA-free protease inhibitor (Roche), as described [28]. Total protein

concentration of lysates was normalized to 2 mg/mL. Serial dilutions were in

untransfected HEK293T cell lysate. Varying concentrations of cell lysates were

incubated with pre-blocked ELISA plates for 1 h at RT. Excess lysate was removed by

washing 3 times with PBS. For detection of FLAG-NLRP11, wells were incubated with

0.8 µg/mL anti-FLAG rabbit (Sigma, F7425-.2MG) was incubated for 1 h at RT, and

after washing with PBS, 0.16 µg/mL goat-anti-rabbit conjugated to horseradish

peroxidase (HRP) (Jackson, 111-035-144). To detect caspase-4 bound to the plate,

wells were incubated with 0.5 µg/mL anti-caspase-4 (Santa Cruz, sc-56056) and, after

washing with PBS, 0.16 µg/mL goat-anti-mouse conjugated to HRP (Jackson, 115-035-

003), each for 1 hr. After 6 washes with PBS, 50 µL of One Step Ultra-TMB (Fisher

Scientific, PI34029) was added to each well and incubated for 5 min before stopping

with 4 N sulfuric acid and reading in a BioTek plate reader.

Bacterial infection. Overnight cultures were started from a single colony taken from a

freshly streaked plate. Infections with *S. flexneri* were performed with exponential phase

bacteria, after back-dilution from overnight cultures. Infections with *L. monocytogenes*

were performed with stationary phase bacteria from overnight cultures. Infections were

with a multiplicity of infection (MOI) of 10 for *S. flexneri* cultures and 1 for *L.*

monocytogenes. Bacteria were centrifuged onto cells at 2000 rpm for 10 min, followed

by incubation at 37°C. After 30 min, the cells were washed twice, and then 25 µg/mL

gentamicin was added to the media to kill extracellular bacteria. Where appropriate *S.*

enterica serovar Minnesota LPS was added to 10 μg/mL at the time of infection and

MCC950 to 1 μM 30 min prior to infection and maintained during infection. For LDH

assays of *S. flexneri* infected cells, 100 μL of cell culture supernatant was collected at 2

h and 30 min and at 5 h and/or 30 min of infection; for detection of caspases and/or

gasdermin D by western blot, 100 μL of cell culture supernatant was collected at the

same time points and was mixed with 35 μL 4x Laemmli buffer supplemented with

cOmplete EDTA-free protease inhibitors (Roche, 5056489001). For parallel analyses of

L. monocytogenes infected cells, gentamicin was used at 15 μg/mL, and samples were

collected at 6 h of infection. For western blot analyses, whole cell lysates were collected

in 2x Laemmli buffer.

Infections with *B. thailandensis* and *S.* Typhimurium were performed with exponential

phase bacteria at a multiplicity of infection (MOI) of 10. Bacteria were centrifuged onto

cells at 2000 rpm for 10 min, followed by incubation at 37°C. For *S.* Typhimurium, after

30 min incubation, the cells were washed twice, and then 25 μg/mL gentamicin was

added to the media to kill extracellular bacteria. For *B. thailandensis*, after 2 h

incubation, the cells were washed twice with cRPMI. For LDH assays with *S.*

Typhimurium infection, 100 μL of cell culture supernatant was collected at 2 h and 30

min and at 5 h and 30 min of infection and, for LDH assays with *B. thailandensis*, 100

μL of cell culture supernatant was collected at 24 h*.*

Isolation of primary human peripheral blood mononuclear cells (PBMCs) and

CD14 monocytes. Cells were isolated from leukopaks using density-gradient

centrifugation. Briefly, blood was diluted 1:2 with RPMI-1640 (Gibco), layered on top of

Ficoll-Paque Plus (GE Healthcare) using SepMate™-50 (IVD) (StemCell Technologies),

and centrifuged at 1,200g for 20 min. The PBMC layer was collected and resuspended

in 50 ml RPMI-1640 (Gibco) and centrifuged again at 300 g for 10 min. The cells were

counted, resuspended in Cryostor CS10 (StemCell Technologies), and aliquoted in

cryopreservation tubes at a concentration of 1×10^8 cells per mL. The tubes were

placed at −80°C overnight, then transferred to liquid nitrogen for long-term storage. To

isolate CD14+ cells, EasySep™ Human CD14 Positive Selection Kit II (StemCell

Technologies) was used, following the manufacturer's instructions. Cells were cultured

in complete RPMI, differentiated with 25 ng/mL m-CSF (StemCell Technologies), and

maintained at 37°C in 5% CO_2.

LPS transfection. Electroporation of cells with LPS was performed using the Neon

Transfection System (Invitrogen) according to the manufacturer's instructions. THP-1

cells were seeded in low-attachment 10 cm dishes with 50 ng/mL PMA in RPMI. After

48 h, the cells were lifted in PBS (without calcium or magnesium). For electroporation,

1.5×10^6 THP-1 cells were pelleted, resuspended in Resuspension Buffer R with 2

μg/mL of LPS or water control. Using a 10 μL Neon tip, cells were pulsed once at 1400

V and 10 ms pulse width. For primary human macrophage cells, 7.5×10^5 CD14

macrophages were pelleted, then resuspended in Resuspension Buffer T with 2 μg/mL

of LPS or water control. Using a 10 μL Neon tip, cells were pulsed 2 times at 2250 V

and 20 ms pulse width. Electroporated cells were immediately resuspended in 2 mL of

RPMI and placed into wells of a 24 well plate. Cells were incubated for 2.5 h at 37°C

before measuring LDH release.

siRNA transient knockdowns. The siRNAs used for NLRP11 knockdowns were

Hs_NALP11_1 FlexiTube siRNA and Hs_NALP11_5 FlexiTube siRNA at 10 nM total (5

nM each siRNA); for Caspase-1 Hs_CASP1_14 FlexiTube siRNA and Hs_CASP1_13

FlexiTube siRNA at 30 nM total (15 nM each siRNA); and for MyD88 Hs_MYD88_5

FlexiTube siRNA and Hs_MYD88_11 FlexiTube siRNA at 30 nM total (15 nM each

siRNA) (Qiagen). AllStars Negative Control siRNAs (Qiagen) were used as negative

controls at 10 nM or 30 nM, accordingly. The siRNAs used for caspase-4 knockdowns

were 3941, 4036 and 4130 at 30 nM total (10 nM each siRNA) (ThermoFisher). siRNAs

were transfected using the HiPerfect reagent (Qiagen) into primary human monocyte-

derived macrophages 72 h before or into THP-1 cells 48 h before experimental assays,

following the manufacturer's protocol for "Transfection of Differentiated Macrophage

Cell Lines." After transfection, cells were maintained at 37°C in 5% CO_2.

Nigericin. PMA differentiated THP-1 cells were washed once with fresh RPMI and

primed with 1 µM O55:B5 *E. coli* LPS for 3 h, at which time 10 µM nigericin or vehicle

control was added. LDH release was assayed at indicated times after treatment.

LDH assay. LDH was measured using an LDH Cytotoxicity Kit (Pierce, 88953) or

Cytotoxicity Detection Kit (Sigma, 11644793001) per the manufacturer's instructions.

Measurements at 490 nm and 680 nm were taken immediately after addition of Stop

Solution using a BioTek Plate Reader. To determine LDH activity, the 680 nm

absorbance value (background) was subtracted from the 490-nm absorbance value,

and cytotoxicity was calculated for each sample as LDH activity as a fraction of the total

LDH activity for cells treated with the Triton X-100 lysis buffer provided by the

manufacturer. For experiments involving primary cells, a different donor was used for

each biological replicate; due to inherent variability from donor to donor in the baseline

levels of death in cell death assays, within-donor data were normalized to the negative

control condition before compilation of the biological replicates.

Statistics and reproducibility. All results are data from at least 3 independent

biological replicates. Statistical analysis was performed with Prism 9.

Data availability. All relevant data are included in the manuscript and will be made

available by request to the corresponding author, marcia.goldberg@mgh.harvard.edu.

Supplementary information

378 The online version contains supplementary material.

Acknowledgements

380 We thank Jonathan Kagan, Hao Wu, Kate Fitzgerald, Sunny Shin, Ann Goldfeld,

381 Margaret Kielian, Peggy Cotter, and Jeffrey Miller for reagents.

382 This work was supported by a Real Colegio Complutense Fellowship from the Real

383 Colegio Complutense at Harvard (to M.L.G.M.) and NIH grants R01 AI081724 (to

384 M.B.G), F32 AI145128 (to A.S.Z.), and T32 AI007061 (to A.S.Z.).

Author contributions

386 M.B.G. conceived the study, supervised the project, and wrote the manuscript with input

387 from all authors; M.R.L. K.A.M., A.C.H., and V.K. performed cell-based experiments;

388 A.S.Z., T.E.W., and M.L.G.M. performed biochemical experiments. All authors

389 organized and analyzed data and edited the manuscript.

Competing interests

391 The authors declare no competing interests.

Materials and correspondence

393 Correspondence and material requests should be addressed to M.B.G.

References

1   Kayagaki, N. *et al.* Non-canonical inflammasome activation targets caspase-11.
Nature **479**, 117-121, doi:10.1038/nature10558 (2011).

2   Kayagaki, N. *et al.* Noncanonical inflammasome activation by intracellular LPS
independent of TLR4. *Science* **341**, 1246-1249, doi:10.1126/science.1240248
(2013).

3   Hagar, J. A., Powell, D. A., Aachoui, Y., Ernst, R. K. & Miao, E. A. Cytoplasmic
LPS activates caspase-11: implications in TLR4-independent endotoxic shock.
Science **341**, 1250-1253, doi:10.1126/science.1240988 (2013).

4   Shi, J. *et al.* Inflammatory caspases are innate immune receptors for intracellular
LPS. *Nature* **514**, 187-192, doi:10.1038/nature13683 (2014).

5   Brewer, S. M., Brubaker, S. W. & Monack, D. M. Host inflammasome defense
mechanisms and bacterial pathogen evasion strategies. *Curr Opin Immunol* **60**,
63-70, doi:10.1016/j.coi.2019.05.001 (2019).

6   Kayagaki, N. *et al.* Caspase-11 cleaves gasdermin D for non-canonical
inflammasome signalling. *Nature* **526**, 666-671, doi:10.1038/nature15541 (2015).

7   Shi, J. *et al.* Cleavage of GSDMD by inflammatory caspases determines
pyroptotic cell death. *Nature* **526**, 660-665, doi:10.1038/nature15514 (2015).

8   Ding, J. *et al.* Pore-forming activity and structural autoinhibition of the gasdermin
family. *Nature* **535**, 111-116, doi:10.1038/nature18590 (2016).

9   Schmid-Burgk, J. L. *et al.* Caspase-4 mediates non-canonical activation of the
NLRP3 inflammasome in human myeloid cells. *Eur J Immunol* **45**, 2911-2917,
doi:10.1002/eji.201545523 (2015).

10   Baker, P. J. *et al.* NLRP3 inflammasome activation downstream of cytoplasmic
LPS recognition by both caspase-4 and caspase-5. *Eur J Immunol* **45**, 2918-
2926, doi:10.1002/eji.201545655 (2015).

11   Liu, X. *et al.* Inflammasome-activated gasdermin D causes pyroptosis by forming
membrane pores. *Nature* **535**, 153-158, doi:10.1038/nature18629 (2016).

12   Russo, B. C. *et al.* Intermediate filaments enable pathogen docking to trigger
type 3 effector translocation. *Nat Microbiol* **1**, 16025,
doi:10.1038/nmicrobiol.2016.25 (2016).

13   Watson, J. L. *et al.* Shigella sonnei O-Antigen Inhibits Internalization, Vacuole
Escape, and Inflammasome Activation. *mBio* **10**, doi:10.1128/mBio.02654-19
(2019).

14 Zychlinsky, A., Prevost, M. C. & Sansonetti, P. J. Shigella flexneri induces
apoptosis in infected macrophages. *Nature* **358**, 167-169, doi:10.1038/358167a0
(1992).

15 Suzuki, T. *et al.* Differential regulation of caspase-1 activation, pyroptosis, and
autophagy via Ipaf and ASC in Shigella-infected macrophages. *PLoS Pathog* **3**,
e111, doi:10.1371/journal.ppat.0030111 (2007).

16 Ellwanger, K. *et al.* The NLR family pyrin domain-containing 11 protein
contributes to the regulation of inflammatory signaling. *The Journal of biological*
chemistry **293**, 2701-2710, doi:10.1074/jbc.RA117.000152 (2018).

17 Wu, C. *et al.* NLRP11 attenuates Toll-like receptor signalling by targeting TRAF6
for degradation via the ubiquitin ligase RNF19A. *Nat Commun* **8**, 1977,
doi:10.1038/s41467-017-02073-3 (2017).

18 Qin, Y. *et al.* NLRP11 disrupts MAVS signalosome to inhibit type I interferon
signaling and virus-induced apoptosis. *EMBO Rep* **18**, 2160-2171,
doi:10.15252/embr.201744480 (2017).

19 Sanada, T. *et al.* The Shigella flexneri effector OspI deamidates UBC13 to
dampen the inflammatory response. *Nature* **483**, 623-626,
doi:10.1038/nature10894 (2012).

20 Gaillard, J. L., Berche, P., Mounier, J., Richard, S. & Sansonetti, P. In vitro model
of penetration and intracellular growth of Listeria monocytogenes in the human
enterocyte-like cell line Caco-2. *Infect Immun* **55**, 2822-2829 (1987).

21 Meunier, E. *et al.* Caspase-11 activation requires lysis of pathogen-containing
vacuoles by IFN-induced GTPases. *Nature* **509**, 366-370,
doi:10.1038/nature13157 (2014).

22 Beuzon, C. R. *et al.* Salmonella maintains the integrity of its intracellular vacuole
through the action of SifA. *The EMBO journal* **19**, 3235-3249,
doi:10.1093/emboj/19.13.3235 (2000).

23 Grenier, J. M. *et al.* Functional screening of five PYPAF family members
identifies PYPAF5 as a novel regulator of NF-kappaB and caspase-1. *FEBS*
letters **530**, 73-78, doi:10.1016/s0014-5793(02)03416-6 (2002).

24 Bitto, N. J. *et al.* Membrane vesicles from Pseudomonas aeruginosa activate the
noncanonical inflammasome through caspase-5 in human monocytes. *Immunol*
Cell Biol **96**, 1120-1130, doi:10.1111/imcb.12190 (2018).

25 Pilla, D. M. *et al.* Guanylate binding proteins promote caspase-11-dependent
pyroptosis in response to cytoplasmic LPS. *Proc Natl Acad Sci U S A* **111**, 6046-
6051, doi:10.1073/pnas.1321700111 (2014).

26   Fisch, D. *et al.* Human GBP1 is a microbe-specific gatekeeper of macrophage
apoptosis and pyroptosis. *The EMBO journal* **38**, e100926,
doi:10.15252/embj.2018100926 (2019).

27   Santos, J. C. *et al.* LPS targets host guanylate-binding proteins to the bacterial
outer membrane for non-canonical inflammasome activation. *The EMBO journal*
37, doi:10.15252/embj.201798089 (2018).

28   Wandel, M. P. *et al.* Guanylate-binding proteins convert cytosolic bacteria into
caspase-4 signaling platforms. *Nat Immunol* **21**, 880-891, doi:10.1038/s41590-
020-0697-2 (2020).

29   Santos, J. C. *et al.* Human GBP1 binds LPS to initiate assembly of a caspase-4
activating platform on cytosolic bacteria. *Nat Commun* **11**, 3276,
doi:10.1038/s41467-020-16889-z (2020).

30   Cheng, Y. S., Colonno, R. J. & Yin, F. H. Interferon induction of fibroblast
proteins with guanylate binding activity. *The Journal of biological chemistry* **258**,
7746-7750 (1983).

31   Horisberger, M. A., Staeheli, P. & Haller, O. Interferon induces a unique protein
in mouse cells bearing a gene for resistance to influenza virus. *Proc Natl Acad*
Sci U S A **80**, 1910-1914, doi:10.1073/pnas.80.7.1910 (1983).

32   Kutsch, M. *et al.* Direct binding of polymeric GBP1 to LPS disrupts bacterial cell
envelope functions. *The EMBO journal* **39**, e104926,
doi:10.15252/embj.2020104926 (2020).

33   Paz, I. *et al.* Galectin-3, a marker for vacuole lysis by invasive pathogens. *Cell*
Microbiol **12**, 530-544, doi:10.1111/j.1462-5822.2009.01415.x (2010).

34   Feeley, E. M. *et al.* Galectin-3 directs antimicrobial guanylate binding proteins to
vacuoles furnished with bacterial secretion systems. *Proc Natl Acad Sci U S A*
114, E1698-E1706, doi:10.1073/pnas.1615771114 (2017).

35   Lo, T. H. *et al.* Galectin-3 promotes noncanonical inflammasome activation
through intracellular binding to lipopolysaccharide glycans. *Proc Natl Acad Sci U*
S A **118**, doi:10.1073/pnas.2026246118 (2021).

36   Taveira da Silva, A. M. *et al.* Brief report: shock and multiple-organ dysfunction
after self-administration of Salmonella endotoxin. *N Engl J Med* **328**, 1457-1460,
doi:10.1056/NEJM199305203282005 (1993).

37   Schaedler, R. W. & Dubos, R. J. The susceptibility of mice to bacterial
endotoxins. *J Exp Med* **113**, 559-570 (1961).

38   Sauter, C. & Wolfensberger, C. Interferon in human serum after injection of
endotoxin. *Lancet* **2**, 852-853 (1980).

39 Reynolds, K. *et al.* Pretreatment with troglitazone decreases lethality during
endotoxemia in mice. *J Endotoxin Res* **8**, 307-314,
doi:10.1179/096805102125000515 (2002).

40 McCuskey, R. S., McCuskey, P. A., Urbaschek, R. & Urbaschek, B. Species
differences in Kupffer cells and endotoxin sensitivity. *Infect Immun* **45**, 278-280
(1984).

41 Glode, L. M., Mergenhagen, S. E. & Rosenstreich, D. L. Significant contribution
of spleen cells in mediating the lethal effects of endotoxin in vivo. *Infect Immun*
14, 626-630 (1976).

42 van der Poll, T. & Lowry, S. F. in *Modulation of the inflammatory response in*
severe sepsis (eds J. M. Tellado, R. A. Forse, & J. S. Solomkin) 18-32 (Karger,
1995).

43 Labrec, E. H., Schneider, H., Magnani, T. J. & Formal, S. B. Epithelial Cell
Penetration as an Essential Step in the Pathogenesis of Bacillary Dysentery. *J*
Bacteriol **88**, 1503-1518, doi:10.1128/jb.88.5.1503-1518.1964 (1964).

44 Wing, H. J., Yan, A. W., Goldman, S. R. & Goldberg, M. B. Regulation of IcsP,
the outer membrane protease of the Shigella actin tail assembly protein IcsA, by
virulence plasmid regulators VirF and VirB. *J Bacteriol* **186**, 699-705,
doi:10.1128/JB.186.3.699-705.2004 (2004).

45 Maurelli, A. T., Blackmon, B. & Curtiss, R., 3rd. Loss of pigmentation in Shigella
flexneri 2a is correlated with loss of virulence and virulence-associated plasmid.
Infect Immun **43**, 397-401, doi:10.1128/iai.43.1.397-401.1984 (1984).

46 Miller, K. A., Garza-Mayers, A. C., Leung, Y. & Goldberg, M. B. Identification of
interactions among host and bacterial proteins and evaluation of their role early
during Shigella flexneri infection. *Microbiology (Reading)* **164**, 540-550,
doi:10.1099/mic.0.000637 (2018).

47 Novem, V. *et al.* Structural and biological diversity of lipopolysaccharides from
Burkholderia pseudomallei and Burkholderia thailandensis. *Clin Vaccine Immunol*
16, 1420-1428, doi:10.1128/CVI.00472-08 (2009).

48 Gedde, M. M., Higgins, D. E., Tilney, L. G. & Portnoy, D. A. Role of listeriolysin O
in cell-to-cell spread of Listeria monocytogenes. *Infect Immun* **68**, 999-1003,
doi:10.1128/IAI.68.2.999-1003.2000 (2000).

49 Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640-
6645, doi:10.1073/pnas.120163297 (2000).

50   Ranjbar, S., Jasenosky, L. D., Chow, N. & Goldfeld, A. E. Regulation of
Mycobacterium tuberculosis-dependent HIV-1 transcription reveals a new role for
NFAT5 in the toll-like receptor pathway. *PLoS Pathog* **8**, e1002620,
doi:10.1371/journal.ppat.1002620 (2012).

51   Gokhale, A., Ryder, P. V., Zlatic, S. A. & Faundez, V. Identification of the
Interactome of a Palmitoylated Membrane Protein, Phosphatidylinositol 4-Kinase
Type II Alpha. *Methods Mol Biol* **1376**, 35-42, doi:10.1007/978-1-4939-3170-5_4
(2016).

52   Kobayashi, T. *et al.* The Shigella OspC3 effector inhibits caspase-4, antagonizes
inflammatory cell death, and promotes epithelial infection. *Cell host & microbe*
13, 570-583, doi:10.1016/j.chom.2013.04.012 (2013).

53   Li, F. *et al.* Shigella evades pyroptosis by arginine ADP-riboxanation of caspase-
11. *Nature*, doi: 10.1038/s41586-41021-04020-41581. Online ahead of print.
(2021).

Figures and figure legends

Fig. 1. In the absence of *NLRP11*, cell death of human-derived macrophage-like cells induced by intracellular gram-negative bacteria is defective.

(**a-b**) Death of human-derived THP-1 macrophage-like cells upon infection with the cytosolic gram-negative pathogens *S. flexneri* (**a**), and *B. thailandensis* (**b**). Cell death in response to *S. flexneri* infection is defective in each of two independent *NLRP11*[-/-] THP-1 cell lines (**a**). (**c**) *NLRP11* is required for efficient cell death of THP-1 cells induced by the pathogen *S.* Typhimurium. Cell death induced by the *sifA* mutant, which is largely

cytosolic, was similar to that of WT *S.* Typhimurium, which is largely vacuolar. (**e-f**)

During *S. flexneri* infection of THP-1 macrophage-like cells, *NLRP11* is required for

processing of caspase-4 (**d**) and gasdermin D (**e**); western blot of culture supernatants.

In the absence of OspC3, processing was similar to induced by WT *S. flexneri*; in

HaCaT cells and SiHa cells, OspC3 inhibits caspase-4 activation, and in mouse bone

marrow-derived macrophages, OspC3 inhibits caspase-11 activation[52,53]. Cell death is

by LDH release, reported as the percent of total cell lysis induced by Triton X-100.

Mean ± S.E.M. Two-way ANOVA with Dunnett *post hoc* test (**a**, **c**) or with Sidak *post*

hoc test (**d-e**), or student T-test (**b**). *, $p<0.05$; **, $p<0.01$; ***; $p<0.001$; ****, $p<0.0001$;

a.u., arbitrary units.

Fig. 2. *NLRP11* potentiates cell death in human macrophages in response to cytosolic LPS.

(**a**) *NLRP11* potentiates death of THP-1 macrophage-like cells induced by infection with cytosolic gram-positive pathogen *L. monocytogenes* that is coated with *S. enterica* serovar Minnesota LPS. Cell death depends on coating of *L. monocytogenes* with LPS and on escape from the vacuole, as *L. monocytogenes LLO⁻* is restricted to the phagosome. (**b-c**) In the absence of *NLRP11* (**b**) or following NLRP11 siRNA (**c**), death of THP-1 macrophage-like cells in response to electroporated LPS is defective. (**d**) NLRP11-dependent cell death of THP-1 macrophage-like cells depends on caspase-4, but not caspase-1. Cell death following siRNA depletion of *CASP4* or *CASP1*. (**e-g**) NLRP11 is required for efficient cell death of primary human monocyte-derived

macrophages in response to cLPS. Cell death in isolated CD14+ cells differentiated to

macrophages, treated with siRNA to *NLRP11* (**e**), *caspase-4* (**f**), or *MyD88* (**g**), and

subsequently electroporated with LPS. Cell death measured as LDH release relative to

complete lysis control; in (**e-g**), normalized to mock transfection following scRNA

treatment. Tfxn, transfection; mock, carrier alone; scRNA, scrambled siRNA; CASP4,

caspase-4, CASP1, caspase-1. Two-way ANOVA with Sidak *post hoc* test (**a**) or Tukey

post hoc test (**b-d**); mixed-effects analyses (**e-g**). Mean ± S.E.M. **, $p < 0.01$; ***;

$p < 0.001$; ****, $p < 0.0001$; n.s., not significant.

Fig. 3. NLRP11 interactions with caspases in reconstituted HEK293T lysates.

(**a-b**) Precipitation of NLRP11 and wildtype caspase-4 (CASP4) by biotin-LPS,

separately (**a**) and together (**b**), and not by biotin-PAM3CSK4 (Pam3) (**a**). PD: SA, pull

down with streptavidin beads. (**c**) NLRP11 precipitates wildtype CASP4. (**d**) Interaction

between NLRP11 and CASP4 increases as a function of the concentration of each

protein. Binding of FLAG-NLRP11 in transfected HEK293T cells to plates coated with

purified wildtype CASP4 with detection of bound FLAG. Dilutions of NLRP11 lysates:

circles, undiluted; squares, 1:2 dilution; triangles, 1:4 dilution; inverted triangles, 1:8

dilution. Diamonds, untransfected (untfxd) lysate. Two-way ANOVA with Tukey *post hoc*

test. a.u., arbitrary units. (**e**) Specificity of NLRP11 for caspase-4 and caspase-1. FLAG-

600   tagged catalytically inactive caspases (CASP1[C285A]-FLAG, CASP5a[C315A]-FLAG,

601   CASP9[C287S]-FLAG) precipitation of Myc-NLRP11. CASP*, catalytically inactive

602   caspase. IP: FLAG, pull down with beads coated with FLAG antibody. Detection by

603   western blot with anti-FLAG or anti-Myc antibody.