1 *Type: Letter/Methods*

3 `selscan 2.0`: scanning for sweeps in unphased data

5 *Zachary A. Szpiech[1,2,*]*

7 [1] *Department of Biology, Pennsylvania State University, University Park, PA 16802, USA*
8 [2] *Institute for Computational and Data Sciences, Pennsylvania State University, University Park,*
9 *PA 16802, USA*
10 [*] *Correspondence: szpiech@psu.edu*

12 Abstract

Haplotype-based scans to identify recent and ongoing positive selection have become

commonplace in evolutionary genomics studies of numerous species across the tree of life.

However, the most widely adopted approaches require phased haplotypes to compute the key

statistics. Here we release a major update to the selscan software that re-defines popular

haplotype-based statistics for use with unphased "multi-locus genotype" data. We provide

unphased implementations of iHS, nSL, XP-EHH, and XP-nSL and evaluate their performance

across a range of important parameters in a generic demographic history. Source code and

executables are available at https://www.github.com/szpiech/selscan.

21 1 Introduction

Haplotype-based summary statistics—such as iHS (Voight, et al. 2006), nSL (Ferrer-

Admetlla, et al. 2014), XP-EHH (Sabeti, et al. 2007), and XP-nSL (Szpiech, et al. 2021)—have

become commonplace in evolutionary genomics studies to identify recent and ongoing positive

selection in populations (e.g.,Colonna, et al. 2014; Zoledziewska, et al. 2015; Nedelec, et al.

2016; Crawford, et al. 2017; Meier, et al. 2018; Lu, et al. 2019; Zhang, et al. 2020; Salmon, et al.

2021). When an adaptive allele sweeps through a population, it leaves a characteristic pattern

of long high-frequency haplotypes and low genetic diversity in the vicinity of the allele. These

statistics aim to capture these signals by summarizing the decay of haplotype homozygosity as

a function of distance from a putatively selected region, either within a single population (iHS

and nSL) or between two populations (XP-EHH and XP-nSL). However, each of these statistics

presumes that haplotype phase is known.

Recent work has shown that converting haplotype data into multi-locus genotype data is

an effective approach for using haplotype-based selection statistics such as G12, LASSI, and

saltiLASSI (Harris, et al. 2018; Harris and DeGiorgio 2020; DeGiorgio and Szpiech 2021) in

unphased data. Recognizing this, we have reformulated the iHS, nSL, XP-EHH, and XP-nSL

statistics to use multi-locus genotypes and provided an easy-to-use implementation in selscan

2.0 (Szpiech and Hernandez 2014). We also evaluate the performance of these unphased

statistics under various generic demographic models.

40   2 New Approaches

41   When the --unphased flag is set in selscan v2.0+, biallelic genotype data is collapsed

42   into multi-locus genotype data by representing the genotype as either 0, 1, or 2—the number of

43   derived alleles observed. In this case, selscan v2.0+ will then compute iHS, nSL, XP-EHH, and

44   XP-nSL as described below. We follow the notation conventions of Szpiech and Hernandez

45   (2014).

46   2.1 Extended Haplotype Homozygosity

In a sample of n diploid individuals, let $\mathcal{C}$ denote the set of all possible genotypes at

locus x_0. For multi-locus genotypes, $\mathcal{C} \coloneqq \{0,1,2\}$, representing the total counts of a derived

allele. Let $\mathcal{C}(x_i)$ be the set of all unique haplotypes extending from site x_0 to site x_i either

upstream or downstream of x_0. If x_1 is a site immediately adjacent to x_0, then $\mathcal{C}(x_1) \coloneqq$

$\{00,01,02,10,11,12,20,21,22\}$, representing all possible two-site multi-locus genotypes. We can

then compute the extended haplotype homozygosity (EHH) of a set of multi-locus genotypes as

53
$$EHH(x_i) \sum_{h \in \mathcal{C}(x_i)} \frac{\binom{n_h}{2}}{\binom{n}{2}},$$

where n_h is the number of observed haplotypes of type h.

If we wish to compute the EHH of a subset of observed haplotypes that all contain the

same 'core' multi-locus genotype, let $\mathcal{H}_c(x_i)$ be the partition of $\mathcal{C}(x_i)$ containing genotype $c \in \mathcal{C}$

at x_0. For example, choosing a homozygous derived genotype ($c = 2$) as the core, $\mathcal{H}_2 :=$

$\{20,21,22\}$. Thus, we can compute the EHH of all individuals carrying a given genotype at site x_0

extending out to site x_i as

60
$$EHH_c(x_i) = \sum_{h \in \mathcal{H}_c(x_i)} \frac{\binom{n_h}{2}}{\binom{n_c}{2}},$$

61   where n_h is the number of observed haplotypes of type h and n_c is the number of observed

62   multi-locus genotypes with core genotype of c. Finally, we can compute the complement EHH of

63   a sample of multi-locus genotypes as

64
$$cEHH_c(x_i) = \sum_{h \in \mathcal{C}(x_i)\backslash\mathcal{H}_c(x_i)} \frac{\binom{n_h}{2}}{\binom{n_{c'}}{2}},$$

65   where $n_{c'}$ is the number of observed multi-locus genotypes with a core genotype of not c.

66   **2.2 iHS and nSL**

67   Unphased iHS and nSL are calculated using the equations above. First, we compute the

68   integrated haplotype homozygosity (iHH) for the homozygous ancestral ($c = 0$) and derived ($c =$

69   2) core genotypes as

70
$$iHH_c = \sum_{i=1}^{|\mathcal{D}|} \frac{1}{2}\big(EHH_c(x_{i-1}) + EHH_c(x_i)\big)\mathrm{d}(x_{i-1}, x_i) + \sum_{i=1}^{|\mathcal{U}|} \frac{1}{2}\big(EHH_c(x_{i-1}) + EHH_c(x_i)\big)\mathrm{d}(x_{i-1}, x_i),$$

71   where $\mathcal{D}$ is the set of downstream sites from the core locus and $\mathcal{U}$ is the set of upstream sites.

72     $d(x_{i-1}, x_i)$ is a measure of genomic distance between to markers and is the genetic distance in

73   centimorgans or physical distance in basepairs for iHS (Voight, et al. 2006) or the number of

74   sites observed for nSL (Ferrer-Admetlla, et al. 2014). We similarly compute the complement

75   integrated haplotype homozygosity (ciHH) for both homozygous core genotypes as

76
$$ciHH_c = \sum_{i=1}^{|\mathcal{D}|} \frac{1}{2}\left(cEHH_c(x_{i-1}) + cEHH_c(x_i)\right)d(x_{i-1}, x_i)$$

77
$$+ \sum_{i=1}^{|\mathcal{U}|} \frac{1}{2}\left(cEHH_c(x_{i-1}) + cEHH_c(x_i)\right)d(x_{i-1}, x_i).$$

78 The (unstandardized) unphased iHS is then calculated as

79
$$iHS = \begin{cases} iHS_2, & \text{if } iHS_2 > iHS_0 \\ -iHS_0, & \text{otherwise} \end{cases}$$

where $iHS_2 = \log_{10}\left(\frac{iHH_2}{ciHH_2}\right)$ and $iHS_0 = \log_{10}\left(\frac{iHH_0}{ciHH_0}\right)$. Unstandardized iHS scores are then

normalized in frequency bins, as previously described (Voight, et al. 2006; Ferrer-Admetlla, et

al. 2014). Unstandardized unphased nSL is computed similarly with the appropriate distance

measure. Large positive scores indicate long high-frequency haplotypes with a homozygous

derived core genotype, and large negative scores indicate long high-frequency haplotypes with

a homozygous ancestral core genotype. Clusters of extreme scores in both directions indicate

evidence for a sweep.

2.3 XP-EHH and XP-nSL

Unphased XP-EHH and XP-nSL are calculated by comparing the iHH between

populations A and B, using the entire sample in each population. iHH in a population P is

computed as

91
$$iHH_P = \sum_{i=1}^{|\mathcal{D}|} \frac{1}{2}\left(EHH(x_{i-1}) + EHH(x_i)\right)d(x_{i-1}, x_i) + \sum_{i=1}^{|\mathcal{U}|} \frac{1}{2}\left(EHH(x_{i-1}) + EHH(x_i)\right)d(x_{i-1}, x_i),$$

where the distance measure is given as centimorgans or basepairs for XP-EHH (Sabeti, et al.

2007) and number of sites observed for XP-nSL (Szpiech, et al. 2021). The XP statistics

between population A and B are then computed as $XP = \log_{10}\left(\frac{iHH_A}{iHH_B}\right)$ and are normalized

genome wide in a single bin. Large positive scores indicate long high-frequency haplotypes in

population A, and large negative scores indicate long high-frequency haplotypes in population

B. Clusters of extreme scores in one direction indicate evidence for a sweep in that population.

98   3 Methods

99   3.1 Simulations

We evaluate the performance of the unphased versions of iHS, nSL, XP-EHH, and XP-

nSL under a generic two-population divergence model using the coalescent simulation program

discoal (Kern and Schrider 2016). We explore five versions of this generic model and name

them Demo 1 through Demo 5 (Table 1). Let N_0 and N_1 be the effective population sizes of

Population 0 and Population 1 after the split from their ancestral population (of size N_A). For

Demo 1, we keep a constant population size post-split and let $N_0 = N_1 = 10{,}000$. For Demo 2,

we keep a constant population size post-split and let $N_0 = 2N_1 = 10{,}000$. For Demo 3, we keep

a constant population size post-split and let $2N_0 = N_1 = 10{,}000$. For Demo 4, we initially set

$N_0 = N_1 = 10{,}000$ and let N_0 grow stepwise exponentially every 50 generations starting at 2,000

generations ago until $N_0 = 5N_1 = 50{,}000$. For Demo 5, we initially set $N_0 = N_1 = 10{,}000$ and let

N_1 grow stepwise exponentially every 50 generations starting at 2,000 generations ago until

$5N_0 = N_1 = 50{,}000$.

For each demographic history we vary the population divergence time $t_d \in$

$\{2000, 4000, 8000\}$ generations ago. For non-neutral simulations, we simulate a sweep in

Population 0 in the middle of the simulated region across a range of selection coefficients $s \in$

$\{0.005, 0.01, 0.02\}$. We vary the frequency at which the adaptive allele starts sweeping as $e \in$

$\{0, 0.01, 0.02, 0.05, 0.10\}$, where $e = 0$ indicates a hard sweep and $e > 0$ indicates a soft sweep,

and we also vary the frequency of the selected allele at time of sampling $f \in \{0.7, 0.8, 0.9, 1.0\}$

as well as $g \in \{50, 100\}$ representing fixation of the sweeping allele g generations ago. For all

simulations we set the genome length to be $L = 500{,}000$ basepairs, the ancestral effective

population size to be $N_A = 10{,}000$, the per site per generation mutation rate at $\mu = 2.35 \times 10^{-8}$,

and the per site per generation recombination rate at $r = 1.2 \times 10^{-8}$. For neutral simulations, we

simulate 1,000 replicates for each parameter set, and for non-neutral simulations we simulate

100 replicates for each parameter set. As iHS and nSL are single population statistics, we only

analyze Demo 1, Demo 3, and Demo 4 with these statistics, as Demo 2 and Demo 5 have a

constant size history identical to Demo 1 for Population 0, where the sweeps are simulated.

For all simulations, we compute the relevant statistics (--ihs, --nsl, --xpehh, or --xpnsl)

with selscan v2.0, using the --unphased and --trunc-ok flags. For iHS and XP-EHH, we also use

the --pmap flag in order to use physical distance instead of a recombination map.

3.2 Power and False Positive Rate

To compute power for iHS and nSL, we follow the approach of Voight et al. (2006). For

these statistics, each non-neutral replicate is individually normalized jointly with all matching

neutral replicates in 1% allele frequency bins. Because extreme values of the statistic are likely

to be clustered along the genome (Voight, et al. 2006), we then compute the proportion of

extreme scores ($|iHS| > 2$ or $|nSL| > 2$) within 100kbp non-overlapping windows. We then bin

these windows into 10 quantile bins based on the number of scores observed in each window

and call the top 1% of these windows as putatively under selection. We calculate the proportion

of non-neutral replicates that fall in this top 1% as the power. To compute the false positive rate,

we compute the proportion of neutral simulations that fall within the top 1%.

To compute power for XP-EHH and XP-nSL, we follow the approach of Szpiech et al.

(2021). For these statistics, each non-neutral replicate is individually normalized jointly with all

matching neutral replicates. Because extreme values of the statistic are likely to be clustered

along the genome (Szpiech, et al. 2021), we then compute the proportion of extreme scores

(XP-EHH > 2 or XP-nSL > 2) within 100kbp non-overlapping windows. We then bin these

windows into 10 quantile bins based on the number of scores observed in each window and call

the top 1% of these windows as putatively under selection. We calculate the proportion of non-

neutral replicates that fall in this top 1% as the power. To compute the false positive rate, we

compute the proportion of neutral simulations that fall within the top 1%.

4 Results

We find that the unphased versions of iHS and nSL have good power (Figures 1, S1-S4,

S13-16, and S21-24) to detect selection prior to fixation of the allele, with nSL generally

outperforming iHS. In smaller populations (Figure 1C and 1D), power does suffer relative to

larger populations (Figure 1A, 1B, 1E, 1F). Each of these statistics also have low false positive

rates hovering around 1% (Table S1).

Similarly, we find that the unphased versions of XP-EHH and XP-nSL have good power

as well (Figures 2, 3, S5-S12, S17-S20, and S25-S32). When the sweep takes place in the

smaller of the two populations (Figure 2C and 2D), we see a similar decrease in power. When

one population is undergoing exponential growth (Figure 3) performance is generally quite

good, likely the result of a larger effective selection coefficient in large populations. These two-

population statistics generally outperform their single-population counterparts, especially for

sweeps that have reached fixation recently. Each of these statistics also have low false positive

rates hovering around 1% (Table S1).

5 Discussion

We introduce multi-locus genotype versions of four popular haplotype-based selection

statistics—iHS (Voight, et al. 2006), nSL (Ferrer-Admetlla, et al. 2014), XP-EHH (Sabeti, et al.

2007), and XP-nSL (Szpiech, et al. 2021)—that can be used when the phase of genotypes is

unknown. We implement these updates in the latest v2.0 update of the program selscan

(Szpiech and Hernandez 2014), with source code and pre-compiled binaries available at

https://www.github.com/szpiech/selscan.

6 Acknowledgements

170   This work was supported by start-up funds from the Pennsylvania State University's

171   Department of Biology. Computations for this research were performed using the Pennsylvania

172   State University's Institute for Computational Data Sciences' Roar supercomputer.

174
175   **Figure 1**. Power curves for unphased implementations of iHS (A, C, and E) and nSL (B, D, and

176   F) under demographic histories Demo 1 (A and B), Demo 3 (C and D), and Demo 4 (E and F). s

177   is the selection coefficient, f is the frequency of the adaptive allele at time of sampling, g is the

178   number of generations at time of sampling since fixation, e is the frequency at which selection

179   began, and t_d is the time in generations since the two populations diverged.

Figure 2. Power curves for unphased implementations of XP-EHH (A, C, and E) and XP-nSL

(B, D, and F) under demographic histories Demo 1 (A and B), Demo 2 (C and D), and Demo 3

(E and F). s is the selection coefficient, f is the frequency of the adaptive allele at time of

sampling, g is the number of generations at time of sampling since fixation, e is the frequency at

which selection began, and t_d is the time in generations since the two populations diverged.

Figure 3. Power curves for unphased implementations of XP-EHH (A and C) and XP-nSL (B

and D) under demographic histories Demo 4 (A and B), and Demo 5 (C and D). s is the

selection coefficient, f is the frequency of the adaptive allele at time of sampling, g is the

number of generations at time of sampling since fixation, e is the frequency at which selection

began, and t_d is the time in generations since the two populations diverged.

Figure S1. Demo 1 iHS $t_d = 2000$ power curves. s is the selection coefficient, f is the frequency of the adaptive allele at time of sampling, g is the number of generations at time of sampling since fixation, e is the frequency at which selection began, and t_d is the time in generations since the two populations diverged.

Figure S2. Demo 1 iHS $t_d = 8000$ power curves. s is the selection coefficient, f is the frequency of the adaptive allele at time of sampling, g is the number of generations at time of sampling since fixation, e is the frequency at which selection began, and t_d is the time in generations since the two populations diverged.

Figure S3. Demo 1 nSL $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S4. Demo 1 nSL $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S5. Demo 1 XP-EHH $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S6. Demo 1 XP-EHH $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S7. Demo 1 XP-nSL $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S8. Demo 1 XP-nSL $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S9. Demo 2 XP-EHH $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S10. Demo 2 XP-EHH $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S11. Demo 2 XP-nSL $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S12. Demo 2 XP-nSL $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S13. Demo 3 iHS $t_d = 2000$ power curves. s is the selection coefficient, f is the frequency of the adaptive allele at time of sampling, g is the number of generations at time of sampling since fixation, e is the frequency at which selection began, and t_d is the time in generations since the two populations diverged.

Figure S14. Demo 3 iHS $t_d = 8000$ power curves. s is the selection coefficient, f is the frequency of the adaptive allele at time of sampling, g is the number of generations at time of sampling since fixation, e is the frequency at which selection began, and t_d is the time in generations since the two populations diverged.

Figure S15. Demo 3 nSL $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S16. Demo 3 nSL $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S17. Demo 3 XP-EHH $t_d = 2000$ power curves. s is the selection coefficient, f is the frequency of the adaptive allele at time of sampling, g is the number of generations at time of sampling since fixation, e is the frequency at which selection began, and t_d is the time in generations since the two populations diverged.

Figure S18. Demo 3 XP-EHH $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S19. Demo 3 XP-nSL $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S20. Demo 3 XP-nSL $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S21. Demo 4 iHS $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S22. Demo 4 iHS $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S23. Demo 4 nSL $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S24. Demo 4 nSL $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S25. Demo 4 XP-EHH $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S26. Demo 4 XP-EHH $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S27. Demo 4 XP-nSL $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S28. Demo 4 XP-nSL $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S29. Demo 5 XP-EHH $t_d = 2000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S30. Demo 5 XP-EHH $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

Figure S31. Demo 5 XP-nSL $t_d = 2000$ power curves. s is the selection coefficient, f is the frequency of the adaptive allele at time of sampling, g is the number of generations at time of sampling since fixation, e is the frequency at which selection began, and t_d is the time in generations since the two populations diverged.

Figure S32. Demo 5 XP-nSL $t_d = 8000$ power curves. s is the selection coefficient, f is the

frequency of the adaptive allele at time of sampling, g is the number of generations at time of

sampling since fixation, e is the frequency at which selection began, and t_d is the time in

generations since the two populations diverged.

361 **Table 1**. Demographic history parameters for simulations.

	N_A	N_0 at split	N_0 at present	N_1 at split	N_1 at present	t_d
Demo 1	10,000	10,000	10,000	10,000	10,000	2,000/4,000/8,000
Demo 2	10,000	10,000	10,000	5,000	5,000	2,000/4,000/8,000
Demo 3	10,000	5,000	5,000	10,000	10,000	2,000/4,000/8,000
Demo 4	10,000	10,000	50,000[†]	10,000	10,000	2,000/4,000/8,000
Demo 5	10,000	10,000	10,000	10,000	50,000[†]	2,000/4,000/8,000

362 [†]The reached via exponential growth starting 2,000 generations ago.

364 **Table S1**. False positive rate computed from neutral simulations for varying t_d and demographic
365 history.

		$t_d = 2000$	$t_d = 4000$	$t_d = 8000$
iHS	Demo 1	0.013	0.1	0.009
	Demo 3	0.007	0.013	0.007
	Demo 4	0.015	0.018	0.008
nSL	Demo 1	0.01	0.015	0.008
	Demo 3	0.008	0.011	0.007
	Demo 4	0.014	0.021	0.014
XP-EHH	Demo 1	0.013	0.013	0.016
	Demo 2	0.017	0.009	0.015
	Demo 3	0.01	0.011	0.012
	Demo 4	0.012	0.014	0.014
	Demo 5	0.011	0.012	0.013
XP-nSL	Demo 1	0.014	0.011	0.013
	Demo 2	0.019	0.011	0.012
	Demo 3	0.011	0.011	0.012
	Demo 4	0.012	0.012	0.014
	Demo 5	0.011	0.012	0.014

366
367

References

Colonna V, Ayub Q, Chen Y, Pagani L, Luisi P, Pybus M, Garrison E, Xue Y, Tyler-Smith C,
Genomes Project C, et al. 2014. Human genomic regions with exceptionally high levels
of population differentiation identified from 911 whole-genome sequences. Genome
Biol 15:R88.
Crawford NG, Kelly DE, Hansen MEB, Beltrame MH, Fan S, Bowman SL, Jewett E, Ranciaro A,
Thompson S, Lo Y, et al. 2017. Loci associated with skin pigmentation identified in
African populations. Science 358.
DeGiorgio M, Szpiech ZA. 2021. A spatially aware likelihood test to detect sweeps from
haplotype distributions. bioRxiv:2021.2005.2012.443825.
Ferrer-Admetlla A, Liang M, Korneliussen T, Nielsen R. 2014. On detecting incomplete soft
or hard selective sweeps using haplotype structure. Mol Biol Evol 31:1275-1291.
Harris AM, DeGiorgio M. 2020. A likelihood approach for uncovering selective sweep
signatures from haplotype data. Mol Biol Evol.

383 Harris AM, Garud NR, DeGiorgio M. 2018. Detection and Classification of Hard and Soft
384  Sweeps from Unphased Genotypes by Multilocus Genotype Identity. Genetics 210:1429-
385  1452.
386 Kern AD, Schrider DR. 2016. Discoal: flexible coalescent simulations with selection.
387  Bioinformatics 32:3839-3841.
388 Lu K, Wei L, Li X, Wang Y, Wu J, Liu M, Zhang C, Chen Z, Xiao Z, Jian H, et al. 2019. Whole-
389  genome resequencing reveals Brassica napus origin and genetic loci involved in its
390  improvement. Nat Commun 10:1154.
391 Meier JI, Marques DA, Wagner CE, Excoffier L, Seehausen O. 2018. Genomics of Parallel
392  Ecological Speciation in Lake Victoria Cichlids. Mol Biol Evol 35:1489-1506.
393 Nedelec Y, Sanz J, Baharian G, Szpiech ZA, Pacis A, Dumaine A, Grenier JC, Freiman A, Sams
394  AJ, Hebert S, et al. 2016. Genetic Ancestry and Natural Selection Drive Population
395  Differences in Immune Responses to Pathogens. Cell 167:657-669 e621.
396 Sabeti PC, Varilly P, Fry B, Lohmueller J, Hostetter E, Cotsapas C, Xie X, Byrne EH, McCarroll
397  SA, Gaudet R, et al. 2007. Genome-wide detection and characterization of positive
398  selection in human populations. Nature 449:913-918.
399 Salmon P, Jacobs A, Ahren D, Biard C, Dingemanse NJ, Dominoni DM, Helm B, Lundberg M,
400  Senar JC, Sprau P, et al. 2021. Continent-wide genomic signatures of adaptation to
401  urbanisation in a songbird across Europe. Nat Commun 12:2983.
402 Szpiech ZA, Hernandez RD. 2014. selscan: an efficient multithreaded program to perform
403  EHH-based scans for positive selection. Mol Biol Evol 31:2824-2827.
404 Szpiech ZA, Novak TE, Bailey NP, Stevison LS. 2021. Application of a novel haplotype-based
405  scan for local adaptation to study high-altitude adaptation in rhesus macaques. Evol
406  Lett 5:408-421.
407 Voight BF, Kudaravalli S, Wen X, Pritchard JK. 2006. A map of recent positive selection in
408  the human genome. Plos Biology 4:e72.
409 Zhang SJ, Wang GD, Ma P, Zhang LL, Yin TT, Liu YH, Otecko NO, Wang M, Ma YP, Wang L, et
410  al. 2020. Genomic regions under selection in the feralization of the dingoes. Nat
411  Commun 11:671.
412 Zoledziewska M, Sidore C, Chiang CWK, Sanna S, Mulas A, Steri M, Busonero F, Marcus JH,
413  Marongiu M, Maschio A, et al. 2015. Height-reducing variants and selection for short
414  stature in Sardinia. Nat Genet 47:1352-1356.
415