1   Rapid decline in Western monarch butterflies leads to phenological and geographic Allee effects

3   Collin Edwards[12]*, Cheryl Schultz[2], Elizabeth Crone[1]

5   [1]Department of Biology, Tufts University; Somerville, MA

6   [2]School of Biological Sciences, Washington State University; Vancouver, WA

7   *Corresponding author. Email: edwards.evoeco@gmail.com

Abstract: Allee effects are processes that become disrupted at low population sizes, causing further declines to eventual extinction. Identifying Allee effects in natural populations is important for both ecological theory and conservation efforts but challenging because populations typically only experience them briefly before collapsing. In 2018, the Western monarch butterfly population fell below the threshold that scientists predicted would trigger Allee effects. Here we show that since 2018, the Western monarch population has expanded more slowly during spring and summer breeding and filled a smaller breeding range. Other aspects of the monarch life cycle (winter survival, fecundity, morphology) have not declined after the crash. However, winter survival is much lower and wing sizes of overwintering monarchs are larger now than in the 20th century, suggesting higher mortality and selection for increased dispersal in recent decades. Delayed arrival to parts of the Western monarch breeding range is a novel Allee effect, which could lead to temporal mismatches between monarchs and their and host plants.

Keywords: *Western monarch butterfly; Danaus plexippus; Allee effects; phenology; population viability; surprises*

Introduction

When populations decline to very low densities, individuals can face additional distinct challenges, such as difficulty finding mates, that decrease population growth (Stephens et al. 1999). These challenges can cause even more rapid declines, possibly contributing to an extinction vortex (Gilpin & Soulé 1986). Positive density dependence at low population size – known as Allee effects – can prevent invasions as well as drive extinctions (Courchamp et al. 1999). Documenting and understanding Allee effects is important both for conservation biology and for broader ecological theory. However, there is only a very narrow window of time when Allee effects are observable before a population is extirpated.

Western monarch butterflies have rapidly declined in the past forty years. Overwintering population estimates were millions of butterflies in the 20th century, but the population declined by an average of 6% per year through 2016 (Schultz et al. 2017). In 2017 we predicted that Allee effects would occur if the Western monarch population fell below 30,000 butterflies (Schultz et al. 2017). The following spring there was a massive decline, and the population dipped below that threshold at Thanksgiving 2018 (27,721 butterflies, Xerces Society 2021).

Here, we test for Allee effects in a range of demographic processes. We measured two broad categories of monarch performance following the 2018 crash: traits of overwintering monarch butterflies, and rates of population expansion during the summer breeding season. We compare these measurements to comparable measures of performance from studies done when monarch populations were higher. We categorized time periods into three eras: "historical", before 2000 when there were millions of monarchs; "recent past", from 2000-2017 when monarchs were in the hundreds of thousands; and "current", from 2018-2020 when winter counts were below the hypothesized threshold for Allee effects. Differences between the current era and

recent past indicate Allee effects. Differences between the historic era and current or recent past

indicate possible drivers of long-term declines.

Methods:

Overwintering monarch butterflies: In February 2019, we collected 35 female monarchs

near the overwintering site at Pismo Beach State Park. We measured wing length and body size,

used the palpitation method to evaluate which had been mated (Van Hook 1997), and measured

daily fecundity of caged individuals (see detailed methods in Appendix S1). We analyzed

previously published Thanksgiving and New Years Count data to estimate overwinter survival

(Xerces Society 2021). We identified studies with comparable demographic measures from

before the 2018 crash (overwinter survival: Tuskes & Brower 1978; body mass: Chaplin and

Wells 1982, Leong et al. 1995; wing size: Beall 1946, Tuskes and Brower 1978, Leong et al.

1995; fecundity: Leong et al. 1995, 2012). For each performance metric, we either obtained

estimates directly or through back-calculation with error propagation (see detailed methods in

Appendix S1).

Summer population expansion: To characterize the yearly population expansion into the

summer breeding grounds, we used community-science observations of monarchs available

through Xerces and iNaturalist (Western Monarch and Milkweed Occurrence database 2018,

GBIF.org 2020). These data included monarch observations across western North America from

the late 1800s through 2020, but required careful attention to address its ad-hoc and presence-

only nature. We used quantile regression (Cade and Noon 2003) to fit distance from

overwintering sites through time as a Gaussian curve (Edwards and Crone 2021) (Figure 2B).

We used the 0.9 quantile, which represents the boundary between the 10% of monarchs that were

most distant and the remaining 90%, as a measure of breeding season range size that is robust to

outliers (we find similar patterns for other quantiles; see Figure S2). From fitted curves for each

era, we estimated the maximum (i.e., 90% quantile) distance from overwintering sites (a measure

of the size of the breeding range), the day of maximum distance from overwintering sites (a

measure of phenology that indicates the shift from population expansion to contraction), and the

day the 90% quantile first reached 100 kilometers (a different measure of phenology that

indicates how quickly they arrived in summer breeding areas away from the coast).

Results and Discussion

There were no Allee effects in overwinter survival (Fig. 1A), body mass (Fig. 1B), wing

length (Fig. 1C), or mating success (Fig. 1D). In contrast, there were strong Allee effects in

breeding range. The historic and recent past eras had similar breeding ranges (Fig. 2B, Movie

S1), with the 0.9 quantile of the population expanding to up to 1070 and 1003 kilometers from

overwintering sites, respectively (Fig. 2F). Starting in 2018, this maximum distance traveled fell

by more than 50%, to 491 kilometers. Western monarchs also experienced a shorter breeding

season after 2018: summer population expansion stopped almost a week earlier (Fig. 2C).

Further, in the current era, monarchs have reached regions within their extant breeding grounds

up to two months later (Fig 2D,E).

Shifts in phenology (timing of life history events) are a conspicuous effect of climate

change (Parmesan 2007), and can lead to temporal mismatches between interacting species,

including insects and their hosts (Forrest 2016). Changes in monarch phenology reported here

differ dramatically from typical climate responses; the average phenological shift across 203

species in the Northern hemisphere is 2.8 days/decade (Parmesan 2007), while between recent

past and current eras, monarch arrival times were delayed between 3.6 and 60.2 days, depending

on distance from overwintering sites. Our findings suggest that later generations of monarchs are

now experiencing very different life stages ("phenophases") of their milkweed hostplants. This

shift is especially concerning because monarch caterpillar survival and growth strongly depend

on their phenology in relation to host plants and predators (Yang and Cenzer 2020).

Although not Allee effects, there were two notable changes between the historic and

current eras: a reduction in overwintering survival (Fig. 1A), and an increase in wing length (Fig.

1C). Reduced winter survival may have contributed to the long-term decline of Western

monarchs. This reduction could reflect reduced habitat quality, climate-driven disruption of

overwintering behavior, or reduced butterfly conditions in the summer breeding ground (Crone

et al. 2019). Larger body size and wing length (measured in February) may indicate biased

overwintering survival, with smaller individuals dying at higher rates. Alternately, (Halsch et al

2020) hypothesized that the Central Valley of California (immediately east of many

overwintering sites) has been substantially degraded as monarch habitat. If selection favored

individuals that flew past the Central Valley, the Western monarch population may have evolved

increased wing size, which is associated with long-range movement (Yang et al. 2016).

The Allee effects we found were not the ones we expected (Schultz et al. 2017),

highlighting the ways in which dynamics become surprising in small populations. Similarly,

while finalizing this manuscript, early reports indicate there will likely be 200,000 or more

individuals in the 2021 Thanksgiving count -- far more than in 2020, but far less than the

millions of individuals of the late 20th century. Preliminary analyses suggest that the Western

monarch breeding range in Summer 2021 was not a return to the fast expansion and larger

breeding range of the recent past or historic eras (Fig. 3). Instead, the population increased this

year *in spite of* reduced breeding season range. This recent change highlights that we do not

know whether reduced colonization of the breeding range is a cause of population declines, or

simply a consequence of unprecedentedly low population sizes. As populations approach

extinction, population growth rates of many animal species become more variable among years

(Fagan and Holmes 2006). One important lesson from this system is that we should expect

surprises in population dynamics as species enter uncharted conditions. It is easy to consider

uncharted conditions in terms of the external environment, but it is equally important to

recognize the importance of how populations respond to changes in their own abundance,

especially to unprecedentedly low population sizes.

Data availability: All custom scripts and data that support the findings of this study will

be deposited to Dryad prior to final publication.

Acknowledgments:

Leone Brown, Norah Warchola, and Chelsea Thomas assisted with the field surveys and

extraction of previously published results; Emma Pelton and Stephanie McKnight provided

guidance and insight in working with Xerces data. California State Parks, California Department

of Fish and Game and USFWS were all important in securing permits to work with monarch

butterflies at Pismo Beach State Park. California State Parks generously provided use of their

greenhouse to conduct studies on reproductive biology of female monarchs at Pismo Beach,

including Danielle Bronson, Stephanie Little and Ronnie Glick. Cheryl Powers, a docent

volunteering at the Pismo Beach site, generously opened her home to us to support lodging needs

by staff throughout February 2019, and Shore Cliff Hotel donated additional lodging during the

peak activity period. We are grateful to Marm Kilpatrick and Sarina Jepsen for discussions

during planning and interpretation of data. Funding for this work was contributed by CREOi

(Conservation Research Education and Opportunities International), WSU Vancouver Fund for

Ecology and Conservation of Northwest Butterflies and Their Habitats as well as private donors

to WSU and UC Santa Cruz, and the National Science Foundation DEB 1920834.

Author contributions:

Conceptualization: CBE, CBS, EEC

Data Curation: CBE, CBS, EEC

Formal Analysis: CBE

Funding acquisition: CBS, EEC

Investigation: CBS, EEC

Methodology: CBE, EEC

Project administration: EEC

Software: CBE

Supervision: CBS, EEC

Visualization: CBE

Writing – original draft: CBE

Writing – review & editing: CBE, CBS, EEC

Competing interests: Authors declare they have no competing interests.

Supplemental Information is available for this paper. Correspondences and requests for materials

should be addressed to CBE.

Reference

G. Beall, Seasonal variation in sex proportion and wing length in the migrant butterfly, *Danaus*

plexippus l. (Lep. Danaidae). Trans. R. Entomol. Soc. Lond. 97, 337–353 (1946). doi:

10.1111/j.1365-2311.1946.tb00268.x

Cade, B. S., and B. R. Noon. 2003. A gentle introduction to quantile regression for ecologists.

Frontiers in ecology and the environment 1:412–420.

S. B. Chaplin, P. H. Wells, Energy reserves and metabolic expenditures of monarch butterflies

overwintering in southern California. Ecol. Entomol. 7, 249–256 (1982). doi:

10.1111/j.1365-2311.1982.tb00664.x

F. Courchamp, T. Clutton-Brock, B. Grenfell, Inverse density dependence and the Allee effect.

Trends Ecol. Evol. 14, 405–410 (1999). doi: 10.1016/S0169-5347(99)01683-3

E. E. Crone, E. M. Pelton, L. M. Brown, C. C. Thomas, C. B. Schultz, Why are monarch

butterflies declining in the West? Understanding the importance of multiple correlated

drivers. Ecol. Appl. 29 (2019). doi: 10.1002/eap.1975

C. B. Edwards, E. E. Crone, Estimating abundance and phenology from transect count data with

GLMs. Oikos (2021). doi: 10.1111/oik.08368.

W. F. Fagan, E. E. Holmes, Quantifying the extinction vortex. Ecol. Lett. 9, 51–60 (2006). doi:

10.1111/j.1461-0248.2005.00845.x

J. R. Forrest, Complex responses of insect phenology to climate change. Curr Opin Insect Sci.

17, 49–54 (2016). doi: 10.1016/j.cois.2016.07.002

GBIF.org. 17 December 2020. GBIF Occurrence Download doi: 10.15468/dl.aq6d3w

M.E. Gilpin & M.E. Soulé, M.E. (1986). Minimum viable populations: processes of extinction.

In: Conservation Biology: The Science of Scarcity and Diversity (ed. Soulé, M.E.). Sinauer

Associates, Sunderland, MA, pp. 19–34.

C. A. Halsch, A. Code, S. M. Hoyle, J. A. Fordyce, N. Baert, M. L. Forister, Pesticide

contamination of milkweeds across the agricultural, urban, and open spaces of low-elevation

Northern California. Frontiers in Ecology and Evolution. 8, 162 (2020). doi:

10.3389/fevo.2020.00162

K. L. H. Leong, E. O'Brien, K. Lowerisen, M. Colleran, Mating activity and status of

overwintering monarch butterflies (Lepidoptera: Danaidae) in Central California. Ann.

Entomol. Soc. Am. 88, 45–50 (1995). doi: 10.1093/aesa/88.1.45

K. H. L. Leong, M. A. Yoshimura, C. Williams, Adaptive significance of previously mated

monarch butterfly females (*Danaus plexippus* (Linneaus)) overwintering at a California

winter site. lepi. 66, 205–210 (2012). doi: 10.18473/lepi.v66i4.a3

P. C. Parmesan, Influences of species, latitudes and methodologies on estimates of phenological

response to global warming. Glob. Chang. Biol. 13, 1860–1872 (2007). doi: 10.1111/j.1365-

2486.2007.01404.x

M. E. Payton, M. H. Greenstone, N. Schenker, Overlapping confidence intervals or standard

error intervals: what do they mean in terms of statistical significance? J. Insect Sci. 3, 34

(2003). doi: 10.1093/jis/3.1.34

C. B. Schultz, L. M. Brown, E. Pelton, E. E. Crone, Citizen science monitoring demonstrates

dramatic declines of monarch butterflies in Western North America. Biol. Conserv. 214,

343–346 (2017). doi: 10.1016/j.biocon.2017.08.019

P. A. Stephens, W. J. Sutherland, R. P. Freckleton, What is the Allee effect? Oikos, 185–190

(1999). doi: 10.2307/3547011

P. M. Tuskes, L. P. Brower, Overwintering ecology of the monarch butterfly, *Danaus plexippus*

L., in California. Ecol. Entomol. 3, 141–153 (1978). doi: 10.1111/j.1365-

2311.1978.tb00912.x

T. Van Hook, in 1997 North American conference on the monarch butterfly (Commission for

Environmental Cooperation, 1999; p. 101.

Western Monarch and Milkweed Occurrence Database. 2018. Data accessed from the Western

Monarch Milkweed Mapper, a project by the Xerces Society, U.S. Fish and Wildlife Service,

Idaho department of Fish and Game, www.monarchmilkweedmapper.org. Accessed: 12/3/20.

Xerces Society. Western Monarch Thanksgiving Count Data, 1997– 2020. Available at

www.westernmonarchcount.org Accessed 23 January 2021. (2021)

L. H. Yang, M. L. Cenzer, Seasonal windows of opportunity in milkweed-monarch interactions.

Ecology. 101, e02880 (2020). doi: 10.1002/ecy.2880

L. H. Yang, D. Ostrovsky, M. C. Rogers, J. M. Welker, Intra-population variation in the natal

origins and wing morphology of overwintering Western monarch butterflies *Danaus*

plexippus. Ecography . 39, 998–1007 (2016). doi: 10.1111/ecog.01994

Fig. 1. Comparison across eras for (**A**) winter survival (from Thanksgiving to New Year's Day), (**B**) body mass, (**C**) wing length, (**D**) proportion of females mated, (**E**) fecundity, and (**F**) breeding season range as measured by maximum distance from overwintering sites. Points correspond to estimated values before 2000 ("historic"), from 2001-2017 ("recent past"), and from 2018-2020 ("current") when the population was below the hypothesized Allee effect threshold. In some cases (B, C, E), no published data was available to for either the historic or recent past eras. Bars represent 84% confidence intervals (nonoverlapping bars represent significant differences) (13). Allee effects imply a decrease in performance for very small populations; for the Western monarch butterfly, this means a change from the recent past to the current era. Only breeding season range (F) shows signs of an Allee effect.

Fig. 2. (**A**) Raw observation data show fewer observations in OR and WA in the current era. Shaded southern CA and NV excluded to avoid resident populations. White triangles show Sacramento, CA (see D). (**B**) Fitted curves of the 0.9 quantile of the distance from observations to the nearest overwintering site (a measure of expansion into the summer breeding range) show substantial reduction in the current era, even after correcting for changes in abundance. (**C**) Comparisons across eras of day of the maximum distance (i.e. day of peak in (B)) show monarchs now leave their summer breeding range sooner. (**D**) Days for the 0.9 quantile to reach 100 km from the nearest overwintering site; arrival day is much later in current era than in recent past or historical era. (**E**) Phenology in the current era is delayed compared to the recent past, with more delay further from overwintering sites. Black line: difference in arrival of 0.9 quantile in current era compared to recent past. In C&D, bars show 84% confidence intervals; in E, gray region shows 95% CI.

Fig. 3 High numbers of monarch observations allow us to the past seven years individually. (A-
C) Breeding range measures analogous to Fig. 2F, Fig. 3C-D. Bars show 84% CI. (A) and (C)
show clear differences between eras that reflect the patterns reported in the main text, although
monarchs had particularly late phenology in 2019. 2021 estimates were fit using observations
through October 27, 2021, and were not part of the main analyses. These preliminary estimates
show that despite higher counts in Thanksgiving 2021, the spatial Allee effect still occurred
during the preceding summer.

Figure 1

Figure 2

Figure 3

Movie S1: Western monarch butterfly yearly range expansion reaches half the distance and

takes up to two months longer since the population fell below the hypothesized Allee effect

threshold. Top: plots of raw data observations separated by era (Historic = 20[th] century, Recent

past = 2000-2017, Current = 2018-2020). Colored circles represent observations in the last 7

days; smaller white diamonds act as placeholders to identify locations of earlier observations.

Note the exclusion of AZ and southern regions of CA and NV to avoid resident populations.

Only points at least 10 kilometers from overwintering sites are plotted. Bottom: the same

observations plotted by day of year and distance from the nearest overwintering sites; curves

show Gaussian curves fit to the 0.9 quantile of observations for each era, representing the yearly

expansion into the summer breeding range.