Accurate estimation of intrinsic biases for improved analysis of chromatin accessibility

sequencing data using SELMA

Shengen Shawn Hu[1], Lin Liu[2], Qi Li[3], Wenjing Ma[1,4], Michael J. Guertin[5], Clifford A. Meyer[6], Ke

Deng[3], Tingting Zhang[7], Chongzhi Zang[1,8,9]*

[1]Center for Public Health Genomics, University of Virginia, Charlottesville, VA 22908, USA

[2]Institute of Natural Sciences, MOE-LSC, School of Mathematical Sciences, and SJTU-Yale

Joint Center for Biostatistics and Data Science, Shanghai Jiao Tong University, Shanghai

200240, China

[3]Center for Statistical Science, Tsinghua University, Beijing 100084, China

[4]Department of Computer Science, Emory University, Atlanta, GA 30322, USA

[5]Center for Cell Analysis and Modeling, Department of Genetics and Genome Sciences,

University of Connecticut, Farmington, CT 06030, USA

[6]Department of Data Science, Dana-Farber Cancer Institute, Boston, MA 02215, USA

[7]Department of Statistics, University of Pittsburgh, Pittsburgh, PA 15260, USA

[8]Department of Public Health Sciences, University of Virginia, Charlottesville, VA 22908, USA

[9]UVA Cancer Center, University of Virginia, Charlottesville, VA 22908, USA

*Correspondence should be addressed to C.Z.: zang@virginia.edu

Abstract

Genome-wide profiling of chromatin accessibility by DNase-seq or ATAC-seq has been widely used to identify regulatory DNA elements and transcription factor binding sites. However, enzymatic DNA cleavage exhibits intrinsic sequence biases that confound chromatin accessibility profiling data analysis. Existing computational tools are limited in their ability to account for such intrinsic biases. Here, we present Simplex Encoded Linear Model for Accessible Chromatin (SELMA), a computational method for systematic estimation of intrinsic cleavage biases from genomic chromatin accessibility profiling data. We demonstrate that SELMA yields accurate and robust bias estimation from both bulk and single-cell DNase-seq and ATAC-seq data. We show that transcription factor binding inference from DNase footprints can be improved by incorporating estimated biases using SELMA. We also demonstrate improved cell clustering of single-cell ATAC-seq data by considering the SELMA-estimated bias effect. SELMA can be applied to existing bioinformatics tools to improve the analysis of chromatin accessibility sequencing data.

Introduction

Cis-regulatory elements in the genome play a critical role in transcription regulation by interaction with protein molecules such as transcription factors (TFs). These DNA elements are usually unwrapped from packed nucleosomes and are accessible in the chromatin structure[1,2]. Genome-wide profiles of chromatin accessibility are a means to measure the global landscapes of active regulatory elements in different cell types. DNaseI hypersensitivity sequencing (DNase-seq) and the assay for transposase-accessible chromatin using sequencing (ATAC-seq) have become widely used for the genomic profiling of chromatin structure and accessibility[3,4]. Signal enrichments, or "peaks", from DNase-seq or ATAC-seq data are considered to represent accessible chromatin regions and can be used for inferring regulatory elements or TF binding sites. In addition, DNase-seq and ATAC-seq data also exhibit footprint patterns, which are fine structures in the accessible chromatin where high-affinity protein-DNA interactions protect the DNA from DNaseI or Tn5-transposase cleavages[4,5]. DNase/ATAC-seq footprint detection has been implicated as an effective approach for identifying accurate TF binding sites at base-pair resolution[6,7]. A few computational tools have been developed for detecting footprints from DNase-seq (RepFootprint[8], Wellington[9], PIQ[10]) or ATAC-seq data (HINT-ATAC[11], ToBIAS[12]). A recent study integrated 243 DNase-seq samples from different human cell and tissue types and reported approximately 4.5 million DNaseI consensus footprints associated with TF occupancy across the human genome as one of the largest maps of human regulatory DNA[7].

The premise of using DNase-seq and ATAC-seq data to profile chromatin accessibility is that enzymatic DNA cleavages represented by sequence reads reflect local chromatin openness only. However, it has been shown that both DNaseI and Tn5 transposase exhibit sequence selection biases in high-throughput sequencing data[13–16]. Such intrinsic enzymatic biases in

sequencing data can potentially confound observed cleavage patterns and footprint detection.

The characterization and correction of intrinsic cleavage biases are essential to DNase/ATAC-

seq data analysis. To characterize intrinsic cleavage biases, most studies use a k-mer model, in

which the k-mer DNA sequence centered at a cleavage site of DNaseI/Tn5 is used as the

signature of this cleavage[14,15,17]. The sequence bias can be estimated by counting the

occurrences of cleavages with each k-mer in one dataset relative to the genome-wide

occurrences of this k-mer. A naïve k-mer model assumes that k-mers are independent of each

other, resulting in an exponential increase in the degree of freedom when k increases.

Therefore, a naïve k-mer model becomes less feasible in practice with a large k, especially with

samples having insufficient sequencing depth. Although most studies use 6-mers with $4^6 = 4096$

parameters[8,10,14,15,17,18], it is unclear whether a different model with a larger k-mer size and more

feasible parameter estimation can achieve better performance.

Several studies have used various computational approaches for intrinsic cleavage bias

estimation[8,15,16,19] and footprint detection with bias correction[15,18–20] for bulk DNase/ATAC-seq

data. Recently, single-cell ATAC-seq (scATAC-seq) has enabled chromatin accessibility

profiling in thousands of individual cells at one time[21–24]. Due to the high sparsity of single-cell

data and because most chromatin accessibility regions contain only one read in one cell, any

potential bias can be substantial in scATAC-seq data compared to bulk data, creating additional

challenges in computational analysis. Incorporating intrinsic cleavage bias effects for improved

scATAC-seq analysis also requires comprehensive assessment and development of innovative

methods.

Here, we present <u>S</u>implex <u>E</u>ncoded <u>L</u>inear <u>M</u>odel for <u>A</u>ccessible Chromatin (SELMA), a

computational framework for the accurate estimation of intrinsic cleavage biases and improved

analysis of DNase/ATAC-seq data for both bulk and single-cell experiments. We demonstrate

that SELMA generates more accurate and robust bias estimation from bulk DNase/ATAC-seq

data than the naïve k-mer model and that SELMA can utilize mitochondrial DNA (mtDNA) for

bias estimation instead of requiring a separate naked DNA sample. We show an improved TF

occupancy inference on ENCODE consensus footprints by including SELMA-estimated biases

for each footprint. Finally, we show that SELMA-estimated biases can be incorporated with

existing scATAC-seq computational tools to generate more accurate cell clustering analysis.

Results

SELMA improves cleavage bias estimation in DNase-seq and ATAC-seq data

We developed SELMA for an accurate and robust estimation of intrinsic cleavage biases from

chromatin accessibility sequencing data. In SELMA, we start with a naïve k-mer model to

calculate the frequency of observed cleavage sites at each k-mer relative to the total

occurrences of this k-mer (Fig. 1a). We further encode each k-mer as a vector in the Hadamard

Matrix H_{16}, derived from a simplex encoding model, in which the k-mer sequences are encoded

as the vertices of a regular 0-centered simplex[25,26]. Intuitively, a k-mer can be represented as k

mononucleotides and $k - 1$ adjacent dinucleotides. Each mononucleotide is encoded as the 3-

dimensional vector of one of the four tetrahedral vertices of the cube of side 2 centered at the

origin. Each dinucleotide is encoded as the outer product of the 2 vectors representing the

associated nucleotides (Fig. 1b). This k-mer simplex encoding model has $1 + 3 \times k +$

$9 \times (k - 1) = 12k - 8$ parameters, much fewer than the naïve k-mer model (4^k). We use a

linear model to fit these $12k - 8$ parameters from the naïve k-mer biases, and use the fitted

values as the SELMA-modeled cleavage biases (Fig. 1c).

As an intrinsic property of the enzyme (DNaseI/Tn5 transposase), the cleavage biases are

expected to be invariant across cell types and independent of chromatin states (Fig. S1a).

Comparing data from two cell types using different 8-mer models as an example, we found that

the biases estimated using SELMA have a higher correlation than those estimated using the

naïve k-mer model, for both DNase-seq (Fig. 1d,e) and ATAC-seq (Fig. 1f,g). Using sequence

reads from genomic regions with different chromatin accessibility levels, the naïve k-mer model

was not able to generate very consistent bias estimations (Fig. S1b,c), but the k-mer biases

estimated by SELMA using the same data were highly consistent (Fig. S1b-e). The sequence

preferences of DNaseI or Tn5 cleavage can be better reflected when the enzymes are applied

to deproteinized naked genomic DNA[15,17]. We found that the k-mer cleavage biases in naked

DNA DNase/ATAC-seq data estimated with the naïve k-mer model can still be improved by

SELMA, obtaining more consistent bias scores between different cell systems (Fig. 1h-k). These

data demonstrated that SELMA can improve the accuracy of estimating k-mer cleavage biases

in DNase-seq and ATAC-seq data.

With fewer parameters, SELMA enabled us to assess the effect of k-mer size on the

performance of bias estimation. Using a "bias expected cleavage" approach[8,14,15,19], we

compared the bias estimation performances measured as the correlation coefficient between

the genome-wide observed cleavages and bias expected cleavages estimated using SELMA

with different k. A higher correlation coefficient indicates a more accurate bias estimation to

calculate the expected cleavages. For both DNase-seq and ATAC-seq data from two different

cell lines, we found that 10-mer outperforms any other k-mer (Fig. 1l-o). We then applied this

analysis to more DNase-seq and ATAC-seq data from a variety of human tissues generated by

ENCODE and found that 10-mer always exhibited the best performance (Fig. S2). The above

empirical evidence suggested that 10-mer is the optimal choice for intrinsic cleavage bias

estimation for both DNaseI (DNase-seq) and Tn5 (ATAC-seq) cleavages.

SELMA improves ATAC-seq bias estimation by considering dimeric Tn5 cleavages

SELMA improves ATAC-seq bias estimation by considering dimeric Tn5 cleavages

Many studies processed DNase-seq and ATAC-seq data in a similar way, treating individual

DNA cleavage sites directly as indications of accessible chromatin[8,16]. However, the

mechanisms of enzymatic DNA cleavage are different between DNaseI and Tn5 transposases.

Unlike DNaseI, the Tn5 transposase binds DNA as a dimer and inserts adaptors on the two

strands separated by 9bp[4,27] (Fig. 2a). As a result, the presence of each observed Tn5 cleavage

in ATAC-seq data should be the consequence of two insertion events induced by the same Tn5

dimer, and the bias estimation of a Tn5 cleavage site should consider both the observed

cleavage and the coupling cleavage 9bp downstream on the reverse strand. Therefore, SELMA

estimated the bias on an ATAC-seq cleavage site as the geometric mean of the bias scores of

the 10-mers at the 5' observed cleavage and at the 3' cleavage 9bp downstream on the reverse

strand (Fig. 2a). To show the dimeric Tn5 cleavage effect, we calculated the cross-correlation

between the genome-wide profiles of the plus strand cleavages and the minus strand

cleavages. As expected, we observed a peak at 9bp of the cross-correlation curve specifically in

the ATAC-seq data but not the DNase-seq data (Fig. 2b,c). Similarly, we observed perfectly

matching aggregate cleavage patterns on TF motif consensus sites between plus strand and

minus strand cleavages shifted by 9bp (Fig. S3a,b).

Using the similar "observed-expected correlation" approach described above, we found that

SELMA considering dimeric cleavages outperformed models considering only 5' cleavage in

generating a more accurate bias expected cleavage pattern for ATAC-seq data (Fig. 2d,e). We

confirmed that this observation was specific to ATAC-seq, as similar analyses for DNase-seq

from the same cell lines did not show a similar level of improvement (Fig. S3c,d). We also

compared SELMA with several existing bias estimation approaches[18] and found that SELMA's

performance was the best for ATAC-seq data from several cell lines as well as different human

tissue types from ENCODE (Fig. 2d,e, Fig. S3e-j). Several new Tn5-based techniques have

recently been developed for improved chromatin accessibility profiling, including THS-seq[28],

Fast-ATAC-seq[29], and Omni-ATAC-seq[30]. Data from these technologies showed intrinsic

cleavage biases similar to those of conventional ATAC-seq (Fig. S3k-n). We found that SELMA

also outperformed other approaches in bias estimation (Fig. 2f-i). Collectively, these data

suggested that SELMA can most accurately estimate intrinsic cleavage biases in data from

ATAC-seq and other Tn5-based techniques.

SELMA enables accurate bias estimation by utilizing mitochondrial DNA (mtDNA) reads

Accurate estimation of enzymatic cleavage biases independent of chromatin usually requires a

control sample of deproteinized naked DNA digested by the same enzyme. Biases estimated

from the naked DNA control data can be used to correct the chromatin accessibility profiling

data[14–18] (Fig. 3a). In conventional DNase/ATAC-seq data analyses, sequence reads mapped to

mitochondrial DNA (mtDNA) are usually discarded[4]. Lacking histones responsible for the

chromatin structure, mtDNA is similar to deproteinized naked DNA[31–33]. Therefore, we sought to

use the mtDNA reads from DNase/ATAC-seq data for cleavage bias estimation as an

alternative to using a naked DNA control sample (Fig. 3b). Likely because of relatively low read

counts and lack of sequence diversity (e.g., human mtDNA is only < 20kb long), the naïve k-mer

model exhibited poor performance on bias estimation from mtDNA reads, using bias scores

estimated from naked DNA as a reference (Fig. 3c). In contrast, SELMA generated a more

accurate bias estimation from the same mtDNA reads, which was highly consistent with the bias

scores estimated from the naked DNA data (Fig. 3d), demonstrating the power of SELMA to use

less input to make accurate bias estimations. We applied this approach to a series of ATAC-seq

and DNase-seq datasets for different human tissues from ENCODE and found that SELMA was

better than the naïve k-mer model in yielding a consistently higher correlation coefficient (>0.9)

between mtDNA-estimated bias and naked DNA-estimated bias for every ATAC-seq and

DNase-seq sample tested (Fig. S4a,b). Many optimized ATAC-seq protocols aimed to reduce

mtDNA reads to increase the fraction of chromatin DNA reads for chromatin accessibility signal

yield[29,30]. We sampled down mtDNA reads to test the performance of SELMA in making robust

bias estimations and found that SELMA could accurately estimate the bias with as few as

50,000 mtDNA reads (Fig. 3e, Fig. S4c), which is approximately 0.2% of the sequencing depth

of a regular ATAC-seq sample and lower than the fraction of mtDNA reads in any existing

ATAC-seq experiment[29,30]. These data demonstrated that by applying SELMA, mtDNA reads

can be utilized to substitute naked DNA control samples for accurate bias estimation.

Considering SELMA-estimated bias improves TF binding inference on ENCODE DNaseI

footprint regions

With an accurate bias estimation model developed, we next sought to examine the potential

effect of intrinsic biases on TF binding footprints in chromatin accessibility profiling data.

Focusing on the 4,465,728 human DNaseI consensus footprints across the human genome, we

plotted the DNaseI cleavages from different human cell lines and observed similar classic

DNaseI cleavage protection ("footprint") patterns across these footprint regions (Fig. 4a, Fig.

S5a). Interestingly, we also observed a recurrent pattern of the SELMA-estimated DNaseI

cleavage bias that is well aligned with the DNaseI cleavage pattern across these footprint

regions (Fig. 4b, Fig. S5b). In the aggregate view of footprint regions of different lengths, the

DNaseI cleavage signals exhibited a clear "cliff-bound valley"-shaped footprint pattern (Fig.

S5c). The DNaseI cleavage bias scores exhibited a pattern of two spikes located inside the

footprint coordinates (Fig. S5d). After normalizing various footprint lengths, we plotted the

aggregate DNaseI cleavage patterns across all consensus footprint regions and found that the

overall "footprint pattern" clearly included the double spike pattern of cleavage biases (Fig. 4c),

which aligned well with the two spikes in the aggregate bias score pattern (Fig. 4d), indicating a

possible contribution of intrinsic biases to the DNaseI footprinting.

To assess the interference of intrinsic biases with DNaseI cleavage patterns at TF binding

footprints, we aligned the 4.5 million consensus footprints with more than 10,000 human TF

ChIP-seq datasets from the Cistrome Data Browser database[34,35] and collected two sets of

footprint regions: "TF binding hotspots", the footprint regions overlapping with TF binding sites

detected from more than 3,000 ChIP-seq datasets, and "TF binding deserts", the footprint

regions that do not overlap with any TF binding sites from any ChIP-seq dataset or any human

TF motif sequence from the HOCOMOCO database[36] (Fig. S5e). We compared the aggregate

DNaseI cleavage patterns and the bias score patterns in these two sets of consensus footprint

regions and found that the DNaseI cleavage pattern in TF binding hotspot regions was

dominated by the classic DNaseI footprint pattern with little interference from the bias (Fig. 4e,f),

while in the TF binding desert regions, the entire cleavage "footprint" pattern was apparently

determined by the two spikes from the intrinsic bias (Fig. 4g,h). These results suggested that

although the overall DNaseI cleavage pattern is indicative of TF binding, the intrinsic cleavage

bias may interfere with the real footprint pattern, and the effect on those footprint regions with

rare TF binding events can be drastic. These observations were reproducible in DNase-seq

data from different cell and tissue types (Fig. S6a-b).

To quantify the level of intrinsic cleavage biases in a consensus footprint region, we designed a

footprint bias score (FBS), namely, the relative SELMA-estimated bias score at the two spikes

compared to the average SELMA bias score across the rest of the region for each footprint.

Consistent with what we observed in the aggregate view, the FBSs for TF binding deserts were

significantly higher while the FBSs for TF binding hotspots were significantly lower than the

background of all consensus footprints (Fig. 4i, $p < 10^{-5}$, by Wilcoxon test), indicating that FBS

might be an informative feature of the consensus footprint regions and might help separate true

TF binding footprints from false-positive DNaseI footprint patterns. Next, we used a model-

based approach to assess the potential power of SELMA-derived FBS in boosting the

performance of TF binding inference from DNase-seq signal patterns on consensus footprint

regions containing the TF motif sequence. For every TF with both the motif and ChIP-seq data

available, we employed a multinomial logistic regression model using different sets of features,

each of which may include DNase-seq read count, DNaseI footprint score produced from an

existing method, and an optional FBS, to infer the TF binding occupancy (determined by peak

occurrence in ChIP-seq data) in the motif-containing footprint regions and compared the

inference performances. We included our previous footprint method[14] and several

representative methods that outperform others, Wellington[9] and HINT[11,19] (with and without bias

correction mode) in this comparison. For each footprint method, we compared the TF binding

inference performance of the original method (read count + footprint score), the original method

plus a randomized naïve k-mer bias score as a control, and the original method plus the SELMA

bias score (FBS), using an integrated rank score strategy. For a total of 48 TFs in the human

cell line GM12878 and 45 TFs in K562, the models with SELMA FBS significantly outperformed

those without SELMA FBS, regardless of the approach used originally to calculate the footprint

score (Fig. 4j, Fig. S6). These results suggested that considering the intrinsic cleavage bias

could always improve the performance of existing footprint computational methods for TF

binding inference.

To assess whether the intrinsic bias has different levels of interference with the footprint regions

for different TFs, we selected two subsets of sequence motif-containing footprint regions for

each TF as the top 10% with the highest FBS or the bottom 10% with the lowest FBS, and

compared the frequencies of observing actual TF binding (overlapping with a ChIP-seq peak) in

these subsets of footprints. We found that different TFs had various preferences for binding at

low-FBS footprints or high-FBS footprints. For most TF samples in the two cell lines, the

footprint regions with a low FBS showed a significantly higher chance of observing actual TF

binding than those with a high FBS (Fig. 4k). For example, HLH family TFs such as MYC and

MAX[17] showed a higher fraction of TF binding occupancies in low-FBS footprint regions than at

high-FBS regions. Zinc finger family TFs including CTCF, REST, and NRF1[6,17], showed a higher

chance of TF binding in high-FBS footprint regions than in low-FBS regions (Fig. S6d). This is

possibly because of the different DNA binding affinities and DNA residence times between

different TFs[15]. TFs with stronger DNA-binding affinity or longer residence time might be less

affected by intrinsic cleavage biases, while TFs with shorter residence time or weaker binding

affinity might be more severely affected by intrinsic biases. These results suggested that

intrinsic cleavage biases might affect different TFs at various levels in divergent directions, and

considering intrinsic bias should improve the footprint-based TF binding inference for most TFs.

SELMA improves the accuracy of cell clustering from single-cell ATAC-seq data

Single-cell ATAC-seq (scATAC-seq) technologies enable the detection of accessible chromatin

regions at single-cell resolution in thousands of cells at a time[21–23]. Due to the scarcity of

cleavage events in an individual cell, most chromatin accessibility regions in a single cell have

only one aligned fragment, making the potential influence of intrinsic cleavage biases more

substantial in scATAC-seq data than in bulk ATAC-seq data. We collected three scATAC-seq

datasets generated using two different platforms, human hematopoietic cells[37], human cell

lines[21], and a mouse primitive gut tube sample, and found that the scATAC-seq data contained

similar intrinsic cleavage biases to bulk data with highly correlated bias scores estimated by

SELMA (Fig. S7a-c). We estimated the average cleavage bias for each individual cell (cell bias

score, CBS) and found that the distribution of CBS was different across cell cluster patterns,

batches, and annotated cell types (Fig. 5a-c, Fig. S7d-i). These observations indicated that,

regardless of experimental platforms and biological systems, the Tn5 intrinsic cleavage biases

can contribute to cellular heterogeneity observed from scATAC-seq data and can affect cell

clustering analysis.

We next assessed whether considering intrinsic cleavage biases can improve cell clustering

based on scATAC-seq data. We used the actual cell-type labels as the clustering ground truth

for the human hematopoietic cell sample and the mixed human cell line sample and used

scRNA-seq data-projected cell labels as a "pseudo" ground truth for the mouse primitive gut

tube sample. We used the adjusted Rand index[38] to quantify the accuracy of an unsupervised

clustering result. As scATAC-seq-based cell clustering can be performed on all chromatin

accessibility regions (ATAC-seq peaks), we sought to address whether removing peaks with

high intrinsic biases can increase the clustering accuracy. We first tested a simple clustering

approach that involved dimensionality reduction using principal component analysis (PCA)

followed by K-means clustering. For all three scATAC-seq datasets, cell clustering using

SELMA-debiased peaks only (excluding 1%-50% of peaks with the highest PBS) is significantly

more accurate than clustering using all peaks under the background of randomly excluding the

same number of peaks (Fig. 5d-f). Then, we tested several representative scATAC-seq analysis

tools for cell clustering, including APEC[39], Seurat[40], scran[41], and snapATAC[42], and found that

using SELMA-debiased peaks can significantly improve the clustering results in most cases

(Fig. 5g-j, Fig. S7j-r). These results suggested that scATAC-seq data contain intrinsic cleavage

biases that can be reduced by excluding SELMA-identified high-bias peaks from the cell

clustering analysis.

Discussion

The existence of enzymatic cleavage biases in DNase-seq and ATAC-seq experiments has

been widely acknowledged in the field, but to what extent such intrinsic biases affect data

analysis has not been systematically assessed. SELMA provides a quantitative approach for the

324 accurate and robust estimation of intrinsic cleavage biases in both bulk and single-cell

325 chromatin accessibility sequencing data and requires fewer sequence reads than the naïve k-

326 mer model. Taking Tn5 dimerization into consideration, SELMA yields more accurate bias

327 estimation specifically for ATAC-seq data by including k-mer sequences at two Tn5 cleavages

328 9bp apart. SELMA can improve functional analysis and interpretation of chromatin accessibility

329 profiles. On bulk DNase-seq data, we showed that considering SELMA-estimated biases can

330 improve TF binding inference from ENCODE DNaseI consensus footprints, with superior

331 performances compared to existing tools, including some that already considered "biases". On

332 single-cell ATAC-seq data, we demonstrated that selecting chromatin accessibility regions

333 based on the SELMA-estimated bias level can greatly increase the accuracy of cell clustering by

334 several existing tools and hence can help researchers obtain more biological insights from the

335 data.

337 SELMA is built on top of the widely used k-mer model by combining simplex encoding and a

338 linear model. Simplex encoding has the unique ability to capture the pairwise symmetry and

339 orthogonality between mononucleotides and interactions within each dinucleotide. It significantly

340 reduces the degrees of freedom without losing any variance information compared to the naïve

341 k-mer model. These properties enable SELMA to make robust estimations from fewer reads or

342 smaller datasets than are required by other approaches, hence enabling *de novo* bias

343 estimation from a smaller DNA molecule, such as mtDNA in a DNase/ATAC-seq sample,

344 without having to generate a separate genomic naked DNA sample. However, SELMA still relies

345 on sufficient read counts for each k-mer for an accurate estimation. Although the performance of

346 SELMA may increase with k, this effect is not unlimited. The performance using 12-mer is not as

347 good as using 10-mer possibly because there are not enough reads in the dataset for many 12-

348 mers. Nevertheless, SELMA works well for most existing DNase/ATAC-seq datasets tested in

349 our study. In addition, the feasibility and superior performance of simplex encoding have also

been demonstrated for TF motif characterization[43]. Following similar encoding strategies,

SELMA can potentially be applied to any k-mer-based model or to any high-throughput

sequencing data for robust sequence bias estimation and pattern recognition.

The k-mer biases estimated by SELMA are consistent across species and cell types, as

expected based on the assumption that the cleavage biases are intrinsic features of the

enzymatic assay. Therefore, one can directly use the SELMA-estimated k-mer biases from

DNaseI or Tn5-digested naked DNA data as universal intrinsic biases for DNase-seq and

ATAC-seq, respectively, and incorporate these precalculated bias scores into the data analysis.

However, this bias dataset is not guaranteed to remain accurate in many species that have not

been profiled, as there might be unknown biases that have not been captured. Although we are

confident that the SELMA-estimated results should still be largely consistent, one can always

use the SELMA package for *de novo* estimation of cleavage biases from one's own datasets.

When applying SELMA-estimated FBS to correct biases in DNase footprints, our data were

limited to ENCODE DNaseI consensus footprints as a proof-of-principle study. While this

ENCODE dataset is thus far the largest DNase footprint repertoire, users might be interested in

de novo detection of footprints from their customized DNase/ATAC-seq data. As many

bioinformatics tools are already available for such tasks using various computational models[20],

SELMA or SELMA-generated bias information can be incorporated into each of those models

for improved performance for more accurate regulatory DNA identification from footprints.

Similarly, while we show significant improvement of scATAC-seq cell clustering by removing

peaks with high PBS under the framework of most existing tools, performance still varies across

these tools due to the different statistical and computational models used in these methods.

Further optimization of existing methods or development of new methods is expected for the

integration of SELMA for the systematic removal of intrinsic biases.

Methods

High-throughput sequencing data collection and processing

Data collection. Publicly available ATAC-seq, single-cell ATAC-seq, DNase-esq and ChIP-seq

data used in this study were collected from Gene Expression Omnibus (GEO) and the ENCODE

consortium. The metadata and accession numbers are listed in Table S1.

Processing of bulk ATAC-seq and DNase-seq data. Raw sequencing reads were aligned to

the GRCh38 (hg38) reference genome with bowtie2 (v2.2.9) (-X 2000 for paired end data)[44].

Low-quality reads (MAPQ < 30) were discarded. For paired-end sequencing data, reads with

two ends aligned to different chromosomes (chimeric reads) were also discarded. For paired-

end data, reads with identical 5' end positions for both ends were regarded as redundant reads

and discarded. The nonredundant reads were separated into chromosomal DNA (chromatin

reads) and mitochondrial DNA (mtDNA reads) based on their genomic location. Peak detection

was performed on the nonredundant chromatin reads using MACS2 (v2.1.2) (-q 0.01, --extsize

50) and +/- 200bp centered on the peak summits was collected as the genome-wide chromatin

accessible regions. The accessible regions in each dataset were separated into 5 groups from

highest accessibility to lowest accessibility based on the read count on each peak (for Fig. S1).

The 5' end nucleotides of each read were piled up to generate the genome-wide observed

cleavage profile.

Processing of single-cell ATAC-seq data. For the human HSC sample, raw sequencing reads

were aligned to the GRCh38 (hg38) reference genome with bowtie2 (-X 2000). Low quality

reads (MAPQ < 30), chimeric reads and duplicate reads for each individual cell were discarded.

For the mouse gut tube sample, scATAC-seq data from the 10x Genomics platform were

preprocessed with Cell Ranger ATAC with the default parameters to generate fragments for

each individual cell. The fragment file from the standard Cell Ranger ATAC output was used as

input for the subsequent analysis. Because the Cell Ranger ATAC pipeline shifted from the Tn5

cleavage sites to +4/-5 bp in generating the fragment file, the coordinates were shifted back to

represent the actual cleavage loci. For both datasets, cells with more than 10,000 reads were

retained for analysis.

Processing of ChIP-seq data. ChIP-seq data were collected and processed as described in a

previous study[45]. In brief, based on the quality measurements from the Cistrome data browser

(CistromeDB)[34,35], we included only the ChIP-seq datasets that pass at least four out of the first

five QC measures (i.e., sequence quality, mapping quality, library complexity, ChIP enrichment,

and signal-to-noise ratio). We also excluded the ChIP-seq datasets with fewer than 1000 peaks

identified by MACS2[46] with an FDR cutoff of 0.01. For TFs with multiple datasets available in the

same cell type, we kept only one dataset with the best quality based on passing most of the 6

CistromeDB QC measures, including sequence quality, mapping quality, library complexity,

ChIP enrichment, signal-to-noise ratio, and regulatory region enrichment. The ChIP-seq peak

files (detected by MACS2) for the selected samples were downloaded from CistromeDB for the

following analysis.

TF motif analysis. The motifs of human TFs were collected from the HOCOMOCO database

(v11) [36], and the genome-wide motif sites of TFs were detected by FIMO (v4.12.0) in the MEME

package[47]. Motif sites located outside of the genome-wide 36bp unique mappable regions were

excluded from the analysis. In total, 25,027,116 motif sites for 66 TF motifs from the human

genome were included in the analysis.

SELMA model

Naïve k-mer bias model for intrinsic cleavage bias estimation. The naïve k-mer biases were

calculated as described in a previous study[14]. In brief, a naïve k-mer bias was estimated as the

observed frequency of the cleavages with the k-mer relative to the frequency of that k-mer in the

background. For each mapped sequence read in a DNase-seq or ATAC-seq dataset, the

enzymatic cleavage site was between genomic positions i and i-1 for the plus (+) strand reads

and between i and i+1 for the minus (–) strand reads, where i represents the genomic position

of the 5' nucleotide of the reads. The associated k-mer sequence was thus assigned as

$\left[i - \frac{k}{2}, i + \frac{k}{2} - 1\right]$ for the plus strand reads and $\left[i - \frac{k}{2} + 1, i + \frac{k}{2}\right]$ for the minus strand reads. The

naïve k-mer bias score for k-mer j is defined as the number of all observed cleavages with that

k-mer divided by the occurrences of that k-mer in the background:

$$S_j = \frac{N_j}{M_j}$$

where N_j is the count of cleavages with k-mer j, and M_j is the total count of occurrences of k-

mer j in the background (Fig. 1a). In the case of chromatin DNA, this background included 400

bp from each chromatin accessible region centered at the peak summit detected by MACS2.

The background in the naked DNA samples included genome-wide 36bp unique mappable

regions. The background in mtDNA included the mitochondrial DNA sequence. The median was

further subtracted from the bias scores for visualization in the scatter plots (e.g., Fig. 1d-k).

In the naïve k-mer model, the bias score for each k-mer is independent and empirically

determined from the data. The number of independent variables in the model is the total number

of k-mers, i.e., 4^k.

SELMA model for intrinsic cleavage bias estimation. In SELMA, a simplex encoding model

followed by a linear model was employed on top of the naïve k-mer model to better estimate the

intrinsic cleavage biases for each k-mer. To efficiently encode a k-mer sequence considering

their intrinsic similarities, a simplex encoding model was adopted from previous studies[25,26], in

which the DNA nucleotides were encoded as vectors representing the four tetrahedral vertices

of a regular 0-centered simplex (Fig. 1b).

$$
\begin{aligned}
A &= [\quad 1 \quad -1 \quad -1 \quad] \\
C &= [\quad -1 \quad 1 \quad -1 \quad] \\
G &= [\quad -1 \quad -1 \quad 1 \quad] \\
T &= [\quad 1 \quad 1 \quad 1 \quad]
\end{aligned}
$$

459   In the simplex encoding, the vectors representing the four nucleotides are of equal length,

460   mutually orthogonal, and equidistant from each other. To account for interactions between

461   adjacent nucleotides, a dinucleotide was additionally encoded as the outer product of the two

462   vectors associated with the two nucleotides:

$$
\begin{aligned}
AA &= [\quad 1 \quad -1 \quad -1 \quad -1 \quad 1 \quad 1 \quad -1 \quad 1 \quad 1 \quad] \\
AC &= [\quad -1 \quad 1 \quad -1 \quad 1 \quad -1 \quad 1 \quad 1 \quad -1 \quad 1 \quad] \\
AG &= [\quad -1 \quad -1 \quad 1 \quad 1 \quad 1 \quad -1 \quad 1 \quad 1 \quad -1 \quad] \\
AT &= [\quad 1 \quad -1 \quad -1 \quad -1 \quad 1 \quad 1 \quad -1 \quad 1 \quad 1 \quad] \\
CA &= [\quad -1 \quad 1 \quad 1 \quad 1 \quad -1 \quad -1 \quad -1 \quad 1 \quad 1 \quad] \\
CC &= [\quad 1 \quad -1 \quad 1 \quad -1 \quad 1 \quad -1 \quad 1 \quad -1 \quad 1 \quad] \\
CG &= [\quad 1 \quad 1 \quad -1 \quad -1 \quad -1 \quad 1 \quad 1 \quad 1 \quad -1 \quad] \\
CT &= [\quad -1 \quad -1 \quad -1 \quad 1 \quad 1 \quad 1 \quad -1 \quad -1 \quad -1 \quad] \\
GA &= [\quad -1 \quad 1 \quad 1 \quad -1 \quad 1 \quad 1 \quad 1 \quad -1 \quad -1 \quad] \\
GC &= [\quad 1 \quad -1 \quad 1 \quad 1 \quad -1 \quad 1 \quad -1 \quad 1 \quad -1 \quad] \\
GG &= [\quad 1 \quad 1 \quad -1 \quad 1 \quad 1 \quad -1 \quad -1 \quad -1 \quad 1 \quad] \\
GT &= [\quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad 1 \quad 1 \quad 1 \quad] \\
TA &= [\quad 1 \quad -1 \quad -1 \quad 1 \quad -1 \quad -1 \quad 1 \quad -1 \quad -1 \quad] \\
TC &= [\quad -1 \quad 1 \quad -1 \quad -1 \quad 1 \quad -1 \quad -1 \quad 1 \quad -1 \quad] \\
TG &= [\quad -1 \quad -1 \quad 1 \quad -1 \quad -1 \quad 1 \quad -1 \quad -1 \quad 1 \quad] \\
TT &= [\quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad]
\end{aligned}
$$

One can show that these vectors for dinucleotide interactions are also of equal length, mutually

orthogonal, and equidistant from each other. In fact, in the simplex encoding, mononucleotides

and dinucleotides were encoded as rows in a Hadamard matrix of order 4 and a Hadamard

matrix of order 16, respectively.

Considering both mononucleotides and dinucleotide interactions, a k-mer can then be encoded

as k mononucleotides and $k-1$ dinucleotides. Therefore, the dimensionality of a k-mer simplex

encoding is

$$p(k) = 1 + 3k + 9(k-1) = 12k - 8$$
In SELMA, the intrinsic k-mer biases were then estimated in a linear model with these $12k-8$

parameters using the observed naïve k-mer bias scores (Fig. 1c). In detail, we had the formula:

$$y \sim \sum_{i=1}^{12k-8} a_i x_i \qquad (1)$$

where each observation y is the naïve k-mer bias score; $x_i \in \{1, -1\}$, is the independent

variable in the simplex encoding vectors; and a_i is the parameter to be estimated. After linear

regression, the model-fitted value $\hat{y}$ was defined as the SELMA bias score for each k-mer.

Genome-wide cross-correlation analysis. For the analysis presented in Fig. 2b,c, reads from

plus (+) and minus (-) strands on chromatin accessible regions (peaks) were collected

separately to generate plus strand observed cleavage profile and minus strand observed

cleavage profile, respectively. The Pearson correlation coefficient between the plus strand

signal and the k-bp shifted minus strand signal is:

$$\rho_k = \frac{\sum_i (P_i - \bar{P})(M_{ik} - \overline{M_k})}{\sqrt{\sum_i (P_i - \bar{P})^2 \sum_i (M_{ik} - \overline{M_k})^2}} \qquad (2)$$

where P_i is the log-scaled plus strand cleavage count at genomic position i (log2(cleavage+1)),

M_{ik} is the log-scaled minus strand cleavage count at genomic position i with a k-bp shift, $\underline{P}$ is

the mean of all the P_i, and $\overline{M_k}$ is the mean of M_{ik} for all i. All genomic positions on the genome-

wide DNase/ATAC-seq peaks were included in the analysis. k is chosen from 1 to 20 (x-axis in

Fig. 2b,c).

Comparison of different bias estimation methods. We use a_i^+ and a_i^- to denote the "5' only"

intrinsic sequence bias score at genomic position i on the plus strand and minus strand,

respectively. We use a'^+_i and a'^-_i to denote the bias score from other bias estimation methods in

this section. Different bias estimation methods used in Fig. 2 and its associated sections are

listed below:

● For the "5' only" method, a_i^+ was calculated based on the k-mer ratio associated with the

k-mer spanning positions $[i - k/2, \ i + k/2 - 1]$ on the plus strand, and a_i^- was

calculated in a similar way based on the nucleotides spanning genomic positions $[i -$

$k/2 + 1, \ i + k/2]$ on the minus strand (reverse complement of the DNA sequence on the

corresponding plus strand). This method was applied to both DNase-seq and ATAC-seq,

while the other methods were applied only to ATAC-seq as they were specifically

designed for ATAC-seq.

● For SELMA (Fig. 2a), the bias score was calculated as the geometric mean of the "5'

only" bias score at the given position and the "5' only" bias score at 9bp downstream of

the other strand, i.e., $a'^+_i = \sqrt{a_i^+ \times a_{i+9}^-}$, and $a'^-_i = \sqrt{a_i^- \times a_{i-9}^+}$

● For the model in Martins et al[16], the bias score at genomic position i was calculated in a

similar way to the "5' only" method but using a gapped 11-mer model. The model could

be represented as $XXXXXXNNNXNXCXXNNXNNNNXNX$, in which position i was

represented by C; positions that were ignored were represented by X and informative

positions were represented by N.

● For the model in Baek et al[18], the bias score at genomic position i was calculated in a

similar way to the "5' only" method but the cleavages were shifted +4/-5 bp for +/- strand

cleavages. In practice, following the description in the "bagfoot" package, the bias score

at position i was calculated as $a'^+_i = a_{i+5}^+$, and $a'^-_i = a_{i-5}^+$.

•   For the model in Calviello et al[8], the bias score at genomic position i was calculated in a

similar way to the "5' only" method but using the 6-mer bias table provided in the study.

Calculation of observed and bias-expected cleavage. The 1bp at 5' end positions for DNase-

seq or ATAC-seq reads were piled up to generate the genome-wide observed cleavage profile.

To generate the bias-expected cleavage profile, for a 50-bp window centered on nucleotide i,

we placed the same number of observed cleavages following the sequence bias contribution in

this window. Let $\widehat{N_i^s}$ represent the bias expected cleavage at position i on strand $s \in \{+, -\}$, N_i^s

represent the observed cleavage at position i on strand s, and a_i^s denote the intrinsic cleavage

bias (estimated with any of the above methods) at position i on strand s. Then we have

$$\widehat{N_i^s} = N_i^s y_i^s \tag{3}$$

$$y_i^s = \frac{2^{a_i^s}}{\sum_{j=i-25}^{i+24} 2^{a_j^s}} \tag{4}$$

We used the Pearson correlation coefficient to compare the observed cleavage profile and the

bias-predicted cleavage profile (Fig. 2). We only considered positions within the accessible

regions to ensure that all positions had sufficient reads in the 50-bp window for accurate

estimation.

DNaseI footprint analysis

Data processing. The genome-wide DNaseI consensus footprint regions were downloaded

from Reference[7]

(https://resources.altius.org/~jvierstra/projects/footprinting.2020/consensus.index/). The

observed DNaseI cleavage profile from a DNase-seq dataset and DNaseI SELMA bias scores

across +/- 50bp centered on the footprint region were plotted as heatmaps (Fig. 4a,b). The

footprint regions were ordered by the footprint lengths, and each 1000 footprint regions with

similar lengths were compressed as one row in the heatmap for better visualization. The +/-

strand signals were plotted separately. We aligned the footprint regions based on the two

observed bias spikes in each footprint region (located 7bp to the right of the left boundary and

7bp to the left of the right boundary of the footprint, labeled as -0 and +0 in Fig. 4c-h). The

center regions between the bias spikes were scaled to 4 bins to align footprint regions with

different lengths.

Footprint bias score. The footprint bias score (FBS, in Fig. 4i-k) was defined as the difference

between "spike bias" and "center bias". The "spike bias" was calculated as the average of the

two SELMA bias scores at the spike positions, while the "center bias" was calculated as the

median SELMA bias score at the rest of the positions in the footprint region. Let FBS_j, b_j, and c_j

denote the footprint bias score, spike bias and center bias of footprint j, respectively. The FBS,

spike bias, and center bias are given by:

$$FBS_j = b_j - c_j \tag{5}$$

$$b_j = \overline{a_{\iota \in B}^{s \in \{+,-\}}} \tag{6}$$

$$c_j = \widetilde{a_{\iota \in C}^{s \in \{+,-\}}} \tag{7}$$

where a_i^s represents the SELMA bias score at genomic position i on the strand $s \in \{+, -\}$,

overbar represents the average, tilde represents the median, and B and C represent spike

positions (within the flanking 1bp of the bias spikes) and the remaining positions of the footprint

j, respectively. To calculate the randomly shuffled FBS (random k-mer bias), the FBS was

calculated in the same way, but the bias score a_i^s for each position was randomly selected from

the SELMA bias score table.

Calculation of footprint score with different methods. The following methods were used to

calculate footprint scores for comparison:

•   Raw footprint: The raw footprint score was calculated following a previous study[14], using

the formula $f = -\left(\log\frac{n_C+1}{n_R+1} + \log\frac{n_C+1}{n_L+1}\right)$, where n_C, n_R and n_L denote the DNase

cleavage count in the motif region, and the flanking regions to the right and left of the

motif, respectively. The flanks are both the same length as the motif.

•   Wellington footprint: We used Wellington[9] (v0.2.0) with default parameters to detect

genome-wide footprints and selected significant footprint regions with p-values < 1e-10.

The output footprint score was assigned to the TF motif overlapping with the footprint

region.

•   HINT footprint: We used HINT[11,19] (v0.12.3) with default parameters to detect genome-

wide footprints. For bias correction mode, we used an additional parameter: --bias-

correction. The footprint score of each footprint region was assigned to the overlapping

TF motifs.

Inference of TF binding with different features. For each TF in Fig. 4j, the TF motifs

overlapping with consensus footprint regions were collected as target regions. DNase-seq read

count in the footprint region ("reads"), footprint score from an existing method, and bias score

were used as features in a multinomial logistic regression model to infer TF occupancy at

footprint-overlapping motif regions. For each available method, model training with cross-

validation and predictions were performed using different combination of features: "original

method" refers to reads + footprint score. An additional feature of either SELMA FBS or a

random k-mer bias was added for different models. We used a performance measure

integration approach[11] to evaluate different models' prediction power. For each model, we

calculated the area under the ROC curve (AUROC) at 100%, 10%, and 1% false positive rate

(FPR). We also calculated the area under the precision-recall curve (AUPRC) at 100%, 10%,

and 1% recall. We then combined these six performance measures as a rank score *S*, defined

as

$$S = \frac{1}{6}\sum_i -\log\frac{r_i}{N+1}$$

where $i = 1, \ldots, 6$ denotes the 6 performance measures, r_i is the rank of a model for

performance measure i, and N is the total number of models.

To calculate the random k-mer bias, we randomly permuted the SELMA k-mer bias table and

generated the k-mer bias table for a "simulated enzyme". We used this "simulated" bias table to

calculate the FBS for each footprint region and performed TF inference modeling similar to what

we did for the DNaseI SELMA FBS. This permutation was performed 100 times to generate 100

performance rank scores for random k-mer bias used as controls.

Motif prediction power of TF binding sites. In Fig. 4k, we collected the genome-wide motif

sites overlapping with consensus footprint regions and the ChIP-seq peaks for each TF with

HOCOMOCO[36] motif and ChIP-seq data available in GM12878 or K562 (Table S1). As a result,

we collected genome-wide motif sites for 65 TFs (48 in GM12878 and 45 in K562), and

35,721,822 motif sites in total. For each TF, we selected two subgroups of its motif sites based

on the FBS of their overlapped footprint regions: the top 10% of motif sites with the highest FBS,

defined as "sites with high-bias footprint"; and the bottom 10% of motif sites with the lowest

FBS, defined as "sites with low-bias footprint". We calculated the proportion of motif sites

overlapping with TF ChIP-seq peaks for each of the two subgroups and plotted on a scatter plot

(Fig. 4k). To test whether low-bias footprints tend to have more TF binding than high-bias

footprints for most TFs, we conducted a t-test comparing the distribution of TF binding log

likelihood ratios of low-bias over high-bias footprints to the standard normal distribution, and the

test p-value is labeled in Fig. 4k.

Single-cell ATAC-seq clustering analysis

Gold standard of single-cell clustering. For single-cell ATAC-seq data in the human

hematopoietic cells and human cell line samples, the cell-type information for each individual

cell was used as the ground truth, or the gold standard. For single-cell ATAC-seq data in the

mouse gut tube sample, the cell-type information was assigned based on label transfer[40] from

the single-cell RNA-seq dataset in the same system[48] (GSE136689), as the "pseudo" ground

truth, or the silver standard. In detail, we integrated scRNA-seq and scATAC-seq data using the

ArchR package[49] (v1.0.1). Individual cells with a high RNA integration score (unconstrained

predicted score < 0.56) were collected as high-quality cells for the analysis. The cutoff of the

RNA integration score (0.56) was determined by maximizing the interclass variance in the RNA

integration score using Otsu's method[50]. Cell types represented by fewer than 10 cells were

discarded.

Average bias for individual cells. The average bias for each individual cell (median SELMA

cell bias score, median CBS, used in Fig. 5) was calculated as the median of the bias for all the

fragments in the individual cell. The bias for each paired-end fragment was calculated as the

mean of the SELMA bias scores for the 5' end and the 3' end.

Peak detection and peak bias score calculation. We first combined all the single-cell ATAC-

seq reads in the dataset and performed MACS2[46] peak calling with a q-value cutoff of 0.1 to

include all the potential accessible regions in the human hematopoietic cell and human cell line

systems (100,456 peaks detected for human hematopoietic cells and 83,318 peaks for human

cell lines). For the 10X scATAC-seq data in mouse gut tube, the q-value cutoff of 0.01 was used

for peak calling (146,098 peaks detected). Potential accessible regions with fewer than 10

covered reads or more than 4,000 covered reads were discarded in the subsequent analysis. To

641 consider the effect of intrinsic cleavage bias in scATAC-seq data, we summarized the bias for

642 each potential accessible region by calculating the peak bias score (PBS), defined as the

643 median SELMA bias score of all reads (from all individual cells) located in the peak region.

645 **Cell clustering with scATAC-seq data.** In the scATAC-seq clustering analysis, as a naïve

646 method, we performed PCA dimensionality reduction on the accessible regions by individual

647 cells matrix of normalized read count and kept 60 PCs, followed by k-means clustering. The

648 number of clusters (k) in k-means clustering was determined as the actual number of cell types

649 in the dataset. To evaluate the performance of cell clustering, we compared the clustering

650 results with the gold standard using adjusted rand index (ARI)[38]. We further repeated the k-

651 means clustering with 100 different random seeds to ensure the stability of the clustering results

652 and the average ARI of the 100 runs was used to estimate the clustering accuracy. We also

653 applied several published methods, including APEC[39], Seurat[40], scran[41], and snapATAC[42] with

654 default parameters, on the same dataset. We selected the top 60 dimensions for those methods

655 at the dimensional reduction step (PCA, Seurat, scran and snapATAC) to keep more variance

656 potentially introduced by intrinsic biases. We used the ArchR package to practice the Seurat

657 and scran clustering methods. To explore the effect of intrinsic cleavage bias on scATAC-seq

658 analysis, we selected different percentages (from 50% to 99%, with a 1% increment) of peaks

659 with the lowest PBS (SELMA debiased peaks in Fig. 5) as input to perform cell clustering. We

660 also randomly selected the same percentage of peaks as a control to estimate the relative rank

661 of the clustering performance from using all peaks and debiased peaks. In detail, for each

662 percentage from 50% to 99%, we randomly sampled peaks 100 times and defined the relative

663 rank as the number of ARIs from random samples that were less than the ARI from the same

664 percentage of debiased peaks or all peaks. The relative ranks for different percentages from

665 50% to 99% are summarized and plotted as boxplots in Fig. 5.

Code and data availability

The SELMA package is available at https://github.com/zang-lab/SELMA. User instructions and

example data files can be found in the README document. Essential annotation data, analysis

results, and scripts are also available in the repository. The mouse gut tube scATAC-seq

dataset has been deposited in the Gene Expression Omnibus (GEO) with accession number

GSE168373. Publicly available data used in this study can be found in the GEO and the

ENCODE project. Accession numbers for all data used in the study are listed in Additional file 2:

Table S1.

Acknowledgements

The authors thank Dr. Tae-Hee Kim for sharing the mouse primitive gut tube scATAC-seq data

and Dr. Xiaole Shirley Liu for helpful discussions. This work was supported by US National

Institutes of Health (NIH) grants R35GM133712 and K22CA204439 (to C.Z.), R35GM128635 (to

M.J.G.), National Science Foundation grant NSF-2048991 (to T.Z.), UVA Cancer Center Farrow

Fellowship and the NCI Cancer Center Support Grant P30 CA44579 (to S.S.H.).

Author contributions

C.Z. conceived and directed the research. S.S.H., C.A.M., and C.Z. designed the method. L.L.,

M.J.G., K.D., and T.Z. contributed to the method design. S.S.H. implemented the method and

analyzed the data. Q.L. and W.M. helped the data analysis. S.S.H. and C.Z. wrote the

manuscript. All authors approved the manuscript.

Competing interests

The authors declare that they have no competing interests.

References

1. McGhee, J. D., Wood, W. I., Dolan, M., Engel, J. D. & Felsenfeld, G. A 200 base pair region

at the 5′ end of the chicken adult β-globin gene is accessible to nuclease digestion. *Cell* **27**, 45–

55 (1981).

2. Keene, M. A., Corces, V., Lowenhaupt, K. & Elgin, S. C. DNase I hypersensitive sites in

Drosophila chromatin occur at the 5' ends of regions of transcription. *Proc National Acad Sci* **78**,

143–146 (1981).

3. Boyle, A. P. *et al.* High-Resolution Mapping and Characterization of Open Chromatin across

the Genome. *Cell* **132**, 311–322 (2008).

4. Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J. Transposition of

native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding

proteins and nucleosome position. *Nature Methods* **10**, 1213–1218 (2013).

5. Hesselberth, J. R. et al. Global mapping of protein-DNA interactions in vivo by digital genomic

footprinting. Nature Methods **6**, 283–289 (2009).

6. Neph, S. et al. An expansive human regulatory lexicon encoded in transcription factor

footprints. Nature **489**, 83–90 (2012).

7. Vierstra, J. et al. Global reference mapping of human transcription factor footprints. Nature

583, 729–736 (2020).

8. Calviello, A. K., Hirsekorn, A., Wurmus, R., Yusuf, D. & Ohler, U. Reproducible inference of

transcription factor footprints in ATAC-seq and DNase-seq datasets using protocol-specific bias

modeling. *Genome Biol* **20**, 42 (2019).

9. Piper, J. et al. Wellington: a novel method for the accurate identification of digital genomic

footprints from DNase-seq data. *Nucleic Acids Research* **41**, e201–e201 (2013).

10. Sherwood, R. I. et al. Discovery of directional and nondirectional pioneer transcription

factors by modeling DNase profile magnitude and shape. *Nat Biotechnol* **32**, **17**1–178 (20**14**).

11. Li, Z. et al. Identification of transcription factor binding sites using ATAC-seq. Genome Biol

20, **45** (20**19**).

12. Bentsen, M. et al. ATAC-seq footprinting unravels kinetics of transcription factor binding

during zygotic genome activation. *Nat Commun* **11**, **42**67 (2020).

13. Meyer, C. A. & Liu, X. S. Identifying and mitigating bias in next-generation sequencing

methods for chromatin biology. *Nat Rev Genet* **15**, 709–**721** (2014).

14. He, H. H. et al. Refined DNase-seq protocol and data analysis reveals intrinsic bias in

transcription factor footprint identification. *Nat Methods* **11**, 73–78 (2014).

15. Sung, M.-H., Guertin, M. J., Baek, S. & Hager, G. L. DNase Footprint Signatures Are

Dictated by Factor Dynamics and DNA Sequence. *Molecular Cell* **56**, 275–285 (2014).

16. Martins, A. L., Walavalkar, N. M., Anderson, W. D., Zang, C. & Guertin, M. J. Universal

correction of enzymatic sequence bias reveals molecular signatures of protein/DNA interactions.

Nucleic Acids Res **46**, gkx1053- (2017).

17. Yardımcı, G. G., Frank, C. L., Crawford, G. E. & Ohler, U. Explicit DNase sequence bias

modeling enables high-resolution transcription factor footprint detection. Nucleic Acids Research

42, 11865–11878 (2014).

18. Baek, S., Goldstein, I. & Hager, G. L. Bivariate Genomic Footprinting Detects Changes in

Transcription Factor Activity. *CellReports* **19**, 1710–1722 (2017).

19. Gusmao, E. G., Allhoff, M., Zenke, M. & Costa, I. G. Analysis of computational footprinting

methods for DNase sequencing experiments. Nature Methods 13, 303–309 (2016).

20. Yan, F., Powell, D. R., Curtis, D. J. & Wong, N. C. From reads to insight: a hitchhiker's guide

to ATAC-seq data analysis. Genome Biol 21, 22 (2020).

21. Buenrostro, J. D. et al. Single-cell chromatin accessibility reveals principles of regulatory

variation. Nature **523**, 486–490 (2015).

22. Cusanovich, D. A. et al. Multiplex single cell profiling of chromatin accessibility by

combinatorial cellular indexing. *Science* **348**, 910–914 (2015).

23. Lake, B. B. et al. Integrative single-cell analysis of transcriptional and epigenetic states in

the human adult brain. *Nature Biotechnology* **36**, 70–80 (2017).

24. Preissl, S. et al. Single-nucleus analysis of accessible chromatin in developing mouse

forebrain reveals cell-type-specific transcriptional regulation. *Nat Neurosci* **21**, 432–439 (2018).

25. Zhang, C.-T. & Zhang, R. Analysis of distribution of bases in the coding sequences by a

digrammatic technique. Nucleic Acids Res 19, 6313–6317 (1991).

26. Stormo, G. D. Maximally Efficient Modeling of DNA Sequence Motifs at All Levels of

Complexity. *Genetics* **187**, **12**19–1224 (2011).

27. Adey, A. et al. Rapid, low-input, low-bias construction of shotgun fragment libraries by high-

density in vitro transposition. Genome Biol 11, R119 (2010).

28. Sos, B. C. et al. Characterization of chromatin accessibility with a transposome

hypersensitive sites sequencing (THS-seq) assay. Genome Biol 17, 20 (2016).

29. Corces, M. R. et al. Lineage-specific and single-cell chromatin accessibility charts human

hematopoiesis and leukemia evolution. *Nature Genetics* **48**, 1193–1203 (2016).

30. Corces, M. R. et al. An improved ATAC-seq protocol reduces background and enables

interrogation of frozen tissues. Nature Methods 14, 959–962 (2017).

31. Kornberg, R. D. Chromatin structure: a repeating unit of histones and DNA. Science **184**,

868–871 (1974).

32. Caron, F., Jacq, C. & Rouvière-Yaniv, J. Characterization of a histone-like protein extracted

from yeast mitochondria. *Proceedings of the National Academy of Sciences* **76**, 4265–4269

(1979).

33. DeFrancesco, L. & Attardi, G. In situ photochemical crosslinking of HeLa cell mitochondrial

DNA by a psoralen derivative reveals a protected region near the origin of replication. Nucleic

Acids Research 9, 6017–6030 (1981).

34. Mei, S. et al. Cistrome Data Browser: a data portal for ChIP-Seq and chromatin accessibility

data in human and mouse. Nucleic Acids Research 45, D658–D662 (2017).

35. Zheng, R. et al. Cistrome Data Browser: expanded datasets and new tools for gene

regulatory analysis. Nucleic Acids Research **47**, D729–D735 (2018).

36. Kulakovskiy, I. V. et al. HOCOMOCO: towards a complete collection of transcription factor

binding models for human and mouse via large-scale ChIP-Seq analysis. Nucleic Acids Res 46,

gkx1106- (2017).

37. Buenrostro, J. D. et al. Integrated Single-Cell Analysis Maps the Continuous Regulatory

Landscape of Human Hematopoietic Differentiation. Cell **173**, 1535-1548.e16 (2018).

38. Rand, W. M. Objective Criteria for the Evaluation of Clustering Methods. *J Am Stat Assoc*

66, 846 (1971).

39. Li, B. et al. APEC: an accesson-based method for single-cell chromatin accessibility

analysis. Genome Biol 21, 116 (2020).

40. Stuart, T. et al. Comprehensive Integration of Single-Cell Data. Cell **177**, 1888-1902.e21

(2019).

41. Lun, A. T. L., McCarthy, D. J. & Marioni, J. C. A step-by-step workflow for low-level analysis

of single-cell RNA-seq data with Bioconductor. *F1000research* **5**, 2122 (2016).

42. Fang, R. et al. Comprehensive analysis of single cell ATAC-seq data with SnapATAC. Nat

Commun 12, 1337 (2021).

43. Zhao, Y., Ruan, S., Pandey, M. & Stormo, G. D. Improved Models for Transcription Factor

Binding Site Identification Using Nonindependent Interactions. Genetics **191**, 781–790 (2012).

44. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. Nature Methods

9, 357–359 (2012).

45. Hu, S. et al. ncHMR detector: a computational framework to systematically reveal non-

classical functions of histone modification regulators. *Genome biology* **21**, 48 (2020).

46. Zhang, Y. et al. Model-based analysis of ChIP-Seq (MACS). 9, R137-9 (2008).

47. Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif.

Bioinformatics **27**, 1017–1018 (2011).

48. Han, L. et al. Single cell transcriptomics identifies a signaling network coordinating

endoderm and mesoderm diversification during foregut organogenesis. Nat Commun 11, 4158

(2020).

49. Granja, J. M. et al. ArchR is a scalable software package for integrative single-cell chromatin

accessibility analysis. *Nat Genet* **53**, 403–411 (2021).

50. Otsu, N. A Threshold Selection Method from Gray-Level Histograms. *Ieee Transactions Syst*

Man Cybern **9**, 62–66 (1979).

 Figure legends

 Figure 1. SELMA framework for cleavage bias estimation.

 (**a**) Schematic of a naïve k-mer model for cleavage bias estimation.

 (**b**) Simplex encoding model. The coordinates of the 4 tetrahedral vertices of the cube encode

 the 4 nucleotides. Each dinucleotide is encoded as the outer product of the 2 mononucleotides.

 p represents the number of parameters in a k-mer simplex encoding model.

 (**c**) SELMA uses k-mer simplex encoding followed by a linear regression for k-mer cleavage

 bias estimation.

 (**d-k**) Comparison between the naïve k-mer model (**d,f,h,j**) and SELMA (**e,g,I,k**) on k-mer

 cleavage bias scores estimated from DNase-seq data (**d,e,h,i**) and ATAC-seq data (**f,g,j,k**).

 Each dot in a scatter plot represents an 8-mer, with its estimated bias score from different

 datasets represented in the x- and y-axes. Chromatin DNA (**d-g**) and naked DNA (**h-k**) from

 different human cell lines (**d-i**) or different species (**j,k**) were compared as labeled. R represents

 the Pearson correlation coefficient.

 (**l-o**) Intrinsic cleavage bias estimation accuracy measured by correlation between genome-wide

 observed (OBS) and bias-expected (EXP) cleavages with different k-mers. A higher Pearson

 correlation coefficient (R) indicates a better prediction of the observed cleavage profile using the

 estimated biases.

 Figure 2. SELMA consideration of dimeric Tn5 cleavages for ATAC-seq.

 (**a**) Schematic of SELMA consideration of dimeric Tn5 cleavages.

 (**b**,**c**) Cross-correlation between 5' cleavages and 3' cleavages across genome-wide accessible

 chromatin regions in the human GM12878 cell line from ATAC-seq (**b**) and DNase-seq (**c**)

 experiments. The x-axis represents the shift distance (in bp) between 5' and 3' cleavages.

 (**d-i**) Comparison of ATAC-seq intrinsic cleavage bias estimation accuracy measured by

Pearson correlation coefficients (R) between genome-wide observed (OBS) and bias-expected

(EXP) cleavages for human GM12878 (**d**) and K562 (**e**) cell lines, as well as several modified

Tn5-based assays (**f-i**). Different bars represent different estimation approaches: gray for

considering the 10-mer at the observed cut only (5' only); orange for SELMA; and blue for

several published approaches. Modified Tn5-based assays include (**f**) THS-seq with standard

Tn5; (**g**) THS-seq with mutated Tn5; (**h**) fast-ATAC-seq; and (**i**) omni-ATAC-seq.

Figure 3. SELMA bias estimation using mtDNA reads.

(**a**) Conventional approach to intrinsic cleavage bias estimation, in which a naked DNA sample

is required for bias estimation, and mtDNA reads from the chromatin DNA ATAC-seq

experiment are discarded.

(**b**) SELMA approach of intrinsic cleavage bias estimation using mtDNA reads. Chromatin reads

and mtDNA reads from the same ATAC-seq dataset were separated for Tn5 cleavage profiling

and bias estimation with SELMA, respectively.

(**c-d**) Scatter plots demonstrating the robustness of 10-mer bias scores estimated from mtDNA

(mtDNA reads from chromatin ATAC-seq data, x-axis) and genomic DNA (reads from naked

DNA ATAC-seq data, y-axis), comparing the naïve k-mer model (**c**) and SELMA (**d**). SELMA

shows a more robust estimation using mtDNA reads with higher correlation than the naïve k-

mer model.

(**e**) Comparison of the correlation coefficients between biases estimated from mtDNA reads and

from naked DNA with different mtDNA read counts.

Figure 4. TF binding inference on DNaseI footprint regions is affected by intrinsic cleavage

biases and improved by SELMA.

(**a,b**) Heatmaps of DNaseI cleavage patterns in the GM12878 cell line (**a**) and SELMA-

estimated bias scores (**b**) around ENCODE DNaseI consensus footprint regions. 5' and 3'

patterns were plotted separately. Footprint regions were consistently ranked by footprint length,

and signals for every 1000 regions with similar lengths were averaged as one row in the

heatmap.

(**c-h**) Aggregate plots of DNaseI cleavage patterns (**c,e,g**) and SELMA-estimated bias scores

(**d,f,h**) at all ENCODE DNaseI consensus footprint regions (**c,d**), footprint regions overlapping

with TF binding hotspots (**e,f**), and footprint regions overlapping with TF binding deserts (**g,h**).

DNaseI cleavages are from GM12878 for (**c** & **e**) and merged from GM12878, K562 and

ENCODE tissues (**g**). Dashed lines represent estimated bias spikes in the footprint regions.

(**i**) SELMA-estimated footprint bias scores (FBS) for different groups of footprint regions. P-

values were calculated by the Wilcoxon signed-rank test. The centerline, bounds of box, top line

and bottom line represent the median, 25^{th} to 75^{th} percentile range, 25^{th} percentile -

1.5*interquartile range (IQR) and 75^{th} percentile + 1.5*IQR, respectively.

(**j**) Performance rank scores for transcription factor binding prediction from DNaseI footprint

using various methods for the human GM12878 cell line. Boxplots with different colors represent

different approaches for prediction as indicated in the legends. Different sets of boxplots

represent footprint scores calculated by different published methods as labeled. P-values were

calculated by the Wilcoxon signed-rank test.

(**k**) Scatter plot showing the heterogeneous effect of intrinsic cleavage biases on different TF

motifs. Each data point represents motif sites for one TF in one cell type. Fractions of the TF-

bound motif sites overlapping with high-FBS footprint regions and with low-FBS footprint regions

are plotted on the x- and y-axes, respectively. More motifs located above the diagonal line

indicate that more TFs are more likely to bind at low-FBS sites than at high-FBS sites. The p-

value was calculated by t-test comparing the distribution of log likelihood ratios (y-axis/x-axis) to

the standard normal distribution.

Figure 5. Intrinsic cleavage biases affect scATAC-seq data analysis.

(**a,b**) UMAP visualization of single cells from scATAC-seq data in human hematopoietic cells.

Cells are colored by cell type (**a**) or by cell bias score (CBS) (**b**).

(**c**) CBS distributions of cells from different cell types and batches. Boxes are colored by cell

type using the same color palette as (**a**).

(**d-f**) Comparisons of cell clustering accuracy before and after considering the peak bias score

(PBS) on scATAC-seq data for human hematopoietic cells (**d**), human cell lines (**e**), and the

mouse primitive gut tube (**f**). K-means clustering was performed after PCA dimensionality

reduction. Blue, clustering using all ATAC-seq peaks; orange, using debiased peaks, i.e., 50%

to 99% of peaks with the lowest peak bias score (PBS). For each percentage from 50% to 99%

with a 1% increment, 100 randomly sampled subsets of peaks were used as the background for

determining the relative ranks of all peaks or debiased peaks. The relative ranks of the adjusted

Rand index (ARI), defined as the rank relative to the 100 randomly sampled subdatasets, for the

50 cases from 50% to 99%, were plotted.

(**g-j**) Comparisons of cell clustering accuracy before and after (orange) considering the peak

bias score (PBS) generated by existing methods on scATAC-seq data for human hematopoietic

cells. Each panel represents the result from a clustering approach: APEC (**g**), Seurat (**h**), scran

(**i**), and snapATAC (**j**). P-values were calculated by the Wilcoxon signed-rank test.

Figure 1

Figure 2

Figure 3

a Conventional approach

b SELMA approach

c naive K-mer

R = 0.586

Human naked DNA Bias score

Bias score GM12878 mtDNA

d SELMA

R = 0.924

Human naked DNA Bias score

Bias score GM12878 mtDNA

e

Bias score correlation mtDNA v.s. naked DNA

GM12878 mtDNA read count (x10^5)

SELMA

naive k-mer

Figure 4

Figure 5