Accurate estimation of intrinsic biases for improved analysis of bulk and single-cell

chromatin accessibility sequencing data using SELMA

Shengen Shawn Hu[1], Lin Liu[2], Qi Li[3], Wenjing Ma[1,4], Michael J. Guertin[5], Clifford A. Meyer[6], Ke

Deng[3], Tingting Zhang[7], Chongzhi Zang[1,8,9]*

[1]Center for Public Health Genomics, University of Virginia, Charlottesville, VA 22908, USA

[2]Institute of Natural Sciences, MOE-LSC, School of Mathematical Sciences, and SJTU-Yale

Joint Center for Biostatistics and Data Science, Shanghai Jiao Tong University, Shanghai

200240, China

[3]Center for Statistical Science, Tsinghua University, Beijing 100084, China

[4]Department of Computer Science, Emory University, Atlanta, GA 30322, USA

[5]Center for Cell Analysis and Modeling, Department of Genetics and Genome Sciences,

University of Connecticut, Farmington, CT 06030, USA

[6]Department of Data Science, Dana-Farber Cancer Institute, Boston, MA 02215, USA

[7]Department of Statistics, University of Pittsburgh, Pittsburgh, PA 15260, USA

[8]Department of Public Health Sciences, University of Virginia, Charlottesville, VA 22908, USA

[9]UVA Cancer Center, University of Virginia, Charlottesville, VA 22908, USA

*To whom correspondence should be addressed: zang@virginia.edu

Abstract

Genome-wide profiling of chromatin accessibility by DNase-seq or ATAC-seq has been widely

used to identify regulatory DNA elements and transcription factor binding sites. However,

enzymatic DNA cleavage exhibits intrinsic sequence biases that confound chromatin

accessibility profiling data analysis. Existing computational tools are limited in their ability to

account for such intrinsic biases and not designed for analyzing single-cell data. Here, we

present Simplex Encoded Linear Model for Accessible Chromatin (SELMA), a computational

method for systematic estimation of intrinsic cleavage biases from genomic chromatin

accessibility profiling data. We demonstrate that SELMA yields accurate and robust bias

estimation from both bulk and single-cell DNase-seq and ATAC-seq data. SELMA can utilize

internal mitochondrial DNA data to improve bias estimation. We show that transcription factor

binding inference from DNase footprints can be improved by incorporating estimated biases

using SELMA. Furthermore, we show strong effects of intrinsic biases in single-cell ATAC-seq

data, and develop the first single-cell ATAC-seq intrinsic bias correction model to improve cell

clustering. SELMA can enhance the performance of existing bioinformatics tools and improve

the analysis of both bulk and single-cell chromatin accessibility sequencing data.

Introduction

Cis-regulatory elements in the genome play a critical role in transcription regulation by interaction with protein molecules such as transcription factors (TFs). These DNA elements are usually unwrapped from packed nucleosomes and are accessible in the chromatin structure[1,2]. Genome-wide profiles of chromatin accessibility are a means to measure the global landscapes of active regulatory elements in different cell types. DNaseI hypersensitivity sequencing (DNase-seq) and the assay for transposase-accessible chromatin using sequencing (ATAC-seq) have become widely used for the genomic profiling of chromatin structure and accessibility[3,4]. Signal enrichments, or "peaks", from DNase-seq or ATAC-seq data are considered to represent accessible chromatin regions and can be used for inferring regulatory elements or TF binding sites. In addition, DNase-seq and ATAC-seq data also exhibit footprint patterns, which are fine structures in the accessible chromatin where high-affinity protein-DNA interactions protect the DNA from DNaseI or Tn5-transposase cleavages[4,5]. DNase/ATAC-seq footprint detection has been implicated as an effective approach for identifying accurate TF binding sites at base-pair resolution[6,7]. A few computational tools have been developed for detecting footprints from DNase-seq (RepFootprint[8], Wellington[9], PIQ[10]) or ATAC-seq data (HINT-ATAC[11], ToBIAS[12]). A recent study integrated 243 DNase-seq samples from different human cell and tissue types and reported approximately 4.5 million DNaseI consensus footprints associated with TF occupancy across the human genome as one of the largest maps of human regulatory DNA[7].

The premise of using DNase-seq and ATAC-seq data to profile chromatin accessibility is that enzymatic DNA cleavages represented by sequence reads reflect local chromatin openness only. However, it has been shown that both DNaseI and Tn5 transposase exhibit sequence selection biases in high-throughput sequencing data[13-16]. Such intrinsic enzymatic biases in

sequencing data can potentially confound observed cleavage patterns and footprint detection.

The characterization and correction of intrinsic cleavage biases are essential to DNase/ATAC-

seq data analysis. To characterize intrinsic cleavage biases, most studies use a k-mer model, in

which the k-mer DNA sequence centered at a cleavage site of DNaseI/Tn5 is used as the

signature of this cleavage[13,16,17]. The sequence bias can be estimated by counting the

occurrences of cleavages with each k-mer in one dataset relative to the genome-wide

occurrences of this k-mer. A naïve k-mer model assumes that k-mers are independent of each

other, resulting in an exponential increase in the degree of freedom when k increases.

Therefore, a naïve k-mer model becomes less feasible in practice with a large k, especially with

samples having insufficient sequencing depth. Although most studies use 6-mers with $4^6 = 4096$

parameters[8,10,13,16-18], it is unclear whether a different model with a larger k-mer size and more

feasible parameter estimation can achieve better performance.

Several studies have used various computational approaches for intrinsic cleavage bias

estimation[8,14,16,19] and footprint detection with bias correction[16,18-20] for bulk DNase/ATAC-seq

data. Recently, single-cell ATAC-seq (scATAC-seq) has enabled chromatin accessibility

profiling in thousands of individual cells at one time[21-24]. Due to the high sparsity of single-cell

data and because most chromatin accessibility regions contain only one read in one cell, any

potential bias can be substantial in scATAC-seq data compared to bulk data, creating additional

challenges in computational analysis. Incorporating intrinsic cleavage bias effects for improved

scATAC-seq analysis also requires comprehensive assessment and development of innovative

methods.

Here, we present Simplex Encoded Linear Model for Accessible Chromatin (SELMA), a

computational framework for the accurate estimation of intrinsic cleavage biases and improved

analysis of DNase/ATAC-seq data for both bulk and single-cell experiments. We demonstrate

that SELMA generates more accurate and robust bias estimation from bulk DNase/ATAC-seq

data than the naïve k-mer model and that SELMA can utilize mitochondrial DNA (mtDNA) for

bias estimation instead of requiring a separate naked DNA sample. We show an improved TF

occupancy inference on ENCODE consensus footprints by including SELMA-estimated biases

for each footprint. Finally, we show that SELMA-estimated biases can be incorporated with

existing scATAC-seq computational tools to generate more accurate cell clustering analysis.

Results

SELMA improves cleavage bias estimation in DNase-seq and ATAC-seq data

We developed SELMA for an accurate and robust estimation of intrinsic cleavage biases from

chromatin accessibility sequencing data. In SELMA, we start with a naïve k-mer model to

calculate the frequency of observed cleavage sites at each k-mer relative to the total

occurrences of this k-mer (Fig. 1a). We further encode each k-mer as a vector in the Hadamard

Matrix H_{16}, derived from a simplex encoding model, in which the k-mer sequences are encoded

as the vertices of a regular 0-centered simplex[25,26]. Intuitively, a k-mer can be represented as k

mononucleotides and $k - 1$ adjacent dinucleotides. Each mononucleotide is encoded as the 3-

dimensional vector of one of the four tetrahedral vertices of the cube of side 2 centered at the

origin. Each dinucleotide is encoded as the outer product of the 2 vectors representing the

associated nucleotides (Fig. 1b). Including an intercept, this k-mer simplex encoding model has

a total of $1 + 3 \times k + 9 \times (k - 1) = 12k - 8$ parameters, much fewer than the naïve k-mer model

(4^k). We use a linear model to fit these $12k - 8$ parameters from the naïve k-mer biases, and

use the fitted values as the SELMA-modeled cleavage biases (Fig. 1c).

As an intrinsic property of the enzyme (DNaseI/Tn5 transposase), the cleavage biases are

expected to be invariant across cell types and independent of chromatin states (Fig. S1a).

Comparing data from two cell types using different 8-mer models as an example, we found that

the biases estimated using SELMA have a higher correlation than those estimated using the

naïve k-mer model, for both DNase-seq (Fig. 1d,e) and ATAC-seq (Fig. 1f,g). Using sequence

reads from genomic regions with different chromatin accessibility levels, the naïve k-mer model

was not able to generate very consistent bias estimations (Fig. S1b,c), but the k-mer biases

estimated by SELMA using the same data were highly consistent (Fig. S1b-e). The sequence

preferences of DNaseI or Tn5 cleavage can be better reflected when the enzymes are applied

to deproteinized naked genomic DNA[16,17]. We found that the k-mer cleavage biases in naked

DNA DNase/ATAC-seq data estimated with the naïve k-mer model can still be improved by

SELMA, obtaining more consistent bias scores between different cell systems (Fig. 1h-k). These

data demonstrated that SELMA can improve the accuracy of estimating k-mer cleavage biases

in DNase-seq and ATAC-seq data.

With fewer parameters, SELMA enabled us to assess the effect of k-mer size on the

performance of bias estimation. Using a "bias expected cleavage" approach[8,13,16,19], we

compared the bias estimation performances measured as the correlation coefficient between

the genome-wide observed cleavages and bias expected cleavages estimated using SELMA

with different k. A higher correlation coefficient indicates a more accurate bias estimation to

calculate the expected cleavages. For both DNase-seq and ATAC-seq data from two different

cell lines, we found that 10-mer outperforms any other k-mer (Fig. 1l-o). We then applied this

analysis to more DNase-seq and ATAC-seq data from a variety of human tissues generated by

ENCODE and found that 10-mer always exhibited the best performance (Fig. S2). The above

empirical evidence suggested that 10-mer is the optimal choice for intrinsic cleavage bias

estimation for both DNaseI (DNase-seq) and Tn5 (ATAC-seq) cleavages.

SELMA improves ATAC-seq bias estimation by considering dimeric Tn5 cleavages

Many studies processed DNase-seq and ATAC-seq data in a similar way, treating individual

DNA cleavage sites directly as indications of accessible chromatin[8,14]. However, the

mechanisms of enzymatic DNA cleavage are different between DNaseI and Tn5 transposases.

Unlike DNaseI, the Tn5 transposase binds DNA as a dimer and inserts adaptors on the two

strands separated by 9bp[4,27] (Fig. 2a). As a result, the presence of each observed Tn5 cleavage

in ATAC-seq data should be the consequence of two insertion events induced by the same Tn5

dimer, and the bias estimation of a Tn5 cleavage site should consider both the observed

cleavage and the coupling cleavage 9bp downstream on the reverse strand. Therefore, SELMA

estimated the bias on an ATAC-seq cleavage site as the geometric mean of the bias scores of

the 10-mers at the 5' observed cleavage and at the 3' cleavage 9bp downstream on the reverse

strand (Fig. 2a). To show the dimeric Tn5 cleavage effect, we calculated the cross-correlation

between the genome-wide profiles of the plus strand cleavages and the minus strand

cleavages. As expected, we observed a peak at 9bp of the cross-correlation curve specifically in

the ATAC-seq data but not the DNase-seq data (Fig. 2b,c). Similarly, we observed perfectly

matching aggregate cleavage patterns on TF motif consensus sites between plus strand and

minus strand cleavages shifted by 9bp (Fig. S3a,b).

Using the similar "observed-expected correlation" approach described above, we found that

SELMA considering dimeric cleavages outperformed models considering only 5' cleavage in

generating a more accurate bias expected cleavage pattern for ATAC-seq data (Fig. 2d,e). We

confirmed that this observation was specific to ATAC-seq, as similar analyses for DNase-seq

from the same cell lines did not show a similar level of improvement (Fig. S3c,d). We also

compared SELMA with several existing bias estimation approaches[18] and found that SELMA's

performance was the best for ATAC-seq data from several cell lines as well as different human

tissue types from ENCODE (Fig. 2d,e, Fig. S3e-j). Several new Tn5-based techniques have

recently been developed for improved chromatin accessibility profiling, including THS-seq[28],

Fast-ATAC-seq[29], and Omni-ATAC-seq[30]. Data from these technologies showed intrinsic

cleavage biases similar to those of conventional ATAC-seq (Fig. S3k-n). We found that SELMA

also outperformed other approaches in bias estimation (Fig. 2f-i). Collectively, these data

suggested that SELMA can most accurately estimate intrinsic cleavage biases in data from

ATAC-seq and other Tn5-based techniques.

SELMA enables accurate bias estimation by utilizing mitochondrial DNA (mtDNA) reads

Accurate estimation of enzymatic cleavage biases independent of chromatin usually requires a

control sample of deproteinized naked DNA digested by the same enzyme. Biases estimated

from the naked DNA control data can be used to correct the chromatin accessibility profiling

data[13,14,16-18] (Fig. 3a). In conventional DNase/ATAC-seq data analyses, sequence reads

mapped to mitochondrial DNA (mtDNA) are usually discarded[4]. Lacking histones responsible for

the chromatin structure, mtDNA is similar to deproteinized naked DNA[31-33]. Therefore, we

sought to use the mtDNA reads from DNase/ATAC-seq data for cleavage bias estimation as an

alternative to using a naked DNA control sample (Fig. 3b). Likely because of relatively low read

counts and lack of sequence diversity (e.g., human mtDNA is only < 20kb long), the naïve k-mer

model exhibited poor performance on bias estimation from mtDNA reads, using bias scores

estimated from naked DNA as a reference (Fig. 3c). In contrast, SELMA generated a more

accurate bias estimation from the same mtDNA reads, which was highly consistent with the bias

scores estimated from the naked DNA data (Fig. 3d), demonstrating the power of SELMA to use

less input to make accurate bias estimations. We applied this approach to a series of ATAC-seq

and DNase-seq datasets for different human tissues from ENCODE and found that SELMA was

better than the naïve k-mer model in yielding a consistently higher correlation coefficient (>0.9)

between mtDNA-estimated bias and naked DNA-estimated bias for every ATAC-seq and

DNase-seq sample tested (Fig. S4a,b). Many optimized ATAC-seq protocols aimed to reduce

mtDNA reads to increase the fraction of chromatin DNA reads for chromatin accessibility signal

yield[29,30]. We sampled down mtDNA reads to test the performance of SELMA in making robust

bias estimations and found that SELMA could accurately estimate the bias with as few as

50,000 mtDNA reads (Fig. 3e, Fig. S4c), which is approximately 0.2% of the sequencing depth

of a regular ATAC-seq sample and lower than the fraction of mtDNA reads in any existing

ATAC-seq experiment[29,30]. These data demonstrated that by applying SELMA, mtDNA reads

can be utilized to substitute naked DNA control samples for accurate bias estimation.

Considering SELMA-estimated bias improves TF binding inference on ENCODE DNaseI

footprint regions

With an accurate bias estimation model developed, we next sought to examine the potential

effect of intrinsic biases on TF binding footprints in chromatin accessibility profiling data.

Focusing on the 4,465,728 human DNaseI consensus footprints across the human genome, we

plotted the DNaseI cleavages from different human cell lines and observed similar classic

DNaseI cleavage protection ("footprint") patterns across these footprint regions (Fig. 4a, Fig.

S5a). Interestingly, we also observed a recurrent pattern of the SELMA-estimated DNaseI

cleavage bias that is well aligned with the DNaseI cleavage pattern across these footprint

regions (Fig. 4b, Fig. S5b). In the aggregate view of footprint regions of different lengths, the

DNaseI cleavage signals exhibited a clear "cliff-bound valley"-shaped footprint pattern (Fig.

S5c). The DNaseI cleavage bias scores exhibited a pattern of two spikes located inside the

footprint coordinates (Fig. S5d). After normalizing various footprint lengths, we plotted the

aggregate DNaseI cleavage patterns across all consensus footprint regions and found that the

overall "footprint pattern" clearly included the double spike pattern of cleavage biases (Fig. 4c),

which aligned well with the two spikes in the aggregate bias score pattern (Fig. 4d), indicating a

possible contribution of intrinsic biases to the DNaseI footprinting.

To assess the interference of intrinsic biases with DNaseI cleavage patterns at TF binding

footprints, we aligned the 4.5 million consensus footprints with more than 10,000 human TF

ChIP-seq datasets from the Cistrome Data Browser database[34,35] and collected two sets of

footprint regions: "TF binding hotspots", the footprint regions overlapping with TF binding sites

detected from more than 3,000 ChIP-seq datasets, and "TF binding deserts", the footprint

regions that do not overlap with any TF binding sites from any ChIP-seq dataset or any human

TF motif sequence from the HOCOMOCO database[36] (Fig. S5e). We compared the aggregate

DNaseI cleavage patterns and the bias score patterns in these two sets of consensus footprint

regions and found that the DNaseI cleavage pattern in TF binding hotspot regions was

dominated by the classic DNaseI footprint pattern with little interference from the bias (Fig. 4e,f),

while in the TF binding desert regions, the entire cleavage "footprint" pattern was apparently

determined by the two spikes from the intrinsic bias (Fig. 4g,h). These results suggested that

although the overall DNaseI cleavage pattern is indicative of TF binding, the intrinsic cleavage

bias may interfere with the real footprint pattern, and the effect on those footprint regions with

rare TF binding events can be drastic. These observations were reproducible in DNase-seq

data from different cell and tissue types (Fig. S6a-b).

To quantify the level of intrinsic cleavage biases in a consensus footprint region, we defined a

footprint bias score (FBS), as the relative SELMA-estimated bias score at the two spikes

compared to the average SELMA bias score across the rest of the region, for each footprint.

Consistent with what we observed in the aggregate view, the FBSs for TF binding deserts were

significantly higher while the FBSs for TF binding hotspots were significantly lower than the

background of all consensus footprints (Fig. 4i, $p < 10^{-5}$, by Wilcoxon test), indicating that FBS

might be an informative feature of the consensus footprint regions and might help separate true

TF binding footprints from false-positive DNaseI footprint patterns. Next, we used a model-

based approach to assess the potential power of SELMA-derived FBS in boosting the

performance of TF binding inference from DNase-seq signal patterns on consensus footprint

regions containing the TF motif sequence. For every TF with a known motif in a cell type with

both ChIP-seq and DNase-seq data available, we employed a multinomial logistic regression

model using different sets of features, each of which may include DNase-seq read count,

DNaseI footprint score produced from an existing method, and an optional FBS, to infer the TF

binding occupancy (determined by peak occurrence in ChIP-seq data) in the motif-containing

footprint regions. For each footprint method, we compared the TF binding inference

performance of the original method (read count + footprint score as features), the original

method plus a randomized naïve k-mer bias score feature as a control, and the original method

plus the SELMA bias score (FBS), using an integrated rank score strategy. We included our

previous footprint method[13] and several representative methods that outperform others,

Wellington[9] and HINT[11,19] (with and without bias correction mode) in this comparison. We

surveyed all human cell types that have both DNase-seq and more than 20 TF ChIP-seq

samples available from ENCODE, including in 7 cell lines and a total of 375 ChIP-seq samples

for 156 different TFs (Table S1, Fig. S7a). For each TF ChIP-seq sample in each cell line, we

calculated differential rank score of the inference performance by adding a bias score feature.

As a result, adding a random k-mer bias score did not change the inference performance. In

contrast, the models with SELMA FBS added can significantly increase the rank scores for the

majority of ChIP-seq samples, regardless of which method was used originally to calculate the

footprint score (277-291, or 74%-78%, varying across different footprint methods, Fig. 4j, Fig.

S7, Table S2). For example, using our previous footprint method[13], 277 ChIP-seq samples

(74%) received a higher inference rank score when considering SELMA FBS, covering 117

(75%) of the 156 TFs (Table S2). Among these, Zinc finger family TFs including CTCF and

REST showed the highest improvement after considering footprint bias (Fig. S8, Table S2),

consistent with previous studies about the positive correlation between footprint strength and

residence time of the TF on DNA[13,16]. Meanwhile, the SOX family (e.g., SOX5) and HLH family

(e.g., MYC) TFs rarely showed improved inference performance after considering footprint bias

(Fig. S8, Table S2). These TFs were those having short residence time on DNA and weak

footprints[15], which were expected not to be affected by footprint biases. These results

demonstrated that considering the intrinsic cleavage bias can improve the performance of

existing footprint computational methods for the binding inference for most TFs with footprints.

To assess whether the intrinsic bias has different levels of interference with the footprint regions

for different TFs, we selected two subsets of sequence motif-containing footprint regions for

each TF as the top 10% with the highest FBS or the bottom 10% with the lowest FBS, and

compared the frequencies of observing actual TF binding (overlapping with a ChIP-seq peak) in

these subsets of footprints. We found that different TFs had various preferences for binding at

low-FBS footprints or high-FBS footprints. Among the 156 TFs included, a significantly larger

amount of TFs can be improved with SELMA FBS than those that cannot be improved by

SELMA (Fig. 4k). These results suggested that intrinsic cleavage biases might affect different

TFs at various levels in divergent directions, and considering intrinsic bias should improve the

footprint-based TF binding inference for most TFs.

SELMA improves the accuracy of cell clustering from single-cell ATAC-seq data

Single-cell ATAC-seq (scATAC-seq) technologies enable the detection of accessible chromatin

regions at single-cell resolution in thousands of cells at a time[21-23]. Due to the scarcity of

cleavage events in an individual cell, most chromatin accessibility regions in a single cell have

only one aligned fragment, making the potential influence of intrinsic cleavage biases more

substantial in scATAC-seq data than in bulk ATAC-seq data. We collected scATAC-seq

datasets generated using different platforms for different biological samples, human

hematopoietic cells[37], human cell lines[21], and mouse primitive gut tube[38], and found that the

scATAC-seq data contained similar intrinsic cleavage biases to bulk data with highly correlated

bias scores estimated by SELMA (Fig. S9a-c). We estimated the average cleavage bias for

each individual cell (cell bias score, CBS) and found that the distribution of CBS was different

across cell cluster patterns, batches, and annotated cell types (Fig. 5a-i). We further surveyed

datasets from the 10x Single Cell Multiome platform for different biological systems including

mouse embryonic brain, human peripheral blood mononuclear cells (PBMC), and human lymph

node, each of which has scATAC-seq and scRNA-seq performed in the same cell and we used

the scRNA-seq derived cell clusters as the "pseudo" ground truth to label the cells. As a result,

we still found similar intrinsic cleavage biases in the scATAC-seq part of the data (Sig. S9d-f)

and CBS affecting different cell clusters (Fig. 5j-r). These observations indicated that, regardless

of experimental platforms and biological systems, the Tn5 intrinsic cleavage biases can

contribute to cellular heterogeneity observed from scATAC-seq data and can affect cell

clustering analysis.

We next assessed whether considering intrinsic cleavage biases can improve cell clustering

based on scATAC-seq data. We used the actual cell-type labels as the clustering ground truth

for the human hematopoietic cell sample and the mixed human cell line sample, and used

scRNA-seq data-projected cell labels as a "pseudo" ground truth for the mouse primitive gut

tube sample. We used the adjusted Rand index (ARI)[39] to quantify the accuracy of an

unsupervised clustering result. As scATAC-seq-based cell clustering can be performed on all

chromatin accessibility regions (ATAC-seq peaks), we sought to address whether removing

peaks with high intrinsic biases can increase the clustering accuracy. We first tested a simple

clustering approach that involved principal component analysis (PCA) dimensionality reduction

followed by K-means clustering. For all 6 scATAC-seq datasets, cell clustering after removing

1%-50% of peaks with the highest PBS largely increased ARI from using all peaks (Fig. S10).

This improvement in clustering accuracy by removing high-bias peaks is significant compared to

the control of randomly removing the same number of peaks (Fig. 6a-f). These data suggested

that scATAC-seq cell clustering could be improved by excluding high-bias peaks that

confounded the analysis.

To correct the intrinsic cleavage bias effect in scATAC-seq data in a systematic manner, we

developed a general model to weight peaks by the percentile of SELMA PBS (Fig. 6g). The

weight function was determined empirically using a Beta distribution based on the relative

contribution of each PBS percentile of peaks to the true cell type classification (Fig. S11,

Methods). We applied this weight function to adjust the peak-by-cell read count matrix for

intrinsic bias correction and used the bias-corrected data matrix for cell clustering analysis. To

evaluate the performance, we tested several commonly used single-cell data analysis tools,

including APEC[40], Seurat[41], scran[42], and snapATAC[43], in addition to K-means, for scATAC-seq

cell clustering, and compared the cell clustering accuracy between using the uncorrected raw

data and using the bias-corrected data. Although different tools showed various performances,

across the 6 biological samples we tested, SELMA-corrected data yielded a higher ARI than

uncorrected data for most cases, and always reached the highest ARI for each sample (Fig. 6h-

m, Table. S4). These results demonstrated that SELMA can reduce the effect of intrinsic

cleavage biases in scATAC-seq data and can be applied to existing single cell data analytical

tools to improve the accuracy of cell clustering analysis.

Discussion

The existence of enzymatic cleavage biases in DNase-seq and ATAC-seq experiments has

been widely acknowledged in the field, but to what extent such intrinsic biases affect data

analysis, especially on the single-cell level, has not been systematically assessed. SELMA

provides a quantitative approach for the accurate and robust estimation of intrinsic cleavage

biases in both bulk and single-cell chromatin accessibility sequencing data and requires fewer

sequence reads than the naïve k-mer model. Taking Tn5 dimerization into consideration,

SELMA yields more accurate bias estimation specifically for ATAC-seq data by including k-mer

sequences at two Tn5 cleavages 9bp apart. SELMA can improve functional analysis and

interpretation of chromatin accessibility profiles. On the bulk level, we showed that considering

SELMA-estimated biases can improve TF binding inference from ENCODE DNaseI consensus

footprints for most TFs, with better performances compared to existing tools, including some

that already considered "biases". On single-cell level, we showed widespread existence of

intrinsic cleavage biases in single-cell ATAC-seq data, and demonstrated that SELMA single-

cell bias reduction model can enhance the performance of existing tools and can increase the

accuracy of cell type clustering. Therefore, SELMA can help researchers obtain more biological

insights from chromatin accessibility data.

SELMA is built on top of the widely used k-mer model by combining simplex encoding and a

linear model. Simplex encoding has the unique ability to capture the pairwise symmetry and

orthogonality between mononucleotides and interactions within each dinucleotide. It significantly

reduces the degrees of freedom without losing any variance information compared to the naïve

k-mer model. These properties enable SELMA to make robust estimations from fewer sequence

reads or smaller datasets than are required by other approaches, hence enabling *de novo* bias

estimation from a smaller DNA molecule, such as mtDNA in a DNase/ATAC-seq sample,

without having to generate a separate genomic naked DNA sample. However, SELMA still relies

on sufficient read counts for each k-mer for an accurate estimation. Although the performance of

SELMA may increase with k, this effect is not unlimited. The performance using 12-mer is not as

good as using 10-mer possibly because there are not enough reads in the dataset for many 12-

mers. Nevertheless, SELMA works well for most existing DNase/ATAC-seq datasets tested in

our study. In addition, the feasibility and superior performance of simplex encoding have also

been demonstrated for TF motif characterization[44]. Following similar encoding strategies,

SELMA can potentially be applied to any k-mer-based model or to any high-throughput

sequencing data for robust sequence bias estimation and pattern recognition.

The k-mer biases estimated by SELMA are consistent across species and cell types, reflecting

the assumption that the cleavage biases are intrinsic features of the enzymatic assays.

Therefore, one can directly use the SELMA-estimated k-mer biases from DNaseI or Tn5-

digested naked DNA data as universal intrinsic biases for DNase-seq and ATAC-seq,

respectively, and incorporate these precalculated bias scores into the data analysis. However,

this bias dataset is not guaranteed to remain accurate in many species that have not been

profiled, as there might be unknown biases that have not been characterized. Although we are

confident that the SELMA-estimated results should still be largely valid, one can always use the

SELMA package for *de novo* estimation of cleavage biases from one's own datasets.

When applying SELMA-estimated FBS to correct biases in DNase footprints, our data were

limited to ENCODE DNaseI consensus footprints as a proof-of-principle study. While this

ENCODE dataset is thus far the largest DNase footprint repertoire, users might be interested in

de novo detection of footprints from their customized DNase/ATAC-seq data. As many

bioinformatics tools are already available for such tasks using various computational models[20],

SELMA or SELMA-generated bias information can be incorporated into each of those models

for improved performance for more accurate regulatory DNA identification from footprints. On

single-cell ATAC-seq analysis, while we show that SELMA single-cell bias correction model can

achieve more accurate cell clustering using a few existing tools, performance still varies across

these tools due to the different statistical and computational models used in these methods.

One potential issue that limits a larger-scale benchmarking is the lack of ground truth for most

existing scATAC-seq data. Except for the human hematopoietic cell sample and the mixed

human cell line sample that have the known cell type labels as ground truth, we had to use

transcriptome profiling scRNA-seq data as a "pseudo" ground truth or solver standard to

determine the cell type identities, which implies the assumption that chromatin accessibility

profiles should characterize the same cell identity as transcriptomic profiles. Nevertheless, the

overall increase on the cell clustering consistency for different biological systems tested

indicates that SELMA bias correction reduces confounding noises in the data while biologically

meaningful variances are retained[45]. In summary, SELMA is a universal and systematic bias

reduction model and can be used to enhance the performance of existing methods and to

improve single-cell chromatin accessibility profiling analysis.

Methods

High-throughput sequencing data collection and processing

Data collection. Publicly available ATAC-seq, single-cell ATAC-seq, DNase-esq and ChIP-seq

data used in this study were collected from Gene Expression Omnibus (GEO) and the ENCODE

consortium. The metadata and accession numbers are listed in Table S1.

Processing of bulk ATAC-seq and DNase-seq data. Raw sequencing reads were aligned to

the GRCh38 (hg38) reference genome with bowtie2 (v2.2.9) (-X 2000 for paired end data)[46].

Low-quality reads (MAPQ < 30) were discarded. For paired-end sequencing data, reads with

two ends aligned to different chromosomes (chimeric reads) were also discarded. For paired-

end data, reads with identical 5' end positions for both ends were regarded as redundant reads

and discarded. The nonredundant reads were separated into chromosomal DNA (chromatin

reads) and mitochondrial DNA (mtDNA reads) based on their genomic location. Peak detection

was performed on the nonredundant chromatin reads using MACS2[47] (v2.1.2) (-q 0.01, --extsize

50) and +/- 200bp centered on the peak summits was collected as the genome-wide chromatin

accessible regions. The accessible regions in each dataset were separated into 5 groups from

highest accessibility to lowest accessibility based on the read count on each peak (for Fig. S1).

The 5' end nucleotides of each read were piled up to generate the genome-wide observed

cleavage profile.

Processing of single-cell ATAC-seq data. For the human hematopoietic cell sample, raw

sequencing reads were aligned to the GRCh38 (hg38) reference genome with bowtie2 (-X

2000). Low quality reads (MAPQ < 30), chimeric reads and duplicate reads for each individual

cell were discarded. For the mouse gut tube sample, scATAC-seq data from the 10x Genomics

platform were preprocessed with Cell Ranger ATAC with the default parameters to generate

fragments for each individual cell. For the 10x Single Cell Multiome datasets, the processed

fragment files were downloaded from the 10x genomics website

(https://www.10xgenomics.com/resources/datasets/fresh-embryonic-e-18-mouse-brain-5-k-1-

standard-2-0-0, https://www.10xgenomics.com/resources/datasets/pbmc-from-a-healthy-donor-

no-cell-sorting-10-k-1-standard-2-0-0, https://www.10xgenomics.com/resources/datasets/fresh-

frozen-lymph-node-with-b-cell-lymphoma-14-k-sorted-nuclei-1-standard-2-0-0). The fragment

files were then used as input for the subsequent analysis. Because the Cell Ranger ATAC/ARC

pipeline shifted from the Tn5 cleavage sites to +4/-5 bp in generating the fragment file, the

coordinates were shifted back to represent the actual cleavage loci. For scATAC-seq datasets,

cells with more than 10,000 reads were retained for analysis. For 10x Single Cell Multiome

datasets, cells pre-selected by Cell Ranger ARC and with more than 10,000 reads in both

scRNA-seq and scATAC-seq parts were retained for analysis.

Processing of ChIP-seq data. ChIP-seq data were collected and processed as described in a

previous study[48]. In brief, based on the quality measurements from the Cistrome data browser

(CistromeDB)[34,35], we included only the ChIP-seq datasets that pass at least four out of the first

five QC measures (i.e., sequence quality, mapping quality, library complexity, ChIP enrichment,

and signal-to-noise ratio). We also excluded the ChIP-seq datasets with fewer than 1000 peaks

identified by MACS2[27] with an FDR cutoff of 0.01. For TFs with multiple datasets available in the

same cell type, we kept only one dataset with the best quality based on passing most of the 6

CistromeDB QC measures, including sequence quality, mapping quality, library complexity,

ChIP enrichment, signal-to-noise ratio, and regulatory region enrichment. The ChIP-seq peak

files (detected by MACS2) for the selected samples were downloaded from CistromeDB for the

following analysis.

TF motif analysis. The motifs of human TFs were collected from the HOCOMOCO database[36]

(v11), and the genome-wide motif sites of TFs were detected by FIMO (v4.12.0) in the MEME

package[49]. Motif sites located outside of the genome-wide 36bp unique mappable regions were

excluded from the analysis. In total, 25,027,116 motif sites for 66 TF motifs from the human

genome were included in the analysis.

SELMA model

Naïve k-mer bias model for intrinsic cleavage bias estimation. The naïve k-mer biases were

calculated as described in a previous study[13]. In brief, a naïve k-mer bias was estimated as the

observed frequency of the cleavages with the k-mer relative to the frequency of that k-mer in the

background. For each mapped sequence read in a DNase-seq or ATAC-seq dataset, the

enzymatic cleavage site was between genomic positions i and i-1 for the plus (+) strand reads

and between i and i+1 for the minus (−) strand reads, where i represents the genomic position

of the 5' nucleotide of the reads. The associated k-mer sequence was thus assigned as

$\left[i - \frac{k}{2}, i + \frac{k}{2} - 1\right]$ for the plus strand reads and $\left[i - \frac{k}{2} + 1, i + \frac{k}{2}\right]$ for the minus strand reads. The

naïve k-mer bias score for k-mer j is defined as the number of all observed cleavages with that

k-mer divided by the occurrences of that k-mer in the background:

484
$$S_j = \frac{N_j}{M_j}$$

where N_j is the count of cleavages with k-mer j, and M_j is the total count of occurrences of k-

mer j in the background (Fig. 1a). In the case of chromatin DNA, this background included 400

bp from each chromatin accessible region centered at the peak summit detected by MACS2.

The background in the naked DNA samples included genome-wide 36bp unique mappable

regions. The background in mtDNA included the mitochondrial DNA sequence. The median was

further subtracted from the bias scores for visualization in the scatter plots (e.g., Fig. 1d-k).

In the naïve k-mer model, the bias score for each k-mer is independent and empirically

determined from the data. The number of independent variables in the model is the total number

of k-mers, i.e., 4^k.

SELMA model for intrinsic cleavage bias estimation. In SELMA, a simplex encoding model

followed by a linear model was employed on top of the naïve k-mer model to better estimate the

intrinsic cleavage biases for each k-mer. To efficiently encode a k-mer sequence considering

their intrinsic similarities, a simplex encoding model was adopted from previous studies[25,26], in

which the DNA nucleotides were encoded as vectors representing the four tetrahedral vertices

of a regular 0-centered simplex (Fig. 1b).

502
$$\begin{aligned} A &= [\quad 1 \quad -1 \quad -1 \quad] \\ C &= [\quad -1 \quad 1 \quad -1 \quad] \\ G &= [\quad -1 \quad -1 \quad 1 \quad] \\ T &= [\quad 1 \quad 1 \quad 1 \quad] \end{aligned}$$

504   In the simplex encoding, the vectors representing the four nucleotides are of equal length,

505   mutually orthogonal, and equidistant from each other. To account for interactions between

506   adjacent nucleotides, a dinucleotide was additionally encoded as the outer product of the two

507   vectors associated with the two nucleotides:

$$
\begin{aligned}
AA &= [\ \ \ 1\ \ -1\ \ -1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ ] \\
AC &= [\ -1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ ] \\
AG &= [\ -1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ ] \\
AT &= [\ \ \ 1\ \ -1\ \ -1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ ] \\
CA &= [\ -1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ -1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ ] \\
CC &= [\ \ \ 1\ \ -1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ ] \\
CG &= [\ \ \ 1\ \ \ \ 1\ \ -1\ \ -1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ ] \\
CT &= [\ -1\ \ -1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ -1\ \ -1\ \ ] \\
GA &= [\ -1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ -1\ \ ] \\
GC &= [\ \ \ 1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ -1\ \ ] \\
GG &= [\ \ \ 1\ \ \ \ 1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ -1\ \ -1\ \ -1\ \ \ \ 1\ \ ] \\
GT &= [\ -1\ \ -1\ \ -1\ \ -1\ \ -1\ \ -1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ ] \\
TA &= [\ \ \ 1\ \ -1\ \ -1\ \ \ \ 1\ \ -1\ \ -1\ \ \ \ 1\ \ -1\ \ -1\ \ ] \\
TC &= [\ -1\ \ \ \ 1\ \ -1\ \ -1\ \ \ \ 1\ \ -1\ \ -1\ \ \ \ 1\ \ -1\ \ ] \\
TG &= [\ -1\ \ -1\ \ \ \ 1\ \ -1\ \ -1\ \ \ \ 1\ \ -1\ \ -1\ \ \ \ 1\ \ ] \\
TT &= [\ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ \ \ 1\ \ ]
\end{aligned}
$$

One can show that these vectors for dinucleotide interactions are also of equal length, mutually

orthogonal, and equidistant from each other. In fact, in the simplex encoding, mononucleotides

and dinucleotides were encoded as rows in a Hadamard matrix of order 4 and a Hadamard

matrix of order 16, respectively.

Considering both mononucleotides and dinucleotide interactions, a k-mer can then be encoded

as k mononucleotides and $k - 1$ dinucleotides, plus an intercept term. Therefore, the

dimensionality of a k-mer simplex encoding is

518
$$p(k) = 1 + 3k + 9(k - 1) = 12k - 8$$

519   In SELMA, the intrinsic k-mer biases were then estimated in a linear model with these $12k - 8$

520   parameters using the observed naïve k-mer bias scores (Fig. 1c). In detail, we had the formula:

521
$$y \sim \sum_{i=1}^{12k-8} a_i x_i \tag{1}$$

where each observation y is the naïve k-mer bias score; $x_i \in \{1, -1\}$, is the independent

variable in the simplex encoding vectors; and a_i is the parameter to be estimated. After linear

regression, the model-fitted value $\hat{y}$ was defined as the SELMA bias score for each k-mer.

Genome-wide cross-correlation analysis. For the analysis presented in Fig. 2b,c, reads from

plus (+) and minus (-) strands on chromatin accessible regions (peaks) were collected

separately to generate plus strand observed cleavage profile and minus strand observed

cleavage profile, respectively. The Pearson correlation coefficient between the plus strand

signal and the k-bp shifted minus strand signal is:

531
$$\rho_k = \frac{\sum_i (P_i - \bar{P})(M_{ik} - \overline{M_k})}{\sqrt{\sum_i (P_i - \bar{P})^2 \sum_i (M_{ik} - \overline{M_k})^2}} \tag{2}$$

where P_i is the log-scaled plus strand cleavage count at genomic position *i* (log2(cleavage+1)),

M_{ik} is the log-scaled minus strand cleavage count at genomic position *i* with a k-bp shift, $\underline{P}$ is

the mean of all the P_i, and $\overline{M_k}$ is the mean of M_{ik} for all *i*. All genomic positions on the genome-

wide DNase/ATAC-seq peaks were included in the analysis. *k* is chosen from 1 to 20 (x-axis in

Fig. 2b,c).

Comparison of different bias estimation methods. We use a_i^+ and a_i^- to denote the "5' only"

intrinsic sequence bias score at genomic position *i* on the plus strand and minus strand,

respectively. We use a'^+_i and a'^-_i to denote the bias score from other bias estimation methods in

this section. Different bias estimation methods used in Fig. 2 and its associated sections are

listed below:

●   For the "5' only" method, a_i^+ was calculated based on the k-mer ratio associated with the

544   k-mer spanning positions $[i - k/2,\ i + k/2 - 1]$ on the plus strand, and a_i^- was

545   calculated in a similar way based on the nucleotides spanning genomic positions $[i -$

546           $k/2 + 1,\ i + k/2]$ on the minus strand (reverse complement of the DNA sequence on the

547   corresponding plus strand). This method was applied to both DNase-seq and ATAC-seq,

548   while the other methods were applied only to ATAC-seq as they were specifically

549   designed for ATAC-seq.

550   ●   For SELMA (Fig. 2a), the bias score was calculated as the geometric mean of the "5'

551   only" bias score at the given position and the "5' only" bias score at 9bp downstream of

552   the other strand, i.e., $a'^+_i = \sqrt{a_i^+ \times a_{i+9}^-}$, and $a'^-_i = \sqrt{a_i^- \times a_{i-9}^+}$

553   ●   For the model in Martins et al[14], the bias score at genomic position i was calculated in a

554   similar way to the "5' only" method but using a gapped 11-mer model. The model could

555   be represented as $XXXXXXNNNXNXCXXNNXNNNNXNX$, in which position i was

556   represented by C; positions that were ignored were represented by X and informative

557   positions were represented by N.

558   ●   For the model in Baek et al[18], the bias score at genomic position i was calculated in a

559   similar way to the "5' only" method but the cleavages were shifted +4/-5 bp for +/- strand

560   cleavages. In practice, following the description in the "bagfoot" package, the bias score

561   at position i was calculated as $a'^+_i = a_{i+5}^+$, and $a'^-_i = a_{i-5}^+$.

562   ●   For the model in Calviello et al[8], the bias score at genomic position i was calculated in a

563   similar way to the "5' only" method but using the 6-mer bias table provided in the study.

Calculation of observed and bias-expected cleavage. The 1bp at 5' end positions for DNase-

seq or ATAC-seq reads were piled up to generate the genome-wide observed cleavage profile.

To generate the bias-expected cleavage profile, for a 50-bp window centered on nucleotide i,

we placed the same number of observed cleavages following the sequence bias contribution in

this window. Let $\widehat{N_i^s}$ represent the bias expected cleavage at position i on strand $s \in \{+,-\}$, N_i^s

represent the observed cleavage at position i on strand s, and a_i^s denote the intrinsic cleavage

bias (estimated with any of the above methods) at position i on strand s. Then we have

$$\widehat{N_i^s} = N_i^s y_i^s \tag{3}$$

$$y_i^s = \frac{2^{a_i^s}}{\sum_{j=i-25}^{i+24} 2^{a_j^s}} \tag{4}$$

We used the Pearson correlation coefficient to compare the observed cleavage profile and the

bias-predicted cleavage profile (Fig. 2). We only considered positions within the accessible

regions to ensure that all positions had sufficient reads in the 50-bp window for accurate

estimation.

DNaseI footprint analysis

Data processing. The genome-wide DNaseI consensus footprint regions were downloaded

from Reference[7]

(https://resources.altius.org/~jvierstra/projects/footprinting.2020/consensus.index/). The

observed DNaseI cleavage profile from a DNase-seq dataset and DNaseI SELMA bias scores

across +/- 50bp centered on the footprint region were plotted as heatmaps (Fig. 4a,b). The

footprint regions were ordered by the footprint lengths, and each 1000 footprint regions with

similar lengths were compressed as one row in the heatmap for better visualization. The +/-

strand signals were plotted separately. We aligned the footprint regions based on the two

observed bias spikes in each footprint region (located 7bp to the right of the left boundary and

7bp to the left of the right boundary of the footprint, labeled as -0 and +0 in Fig. 4c-h). The

center regions between the bias spikes were scaled to 4 bins to align footprint regions with

different lengths.

Footprint bias score. The footprint bias score (FBS, in Fig. 4i-k) was defined as the difference

between "spike bias" and "center bias". The "spike bias" was calculated as the average of the

two SELMA bias scores at the spike positions, while the "center bias" was calculated as the

median SELMA bias score at the rest of the positions in the footprint region. Let FBS_j, b_j, and c_j

denote the footprint bias score, spike bias and center bias of footprint j, respectively. The FBS,

spike bias, and center bias are given by:

$$FBS_j = b_j - c_j \tag{5}$$

$$b_j = \overline{a_{i \in B}^{s \in \{+,-\}}} \tag{6}$$

$$c_j = \widetilde{a_{i \in C}^{s \in \{+,-\}}} \tag{7}$$

where a_i^s represents the SELMA bias score at genomic position i on the strand $s \in \{+, -\}$,

overbar represents the average, tilde represents the median, and B and C represent spike

positions (within the flanking 1bp of the bias spikes) and the remaining positions of the footprint

j, respectively. To calculate the randomly shuffled FBS (random k-mer bias), the FBS was

calculated in the same way, but the bias score a_i^s for each position was randomly selected from

the SELMA bias score table.

Calculation of footprint score with different methods. The following methods were used to

calculate footprint scores for comparison:

- Raw footprint: The raw footprint score was calculated following a previous study[13], using

  the formula $f = -\left(\log\frac{n_C+1}{n_R+1} + \log\frac{n_C+1}{n_L+1}\right)$, where n_C, n_R and n_L denote the DNase

  cleavage count in the motif region, and the flanking regions to the right and left of the

  motif, respectively. The flanks are both the same length as the motif.

● Wellington footprint: We used Wellington[9] (v0.2.0) with default parameters to detect

genome-wide footprints and selected significant footprint regions with p-values < 1e-10.

The output footprint score was assigned to the TF motif overlapping with the footprint

region.

● HINT footprint: We used HINT[11,19] (v0.12.3) with default parameters to detect genome-

wide footprints. For bias correction mode, we used an additional parameter: --bias-

correction. The footprint score of each footprint region was assigned to the overlapping

TF motifs.

Inference of TF binding with different features. For each TF in Fig. 4j, the TF motifs

overlapping with consensus footprint regions were collected as target regions. DNase-seq read

count in the footprint region ("reads"), footprint score from an existing method, and bias score

were used as features in a multinomial logistic regression model to infer TF occupancy at

footprint-overlapping motif regions. For each available method, model training with cross-

validation and predictions were performed using different combination of features: "original

method" refers to reads + footprint score. An additional feature of either SELMA FBS or a

random k-mer bias was added for different models. We used a performance measure

integration approach[11] to evaluate different models' prediction power. For each model, we

calculated the area under the ROC curve (AUROC) at 100%, 10%, and 1% false positive rate

(FPR). We also calculated the area under the precision-recall curve (AUPRC) at 100%, 10%,

and 1% recall. We then combined these six performance measures as a rank score S, defined

as

$$S = \frac{1}{6} \sum_i -\log \frac{r_i}{N+1}$$

where $i = 1, \ldots, 6$ denotes the 6 performance measures, r_i is the rank of a model for

performance measure i, and N is the total number of models.

To calculate the random k-mer bias, we randomly permuted the SELMA k-mer bias table and

generated the k-mer bias table for a "simulated enzyme". We used this "simulated" bias table to

calculate the FBS for each footprint region and performed TF inference modeling similar to what

we did for the DNaseI SELMA FBS. This permutation was performed 100 times to generate 100

performance rank scores for random k-mer bias used as controls.

Motif prediction power of TF binding sites. In Fig. 4j-k, we collected the genome-wide motif

sites overlapping with consensus footprint regions and the ChIP-seq peaks for each TF with

HOCOMOCO[36] motif and ChIP-seq data available in human cell lines. We collected data from

all human cell lines with both DNase-seq and more than 20 TF ChIP-seq samples available

from ENCODE, resulting in 7 cell lines, 375 TF ChIP-seq samples, and 156 TFs (Table S1). In

total, we surveyed genome-wide motif sites for 156 TFs, and 61,531,309 motif sites in total. In

Fig. 4k, for each TF, we selected two subgroups of its motif sites based on the FBS of their

overlapped footprint regions: the top 10% of motif sites with the highest FBS, defined as "sites

with high-bias footprint"; and the bottom 10% of motif sites with the lowest FBS, defined as

"sites with low-bias footprint". We calculated the proportion of motif sites overlapping with TF

ChIP-seq peaks for each of the two subgroups and plotted on a scatter plot (Fig. 4k). To test

whether low-bias footprints tend to have more TF binding than high-bias footprints for most TFs,

we conducted a t-test comparing the distribution of TF binding log likelihood ratios of low-bias

over high-bias footprints to the standard normal distribution, and the test p-value is labeled in

Fig. 4k.

Single-cell ATAC-seq clustering analysis

Gold standard of single-cell clustering. For single-cell ATAC-seq data in the human

hematopoietic cells and human cell line samples, the cell-type information for each individual

cell was used as the ground truth, or the gold standard. For single-cell ATAC-seq data in the

mouse gut tube sample, the cell-type information was assigned based on label transfer[40] from

the single-cell RNA-seq dataset in the same system[48] (GSE136689), as the "pseudo" ground

truth, or the silver standard. In detail, we integrated scRNA-seq and scATAC-seq data using the

ArchR package[50] (v1.0.1). Individual cells with a high RNA integration score (unconstrained

predicted score < 0.56) were collected as high-quality cells for the analysis. The cutoff of the

RNA integration score (0.56) was determined by maximizing the interclass variance in the RNA

integration score using Otsu's method[51]. Cell types represented by fewer than 10 cells were

discarded. For the 10x Single Cell Multiome datasets, scRNA-seq parts from each sample were

separated and clustered using Seurat (v4.0)[41] with default parameters. The scRNA-seq

clustering results were used as the "pseudo" ground truth for scATAC-seq cell clusters.

Average bias for individual cells. The average bias for each individual cell (median SELMA

cell bias score, median CBS, used in Fig. 5) was calculated as the median of the bias for all the

fragments in the individual cell. The bias for each paired-end fragment was calculated as the

mean of the SELMA bias scores for the 5' end and the 3' end.

Peak detection and peak bias score calculation. We first combined all the single-cell ATAC-

seq reads in the dataset and performed MACS2[47] peak calling with a q-value cutoff of 0.1 to

include all the potential accessible regions in the human hematopoietic cell and human cell line

systems (100,456 peaks detected for human hematopoietic cells and 83,318 peaks for human

cell lines). For the 10X scATAC-seq data in mouse gut tube, the q-value cutoff of 0.01 was used

for peak calling (146,098 peaks detected). Potential accessible regions with fewer than 10

covered reads or more than 4,000 covered reads were discarded in the subsequent analysis. To

consider the effect of intrinsic cleavage bias in scATAC-seq data, we summarized the bias for

each potential accessible region by calculating the peak bias score (PBS), defined as the

median SELMA bias score of all reads (from all individual cells) located in the peak region.

Cell clustering with scATAC-seq data. In the scATAC-seq clustering analysis, as a naïve

method, we performed PCA dimensionality reduction on the accessible regions by individual

cells matrix of normalized read count and kept 60 PCs, followed by k-means clustering. The

number of clusters (k) in k-means clustering was determined as the actual number of cell types

in the dataset. To evaluate the performance of cell clustering, we compared the clustering

results with the gold standard using adjusted rand index (ARI)[39]. We further repeated the k-

means clustering with 100 different random seeds to ensure the stability of the clustering results

and the average ARI of the 100 runs was used to estimate the clustering accuracy. We also

applied several published methods, including APEC[40], Seurat[41], scran[42], and snapATAC[43] with

default parameters, to the same dataset. We selected the top 60 dimensions for those methods

at the dimensional reduction step (PCA, Seurat, scran and snapATAC) to keep more variance

potentially introduced by intrinsic biases. We used the ArchR package to practice the Seurat

and scran clustering methods. To explore the effect of intrinsic cleavage bias on scATAC-seq

analysis, we selected different percentages (from 50% to 99%, with a 1% increment) of peaks

with the lowest PBS (i.e., removing 50%-1% of peaks with the highest PBS) as input to perform

cell clustering (Fig. 6a-f). We also randomly selected the same percentage of peaks as a control

to estimate the relative rank of the clustering performance from using all peaks and using

retained peaks. In detail, for each percentage from 50% to 99%, we randomly sampled peaks

100 times and defined the relative rank as the number of ARIs from random samples that were

less than the ARI from the same percentage of retained peaks or all peaks. The relative ranks

for different percentages from 50% to 99% are summarized and plotted as boxplots in Fig. 6a-f.

SELMA single-cell peak bias correction model for scATAC-seq cell clustering. The goal of

this model is to give a different weight to all scATAC-seq peaks based on the PBS, so that

peaks heavily affected by intrinsic biases have a lower weight while peaks less affected by

intrinsic biases have a higher weight when they are used to perform cell clustering analysis. To

estimate the relative contribution of each peak in a scATAC-seq dataset to the clustering result,

we conducted an analysis of variance (ANOVA) for each identified peak, using the scATAC-seq

read count/signal across different cells as the variable and the known cell-type labels as the

target group labels. The F score from the ANOVA output was used to quantify the contribution

of each peak to the clustering. Within each percentile of peaks based on their PBS ranks, the

median F score of the peaks with ANOVA p-value < 0.05 was used to represent the relative

level of contribution for this percentile of peaks (Fig. S11a-f). Based on the contribution patterns

of the scATAC-seq samples tested, a standard beta distribution was used to model the

percentile weight function:

$$W(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{\mathrm{B}(\alpha, \beta)}$$

730   where x = 0, 0.01, 0.02, ..., 0.99 is the percentile of the peak ranked by PBS; α, β > 1 are the

731   shape parameters; B(α, β) is the beta function:

$$\mathrm{B}(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$$

733   where $\Gamma()$ is the gamma function. The beta distribution was chosen because its probability

734   density function has the following properties: 1) constraints: $W(0)$ = 0 and $W(1)$ = 0; 2)

735   normalization: $\int_0^1 W(x)dx = 1$; 3) asymmetry with mode at $x = \frac{\alpha-1}{\alpha+\beta-2}$. Based on the

736   relative contribution pattern in Fig. S11a-f, the parameters were empirically determined as α = 2,

737   β = 3. Therefore, the weight function becomes

$$W(x) = 12x(1-x)^2$$

739 After the read count in each peak in each cell was weighted using this weight function, the

740 whole read count matrix was scaled back to keep the total read count in the matrix unchanged

741 from the raw data matrix. The adjusted read count matrix then underwent the clustering

742 analysis. For the single-cell analysis tools that require raw scATAC-seq reads as input, including

743 Scran and Seurat in the ArchR package and snapATAC, the adjusted number of reads for each

744 peak in each cell was synthesized from each peak region and assigned to the cell.

746 **Code and data availability**

747 The SELMA package is available at https://github.com/zang-lab/SELMA. User instructions and

748 example data files can be found in the README document. Essential annotation data, analysis

749 results, and scripts are also available in the repository. The mouse gut tube scATAC-seq

750 dataset has been deposited in the Gene Expression Omnibus (GEO) with accession number

751 GSE168373. The 10x Single Cell Multiome datasets are downloaded from 10X Genomics

752 website(https://www.10xgenomics.com/resources/datasets/fresh-embryonic-e-18-mouse-brain-

753 5-k-1-standard-2-0-0, https://www.10xgenomics.com/resources/datasets/pbmc-from-a-healthy-

754 donor-no-cell-sorting-10-k-1-standard-2-0-0,

755 https://www.10xgenomics.com/resources/datasets/fresh-frozen-lymph-node-with-b-cell-

756 lymphoma-14-k-sorted-nuclei-1-standard-2-0-0). Publicly available data used in this study can

757 be found in the GEO and the ENCODE project. Accession numbers for all the GEO and

758 ENCODE data used in the study are listed in Table S1.

760 **Acknowledgements**

761 The authors thank Dr. Tae-Hee Kim for sharing the mouse primitive gut tube scATAC-seq data

762 and Dr. Xiaole Shirley Liu for helpful discussions. The authors acknowledge 10X Genomics for

763 sharing the 10x Single Cell Multiome datasets. This work was supported by US National

764 Institutes of Health (NIH) grants R35GM133712 and K22CA204439 (to C.Z.), R35GM128635 (to

M.J.G.), National Science Foundation grant NSF-2048991 (to T.Z.), UVA Cancer Center Farrow

Fellowship and the NCI Cancer Center Support Grant P30 CA44579 (to S.S.H.).

Author contributions

C.Z. conceived and directed the research. S.S.H., C.A.M., and C.Z. designed the method. L.L.,

M.J.G., K.D., and T.Z. contributed to the method design. S.S.H. implemented the method and

analyzed the data. Q.L. and W.M. helped with the data analysis. S.S.H. and C.Z. wrote the

manuscript. All authors approved the manuscript.

Competing interests

775 The authors declare that they have no competing interests.

Supplementary information

779 Supplementary figures (Fig. S1-S11).

References

1   Keene, M. A., Corces, V., Lowenhaupt, K. & Elgin, S. C. DNase I hypersensitive sites in

Drosophila chromatin occur at the 5' ends of regions of transcription. *Proc Natl Acad Sci*

U S A **78**, 143-146, doi:10.1073/pnas.78.1.143 (1981).

2   McGhee, J. D., Wood, W. I., Dolan, M., Engel, J. D. & Felsenfeld, G. A 200 base pair

region at the 5' end of the chicken adult beta-globin gene is accessible to nuclease

digestion. *Cell* **27**, 45-55, doi:10.1016/0092-8674(81)90359-7 (1981).

3   Boyle, A. P. *et al.* High-resolution mapping and characterization of open chromatin

across the genome. *Cell* **132**, 311-322, doi:10.1016/j.cell.2007.12.014 (2008).

4   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J.

Transposition of native chromatin for fast and sensitive epigenomic profiling of open

chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* **10**, 1213-1218,

doi:10.1038/nmeth.2688 (2013).

5   Hesselberth, J. R. *et al.* Global mapping of protein-DNA interactions in vivo by digital

genomic footprinting. *Nat Methods* **6**, 283-289, doi:10.1038/nmeth.1313 (2009).

6   Neph, S. *et al.* An expansive human regulatory lexicon encoded in transcription factor

footprints. *Nature* **489**, 83-90, doi:10.1038/nature11212 (2012).

7   Vierstra, J. *et al.* Global reference mapping of human transcription factor footprints.

Nature **583**, 729-736, doi:10.1038/s41586-020-2528-x (2020).

8   Karabacak Calviello, A., Hirsekorn, A., Wurmus, R., Yusuf, D. & Ohler, U. Reproducible

inference of transcription factor footprints in ATAC-seq and DNase-seq datasets using

protocol-specific bias modeling. *Genome Biol* **20**, 42, doi:10.1186/s13059-019-1654-y

(2019).

9   Piper, J. *et al.* Wellington: a novel method for the accurate identification of digital

genomic footprints from DNase-seq data. *Nucleic Acids Res* **41**, e201,

doi:10.1093/nar/gkt850 (2013).

10 Sherwood, R. I. *et al.* Discovery of directional and nondirectional pioneer transcription

factors by modeling DNase profile magnitude and shape. *Nat Biotechnol* **32**, 171-178,

doi:10.1038/nbt.2798 (2014).

11 Li, Z. *et al.* Identification of transcription factor binding sites using ATAC-seq. *Genome*

Biol **20**, 45, doi:10.1186/s13059-019-1642-2 (2019).

12   Bentsen, M. *et al.* ATAC-seq footprinting unravels kinetics of transcription factor binding

during zygotic genome activation. *Nat Commun* **11**, 4267, doi:10.1038/s41467-020-

18035-1 (2020).

13   He, H. H. *et al.* Refined DNase-seq protocol and data analysis reveals intrinsic bias in

transcription factor footprint identification. *Nat Methods* **11**, 73-78,

doi:10.1038/nmeth.2762 (2014).

14   Martins, A. L., Walavalkar, N. M., Anderson, W. D., Zang, C. & Guertin, M. J. Universal

correction of enzymatic sequence bias reveals molecular signatures of protein/DNA

interactions. *Nucleic Acids Res* **46**, e9, doi:10.1093/nar/gkx1053 (2018).

15   Meyer, C. A. & Liu, X. S. Identifying and mitigating bias in next-generation sequencing

methods for chromatin biology. *Nat Rev Genet* **15**, 709-721, doi:10.1038/nrg3788

(2014).

16   Sung, M. H., Guertin, M. J., Baek, S. & Hager, G. L. DNase footprint signatures are

dictated by factor dynamics and DNA sequence. *Mol Cell* **56**, 275-285,

doi:10.1016/j.molcel.2014.08.016 (2014).

17   Yardimci, G. G., Frank, C. L., Crawford, G. E. & Ohler, U. Explicit DNase sequence bias

modeling enables high-resolution transcription factor footprint detection. *Nucleic Acids*

Res **42**, 11865-11878, doi:10.1093/nar/gku810 (2014).

18   Baek, S., Goldstein, I. & Hager, G. L. Bivariate Genomic Footprinting Detects Changes

in Transcription Factor Activity. *Cell Rep* **19**, 1710-1722,

doi:10.1016/j.celrep.2017.05.003 (2017).

19   Gusmao, E. G., Allhoff, M., Zenke, M. & Costa, I. G. Analysis of computational

footprinting methods for DNase sequencing experiments. *Nat Methods* **13**, 303-309,

doi:10.1038/nmeth.3772 (2016).

20   Yan, F., Powell, D. R., Curtis, D. J. & Wong, N. C. From reads to insight: a hitchhiker's

guide to ATAC-seq data analysis. *Genome Biol* **21**, 22, doi:10.1186/s13059-020-1929-3

(2020).

21   Buenrostro, J. D. *et al.* Single-cell chromatin accessibility reveals principles of regulatory

variation. *Nature* **523**, 486-490, doi:10.1038/nature14590 (2015).

22   Cusanovich, D. A. *et al.* Multiplex single cell profiling of chromatin accessibility by

combinatorial cellular indexing. *Science* **348**, 910-914, doi:10.1126/science.aab1601

(2015).

23   Lake, B. B. *et al.* Integrative single-cell analysis of transcriptional and epigenetic states

in the human adult brain. *Nat Biotechnol* **36**, 70-80, doi:10.1038/nbt.4038 (2018).

24   Preissl, S. *et al.* Single-nucleus analysis of accessible chromatin in developing mouse

forebrain reveals cell-type-specific transcriptional regulation. *Nat Neurosci* **21**, 432-439,

doi:10.1038/s41593-018-0079-3 (2018).

25   Zhang, C. T. & Zhang, R. Analysis of distribution of bases in the coding sequences by a

diagrammatic technique. *Nucleic Acids Res* **19**, 6313-6317, doi:10.1093/nar/19.22.6313

(1991).

26   Stormo, G. D. Maximally efficient modeling of DNA sequence motifs at all levels of

complexity. *Genetics* **187**, 1219-1224, doi:10.1534/genetics.110.126052 (2011).

27   Adey, A. *et al.* Rapid, low-input, low-bias construction of shotgun fragment libraries by

high-density in vitro transposition. *Genome Biol* **11**, R119, doi:10.1186/gb-2010-11-12-

r119 (2010).

28   Sos, B. C. *et al.* Characterization of chromatin accessibility with a transposome

hypersensitive sites sequencing (THS-seq) assay. *Genome Biol* **17**, 20,

doi:10.1186/s13059-016-0882-7 (2016).

29   Corces, M. R. *et al.* Lineage-specific and single-cell chromatin accessibility charts

863   human hematopoiesis and leukemia evolution. *Nat Genet* **48**, 1193-1203,

864   doi:10.1038/ng.3646 (2016).

865   30   Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables

866   interrogation of frozen tissues. *Nat Methods* **14**, 959-962, doi:10.1038/nmeth.4396

867   (2017).

868   31   Caron, F., Jacq, C. & Rouviere-Yaniv, J. Characterization of a histone-like protein

869   extracted from yeast mitochondria. *Proc Natl Acad Sci U S A* **76**, 4265-4269,

870   doi:10.1073/pnas.76.9.4265 (1979).

871   32   DeFrancesco, L. & Attardi, G. In situ photochemical crosslinking of HeLa cell

872   mitochondrial DNA by a psoralen derivative reveals a protected region near the origin of

873   replication. *Nucleic Acids Res* **9**, 6017-6030, doi:10.1093/nar/9.22.6017 (1981).

874   33   Kornberg, R. D. Chromatin structure: a repeating unit of histones and DNA. *Science* **184**,

875   868-871, doi:10.1126/science.184.4139.868 (1974).

876   34   Mei, S. *et al.* Cistrome Data Browser: a data portal for ChIP-Seq and chromatin

877   accessibility data in human and mouse. *Nucleic Acids Res* **45**, D658-D662,

878   doi:10.1093/nar/gkw983 (2017).

879   35   Zheng, R. *et al.* Cistrome Data Browser: expanded datasets and new tools for gene

880   regulatory analysis. *Nucleic Acids Res* **47**, D729-D735, doi:10.1093/nar/gky1094 (2019).

881   36   Kulakovskiy, I. V. *et al.* HOCOMOCO: towards a complete collection of transcription

882   factor binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic*

883   *Acids Res* **46**, D252-D259, doi:10.1093/nar/gkx1106 (2018).

884   37   Buenrostro, J. D. *et al.* Integrated Single-Cell Analysis Maps the Continuous Regulatory

885   Landscape of Human Hematopoietic Differentiation. *Cell* **173**, 1535-1548 e1516,

886   doi:10.1016/j.cell.2018.03.074 (2018).

38   Smith, R. J. *et al.* Single-cell chromatin profiling of the primitive gut tube reveals

regulatory dynamics underlying lineage fate decisions. *Nature Communications* **13**,

2965, doi:10.1038/s41467-022-30624-w (2022).

39   Rand, W. M. Objective Criteria for the Evaluation of Clustering Methods. *Journal of the*

America Statistical Association **66**, 846-850, doi:10.2307/2284239 (1971).

40   Li, B. *et al.* APEC: an accesson-based method for single-cell chromatin accessibility

analysis. *Genome Biol* **21**, 116, doi:10.1186/s13059-020-02034-y (2020).

41   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902

e1821, doi:10.1016/j.cell.2019.05.031 (2019).

42   Lun, A. T., McCarthy, D. J. & Marioni, J. C. A step-by-step workflow for low-level

analysis of single-cell RNA-seq data with Bioconductor. *F1000Res* **5**, 2122,

doi:10.12688/f1000research.9501.2 (2016).

43   Fang, R. *et al.* Comprehensive analysis of single cell ATAC-seq data with SnapATAC.

Nat Commun **12**, 1337, doi:10.1038/s41467-021-21583-9 (2021).

44   Zhao, Y., Ruan, S., Pandey, M. & Stormo, G. D. Improved models for transcription factor

binding site identification using nonindependent interactions. *Genetics* **191**, 781-790,

doi:10.1534/genetics.112.138685 (2012).

45   Zang, C. *et al.* High-dimensional genomic data bias correction and data integration using

MANCIE. *Nat Commun* **7**, 11305, doi:10.1038/ncomms11305 (2016).

46   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

9, 357-359, doi:10.1038/nmeth.1923 (2012).

47   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137,

doi:10.1186/gb-2008-9-9-r137 (2008).

48   Hu, S. *et al.* ncHMR detector: a computational framework to systematically reveal non-

classical functions of histone modification regulators. *Genome Biol* **21**, 48,

doi:10.1186/s13059-020-01953-0 (2020).

49   Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given

motif. *Bioinformatics* **27**, 1017-1018, doi:10.1093/bioinformatics/btr064 (2011).

50   Granja, J. M. *et al.* ArchR is a scalable software package for integrative single-cell

chromatin accessibility analysis. *Nat Genet* **53**, 403-411, doi:10.1038/s41588-021-

00790-6 (2021).

51   Nobuyuki, O. A Threshold Selection Method from Gray-Level Histograms. *IEEE*

Transactions on Systems, Man, and Cybernetics **9**, 62-66,

doi:10.1109/TSMC.1979.4310076. (1979).

a

accessible chromatin region

DNA

ACAAC**GTCA**TTAGCAT

k-mer sequence
at cleavage site

(naive) cleavage bias for k-mer j

$= \log_2 \frac{N_j}{M_j}$

N_j : number of cleavages on k-mer j

M_j : number of occurrences for k-mer j

— nucleosome

— transcription factor

↓ — DNaseI/Tn5 cleavage

b

$AA = A \otimes A = \quad 1 \ -1 \ -1 \quad -1 \quad 1 \quad 1 \quad -1 \quad 1 \quad 1$
$AC = A \otimes C = \ -1 \quad 1 \ -1 \quad 1 \ -1 \quad 1 \quad 1 \ -1 \quad 1$
$AG = A \otimes G = \ -1 \ -1 \quad 1 \quad 1 \quad 1 \ -1 \quad 1 \quad 1 \ -1$
$AT = A \otimes T = \quad 1 \quad 1 \quad 1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1$
$CA = C \otimes A = \ -1 \quad 1 \quad 1 \quad 1 \ -1 \ -1 \ -1 \quad 1 \quad 1$
$CC = C \otimes C = \quad 1 \ -1 \quad 1 \ -1 \quad 1 \ -1 \quad 1 \ -1 \quad 1$
$CG = C \otimes G = \quad 1 \quad 1 \ -1 \ -1 \ -1 \quad 1 \quad 1 \quad 1 \ -1$
$CT = C \otimes T = \ -1 \ -1 \ -1 \quad 1 \quad 1 \quad 1 \ -1 \ -1 \ -1$
$GA = G \otimes A = \ -1 \quad 1 \quad 1 \ -1 \quad 1 \quad 1 \quad 1 \ -1 \ -1$
$GC = G \otimes C = \quad 1 \ -1 \quad 1 \quad 1 \ -1 \quad 1 \ -1 \ -1 \ -1$
$GG = G \otimes G = \quad 1 \quad 1 \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \quad 1$
$GT = G \otimes T = \ -1 \ -1 \ -1 \ -1 \ -1 \ -1 \quad 1 \quad 1 \quad 1$
$TA = T \otimes A = \quad 1 \ -1 \ -1 \quad 1 \ -1 \ -1 \quad 1 \ -1 \ -1$
$TC = T \otimes C = \ -1 \quad 1 \ -1 \ -1 \quad 1 \ -1 \ -1 \quad 1 \ -1$
$TG = T \otimes G = \ -1 \ -1 \quad 1 \ -1 \ -1 \quad 1 \ -1 \ -1 \quad 1$
$TT = T \otimes T = \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1$

$A = \quad 1 \ -1 \ -1$
$C = \ -1 \quad 1 \ -1$
$G = \ -1 \ -1 \quad 1$
$T = \quad 1 \quad 1 \quad 1$

$p = 1 + 3k + 9(k-1) = 12k - 8$

c

linear regression

naive k-mer
bias (y)

encoding
parameters (X)

$\begin{pmatrix} -0.04 \\ -1.01 \\ 0.16 \\ \vdots \\ -1.61 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 & -1 & \cdots & 1 \\ 1 & -1 & 1 & \cdots & 1 \\ 1 & -1 & 1 & \cdots & -1 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 1 & 1 & 1 & \cdots & 1 \end{pmatrix}$

$p = 12k - 8$

$y = \sum_{i=1}^{p} a_i X_i$

SELMA cleavage bias
$= \hat{y}$ (fitted value of the regression)

DNase-seq on chromatin DNA

d naive K-mer

R = 0.947

GM12878 Bias score

Bias score K562

e SELMA

R = 0.981

GM12878 Bias score

Bias score K562

ATAC-seq on chromatin DNA

f naive K-mer

R = 0.947

GM12878 Bias score

Bias score K562

g SELMA

R = 0.997

GM12878 Bias score

Bias score K562

DNase-seq on naked DNA

h naive K-mer

R = 0.958

MCF7 Bias score

Bias score K562

i SELMA

R = 0.993

MCF7 Bias score

Bias score K562

ATAC-seq on naked DNA

j naive K-mer

R = 0.937

Fruit fly Bias score

Bias score Human

k SELMA

R = 0.963

Fruit fly Bias score

Bias score Human

l GM12878 DNase-seq

R (OBS vs. EXP)

4mer 6mer 8mer 10mer 12mer

m K562 DNase-seq

R (OBS vs. EXP)

4mer 6mer 8mer 10mer 12mer

n GM12878 ATAC-seq

R (OBS vs. EXP)

4mer 6mer 8mer 10mer 12mer

o K562 ATAC-seq

R (OBS vs. EXP)

4mer 6mer 8mer 10mer 12mer

Figure 1. SELMA framework for cleavage bias estimation. (**a**) Schematic of a naïve k-mer model for cleavage bias estimation. (**b**) Simplex encoding model. The coordinates of the 4 tetrahedral vertices of the cube encode the 4 nucleotides. Each dinucleotide is encoded as the outer product of the 2 mononucleotides. *p* represents the number of parameters in a k-mer simplex encoding model. (**c**) SELMA uses k-mer simplex encoding followed by a linear regression for k-mer cleavage bias estimation. (**d-k**) Comparison between the naïve k-mer model (**d,f,h,j**) and SELMA (**e,g,l,k**) on k-mer cleavage bias scores estimated from DNase-seq data (**d,e,h,i**) and ATAC-seq data (**f,g,j,k**). Each dot in a scatter plot represents an 8-mer, with its estimated bias score from different datasets represented in the x- and y-axes. Chromatin DNA (**d-g**) and naked DNA (**h-k**) from different human cell lines (**d-i**) or different species (**j,k**) were compared as labeled. R represents the Pearson correlation coefficient. (**l-o**) Intrinsic cleavage bias estimation accuracy measured by correlation between genome-wide observed (OBS) and bias-expected (EXP) cleavages with different k-mers. A higher Pearson correlation coefficient (R) indicates a better prediction of the observed cleavage profile using the estimated biases.

Figure 2

Figure 2. SELMA consideration of dimeric Tn5 cleavages for ATAC-seq. (**a**) Schematic of SELMA consideration of dimeric Tn5 cleavages. (**b**,**c**) Cross-correlation between 5' cleavages and 3' cleavages across genome-wide accessible chromatin regions in the human GM12878 cell line from ATAC-seq (**b**) and DNase-seq (**c**) experiments. The x-axis represents the shift distance (in bp) between 5' and 3' cleavages. (**d-i**) Comparison of ATAC-seq intrinsic cleavage bias estimation accuracy measured by Pearson correlation coefficients (R) between genome-wide observed (OBS) and bias-expected (EXP) cleavages for human GM12878 (**d**) and K562 (**e**) cell lines, as well as several modified Tn5-based assays (**f-i**). Different bars represent different estimation approaches: gray for considering the 10-mer at the observed cut only (5' only); orange for SELMA; and blue for several published approaches. Modified Tn5-based assays include (**f**) THS-seq with standard Tn5; (**g**) THS-seq with mutated Tn5; (**h**) fast-ATAC-seq; and (**i**) omni-ATAC-seq.

Figure 3

Figure 3. SELMA bias estimation using mtDNA reads. (**a**) Conventional approach to intrinsic cleavage bias estimation, in which a naked DNA sample is required for bias estimation, and mtDNA reads from the chromatin DNA ATAC-seq experiment are discarded. (**b**) SELMA approach of intrinsic cleavage bias estimation using mtDNA reads. Chromatin reads and mtDNA reads from the same ATAC-seq dataset were separated for Tn5 cleavage profiling and bias estimation with SELMA, respectively. (**c-d**) Scatter plots demonstrating the robustness of 10-mer bias scores estimated from mtDNA (mtDNA reads from chromatin ATAC-seq data, x-axis) and genomic DNA (reads from naked DNA ATAC-seq data, y-axis), comparing the naïve k-mer model (**c**) and SELMA (**d**). SELMA shows a more robust estimation using mtDNA reads with higher correlation than the naïve k-mer model. (**e**) Comparison of the correlation coefficients between biases estimated from mtDNA reads and from naked DNA with different mtDNA read counts.

Figure 4

Figure 4. TF binding inference on DNaseI footprint regions is affected by intrinsic cleavage biases and improved by SELMA. (**a,b**) Heatmaps of DNaseI cleavage patterns in the GM12878 cell line (**a**) and SELMA-estimated bias scores (**b**) around ENCODE DNaseI consensus footprint regions. 5' and 3' patterns were plotted separately. Footprint regions were consistently ranked by footprint length, and signals for every 1000 regions with similar lengths were averaged as one row in the heatmap. (**c-h**) Aggregate plots of DNaseI cleavage patterns (**c,e,g**) and SELMA-estimated bias scores (**d,f,h**) at all ENCODE DNaseI consensus footprint regions (**c,d**), footprint regions overlapping with TF binding hotspots (**e,f**), and footprint regions overlapping with TF binding deserts (**g,h**). DNaseI cleavages are from GM12878 for (**c & e**) and merged from GM12878, K562 and ENCODE tissues (**g**). Dashed lines represent estimated bias spikes in the footprint regions. (**i**) SELMA-estimated footprint bias scores (FBS) for different groups of footprint regions. P-values were calculated by the Wilcoxon signed-rank test. The centerline, bounds of box, top line and bottom line represent the median, 25th to 75th

percentile range, 25th percentile - 1.5*interquartile range (IQR) and 75th percentile + 1.5*IQR, respectively. (**j**) Difference of performance rank scores for transcription factor binding inference from DNaseI footprint using various methods for the human GM12878 cell line. Boxplots with different colors represent different approaches as indicated in the legends. Different sets of boxplots represent footprint scores calculated by different published methods as labeled. P-values were calculated by the one-sided Wilcoxon signed-rank test. (**k**) Scatter plot showing the heterogeneous effect of intrinsic cleavage biases on different TF motifs. Each data point represents motif sites for one TF in one cell type. Fractions of the TF-bound motif sites overlapping with high-FBS footprint regions and with low-FBS footprint regions are plotted on the x- and y-axes, respectively. More motifs located above the diagonal line indicate that more TFs are more likely to bind at low-FBS sites than at high-FBS sites. The p-value was calculated by one-tailed t-test comparing the distribution of log likelihood ratios (y-axis/x-axis) to the standard normal distribution.

Figure 5

Figure 5. Intrinsic cleavage biases affect single-cell ATAC-seq data analysis. Visualization of intrinsic cleavage bias effect in different cell clusters derived from scATAC-seq data for different biological samples and different experimental platforms: human hematopoietic cells (**a-c**), human cell lines (**d-f**), mouse primitive gut tube (**g-i**), and 10x Single Cell Multiome data for mouse embryonic brain (**j-l**), human peripheral blood mononuclear cells (PBMC) (**m-o**), and human lymph node (**p-r**). (**a,d,g,j,m,p**) UMAP visualization where cells are colored by cell type labels/clusters. (**b,e,h,k,n,q**) Same UMAP visualization but cells are colored by cell bias score (CBS). (**c,f,i,l,o,r**) CBS distributions of cells from different cell types/batches/clusters. Boxes are colored by cell clusters using the same color palette as the first column.

Figure 6

Figure 6. SELMA bias correction model improves single-cell ATAC-seq cell clustering. (**a-f**) Comparisons of cell clustering accuracy before and after considering the peak bias score (PBS) on scATAC-seq data for human hematopoietic cells (**a**), human cell lines (**b**), mouse primitive gut tube (**c**), and 10x Single Cell Multiome data for mouse embryonic brain (**d**), human PBMC (**e**), and human lymph node (**f**). K-means clustering was performed after PCA dimensionality reduction. Blue, using all ATAC-seq peaks; orange, after removing 1%-50% of peaks with the highest peak bias score (PBS). For each percentage of peaks retained from 50% to 99% with a 1% increment, 100 randomly sampled subsets of peaks were used as the background for determining the relative ranks of all peaks or retained peaks. The relative ranks of the adjusted Rand index (ARI), defined as the rank relative to the 100 randomly sampled sub-datasets, for the 50 cases from 50% to 99%, were plotted. (**g**) Schematic of SELMA single-cell peak bias correction model. Peaks are weighted and adjusted based on PBS percentile using an empirically determined weight function. (**h-m**) Comparisons of the accuracy (measured by ARI) of single cell clustering generated using different existing tools on scATAC-seq data with (orange) or without (blue) SELMA single-cell peak bias correction. Each panel represents the result for an scATAC-seq sample from a different biological system or experimental platform: human hematopoietic cells (**h**), human cell lines (**i**), mouse gut tube (**j**), mouse embryonic brain (**k**), human PBMC (**l**) and human lymph node (**m**). Each dot in each panel represents the ARI for a different method labeled on the x-axis. Blue, clustering with uncorrected raw data; orange, clustering with SELMA-corrected data.

Supplementary Figure S1

Supplementary Figure S1. Improvement of bias estimation of the SELMA model. (**a**) Comparison of intrinsic cleavage bias estimated in different data types. Each dot in a scatter plot represents an 8-mer, with its estimated bias score from K562 and GM12878 cell lines represented in x and y axes, respectively. Different plots represent bias estimated using data from different experiments (ATAC-seq, DNase-seq, Faire-seq, and ChIP-seq). R represents the Pearson correlation coefficient. (**b**) Comparison of Tn5 intrinsic cleavage bias estimated from different groups of fragments. The top panel was for the 8-mer bias estimated from the naïve k-mer model, and the bottom panel was from the SELMA. The fragments were separated based on the region's chromatin accessibility, illustrated in the schematic on the right, and labeled at the axes. Fragments located in the top 20% chromatin accessible regions with the highest accessibility were assigned as group1 (G1). Then fragments located in the top 20%~40%, 40%~60%, 60%~80%, and bottom 20% accessible regions were assigned as G2, G3, G4, and G5, respectively. (**c**) Similar to (b) but for DNase-seq bias score. (**d, e**) Barplots compared the Pearson correlation coefficients (R) calculated in (b, c), respectively. The blue bars were for the naïve k-mer model, and the red bars were for the SELMA model.

Supplementary Figure S2

Supplementary Figure S2. Intrinsic cleavage bias estimation accuracy with different k-mer. Intrinsic cleavage bias estimation accuracy is measured by the correlation between genome-wide observed (OBS) and bias-expected (EXP) cleavages with different k-mers for ENCODE DNase-seq data (**a**) and ENCODE ATAC-seq data (**b**), respectively. Different rows represent data from different tissues. The Pearson correlation coefficients (R) in the same cell type (each row) were z-normalized and plotted as heat to compare the relative performance of different SELMA k-mer models. The higher the normalized score, the better performance the model had.

Supplementary Figure S3

Supplementary Figure S3. SELMA consideration of dimeric Tn5 cleavages for ATAC-seq. (**a**) The average profile of the ATAC-seq cleavage pattern around the CTCF motif center. +/- strand reads were separated and plotted in red and blue lines, respectively. The left panel is for unmodified observed Tn5 cleavage. In the right panel, the - strand cleavage pattern was shifted 9bp towards upstream. (**b**) similar to (a) but for the NRF1 motif. A perfect match of + and – strand profiles can be observed for the 9bp shift. (**c-d**) Comparison of DNase-seq intrinsic cleavage bias estimation accuracy measured by Pearson correlation coefficient (R) between genome-wide observed and bias-expected cleavages for human GM12878 (c) and K562 (d) cell lines. Different bars represent different estimation approaches: grey for considering the 10-mer at the observed cut only (5' only); orange for SELMA. The published approaches (blue bars in Fig. 2d-e) were not shown for DNase-seq since they were specifically designed for ATAC-seq. (**e-j**) Pearson correlation coefficient between genome-wide observed and bias-expected cleavages for ENCODE ATAC-seq data from different tissues. (**k-n**) Comparison between bias scores estimated from ATAC-seq and modified Tn5 based techniques THS-seq with standard Tn5 (k); THS-seq with mutated Tn5 (i); fast-ATAC-seq (m); and omni-ATAC-seq (n).

Supplementary Figure S4

Supplementary Figure S4. SELMA bias estimation using mtDNA reads. (**a**) Bar plots demonstrating the Pearson correlation coefficients (y-axis) between 10-mer bias scores estimated from mtDNA (mtDNA reads from chromatin ATAC-seq data generated from ENCODE project in different cell types) and genomic DNA (reads from naked DNA ATAC-seq data), comparing the naïve k-mer model (blue bars) and SELMA (red bars). (**b**) Similar to (a) but for ENCODE DNase-seq data in the same cell types. (**c**) Scatter plots demonstrating the consistence of 10-mer bias scores estimated from mtDNA (with different fragments count, x-axis) and genomic DNA (reads from naked DNA ATAC-seq data, y-axis). The Pearson correlation coefficients were labeled in the top left corner and used in Fig. 3e.

Supplementary Figure S5

Supplementary Figure S5. TF binding inference on DNaseI footprint regions is affected by intrinsic cleavage biases and improved by SELMA. (**a-b**) Heatmaps of DNaseI cleavage patterns in the K562 cell line. (**c**) Aggregate plots of DNase-seq cleavage patterns at ENCODE DNaseI footprint regions with different length categories (15, 20, 25, 30, and 35bp). The dotted lines represented the boundary of the footprint regions. (**d**) similar to (c) but for SELMA-estimated bias scores. (**e**) TF binding hotspots and TF binding deserts were defined based on the number of human TF ChIP-seq samples that have peaks overlapping with the footprint regions. The footprint regions overlapping with peaks from >= 3000 ChIP-seq samples were defined as TF binding hotspots, while the footprint regions that do not overlap with any peak from any TF ChIP-seq sample nor any motif sites were defined as TF binding deserts.

Supplementary Figure S6

Supplementary Figure S6. Aggregate plots of DNaseI cleavage patterns of all ENCODE DNaseI consensus footprint regions (**a**) and footprint regions overlapping with TF binding hotspots (**b**). Different columns represent ENCODE DNaseI-seq data from K562 cell lines and tissues.

Supplementary Figure S7

Supplementary Figure S7. Performance rank scores for transcription factor binding prediction from DNaseI footprint using various methods for the human cell lines. (**a**) Number of transcription factors (TFs) covered vs. Number of cell lines included in the analysis. Cell lines are ranked by the number of ChIP-seq samples available. (**b-h**) Difference of performance rank scores for transcription factor binding inference from DNaseI footprint using various methods for DNaseI-seq data from different cell lines K562 (b), GM12878 (c), HepG2 (d), A549 (e), H1 (f), HeLaS3 (g), and MCF7 (h). Boxplots with different colors represent different approaches as indicated in the legends. Different sets of boxplots represent footprint scores calculated by different published methods as labeled. P-values were calculated by the one-sided Wilcoxon signed-rank test. The detailed list of rank scores can be found in Supplementary Table 2.

Supplementary Figure S8

Supplementary Figure S8. Rank score changes for all TFs in each cell line. Barplots showed the delta relative rank of TFs from data used in Fig. 4j and Supplementary Fig. S7. Different panels were for TF ChIP-seq data in different cell lines: K562 (**a**), GM12878 (**b**), HepG2 (**c**), A549 (**d**), H1 (**e**), HeLaS3 (**f**), and MCF7 (**g**). The full list can be found in Supplementary Table 2.

Supplementary Figure S9

Supplementary Figure S9. Scatter plots comparing the SELMA bias estimated from bulk naked DNA ATAC-seq data (y-axis) and from different scATAC-seq datasets (x-axis) for human hematopoietic cells (**a**), human cell lines (**b**), mouse primitive gut tube (**c**), and 10x Single Cell Multiome data for mouse embryonic brain (**d**), human PBMC (**e**), and human lymph node (**f**). The bias scores for (a-c) were calculated using mtDNA reads. The bias scores for (d-f) were calculated using chromatin reads because the mtDNA reads were automatically excluded in the fragment data files preprocessed by the 10x Cell Ranger ARC pipeline. R represents the Pearson correlation coefficient.

Supplementary Figure S10

Supplementary Figure S10. Improvement of cell clustering accuracy after excluding scATAC-seq peaks with high PBS. Y-axis represents the percentage changes of the adjusted rand index (ARI) between K-means clustering results and the ground truth cell labels. X-axis represents the percent of peaks retained for clustering (50%~99%) after the peaks with highest PBS were removed. Different panels represent data from different biological systems: human hematopoietic cells (**a**), human cell lines (**b**), mouse gut tubes (**c**), mouse embryonic brain (**d**), human PBMC (**e**), and human lymph node (**f**).

Supplementary Figure S11

Supplementary Figure S11. SELMA single-cell bias correction model development. (**a-f**) ANOVA results for predicting cell type labels with the scATAC-seq signal. An F score was calculated for each peak to measure the importance of the peak to cell type classification. Peaks were then ranked by PBS score, and the median F score of the peaks in each PBS percentile plotted. Different panels represent data from different biological systems: human hematopoietic cells (**a**), human cell lines (**b**), mouse gut tubes (**c**), mouse embryonic brain (**d**), human PBMC (**e**), and human lymph node (**f**). (**g**) PBS percentile weight function used in the SELMA single cell bias correction model.