Fatigue induces behavioural improvements in the unfatigued hand via altered functional connectivity and neurochemicals in cortical motor areas

Justin W. Andrushko[1,2], Jacob M. Levenstein[2,3], Catharina Zich[2,4], Evan C. Edmond[2], Jon Campbell[2], William T. Clarke[2], Uzay Emir[5], Jonathan P. Farthing[1*], Charlotte J. Stagg[2*]

1. College of Kinesiology, University of Saskatchewan, Canada
2. Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, United Kingdom
3. Section on Functional Imaging Methods, National Institutes of Mental Health, National Institutes of Health, Bethesda, Maryland, United States
4. Department of Clinical and Movement Neurosciences, University College London, United Kingdom
5. School of Health Sciences, College of Health and Human Sciences, Purdue University, United States

* Co-senior authors

Correspondence

Dr. Justin W. Andrushko, PhD

Justin.andrushko@usask.ca

ORCID: https://orcid.org/0000-0003-2258-1689

Keywords

resting-state functional connectivity, handgrip, serial reaction time, cross-education, motor learning

Key points summary

- Right handgrip contractions at 50% MVC improve left hand response times.

- Repeated handgrip contractions at 50% MVC alter interhemispheric functional connectivity.

- Unilateral exercise may be an effective method for enhancing contralateral limb motor performance in rehabilitation settings.

32 **Data sharing**

33 All data will be available from the authors upon reasonable request.

Abstract

In humans, motor learning is underpinned by changes in functional connectivity (FC) across the

sensorimotor network. Unilateral exercise-induced fatigue increases FC in the ipsilateral

primary motor cortex (M1) and supplementary motor area (SMA); areas involved in motor

planning and execution of the contralateral hand. Unilateral fatiguing exercise is therefore a

promising potential approach to augment motor performance in the non-fatigued,

contralateral, hand. In a within-participant, controlled, randomized, cross-over design, 15 right-

handed adults had two magnetic resonance imaging (MRI) sessions, where functional MRI and

MR Spectroscopic Imaging were acquired before and after repeated right-hand contractions at

either 5% or 50% maximum voluntary contraction (MVC). Before and after scanning, response

times (RTs) were determined in both hands, and after scanning, participants performed a serial

reaction time task (SRTT) with their left, unfatigued, hand. Nine minutes of 50% MVC

contractions resulted in fatigue. This unimanual fatigue improved motor performance, as

indexed by decreased RTs, in the contralateral hand. Although fatigue had no significant effects

on sequence learning, fatigue led to a significant increase in the transfer of the learned skill to

the untrained hand. These behavioural effects were supported by significant neural changes:

an increase in SMA-SMA functional connectivity, and increased connectivity between right M1

and right Orbitofrontal Cortex. At a neurochemical level, the degree of fatigue-induced

decrease in GABA in left M1, left and right SMA correlated with subsequent behavioural

improvements in the left-hand. These results support unilateral fatiguing exercise as a

potential therapeutic intervention in a range of neurological and orthopedic conditions.

Introduction

The acquisition of motor skill is essential in our daily lives. Able-bodied individuals may take for granted the abilities to grasp and manipulate objects, or even to safely complete essential activities of daily living. However, many people lose their ability to perform these tasks through a variety of injuries. Restoring function in these individuals is of clear clinical importance, but *how* we can optimally improve behaviour is an open scientific question: both in terms of restoring motor function in orthopedic or neurologically impaired individuals such as stroke survivors, and in healthy populations, such as athletes, looking to maximize motor performance.

One approach showing particular promise to enhance motor skill acquisition is cross-education (CE). CE describes the phenomenon by which unilateral exercise enhances muscle activity, force generation, or skill in the contralateral, homologous muscle (Manca *et al.*, 2021). CE is effective in preserving muscle strength and size in an immobilized limb (Farthing *et al.*, 2009, 2011; Magnus *et al.*, 2010; Pearce *et al.*, 2013; Andrushko *et al.*, 2018; Valdes *et al.*, 2021), and strength and range of motion of a fractured arm (Magnus *et al.*, 2013) when the unaffected limb is trained. Studies have also found CE to be effective for motor recovery in chronic stroke survivors (Dragert & Zehr, 2013; Urbin *et al.*, 2015; Sun *et al.*, 2018; Dehno *et al.*, 2021).

Current evidence suggests cortical mechanisms likely drive CE effects (Ruddy & Carson, 2013; Manca *et al.*, 2018), although it is not yet clear *how* these inter-hemispheric behavioural transfer effects occur. At a systems level studies suggest fatiguing exercise increases functional connectivity in the sensorimotor network, particularly in the ipsilateral (unfatigued) M1 (iM1), ipsilateral SMA, and contralateral SMA (Jiang *et al.*, 2012), and enhances muscle activity in the contralateral unfatigued homologous muscle (Carr *et al.*, 2021). Similarly, high force exercise, which is known to produce large CE effects (Urbin *et al.*, 2015; Pelet & Orsatti, 2021), also increases functional connectivity in the ipsilateral hemisphere (Andrushko *et al.*, 2021).

At a neurochemical level, unilateral fatiguing exercise decreases iM1 $GABA_A$ activity, as

quantified by the Transcranial Magnetic Stimulation protocol Short Intracortical Inhibition (SICI;

(Maruyama *et al.*, 2006; Ni *et al.*, 2007; Takahashi *et al.*, 2009)). GABAergic inhibition decreases

during motor learning (Floyer-Lea *et al.*, 2006; Kolasinski *et al.*, 2019), and the magnitude of this

GABA decrease correlates with the extent of subsequent motor learning (Stagg *et al.*, 2011*a*).

Indeed, decreasing M1 GABA using non-invasive brain stimulation techniques increases motor

skill acquisition, both in healthy adults (Reis *et al.*, 2009; Stagg *et al.*, 2011*b*) and in functional

recovery after stroke (Kim *et al.*, 2010; Allman *et al.*, 2016).

Considering motor network connectivity is at least in part controlled by M1 inhibition (Stagg *et*

al., 2014; Bachtiar *et al.*, 2015), here we acquire two independent measures to test the

hypothesis that fatigue increases motor performance in the unfatigued hand by increasing

interhemispheric homologous connectivity of M1 and SMA, via an altered inhibitory/excitatory

state within these cortical motor regions by increasing glutamate and/or decreasing GABA

(figure 1A).

Methods

This study conforms to the Declaration of Helsinki and was approved by the Oxford Central

University Research Ethics Committee (MSD-IDREC-C1-2014-100 and MSD-IDREC-C1-2014-090).

Using an effect of η^2 = 0.250 based on a previous work (Bachtiar *et al.*, 2018), we computed an

estimated total sample size of 12 (G*Power 3.1.9.2; 1-β = 0.95, α = 0.05). To allow for dropout

and data loss, we therefore recruited fifteen right-handed participants aged between 18-35

years (28.92 ± 3.30 years; 7 Female). Exclusion criteria were: playing a musical instrument to

Associated Board of the Royal Schools of Music Grade 5 or higher, any contraindication to MRI,

any history of neurological or psychiatric disorders.

Experimental Outline

Participants had two experimental sessions where they performed a right handgrip force-
matching task at either 5% or 50% of their maximum voluntary contraction (MVC). During each
experimental session, response times (RTs) of each hand were assessed before and after the
MRI. Resting-state functional MRI (fMRI) and magnetic resonance spectroscopic imaging
(MRSI) scans were acquired before and after a nine minute right handgrip force-matching task
at either 5% or 50% MVC. Finally, outside the scanner participants performed a serial reaction
time task (SRTT) with their left hand, which consisted of three separate segments, interspersed
with two minutes of the same right handgrip force-matching task as in the scanner. This study
had a within-participant repeated measures design, with the order of conditions stratified-
randomized for sex across the group (figure 1B).

Handgrip force-matching task

The handgrip force-matching task was performed using an MRI-compatible hand dynamometer
(Biopac Systems Inc. Aero Camino Goleta, CA) and implemented using in-house code
(Psychtoolbox-3 (Kleiner *et al.*, 2007; The Mathworks Inc, 2018)). Participants performed a
visually cued 0.5 Hz (1 second contraction, 1 second rest) repeated force-matching task at
either 5% or 50% of their MVC, depending on the session. The target force was displayed
halfway up the vertical axis, and participants were instructed to squeeze the dynamometer to
guide a cursor up the screen until it reached the target. During scanning, participants
performed 273 handgrip contractions over the nine-minute task. During the post-MRI motor
learning component of the study, participants performed 4 blocks of 62 contractions each,
which were interleaved with the SRTT task.

Handgrip force-matching task analysis

The area under the curve (AUC) of the contraction force was calculated over a two second
window for every contraction separately. For each participant (and session) a regression line
was then fitted to the AUC measures across all contractions for each participant and session

separately. The β of the regression from each session was used to quantify motor performance,

whereby a β < 0 indicates a decrement in motor performance over time, consistent with

fatigue.

Serial reaction time task

Participants performed a visually-cued serial reaction time task before and after MRI scanning,

implemented in PsychoPy3 (Peirce et al., 2019). Participants responded via a button box (4-

Button Inline, HHSC-1x4-L; Current Designs Inc. Philadelphia, PA USA) with their right or left

hand depending on the block. The task was divided into blocks of 64 self-paced visual cues,

which participants were asked to respond to by pressing the corresponding button on the

button box as quickly and accurately as possible.

In random blocks, cues were presented in a pseudo-random order. In sequence blocks, the

cues consisted of a 16-button sequence repeated four times. Participants were not told about

the presence of a sequence. Different sequences were used for each session and the order was

counter-balanced across the participants.

With each hand separately, participants performed two random blocks at the Pre-MRI Baseline

and Post-MRI timepoints. They then performed three learning segments with their left hand

each following the two-minute submaximal handgrip force-matching task with the right hand.

Each learning segment started with a random block, followed by four sequence blocks. The

post-learning timepoint consisted of three blocks (Random, Sequence, Random), performed

separately with their right and left hands. Finally, following a 15-minute break, retention of the

trained motor sequence was assessed identically to the post-learning timepoint (figure 1B).

SRTT analysis

Incorrect button-press responses, and RTs < 50 ms or > 700 ms were removed. The median re-

sponse time of the remaining button presses was then calculated for each block separately. To

investigate changes in RT due to fatigue, the percent change in RTs for each hand was calcu-

lated using the mean of the medians from the two baseline blocks as the Pre-MRI value, and

similarly, the mean of the medians from the two Post-MRI blocks was used as the Post-MRI

value.

To assess learning, we performed a RM-ANOVA with one factor of condition (5% MVC, 50%

MVC) and one factor of time (15 blocks). At the post-learning and retention timepoints, we

compared the median RT in the sequence block to the mean of the median RTs of the two ran-

dom blocks.

To investigate any significant change in RT between Baseline and Post-MRI, we performed a

RM-ANOVA using the percent change in RT data with one factor of condition and another of

hand.

Magnetic Resonance Imaging

MRI sessions were acquired using a Siemens 3-tesla Prisma whole-body MRI scanner with a 32-

channel head array receive coil (Siemens, Erlangen, Germany). T1-weighted MPRAGE scans

(voxel size 1 mm^3, TR = 1900 ms, TI = 912 ms TE = 3.96 ms, FOV = 232 × 256 × 192 mm^3, flip an-

gle = 8°, TA = 7:21) were acquired for registration purposes and to guide placement of the MRSI

slab, prioritising left M1. Pre- and Post-task resting-state semi-LASER localised density-

weighted concentric-ring trajectory MRSI (voxel size 5 mm × 5 mm × 15 mm, semi-LASER Vol 85

mm x 35 mm x 15 mm, TR = 1400 ms, TE = 32 ms, FOV = 240 mm × 240 mm × 15 mm, TA =

4:30) (Steel et al., 2018), and fMRI (490 volumes, voxel size 2.4 mm^3 with zero gap, TR = 735 ms,

TE = 39 ms, FOV = 210 × 210 × 154 mm, flip angle = 52°, TA = 6:10) (Miller et al., 2016) data

were acquired. Participants were asked to fixate on a grey cross presented on a black screen

and to blink normally. To quantify any potential change in head position after task performance

a single localiser (voxel size 1.4 mm, TR = 3.15 ms, TE = 1.37 ms, FOV = 350 mm × 350 mm × 263

mm, 128 slices, flip angle = 8°, TA = 0:22) was collected prior to Post-task MRSI. Finally, B$_0$ field

maps (voxel size 2 mm^3, TR = 482 ms, TE1 = 4.92 ms, TE2 = 7.38 ms, FOV = 216 mm × 216 mm ×

146 mm, 49 slices, flip angle = 46°, TA = 1:45) were acquired to correct for field distortion in the

resting-state fMRI data.

<u>MRI analysis</u>

fMRI preprocessing

Functional MRI analyses were performed using tools from the FMRIB Software Library (FSL

v6.0.3; (Jenkinson *et al.*, 2012)). Standard preprocessing steps were performed, including

removal of non-brain tissue (BET; (Smith, 2002)), removal of the initial two volumes, motion

correction (MCFLIRT; (Jenkinson *et al.*, 2002)), high-pass temporal filtering at 0.01Hz, and

distortion correction with the implementation of field-maps.

Following single-participant MELODIC pre-processing (v3.15), FMRIB's Independent Component

Analysis (ICA)-based Xnoiseifier (ICA-FIX) was used to automatically denoise the data (Griffanti

et al., 2014; Salimi-Khorshidi *et al.*, 2014).   The UK Biobank training-weights file

(UKBiobank.RData) was used with a threshold value of 20, and 0.01Hz high-pass filtered motion

confound cleanup. Following the automated denoising, all components were manually

inspected before the cleaned data were smoothed with a 5 mm full-width half maximum

(FWHM) kernel.

Individual resting state fMRI scans were first registered to the respective T1 structural scan

using boundary-based registration as implemented in FMRIB's Linear Image Registration Tool

(FLIRT; (Jenkinson & Smith, 2001; Jenkinson *et al.*, 2002)), and then to a standard space

template (MNI152 2 mm) using non-linear registration (FNIRT; (Andersson *et al.*, 2007*a*,

2007*b*)).

fMRI analysis

We assessed resting state functional connectivity using two different approaches: (1) a seed-

based functional connectivity approach to test the hypothesis that unimanual handgrip

contractions lead to changes in connectivity of the ipsilateral right M1 (rM1); and (2) a region of

interest (ROI)-based functional connectivity analysis to specifically address our *a priori*

hypothesis that unimanual fatigue would increase M1-M1 and SMA-SMA connectivity.

216   *(1) Seed-based, whole-brain functional connectivity*

The hand area of the unfatigued, rM1 was defined from previous functional MRI data of hand

movements (Weinrich *et al.*, 2017), and the mean timeseries was extracted from this region for

each participant and each session independently. This was then entered as a regressor into a

lower-level FEAT analysis (Woolrich *et al.*, 2001), and task-related changes in functional

connectivity of rM1 were investigated via a higher level mixed-effects analysis using a cluster

forming threshold of z = 3.1 and *p* = 0.05 (Woolrich *et al.*, 2004).

223   *(2) ROI-ROI-based functional connectivity*

Next, we wanted to specifically assess changes in M1-M1 and SMA-SMA functional

connectivity. The hand area of the M1s was functionally defined as above, and SMA was

defined from tractography-based parcellation (Johansen-Berg *et al.*, 2004).   The mean

timeseries from within each region was extracted for each participant and session separately.

We calculated the Pearson's r values between the homologous pairs using custom in-house

MATLAB scripts, which were then converted to z-scores using a Fishers *r* to z transformation.

The z-scores for the correlation between the homologous pairs were then used in a 2 × 2 × 2

(condition (5%, 50% MVC) × ROI (M1, SMA) × time (pre-task, post-task)) RM-ANOVA.

MRSI

233   MRSI data was reconstructed and pre-processed according to Steel et al. (2018), using

in-house scripts. Reconstruction and pre-processing included: i. metabolite cycling

reconstruction (Emir *et al.*, 2017), ii. coil-combination (Walsh *et al.*, 2000), iii. frequency and iv.

phase shift correction, v. HLSVD for residual water removal (Cabanes *et al.*, 2001), and vi. eddy

current correction using the unsuppressed water signal (Klose, 1990). Concentration of

neurochemicals was quantified as in Steel et al. (2018) using LCModel (Provencher, 2001, 2016).

A chemical shift of 0.5 to 4.2 ppm was evaluated with a basis set containing 20 metabolites and

default LCModel macromolecules with disabled soft constraints on metabolites (NRATIO set to

0), and a baseline stiffness setting (DKMTM) of 0.25 (see Figure 5A for example of raw spectra

and model fits). Each voxel spectrum was independently fit with spectral quality and

assessment of model fitting performed using in-house scrips. For the metabolite maps of

glutamate + glutamine (Glx) and GABA, MRSI voxels with CRLB values > 50 or SNR < 40 were

excluded from further analysis. All metabolite measurements are expressed as a ratio over total

Creatine (tCr).

Metabolite maps were next upsampled to $1mm^3$ resolution using nearest neighbour

interpolation to preserve the original sampling grid. Metabolite maps were then aligned to their

native T1 coordinate space. To correct for potential shift in head position, structural scans

acquired prior to each MRSI sequence were aligned using McFLIRT (Jenkinson et al., 2002). The

metabolite maps were then transformed using the McFLIRT generated registration matrix. Only

shared MRSI voxels from within an intersection mask of QA-passed and head position-aligned

MRSI voxels across each timepoint were included in subsequent analyses. For the ROI analysis,

the four binary ROI masks [left M1 (lM1), right (rM1), left SMA (lSMA) and right SMA (rSMA)]

were non-linearly aligned to native T1 space by inverse warping (3dNwarpApply) the native to

standard space alignment, generated with @SSwarper (Cox, 1996; Saad et al., 2009; Cox &

Glen, 2013). The native space ROIs were thresholded at 0.5 to mitigate partial volume effects.

Within the intersection mask, mean concentration of Glx/tCr and GABA/tCr were then

extracted for each of the four ROIs.

Statistical Analyses

Statistical analyses were run using Jamovi v1.6.9 (The Jamovi Project, 2020). Hedges' g effect

sizes are reported for t-tests and partial eta squared (η_p^2) effect sizes are reported for ANOVA

results. Greenhouse-Geisser corrections were applied as necessary where violations of

sphericity were present.

Results

Performance of the 50% MVC handgrip force-matching task induced fatigue

We first wanted to determine if the 50% MVC handgrip force-matching task induced fatigue. We reasoned that the total grip force would be reduced over time during our 50% MVC fatiguing condition, but not during non-fatiguing movement in our 5% MVC control condition. We therefore quantified the AUC of the force profile for each condition separately and calculated a line of best fit for each participant for each session. In line with our hypothesis, 50% MVC showed a significant decrement in performance over time compared with 5% MVC (50% MVC: β = -0.912 ± 1.070; 5% MVC: β = -0.022 ± 0.107; slope of 50% MVC compared with zero: $t(14)$= -3.302, p = 0.005, g = -0.824; paired t-test $t(14)$= 3.211, p = 0.006, g = 0.801; figure 2A).

Right hand fatigue decreased RTs in the left, unfatigued hand

We next investigated whether performance of a 50% MVC fatiguing force-matching task with the right-hand would induce a behavioural improvement of the unfatigued, left-hand. After running a condition × hand RM-ANOVA with the RT percent change data, we observed a significant condition × hand interaction ($F(1,14)$= 4.84, p = 0.045, η_p^2 = 0.257), but no significant main effects. Next, to further understand the interaction we split the data by hand and ran paired sample t-tests between conditions for each hand. This revealed a significant difference between conditions for the left-hand ($t(14)$= 2.30, p = 0.037, g = 0.574), indicating the improvement in RTs for the 50% MVC condition (Δ = -5.85 ± 7.05%) was different than the 5% MVC condition (Δ = 0.08 ± 5.99%). Further, a one-sample t-test was used to determine the RT improvement in the left-hand for the 50% MVC condition was significantly different from zero ($t(14)$= -3.21, p = 0.006, g = -0.801). There were no significant differences for the right-hand between

the 50% MVC (Δ = -2.02 ± 4.35%) and 5% MVC (Δ = -0.73 ± 3.46%) conditions (t(14)= 0.788, p =

0.441, g = 0.196; figure 2B).

<u>Fatigue increased transfer of learned sequence to the untrained hand</u>

Next, we went on to address the hypothesis that right-hand fatigue would lead to CE, i.e., an

increased rate of skill acquisition in the right, untrained hand after left-hand SRTT motor

training.

As expected, participants were able to learn the SRTT with their left-hand, as shown by a

decrease in RTs over time [RM-ANOVA with one factor of condition (5% MVC, 50% MVC) and

one factor of block (1-15); significant main effect of block (F(1.9,26.5) = 12.035, p < 0.001, η_p^2 =

0.462); figure 3A]. However, there were no differences in learning between the two conditions

[no significant main effect of condition (F(1,14) = 0.595, p = 0.453, η_p^2 = 0.041), no condition ×

block interaction (F(4.2,59.1) = 0.836, p = 0.513, η_p^2 = 0.056)]. At post-learning and retention

separate condition × block type (random, sequence) RM-ANOVA tests were run for the left-

hand. These analyses revelated significant main effects of block type at post-learning (F(1,14) =

14.556, p = 0.002, η_p^2 = 0.510) and retention (F(1,14) = 12.739, p = 0.003, η_p^2 = 0.476). However,

the condition × block type interactions at post-learning (F(1,14) = 0.815, p = 0.382, η_p^2 = 0.055)

and retention (F(1,14) = 0.909, p = 0.357, η_p^2 = 0.061), in addition to the main effect of

condition at Post-Learning (F(1,14) = 2.892, p = 0.111, η_p^2 = 0.171) and Retention (F(1,14) =

0.013, p = 0.909, η_p^2 = 0.001) all were not significant. These data suggest that left-hand

sequence learning occurred for both conditions, but no differences were observed between

conditions (figure 3B).

Next, we asked whether the right-hand fatigue affected the ability to transfer the learned

sequence from the trained left-hand to the untrained right-hand. We performed additional

condition × block type RM-ANOVA tests at post-learning and retention for the untrained right-

hand. As would be expected, participants were faster when performing the learned sequence

than the random sequence [significant main effect of block type, (F(1,14) = 8.155, p = 0.013, η_p^2

= 0.368)].   However, there was a significant difference between the 50% and 5% MVC

conditions in the degree of sequence transfer [significant condition × block type interaction

($F(1,14)$ = 6.325, p = 0.025, η_p^2 = 0.311)], which was driven by faster response times in

sequence, but not random, blocks in the 50% condition (50% MVC: sequence vs random blocks:

$t(14)$ = 2.881, p = 0.012, g = 0.719).   However, by the retention timepoint there was no

difference between the conditions (significant main effect of block type ($F(1,14)$ = 8.419, p =

0.012, η_p^2 = 0.376), no significant main effect of condition ($F(1,14)$ = 0.377, p = 0.549, η_p^2 =

0.026) and no significant condition × block type interaction ($F(1,14)$ = 0.797, p = 0.387, η_p^2 =

0.054; figure 3B).

Fatigue increased functional connectivity between the ipsilateral rM1 and right orbitofrontal cortex

328 To investigate the neural changes associated with our observed fatigue-induced behavioural

329 improvements in the unfatigued hand, we performed a voxel-wise seed-based analysis from the

330 hand area of ipsilateral, rM1.   There was a significant increase in functional connectivity with

331 the ipsilateral, right orbitofrontal cortex (rOFC) in the 50% MVC condition (table 1; figure 4A).

SMA-SMA connectivity increased after fatiguing contractions

To address our *a priori* hypothesis that the 50% MVC condition would alter functional

connectivity between M1 and SMA, we performed an ROI-ROI connectivity analysis between

left and right M1 and SMA.   A RM-ANOVA with one factor of condition (5% MVC, 50% MVC),

one factor of ROI (M1, SMA), and one factor of time (Pre-Task, Post-Task) revealed a significant

three-way (condition × ROI × time) interaction [($F(1,14)$ = 10.154, p = 0.007, η_p^2 = 0.420; a

significant main effect of ROI ($F(1,14)$ = 118.113, p < 0.001, η_p^2 = 0.894), but no significant main

effect of condition ($F(1,14)$ = 0.508, p = 0.488, η_p^2 = 0.8035), or time ($F(1,14)$ = 0.797, p = 0.387,

η_p^2 = 0.054)].

To understand this three-way interaction, we then ran separate RM-ANOVA tests for the 50%

and 5% MVC conditions, which revealed a significant ROI × time interaction for the 50% MVC

condition ($F(1,14)$ = 11.970, p = 0.004, η_p^2 = 0.461) but not for the 5% MVC condition ($F(1,14)$ <

0.001, p = 0.997, η_p^2 < 0.001).   Follow-up tests revealed that this interaction was driven by a

significant change in SMA-SMA connectivity in the 50% MVC condition ($t(14)$ = -2.203, p =

0.045, g = -0.550), figure 4B).

Given the 50% MVC force-matching task led to both fatigue and an increase in SMA-SMA

connectivity, we next wanted to investigate whether these two effects were related.   We

demonstrated a negative correlation between the degree of fatigue in the 50% MVC condition

and change in SMA-SMA connectivity, such that greater fatigue, indexed by greater decrease in

the force output, correlated with an increase in SMA-SMA connectivity ($r^2_{adjusted}$ = 0.208, p =

0.050, β = -0.1572; figure 4C). This correlation was both anatomically and behaviourally specific,

such that this relationship was not observed for M1-M1 connectivity (50% MVC: M1-M1

connectivity and fatigue $r^2_{adjusted}$ = -0.077, p = 0.945, β = -0.076) or in SMA-SMA connectivity in

the 5% MVC condition (5% MVC: SMA-SMA connectivity and fatigue: $r^2_{adjusted}$ = -0.024, p =

0.427, β = -0.863).   Fishers r-to-z analysis determined that the correlations between M1-M1 (z =

-0.45, p = 0.65) connectivity and SMA-SMA (z = 0.77, p = 0.44) connectivity with fatigue did not

differ between conditions.

Decrease in SMA and M1 GABA correlated with decrease in left-hand RTs

after fatigue

Finally, given our *a priori* hypothesis that the behavioural effects of fatigue are driving by

changes in inhibition, we investigated changes in GABA within each ROI.   We ran RM-ANOVA

tests for M1 and SMA separately, with one factor of condition (50% MVC, 5% MVC), one factor

of hemisphere (left, right) and one factor of time (pre, post). There were no significant main

effects or interactions with condition.   However, in line with our *a priori* hypothesis that GABA

would decrease and Glx would increase on a participant-by-participant basis for the 50% MVC

condition only, there were significant positive relationships between a decrease in lM1 GABA

(one-tailed: $r^2_{adjusted}$ = 0.196, p = 0.028, β = 4.79), lSMA GABA (one-tailed: $r^2_{adjusted}$ = 0.143, p =

0.045, β = 3.68), rSMA GABA (one-tailed: $r^2_{adjusted}$ = 0.172, p = 0.035, β =6.11) and the decrease

in RTs for the unfatigued left-hand. A significant negative relationship was also observed for

the increase in rM1 Glx with the decrease in RTs for the unfatigued left-hand (one-tailed:

$r^2_{adjusted}$ = 0.167, p = 0.036, β = -2.69). These findings suggest that bilateral M1 and SMA

alterations in inhibition and excitation predict the degree of RT improvement in the unfatigued

hand (figure 5).

Discussion

The objective of this study was to determine the neural correlates of unimanual fatigue and the

impact of fatigue on CE. Specifically, we hypothesised that fatigue would induce increased

sensorimotor functional connectivity (Jiang *et al.*, 2012) and disinhibition of ipsilateral motor

areas (Maruyama *et al.*, 2006, 2012; Takahashi *et al.*, 2009), neural effects that lead to

enhanced performance of the contralateral, unfatigued, hand.

We showed that right handgrip contractions at 50% MVC, but not 5% MVC, resulted in fatigue.

In line with the expected CE effects, this unilateral fatigue had behavioural effects: we observed

decreased RTs in the left, unfatigued hand. These behavioural changes were accompanied by

neural changes. We observed increased SMA-SMA connectivity, which correlated with the

degree of fatigue on a participant-by-participant basis. Additionally, increased rM1-rOFC

connectivity was observed on a voxel-wise analysis, which correlated with the behavioural

change in right hand RTs after fatigue. Finally, we used MRSI to quantify potential changes in

GABA and Glx. We observed no group mean changes, but on a participant-by-participant basis,

GABA decreases in the lM1 and bilateral SMAs correlated with behavioural improvements.

Fatigue increases cross-education

We demonstrated, in line with our hypothesis, that fatiguing contractions performed with the

right-hand led to significant behavioural improvements in the left, unfatigued hand. We

observed a significant improvement in left hand RTs, something that was not seen either in the

fatigued hand, nor with either hand in the non-fatiguing 5% MVC condition.   As would be

expected, participants learned the SRTT sequence, as evidenced by a decrease in RTs in

sequence blocks in the trained, left-hand.   Although against our hypothesis, there was no

difference in learning rates between the two conditions, we did observe a fatigue-induced

increase in transfer of the learned sequence to the right-hand after left-hand SRTT motor

learning, that was not present in the random button presses. There were no additional benefits

to the transfer of motor performance (i.e., random block RTs) after fatiguing contractions, yet,

both conditions did experience significant CE effects with the untrained right-hand improving

RTs in sequence and random blocks.

One plausible explanation for this behavioural effect of transfer of sequence learning is that the

increased interhemispheric SMA-SMA connectivity observed after fatigue may facilitate

transfer of the motor sequence learned with the left-hand to the right.   This-fatigue induced

increase in SMA-SMA connectivity may putatively impact neural circuits specifically

underpinning sequence learning (Perez *et al.*, 2007), rather than simply motor performance, as

no corresponding fatigue-induced improvement in right RTs is observed for the random blocks.

Indeed, SMA is known to be involved in consolidation of a learned sequence (Dahms *et al.*,

2020), plays a major role in the planning and self-initiation of voluntary movement, and serves

to integrate multimodal information to ensure motor and visual systems are in agreement

(Hikosaka *et al.*, 1999; Nakahara *et al.*, 2001). The increased connectivity between left and right

SMA in the present study may reflect a mechanism by which fatigue-induced neural changes

improve sequence memory for the untrained hand.

At retention there was no difference between the 5% MVC and 50% MVC conditions.   This

perhaps suggests that fatigue speeds learning in the untrained hand rather than increasing the

end magnitude of learning.   However, despite not necessarily increasing the total amount of

skill acquisition, the finding that fatigue speeds acquisition has implications for motor

rehabilitation, where CE is seemingly enhanced in clinical situations (Dragert & Zehr, 2013;

Urbin *et al.*, 2015; Sun *et al.*, 2018), and could therefore be putatively used to speed gains from

rehabilitation, something that is of major importance in recovery (Stinear & Byblow, 2014).

Increased functional connectivity between ipsilateral rM1 and rOFC after fatigue

We observed increased functional connectivity between our rM1 seed and the rOFC after

fatigue. Increased OFC activity has been associated with faster response times (Bode *et al.*,

2018) and hand motor learning (Alves Heinze *et al.*, 2019). The increased rM1-rOFC functional

connectivity observed here may therefore reflect the OFC's role as a top-down motor control

region (Ono *et al.*, 2014) which is involved in the regulation of motor responses and error

monitoring (Alves Heinze *et al.*, 2019). In line with this hypothesis, Jackson et al. (2003) showed

increased activation in the right OFC in participants who significantly improved their

performance in a right-hand motor task, either having physically executed the task or having

performed motor imagery. Taken together, our results therefore may suggest that increased

functional connectivity between the rM1 and the rOFC is functionally relevant for improved RTs

in the contralateral, unfatigued left-hand.

Fatigue increased SMA-SMA connectivity

Our ROI-based functional connectivity analysis examined how the temporal correlations

between left and right M1 and SMA homologs changed after the right handgrip force-matching

task. After fatigue, SMA-SMA functional connectivity increased, and the degree of change was

negatively correlated with the degree of fatigue, such that participants who exhibited greater

fatigue showed greater increase in SMA-SMA functional connectivity.

The physiological basis of the role of the SMA in CE is not yet clear, but SMA has been described

as a 'phylogenetically older' M1 (Goldberg, 1985) which has direct excitatory connections not

only onto the M1 circuitry (Shirota *et al.*, 2012; Arai *et al.*, 2012), but onto alpha-motoneurons

that innervate the hand and fingers (Maier *et al.*, 2002; Dum & Strick, 2005). The SMA is also an

important region for consolidation and retention of motor sequence learning (Dahms *et al.*,

2020). Therefore, it is plausible that the fatigue-induced increase in SMA-SMA functional

connectivity improved CE sequence learning either by influencing rM1 through rSMA-rM1

excitatory connections or through direct corticospinal connections to upper limb alpha-

motoneurons (Dum & Strick, 2005). Transcallosal homologous connectivity between motor

regions primarily reflects inhibitory processes (Wahl *et al.*, 2007). Therefore, an increase in

SMA-SMA functional connectivity may reflect an increase in interhemispheric inhibition

between these premotor regions. Previous research has identified the SMA as a critical region

for CE (Perez *et al.*, 2007; Ruddy *et al.*, 2017): inhibiting SMA with repetitive TMS abolished CE

without impacting SRTT learning with the trained hand. Further, SMA-SMA interhemispheric

structural connectivity negatively correlates with the magnitude of CE (Ruddy *et al.*, 2017), such

that individuals with greater structural SMA-SMA connectivity exhibit a lower degree of skill

transfer. Given this, it is possible that an increase in interhemispheric SMA-SMA functional

connectivity, reflecting interhemispheric inhibition, may aid in silencing unwanted cortical

activity that facilitates a more focal excitatory process in the opposite hemisphere (Carson,

2020) giving rise to an enhanced CE effect. These interhemispheric network-level effects from

the SMA may also be important in the faster response times observed with the left, unfatigued

hand following 50% MVC right-hand contractions.

A decrease in lM1 and bilateral SMA GABA after fatigue correlates with

improvements in left-hand RTs

To address our hypothesis that fatigue would lead to decreased inhibition in motor regions, we

used a novel MRSI sequence to quantify neurochemical concentrations across the sensorimotor

network. At a group level, there were no significant GABA changes in M1 and SMA in either

hemisphere after fatigue compared with 5% MVC control condition. However, in line with our

a priori hypothesis, we did observe a significant correlation on a participant-by-participant basis

between the decrease in lM1, lSMA, and rSMA GABA after fatigue with the improved behaviour

in the unfatigued hand. While the mechanism by which lM1 and bilateral SMA GABA decreases

lead to behavioural improvement in the ipsilateral hand is unclear, a combination of decussated
and non-decussated pathways may play an important role (Maruyama *et al.*, 2006; Aboodarda
et al., 2016).

Conclusions

This study identified that fatiguing right handgrip contractions led to behavioural improvements
in both hands which were accompanied by increased SMA-SMA functional connectivity. In
addition, we demonstrated an increase in functional connectivity between rM1 and the rOFC.
Speaking to the neurochemical mechanisms underpinning these behavioural improvements in
the unfatigued hand, we also demonstrated positive relationships between change in inhibition
(GABA) in lM1 and bilateral SMA and a negative relationship between the change in rM1 Glx
with the behavioural improvements in the unfatigued left-hand. Taken together, these results
suggest that unimanual fatiguing exercise may be a potentially promising adjunct therapy for
rehabilitation in a range of neurological or orthopedic conditions.

Author contributions

492 J.W.A., J.M.L, J.P.F., & C.J.S. conceived and designed the experiment. C.Z. & J.M.L developed the
493 software to run the behavioural experiment. J.W.A., & J.M.L. collected the data and J.W.A.,
494 E.C.M., J.M.L., J.P.F., & C.J.S. analyzed the data. All authors were involved in data interpretation.
495 J.W.A. drafted the paper, and all authors revised it and approve the final version.

Conflict of interest

497 None to report

Funding sources

C.J.S. holds a Sir Henry Dale Fellowship, funded by the Wellcome Trust and the Royal Society
(102584/Z/13/Z). This work was also supported by the National Institute for Health Research
(NIHR) Oxford Health Biomedical Research Centre, and the John Fell Fund. The Wellcome
Centre for Integrative Neuroimaging is supported by core funding from the Wellcome Trust

(203139/Z/16/Z). J.P.F. holds a Natural Sciences and Engineering Research Council of Canada

(NSERC) Discovery Grant: RGPIN 2016-0529 at the University of Saskatchewan. J.M.L was

supported by a National Institutes of Health Oxford-Cambridge Scholar Fellowship, and the

International Biomedical Research Alliance. J.W.A. received an NSERC Alexander Graham Bell

Canada Graduate Scholarship – Doctoral (CGS D) Award and a MITACS Globalink Research

Award to complete this work.

Acknowledgements

The authors would like to acknowledge Dr. Jason DeFreitas for his valuable insight on the
manuscript.

512 References

Aboodarda SJ, Šambaher N & Behm DG (2016). Unilateral elbow flexion fatigue modulates

corticospinal responsiveness in non-fatigued contralateral biceps brachii. *Scand J Med Sci*

Sport **26,** 1301–1312.

Allman C, Amadi U, Winkler AM, Wilkins L, Filippini N, Kischka U, Stagg CJ & Johansen-Berg H

(2016). Ipsilesional anodal tDCS enhances the functional benefits of rehabilitation in

patients after stroke. *Sci Transl Med* **8,** 330re1-330re1.

Alves Heinze R, Vanzella P, Zimeo Morais GA & Sato JR (2019). Hand motor learning in a musical

context and prefrontal cortex hemodynamic response: a functional near-infrared

spectroscopy (fNIRS) study. *Cogn Process*; DOI: 10.1007/s10339-019-00925-y.

Andersson JLR, Jenkinson M & Smith S (2007*a*). *Non-linear registration aka spatial*

normalisation FMRIB Technial Report TR07JA2.

Andersson JLR, Jenkinson M & Smith S (2007*b*). *Non-linear optimisation FMRIB Technial Report*

TR07JA1.

Andrushko JW, Gould LA, Renshaw DW, Ekstrand C, Hortobágyi T, Borowsky R & Farthing JP

(2021). High force unimanual handgrip contractions increase ipsilateral sensorimotor

activation and functional connectivity. *Neuroscience* **452,** 111–125.

Andrushko JW, Lanovaz JL, Björkman KM, Kontulainen SA & Farthing JP (2018). Unilateral

strength training leads to muscle-specific sparing effects during opposite homologous limb

immobilization. *J Appl Physiol* **124,** 866–876.

Arai N, Lu MK, Ugawa Y & Ziemann U (2012). Effective connectivity between human

supplementary motor area and primary motor cortex: A paired-coil TMS study. *Exp Brain*

Res **220,** 79–87.

Bachtiar V, Johnstone A, Berrington A, Lemke C, Johansen-Berg H, Emir U & Stagg CJ (2018).

Modulating regional motor cortical excitability with noninvasive brain stimulation results

in neurochemical changes in bilateral motor cortices. *J Neurosci* **38,** 7327–7336.

Bachtiar V, Near J, Johansen-Berg H & Stagg CJ (2015). Modulation of GABA and resting state

functional connectivity by transcranial direct current stimulation. *Elife* **4,** e08789.

Bode S, Bennett D, Sewell DK, Paton B, Egan GF, Smith PL & Murawski C (2018). Dissociating

neural variability related to stimulus quality and response times in perceptual decision-

making. *Neuropsychologia* **111,** 190–200.

Cabanes E, Confort-Gouny S, Le Fur Y, Simond G & Cozzone PJ (2001). Optimization of residual

water signal removal by HLSVD on simulated short echo time proton MR spectra of the

human brain. *J Magn Reson* **150,** 116–125.

Carr JC, Bemben M, Stock MS & DeFreitas JM (2021). Ipsilateral and contralateral responses

following unimanual fatigue with and without illusionary mirror visual feedback. *J*

Neurophysiol jn.00077.2021.

Carson RG (2020). Inter-hemispheric inhibition sculpts the output of neural circuits by co-opting

the two cerebral hemispheres. *J Physiol* JP279793.

Cox RW (1996). AFNI: software for analysis and visualization of functional magnetic resonance

neuroimages. *Comput Biomed Res* **29,** 162–173.

Cox RW & Glen DR (2013). Nonlinear warping in AFNI. *Present 19th Annu Meet Organ Hum*

Brain Mapp. Available at:

https://afni.nimh.nih.gov/pub/dist/HBM2013/Cox_Poster_HBM2013.pdf.

Dahms C, Brodoehl S, Witte OW & Klingner CM (2020). The importance of different learning

stages for motor sequence learning after stroke. *Hum Brain Mapp* **41,** 270–286.

Dehno SN, Kamali F, Shariat A & Jaberzadeh S (2021). Unilateral strength training of the less

affected hand improves cortical excitability and clinical outcomes in patients with subacute

stroke: A randomized controlled trial. *Arch Phys Med Rehabil*; DOI:
10.1016/j.apmr.2020.12.012.

Dragert K & Zehr EP (2013). High-intensity unilateral dorsiflexor resistance training results in
bilateral neuromuscular plasticity after stroke. *Exp Brain Res* **225,** 93–104.

Dum RP & Strick PL (2005). Frontal lobe inputs to the digit representations of the motor areas
on the lateral surface of the hemisphere. *J Neurosci* **25,** 1375–1386.

Emir UE, Burns B, Chiew M, Jezzard P & Thomas MA (2017). Non-water-suppressed
short-echo-time magnetic resonance spectroscopic imaging using a concentric ring k-space
trajectory. *Nmr Biomed*; DOI: 10.1002/NBM.3714.

Farthing JP, Krentz JR & Magnus CR (2009). Strength training the free limb attenuates strength
loss during unilateral immobilization. *J Appl Physiol* **106,** 830–836.

Farthing JP, Krentz JR, Magnus CR, Barss TS, Lanovaz JL, Cummine J, Esopenko C, Sarty GE &
Borowsky R (2011). Changes in functional magnetic resonance imaging cortical activation
with cross education to an immobilized limb. *Med Sci Sports Exerc* **43,** 1394–1405.

Floyer-Lea A, Wylezinska M, Kincses T & Matthews PM (2006). Rapid modulation of GABA
concentration in human sensorimotor cortex during motor learning. *J Neurophysiol* **95,**
1639–1644.

Goldberg G (1985). Supplementary motor area structure and function: Review and hypotheses.
Behav Brain Sci **8,** 567–588.

Griffanti L, Salimi-Khorshidi G, Beckmann CF, Auerbach EJ, Douaud G, Sexton CE, Zsoldos E,
Ebmeier KP, Filippini N, Mackay CE, Moeller S, Xu J, Yacoub E, Baselli G, Ugurbil K, Miller KL
& Smith SM (2014). ICA-based artefact removal and accelerated fMRI acquisition for
improved resting state network imaging. *Neuroimage*; DOI:
10.1016/j.neuroimage.2014.03.034.

Hikosaka O, Sakai K, Lu X, Nakahara H, Rand MK, Nakamura K, Miyachi S & Doya K (1999).
Parallel neural networks for learning sequential procedures. *Trends Neurosci* **22,** 464–471.
Available at: http://www.cell.com/article/S0166223699014393/fulltext [Accessed April 15,
2021].

Jackson PL, Lafleur MF, Malouin F, Richards CL & Doyon J (2003). Functional cerebral

589 reorganization following motor sequence learning through mental practice with motor
590 imagery. *Neuroimage*; DOI: 10.1016/S1053-8119(03)00369-0.

591 Jenkinson M, Bannister P, Brady M & Smith S (2002). Improved optimization for the robust and
592 accurate linear registration and motion correction of brain images. *Neuroimage* **17,** 825–
593 841.

594 Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW & Smith SM (2012). FSL. *Neuroimage*
595 **62,** 782–790.

596 Jenkinson M & Smith S (2001). A global optimisation method for robust affine registration of
597 brain images. *Med Image Anal* **5,** 143–156.

598 Jiang Z, Wang X-F, Kisiel-Sajewicz K, Yan JH & Yue GH (2012). Strengthened functional
599 connectivity in the brain during muscle fatigue. *Neuroimage* **60,** 728–737.

600 Johansen-Berg H, Behrens TEJ, Robson MD, Drobnjak I, Rushworth MFS, Brady JM, Smith SM,
601 Higham DJ & Matthews PM (2004). Changes in connectivity profiles define functionally
602 distinct regions in human medial frontal cortex. *Proc Natl Acad Sci U S A* **101,** 13335–
603 13340.

604 Kim D-Y, Lim J-Y, Kang EK, You DS, Oh M-K, Oh B-M & Paik N-J (2010). Effect of transcranial
605 direct current stimulation on motor recovery in patients with subacute stroke. *Am J Phys*
606 *Med Rehabil* **89,** 879–886.

607 Kleiner M, Brainard DH, Pelli DG, Broussard C, Wolf T & Niehorster D (2007). What's new in
608 Psychtoolbox-3? *Perception*.

609 Klose U (1990). In vivo proton spectroscopy in presence of eddy currents. *Magn Reson Med* **14,**
610 26–30.

611 Kolasinski J, Hinson EL, Divanbeighi Zand AP, Rizov A, Emir UE & Stagg CJ (2019). The dynamics
612 of cortical GABA in human motor learning. *J Physiol*; DOI: 10.1113/JP276626.

613 Magnus CR, Arnold CM, Johnston G, Dal-Bello Haas V, Basran J, Krentz JR & Farthing JP (2013).
614 Cross-education for improving strength and mobility after distal radius fractures: A
615 randomized controlled trial. *Arch Phys Med Rehabil* **94,** 1247–1255.

616 Magnus CR, Barss TS, Lanovaz JL & Farthing JP (2010). Effects of cross-education on the muscle
617 after a period of unilateral limb immobilization using a shoulder sling and swathe. *J Appl*

Physiol **109,** 1887–1894.

Maier MA, Armand J, Kirkwood PA, Yang H-W, Davis JN & Lemon R.N. (2002). Differences in the
corticospinal projection from primary motor cortex and supplementary motor area to
macaque upper limb motoneurons: An anatomical and electrophysiological study. *Cereb*
Cortex **12,** 281–296.

Manca A, Hortobágyi T, Carroll TJ, Enoka RM, Farthing JP, Gandevia SC, Kidgell DJ, Taylor JL &
Deriu F (2021). Contralateral effects of unilateral strength and skill training: Modified
delphi consensus to establish key aspects of cross-education. *Sport Med* **51,** 11–20.

Manca A, Hortobágyi T, Rothwell J & Deriu F (2018). Neurophysiological adaptations in the
untrained side in conjunction with cross-education of muscle strength: a systematic review
and meta-analysis. *J Appl Physiol* **124,** 1502–1518.

Maruyama A, Matsunaga K, Tanaka N & Rothwell JC (2006). Muscle fatigue decreases short-
interval intracortical inhibition after exhaustive intermittent tasks. *Clin Neurophysiol* **117,**
864–870.

Maruyama A, Nuruki A, Etoh S & Rothwell JC (2012). Could fatigue be used as a paradoxical
intervention to improve motor learning after stroke? In *International Conference on*
Complex Medical Engineering, pp. 298–301. IEEE. Available at:
http://ieeexplore.ieee.org/document/6275681/.

Miller KL et al. (2016). Multimodal population brain imaging in the UK Biobank prospective
epidemiological study. *Nat Neurosci 2016 1911* **19,** 1523–1536.

Nakahara H, Doya K & Hikosaka O (2001). Parallel cortico-basal ganglia mechanisms for
acquisition and execution of visuomotor sequences - A computational approach. *J Cogn*
Neurosci **13,** 626–647.

Ni Z, Gunraj C & Chen R (2007). Short interval intracortical inhibition and facilitation during the
silent period in human. *J Physiol* **583,** 971–982.

Ono Y, Nomoto Y, Tanaka S, Sato K, Shimada S, Tachibana A, Bronner S & Noah JA (2014).
Frontotemporal oxyhemoglobin dynamics predict performance accuracy of dance
simulation gameplay: Temporal characteristics of top-down and bottom-up cortical
activities. *Neuroimage* **85,** 461–470.

Pearce AJ, Hendy AM, Bowen WA & Kidgell DJ (2013). Corticospinal adaptations and strength
maintenance in the immobilized arm following 3 weeks unilateral strength training. *Scand*
J Med Sci Sport **23,** 740–748.

Peirce J, Gray JR, Simpson S, MacAskill M, Höchenberger R, Sogo H, Kastman E & Lindeløv JK
(2019). PsychoPy2: Experiments in behavior made easy. *Behav Res Methods* **51,** 195–203.

Pelet DCS & Orsatti FL (2021). Effects of resistance training at different intensities of load on
cross-education of muscle strength. *Appl Physiol Nutr Metab*apnm-2021-0088.

Perez MA, Tanaka S, Wise SP, Sadato N, Tanabe HC, Willingham DT & Cohen LG (2007). Neural
substrates of intermanual transfer of a newly acquired motor skill. *Curr Biol* **17,** 1896–
1902.

Provencher SW (2001). Automatic quantitation of localized in vivo 1H spectra with LCModel.
NMR Biomed **14,** 260–264.

Provencher SW (2016). LCModel and LCMgui User's Manual. Available at: http://s-
provencher.com/lcm-manual.shtml.

Reis J, Schambra HM, Cohen LG, Buch ER, Fritsch B, Zarahn E, Celnik PA & Krakauer JW (2009).
Noninvasive cortical stimulation enhances motor skill acquisition over multiple days
through an effect on consolidation. *Proc Natl Acad Sci U S A* **106,** 1590–1595.

Ruddy KL & Carson RG (2013). Neural pathways mediating cross education of motor function.
Front Hum Neurosci **7,** 1–22.

Ruddy KL, Leemans A, Woolley DG, Wenderoth N & Carson RG (2017). Structural and functional
cortical connectivity mediating cross education of motor function. *J Neurosci* **37,** 2555–
2564.

Saad ZS, Glen DR, Chen G, Beauchamp MS, Desai R & Cox RW (2009). A New Method for
Improving Functional-to-Structural MRI Alignment using Local Pearson Correlation.
Neuroimage **44,** 839.

Salimi-Khorshidi G, Douaud G, Beckmann CF, Glasser MF, Griffanti L & Smith SM (2014).
Automatic denoising of functional MRI data: Combining independent component analysis
and hierarchical fusion of classifiers. *Neuroimage* **90,** 449–468.

Shirota Y, Hamada M, Terao Y, Ohminami S, Tsutsumi R, Ugawa Y & Hanajima R (2012).

Increased primary motor cortical excitability by a single-pulse transcranial magnetic

stimulation over the supplementary motor area. *Exp Brain Res* **219,** 339–349.

Smith SM (2002). Fast robust automated brain extraction. *Hum Brain Mapp* **17,** 143–155.

Stagg CJ, Bachtiar V, Amadi U, Gudberg CA, Ilie AS, Sampaio-Baptista C, O'Shea J, Woolrich M,

Smith SM, Filippini N, Near J & Johansen-Berg H (2014). Local GABA concentration is

related to network-level resting functional connectivity. *Elife* **3,** 1–9.

Stagg CJ, Bachtiar V & Johansen-Berg H (2011*a*). The role of GABA in human motor learning.

Curr Biol **21,** 480–484.

Stagg CJ, Jayaram G, Pastor D, Kincses ZT, Matthews PM & Johansen-Berg H (2011*b*). Polarity

and timing-dependent effects of transcranial direct current stimulation in explicit motor

learning. *Neuropsychologia* **49,** 800–804.

Steel A, Chiew M, Jezzard P, Voets NL, Plaha P, Thomas MA, Stagg CJ & Emir UE (2018).

Metabolite-cycled density-weighted concentric rings k-space trajectory (DW-CRT) enables

high-resolution 1 H magnetic resonance spectroscopic imaging at 3-Tesla. *Sci Rep* **8,** 7792.

Stinear CM & Byblow WD (2014). Predicting and accelerating motor recovery after stroke. *Curr*

Opin Neurol **27,** 624–630.

Sun Y, Ledwell NMH, Boyd LA & Zehr EP (2018). Unilateral wrist extension training after stroke

improves strength and neural plasticity in both arms. *Exp Brain Res* **236,** 2009–2021.

Takahashi K, Maruyama A, Maeda M, Etoh S, Hirakoba K, Kawahira K & Rothwell JC (2009).

Unilateral grip fatigue reduces short interval intracortical inhibition in ipsilateral primary

motor cortex. *Clin Neurophysiol* **120,** 198–203.

The Jamovi Project (2020). Jamovi (Version 1.2). Available at: https://www.jamovi.org.

The Mathworks Inc (2018). MATLAB 2018a. *WwwMathworksCom/Products/Matlab*.

Urbin MA, Harris-Love ML, Carter AR & Lang CE (2015). High-intensity, unilateral resistance

training of a non-paretic muscle group increases active range of motion in a severely

paretic upper extremity muscle group after stroke. *Front Neurol* **6,** 119.

Valdes O, Ramirez C, Perez F, Garcia-Vicencio S, Nosaka K & Penailillo L (2021). Contralateral

effects of eccentric resistance training on immobilized arm. *Scand J Med Sci Sports* **31,** 76–

90.

Wahl M, Lauterbach-Soon B, Hattingen E, Jung P, Singer O, Volz S, Klein JC, Steinmetz H &
Ziemann U (2007). Human motor corpus callosum: Topography, somatotopy, and link
between microstructure and function. *J Neurosci* **27,** 12132–12138.

Walsh DO, Gmitro AF & Marcellin MW (2000). Adaptive reconstruction of phased array MR
imagery. *Magn Reson Med*; DOI: 10.1002/(SICI)1522-2594(200005)43:5.

Weinrich CA, Brittain JS, Nowak M, Salimi-Khorshidi R, Brown P & Stagg CJ (2017). Modulation
of long-range connectivity patterns via frequency-specific stimulation of human cortex.
Curr Biol **27,** 3061-3068.e3.

Woolrich MW, Behrens TEJ, Beckmann CF, Jenkinson M & Smith SM (2004). Multilevel linear
modelling for FMRI group analysis using Bayesian inference. *Neuroimage* **21,** 1732–1747.

Woolrich MW, Ripley BD, Brady M & Smith SM (2001). Temporal autocorrelation in univariate
linear modeling of FMRI data. *Neuroimage* **14,** 1370–1386.

Figure legends

Figure 1. A) Theoretical physiological model for the present study, **B)** Schematic outlining the

study design. Both aspects of this figure were created with BioRender.com.

Figure 2. Behavioural changes. **A)** Beta values for the area under the curve of the force profile

from the nine-minute right handgrip force-matching task in experiment one. * = 50% MVC

experienced a significant decline in motor performance (one-sample *t*-test, p = 0.005), ** =

significant difference between conditions (paired-samples *t*-test, p = 0.006), and **B)** Percent

change in response times from Pre-MRI to Post-MRI for the left (purple data points) and right

(green data points) hands after the right-hand force-matching task at 5% MVC (circles) and 50%

MVC (triangles). † = significant condition × hand interaction (p = 0.045),** = significant paired

sample *t*-test (p = 0.037),* = significant one-sample *t*-test change in response times (p = 0.009).

Schematic of study design was created with BioRender.com.

Figure 3. A) SRTT motor learning blocks across time with the trained left-hand. R = random

trials, S = sequence trials, error bars = 95% confidence intervals, and **B)** Difference in response

times between random blocks and the sequence block for the left (left side; purple data points)

and right hands (right side; green data points) at Post-Learning (top row) and Retention

(bottom row) for the 5% MVC (circles) and 50% MVC (triangles) conditions. * = significant

difference between sequence and random blocks (all p < 0.05), † = significant interaction

between conditions at Post Learning for the right hand (p = 0.025). Schematic of study design

was created with BioRender.com.

Figure 4. Functional connectivity analyses for **A)** Seed-based functional connectivity analysis

between the right ipsilateral primary motor cortex (seed) and the rest of the brain in the 50%

MVC condition, data is a Z-statistic threshold map in standard space. Z-threshold = 3.1. Figure is

in radiological view (left on right, right on left). **B)** Change in interhemispheric functional

connectivity between contralateral and ipsilateral primary motor cortices (M1-M1; red data

points), and the contralateral and ipsilateral supplementary motor areas (SMA-SMA; orange

data points) for 5% MVC (circles) and 50% MVC (triangles). Back horizonal bar represents the

mean change. Grey shaded area represents the standard deviation. † = significant condition ×

ROI × time interaction (p = 0.007), ** = significant ROI × time interaction for 50% MVC condition

(p = 0.004) * = significant paired-sample t-test against (SMA-SMA; p = 0.045), and **C)** Scatter plot

correlation between X-axis: The change in interhemispheric functional connectivity z-scores

between the contralateral left and ipsilateral right supplementary motor areas (SMA-SMA) and

Y-axis: beta values representing the decline in motor performance over time (negative slope =

performance decline over time) ($r^2_{adjusted}$ = 0.208, p = 0.050, β = -0.1572). Schematic of study

design was created with BioRender.com.

Figure 5. A) Magnetic resonance spectroscopic imaging spectrum from a representative

participant, with spectra from M1 voxels in each hemisphere. **B)** Scatter plot correlations for

the right hemisphere (purple) and left hemisphere (green) between X-axes: SMA (top scatter

plots), M1 (bottom scatter plots) for GABA/tCr (rSMA, lSMA, lM1) and Glx/tCr (rM1) ratios with

Y-axes: the change in left-hand response times (ms) after performing the right-hand target

matching task at either 5% MVC (circles) or 50% MVC (triangles) conditions. Schematic of study

design was created with BioRender.com.

Table 1. Post > Pre-force-matching task rs-fMRI contrasts from iM1 seed-based functional connectivity analysis

50% MVC						
			MNI Coordinates (mm)			
Voxels	*p*-value	z-max	X	Y	Z	Label
104	0.0348	5.12	34	26	-6	Right Orbitofrontal Cortex
5% MVC						
			MNI Coordinates (mm)			
Voxels	*p*-value	z-max	X	Y	Z	Label
182	0.00187	4.44	28	-50	-26	Right Cerebellum VI
104	0.0363	4.78	-42	-54	-4	Left Inferior Temporal Gyrus

A

(4) decreased [GABA] ↓

(2) decreased [GABA] ↓

(3) increased
interhemispheric
connectivity ↑

↑ (1) fatigue

(5) increased transfer

(5) improved behaviour
& learning ↑

Right-Hand

Left-Hand

B

Baseline   Post-MRI   ← Learning →   Post-learning   Retention

⚫⚫ Random   ⚫⚫ SRTT   ▢ Left-Hand   ▢ Right-Hand