TITLE

The challenges of independence: ontogeny of at-sea behaviour in a long-lived

seabird

KARINE DELORD[1]*, HENRI WEIMERSKIRCH[1], CHRISTOPHE BARBRAUD[1]

[1] Centre d'Etudes Biologiques de Chizé UMR 7372, CNRS, F-79360 Villiers en Bois, France

Abstract

The transition to independent foraging represents an important developmental stage in the life cycle of

most vertebrate animals. Juveniles differ from adults in various life history traits and tend to survive less

well than adults in most long-lived animals. Several hypotheses have been proposed to explain higher

mortality including that of inadequate/inferior foraging skills compared to adults, young naïve

individuals combining lack of experience and physical immaturity. Thus a change in behaviour, resulting

in an improvement of skills acquired from growing experience, is expected to occur during a period of

learning through the immaturity phase. Very few studies have investigated the ontogeny of foraging

behaviour over long periods of time, particularly in long-lived pelagic seabirds, due to the difficulty of

obtaining individual tracking data over several years. We investigated the foraging behaviour, through

activity patterns, during the three life stages of the endangered Amsterdam albatross by using

miniaturized activity loggers on naïve juveniles, immatures and adults. Naïve juveniles during their first

month at sea after leaving their colony exhibited lower foraging effort (greater proportion of time spent

sitting on water, longer and more numerous bouts on water, shorter and fewer flying bouts). Patterns of

activity parameters in juveniles after independence suggested a progressive change of foraging

performances during the first two months after fledging. We found sex differences in activity

parameters according to time since departure from the colony and month of the year, consistent with

the important sexual dimorphism in the Amsterdam albatross. Regardless of life stage considered,

activity parameters exhibited temporal variability reflecting the modulation of foraging behaviour. This

variability is discussed in light of both extrinsic (i.e. environmental conditions such as variability in food

resources or in wind) and intrinsic (i.e. energetic demands linked to plumage renew during moult)

factors.

Keywords: activity data loggers; foraging behaviour; southern Indian Ocean; Amsterdam albatross;

Diomedea amsterdamensis

Introduction

The transition from parental food dependency to independent foraging represents an important developmental stage in the life cycle of most vertebrate animals (Mushinsky et al. 1982; Margrath and Lill 1985; Martin and Bateson 1985; Marchetti and Price 1989; Langen 1996; Burns et al. 2004) and is increasingly documented in a wide range of taxa (reptiles, birds, and some mammals). A widely accepted hypothesis is inadequate/inferior foraging skills of juveniles compared to adults, young naïve individuals combining lack of experience and physical immaturity (Lack 1954; Daunt et al. 2007). Thus, a change in behaviour, resulting from an improvement of skills acquired from increasing experience is expected to occur during a period of learning through the immaturity phase. Learning often refers to stimulus-response associative learning ('trial and error'; Ruaux et al. 2020), although other forms of learning (such as social learning or imprinting) are also taken into account when considering the ontogeny of complex behaviours (Heyes 1994; Wynn et al. 2020). Such a learning process has been studied on various taxa from insects to primates (Bruner 1972; Caubet et al. 1992; Dukas 2006; Rapaport and Brown 2008).

Juvenile birds are known to undertake vagrant erratic journeys during the post-fledging period in passerines (Naef-Daenzer and Grüebler 2008; Becker 2014; Evans 2018; Boynton et al. 2020), in raptors (Urios et al. 2010; Krüger et al. 2014; Harel et al. 2016) and in seabirds (Riotte-Lambert and Weimerskirch 2013; Collet et al. 2020). Recent studies highlighted that the flight capacities and foraging behaviour of juveniles differed from those of adults in storks (Rotics et al. 2016), raptors (Harel et al. 2016; Nourani et al. 2020) or seabirds (Ydenberg 1989; Péron and Grémillet 2013; de Grissac et al. 2017; Corbeau et al. 2020). Most flight components were found to improve over time to tend towards those of adults (Riotte-Lambert and Weimerskirch 2013; de Grissac et al. 2017; Corbeau et al. 2020).

However, studies focusing on the foraging behaviour of juveniles remain scarce because of the difficulty to obtain individual tracking data for long periods, especially for long-lived pelagic seabirds with deferred maturity. Moreover, existing studies comparing flight capacities and foraging behaviour

between juveniles and adults in such species only collected data during the first few months that

juveniles spent at sea. Since juveniles may spend several years at sea before returning to a colony to

breed, our understanding of the ontogeny of flight capacities and foraging behaviour remains

fragmentary.

The Amsterdam albatross *Diomedea amsterdamensis* is a large and long-lived pelagic seabird with an

extended immaturity stage (~ 9 years, Rivalan et al. 2010). Similarly to a closely related species, the

wandering albatross *D. exulans*, their foraging strategy relies on very low flight costs as a result of their

dynamic soaring flight, whereby individuals optimize the orientation of their movement with wind

direction to maximize the daily distance covered (Pennycuick 1982). During initial post-fledging

movements juveniles wander alone over very long distances from their colony. At sea distribution during

every stage of the life-cycle of Amsterdam albatross was studied by Thiebot et al. (2014) and de Grissac

et al. (2016) who compared flight trajectories (i.e. departure direction or orientation toward specific

areas) of juveniles and adults. Both studies concluded on slight differences among stages in distribution

due to the extensive area they used. However, foraging behaviour is known to be constrained by

intrinsic factors such as sex, age, reproductive status and body size across a wide range of taxa and

hence play a key role in shaping activity (King 1974; Alerstam and Lindström 1990; Wearmouth and Sims

2008). To understand the changes in foraging proficiency according to experience (life-history stages),

longitudinal studies of individuals spanning critical periods of their lives are thus required. Advances in

animal-borne instrumentation enable key component of foraging behaviour such as foraging effort and

activity to be recorded over long periods.

In this study, we benefited from a unique dataset of different life stages (juveniles, immatures and

adults) and a remarkable duration (up to 28 months for juveniles) to characterise and compare the

changes in behaviour at sea when birds leave the colony (for several months: immatures and adults, or

years: juveniles before returning to land). We analyse the foraging behaviour, through activity patterns,

of naïve juveniles (first years of independence at sea), immatures (individuals that never bred, age 2-10

years) and adults (individuals that bred at least once, age 8-28 years) of Amsterdam albatross (Table 1).

By using miniaturized activity loggers (Global Location Sensing; GLS) to infer foraging behaviour (activity)

throughout the successive life stages we addressed the following questions: i) do individuals belonging

to different life-stages behave differently? ii) are there detectable progressive changes in activity

patterns? It is noteworthy that the loggers used do not yet allow to have longitudinal data (maximum 2-

3 years of recorded data) and to cover the entire period until an individual is recruited into the

population as a breeding adult, i.e. at least 8 years.

Previous knowledge of the ecology of large albatrosses and Amsterdam albatross described above

provides a practical framework for testing predictions about variability in foraging behaviour associated

with stage, time elapsed since departure from the colony, seasons and sex which are summarised in

Table 2. Given the overlap of spatial distribution between life-stages (not presented here but see

Thiebot et al. 2014; de Grissac et al. 2016; Pajot et al. 2021) we predicted that juveniles would

compensate for any lack of foraging proficiency by increasing foraging effort and time (i.e. lower time

spent on water and longer flying bouts, in other words decreasing time sitting on water and increasing

number and duration of flight bouts; Hypothesis (A), Table 2). We also predicted changes in activity of

juveniles early in post-fledging followed by more progressive changes. Based on results found on

wandering albatross fledglings (Riotte-Lambert and Weimerskirch 2013; Pajot et al. 2021) showing that

juveniles reached some adult foraging performances in less than two months, we predicted that

changes should be detected in activity parameters early after the juvenile left the colony (within few

first months). Overall, juveniles should show contrasted foraging effort (i.e. longer time spent on water,

shorter flying effort with fewer and shorter flying bouts) early in post-fledging compared to other life-

stages. Due to seasonal changes in food availability individuals will face at sea after leaving the colony

and the alleviation of energetic constraints linked to reproduction (for breeding adults) or to alternate

foraging trips at sea and period on land for pair bonding and mating display (for immature birds), we

predicted that adjustments of activity will occur according to the time spent (i.e. in months elapsed)

since the departure from the colony (Hypothesis (B), Table 2). In juveniles, we predicted early and rapid

changes during post-fledging and then more progressive changes. The two hypotheses (A & B) are not

mutually exclusive. While our main objective was to study post-fledging foraging behaviour activity as

described above, we also accounted for other sources of changes in foraging behaviour. These included

temporal (i.e. related to the month of the year) changes in activity parameters for all life-stages due (i)

to environmental changes occurring throughout the seasons, (ii) to partial moulting which is suspected

to occur outside the breeding period and to result in reduced activity for adults and immatures (i.e.

more time spent on the water; Weimerskirch et al. 2015, 2020), or (iii) to sex differences in flight

performances (Shaffer et al. 2001; Riotte-Lambert and Weimerskirch 2013; Clay et al. 2020) .

Materials and methods

Study species and data loggers

Amsterdam Island (37° 50' S; 77° 33' E) is located in the subtropical part of the southern Indian Ocean.

The Amsterdam albatross, like other great albatrosses, is a biennial breeder (Roux et al. 1983; Jouventin

et al. 1989), with high survival during juvenile, immature and adult phase (Rivalan et al. 2010). The

adults that raised a chick successfully do not start a new breeding cycle after chick fledging, but remain

at sea for a sabbatical period (~1 yr; Table 1; Rivalan et al. 2010). However, early failed breeders may

start to breed the following year (Rivalan et al. 2010). Immature birds may visit the colony when they

are 4–7 yrs old, but generally only start breeding at 9 yrs old (Table 1; Weimerskirch et al. 1997a).

Juvenile birds fledge and migrate independently from the adults in January (Table 1). Exact fledging

dates were not known for juveniles but were assessed from activity pattern as juvenile birds land on

water quickly after leaving the colony (Weimerskirch et al. 2006). Amsterdam albatrosses were

monitored annually since 1983 and all individuals were individually marked (numbered stainless steel

and plastic engraved colour bands; see Rivalan et al. (2010) for details). Unbanded birds of unknown age

(79 individuals since the beginning of the study) and chicks of the year were banded, weighed (body

mass ± 50 g using a Pesola® spring balance) and measured (wing length ± 1 mm with a ruler, tarsus

length, bill length, and bill depth ± 0.1 mm with calipers).

In Amsterdam Island oceanic area, the southern subtropical front (SSTF) delimits the warmer subtropical

from the colder sub-Antarctic waters (Belkin & Gordon 1996). Though the diet and foraging strategy of

Amsterdam albatross remains poorly known, it is presumed to have very similar foraging behaviour

compared to that of the wandering albatross, although subtle differences can appear (Pajot et al. 2021;

see Supplementary for species biological aspects). The wandering albatross is known to forage over

extensive distances, detecting prey visually or by olfaction during the day (Nevitt et al. 2008) referred as

'foraging-in-flight', the lowest energy consuming feeding strategy. However, the strategy tends to

change depending on breeding stage (Phalan et al. 2007; Louzao et al. 2014), and could result in more

frequent and shorter bouts on the water in a 'sit and wait' technique.

Thiebot et al. (2014) showed that adult Amsterdam albatrosses during their post-breeding sabbatical

period moved widely (31° to 115° E), mostly exhibiting westwards wider-scale migratory movements

(sensu Weimerskirch et al. 2015a) reaching >4000 km from the colony exploiting continuously warm

waters (~18°C; see Supplementary). The immature birds moved widely in longitude (0° to 135° E),

exploiting exclusively warm waters 17°-18° C. Juveniles exhibited very large migratory capacities over

the southern Indian Ocean after fledging (15° to 135° E, ~ 4500 km from the colony), through a large

range of latitudinal gradient (27° to 47° S). De Grissac et al. (2016) compared trajectories (i.e. departure

direction or orientation toward specific areas) of juveniles and adults and showed that juveniles

performed an initial rapid movement taking all individuals away from the vicinity of their native colony,

and secondly performed large-scale movements similar to those of adults during the sabbatical period.

GLS are archival light-recording loggers used to study activity of birds over periods lasting up to ~ 2

years. GLSs record the ambient light level every 10 min, from which local sunrise and sunset hours can

be inferred to estimate location every 12 h (Wilson et al. 1992). GLS also recorded saltwater immersion

data by testing for saltwater immersion at regular interval, storing the number of samples wet (>0) at

the end of each 10 min period. We used saltwater immersion to estimate daily activity budget. Despite

the higher mean spatial error of location estimates with these devices (over 100 km; Phillips et al.

2004a), GLS loggers allowed us to track the birds for prolonged periods with minimal disturbance to

them. We considered the following stages with respect to the ages when GLS were deployed (see Table

1): juvenile, as a fledgling equipped with a GLS just before leaving the colony for the first time;

immature, as a non-breeding young bird that had never bred equipped with a GLS when visiting the

colony; adult, as a breeding adult equipped with a GLS during the incubation or brooding period which

successfully fledged a chick and thereafter took a sabbatical year. To date, we have retrieved 40 of the

50 GLS loggers deployed in total over 4 years, from which 33 individual tracks were estimated (Table 1).

Our original aim was to collect activity data on the three stages over a long period of time (>1 year).

These data are available from a total of 10 adults tracked throughout their sabbatical period, 13

immature birds and 10 juvenile birds (up to 3.2 years).

Data processing

The raw immersion data were values from 0 (no immersion or dry, in flight or sitting on the ground) to

200 (permanently immersed in sea water or wet, indicating the number of 3 s periods during 10 min

blocks when the sensor was immersed in saltwater). Loggers recorded the proportion of time in

seawater at 10 min intervals, which we summarized as hours in the water per day (hereafter time spent

on water; 10 min blocks immersion data > 0). This measure is a reliable proxy of foraging effort linked to

foraging behaviour of the species which enters the water principally to forage (Weimerskirch and

Guionnet 2002). Additionally, the duration of the bouts spent entirely immersed (10 min blocks

immersion data = 200) was calculated daily (hereafter referred as wet bouts duration). Conversely,

when birds are not on land, the time spent dry was interpreted as flying (and thus not feeding). The

duration of the bouts spent entirely dry (10 min blocks immersion data = 0) was calculated daily

(hereafter referred as dry bouts duration). Additionally the numbers of bouts (number of wet bouts -

sitting on water-and of dry bouts -flying) were obtained daily. Although the loggers integrated activity

within each 10 min block and so did not provide the exact timing of landings and take-offs, Phalan et al.

(2007) found for comparative purposes that bouts defined as a continuous sequence of 0 values for

flight (dry) and a sequence of values of 1 or greater for wet bouts, were suitable proxies for activity.

To select the data corresponding to periods spent at sea after leaving the breeding site, we used the

following criteria on activity to define the departure time from the colony for each stage: 1) juveniles,

the first bout spent on seawater (wet bouts duration) > 1h based on Argos Platform Transmitters

Terminals (PTT) tracking data (data obtained in a other project and not shown here, Weimerskirch et al.

unpublished data); 2) immatures and adults, the last bout spent flying (dry bouts duration) > 12h based

on PTT tracking data (Weimerskirch et al. unpublished data). Using these criteria we obtained departure

months as follows: 1) the juveniles fledged from the colony from January to March, 2) the immatures

left between April and August, and 3) the departures of sabbatical adults were spread over two periods,

first between December and February and then from May to July.

Statistical analyses

Variation in activity parameters

The aim was to determine whether distinct foraging behaviours could be detected across the patterns of

variation of wet/dry data, and then to appraise how these behaviours varied over time and among

individuals. First, to deal with the fact that wet/dry metrics were interrelated (number of wet bouts

sitting on water and time spent on water, wet bouts duration and dry bouts duration, wet bouts number

and dry bouts number) and to avoid redundancy, we ran principal components analyses (PCA built with

the 'PCA' function, FactoMineR package Lê et al. 2008) to circumvent collinearity issues. To describe

changes in behaviour over time and stages using gradients of activity we ran PCA for i) all stages (PCS;

based on activity data collected during the first ten months post-departure) and for ii) juveniles only, as

an additional goal was to determine changes in activity patterns during the first two years of life (PCJ;

based on activity data collected during the first twenty-nine months post-departure).

Considering all stages, the first three principal components (PCS) explained 94.2% of the total variance.

For juveniles, the first three principal components (PCJ) explained 92.2% of the total variance. The

detailed results of PCA and the variables retained for each axe are summarised in Table 3.

Second, we used generalized additive mixed models (GAMMs, built with the 'gam' function, itsadug and

mgcv package Lin and Zhang 1999; Wood 2015) with the values associated with each of the three first

axes of the PCA as the dependent variables. We ran separate models testing for variability in activity

parameters i) for all stages combined (PCS) and ii) for juveniles (PCJ), based on different duration of

datasets (28 months since departure for juveniles and 9 months since departure for immatures and

adults; see Supplementary; Table S1). Thus, for i) we considered the lowest number of months elapsed

since departure available (9 months since departure). Months elapsed since departure (the duration

elapsed since fledging expressed in month, i.e. the first month after fledging and so on), month of the

year (i.e. January and so on), sex, and stage (only for i)) were included as fixed effects. The interactions

between stage and time were included as fixed effects to test for the prediction that differences should

vanish with time passed since fledging. To test for the importance of individual variability in our results

we built models with or without random effects. We compared models without random effect, models

with random intercepts, and models with random slopes and intercepts to test whether the rate of

change of activity parameters as a function of time elapsed since departure varied between individuals

(Zuur 2009a). Models included month elapsed since departure as a continuous covariate modelled with

non-parametric smoothing functions (Wood 2017). We limited the amount of smoothing (k) with the

'gam.check' function following Wood (2017) for each spline to avoid excessive flexibility and model

overfitting that would have no ecological meaning. Models including all combinations of explanatory

variables and random effects were then tested and ranked using their Akaike Information Criterion (AIC)

values and Akaike weights following the Information-Theoretic Approach (Burnham and Anderson

2002). The model with the lowest AIC was considered as the best model. Two models separated by a

difference in AIC values of less than 2 were assumed to fit the data similarly. Using AICc to account for

small sample sizes did not change model selection.

Although sexes and stages differed for some body size measurements (see details in Supplementary), we

could not include body size as an additional explanatory variable in GAMMs testing for factors of

variation in activity patterns due to small sample sizes in each sex and stage category (see Table 1).

Spatial and statistical analyses were performed using R (R Core Team 2021). Values are means ± SD.

Results

The most parsimonious models explaining variations in activity parameters in the Amsterdam albatross

included time elapsed since departure from the colony, month of the year, stages and sexes (Tables 4

and 5; Supplementary Figures S1 - S5; Tables S1), whatever the synthetic activity variables considered

(PC1S, PC2S and PC3S; Table 4). The interaction between stage and time elapsed was significant for the

first synthetic activity variable (PC1S). Selected models also included random effects on intercepts and

slopes, indicating inter-individual variability in activity and in the rate of change of activity as a function

of time elapsed since departure from the colony.

In juvenile Amsterdam albatrosses, the most parsimonious models explaining variations in activity

included time elapsed since departure from the colony, month of the year for all three activity variables

considered (Table 5 and 6; PC1J, PC2J and PC3J), and sex was retained only for two variables (PC2J and

PC3J). Selected models also included random effects on intercepts and slopes, indicating inter-individual

variability in activity and in the rate of change of activity as a function of time elapsed since departure

from the colony (Supplementary Figures S6).

Changes in activity for all stages

The two synthetic activity variables (PC1S, PC2S) varied significantly with time exhibiting clear nonlinear

temporal patterns (Figure 1). These variations were related to the time elapsed since their departure

from the colony and showed seasonal changes (indicated by the month of the year; Supplementary

Figures S1 - S5; Tables S1 and S2). With increasing time since departure, birds spent lower percentage of

time on water and made shorter wet bouts. They spent less percentage of time on water during the

period March to July compared to rest of the year (PC1S, Supplementary Figures S1 - S5). They made

longer and fewer bouts on water during the period April to November, and shorter flying bouts during

the period November to February. Juveniles showed strong temporal changes in activity linked to the

time elapsed since departure from the colony in the first two months after fledging (Supplementary,

Figure 2). In immatures and adults the temporal pattern appeared reversed compared to juveniles

(Supplementary, Figure 2).

Compared to adults, immatures and even more so juveniles, spent a lower percentage of time on water

(Table 5, Supplementary Figures S1) and made more flying bouts (PC1S; Supplementary Figures S2),

made shorter and fewer bouts on water (PC2S; Supplementary Figures S4-S5), and made longer flying

bouts (PC3S; Supplementary Figures S2). Males spent a higher percentage of time on water and made

fewer flying bouts (PC1S), longer and more numerous bouts on water (PC2S) and shorter flying bouts

(PC3S) compared to females.

Changes in activity of juveniles during the first two years after fledging

PC1J and PC2J varied significantly with time exhibiting clear nonlinear temporal patterns (Figure 3;

Supplementary Figures S7 - S11; Tables S1 and S3a, 3b). Juveniles seemed to alternate periods of lower

percentage of time spent on water combined with more numerous flying bouts (April) with periods of

higher percentage of time on water combined with fewer flying bouts (February, July-October; PC1J, not

illustrated). The seasonal change was also observed through longer and fewer bouts spent on water and

shorter flying bouts at the end of the year (PC2J: September-December). Juveniles, during the first 28

months after fledging, increased the time spent on water while decreasing the number of flying bouts

(Figure 3a).

PC2J and PC3J varied significantly with sex (Supplementary Figures S7 - S10; Tables S5b, 5c), indicating

that juvenile males made shorter and more numerous bouts on water (PC2J) and shorter flying bouts

(PC3J) compared to females (Supplementary Figures S7 - S10; Tables S5b, 5c).

Discussion

In this study, we benefited from a unique comprehensive dataset of remarkable duration (up to 28

months) to characterise the post-fledging behaviour of naïve seabirds. Using miniaturized activity

loggers (GLS), we showed clear differences and changes in activity characteristics depending on life-

stages. By comparing changes in behaviour at sea and foraging parameters of juveniles after their

departure at sea with those of immatures and adults in the Amsterdam albatross, we showed that

juveniles differed from immatures and adults in their activity values and patterns. Activity also varied

according to time and sex. Our study allows us to compare foraging behaviour among life stages in a

long-lived endangered seabird species, while also providing new insights into the development of

foraging patterns in naïve individuals over a multi-year period.

Stage specific changes

The birds were found to behave differently according to their stage whatever the activity variables

considered, indicating differences in foraging behaviour. Overall, juveniles spent lower percentage of

time on water compared to immatures and adults. During the first months following their departure

from the colony the proportion of time spent on water by immatures and adults showed a dome-shaped

curve peaking three to five months after departure. During the same period of time, the proportion of

time spent on water by the juveniles changed strongly, with values dropping off in the first two months

and then remaining low and overall lower than in adults and immatures. This might indicate a lower

foraging activity in naïve birds. During the same period, the duration and number of water bouts also

exhibited progressive change. These patterns suggest an early and gradual change in foraging behaviour

so that juveniles progressively could behaved similarly to immatures and adults (reaching similar values

in activity covariates). This suggest a progressive behavioural change in movements during the first two

months after fledging. It is noteworthy that the multi-monthly bell-shaped pattern observed during the

first 10 months after departure in immatures and adults appears to be mirrored in juveniles 15-16

months later (see Figure S11). Despite such possible similarities (% time spent on water), there are still

some differences between stages (see Supplementary). Since % time spent on water partly reflects

foraging, this seems to indicate that juvenile individuals may have weaker foraging skills during their first

two months at sea. Although behavioural changes can often equate to improved performance (e.g.

Campioni et al. 2020) this is not always the case. The emergence of juvenile birds as more 'adult like' in

their foraging/flight behavioural patterns is not necessarily a sign of improvement. For example, it could

be partly due to individual differences in area use with different environmental conditions encountered

(food abundance, wind regimes).

Since all stages of the Amsterdam albatross forage in the same water masses (see Thiebot et al. 2014),

differences in foraging behaviour were presumably not due to different oceanographic characteristics as

observed in other species (Thiers et al. 2014; Weimerskirch et al. 2014; Frankish et al. 2020b). These

differences could be due to a combination of lack of experience, poor knowledge of the environment,

use of distinct cues and/or physical immaturity leading to sub-optimal behaviour (Shaffer et al. 2001;

Frankish et al. 2020a, 2022). It is likely that increasing exposure to diverse foraging situations allows

juveniles to rapidly accumulate foraging experience and improve various aspects of foraging.

Results suggest that immatures may differ from both adults and juveniles in some aspects of their

behaviour. While most of the activity parameters and the temporal patterns showed similarities with

adults when considering the time elapsed since departure, they seemed rather comparable to juveniles

when considering seasonal aspects (month of the year). Such differences can be explained by several

non-exclusive explanations: i) similar management of energy constraints than adults, as post-breeding

adults and immatures are less constrained in their central-place foraging strategies (Campioni et al.

2016), ii) comparable capacity to respond to local resource availability in their foraging behaviour than

juveniles (Frankish et al. 2022), and iii) incomplete acquisition of more long-term learning of complex

movement strategies (Thorup et al. 2003; Votier et al. 2011; Rotics et al. 2016). Disentangling these

hypotheses can be achieved by combining higher resolution movement data with longer longitudinal

studies covering all three life stages for the same individuals.

What might be designated as '*lower performance*' of juveniles found in our study is consistent with

studies on wandering albatrosses and Amsterdam albatrosses (Riotte-Lambert and Weimerskirch 2013;

de Grissac et al. 2017; Pajot et al. 2021) during the first weeks at sea. Juvenile albatrosses behaved

differently in first month after fledging (i.e. speed and sinuosity of movements) and readily use similar

foraging strategies as adults (Frankish et al. 2022). Additional skills (such as detection of prey at the

surface, detection of other foraging seabirds, navigational skills...) need to be acquired during the

immature period before the efficiency of these behaviours matches that of adults. This is also typical of

other seabird taxa, which show progressive improvement in flight performance with the numbers of

days since fledging (Yoda et al. 2004; Mendez et al. 2017; Collet et al. 2020; Corbeau et al. 2020;

Frankish et al. 2022). For example juvenile brown boobies *Anous stolidus* improved their flight abilities

(Yoda et al. 2004). In contrast, flight capability (flight speed and sinuosity) comparable to that of adults

allows juvenile white-chinned petrels *Procellaria aequinoctialis* to rapidly fly over long distances away

from the colony (Frankish et al. 2020).

The progressive change of movement behaviours (foraging parameters estimated from activity

parameters improved with time elapsed) quantified in juvenile Amsterdam albatrosses could be either

due to physical development and/or experience gain. Elucidating the mechanisms of the transition to

independence in early life stages is however crucial for understanding the causes of higher juvenile

mortality in long-lived species (Fay et al. 2015; Payo-Payo et al. 2016).

Temporal changes and sex differences in activity

The temporal variability of activity was found whatever the life-stage considered. Part of the activity

changes observed following the departure of juvenile Amsterdam albatrosses may illustrate the swift

change in travel and movement behaviour, reflecting a more '*adult-like*' behaviour, not indicating

necessarily an improvement of flight performances and of the ability to cope with changing (i.e.

increasing wind speed) wind conditions (Sergio et al. 2014), a key parameter for soaring seabirds such as

albatrosses. Both extrinsic (i.e. environmental conditions) and intrinsic (i.e. energetic demands linked to

plumage moult) factors could be involved in the modulation of foraging behaviour, which can be

reflected in the temporal variability.

Moult is an intrinsically costly process requiring time, energy and nutrients (Langston and Rohwer 1996;

Ellis and Gabrielsen 2002), and the annual replacement of flight feathers is crucial to ensure efficiency in

both flight and thermoregulation (Murphy 1996; Peery et al. 2008; Gutowsky et al. 2014). Stage-specific

and sex-specific differences in moult extent occur in wandering albatross, suggesting important

constraints (Weimerskirch 1991; see Supplementary). Adult birds during the non-breeding season

appear to spend much more time on the water during winter, suggesting that partial moult may occur at

this time (Weimerskirch et al. 2015b, 2020). Interestingly, immature individuals appear to have a similar

peak in time spent on the water in spring, suggesting different timing of moult.

Consistently, we found that males flew for longer periods (dry bouts duration) compared to females.

When considering all stages, males spent a higher percentage of time on water compared to females.

Males in all stages did more bouts on water and juvenile males shorter wet bouts, compared to females.

Contrary to the wandering albatross (Weimerskirch et al. 2014), male and female Amsterdam

albatrosses forage in similar oceanic water masses and encounter comparable wind conditions (Jaeger

et al. 2013; Thiebot et al. 2014). Therefore, it is unlikely that sex differences in activity parameters were

caused by differences in foraging habitats.   Sex-specific behavioural differences are common in sexually

dimorphic seabirds as reviewed in Wearmouth and Sims (2008), where the smaller sex usually

undertakes longer trips. Sexual size dimorphism can result in differences in aerial agility, foraging area

and behaviour, and provisioning rate and preferred prey (Gonzales-Solis et al. 2000; Phillips et al. 2004b,

2011; Weimerskirch et al. 2009; Austin et al. 2019; Barbraud et al. 2021). It has also been suggested that

size matters probably because the smaller and lighter sex has a higher foraging and flight efficiency

(Shaffer et al. 2001; Clay et al. 2020), suggesting that lighter and lower wing loaded female wandering

albatrosses, compared to males, are probably better able to exploit subtropical and tropical waters

where winds are lighter. Following this, it can be hypothesized that female Amsterdam albatrosses have

a greater advantage in foraging in the subtropical environment than males. Sex differences in the

acquisition of foraging performance during the first months after fledging yet remain to be explored.

Individual variability in activity

There was inter-individual variability in almost all activity parameters whatever the stage considered. In

juveniles, models indicated inter-individual variability in activity and in the rate of change of activity as a

function of time elapsed since departure from the colony. Since the intercept terms in the models were

significant, it seems as though individual variability (i.e., specialization on different foraging strategies)

was a contributor to observed variability. However, the rate of change of intra-individual variation for

some foraging strategies (percentage of time on water-number of flying bouts axis) oscillated during the

juvenile period with a seemingly remarkable synchrony (see Fig S7). This suggests that changes in

foraging behaviours occurred at the individual level during the juvenile period without stabilizing, at

least during the first two years after fledging. This individual variability suggests development of

specialized individual foraging behaviours (Harel et al. 2016; Rotics et al. 2016, 2021; Phillips et al. 2017).

Nonetheless, given the small sample sizes these results should be interpreted with caution.

Conclusion

Very few studies have investigated the ontogeny of foraging behaviour over such a long period of time,

particularly in long-lived pelagic seabirds, due to the difficulty of obtaining individual tracking data over

several years. We investigated the foraging behaviour, through activity patterns, during the three life

stages of the endangered Amsterdam albatross by using miniaturized activity loggers. Naïve juveniles

during their first month at sea after leaving their colony exhibited lower foraging activity (greater

proportion of time spent sitting on water, longer and more numerous bouts on water, and shorter and

fewer flying bouts). Patterns of activity parameters in juveniles after independence suggested a

progressive change of foraging performances during the first two months after fledging. Regardless of

life stage considered, activity parameters exhibited temporal variability reflecting the modulation of

foraging behaviour presumably linked to both extrinsic (i.e. environmental conditions such as variability

in food resources or in wind) and intrinsic (i.e. energetic demands linked to plumage renew during

moult) factors. Sex differences in activity parameters according to time since departure from the colony

and season were consistent with the sexual dimorphism in the Amsterdam albatross. Overall, an

expected change in behaviour, resulting from the experience gained, may reflect an improvement in

skills occurring during a period of learning through the immaturity phase, which would still need to be

confirmed by directly assessing foraging performance. Results from our study suggest that the lower

foraging activity of juvenile Amsterdam albatrosses may partly explain their lower survival compared to

adults. However, despite foraging metrics of juveniles investigated here reached values observed for

adults quite rapidly, immature survival remains lower than adult survival during several years (Rivalan et

al. 2010). Therefore, more detailed information such as foraging performance realized (the amount or

quality of preys obtained by juveniles) or energetic expenditure are needed to better understand age-

specific differences in life history traits such as survival and the dynamics of the species.

Ethics

All work was carried out was approved by the French Polar Institute (IPEV) ethics committee.

Acknowledgements

This study was made possible thanks to all the fieldworkers involved in the monitoring program on
Amsterdam albatross, namely Jean-Baptiste Thiebot, Jérémy Demay, Rémi Bigonneau, Romain Bazire,
Hélène Le Berre, Marine Quintin, Marine Devaud, Chloé Tanton, Jérémy Dechartre and Anthony Le
Nozahic. We acknowledge Dominique Joubert for the management of the demographic CEBC Seabirds
database. We are grateful to Richard Phillips, British Antarctic Survey, Cambridge for providing GLS
loggers. We thank Bertille Mohring for statistical advices on principal components analyses. We thank
our PCI recommender Blandine Doligez and reviewers, Juliet Lamb and an anonymous reviewer for
constructive comments on an earlier version of the manuscript.

Funding

This monitoring program was supported financially and logistically by the French Polar Institute IPEV
(program 109, PI C. Barbraud/H. Weimerskirch), the Zone Atelier Antarctique (CNRS-INEE), Terres
Australes et Antarctiques Françaises.

This study is a contribution to the National Plan of Actions for Amsterdam albatross.

Authors contribution

K.D. and C.B. conceived the study. H.W. secured funding. K.D. prepared and analysed the data. C.B.
provided feedback on the analyses. K.D. wrote the first draft and all authors contributed to editing
versions of the manuscript.

Table 1 Chronological characteristics of life-cycle stages (adapted from Thiebot et al. 2014) and sample sizes of birds tracked using Global

Location Sensing (GLS) of Amsterdam albatross

Stage[1]	Definition	Age[1]	Tracking duration	Behaviour	Years of deployment	Deployed (n)	Recovered (n)	Recovery rate (%)	GLS with data (n)
Juvenile	Following chick fledging in January	1st year	~2.5 years	Chicks disperse at sea after leaving the colony for the first time	2011	21	12	57 (t+9)	10 (4 F - 6 M)[2]
Immature	After juvenile movements, until first breeding attempt (at an average age of 9 years old)	~2-10 years	~1 year	Non-breeding young birds forage at sea and occasionally visit the colony for mating	2011-2012	18	17	94	13 (3 F - 9 M – 1 NK)
Adult sabbatical	Between two successive breeding periods (~ 15 January year t to the following 15 January year $t+1$)	~8-28 years	~1 year	Breeding adults at the end of reproductive cycle and leave the colony to forage at sea	2006, 2009	11	11	100	10 (6 F - 4 M)

460 [1] Stage/Age at which the individuals were equipped with loggers in our study; [2] number of females F and males M, or not known NK for each stage

461 Table 2 Hypotheses and predictions about the factors driving differences in activity (time spent on water, number and duration of

462 flying bouts, number and duration of water bouts) year-round in Amsterdam albatrosses. The two hypotheses are not mutually

463 exclusive.

Hypothesis	Predictions		
	(1) Time spent on water (%)	(2) Flying bouts (number/duration)	(3) Water bouts (number/duration)
(A) Age and stage specific	Juveniles: increased foraging time/effort and thus lower time spent on water (1), longer flying bouts (2) and shorter water bouts (3) than other stages		
(B) Temporal changes - internal requirements: moult/energetic effects	Adults/immatures: two-periods pattern including one with lowering activity Juveniles: change in foraging skills corresponding to gradual change with less time sitting on water (lower time spent on water (1)), increasing flying bouts duration and number (2) and decreasing water bouts duration and number (3), during the 1st month after fledging Adjustment in foraging effort to energetic requirements or moult constraints according to time elapsed since departure: higher time spent on water (1), lower flying bouts duration and number (2) and higher water bouts duration and number (3) during moulting		

464 Table 3 Results of principal components analyses (PCA) on six wet/dry metrics on Amsterdam albatross

Life-stages	Principal components	Total variance explained (%)	Time spent on water	Dry bouts duration	Dry bouts number	Wet bouts duration	Wet bouts number
All	First (PC1S)	41.5	+ (r = 0.97)[1]		- (r = -0.79)		
	Second (PC2S)	32.5				+ (r = 0.79)	- (r = -0.75)
	Third (PC3S)	20.2		+ (r = 0.74)	- (r = -0.44)		
Juveniles	First (PC1J)	42.3	+ (r = 0.98)		- (r = -0.76)		
	Second (PC2J)	32.2				+ (r = 0.72)	- (r = -0.75)
	Third (PC3J)	20.7		+ (r = 0.48)	- (r = -0.46)	- (r = -0.46)	

465 [1] the symbol used gives the sign of the correlation (+: positive, -: negative); the number in brackets indicates the value of the correlation
466 coefficient r

469   Table 4 Model selection for variation in activity parameters of Amsterdam albatrosses in relation to sex, stage, number of months spent since

470   departure (month elapsed: duration elapsed since fledging expressed in month, i.e. the first month after fledging and so on) and month of the

471   year (i.e. January and so on)

Models	Fixed effects	Random effects	AIC	ΔAIC
Proportion of time spent on water (PC1S)				
M_5	Month elapsed + Month + Stage + Sex + Month elapsed: Stage	Month elapsed: Individual	26461.62	
M_4	Month elapsed + Month + Stage + Sex	Month elapsed: Individual	27224.79	-763.17
M_3	Month elapsed + Month + Stage	Month elapsed: Individual	27276.18	-814.56
M_2	Month elapsed + Month	Month elapsed: Individual	28322.25	-1860.63
M_1	Month elapsed	Month elapsed: Individual	28591.69	-2130.07
M_0	Null model		28874.42	-2412.80
Bouts spent on water (PC2S)				
M_3	Month elapsed + Month + Stage	Month elapsed: Individual	25751.47	
M_4	Month elapsed + Month + Stage + Sex	Month elapsed: Individual	25752.62	-1.15
M_2	Month elapsed + Month	Month elapsed: Individual	25756.37	-4.90
M_1	Month elapsed	Month elapsed: Individual	25803.80	-52.33
M_5	Month elapsed + Month + Stage + Month elapsed: Stage	Month elapsed: Individual	26750.55	-999.08
M_0	Null model		26903.12	-1151.65

Bouts spent dry -flying (PC3S)				
M_4	Month + Stage + Sex	Month elapsed: Individual	22427.29	
M_3	Month	Month elapsed: Individual	22509.79	-8.14
M_2	Month elapsed	Month elapsed: Individual	22539.75	-82.50
M_1	Null model	Month elapsed: Individual	22540.25	-112.96
M_0	Null model		23042.26	-614.97

472   Models are ranked according to decreasing statistical support, as indicated by AIC. The first best models are shown

473   Table 5 Values of activity parameters (mean ± sd) recorded using Global Location Sensing (GLS) depending on stage and sex of Amsterdam

474   albatross

	Juvenile[1]		Juvenile[2]		Immature		Adult sabbatical	
	female	male	female	male	female	male	female	male
Time spent on water (%)	55.04 ± 20.46	58.18 ± 21.11	51.41 ± 19.18	52.88 ± 20.39	59.25 ± 21.53	63.31 ± 21.17	64.89 ± 20.90	69.98 ± 18.10
Wet bouts (sitting on water) duration (h)	1.21 ± 1.74	1.24 ± 1.76	1.16 ± 1.73	1.12 ± 1.59	1.07 ± 1.31	1.48 ± 2.12	1.47 ± 1.95	1.33 ± 1.96
Dry bouts duration (h)	1.29 ± 1.37	1.21 ± 1.32	1.34 ± 1.41	1.26 ± 1.40	1.32 ± 1.42	1.28 ± 1.55	1.44 ± 1.56	1.31 ± 1.42
Wet bouts (sitting on water) number	8.71 ± 4.01	8.76 ± 4.09	8.14 ± 3.85	8.48 ± 4.11	10.34 ± 4.29	8.59 ± 4.24	8.96 ± 3.98	10.28 ± 5.33
Dry bouts number	7.06 ± 3.20	7.27 ± 3.52	7.57 ± 3.21	7.85 ± 3.50	6.31 ± 3.21	5.75 ± 2.99	5.01 ± 2.64	4.64 ± 2.48

476   [1] calculated during 28 months following departure; [2] calculated during 9 months following departure

477 Table 6 Model selection for variation activity parameters for juvenile Amsterdam albatrosses in relation to sex, number of months spent since

478 departure (month elapsed: duration elapsed since fledging expressed in month, i.e. the first month after fledging and so on) and month of the

479 year (i.e. January and so on)

Models	Fixed effects	Random effects	AIC	ΔAIC
Proportion of time spent on water (PC1J)				
M_2	Month elapsed + Month	Month elapsed: Individual	21625.69	
M_1	Month elapsed	Month elapsed: Individual	21864.11	-238.42
M_0	Null model		22109.52	-483.83
Bouts spent on water (PC2J)				
M_3	Month elapsed + Month + Sex	Month elapsed: Individual	19999.00	
M_2	Month elapsed + Month	Month elapsed: Individual	20004.65	-5.65
M_1	Month elapsed	Month elapsed: Individual	20072.42	-73.42
M_0	Null model		20417.76	-418.76
Bouts spent dry -flying (PC3J)				
M_3	Month + Sex	Month elapsed: Individual	17541.02	
M_2	Month elapsed	Month elapsed: Individual	17549.00	-7.98
M_1	Null model	Month elapsed: Individual	17548.75	-7.73
M_0	Null model		17708.47	-167.45

Models are ranked according to decreasing statistical support, as indicated by AIC. The first best models are show

FIGURES

Figure 1 Modeled a) first and b) second axis of principal components analysis of activity parameters for

all stages (e.g. adult, immature and juvenile) of Amsterdam albatrosses according to time elapsed (e.g.

duration elapsed since departure from the colony expressed in month). Plain line corresponds to

estimated smoother from the GAMM model. Dotted lines indicate 95% confidence interval. Boxplot

represent raw data. The first axis correlated positively with time spent on water and negatively with dry

bouts number and the second axis correlated positively with wet bouts duration and negatively with wet

bouts number.

Figure 2 Modeled a) first, b) second and c) third axis of principal components analysis of activity

parameters for all stages i.e. adult (plain black line), immature (dashed red line) and juvenile (dotted

green line) in Amsterdam albatrosses according to time elapsed (e.g. duration elapsed since departure

from the colony expressed in month). Plain line corresponds to estimated smoother from the GAMM

model

Figure 3 Modeled a) first and b) second axis of principal components analysis of activity parameters of

juveniles of Amsterdam albatrosses according to time elapsed (e.g. duration elapsed since departure

from the colony expressed in month). Plain line corresponds to estimated smoother from the GAMM

model. Dotted lines indicate 95% confidence interval. Boxplot represent raw data.

Figure 1a

Figure 1b

Figure 2a

Figure 2b

Figure 2c

Figure 3a

Conflict of interest disclosure

We, the authors of this article declare that we have no financial conflict of interest with the content of
this article.

REFERENCES

Alerstam T, Lindström Å (1990) Optimal bird migration: the relative importance of time, energy,
and safety. In: Bird migration. Springer, pp 331–351

Austin RE, De Pascalis F, Arnould JP, et al (2019) A sex-influenced flexible foraging strategy in
a tropical seabird, the magnificent frigatebird. Marine Ecology Progress Series 611:203–
214

Barbraud C, Delord K, Kato A, et al (2019) Sexual segregation in a highly pagophilic and
sexually dimorphic marine predator. bioRxiv 472431

Becker J (2014) Dispersal and Genetic Variation in a Naturally Fragmented Environment:
Insights from a Long□term Study of White□throated Dippers (Cinclus cinclus)

Boynton CK, Mahony NA, Williams TD (2020) Barn Swallow (Hirundo rustica) fledglings use
crop habitat more frequently in relation to its availability than pasture and other habitat
types. The Condor 122:duz067

Bruner JS (1972) Nature and uses of immaturity. American psychologist 27:687

Burnham KP, Anderson DR (2002) Model Selection and Multimodel Inference: A Practical
Information-Theoretic Approach. NewYork,NY.

Burns JM, Clark CA, Richmond JP (2004) The impact of lactation strategy on physiological
development of juvenile marine mammals: implications for the transition to independent
foraging. Elsevier, pp 341–350

Campioni L, Dias MP, Granadeiro JP, Catry P (2020) An ontogenetic perspective on migratory
strategy of a long□lived pelagic seabird: Timings and destinations change progressively
during maturation. Journal of Animal Ecology 89:29–43

Campioni L, Granadeiro JP, Catry P (2016) Niche segregation between immature and adult
seabirds: does progressive maturation play a role? Behavioral Ecology 27:426–433

Caubet Y, Jaisson P, Lenoir A (1992) Preimaginal induction of adult behaviour in insects. The
Quarterly Journal of Experimental Psychology Section B 44:165–178

Cherel Y, Quillfeldt P, Delord K, Weimerskirch H (2016) Combination of at-sea activity,
geolocation and feather stable isotopes documents where and when seabirds molt.
Frontiers in Ecology and the Environment 4:1–16

Clay TA, Joo R, Weimerskirch H, et al (2020) Sex☐specific effects of wind on the flight
decisions of a sexually dimorphic soaring bird. Journal of Animal Ecology 89:1811–1823

Collet J, Prudor A, Corbeau A, et al (2020) First explorations: ontogeny of central place foraging
directions in two tropical seabirds. Behavioral Ecology 31:815–825

Corbeau A, Prudor A, Kato A, Weimerskirch H (2020a) Development of flight and foraging
behaviour in a juvenile seabird with extreme soaring capacities. Journal of Animal
Ecology 89:20–28

Corbeau A, Prudor A, Kato A, Weimerskirch H (2020b) Development of flight and foraging
behaviour in a juvenile seabird with extreme soaring capacities. Journal of Animal
Ecology 89:20–28. https://doi.org/10.1111/1365-2656.13121

Daunt F, Afanasyev V, Adam A, et al (2007) From cradle to early grave: juvenile mortality in
European shags Phalacrocorax aristotelis results from inadequate development of
foraging proficiency. Biology Letters 3:371–374. https://doi.org/10.1098/rsbl.2007.0157

de Grissac S, Bartumeus F, Cox SL, Weimerskirch H (2017) Early☐life foraging: Behavioral
responses of newly fledged albatrosses to environmental conditions. Ecology and
evolution 7:6766–6778

de Grissac S, Börger L, Guitteaud A, Weimerskirch H (2016) Contrasting movement strategies
among juvenile albatrosses and petrels. Scientific reports 6:26103

Delord K, Barbraud C, Bost CA, et al (2013) Atlas of top predators from the French Southern
Territories in the Southern Indian Ocean.
http://www.cebc.cnrs.fr/ecomm/Fr_ecomm/ecomm_ecor_OI1.html

Dukas R (2006) Learning in the context of sexual behaviour in insects. Animal Biology 56:125–
141

Ellis HI, Gabrielsen GW (2002) Energetics of free-ranging seabirds. In: Seabirds energetics,
Whittow, G. C. and Rahn, H. Plenum Press, N.Y., pp 359–408

Evans DR (2018) The post-fledging survival and movements of juvenile Barn Swallows
(Hirundo rustica): an automated telemetry approach

Fay R, Weimerskirch H, Delord K, Barbraud C (2015) Population density and climate shape
early☐life survival and recruitment in a long☐lived pelagic seabird. Journal of Animal
Ecology 84:1423–1433

Frankish CK, Manica A, Clay TA, et al (2022) Ontogeny of movement patterns and habitat
selection in juvenile albatrosses. Oikos e09057

Frankish CK, Manica A, Phillips RA (2020a) Effects of age on foraging behavior in two closely
related albatross species. Movement ecology 8:7

Frankish CK, Phillips RA, Clay TA, et al (2020b) Environmental drivers of movement in a
threatened seabird: insights from a mechanistic model and implications for conservation.
Diversity and Distributions 26:1315–1329

Furness RW (1988) Influences of status and recent breeding experience on the moult strategy of
the Yellow-nosed albatross Diomedea chlororhynchos. J Zool 215:719–727

Gonzales-Solis J, Croxall JP, Wood AG (2000) Sexual dimorphism and sexual segregation in
foraging strategies of northern giant petrels Macronectes halli during incubation. Oikos
90:390–398

Gutowsky SE, Gutowsky LF, Jonsen ID, et al (2014) Daily activity budgets reveal a quasi-
flightless stage during non-breeding in Hawaiian albatrosses. Movement ecology 2:23–23

Harel R, Horvitz N, Nathan R (2016a) Adult vultures outperform juveniles in challenging
thermal soaring conditions. Scientific reports 6:27865

Harel R, Horvitz N, Nathan R (2016b) Adult vultures outperform juveniles in challenging
thermal soaring conditions. Scientific reports 6:27865

Heyes CM (1994) Social learning in animals: categories and mechanisms. Biological Reviews
69:207–231

Jaeger A, Jaquemet S, Phillips RA, et al (2013) Stable isotopes document inter- and intra-
specific variation in feeding ecology of nine large southern Procellariiformes. Mar Ecol
Prog Ser 490:255–266

Jouventin P, Martinez J, Roux J-P (1989) Breeding biology and current status of the Amsterdam
Island Albatross *Diomedea amsterdamensis*. Ibis 131:182-

King JR (1974) Seasonal allocation of time and energy resources in birds. Avian energetics 4–85

Krüger S, Reid T, Amar A (2014) Differential range use between age classes of southern African
Bearded Vultures Gypaetus barbatus. PLoS One 9:

Lack D (1954) The natural regulation of animal numbers. The Natural Regulation of Animal
Numbers

Langen TA (1996) Skill acquisition and the timing of natal dispersal in the white-throated
magpie-jay, Calocitta formosa. Animal Behaviour 51:575–588

Langston NE, Rohwer S (1996) Molt-breeding tradeoffs in albatrosses: life history
implications for big birds. Oikos 76:498–510

Lê S, Josse J, Husson F (2008) FactoMineR: an R package for multivariate analysis. Journal of
statistical software 25:1–18

Lin X, Zhang D (1999) Inference in generalized additive mixed modelsby using smoothing splines. Journal of the royal statistical society: Series b (statistical methodology) 61:381–400

Louzao M, Wiegand T, Bartumeus F, Weimerskirch H (2014) Coupling instantaneous energy-budget models and behavioural mode analysis to estimate optimal foraging strategy: an example with wandering albatrosses. Movement ecology 2:1–15

Marchetti K, Price T (1989) Differences in the foraging of juvenile and adult birds: the importance of developmental constraints. Biological Reviews 64:51–70

Margrath R, Lill A (1985) Age-related differences in behaviour and ecology of crimson rosellas, Platycercus elegans, during the non-breeding season. Wildlife Research 12:299–306

Martin P, Bateson P (1985) The ontogeny of locomotor play behaviour in the domestic cat. Animal behaviour 33:502–510

Mendez L, Prudor A, Weimerskirch H (2017) Ontogeny of foraging behaviour in juvenile red-footed boobies (Sula sula). Scientific reports 7:1–11

Murphy ME (1996) Energetics and nutrition of molt. In: Avian energetics and nutritional ecology. Springer, pp 158–198

Mushinsky HR, Hebrard JJ, Vodopich DS (1982) Ontogeny of water snake foraging ecology. Ecology 63:1624–1629

Naef-Daenzer B, Grüebler MU (2008) Post-fledging range use of Great Tit Parus major families in relation to chick body condition. Ardea 96:181–190

Nevitt GA, Losekoot M, Weimerskirch H (2008) Evidence for olfactory search in wandering albatross, Diomedea exulans. Proceedings of the National Academy of Sciences 105:4576–4581

Nourani E, Vansteelant WMG, Byholm P, Safi K (2020) Dynamics of the energy seascape can explain intra-specific variations in sea-crossing behaviour of soaring birds. Biology Letters 16:20190797. https://doi.org/10.1098/rsbl.2019.0797

Pajot A, Corbeau A, Jambon A, Weimerskirch H (2021) Diel at□sea activity of two species of great albatrosses: the ontogeny of foraging and movement behaviour. Journal of Avian Biology 52:

Payo-Payo A, Genovart M, Bertolero A, et al (2016) Consecutive cohort effects driven by density-dependence and climate influence early-life survival in a long-lived bird. Proceedings of the Royal Society B: Biological Sciences 283:20153042

Peery MZ, Henkel LA, Newman SH, et al (2008) Effects of rapid flight-feather molt on postbreeding dispersal in a pursuit-diving seabird. The Auk 125:113–123

Pennycuick CJ (1982) The flight of petrels and albatrosses (Procellariiformes), observed in
South Georgia and its vicinity. Phil Trans R Soc Lond 300:75–106

Peron C, Gremillet D (2013) Tracking through Life Stages: Adult, Immature and Juvenile
Autumn Migration in a Long-Lived Seabird. Plos One 8:

Phalan B, Phillips RA, Silk JR, et al (2007) Foraging behaviour of four albatross species by night
and day. Marine Ecology Progress Series 340:271–286

Phillips RA, Lewis S, González-Solís J, Daunt F (2017) Causes and consequences of individual
variability and specialization in foraging and migration strategies of seabirds. Marine
Ecology Progress Series 578:117–150

Phillips RA, Mcgill RAR, Dawson DA, Bearhop S (2011) Sexual segregation in distribution, diet
and trophic level of seabirds: insights from stable isotope analysis. Marine Biology
158:2199–2208

Phillips RA, Silk JRD, Croxall JP, et al (2004a) Accuracy of geolocation estimates for flying
seabirds. Marine Ecology-Progress Series 266:265–272

Phillips RA, Silk JRD, Phalan B, et al (2004b) Seasonal sexual segregation in two *Thalassarche*
albatross species: competitive exclusion, reproductive role specialization or foraging
niche divergence? Proceedings of the Royal Society of LondonSeries B: Biological
Sciences 271:1283–1291

Rapaport LG, Brown GR (2008) Social influences on foraging behavior in young nonhuman
primates: learning what, where, and how to eat. Evolutionary Anthropology: Issues,
News, and Reviews: Issues, News, and Reviews 17:189–201

Riotte-Lambert L, Weimerskirch H (2013) Do naive juvenile seabirds forage differently from
adults? Proceedings of the Royal Society B: Biological Sciences 280:20131434

Rivalan P, Barbraud C, Inchausti P, Weimerskirch H (2010) Combined impact of longline
fisheries and climate on the persistence of the Amsterdam albatross. Ibis 152:6–18

Rotics S, Kaatz M, Resheff YS, et al (2016a) The challenges of the first migration: movement
and behaviour of juvenile vs. adult white storks with insights regarding juvenile
mortality. Journal of Animal Ecology 85:938–947

Rotics S, Kaatz M, Resheff YS, et al (2016b) The challenges of the first migration: movement
and behaviour of juvenile vs. adult white storks with insights regarding juvenile
mortality. Journal of Animal Ecology 85:938–947

Rotics S, Turjeman S, Kaatz M, et al (2021) Early-life behaviour predicts first-year survival in a
long-distance avian migrant. Proceedings of the Royal Society B 288:20202670

Roux JP, Jouventin P, Mougin JL, et al (1983) Un nouvel albatros Diomedea amsterdamensis n.
sp. découvert sur l'île Amsterdam (37°50'S, 77°35'E). L'Oiseau et R F O 53:1–11

Ruaux G, Lumineau S, de Margerie E (2020) The development of flight behaviours in birds.
Proceedings of the Royal Society B 287:20200668

Sergio F, Tanferna A, De Stephanis R, et al (2014) Individual improvements and selective
mortality shape lifelong migratory performance. Nature 515:410–413

Shaffer SA, Weimerskirch H, Costa D (2001) Functional significance of sexual dimorphism in
wandering albatrosses, Diomedea exulans. Functional Ecology 15:203–210

Team RC (2019) R: A Language and Environment for Statistical Computing (Version 3.5. 2, R
Foundation for Statistical Computing, Vienna, Austria, 2018). There is no corresponding
record for this reference[Google Scholar]

Thiebot J-B, Delord K, Marteau C, Weimerskirch H (2014) Stage-dependent distribution of the
critically endangered Amsterdam albatross in relation to Economic Exclusive Zones.
Endangered Species Research 23:263–276

Thiers L, Delord K, Barbraud C, et al (2014) Foraging zones of the two sibling species of giant
petrels in the Indian Ocean throughout the annual cycle: implication for their
conservation. Mar Ecol Prog Ser 499:233–248

Thorup K, Alerstam T, Hake M, Kjellén N (2003) Bird orientation: compensation for wind drift
in migrating raptors is age dependent. Proceedings of the Royal Society of London Series
B: Biological Sciences 270:S8–S11

Urios V, López-López P, Limiñana R, Godino A (2010) Ranging behaviour of a juvenile
Bearded Vulture (Gypaetus barbatus meridionalis) in South Africa revealed by GPS
satellite telemetry

Votier SC, Grecian WJ, Patrick S, Newton J (2011) Inter-colony movements, at-sea behaviour
and foraging in an immature seabird: results from GPS-PPT tracking, radio-tracking and
stable isotope analysis. Marine biology 158:355–362

Wearmouth VJ, Sims DW (2008) Sexual segregation in marine fish, reptiles, birds and
mammals: behaviour patterns, mechanisms and conservation implications. Advances in
marine biology 54:107–170

Weimerskirch H (1991) Sex-specific differences in molt strategy in relation to breeding in the
wandering albatross. Condor 93:731–737

Weimerskirch H, Akesson S, Pinaud D (2006) Postnatal dispersal of wandering albatrosses
Diomedea exulans: implications for the conservation of the species. J Avian Biol 37:23–
28

Weimerskirch H, Brothers N, Jouventin P (1997a) Population dynamics of Wandering Albatross
Diomedea exulans and Amsterdam Albatross D. amsterdamensis in the Indian Ocean and
their relationship with long-line fisheries: conservation implications. Biological
conservation 79:257–270

Weimerskirch H, Cherel Y, Delord K, et al (2014) Lifetime foraging patterns of the wandering
albatross: Life on the move! Journal of Experimental Marine Biology and Ecology
450:68–78

Weimerskirch H, de Grissac S, Ravache A, et al (2020) At-sea movements of wedge-tailed
shearwaters during and outside the breeding season from four colonies in New Caledonia.
Marine Ecology Progress Series 633:225–238

Weimerskirch H, Delord K, Guitteaud A, et al (2015a) Extreme variation in migration strategies
between and within wandering albatross populations during their sabbatical year, and
their fitness consequences. Scientific Reports 5:8853-

Weimerskirch H, Guionnet T (2002) Comparative activity pattern during foraging of four
albatross species. Ibis 144:40–50

Weimerskirch H, Shaffer SA, Tremblay Y, et al (2009) Species-and sex-specific differences in
foraging behaviour and foraging zones in blue-footed and brown boobies in the Gulf of
California. Marine Ecology Progress Series 391:267–278

Weimerskirch H, Tarroux A, Chastel O, et al (2015b) Population-specific wintering distributions
of adult south polar skuas over the three oceans. Mar Ecol Prog Ser 538:229–237

Weimerskirch H, Wilson RP, Lys P (1997b) Activity pattern of foraging in the wandering
albatross: a marine predator with two modes of prey searching. Marine Ecology Progress
Series 151:245–254

Wilson RP, Ducamp JJ, Rees G, et al (1992) Estimation of location: global coverage using light
intensity. In: Priede IMSS (ed). Ellis Horward, Chichester, pp 131–134

Wood S (2015) Package 'mgcv.' R package version 1:729

Wood S (2017) Generalized Additive Models: An Introduction with R, 2nd Edition Chapman
and Hall//CRC, Boca Raton. Chapman and Hall/CRC

Wynn J, Padget O, Mouritsen H, et al (2020) Natal imprinting to the Earth's magnetic field in a
pelagic seabird. Current Biology 30:2869–2873

Ydenberg RC (1989) Growth-mortality trade-offs and the evolution of juvenile life histories in
the alcidae. Ecology 70:1494–1506

Yoda K, Kohno H, Naito Y (2004) Development of flight performance in the brown booby.
Proceedings of the Royal Society of London Series B: Biological Sciences 271:S240–
S242

Zuur AF (2009a) Mixed effects models and extensions in ecology with R, NY: Springer.
Springer, New York

Supplementary

Species biological aspects

Though the diet and foraging strategy of Amsterdam albatross remains poorly known, it is presumed to

have very similar foraging behaviour compared to that of the wandering albatross, although subtle

differences can appear (Pajot et al. 2021). Like other large albatross species (*Diomedea spp.*), the

Amsterdam albatross is likely to prey on large squid, fish and carrion found on the sea surface (Delord et

al. 2013, Cherel et al. unpublished data). The wandering albatross is known to forage over extensive

distances, detecting prey visually or by olfaction during the day (Nevitt et al. 2008). This strategy

referred as *'foraging-in-flight'* is the lowest energy consuming feeding strategy for the wandering

albatross (Weimerskirch et al. 1997b). However, this strategy tends to change depending on breeding

stage (Phalan et al. 2007; Louzao et al. 2014) leading to a more important utilization of the *'sit-and-wait'*

technique and possibly to vary depending on sites suggesting considerable behavioural plasticity (Phalan

et al. 2007). This switch in foraging techniques could result in more frequent and shorter bouts on the

water in the former technique (compared to *'foraging-in-flight'*).

Thiebot et al. (2014) showed that adult Amsterdam albatrosses during their post-breeding sabbatical

period moved widely (31° to 115° E), mostly exhibiting westwards wider-scale migratory movements

(*sensu* Weimerskirch et al. 2015a) reaching >4000 km from the colony exploiting continuously warm

waters (~18°C). No clear longitudinal seasonality existed in the movements of adults, nonetheless they

tended to move westwards in June/July and eastwards in November. The immature birds moved widely

in longitude (0° to 135° E), exploiting exclusively warm waters 17°-18° C. Similarly to adults no clear

longitudinal seasonality synchronicity existed in the movements, except that they also tended to move

westwards in June and eastwards in November. Juveniles exhibited very large post-fledging movement

capacities over the southern Indian Ocean after fledging (15° to 135° E, ~ 4500 km from the colony),

through a large range of latitudinal gradient (27° to 47° S). Juveniles birds tended to move westwards

first in March-April and temporarily exhibited synchronous individual movements. De Grissac et al.

(2016) compared trajectories (i.e. departure direction or orientation toward specific areas) of juveniles

and adults and showed that juveniles performed an initial rapid movement taking all individuals away

from the vicinity of their native colony, and in a second time performed large-scale movements similar

to those of adults during the sabbatical period. High individual variability and no clear differences

between juveniles and adults patterns were found, except that adults foraged at significantly higher

latitudes. De Grissac et al. (2016) concluded in an overlap in distribution between adults and juveniles

due to the extensive area they used and their differences in latitudinal distribution compared to other

Procellariiformes species.

Moult is an intrinsically costly process requiring time, energy and nutrients (Langston and Rohwer 1996;

Ellis and Gabrielsen 2002), and the annual replacement of flight feathers is crucial to ensure efficiency in

both flight and thermoregulation (Murphy 1996; Peery et al. 2008; Gutowsky et al. 2014). In large-sized

albatrosses like Amsterdam albatross, replacement of primary feathers lasts for more than one breeding

season, and the moult of primaries never occurs during the breeding season (Furness 1988;

Weimerskirch 1991). Stage-specific and sex-specific differences in moult extent occur in wandering

albatross, suggesting important constraints that could compete with breeding (immature birds tend to

renew fewer feathers compared to adult breeders), and particularly in females (Weimerskirch 1991). In

smaller sized seabirds, a link between moulting pattern and activity parameters was evidenced, resulting

in a clear temporal pattern partly explained by moult (Cherel et al. 2016). Recently Gutowsky et al.

(2014) suggested that tropical albatrosses (i.e. Laysan *Phoebastria immutabilis* and black-footed *P.*

nigripes albatrosses) could compromise flight from active wing moult during the nonbreeding period

and induce changes in daily activity budget during a 'quasi-flightless' stage. However, there is no such

data for southern albatrosses. Furthermore for large sized species (*Diomedea spp.*) the activity data

recorded using GLS never suggested it such a compromise. However, adult birds during the non-

breeding season appear to spend much more time on the water during winter, suggesting that partial

moult may occur at this time, as observed in many other seabird species that have to moult during the

non-breeding season and show reduced activity during specific periods that may correspond to moulting

(Weimerskirch et al. 2015b, 2020).

Statistical analyses

Variation in activity parameters between stages with time-lag

The visual comparison shown on Figure S11 was statistically tested using generalized additive mixed

models (GAMMs, built with the 'gam' function, itsadug and mgcv package, (Lin and Zhang 1999; Wood

2015)) with the values associated with the first axe of the PCA as the dependent variable. We ran model

testing for variability in activity parameters for all stages combined (PC1Slag; Table S4). We applied time

lag as illustrated in Figure S11, the first axe was modelled as a function of months spent since departure

from the colony (monthelap.lag) with a delay of 16 months.

Variation in body size

Differences between sexes in body size measurements were tested using Student's t-tests and Wilcoxon

rank tests. We tested independently if each measurement (wing length, tarsus length, bill length, bill

depth and body mass) varied according to sex and stage (juvenile and adult). The effects were tested

using generalised linear models (GLMs) with a Gaussian family and identity link function (Zuur 2009b).

Model validation and model selection were performed following (Zuur 2009b). GLMs tested for effect of

sex and stage and T-tests tested the differences of body size measurements between males and

females. Although sexes and stages differed for some body size measurements, we could not include

body size as an additional explanatory variable in GAMMs testing for factors of variation in activity

patterns due to small sample sizes in each sex and stage category.

Male Amsterdam albatrosses were larger than females, particularly for tarsus length and bill length and

bill depth whatever the stage (juvenile or adult; Tables S5). In juveniles, males were ~13% heavier than

females, while the difference was not significant in adults (Table S5). The most sexually dimorphic

phenotypic traits were body mass, bill depth and tarsus length in juveniles while in adults they were

body mass, tarsus length and bill length.

Table S1 Selected models testing for the effects of sex, stage, number of months spent since departure (monthelap: duration elapsed since

fledging expressed in month, i.e. the first month after fledging and so on) and month of the year (i.e. January and so on) on activity parameters

of Amsterdam albatrosses

	Model #	Study variable[1]	Model structure	Sample size
All stages	gamm1	PC1S	~s(monthelap, by=stage, k = 2) + monthf + stade + sex + s(monthelap, device_code[2], bs = "re")	8094
All stages	gamm2	PC2S	~ s(monthelap, k = 3) + monthf + stade + s(monthelap, device_code, bs = "re")	8094
All stages	gamm3	PC3S	~monthf+stade+sex+s(monthelap,device_code, bs='re')	8094
Juveniles	gamm4	PC1J	~ s(monthelap,k=2)+monthf+s(monthelap,device_code, bs='re')	6161
Juveniles	gamm5	PC2J	~ s(monthelap, k = 2)+monthf+sex+s(monthelap, device_code, bs = "re")	6161
Juveniles	gamm6	PC3J	~monthf+sex+s(monthelap,device_code, bs='re')	6161

841   [1] First, second and third principal component issued from principal components analyses considering i) all stages combined (PCS) and ii) only

842   juveniles (PCJ); [2] Individuals

843   Table S2a GAMM results for the first principal components (PC1S; gamm1 see Table S2) of Amsterdam albatross modelled as a function of

844   months spent since departure from the colony (monthelap), month of the year, stage and sex. Reference values are January, adults and females.

Variable	Smoother edf	F-test	p-value
s(monthelap):stadeAdult	5.178	24.22	0.000
s(monthelap):stadeimm	7.437	45.66	0.000
s(monthelap):stadejuv	5.568	22.32	0.000

	Estimate	Std.Error	t-value	p-value
(Intercept)	0.66	0.08	8.34	0.000
February	-0.28	0.08	-3.43	0.000
March	-0.41	0.09	-4.73	0.000
April	-0.60	0.10	-6.01	0.000
May	-0.33	0.10	-3.19	0.001
June	-0.43	0.10	-4.12	0.000
July	-0.19	0.10	-1.88	0.060
August	0.00	0.10	-0.02	0.985
September	0.02	0.09	0.17	0.862
October	-0.10	0.09	-1.10	0.272
November	-0.18	0.09	-2.03	0.043
December	-0.06	0.08	-0.69	0.489
Immatures	-0.63	0.04	-15.70	0.000
Juveniles	-1.21	0.04	-34.00	0.000
Males	0.23	0.03	7.69	0.000

848   Table S2b GAMM results for the second principal components (PC2S; gamm2 see Table S2) of Amsterdam albatross modelled as a function of

849   months spent since departure from the colony (monthelap), month of the year, stage and sex. Reference values are January, adults and females.

Variable	Smoother edf	F-test	p-value
s(monthelap)	1.001	0.504	0.478
s(monthelap,device_code)	27.107	39.991	0.000

	Estimate	Std.Error	t-value	p-value
(Intercept)	-0.15	0.10	-1.53	0.126
February	0.13	0.08	1.71	0.088
March	0.16	0.08	2.00	0.046
April	0.42	0.08	5.03	0.000
May	0.40	0.08	4.99	0.000
June	0.25	0.08	3.16	0.002
July	0.23	0.08	2.92	0.004
August	0.26	0.08	3.40	0.001
September	0.48	0.08	6.22	0.000
October	0.35	0.08	4.57	0.000
November	0.34	0.08	4.41	0.000
December	0.19	0.08	2.49	0.013
Immatures	-0.12	0.08	-1.57	0.116
Juveniles	-0.18	0.06	-2.96	0.003

853 Table S2c GAMM results for the third principal components (PC3S; gamm3 see Table S2) of Amsterdam albatross modelled as a function of

854 months spent since departure from the colony (monthelap), month of the year, stage and sex. Reference values are January, adults and females.

Variable	Smoother edf	F-test	p-value
s(monthelap,device_code)	26.52	16.58	0.000

	Estimate	Std.Error	t-value	p-value
(Intercept)	0.34	0.06	5.37	0.000
February	-0.22	0.06	-3.43	0.000
March	-0.07	0.06	-1.08	0.279
April	-0.10	0.07	-1.53	0.127
May	0.00	0.06	0.05	0.958
June	0.05	0.06	0.87	0.385
July	0.02	0.06	0.39	0.694
August	-0.04	0.06	-0.70	0.483
September	-0.06	0.06	-0.93	0.355
October	-0.10	0.06	-1.58	0.012
November	-0.16	0.06	-2.57	0.010
December	-0.23	0.06	-3.70	0.000
Immatures	-0.27	0.06	-4.61	0.000
Juveniles	-0.45	0.05	-9.12	0.000
Males	-0.14	0.04	-3,39	0.000

858   Table S3a GAMM results for the first principal components (PC1J; gamm4 see Table S2) of juveniles Amsterdam albatross modelled as a function

859   of months spent since departure from the colony (monthelap) and month of the year. Reference value is January.

Variable	Smoother edf	F-test	p-value
s(monthelap)	1.922	24.12	0.000
s(monthelap,device_code)	6.506	12.20	0.000

	Estimate	Std.Error	t-value	p-value
(Intercept)	-0.10	0.09	-1.16	0.247
February	0.20	0.09	2.30	0.021
March	-0.13	0.09	-1.51	0.132
April	-0.27	0.09	-3.12	0.002
May	-0.16	0.09	-1.91	0.056
June	0.06	0.09	0.71	0.475
July	0.59	0.09	6.66	0.000
August	0.68	0.09	7.66	0.000
September	0.47	0.09	5.26	0.000
October	0.22	0.09	2.47	0.013
November	-0.05	0.09	-0.50	0.616
December	-0.04	0.09	-0.45	0.653

863   Table S3b GAMM results for the second principal components (PC2J; gamm5 see Table S2) of juveniles Amsterdam albatross modelled as a

864   function of months spent since departure from the colony (monthelap) and month of the year. Reference value is January.

Variable	Smoother edf	F-test	p-value
s(monthelap)	1.000	2.40	0.122
s(monthelap,device_code)	6.813	55.91	0.000

	Estimate	Std.Error	t-value	p-value
(Intercept)	-0.05	0.12	-0.43	0.668
February	-0.27	0.08	-3.54	0.000
March	-0.14	0.08	-1.88	0.061
April	0.12	0.07	1.61	0.107
May	0.08	0.08	1.02	0.306
June	0.01	0.08	0.14	0.886
July	0.08	0.08	1.05	0.293
August	0.12	0.08	1.47	0.141
September	0.30	0.08	3.84	0.000
October	0.18	0.08	2.33	0.020
November	0.19	0.08	2.44	0.015
December	0.20	0.08	2.49	0.013
Male	-0.17	0.06	-2.78	0.005

868   Table S3c GAMM results for the third principal components (PC3J; gamm5 see Table S2) of juveniles Amsterdam albatross modelled as a

869   function of months spent since departure from the colony (monthelap), month of the year and sex. Reference value are January and females.

Variable	Smoother edf	F-test	p-value
s(monthelap,device_code)	7.724	19.37	0.000

	Estimate	Std.Error	t-value	p-value
(Intercept)	-0.04	0.05	-0.77	0.444
February	-0.10	0.06	-1.59	0.113
March	-0.13	0.06	-2.19	0.029
April	-0.06	0.06	-0.94	0.349
May	0.02	0.06	0.37	0.715
June	0.13	0.06	2.11	0.035
July	0.02	0.06	0.25	0.802
August	0.00	0.06	0.01	0.992
September	0.00	0.06	0.01	0.996
October	-0.04	0.06	-0.59	0.556
November	-0.06	0.07	-0.85	0.395
December	0.02	0.06	0.34	0.735
Male	0.10	0.05	1.89	0.059

873   Table S4 GAMM results for the first principal components (PC1Slag) of Amsterdam albatross modelled as a function of months spent since

874   departure from the colony (monthelap.lag) with a delay of 16 months (see Figure S11), month of the year, stage and sex. Reference values are

875   January, adults and females.

Variable	Smoother edf	F-test	p-value
s(monthelap.lag):stadeAdult	5.001	49.37	0.000
s(monthelap.lag):stadeimm	4.810	19.39	0.000
s(monthelap.lag):stadejuv	7.643	53.53	0.000

	Estimate	Std.Error	t-value	p-value
(Intercept)	0.99	0.08	11.95	0.000
February	-0.53	0.09	-5.79	0.000
March	-1.08	0.09	-11.74	0.000
April	-1.48	0.09	-15.75	0.000
May	-1.23	0.09	-13.59	0.001
June	-1.03	0.09	-11.82	0.000
July	-0.42	0.08	-4.92	0.060
August	-0.07	0.08	-0.90	0.985
September	0.02	0.08	0.28	0.862
October	-0.05	0.08	-0.70	0.272
November	-0.15	0.08	-1.96	0.043
December	0.05	0.08	0.67	0.489
Immatures	-0.81	0.06	-14.24	0.000
Juveniles	-0.74	0.05	-14.71	0.000
Males	0.20	0.03	6.96	0.000

877   Table S5 Body measurements of juveniles and adults Amsterdam albatross and percentage of differences between sexes for each measurement.

878   Δ is the difference in %, p values are reported

	Juvenile		Adult		Δ		Dimorphism ratio[1]		t-test[2]	
	Female (n=159)	Male (n=162)	Female (n=30)	Male (n=45)	Juvenile	Adult	Juvenile	Adult	Juvenile	Adult
Wing length (mm)	532.3 ± 62.1	543.6 ± 72.5	637.0 ± 13.3	654.7 ± 14.7	2.1	2.7	1.021	1.026	**W=10554** **P<0.01**	**t_{64}=-4.882** **p<0.001**
Tarsus length (mm)	113.1 ± 3.8	118.4 ± 3.8	112.4 ± 3.9	116.4 ± 4.0	4.5	3.5	1.047	1.077	**t_{229}= -10.54** **p<0.001**	**t_{28}= -3.982** **p<0.001**
Bill length (mm)	137.1 ± 4.0	143.1 ± 3.8	140.9 ± 4.7	145.0 ± 3.5	4.2	2.8	1.044	1.034	**t_{319}= -13.89** **p<0.001**	**t_{72}= -4.268** **p<0.001**
Bill depth (mm)	34.8 ± 3.4	36.9 ± 3.4	36.0 ± 1.5	38.0 ± 1.5	5.8	5.2	1.061	1.000	**t_{318}= -5.629** **p<0.001**	**t_{58}= -4.821** **p<0.001**
Body mass (g)	7719 ± 1228	8859 ± 1546	7509 ± 1561	7257 ± 1207	12.9	3.4	1.148	1.077	**t_{193}=-5.846** **p<0.001**	t_{32}=0.512 p=0.612

879   [1] ratio calculated as Male/Female following Lovich and Gibbons (1992) for species for which males are known to be larger than females cited in Smith (1999); [2]
880   Student's t-tests used after check for normality of the data for all measurements except for wing length of juveniles (Wilcoxon rank test)

 FIGURES

883   Figure S1 Daily proportions of time spent on water depending on stage (juveniles, immatures and adults)
884   for every month since departure from the colony (upper panel) and for each month of the year (lower
885   panel). Error bars represent ± 1 sd

886 Figure S2 Daily flying bouts duration (dry bouts in hours) depending and on sex (females and males) and
887

on stage (juveniles, immatures and adults) for a) time elapsed since departure from the colony
expressed in month (left panel) and for b) each month of the year (right panel). One side error bars
represent ± 1 sd

Figure S3 Daily flying bouts number (dry bouts) for every month since departure from the colony for
juveniles, immatures and adults for females (upper panel) and males (lower panel). Error bars represent
± 1 sd

Figure S4 Daily wet bouts duration (bouts on water in hours) depending on stage (juveniles, immatures
and adults) and on sex (females and males) for every month since departure of the colony (upper panel)
and for each month of the year (lower panel). Error bars represent ± 1 sd

Figure S5 Daily wet bouts number (bouts on water) for every month since departure from the colony for

juveniles, immatures and adults for females (upper panel) and males (lower panel). Error bars represent

± 1 sd

Figure S6 Modeled first (left panel) and second (right panel) axis of principal components analysis of activity parameters of juveniles of
Amsterdam albatrosses according to time elapsed (e.g. duration elapsed since departure from the colony expressed in month). Models outputs

907   obtained using random intercepts and slopes (each coloured line representing an individual). Line corresponds to estimated smoother from the

908   GAMM models

911 Figure S7 Daily flying bouts duration (dry bouts in hours) for every month since departure of the colony
912 for juveniles, immatures and adults for females (upper panel) and males (lower panel). Error bars
913 represent ±1 sd

Figure S8 Daily flying bouts number (dry bouts) for every month since departure of the colony for juveniles, immatures and adults for females (upper panel) and males (lower panel). Error bars represent ± 1 sd

915
916
917

Figure S9 Daily wet bouts duration (bouts on water in hours) for every month since departure of the

colony for juveniles, immatures and adults for females (upper panel) and males (lower panel). Error bars

represent ±1 sd

Figure S10 Daily wet bouts number (bouts on water) for every month since departure of the colony for
juveniles, immatures and adults for females (upper panel) and males (lower panel). Error bars represent
± 1 sd

Figure S11 Daily proportions of time spent on water for every month since departure of the colony for juveniles-during the first 28 months spent at sea (after departure), immatures and adults (upper panel) and with a 15-16 months of delay for immatures and adults compared to juveniles (lower panel). Error bars represent $\pm$ 1 sd

REFERENCES

Alerstam T, Lindström Å (1990) Optimal bird migration: the relative importance of time, energy, and
safety. In: Bird migration. Springer, pp 331–351

Austin RE, De Pascalis F, Arnould JP, et al (2019) A sex-influenced flexible foraging strategy in a tropical
seabird, the magnificent frigatebird. Marine Ecology Progress Series 611:203–214

Barbraud C, Delord K, Kato A, et al (2019) Sexual segregation in a highly pagophilic and sexually
dimorphic marine predator. bioRxiv 472431

Becker J (2014) Dispersal and Genetic Variation in a Naturally Fragmented Environment: Insights from a
Long-term Study of White-throated Dippers (Cinclus cinclus)

Boynton CK, Mahony NA, Williams TD (2020) Barn Swallow (Hirundo rustica) fledglings use crop habitat
more frequently in relation to its availability than pasture and other habitat types. The Condor
122:duz067

Bruner JS (1972) Nature and uses of immaturity. American psychologist 27:687

Burnham KP, Anderson DR (2002) Model Selection and Multimodel Inference: A Practical Information-
Theoretic Approach. NewYork,NY.

Burns JM, Clark CA, Richmond JP (2004) The impact of lactation strategy on physiological development
of juvenile marine mammals: implications for the transition to independent foraging. Elsevier,
pp 341–350

Campioni L, Dias MP, Granadeiro JP, Catry P (2020) An ontogenetic perspective on migratory strategy of
a long-lived pelagic seabird: Timings and destinations change progressively during maturation.
Journal of Animal Ecology 89:29–43

Campioni L, Granadeiro JP, Catry P (2016) Niche segregation between immature and adult seabirds:
does progressive maturation play a role? Behavioral Ecology 27:426–433

Caubet Y, Jaisson P, Lenoir A (1992) Preimaginal induction of adult behaviour in insects. The Quarterly
Journal of Experimental Psychology Section B 44:165–178

Cherel Y, Quillfeldt P, Delord K, Weimerskirch H (2016) Combination of at-sea activity, geolocation and
feather stable isotopes documents where and when seabirds molt. Frontiers in Ecology and the
Environment 4:1–16

Clay TA, Joo R, Weimerskirch H, et al (2020) Sex-specific effects of wind on the flight decisions of a
sexually dimorphic soaring bird. Journal of Animal Ecology 89:1811–1823

Collet J, Prudor A, Corbeau A, et al (2020) First explorations: ontogeny of central place foraging
directions in two tropical seabirds. Behavioral Ecology 31:815–825

Corbeau A, Prudor A, Kato A, Weimerskirch H (2020a) Development of flight and foraging behaviour in a
juvenile seabird with extreme soaring capacities. Journal of Animal Ecology 89:20–28

Corbeau A, Prudor A, Kato A, Weimerskirch H (2020b) Development of flight and foraging behaviour in a
juvenile seabird with extreme soaring capacities. Journal of Animal Ecology 89:20–28.
https://doi.org/10.1111/1365-2656.13121

Daunt F, Afanasyev V, Adam A, et al (2007) From cradle to early grave: juvenile mortality in European
shags Phalacrocorax aristotelis results from inadequate development of foraging proficiency.
Biology Letters 3:371–374. https://doi.org/10.1098/rsbl.2007.0157

de Grissac S, Bartumeus F, Cox SL, Weimerskirch H (2017) Early-life foraging: Behavioral responses of
newly fledged albatrosses to environmental conditions. Ecology and evolution 7:6766–6778

de Grissac S, Börger L, Guitteaud A, Weimerskirch H (2016) Contrasting movement strategies among
juvenile albatrosses and petrels. Scientific reports 6:26103

Delord K, Barbraud C, Bost CA, et al (2013) Atlas of top predators from the French Southern Territories in
the Southern Indian Ocean. http://www.cebc.cnrs.fr/ecomm/Fr_ecomm/ecomm_ecor_OI1.html

Dukas R (2006) Learning in the context of sexual behaviour in insects. Animal Biology 56:125–141

Ellis HI, Gabrielsen GW (2002) Energetics of free-ranging seabirds. In: Seabirds energetics, Whittow, G. C.
and Rahn, H. Plenum Press, N.Y., pp 359–408

Evans DR (2018) The post-fledging survival and movements of juvenile Barn Swallows (Hirundo rustica):
an automated telemetry approach

Fay R, Weimerskirch H, Delord K, Barbraud C (2015) Population density and climate shape early-life
survival and recruitment in a long-lived pelagic seabird. Journal of Animal Ecology 84:1423–1433

Frankish CK, Manica A, Clay TA, et al (2022) Ontogeny of movement patterns and habitat selection in
juvenile albatrosses. Oikos e09057

Frankish CK, Manica A, Phillips RA (2020a) Effects of age on foraging behavior in two closely related
albatross species. Movement ecology 8:7

Frankish CK, Phillips RA, Clay TA, et al (2020b) Environmental drivers of movement in a threatened
seabird: insights from a mechanistic model and implications for conservation. Diversity and
Distributions 26:1315–1329

Furness RW (1988) Influences of status and recent breeding experience on the moult strategy of the
Yellow-nosed albatross Diomedea chlororhynchos. J Zool 215:719–727

Gonzales-Solis J, Croxall JP, Wood AG (2000) Sexual dimorphism and sexual segregation in foraging
strategies of northern giant petrels Macronectes halli during incubation. Oikos 90:390–398

Gutowsky SE, Gutowsky LF, Jonsen ID, et al (2014) Daily activity budgets reveal a quasi-flightless stage
during non-breeding in Hawaiian albatrosses. Movement ecology 2:23–23

Harel R, Horvitz N, Nathan R (2016a) Adult vultures outperform juveniles in challenging thermal soaring
conditions. Scientific reports 6:27865

Harel R, Horvitz N, Nathan R (2016b) Adult vultures outperform juveniles in challenging thermal soaring
conditions. Scientific reports 6:27865

Heyes CM (1994) Social learning in animals: categories and mechanisms. Biological Reviews 69:207–231

Jaeger A, Jaquemet S, Phillips RA, et al (2013) Stable isotopes document inter- and intra-specific
variation in feeding ecology of nine large southern Procellariiformes. Mar Ecol Prog Ser
490:255–266

Jouventin P, Martinez J, Roux J-P (1989) Breeding biology and current status of the Amsterdam Island
Albatross *Diomedea amsterdamensis*. Ibis 131:182-

King JR (1974) Seasonal allocation of time and energy resources in birds. Avian energetics 4–85

Krüger S, Reid T, Amar A (2014) Differential range use between age classes of southern African Bearded
Vultures Gypaetus barbatus. PLoS One 9:

Lack D (1954) The natural regulation of animal numbers. The Natural Regulation of Animal Numbers

Langen TA (1996) Skill acquisition and the timing of natal dispersal in the white-throated magpie-jay,
Calocitta formosa. Animal Behaviour 51:575–588

Langston NE, Rohwer S (1996) Molt-breeding tradeoffs in albatrosses⊠: life history implications for big
birds. Oikos 76:498–510

Lê S, Josse J, Husson F (2008) FactoMineR: an R package for multivariate analysis. Journal of statistical
software 25:1–18

Lin X, Zhang D (1999) Inference in generalized additive mixed modelsby using smoothing splines. Journal
of the royal statistical society: Series b (statistical methodology) 61:381–400

Louzao M, Wiegand T, Bartumeus F, Weimerskirch H (2014) Coupling instantaneous energy-budget
models and behavioural mode analysis to estimate optimal foraging strategy: an example with
wandering albatrosses. Movement ecology 2:1–15

Lovich JE, Gibbons JW (1992) A review of techniques for quantifying sexual size dimorphism. Growth
Development and Aging 56:269–269

Marchetti K, Price T (1989) Differences in the foraging of juvenile and adult birds: the importance of
developmental constraints. Biological Reviews 64:51–70

Margrath R, Lill A (1985) Age-related differences in behaviour and ecology of crimson rosellas,
Platycercus elegans, during the non-breeding season. Wildlife Research 12:299–306

Martin P, Bateson P (1985) The ontogeny of locomotor play behaviour in the domestic cat. Animal
behaviour 33:502–510

Mendez L, Prudor A, Weimerskirch H (2017) Ontogeny of foraging behaviour in juvenile red-footed
boobies (Sula sula). Scientific reports 7:1–11

Murphy ME (1996) Energetics and nutrition of molt. In: Avian energetics and nutritional ecology.
Springer, pp 158–198

Mushinsky HR, Hebrard JJ, Vodopich DS (1982) Ontogeny of water snake foraging ecology. Ecology
63:1624–1629

Naef-Daenzer B, Grüebler MU (2008) Post-fledging range use of Great Tit Parus major families in relation
to chick body condition. Ardea 96:181–190

Nevitt GA, Losekoot M, Weimerskirch H (2008) Evidence for olfactory search in wandering albatross,
Diomedea exulans. Proceedings of the National Academy of Sciences 105:4576–4581

Nourani E, Vansteelant WMG, Byholm P, Safi K (2020) Dynamics of the energy seascape can explain
intra-specific variations in sea-crossing behaviour of soaring birds. Biology Letters 16:20190797.
https://doi.org/10.1098/rsbl.2019.0797

Pajot A, Corbeau A, Jambon A, Weimerskirch H (2021) Diel at-sea activity of two species of great
albatrosses: the ontogeny of foraging and movement behaviour. Journal of Avian Biology 52:

Payo-Payo A, Genovart M, Bertolero A, et al (2016) Consecutive cohort effects driven by density-
dependence and climate influence early-life survival in a long-lived bird. Proceedings of the
Royal Society B: Biological Sciences 283:20153042

Peery MZ, Henkel LA, Newman SH, et al (2008) Effects of rapid flight-feather molt on postbreeding
dispersal in a pursuit-diving seabird. The Auk 125:113–123

Pennycuick CJ (1982) The flight of petrels and albatrosses (Procellariiformes), observed in South Georgia
and its vicinity. Phil Trans R Soc Lond 300:75–106

Peron C, Gremillet D (2013) Tracking through Life Stages: Adult, Immature and Juvenile Autumn
Migration in a Long-Lived Seabird. Plos One 8:

Phalan B, Phillips RA, Silk JR, et al (2007) Foraging behaviour of four albatross species by night and day.
Marine Ecology Progress Series 340:271–286

Phillips RA, Lewis S, González-Solís J, Daunt F (2017) Causes and consequences of individual variability
and specialization in foraging and migration strategies of seabirds. Marine Ecology Progress
Series 578:117–150

Phillips RA, Mcgill RAR, Dawson DA, Bearhop S (2011) Sexual segregation in distribution, diet and trophic
level of seabirds: insights from stable isotope analysis. Marine Biology 158:2199–2208

Phillips RA, Silk JRD, Croxall JP, et al (2004a) Accuracy of geolocation estimates for flying seabirds.
Marine Ecology-Progress Series 266:265–272

Phillips RA, Silk JRD, Phalan B, et al (2004b) Seasonal sexual segregation in two *Thalassarche* albatross
species: competitive exclusion, reproductive role specialization or foraging niche divergence?
Proceedings of the Royal Society of LondonSeries B: Biological Sciences 271:1283–1291

Rapaport LG, Brown GR (2008) Social influences on foraging behavior in young nonhuman primates:
learning what, where, and how to eat. Evolutionary Anthropology: Issues, News, and Reviews:
Issues, News, and Reviews 17:189–201

Riotte-Lambert L, Weimerskirch H (2013) Do naive juvenile seabirds forage differently from adults?
Proceedings of the Royal Society B: Biological Sciences 280:20131434

Rivalan P, Barbraud C, Inchausti P, Weimerskirch H (2010) Combined impact of longline fisheries and
climate on the persistence of the Amsterdam albatross. Ibis 152:6–18

Rotics S, Kaatz M, Resheff YS, et al (2016a) The challenges of the first migration: movement and
behaviour of juvenile vs. adult white storks with insights regarding juvenile mortality. Journal of
Animal Ecology 85:938–947

Rotics S, Kaatz M, Resheff YS, et al (2016b) The challenges of the first migration: movement and
behaviour of juvenile vs. adult white storks with insights regarding juvenile mortality. Journal of
Animal Ecology 85:938–947

Rotics S, Turjeman S, Kaatz M, et al (2021) Early-life behaviour predicts first-year survival in a long-
distance avian migrant. Proceedings of the Royal Society B 288:20202670

Roux JP, Jouventin P, Mougin JL, et al (1983) Un nouvel albatros Diomedea amsterdamensis n. sp.
découvert sur l'île Amsterdam (37°50'S, 77°35'E). L'Oiseau et R F O 53:1–11

Ruaux G, Lumineau S, de Margerie E (2020) The development of flight behaviours in birds. Proceedings
of the Royal Society B 287:20200668

Sergio F, Tanferna A, De Stephanis R, et al (2014) Individual improvements and selective mortality shape
lifelong migratory performance. Nature 515:410–413

Shaffer SA, Weimerskirch H, Costa D (2001) Functional significance of sexual dimorphism in wandering
albatrosses, Diomedea exulans. Functional Ecology 15:203–210

Smith RJ (1999) Statistics of sexual size dimorphism. Journal of Human Evolution 36:423–458.
https://doi.org/10.1006/jhev.1998.0281

Team RC (2019) R: A Language and Environment for Statistical Computing (Version 3.5. 2, R Foundation
for Statistical Computing, Vienna, Austria, 2018). There is no corresponding record for this
reference[Google Scholar]

Thiebot J-B, Delord K, Marteau C, Weimerskirch H (2014) Stage-dependent distribution of the critically
1104   endangered Amsterdam albatross in relation to Economic Exclusive Zones. Endangered Species
1105   Research 23:263–276

1106   Thiers L, Delord K, Barbraud C, et al (2014) Foraging zones of the two sibling species of giant petrels in
1107   the Indian Ocean throughout the annual cycle: implication for their conservation. Mar Ecol Prog
1108   Ser 499:233–248

1109   Thorup K, Alerstam T, Hake M, Kjellén N (2003) Bird orientation: compensation for wind drift in
1110   migrating raptors is age dependent. Proceedings of the Royal Society of London Series B:
1111   Biological Sciences 270:S8–S11

1112   Urios V, López-López P, Limiñana R, Godino A (2010) Ranging behaviour of a juvenile Bearded Vulture
1113   (Gypaetus barbatus meridionalis) in South Africa revealed by GPS satellite telemetry

1114   Votier SC, Grecian WJ, Patrick S, Newton J (2011) Inter-colony movements, at-sea behaviour and
1115   foraging in an immature seabird: results from GPS-PPT tracking, radio-tracking and stable
1116   isotope analysis. Marine biology 158:355–362

1117   Wearmouth VJ, Sims DW (2008) Sexual segregation in marine fish, reptiles, birds and mammals:
1118   behaviour patterns, mechanisms and conservation implications. Advances in marine biology
1119   54:107–170

1120   Weimerskirch H (1991) Sex-specific differences in molt strategy in relation to breeding in the wandering
1121   albatross. Condor 93:731–737

1122   Weimerskirch H, Akesson S, Pinaud D (2006) Postnatal dispersal of wandering albatrosses *Diomedea*
1123   *exulans*: implications for the conservation of the species. J Avian Biol 37:23–28

1124   Weimerskirch H, Brothers N, Jouventin P (1997a) Population dynamics of Wandering Albatross
1125   *Diomedea exulans* and Amsterdam Albatross *D. amsterdamensis* in the Indian Ocean and their
1126   relationship with long-line fisheries: conservation implications. Biological conservation 79:257–
1127   270

1128   Weimerskirch H, Cherel Y, Delord K, et al (2014) Lifetime foraging patterns of the wandering albatross:
1129   Life on the move! Journal of Experimental Marine Biology and Ecology 450:68–78

1130   Weimerskirch H, de Grissac S, Ravache A, et al (2020) At-sea movements of wedge-tailed shearwaters
1131   during and outside the breeding season from four colonies in New Caledonia. Marine Ecology
1132   Progress Series 633:225–238

1133   Weimerskirch H, Delord K, Guitteaud A, et al (2015a) Extreme variation in migration strategies between
1134   and within wandering albatross populations during their sabbatical year, and their fitness
1135   consequences. Scientific Reports 5:8853-

1136   Weimerskirch H, Guionnet T (2002) Comparative activity pattern during foraging of four albatross
1137   species. Ibis 144:40–50

Weimerskirch H, Shaffer SA, Tremblay Y, et al (2009) Species-and sex-specific differences in foraging
behaviour and foraging zones in blue-footed and brown boobies in the Gulf of California. Marine
Ecology Progress Series 391:267–278

Weimerskirch H, Tarroux A, Chastel O, et al (2015b) Population-specific wintering distributions of adult
south polar skuas over the three oceans. Mar Ecol Prog Ser 538:229–237

Weimerskirch H, Wilson RP, Lys P (1997b) Activity pattern of foraging in the wandering albatross: a
marine predator with two modes of prey searching. Marine Ecology Progress Series 151:245–
254

Wilson RP, Ducamp JJ, Rees G, et al (1992) Estimation of location: global coverage using light intensity.
In: Priede IMSS (ed). Ellis Horward, Chichester, pp 131–134

Wood S (2015) Package 'mgcv.' R package version 1:729

Wood S (2017) Generalized Additive Models: An Introduction with R, 2nd Edition Chapman and
Hall//CRC, Boca Raton. Chapman and Hall/CRC

Wynn J, Padget O, Mouritsen H, et al (2020) Natal imprinting to the Earth's magnetic field in a pelagic
seabird. Current Biology 30:2869–2873

Ydenberg RC (1989) Growth-mortality trade-offs and the evolution of juvenile life histories in the
alcidae. Ecology 70:1494–1506

Yoda K, Kohno H, Naito Y (2004) Development of flight performance in the brown booby. Proceedings of
the Royal Society of London Series B: Biological Sciences 271:S240–S242

Zuur AF (2009a) Mixed effects models and extensions in ecology with R, NY: Springer. Springer, New
York

Zuur AF (ed) (2009b) Mixed effects models and extensions in ecology with R. Springer, New York, NY

1160

1161

PCIEcology

Decision concerning your submission

───────────────────────────────

Dear Karine Delord,

Your article, entitled **The challenges of independence: ontogeny of at-sea behaviour in a long-**
lived seabird, has now been reviewed.

The referees' comments and the recommender's decision are shown below. As you can see, the
recommender found your article very interesting but suggests certain revisions.

We shall, in principle, be happy to recommend your article as soon as it has been revised in
response to the points raised by the referees.

When revising your article, we remind you that your article must contain the following sections
(see our Guide for Authors in the Help section of the PCIEcology website):

1) **Data, script and code availability (if applicable)**

• **Data, statistical scripts, command lines and simulation code must be made available**
to readers. They should either be included in the article or deposited **in** an open
repository such as Zenodo **with a DOI**. A perennial URL can be provided if no DOI is
available; please note that GitHub URL are not perennial.
• **If deposited in** an open repository, a reference to **Data, statistical scripts, command**
lines and simulation code, with a DOI or a perennial URL, must be provided in the
reference list and in the "Data, script and code availability" section
• The "Data, script and code availability" section must clearly indicate **where and**
how data can be accessed.
• Wherever possible, data, scripts and code should be provided in machine-readable
formats. Avoid PDFs other than for textual supplementary information.
• Metadata should accompany the data, to make the data understandable and reusable
by the reader.

2) **Supplementary information (if applicable)**

• Supplementary information (text, tables, figures, videos, etc.) can be referred to in the
article. It must be available in an open repository (such as Zenodo, Dryad, OSF, Figshare,
Morphobank, Morphosource, Github, MorphoMuseuM, Phenome10k, etc. or any
institutional repository, etc...) with a DOI. A perennial URL can be provided if no DOI is
available.
• A reference to the supplementary information, with a DOI or a perennial URL, must be
provided in the reference list and in the "Supplementary information" section.
• List all documents attached to the manuscript as Supplementary Information in the
"Supplementary Information" section.

3) **Funding (mandatory)**

• All sources of funding must be listed in a separate "Funding section". The absence of
funding must be clearly indicated in this section.

4) **Conflict of interest disclosure (mandatory)**

• Authors should declare any potential non-financial conflict of interest (financial conflicts
of interest are forbidden, see the PCI code of conduct).
• In the absence of competing interests, the authors should add the following sentence to
the "Conflict of interest disclosure" section: "The authors declare they have no conflict
of interest relating to the content of this article." If appropriate, this disclosure may be
completed by a sentence indicating that some of the authors are PCI recommenders:
"XXX is a recommender for PCI XX."

5) **Materials and methods (mandatory)**

• Details of experimental procedures and quantitative analyses must be made **fully
available** to readers, in the text, as appendices, or as Supplementary Information
deposited in an open repository, such as Zenodo, Dryad or institutional repositories with
a DOI.
• For specimen-based studies, **complete repository information** should be provided and
institutional abbreviations should be listed in a dedicated subsection (if applicable).
Specimens on which conclusions are based **must be deposited in an accessible and
permanent repository**.

When your revised article is ready, please:

1) Upload the new version of your manuscript onto your favorite open archive and **wait until it
appears online**;

2) Follow this link https://ecology.peercommunityin.org/user/my_articles or logging onto the
PCIEcology website and go to 'For Contributors -> Your submitted preprints' in the top
menu and **click on the blue 'VIEW/EDIT' button at the right end of the line** referring to the
preprint in question.

3) Click on the black 'EDIT YOUR ARTICLE DATA' button (mandatory step). You can then edit the
title, authors, DOI, abstract, keywords, disciplines, and DOI/URL of data, scripts and code. Do
not forget to save your modifications by clicking on the green button.

4) Click on the blue 'EDIT YOUR REPLY TO THE RECOMMENDER' button (mandatory step). You
could then write or paste your text, upload your reply as a PDF file, and upload a document
with the modifications marked in TrackChange mode. If you are submitting the final formatted
version ready to be recommended, you should only add a sentence indicating that you posted
the final version on the preprint server. Do not forget to **save your modifications by clicking on**
the green button.

5) Click on the green 'SEND RESUBMISSION' button. This will result in your submission being
sent to the recommender.

Once the recommender has read the revised version, they may decide to recommend it
directly, in which case the editorial correspondence (reviews, recommender's decisions,
authors' replies) and a recommendation text will be published by PCIEcology under the license
CC-BY.

Alternatively, other rounds of reviews may be needed before the recommender reaches a
favorable conclusion. They may also reject your article, in which case the reviews and decision
will be sent to you, but they will not be published or publicly released by PCIEcology. They will
be safely stored in our database, to which only the Managing Board has access. You will be
notified by e-mail at each stage in the procedure.

We thank you in advance for submitting your revised version.

Yours sincerely,

The Managing Board of PCIEcology

1246

The challenges of independence: ontogeny of at-sea behaviour in a long-lived seabird

Karine Delord, Henri Weimerskirch, Christophe Barbraud

Abstract

The transition to independent foraging represents an important developmental stage in the life
cycle of most vertebrate animals. Juveniles differ from adults in various life history traits and
tend to survive less well than adults in most long-lived animals. Several hypotheses have been
proposed to explain higher mortality including that of inadequate/inferior foraging skills
compared to adults, young naïve individuals combining lack of experience and physical
immaturity. Thus a change in behaviour, resulting in an improvement of skills acquired from
growing experience, is expected to occur during a period of learning through the immaturity
phase. Very few studies have investigated the ontogeny of foraging behaviour over long periods
of time, particularly in long-lived pelagic seabirds, due to the difficulty of obtaining individual
tracking data over several years. We investigated the foraging behaviour, through activity
patterns, during the three life stages of the endangered Amsterdam albatross by using
miniaturized activity loggers on naïve juveniles, immatures and adults. Naïve juveniles during
their first month at sea after leaving their colony exhibited lower foraging effort (greater
proportion of time spent sitting on water, longer and more numerous bouts on water, shorter
and fewer flying bouts). Patterns of activity parameters in juveniles after independence
suggested a progressive change of foraging performances during the first two months after
fledging. We found sex differences in activity parameters according to time since departure
from the colony and month of the year, consistent with the important sexual dimorphism in the
Amsterdam albatross. Regardless of life stage considered, activity parameters exhibited
temporal variability reflecting the modulation of foraging behaviour. This variability is discussed
in light of both extrinsic (i.e. environmental conditions such as variability in food resources or in
wind) and intrinsic (i.e. energetic demands linked to plumage renew during moult) factors.

Keywords: activity data loggers; foraging behaviour; southern Indian Ocean; Amsterdam
albatross; Diomedea amsterdamensis

Round #4

───────────────────────────────

by Blandine Doligez, 02 Oct 2023 15:20
Manuscript: **https://doi.org/10.1101/2021.10.23.465439** version 5

Very minor revision

1281

1282 Dear authors,

1283

Thank you for carefully editing the last version with the corrections suggested. I have still
spotted a few additional ones that would need to be dealt with before final recommendation is
made (as no further changes can be done later on). This should however be done very quickly...
Sorry for the additional delay but this time it should be the last!

R: thank you for your careful checking of the manuscript. All the corrections suggest have been
addressed in the revised version

Here are the edits needed:

1291

l. 114-119: the separators need adjustment: the sentence should read as:

These included temporal (i.e. related to the month of the year) changes in activity parameters
*for all life-stages due **(i) to** environmental changes occurring throughout the seasons, **(ii) to***
partial moulting which is suspected to occur outside the breeding period and to result in reduced
activity for adults and immatures (i.e. more time 115 spent on the water; Weimerskirch et al.
*2015, 2020), or **(iii) to** sex differences in flight performances (Shaffer 116 et al. 2001; Riotte-*
Lambert and Weimerskirch 2013; Clay et al. 2020).

R: this was changed accordingly

There are still a number of places where the double parentheses should be removed when a
reference appears in a section that is already in parentheses (e.g. l. 66, l. 219). In these cases,
the parentheses around the reference should be dropped. In the same way, l. 879, parentheses
for the Smith reference should be around the year only.

R: following this comment double parentheses were removed throughout the ms in the revised version

Regarding the use of AICc: thanks for checking that results were unchanged with AICc instead of
AIC. Thus I think there is no need to replace AIC by AICc in the ms. However, because readers
may ask themselves the same question, I would simply add a short sentence to mention this
clearly: "Using AICc to account for small sample sizes did not change model selection" (or
something similar, when you mention AIC in methods, l. 238).

R: this was added in the revised version

l. 834: there is no Table S6 (?)

R: this was modified in the revised version

l. 252: I would delete the second "inter-individual variability"

R: this was deleted

l. 258: I would delete the second "variability"

R: this was deleted

l. 438: seems like permit number has been forgotten, but I am not sure whether this is
compulsory (would be better to mention them I think).

R: we do not have a permit number for our projetc, but simply the fact that our demands are approved
by the ethics committee

l. 348: add a "." after "sea"

R: this was added

l. 360: remove the "," after "albatrosses"

R: this was removed

l. 428: add a "," before "which", and "which would still need" instead of "which still would
need"

R: this was changed in the revised version

l. 490: remove the very first parenthesis (before i.e.)

R: this was removed

l. 491; replace "of" by "in"

R: this was modified

1332

1333

1334

1335