Retraction ATPase motors are promiscuous among type IVa pilus systems

Evan Couser[1], Jennifer L. Chlebek[1], and Ankur B. Dalia[1,*]

[1]Department of Biology, Indiana University, Bloomington, IN 47405.
*Correspondence to: ankdalia@indiana.edu.

Running title: T4aP retraction ATPase motors are promiscuous

ABSTRACT

Bacterial surface appendages called type IVa pili (T4aP) promote diverse activities including DNA uptake, twitching motility, and virulence. These activities rely on the ability of T4aP to dynamically extend and retract from the cell surface. Dynamic extension relies on a motor ATPase commonly called PilB. Most T4aP also rely on specific motor ATPases, commonly called PilT and PilU, to dynamically and forcefully retract. Here, we systematically assess whether motor ATPases from 4 distinct T4aP could functionally complement *Vibrio cholerae* mutants that lacked their endogenous motors. We found that the retraction ATPases PilT and PilU are highly promiscuous and promote retraction of the *V. cholerae* competence T4aP despite a high degree of sequence divergence. In contrast, orthologous extension ATPases were not able to mediate extension of the *V. cholerae* competence T4aP despite a similar degree of sequence divergence. Also, we show that one of the PilT orthologs characterized does not support PilU-dependent retraction and we attributed this loss of activity to the 3' end of the gene, which suggests that the C-terminus of PilT plays an important role in promoting PilU-dependent retraction. Together, our data suggest that retraction ATPases have maintained a high degree of promiscuity for promoting retraction of diverse T4aP, while extension ATPases have evolved to become highly specific for their cognate systems.

IMPORTANCE

One way that bacteria interact with their environments is via hair-like appendages called type IVa pili (T4aP). These appendages dynamically extend and retract from the cell surface via the action of distinct ATPase motors. T4aP are present in diverse bacterial species. Here, we demonstrate that retraction motors from T4aP are highly promiscuous, and capable of promoting retraction of heterologous T4aP systems. By contrast, extension ATPase motors are highly specific and cannot promote extension of heterologous T4aP. Thus, these results suggest that T4aP extension may be more tightly regulated compared to T4aP retraction.

INTRODUCTION

Type IVa pilus systems (T4aP) are encoded by diverse bacterial species. These systems promote the dynamic extension and retraction of filamentous appendages called pili from the bacterial surface. Through this dynamic activity, T4aP carry out a variety of functions including twitching motility, surface attachment, biofilm formation, virulence, and DNA uptake (1). Dynamic extension of all T4aP relies on the activity of a dedicated motor ATPase, which is commonly called PilB (2). Some systems also have an additional accessory extension ATPase motor called TfpB (3). Dynamic retraction of T4aP canonically relies on a dedicated retraction ATPase called PilT. Additionally, many T4aP systems also contain an accessory retraction

ATPase called PilU. PilU is a PilT-dependent retraction ATPase that is required for forceful
retraction of the competence T4aP in *V. cholerae* (4) and is required for twitching motility in
other bacterial species (5). These hexameric motors are thought to function via a mechanical
interaction with the pilus platform, which together take pilin subunits from the inner membrane
and polymerize them into a filament during pilus extension or depolymerize these subunits and
return them to the inner membrane during pilus retraction. Conformational changes in the
ATPase motors due to ATP binding and hydrolysis are thought to power these processes (6).
Previous studies have demonstrated that retraction motors can be interchangeable between
some T4aP systems (7). However, a systematic analysis to assess motor promiscuity among
T4aP is lacking.

Vibrio cholerae is a facultative pathogen that encodes a competence T4aP, which promotes
DNA uptake for natural transformation (8, 9). *Acinetobacter baylyi* and *Pseudomonas stutzeri*
encode functionally analogous competence T4aP that also promote horizontal gene transfer (3,
10-12). All of these T4aP systems encode canonical PilB, PilT, and PilU motor ATPases. The
T4aP of *A. baylyi* also encodes an additional extension motor called TfpB that is required for
initiating pilus extension (3). Some strains of *Escherichia coli* also encode an ortholog of PilT
(also referred to as *yggR*) though these strains are not predicted to encode a PilU ortholog.

Here, we sought to formally test promiscuity among T4aP extension and retraction ATPases by
using the motors from the systems discussed above. We and others have demonstrated that the
competence pilus of *Vibrio cholerae* is a highly genetically tractable system for studying T4aP
dynamic activity (4, 7, 8, 13). Thus, we assessed promiscuity among T4aP motors by
determining if ectopic expression of heterologous motors could functionally complement *V.*
cholerae mutants that lack their endogenous motor ATPases. Our data suggests that while the
retraction ATPases PilT and PilU may be promiscuous in their ability to mediate retraction of
T4aP systems, extension ATPases display a high level of specificity for their cognate T4aP.
Furthermore, our data suggest that the C-terminus of PilT may play an important role in
supporting PilU-dependent retraction.

RESULTS

Retraction ATPase orthologs are promiscuous among T4aP systems, while extension ATPase
orthologs are not.
Because previous studies have shown that retraction ATPases from T4aP systems may be
promiscuous (7), we first wanted to determine if this characteristic is conserved among other
bacterial species. To test this, we ectopically expressed *pilT* from *A. baylyi* ($pilT_{Ab}$), *P. stutzeri*
($pilT_{Ps}$), and *E. coli* ($pilT_{Ec}$) in *V. cholerae* strains where the native *pilT* or *pilTU* genes were
inactivated (i.e., Δ*pilT* and Δ*pilTU* backgrounds). As a control, we also ectopically expressed *V.*
cholerae pilT ($pilT_{Vc}$). Competence pili in *V. cholerae* promote DNA uptake during natural
transformation, a function that requires both pilus extension and retraction (8, 14). Thus, we
functionally assessed competence pilus dynamic activity by performing natural transformation
assays. As expected, ectopic expression of $pilT_{Vc}$ rescued the transformation defect of the Δ*pilT*
and Δ*pilTU* strains (**Fig. 1A**). Furthermore, we observed that ectopic expression of $pilT_{Ab}$, $pilT_{Ps}$,
and $pilT_{Ec}$, all fully complemented natural transformation of Δ*pilT* and Δ*pilTU V. cholerae* (**Fig.**
1A) These data indicate that all of these retraction ATPases are capable of mediating retraction
of the *V. cholerae* competence T4aP, suggesting that the promiscuity of PilT retraction ATPases
is conserved across these species.

In addition to the dominant retraction ATPase PilT, many T4aP also encode an accessory
retraction ATPase called PilU that can promote pilus retraction in a PilT-dependent manner (4,

7). So next, we wanted to test whether PilU from *A. baylyi* and *P. stutzeri* could complement a
V. cholerae pilU mutant (the *E. coli* strain does not have a *pilU* ortholog). To test this, we
ectopically expressed *pilU_{Ab}* and *pilU_{Ps}* in a *V. cholerae* strain where native *pilT* contained a
Walker A mutation ($pilT^{K136A}$ = $pilT^{WA}$) and the native *pilU* gene was inactivated (i.e., Δ*pilU*).
Mutation of the invariantly conserved lysine in the Walker A domain (K136A) inhibits the
ATPase activity of *pilT* (15), thus, efficient pilus retraction in this background (and
correspondingly a high frequency of natural transformation) is reliant on PilU ATPase activity (4,
7). As expected, ectopic expression of *pilU_{Vc}* recovered the transformation defect of the $pilT^{WA}$
Δ*pilU* strain to parental levels (**Fig. 1B**). The expression of *pilU_{Ab}* also showed almost full
recovery of transformation when compared to the parent strain (**Fig. 1B**). Interestingly, ectopic
expression of *pilU_{Ps}* only partially complemented the transformation defect of this strain (**Fig.**
1B). This suggested that PilU_{Ps} was either (1) generally a less functional PilU motor, or perhaps
(2) specifically less active at promoting pilus retraction in conjunction with PilT_{Vc}. To test this
further, we ectopically expressed *pilU_{Ps}* in a background that also ectopically expressed a
Walker A mutant of its cognate PilT ($pilT_{Ps}{}^{WA}$). We found that in the presence of PilT_{Ps}{}^{WA}, PilU_{Ps}
was able to completely rescue natural transformation (**Fig. S1**). Together, this suggests that
while PilU is also promiscuous between T4aP, it is perhaps less promiscuous than PilT.

Because we found that retraction ATPases are, indeed, promiscuous between these T4aP, we
next wanted to test whether the extension ATPases from these same organisms were similarly
promiscuous. To test this, we ectopically expressed *pilB* orthologs from *A. baylyi* (*pilB_{Ab}* and
tfpB_{Ab}) and *P. stutzeri* (*pilB_{Ps}*) in a *V. cholerae* Δ*pilB* mutant and assessed natural transformation
(the *E. coli* PilB ortholog, HofB, was not tested). While *pilB_{Vc}* recovered natural transformation
as expected, ectopic expression of *pilB_{Ab}*, *tfpB_{Ab}*, and *pilB_{Ps}* failed to complement the Δ*pilB*
mutant (**Fig. 1C**). This suggests that extension ATPase orthologs from these species are not
able to mediate extension of the *V. cholerae* competence pilus. These data suggest that there
may be a high degree of specificity among extension ATPases, which is in stark contrast to the
observed promiscuity among retraction ATPases.

Extension and retraction ATPases likely interact with the pilus machinery platform protein PilC
to promote dynamic pilus activity (16). Thus, the observed promiscuity and/or specificity among
these motor ATPases may be attributed to their ability and/or inability to interact with PilC. To
test this, we assessed protein-protein interactions genetically using Bacterial Adenylate Cyclase
Two-Hybrid (BACTH) assays (17). First, looking at PilT, we found that each of the PilT orthologs
could self-interact, which is consistent with the hexameric conformation that likely all pilus
motors adopt (**Fig. 1D**). Also, consistent with the observation that all of the PilT orthologs tested
support natural transformation, we found that all of them could interact with PilC_{Vc} (**Fig. 1D**).
Because PilU is a PilT-dependent retraction ATPase, it may need to interact with both PilT and
PilC to support pilus retraction. Indeed, all of the PilU orthologs could interact with PilT_{Vc},
PilT_{Vc}{}^{WA}, and PilC_{Vc}, which is consistent with the ability of each PilU ortholog to support some
degree of natural transformation (**Fig. 1E**). For the PilB orthologs, we found that PilB_{Vc} could
interact with PilC_{Vc} as expected (**Fig. 1F**). Additionally, each of the PilB orthologs could self-
interact, which is consistent with the hexameric conformation that these proteins likely adopt
(**Fig. 1F**). Interestingly, PilB_{Ab}, TfpB_{Ab}, and PilB_{Ps} varied in their ability to interact with PilC_{Vc}.
PilB_{Ab} and PilB_{Ps} could not interact with PilC_{Vc} (**Fig. 1F**), which could suggest that this interaction
is required for extension of the competence pilus. However, TfpB_{Ab} could interact with PilC (**Fig.**
1F), but it could not support natural transformation (**Fig. 1C**). These data indicate that while
interactions between PilB and PilC may be required for extension of the competence pilus,
these interactions are not sufficient to promote extension.

The PilT ortholog from E. coli cannot support PilU-mediated retraction.

Above, our results indicate that PilT orthologs from distinct T4aP can promote retraction of the
V. cholerae competence pilus in the absence of native PilT$_{Vc}$ and PilU$_{Vc}$. PilT, however, can also
support PilU-dependent retraction, in a manner that is not dependent on PilT ATPase activity
(4). So next, we wanted to test if the PilT orthologs among the studied T4aP could support PilU-
dependent retraction in conjunction with PilU$_{Vc}$. To test this, Walker A mutants of each *pilT* gene
(*pilT$_{Vc}$*^{WA}, *pilT$_{Ab}$*^{WA}, *pilT$_{Ps}$*^{WA}, *pilT$_{Ec}$*^{WA}) were ectopically expressed in a Δ*pilT* background (i.e.,
where *pilU$_{Vc}$* is intact) and a Δ*pilTU* background. As expected, the ectopic expression of *pilT$_{Vc}$*^{WA}
fully restored the transformation efficiency of a Δ*pilT* background (since *pilU$_{Vc}$* is intact) (**Fig.**
2A). Importantly, transformation was not restored by ectopic expression of any Walker A *pilT*
alleles in a Δ*pilTU* background (**Fig. 2A**), which indicates that these ATPase defective alleles of
PilT only support PilU-dependent retraction and cannot promote retraction on their own. When
pilT$_{Ab}$^{WA} and *pilT$_{Ps}$*^{WA} were ectopically expressed in the Δ*pilT* background, transformation was
rescued to parental levels (**Fig. 2A**). These data suggest that PilU$_{Vc}$ is able to mediate retraction
of the *V. cholerae* competence pilus in a PilT$_{Ab}$- and PilT$_{Ps}$-dependent manner, which further
demonstrates the promiscuity among these retraction ATPases. Ectopic expression of the
Walker A mutant allele of *pilT$_{Ec}$*, however, did not restore natural transformation in the Δ*pilT*
background (**Fig. 2A**). This suggests that PilT$_{Ec}$^{WA} is incapable of promoting PilU-dependent
retraction in the presence of PilU$_{Vc}$.

The ability of PilU to interact with PilT is likely necessary for PilU-dependent retraction of the
competence pilus. As expected, PilT$_{Vc}$, PilT$_{Ab}$, PilT$_{Ps}$, and PilT$_{Ec}$ could self-interact (**Fig. 2B**) as
seen previously (**Fig. 1D**), and each allele could also interact with PilU$_{Vc}$ (**Fig. 2B**). Because
PilU$_{Vc}$-mediated retraction was possible when *pilT$_{Vc}$*^{WA}, *pilT$_{Ab}$*^{WA}, and *pilT$_{Ps}$*^{WA} were ectopically
expressed, we hypothesized that each of these PilT alleles would be able to self-interact and
also interact with PilU. PilT$_{Vc}$^{WA} and PilT$_{Ab}$^{WA} were able to self-interact (**Fig. 2B**), which suggests
that the Walker A mutation does not inhibit the ability of these proteins to self-assemble into
hexamers. This is consistent with our previous results which demonstrate that purified PilT$_{Vc}$^{WA}
can still form hexameric rings (4). Unexpectedly, PilT$_{Ps}$^{WA} and PilT$_{Ec}$^{WA} did not exhibit self-
interactions in these BACTH assays (**Fig. 2B**). We also observed that PilU$_{Vc}$ could interact with
PilT$_{Vc}$^{WA} and PilT$_{Ab}$^{WA}, however, it did not interact with PilT$_{Ec}$^{WA} or PilT$_{Ps}$^{WA} (**Fig. 2B**). The
absence of interactions between PilU$_{Vc}$ and PilT$_{Ec}$^{WA} (**Fig. 2B**) is consistent with the inability of
PilT$_{Ec}$^{WA} to promote PilU-dependent retraction (**Fig. 2A**). However, the absence of interactions
between PilU$_{Vc}$ and PilT$_{Ps}$^{WA} was particularly surprising since this PilT allele can support PilU-
dependent retraction when expressed in *V. cholerae* (**Fig. 2A**). Because PilT$_{Ps}$^{WA} supports PilU-
dependent retraction in *V. cholerae*, but does not exhibit self-interactions in BACTH assays, this
may suggest that self-assembly or hexamerization of PilT is not required for PilU-dependent
retraction. However, the inability of PilT$_{Ps}$^{WA} to interact with PilU$_{Vc}$ in these same BACTH assays
(**Fig. 2B**), despite the ability of this allele to support PilU-dependent retraction (**Fig. 2A**) makes
this difficult to conclude. The observed inability of PilT$_{Ec}$^{WA} and PilT$_{Ps}$^{WA} to self-interact and/or
interact with PilU$_{Vc}$ in BACTH assays could either be due to an absence of interactions between
these proteins, or could be due to a poor affinity interaction between these proteins that are
below the detection limit for the assay.

The molecular mechanism underlying PilU-mediated retraction in a PilT-dependent manner
remains unclear. The ability of PilT to self-assemble and form a hexamer may be a prerequisite
for PilU-mediated retraction. However, other characteristics of PilT may also be responsible for
supporting PilU-mediated retraction. Because *pilT$_{Ec}$*^{WA} lacked self-interactions, we constructed
an alternative mutant that also disrupted the ATPase activity of PilT$_{Ec}$ by mutating an invariantly
conserved glutamate in the Walker B domain of *pilT$_{Ec}$* (*pilT$_{Ec}$*^{E201A} = *pilT$_{Ec}$*^{WB}). Importantly,
PilT$_{Ec}$^{WB} exhibited self-interactions in BACTH assays (**Fig. 2B**), suggesting that the Walker B
mutation allows for self-assembly and/or a higher affinity between subunits compared to the

Walker A mutant. When we ectopically expressed the $pilT_{Ec}^{WB}$ in a $\Delta pilT$ background (where
$pilU$ is intact), however, we observed that this mutant did not restore transformation (**Fig. 2A**),
which is similar to the $pilT_{Ec}^{WA}$ allele. This indicates that self-assembly of an ATPase defective
$PilT_{Ec}$ is not sufficient to promote PilU-dependent retraction. Together, these data suggest that
$PilT_{Ec}$ may have unique features that prevents it from promoting PilU-dependent retraction. We
hypothesized that analyzing the differences between $PilT_{Ec}$ and the other PilT alleles ($PilT_{Vc}$,
$PilT_{Ab}$, and $PilT_{Ps}$) may help define the features of PilT that allow for PilU-dependent retraction.

The C-terminus of $PilT_{Vc}$ contributes to PilU-mediated retraction.
In order to narrow down the region of $PilT_{Vc}$ that is necessary and/or sufficient to support PilU-
dependent retraction, we constructed defined chimeras of $pilT_{Vc}$ and $pilT_{Ec}$. These chimeras
were first ectopically expressed in a $\Delta pilTU$ background to test their ability to mediate retraction
of the competence pilus in a manner that is dependent on their ATPase activity – a property that
both $PilT_{Vc}$ and $PilT_{Ec}$ shared (**Fig. 1A**). Surprisingly, only three of the chimeras were able to
restore parental levels of transformation in the $\Delta pilTU$ background (**Fig. 3A**). These are the only
chimeras of PilT that we can be reasonably confident could potentially support PilU-dependent
retraction, thus, we only proceeded with characterizing these three chimeras of PilT moving
forward. Next, we introduced a Walker A mutation into these chimeras to determine if they were
able to support PilU-mediated retraction. Consistent with the idea that PilT self-assembly is
required for PilU-mediated retraction, none of the $pilT_{Vc}$ / $pilT_{Ec}$ chimeras with the Walker A
mutation were able to rescue transformation efficiency in a $\Delta pilT$ background (**Fig. 3B**).
Separately, a Walker B mutation was introduced into these chimeras. Again, none of the 3
chimeras were able to restore transformation in a $\Delta pilT$ background (**Fig. 3B**). Because the
chimera that contained the N-terminal ~200 amino acids of $PilT_{Vc}$ and C-terminal ~150 amino
acids of $PilT_{Ec}$ did not support PilU-dependent retraction, we reasoned that the C-terminus of
$PilT_{Vc}$ is important for PilU-dependent retraction.

In order to further investigate potential components of PilT that are required for supporting PilU-
mediated retraction, we constructed an alignment of the 4 PilT orthologs that were studied here.
The alignment revealed that the middle section of $PilT_{Ec}$ was conserved relative to all of the
other PilT orthologs (**Fig. S2**). However, the N-terminus and the C-terminus of $PilT_{Ec}$ were
poorly conserved relative to the other PilT orthologs. Because our data suggests that the C-
terminus of $PilT_{Vc}$ is required for supporting PilU-mediated retraction, that is where we focused
our investigation. In particular, we observed that $pilT_{Ec}$ contains two substitutions in the highly
conserved AIRNLIRE domain of PilT (18). To determine if these substitutions were responsible
for the inability of $PilT_{Ec}$ to support PilU-mediated retraction (7), we mutated the two residues in
$pilT_{Ec}$ to the residues present in the other PilT orthologs ($PilT_{Ec}^{AVG \rightarrow AIR}$). Conversely, to determine
if these changes in $PilT_{Ec}$ contributed to its inability to support PilU-mediated retraction, the
residues in $pilT_{Vc}$ were mutated to the residues present in $pilT_{Ec}$ ($PilT_{Vc}^{AIR \rightarrow AVG}$). These mutations
were introduced into Walker A (WA) or Walker B (WB) mutant alleles of PilT so that we could
assess PilU-mediated retraction. The introduction of these mutations into $pilT_{Vc}$ did not result in
the inability of $PilT_{Vc}$ to facilitate PilU-mediated retraction (**Fig. 3C**). Moreover, the restoration of
the two AIRNLIRE substitutions in $pilT_{Ec}$ did not rescue PilU-mediated retraction (**Fig. 3C**).
These data indicate that the substitutions in the AIRNLIRE domain of $pilT_{Ec}$ are not responsible
for the inability of $PilT_{Ec}$ to support PilU-mediated retraction.

The alignment of the 4 PilT orthologs also revealed that 15 amino acids in the C-terminal
domain of $PilT_{Ec}$ were missing when compared to the other PilT orthologs that did support PilU-
mediated retraction ($PilT_{Vc}$, $PilT_{Ab}$, and $PilT_{Ps}$). We hypothesized that the deletion of the last 15
amino acids in $PilT_{Ec}$ could be responsible for the inability of this protein to support PilU-
mediated retraction. To test this, we deleted the last 15 amino acids from $PilT_{Vc}$ ($pilT_{Vc}^{\Delta 15aa}$).
Additionally, we fused the last 15 amino acids present at the C-terminus of $pilT_{Vc}$ to the end of

PilT$_{Ec}$ (*pilT$_{Ec}$$^{+15aa}$*). Again, these mutations were generated in Walker A and Walker B mutant
alleles of PilT so that we could assess PilU-mediated retraction. The *pilT$_{Vc}$$^{Δ15aa, WA}$* mutant
displayed no reduction in transformation (**Fig. 3C**), suggesting that PilT$_{Vc}$$^{Δ15aa, WA}$ is still able to
facilitate PilU-mediated retraction. Conversely, the *pilT$_{Ec}$$^{+15aa, WA}$* and *pilT$_{Ec}$$^{+15aa, WB}$* mutations did
not restore transformation to parental levels (**Fig. 3C**), which suggests that PilU$_{Vc}$ cannot
mediate retraction in the presence of these PilT$_{Ec}$ alleles. Together, these data suggest that the
last 15 amino acids present at the C-terminus of PilT$_{Vc}$ is not necessary or sufficient to promote
PilU-mediated retraction.

Our results have suggested that, individually, the substitutions present in the conserved
AIRNLIRE domain of PilT$_{Ec}$ and the deletion of the last 15 amino acids in the C-terminus are not
responsible for the inability of this PilT allele to support PilU-mediated retraction. However, it is
possible that the combination of these two factors diminishes PilU-mediated retraction. To test
this, we constructed *pilT$_{Vc}$ alleles* that contained both the AIRNLIRE domain substitutions and
the C-terminal deletion of 15 amino acids (*pilT$_{Vc}$$^{AIR→AVG, Δ15aa}$*). Also, we generated a *pilT$_{Ec}$* allele
where we restored the AIRNLIRE domain and inserted the terminal 15 amino acids from *pilT$_{Vc}$*
onto the C-terminus (*pilT$_{Ec}$$^{AVG→AIR, +15aa}$*). The combination of mutations in *pilT$_{Vc}$* resulted in a loss
of function, even in a strain where the ATPase activity of the allele was intact (**Fig. 3C**).
Additionally, the *pilT$_{Ec}$$^{AVG→AIR, +15aa, WA}$* and *pilT$_{Ec}$$^{AVG→AIR, +15aa, WB}$* alleles did not restore
transformation to parental levels in the Δ*pilT* background, suggesting that PilU was unable to
mediate retraction in these backgrounds. This suggests that the combination of the 15 amino
acid deletion and the substitutions in the AIRNLIRE domain are not solely responsible for the
inability of PilT$_{Ec}$ to support PilU-mediated retraction.

DISCUSSION

This work demonstrates that the retraction ATPases PilT and PilU are highly promiscuous
among T4aP systems, which contrasts with the fidelity displayed by the extension ATPases
from the same systems. Additionally, our data suggests that the C-terminus of PilT$_{Vc}$ likely plays
an important role in supporting PilU-mediated retraction. However, the exact residues in the PilT
C-terminus that contribute to PilU-mediated retraction remain elusive.

One possible explanation for the discrepancy in the promiscuity / specificity among motor
ATPases could be due to differences in sequence divergence; where retraction motor orthologs
are more similar to one another compared to the extension motor orthologs. When we
performed this analysis, however, we found that the PilT orthologs ranged from ~62-80% similar
(with PilT$_{Ec}$ being the most divergent), the PilU orthologs ranged from ~70-74% similar, and the
PilB orthologs ranged from ~64-66% similar (and the TfpB ortholog was ~53% similar to PilB$_{Vc}$)
(**Fig. S3**). These data indicate that overall sequence divergence among PilT and PilB orthologs
is fairly similar. Thus, suggesting that sequence divergence alone cannot account for the
promiscuity / specificity among these motor ATPases.

Why are retraction ATPases promiscuous and extension ATPases specific? One possibility is
that it is advantageous for T4aP systems to maintain promiscuous retraction ATPases. T4aP
are highly diverse and retraction ATPases are generally expressed at distinct loci from the other
T4aP components. Some T4aP systems are likely transferred between organisms through
horizontal gene transfer (19). The inability to retract T4aP is likely disadvantageous in many
circumstances since these surface appendages are highly immunogenic and can also be
receptors for diverse phages (20). Thus, it is tempting to speculate that one potential advantage
to retraction ATPase promiscuity is that this would allow cells that acquire a new T4aP the
ability to retract pili from the surface.

Some organisms, like *V. cholerae*, encode multiple T4P - competence pili, mannose sensitive hemagglutinin (MSHA) pili, and toxin co-regulated pili (TCP) - and there are distinct environmental conditions under which each pilus system is likely needed. Competence pili and MSHA pili are likely critical for the survival of *V. cholerae* in its environmental reservoir, while TCP are required for infection of its human host. In fact, it has previously been shown that the inappropriate expression of MSHA pili during the context of an infection attenuates *V. cholerae* (21). A high degree of specificity among the extension ATPase motors for these distinct T4P may represent one mechanism by which cells can exhibit tight control over when these distinct pili are made. In contrast, the retraction of these pili may not require the same level of regulation because pilus extension is so tightly regulated. Indeed, in *V. cholerae*, PilT and PilU promote retraction of both competence pili and MSHA pili (8, 22).

The molecular mechanism underlying the specificity of extension ATPase motors remains unclear. One possibility is that the N-terminal domain of PilB provides specificity. It is thought that the N-terminal domain of PilB is responsible for interactions with PilM, which is likely required for proper pilus extension (6). An alignment of the extension and retraction motor orthologs showed that the extension motors contain an extended N-terminal domain (~184 residues) that distinguishes them from the retraction ATPase motors (**Fig. S4**), and we found that much of the sequence divergence among extension ATPases was attributed to this N-terminal domain (**Fig. S3**). Thus, it is tempting to speculate that the N-terminal domain of extension ATPases plays a critical role in defining the specificity of these motors for their cognate T4aP system.

While our results indicate that retraction motors are promiscuous and extension motors are specific, it remains unclear which of these phenotypes is the exception vs the rule. To distinguish between these possibilities, it would be interesting to test the degree of specificity / promiscuity among other T4aP components. Addressing this question may continue to shed light on the evolutionary pressures that have helped shaped these nearly ubiquitous molecular machines, and will be the focus of future research.

METHODS

Bacterial Strains and Growth Conditions
Vibrio cholerae strains were grown in LB Miller broth and LB Miller agar plates that contained erythromycin (10 μg/mL), carbenicillin (20 μg/mL), and/or kanamycin (50 μg/mL) when necessary. A list of all strains used in this study can be found in **Table S1**.

Construction of Mutant Strains and Constructs
All of the *V. cholerae* strains used in this study are derivatives of the El tor isolate E7946. The motor ATPase orthologs from other species were amplified from genomic DNA of the following strains: *Acinetobacter baylyi* ADP1 (23), *Pseudomonas stutzeri* JM300 (24), and *Escherichia coli* MG1655 (25). All mutant constructs were made via splicing-by-overlap extension (SOE) PCR, exactly as previously described (13, 26). A detailed list of all primers used to generate mutant constructs can be found in **Table S2**. Mutant constructs were introduced onto the *V. cholerae* genome by natural transformation and natural cotransformation exactly as previously described (26).

Natural Transformation Assays
In this study, we aimed to address the role of distinct T4aP components on competence pilus function. Thus, we opted to bypass the natural mechanism of competence induction in *V. cholerae* by ectopically expressing the master competence regulator TfoX (P_{tac}-*tfoX*) and

inactivating the quorum sensing negative regulator LuxO (Δ*luxO*) as previously described (8).
Chitin-independent natural transformation assays were performed exactly as previously
described (8, 14). Briefly, strains were grown overnight in 3 mL of LB broth containing 100 μM
IPTG at 30 °C rolling. Next, cells were subcultured into 3 mL of LB supplemented with 20 mM
$MgCl_2$, 10 mM $CaCl_2$, 100 μM IPTG, and arabinose (when appropriate; 0.05-0.2%) and grown to
late-log phase at 30 °C rolling. Next, ~10^8 cells were transferred to Instant Ocean medium (7
g/L; Aquarium Systems) containing 100 μM IPTG, arabinose (when appropriate; 0.05-0.2%),
and 200 ng of a transforming DNA (tDNA) PCR product. Reactions were then incubated
statically overnight at 30 °C. The tDNA product used throughout this study targeted a "neutral
gene" (the frame-shifted transposase, VC1807) for inactivation (ΔVC1807::Erm^R) as previously
described (27). A negative control, containing no tDNA, was included for each strain in all
transformation assays. After the overnight incubation with tDNA, reactions were outgrown by
adding 1 mL of LB to each reaction and incubating at 37 °C shaking for 2 hours. After the
outgrowth, reactions were plated for quantitative culture to distinguish between transformants
(plated on LB + erythromycin) and total viable counts (plated on plain LB). The transformation
frequency for each reaction was calculated by dividing the number of transformants by the total
viable counts.

Bacterial Adenylate Cyclase Two-Hybrid Assay
Using PCR, the desired motor ATPase genes were amplified and cloned into the BACTH
vectors pKT25 and pUT18C in order to produce N-terminal fusions between the motor ATPase
and the T25 and T18 fragments of adenylate cyclase, respectively. Miniprepped plasmids
(Qiagen) were then co-transformed into *E. coli* BTH101 cells. As a positive control, T25-zip and
T18-zip vectors were cotransformed into BTH101. And as a negative control, the pKT25 and
pUT18C empty vectors were cotransformed into BTH101. Transformations were plated onto LB
+ kanamycin (50 μg/mL) + carbenicillin (100 μg/mL) plates in order to select for transformants
that received both plasmids. These transformants were then selected and grown overnight at 30
°C static in LB + kanamycin (50 μg/mL) + carbenicillin (100 μg/mL). Finally, 3 μL of the overnight
cultures were spotted onto LB agar plates supplemented with 500 μM IPTG, kanamycin (50
μg/mL), carbenicillin (100 μg/mL), and X-gal (40 μg/mL). These plates were allowed to incubate
statically at 30 °C for ~48 hours prior to imaging on a HP Scanjet G4010 flatbed scanner.

Alignments
The sequences of the genes that encode for the desired proteins were downloaded from
GenBank. Sequences were aligned using T-Coffee (28) and the resulting alignments were run
through Boxshade to generate the final figures.

Pairwise comparisons to assess similarity among ATPase motor orthologs
Pairwise comparisons between ATPase motor orthologs were performed via global alignment
using the Needleman-Wunsch algorithm (29). The percent similarities generated by these
pairwise comparisons are reported in **Fig. S3**.

ACKNOWLEDGEMENTS

We would like to thank David Kysela for helpful discussions. This work was supported by grant
R35GM128674 from the National Institutes of Health (to ABD).

References

1. Craig L, Forest KT, Maier B. 2019. Type IV pili: dynamics, biophysics and functional consequences. Nat Rev Microbiol 17:429-440.

2. Jakovljevic V, Leonardy S, Hoppert M, Sogaard-Andersen L. 2008. PilB and PilT are ATPases acting antagonistically in type IV pilus function in Myxococcus xanthus. J Bacteriol 190:2411-21.

3. Ellison CK, Dalia TN, Klancher CA, Shaevitz JW, Gitai Z, Dalia AB. 2021. Acinetobacter baylyi regulates type IV pilus synthesis by employing two extension motors and a motor protein inhibitor. Nat Commun 12:3744.

4. Chlebek JL, Hughes HQ, Ratkiewicz AS, Rayyan R, Wang JC, Herrin BE, Dalia TN, Biais N, Dalia AB. 2019. PilT and PilU are homohexameric ATPases that coordinate to retract type IVa pili. PLoS Genet 15:e1008448.

5. Whitchurch CB, Mattick JS. 1994. Characterization of a gene, pilU, required for twitching motility but not phage sensitivity in Pseudomonas aeruginosa. Mol Microbiol 13:1079-91.

6. McCallum M, Tammam S, Khan A, Burrows LL, Howell PL. 2017. The molecular mechanism of the type IVa pilus motors. Nat Commun 8:15091.

7. Adams DW, Pereira JM, Stoudmann C, Stutzmann S, Blokesch M. 2019. The type IV pilus protein PilU functions as a PilT-dependent retraction ATPase. PLoS Genet 15:e1008393.

8. Ellison CK, Dalia TN, Vidal Ceballos A, Wang JC, Biais N, Brun YV, Dalia AB. 2018. Retraction of DNA-bound type IV competence pili initiates DNA uptake during natural transformation in Vibrio cholerae. Nat Microbiol 3:773-780.

9. Seitz P, Blokesch M. 2013. DNA-uptake machinery of naturally competent Vibrio cholerae. Proc Natl Acad Sci U S A 110:17987-92.

10. Leong CG, Bloomfield RA, Boyd CA, Dornbusch AJ, Lieber L, Liu F, Owen A, Slay E, Lang KM, Lostroh CP. 2017. The role of core and accessory type IV pilus genes in natural transformation and twitching motility in the bacterium Acinetobacter baylyi. PLoS One 12:e0182139.

11. Graupner S, Weger N, Sohni M, Wackernagel W. 2001. Requirement of novel competence genes pilT and pilU of Pseudomonas stutzeri for natural transformation and suppression of pilT deficiency by a hexahistidine tag on the type IV pilus protein PilAI. J Bacteriol 183:4694-701.

12. Graupner S, Frey V, Hashemi R, Lorenz MG, Brandes G, Wackernagel W. 2000. Type IV pilus genes pilA and pilC of Pseudomonas stutzeri are required for natural genetic transformation, and pilA can be replaced by corresponding genes from nontransformable species. J Bacteriol 182:2184-90.

13. Dalia TN, Chlebek JL, Dalia AB. 2020. A modular chromosomally integrated toolkit for ectopic gene expression in Vibrio cholerae. Sci Rep 10:15398.

14. Chlebek JL, Dalia TN, Biais N, Dalia AB. 2021. Fresh Extension of Vibrio cholerae Competence Type IV Pili Predisposes Them for Motor-Independent Retraction. Appl Environ Microbiol 87:e0047821.

15. Chiang P, Sampaleanu LM, Ayers M, Pahuta M, Howell PL, Burrows LL. 2008. Functional role of conserved residues in the characteristic secretion NTPase motifs of the Pseudomonas aeruginosa type IV pilus motor proteins PilB, PilT and PilU. Microbiology (Reading) 154:114-126.

16.   Bischof LF, Friedrich C, Harms A, Sogaard-Andersen L, van der Does C. 2016. The
Type IV Pilus Assembly ATPase PilB of Myxococcus xanthus Interacts with the Inner
Membrane Platform Protein PilC and the Nucleotide-binding Protein PilM. J Biol Chem
291:6946-57.
17.   Karimova G, Pidoux J, Ullmann A, Ladant D. 1998. A bacterial two-hybrid system based
on a reconstituted signal transduction pathway. Proc Natl Acad Sci U S A 95:5752-6.
18.   Aukema KG, Kron EM, Herdendorf TJ, Forest KT. 2005. Functional dissection of a
conserved motif within the pilus retraction protein PilT. J Bacteriol 187:611-8.
19.   Denise R, Abby SS, Rocha EPC. 2019. Diversification of the type IV filament
superfamily into machines for adhesion, protein secretion, DNA uptake, and motility.
PLoS Biol 17:e3000390.
20.   Giltner CL, Nguyen Y, Burrows LL. 2012. Type IV pilin proteins: versatile molecular
modules. Microbiol Mol Biol Rev 76:740-72.
21.   Hsiao A, Liu Z, Joelsson A, Zhu J. 2006. Vibrio cholerae virulence regulator-coordinated
evasion of host immunity. Proc Natl Acad Sci U S A 103:14542-7.
22.   Floyd KA, Lee CK, Xian W, Nametalla M, Valentine A, Crair B, Zhu S, Hughes HQ,
Chlebek JL, Wu DC, Hwan Park J, Farhat AM, Lomba CJ, Ellison CK, Brun YV,
Campos-Gomez J, Dalia AB, Liu J, Biais N, Wong GCL, Yildiz FH. 2020. c-di-GMP
modulates type IV MSHA pilus retraction and surface attachment in Vibrio cholerae. Nat
Commun 11:1549.
23.   Barbe V, Vallenet D, Fonknechten N, Kreimeyer A, Oztas S, Labarre L, Cruveiller S,
Robert C, Duprat S, Wincker P, Ornston LN, Weissenbach J, Marliere P, Cohen GN,
Medigue C. 2004. Unique features revealed by the genome sequence of Acinetobacter sp.
ADP1, a versatile and naturally transformation competent bacterium. Nucleic Acids Res
32:5766-79.
24.   Busquets A, Pena A, Gomila M, Bosch R, Nogales B, Garcia-Valdes E, Lalucat J,
Bennasar A. 2012. Genome sequence of Pseudomonas stutzeri strain JM300 (DSM
10701), a soil isolate and model organism for natural transformation. J Bacteriol
194:5477-8.
25.   Blattner FR, Plunkett G, 3rd, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J,
Glasner JD, Rode CK, Mayhew GF, Gregor J, Davis NW, Kirkpatrick HA, Goeden MA,
Rose DJ, Mau B, Shao Y. 1997. The complete genome sequence of Escherichia coli K-
12. Science 277:1453-62.
26.   Dalia AB. 2018. Natural Cotransformation and Multiplex Genome Editing by Natural
Transformation (MuGENT) of Vibrio cholerae. Methods Mol Biol 1839:53-64.
27.   Dalia AB, McDonough E, Camilli A. 2014. Multiplex genome editing by natural
transformation. Proc Natl Acad Sci U S A 111:8937-42.
28.   Notredame C, Higgins DG, Heringa J. 2000. T-Coffee: A novel method for fast and
accurate multiple sequence alignment. J Mol Biol 302:205-17.
29.   Needleman SB, Wunsch CD. 1970. A general method applicable to the search for
similarities in the amino acid sequence of two proteins. J Mol Biol 48:443-53.

Fig. 1. *Retraction ATPase orthologs are promiscuous among T4aP systems, while extension ATPase orthologs are not*. Natural transformation assays were performed to test whether (**A**) PilT, (**B**) PilU, or (**C**) PilB/TfpB orthologs from 3 heterologous T4aP systems could complement *V. cholerae* mutants lacking their endogenous motor ATPases. The relevant genotypes of strains are indicated below each bar, which includes the status of the native *V. cholerae pilT/pilU/pilB* genes (top) as well as the ectopically expressed motor ATPase allele (induced via P_{BAD}) (bottom). The color of the bar corresponds to the species from which the ectopically expressed *motor ATPase* was derived (green bars = *Vc motors*, yellow bars = *Ab* motors, orange bars = *Ps* motors, and purple bars = *Ec* motors). Data are from at least 3 independent biological replicates and shown as the mean ± SD. The limit of detection for transformation frequency assays is ~10^{-7}. In **C**, Ab_1 denotes *pilB*$_{Ab}$ and Ab_2 denotes *tfpB*$_{Ab}$. In **C**, for all samples where the transformation frequency is ~10^{-7}, at least 1 replicate contained zero transformants and the limit of detection for that sample is displayed. BACTH assays to test whether the indicated (**D**) PilT, (**E**) PilU, or (**F**) PilB/TfpB alleles can interact with themselves ("self") or the indicated T4P components (PilC or PilT). "^{WA}" denotes a Walker A mutation in the tested allele. "+" and "−" denote positive and negative controls for the BACTH assay. "E.V." denotes an empty vector. Data are representative of at least 2 independent experiments.

Fig. 2. *The PilT homolog from E. coli cannot support PilU-mediated retraction*. (A) Natural
transformation assays were performed to test whether retraction ATPase PilT mutants from 3
heterologous T4aP systems could complement *V. cholerae* mutants lacking their endogenous motor
ATPases. The relevant genotypes of strains are indicated below each bar, which includes the status of
the native *pilT/pilU* genes as well as the ectopically expressed motor ATPase allele (induced via P_{BAD}).
The color of the bar corresponds to the species from which the ectopically expressed *pilT* was derived
(green bars = *Vc pilT*, yellow bars = *Ab pilT*, orange bars = *Ps pilT*, and purple bars = *Ec pilT*). Data are
from at least 3 independent biological replicates and shown as the mean ± SD. (**B**) BACTH assays to test
whether the indicated PilT alleles can interact with themselves ("self") or PilU$_{Vc}$. "WA" denotes a mutation in
the Walker A domain of the tested allele, while "WB" denotes a mutation in the Walker B domain of the
tested allele. "+" and "−" denote positive and negative controls for the BACTH assay. "E.V." denotes an
empty vector. Data are representative of at least 2 independent experiments.

Fig. 3. *The C-terminus of PilT_{Vc} contributes to PilU-mediated retraction*. Natural transformation assays were performed to test whether retraction ATPase chimeras between PilT_{Vc} and PilT_{Ec} can complement *V. cholerae* mutants lacking their endogenous motor ATPases. PilT chimeras either have (**A**) their ATPase activity intact to assess ATPase-dependent retraction, or (**B**) their ATPase activity inactivated via Walker A or Walker B mutations to assess PilU-mediated retraction. The relevant genotypes of strains are indicated to the left of each bar, which includes the status of the native *pilT/pilU* genes as well as the ectopically expressed motor ATPase chimera (induced via P_{BAD}). Diagrams to the left of bars denote the composition of each chimera with the fragments of PilT_{Vc} denoted in green and PilT_{Ec} denoted in pink. The number in parentheses indicates the number of N-terminal amino acids from the PilT allele indicated fused to its PilT ortholog. (**C**) Natural transformation assays were performed to test whether two elements at the C-terminus of PilT_{Vc} (the AIRNLIRE domain and a 15-residue domain at the C-terminus) contribute to PilU-mediated retraction. The relevant genotypes of strains are indicated below each bar, which includes the status of the native *pilT/pilU* genes as well as the ectopically expressed motor ATPase chimera (induced via P_{BAD}). Four attributes of each ectopically expressed allele are indicated: the organism from which the PilT allele is derived, the ATPase status of the allele ("+" denotes ATPase intact, "WA" denotes Walker A mutant, and "WB" denotes Walker B mutant), the sequence of the AXXNLIRE domain (IR = AIRNLIRE, VG = AVGNLIRE), and the presence / absence of the C-terminal 15 amino acids. If the attribute changes the natural allele, it is displayed in red text. The color of the bar corresponds to the species from which the ectopically expressed *pilT* was derived (green bars = *Vc pilT* and purple bars = *Ec pilT*). All data are from at least 3 independent biological replicates and shown as the mean ± SD.

pilTU	+	Δ	Δ	Δ
P_{tac}-$pilT_{XX}$^{WA}	-	-	Vc	Ps
P_{BAD}-$pilU_{XX}$	-	-	Ps	Ps

Fig. S1. *$PilU_{Ps}$ is a fully functional retraction ATPase motor.* Natural transformation assays
were performed to assess whether $PilU_{Ps}$ can support pilus retraction in conjunction with its
cognate $PilT_{Ps}$^{WA} better than with the heterologous $PilT_{Vc}$^{WA}. The relevant genotypes of each
strain are indicated below each bar, which includes the status of the native *pilT/pilU* genes as
well as the ectopically expressed retraction motor ATPases. Data are from at least 3
independent biological replicates and shown as the mean ± SD.

Fig. S2. *Alignment of PilU and PilT sequences from the four T4aP studied.* Alignments of
(**A**) PilU orthologs and (**B**) PilT orthologs. Residues highlighted in black are identical, while
those highlighted in gray are similar. The AIRNLIRE and C-terminal 15 residue domains
experimentally tested in this study are boxed in red. The GenBank Accession numbers for the
protein sequences used are as follows: Vc, *Vibrio cholerae* PilT (AAF93635); Ab, *Acinetobacter
baylyi* PilT (CAG67808); Ps, *Pseudomonas stutzeri* PilT (AFN76400); Ec, *Escherichia coli* PilT
(NP_417425); Vc, *V. cholerae* PilU (AAF93636); Ab, *A. baylyi* PilU (CAG67807); Ps, *P. stutzeri*
PilU (AFN76401).

A

	PilT_Vc	PilT_Ab	PilT_Ps	PilT_Ec
PilT_Vc	100%	83%	80%	62%
PilT_Ab	83%	100%	91%	60%
PilT_Ps	80%	91%	100%	59%
PilT_Ec	62%	60%	59%	100%

B

	PilU_Vc	PilU_Ab	PilU_Ps
PilU_Vc	100%	70%	74%
PilU_Ab	70%	100%	78%
PilU_Ps	74%	78%	100%

C

full length

	PilB_Vc	PilB_Ab	TfpB_Ab	PilB_Ps
PilB_Vc	100%	64%	53%	66%
PilB_Ab	64%	100%	55%	75%
TfpB_Ab	53%	55%	100%	54%
PilB_Ps	66%	75%	54%	100%

CTD

	PilB_Vc	PilB_Ab	TfpB_Ab	PilB_Ps
PilB_Vc	100%	75%	64%	75%
PilB_Ab	75%	100%	64%	86%
TfpB_Ab	64%	64%	100%	61%
PilB_Ps	75%	86%	61%	100%

NTD

	PilB_Vc	PilB_Ab	TfpB_Ab	PilB_Ps
PilB_Vc	100%	41%	32%	49%
PilB_Ab	41%	100%	38%	52%
TfpB_Ab	32%	38%	100%	41%
PilB_Ps	49%	52%	41%	100%

Fig. S3. *Pairwise comparisons of the indicated motor ATPases.* Percent similarities are reported for pairwise comparisons of the indicated (**A**) PilT, (**B**) PilU, and (**C**) PilB/TfpB orthologs. PilB/TfpB orthologs were further analyzed for pairwise comparisons of the C-terminal domains (CTD) and N-terminal domains (NTD) (see **Fig. S4** for details on what portion of PilB represents the NTD vs CTD). The GenBank Accession numbers for the protein sequences used in these comparisons are as follows: Vc, *Vibrio cholerae* PilT (AAF93635); Ab, *Acinetobacter baylyi* PilT (CAG67808); Ps, *Pseudomonas stutzeri* PilT (AFN76400); Ec, *Escherichia coli* PilT (NP_417425); Vc, *V. cholerae* PilU (AAF93636); Ab, *A. baylyi* PilU (CAG67807); Ps, *P. stutzeri* PilU (AFN76401); Vc, *V. cholerae* PilB (AAF95567); Ab, *A. baylyi* PilB (CAG67312); Ab, *A. baylyi* TfpB (CAG69034); Ps, *P. stutzeri* PilB (AFN76867).

```
TfpB Ab   342  TGPTGSGKTTTLYSSLKQLATD-QVNVCTIEDPIDMLESS----FNQMQVNAAIDLGFAD
PilB Ps   326  TGPTGSGKLVSLYTGLNILNTV-DVNISTAEDPVPINLEG----INQVNVNPRQGVDFAQ
                                              CTD
PilT Vc   188  ALRSALREDPDVIILVGELRDQETISLALTAAETGHLVFGTLHTSSAAKTIDRIIDVFPGS
PilT Ab   188  ALRSALREDPDIIILVGEMRDLETIRLALTAAETGHLVFGTLHTTSAAKTIDRIIDVFPAE
PilT Ps   188  ALRSALREDPDIIILVGEMRDLETIRLALTAAETGHLVFGTLHTTSAAKTIDRVVDVFPAE
PilT Ec   185  GLRAALREDPDVILLGELRDSETIRLALTAAETGHLVIATLHTRGAAQAVERLVDSFPAQ
PilB Vc   378  ALRSFLRQDPDVVMVGEIRDLETAEIAIVKAAQTGHLVISTLHTNSAAETVIRIANMGVE-
PilB Ab   391  ALRSFLRQDPDIIMVGEIRDLETAEIAIKAAQTGHLVMSTLHTNNAAETLTRIRNMGVA-
TfpB Ab   397  GVRALMRQDPDIIMVGEIRDQDTANVAIQAALTGHLVISTLHTNDAPSSLTRIHDLGVQ-
PilB Ps   381  ALRAFLRQDPDVIMVGEIRDLETAEIAIKAAQTGHMVVSTLHTNSAAETLTRIRNMGVA-
                                              CTD
PilT Vc   248  DKDMVRSMLSESLRAVIAQKLLKRV------------------------------------
PilT Ab   248  EKDMVRTMLSESLQAVISQTLLKKE------------------------------------
PilT Ps   248  EKSMVRSMLSESLQAVISQTLLKKI------------------------------------
PilT Ec   245  EKDPVRNQLAGSLRAVISQKLEVDK-----------------------------------
PilB Vc   437  -----PFNLASSLSLIIAQRLARRLCKHCKIAVR--PSALLQSQFAFQPN-E-ILYEANA
PilB Ab   450  -----SFNIATSVNLVIAQRLARRLCSQCKRPIQVPERSLLEMGFTPEDLAQPEFQIFEP
TfpB Ab   456  -----PFLTAATLLGVIAQRLVRQLCPHCKVLSDIDKQQWQHLT-QDYEIAM-PEQIYGP
PilB Ps   440  -----AFNIATSVNLIIAQRLARRLCSSCKKEVEVPRDALLKEGFPEESI-G-TFKIYGP
                                              CTD
PilT Vc   273  ----------GGGRVACHEIMIATPAIRNLIREDKVAQVYSIIQTGAAHGMQTMEQNAKQ
PilT Ab   273  ----------GGGRVAAHEIMIGIPAIRNLIREGKVAQVYSSIQTGATHGMTTLDQSLKQ
PilT Ps   273  ----------GGGRVAAHEIMIGTPAIRNLIREDKVAQVYSAIQTGGSLGMQTLDMCLKG
PilT Ec   270  ----------QEGRVALFELLINTPAVGNLIREGKTHQLPHVIQTGQQVGMITFQQSYQH
PilB Vc   488  AGCNECT-GGYSGRVGIYEVMAFNTELAEAIMQRASI--HQIERLAKANGMQTLQESGLE
PilB Ab   505  VGCHDCR-EGYKGRVGIYEVMKITPFISKIIMEDGNA--LEIAATAETLGFNNLRRSGLK
TfpB Ab   509  VGCDHCRQTGYKGRIGIYEFMSLDNLKQLVSSPANL--NQLKSEAKKQGICPLRIAGAR
PilB Ps   493  VGCENCV-GGYKGRVGIYEVVKNTPALARIIMEDGNS--IDISTQMRKDGFNDLRTSALI
                                              CTD
PilT Vc   323  LTARGVVDAQEVQSKIELDLKAF
PilT Ab   323  LLAKGTINLPTARAVA-KSPESF
PilT Ps   323  LLAKGLISRDSAREKA-KQPENF
PilT Ec   320  RVGEGRL----------------
PilB Vc   545  KIREGITSFAELQRVLY-----F
PilB Ab   562  KVMQGVTSLQEINRVTS-----E
TfpB Ab   567  KILEGVTSLEEVLRVVP-L---T
PilB Ps   550  KAMQGVTSLEEVNRVTK-----D
```

Fig. S4. *Alignment of the PilB/TfpB and PilT sequences from the T4aP studied.* Alignments
of PilB/TfpB orthologs and PilT orthologs. Residues highlighted in black are identical, while
those highlighted in gray are similar. The N-terminal domain and C-terminal domain of PilB are
denoted by the "NTD" and "CTD" bars above the sequence. The GenBank Accession numbers
for the protein sequences used are as follows: Vc, *Vibrio cholerae* PilT (AAF93635); Ab,
Acinetobacter baylyi PilT (CAG67808); Ps, *Pseudomonas stutzeri* PilT (AFN76400); Ec,
Escherichia coli PilT (NP_417425); Vc, *V. cholerae* PilB (AAF95567); Ab, *A. baylyi* PilB
(CAG67312); Ab, *A. baylyi* TfpB (CAG69034); Ps, *P. stutzeri* PilB (AFN76867).

 Table S1. Strains used in this study

Strain name in manuscript	Genotype and antibiotic resistance	Figure	Strain #
Parent	E7946 SmR, ΔluxO::miniFRT, lacZ::lacIq, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*	Fig. 1, 2, 3, S1	TND0905 (SAD2468)
Δ*pilT*	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔpilT::TmR	Fig. 1, 2, 3	TND1035 (SAD2469)
Δ*pilTU*	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔpilTU::TmR	Fig. 1, 2, 3, S1	JLC227 (SAD2470)
P_{BAD}-*pilT*$_{Vc}$, Δ*pilT*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Vc}$ CarbR, ΔpilT::TmR	Fig. 1, 3	TND1831 (SAD3199)
P_{BAD}-*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 1, 3	JLC1155 (SAD3222)
P_{BAD}-*pilT*$_{Ab}$, Δ*pilT*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ab}$ CarbR, ΔpilT::TmR	Fig. 1	TND2148 (SAD3200)
P_{BAD}-*pilT*$_{Ab}$, Δ*pilTU*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ab}$ CarbR, ΔpilTU::TmR	Fig. 1	JLC1153 (SAD3223)
P_{BAD}-*pilT*$_{Ps}$, Δ*pilT*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ps}$ CarbR, ΔpilT::TmR	Fig. 1	TND2125 (SAD3201)
P_{BAD}-*pilT*$_{Ps}$, Δ*pilTU*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ps}$ CarbR, ΔpilTU::TmR	Fig. 1	JLC1157 (SAD3224)
P_{BAD}-*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 1	JLC1473 (SAD3225)
P_{BAD}-*pilT*$_{Ec}$, Δ*pilTU*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 1	JLC1474 (SAD3226)
pilT^{WA}, Δ*pilU*	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, pilT^{K136A}, ΔpilU::TmR	Fig. 1, 2	JLC444 (SAD2472)
P_{BAD}-*pilU*$_{Vc}$, *pilT*^{WA}, Δ*pilU*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}$^{K136A}, ΔlacZ:: P_{BAD}-PilU$_{Vc}$ CarbR, ΔpilU::TmR	Fig. 1	EHC062 (SAD3239)
P_{BAD}-*pilU*$_{Ab}$, *pilT*^{WA}, Δ*pilU*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}$^{K136A}, ΔlacZ:: P_{BAD}-PilU$_{Ab}$ CarbR, ΔpilU::TmR	Fig. 1	EHC060 (SAD3240)
P_{BAD}-*pilU*$_{Ps}$, *pilT*^{WA}, Δ*pilU*	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}$^{K136A}, ΔlacZ:: P_{BAD}-PilU$_{Ps}$ CarbR, ΔpilU::TmR	Fig. 1	EHC061 (SAD3241)
Parent (P_{const}-*tfoX*)	E7946 SmR, ΔlacZ::lacIq, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C	Fig. 1	TND1076 (SAD2513)
Δ*pilB*	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT	Fig. 1	TND2373 (SAD2787)

P_{BAD}-$pilB_{Vc}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P_{BAD}-pilB$_{Vc}$ CarbR	Fig. 1	TND2403 (SAD2789)
P_{BAD}-$pilB_{Ab}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P_{BAD}-pilB$_{Ab}$ CarbR	Fig. 1	TND2354 (SAD3202)
P_{BAD}-$tfpB_{Ab}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P_{BAD}-tfpB$_{Ab}$ CarbR	Fig. 1	TND2372 (SAD3203)
P_{BAD}-$pilB_{Ps}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P_{BAD}-pilB$_{Ps}$ CarbR	Fig. 1	TND2371 (SAD3204)
$pilT^{WA}$	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT^{K136A}	Fig. 2	TND1077 (SAD2471)
P_{BAD}-$pilT_{Vc}^{WA}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Vc}^{K136A}$ CarbR, ΔpilT::TmR	Fig. 2, 3	JLC1609 (SAD3227)
P_{BAD}-$pilT_{Vc}^{WA}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Vc}^{K136A}$ CarbR, ΔpilTU::TmR	Fig. 2	EHC031 (SAD3242)
P_{BAD}-$pilT_{Vc}^{WB}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Vc}^{E204A}$ CarbR, ΔpilT::TmR	Fig. 2	EHC075 (SAD3243)
P_{BAD}-$pilT_{Ab}^{WA}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ab}^{K136A}$ CarbR, ΔpilT::TmR	Fig. 2	EHC015 (SAD3244)
P_{BAD}-$pilT_{Ab}^{WA}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ab}^{K136A}$ CarbR, ΔpilTU::TmR	Fig. 2	EHC019 (SAD3245)
P_{BAD}-$pilT_{Ps}^{WA}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ps}^{K136A}$ CarbR, ΔpilT::TmR	Fig. 2	EHC016 (SAD3246)
P_{BAD}-$pilT_{Ps}^{WA}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ps}^{K136A}$ CarbR, ΔpilTU::TmR	Fig. 2	EHC020 (SAD3247)
P_{BAD}-$pilT_{Ec}^{WA}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ec}^{K133A}$ CarbR, ΔpilT::TmR	Fig. 2	JLC1499 (SAD3228)
P_{BAD}-$pilT_{Ec}^{WA}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ec}^{K133A}$ CarbR, ΔpilTU::TmR	Fig. 2	JLC1498 (SAD3229)
P_{BAD}-$pilT_{Ec}^{WB}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ec}^{E201A}$ CarbR, ΔpilT::TmR	Fig. 2, 3	EHC076 (SAD3248)
P_{BAD}-$pilT_{Vc}$(50)/$pilT_{Ec}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}$(50)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3069 (SAD3205)
P_{BAD}-$pilT_{Vc}$(100)/$pilT_{Ec}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}$(100)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3070 (SAD3206)
P_{BAD}-$pilT_{Vc}$(150)/$pilT_{Ec}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}$(150)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3071 (SAD3207)

P$_{BAD}$-*pilT*$_{Vc}$(200)/*pilT*$_{Ec}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Vc}$(200)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3073 (SAD3208)
P$_{BAD}$-*pilT*$_{Vc}$(300)/*pilT*$_{Ec}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Vc}$(300)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3072 (SAD3209)
P$_{BAD}$-*pilT*$_{Ec}$(50)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Ec}$(50)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3074 (SAD3210)
P$_{BAD}$-*pilT*$_{Ec}$(100)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Ec}$(100)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3075 (SAD3211)
P$_{BAD}$-*pilT*$_{Ec}$(150)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Ec}$(150)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3076 (SAD3212)
P$_{BAD}$-*pilT*$_{Ec}$(200)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Ec}$(200)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3077 (SAD3213)
P$_{BAD}$-*pilT*$_{Ec}$(250)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Ec}$(250)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3078 (SAD3214)
P$_{BAD}$-*pilT*$_{Ec}$(300)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Ec}$(300)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3079 (SAD3215)
P$_{BAD}$-*pilT*$_{Vc}$^{WA}(100)/*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Vc}$^{K136A}(100)/pilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 3	TND3008 (SAD3216)
P$_{BAD}$-*pilT*$_{Vc}$^{WA}(200)/*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Vc}$^{K136A}(200)/pilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 3	TND3010 (SAD3217)
P$_{BAD}$-*pilT*$_{Ec}$^{WA}(300)/*pilT*$_{Vc}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Ec}$^{K133A}(300)/pilT$_{Vc}$ CarbR, ΔpilT::TmR	Fig. 3	TND3018 (SAD3218)
P$_{BAD}$-*pilT*$_{Vc}$^{WB}(100)/*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Vc}$^{E204A}(100)/pilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 3	EHC106 (SAD3249)
P$_{BAD}$-*pilT*$_{Vc}$^{WB}(200)/*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Vc}$^{E204A}(200)/pilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 3	EHC108 (SAD3250)
P$_{BAD}$-*pilT*$_{Ec}$^{WB}(300)/*pilT*$_{Vc}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-3X Flag-pilT$_{Ec}$^{E201A}(300)/pilT$_{Vc}$ CarbR, ΔpilT::TmR	Fig. 3	EHC115 (SAD3251)
P$_{BAD}$-*pilT*$_{Vc}$$^{AIR→AVG}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-pilT$_{Vc-}$$^{AIR→AVG}$ CarbR, ΔpilTU::TmR	Fig. 3	JLC1622 (SAD3230)
P$_{BAD}$-*pilT*$_{Vc}$$^{WA, AIR→AVG}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-pilT$_{Vc}$^{K136A,} $^{AIR→AVG}$ CarbR, ΔpilT::TmR	Fig. 3	JLC1625 (SAD3231)
P$_{BAD}$-*pilT*$_{Vc}$^{Δ15aa}, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-pilT$_{Vc}$^{Δ15aa} CarbR, ΔpilTU::TmR	Fig. 3	TND2774 (SAD3219)
P$_{BAD}$-*pilT*$_{Vc}$$^{WA, Δ15aa}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P$_{tac}$-*tfoX*, ΔlacZ:: P$_{BAD}$-pilT$_{Vc}$^{K136A,} ^{Δ15aa} CarbR, ΔpilT::TmR	Fig. 3	JLC1611 (SAD3232)

P_{BAD}-$pilT_{Vc}$[AIR→AVG, Δ15aa], $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Vc}$-[AIR→AVG, Δ15aa] CarbR, ΔpilTU::TmR	Fig. 3	JLC1624 (SAD3233)
P_{BAD}-$pilT_{Vc}$[WA, AIR→AVG, Δ15aa], $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Vc}$[K136A, AIR→AVG, Δ15aa] CarbR, ΔpilT::TmR	Fig. 3	JLC1627 (SAD3234)
P_{BAD}-$pilT_{Ec}$[AVG→AIR], $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-[AVG→AIR] CarbR, ΔpilTU::TmR	Fig. 3	JLC1614 (SAD3235)
P_{BAD}-$pilT_{Ec}$[WA, AVG→AIR], $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$[K133A, AVG→AIR] CarbR, ΔpilT::TmR	Fig. 3	JLC1615 (SAD3236)
P_{BAD}-$pilT_{Ec}$[WB, AVG→AIR], $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$[E201A, AVG→AIR] CarbR, ΔpilT::TmR	Fig. 3	EHC095 (SAD3252)
P_{BAD}-$pilT_{Ec}$[+15aa], $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$[+15aa] CarbR, ΔpilTU::TmR	Fig. 3	TND2730 (SAD3220)
P_{BAD}-$pilT_{Ec}$[WA, +15aa], $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$[K133A, +15aa] CarbR, ΔpilT::TmR	Fig. 3	TND2727 (SAD3221)
P_{BAD}-$pilT_{Ec}$[WB, +15aa], $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$[E201A, +15aa] CarbR, ΔpilT::TmR	Fig. 3	EHC093 (SAD3253)
P_{BAD}-$pilT_{Ec}$[AVG→AIR, +15aa], $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-[AVG→AIR, +15aa] CarbR, ΔpilTU::TmR	Fig. 3	JLC1620 (SAD3237)
P_{BAD}-$pilT_{Ec}$[WA, AVG→AIR, +15aa], $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$[K133A, AVG→AIR, +15aa] CarbR, ΔpilT::TmR	Fig. 3	JLC1617 (SAD3238)
P_{BAD}-$pilT_{Ec}$[WB, AVG→AIR, +15aa], $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$[E201A, AVG→AIR, +15aa] CarbR, ΔpilT::TmR	Fig. 3	EHC097 (SAD3254)
P_{tac}-$pilT_{Vc}$[WA], P_{BAD}-$pilU_{Ps}$, $\Delta pilTU$	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, ΔlacZ::P_{BAD}-PilU$_{Ps}$ CarbR, ΔVC0692::P_{tac}-PilT$_{Vc}$[K136A] SpecR, ΔpilTU::TmR	Fig. S1	EHC099 (SAD3255)
P_{tac}-$pilT_{Ps}$[WA], P_{BAD}-$pilU_{Ps}$, $\Delta pilTU$	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, ΔlacZ::P_{BAD}-PilU$_{Ps}$ CarbR, ΔVC0692::P_{tac}-PilT$_{Ps}$[K136A] SpecR, ΔpilTU::TmR	Fig. S1	EHC100 (SAD3256)
T25-PilT$_{Vc}$	TG1, pkT25-PilT$_{Vc}$ KanR	Fig. 1, 2	SAD2248
T25-PilT$_{Ab}$	TG1, pkT25-PilT$_{Ab}$ KanR	Fig. 1, 2	SAD2659
T25-PilT$_{Ps}$	TG1, pkT25-PilT$_{Ps}$ KanR	Fig. 1, 2	EHC043 (SAD3257)
T25-PilT$_{Ec}$	TG1, pkT25-PilT$_{Ec}$ KanR	Fig. 1, 2	EHC047 (SAD3258)
T18-PilT$_{Vc}$	TG1, pUT18C-PilT$_{Vc}$ CarbR	Fig. 1, 2	SAD2249
T18-PilT$_{Ab}$	TG1, pUT18C-PilT$_{Ab}$ CarbR	Fig. 1, 2	SAD2660
T18-PilT$_{Ps}$	TG1, pUT18C-PilT$_{Ps}$ CarbR	Fig. 1, 2	EHC044 (SAD3259)
T18-PilT$_{Ec}$	TG1, pUT18C-PilT$_{Ec}$ CarbR	Fig. 1, 2	EHC042 (SAD3260)
T25-PilU$_{Vc}$	TG1, pkT25-PilU$_{Vc}$ KanR	Fig. 1, 2	SAD2250

T18-PilU$_{Vc}$	TG1, pUT18C-PilU$_{Vc}$ CarbR	Fig. 1, 2	SAD2251
T25-PilC$_{Vc}$	TG1, pkT25-PilC$_{Vc}$ KanR	Fig. 1	SAD2260
T18-PilC$_{Vc}$	TG1, pUT18C-PilC$_{Vc}$ CarbR	Fig. 1	SAD2261
T25-PilU$_{Ab}$	TG1, pkT25-PilU$_{Ab}$ KanR	Fig. 1	EHC053 (SAD3261)
T25-PilU$_{Ps}$	TG1, pkT25-PilU$_{Ps}$ KanR	Fig. 1	EHC048 (SAD3262)
T18-PilU$_{Ab}$	TG1, pUT18C-PilU$_{Ab}$ CarbR	Fig. 1	SAD2664
T18-PilU$_{Ps}$	TG1, pUT18C-PilU$_{Ps}$ CarbR	Fig. 1	EHC064 (SAD3263)
T25-PilB$_{Vc}$	TG1, pkT25-PilB$_{Vc}$ KanR	Fig. 1	EHC052 (SAD3264)
T25-PilB$_{Ab}$	TG1, pkT25-PilB$_{Ab}$ KanR	Fig. 1	SAD2647
T25-TfpB$_{Ab}$	TG1, pkT25-TfpB$_{Ab}$ KanR	Fig. 1	SAD2878
T25-PilB$_{Ps}$	TG1, pkT25-PilB$_{Ps}$ KanR	Fig. 1	SAD2959
T18-PilB$_{Vc}$	TG1, pUT18C-PilB$_{Vc}$ CarbR	Fig. 1	EHC050 (SAD3265)
T18-PilB$_{Ab}$	TG1, pUT18C-PilB$_{Ab}$ CarbR	Fig. 1	SAD2648
T18-TfpB$_{Ab}$	TG1, pUT18C-TfpB$_{Ab}$ CarbR	Fig. 1	SAD2879
T18-PilB$_{Ps}$	TG1, pUT18C-PilB$_{Ps}$ CarbR	Fig. 1	SAD2960
T25-PilT$_{Vc}$^{WA}	TG1, pkT25-PilT$_{Vc}$^{K136A} KanR	Fig. 1, 2	SAD2285
T25-PilT$_{Ab}$^{WA}	TG1, pkT25-PilT$_{Ab}$^{K136A} KanR	Fig. 2	EHC054 (SAD3266)
T25-PilT$_{Ps}$^{WA}	TG1, pkT25-PilT$_{Ps}$^{K136A} KanR	Fig. 2	EHC056 (SAD3267)
T25-PilT$_{Ec}$^{WA}	TG1, pkT25-PilT$_{Ec}$^{K133A} KanR	Fig. 2	EHC057 (SAD3268)
T18-PilT$_{Vc}$^{WA}	TG1, pUT18C-PilT$_{Vc}$^{K136A} CarbR	Fig. 1, 2	SAD2286
T18-PilT$_{Ab}$^{WA}	TG1, pUT18C-PilT$_{Ab}$^{K136A} CarbR	Fig. 2	EHC055 (SAD3269)
T18-PilT$_{Ps}$^{WA}	TG1, pUT18C-PilT$_{Ps}$^{K136A} CarbR	Fig. 2	EHC059 (SAD3270)
T18-PilT$_{Ec}$^{WA}	TG1, pUT18C-PilT$_{Ec}$^{K133A} CarbR	Fig. 2	EHC058 (SAD3271)
T25-PilT$_{Vc}$^{WB}	TG1, pkT25-PilT$_{Vc}$^{E204A} KanR	Fig. 2	SAD2287
T25-PilT$_{Ec}$^{WB}	TG1, pkT25-PilT$_{Ec}$^{E201A} KanR	Fig. 2	EHC078 (SAD3272)
T18-PilT$_{Vc}$^{WB}	TG1, pUT18C-PilT$_{Vc}$^{E204A} CarbR	Fig. 2	SAD2288
T18-PilT$_{Ec}$^{WB}	TG1, pUT18C-PilT$_{Ec}$^{E201A} CarbR	Fig. 2	EHC079 (SAD3273)
T25-Zip	TG1, pkT25-leucine zipper KanR	Fig. 1, 2	SAD2238
T18-Zip	TG1, pUT18C-leucine zipper CarbR	Fig. 1, 2	SAD2241
T25-vector	TG1, pkT25 vector KanR	Fig. 1, 2	SAD2236
T18-vector	TG1, pUT18C vector CarbR	Fig. 1, 2	SAD2240
Acinetobacter baylyi	ADP1	Purified gDNA	SAD631
Pseudomonas stutzeri	JM300	Purified gDNA	SAD2178
Escherichia coli	MG1655	Purified gDNA	SAD720

 Table S2. Primers used in this study

Primer name	Sequence (5' to 3')	Description
BBC374	TGGCAAAAAGCGAGAGAAGAAG	ΔluxO F1
BBC375	gtcgacggatccccggaatCATGAGGACATATTTTGTTTTCTGC	ΔluxO R1
BBC376	gaagcagctccagcctacaTAAGCGATGAGAGAATGGATCAAC	ΔluxO F2
BBC377	TCACACCCGAATTTCCATCATGC	ΔluxO R2
ABD344	GATTAGCAACGATTCTAGCGCAGGAG	ΔVC1807 F1
ABD340	gtcgacggatccccggaatACGTTTCATTAGTCACCTCTATTGT TAACTTGTTC	ΔVC1807 R1
ABD341	gaagcagctccagcctacaTAGTCGAAAATAAAAAAAGAGG CTCGCCTC	ΔVC1807 F2
ABD345	CTTGCTAACCGTTGGTGTTACCAGC	ΔVC1807 R2
DOG0400	ACTTCTGGCTGAAGGTCAATTTTC	Amplify ΔpilT/U mutations F1
DOG0403	TCACGTGTTCGGCCAAAATC	Amplify ΔpilT/U mutations R2
ABD332	GGCTGAACGTGGTTGTCGAAAATGAC	P_{BAD}-XX @ lacZ F1
ABD256	CCCAAATACGGCAACTTGGCG	P_{BAD}-XX @ lacZ R2
BBC832	GCTTTTTGCTACAACGACCG	P_{tac}-XX @ VCA0692 F1
BBC834	CACAATTTCTCGCTTAAAATGTCC	P_{tac}-XX @ VCA0692 R2
BBC1723	caatttcacacaggatcccgggAGGAGGTATTTAAATGGATATCGCTGAGTTAC	Amplify $pilT_{Vc}$ for P_{BAD}-pilT or P_{tac}-pilT F
BBC1724	tgtaggctggagctgcttcCTAAAATGCTTTTAAATCCAACTCG	Amplify $pilT_{Vc}$ for P_{BAD}-pilT or P_{tac}-pilT R
BBC2807	caatttcacacaggatcccgggAGGAGGTGAAAATATGGATATAACAGAATTACTCG	Amplify $pilT_{Ab}$ for P_{BAD}-pilT F
BBC2808	tgtaggctggagctgcttcTTAGAACGACTCTGGAGATTTTG	Amplify $pilT_{Ab}$ for P_{BAD}-pilT R
BBC3143	caatttcacacaggatcccgggAGGAGGTAGTCCCATGGATATCACAGAAC	Amplify $pilT_{Ps}$ for P_{BAD}-pilT or P_{tac}-pilT F
BBC3144	tgtaggctggagctgcttcTTATCGGTTGTTAAAAATTTTCCGG	Amplify $pilT_{Ps}$ for P_{BAD}-pilT or P_{tac}-pilT R
BBC3140	caatttcacacaggatcccgggAGGAGGTCTTAGTATGAATATGGAAGAAATTGTGG	Amplify $pilT_{Ec}$ for P_{BAD}-pilT F
BBC3141	tgtaggctggagctgcttcCATGGCAACGTCACAAACG	Amplify $pilT_{Ec}$ for P_{BAD}-pilT R
HPQ0001	GGGCTCGGGTgctTCGACGACCTTGGCG	$pilT_{Vc}^{K136A}$ R1
HPQ0002	CGCCAAGGTCGTCGAagcACCCGAGCCC	$pilT_{Vc}^{K136A}$ F2
BBC2403	GCTGCAAGGGTGGTTGAagcACCGGAACCAGTGG	$pilT_{Ab}^{K136A}$ R1
BBC2404	CCACTGGTTCCGGTgctTCAACCACCCTTGCAGC	$pilT_{Ab}^{K136A}$ F2
JCP135	CGCCAGAGTGGTGGAcgcGCCCGAACCGGTCGG	$pilT_{Ps}^{K136A}$ R1
JCP134	CCGACCGGTTCGGGCgcgTCCACCACTCTGGCG	$pilT_{Ps}^{K136A}$ F2
JCP132	CGCCAGCGTGGTAGAcgcGCCACTCCCCGTCGC	$pilT_{Ec}^{K133A}$ R1
JCP131	GCGACGGGGAGTGGCgcgTCTACCACGCTGGCG	$pilT_{Ec}^{K133A}$ F2
BBC2508	GTTTCTTGGTCACGCAGagcACCGACGAGGATCAC	$pilT_{Vc}^{E204A}$ R1

BBC2509	GTGATCCTCGTCGGTGctCTGCGTGACCAAGAAAC	$pilT_{Vc}^{E204A}$ F2
BBC4065	CTCGCTGTCACGCAGagCTCCGAGCAAAATCACATCAGGATCTTCCCG	$pilT_{Ec}^{E201A}$ R1
BBC4066	GCTCGGAGctCTGCGTGACAGCGAGACAATCCGTCTGGCACTGAC	$pilT_{Ec}^{E201A}$ F2
BBC3897	CAAGCGATGTACATCAGAATGGG	$pilT_{Vc}$(50)/$pilT_{Ec}$ R1
BBC3898	cccattctgatgtacatcgcttgCTGCGGGAGTGGCTGGATG	$pilT_{Vc}$(50)/$pilT_{Ec}$ F2
BBC3899	GGGAATGGTACGAAACACCGCC	$pilT_{Vc}$(100)/$pilT_{Ec}$ R1
BBC3900	ggcggtgtttcgtaccattcccTCGCACTGCCCGCAGCTCGAAC	$pilT_{Vc}$(100)/$pilT_{Ec}$ F2
BBC3901	ATGAGCATTGACATAGTTCACC	$pilT_{Vc}$(150)/$pilT_{Ec}$ R1
BBC3902	ggtgaactatgtcaatgctcatGCCGATGCGCATATTCTGACG	$pilT_{Vc}$(150)/$pilT_{Ec}$ F2
BBC3903	GATCACATCTGGGTCTTCACGC	$pilT_{Vc}$(200)/$pilT_{Ec}$ R1
BBC3904	gcgtgaagacccagatgtgatcTTGCTCGGAGAGCTGCGTG	$pilT_{Vc}$(200)/$pilT_{Ec}$ F2
BBC3907	GTCTTTGTCGCTGCCCGGAAAAAC	$pilT_{Vc}$(300)/$pilT_{Ec}$ R1
BBC3908	gttttccgggcagcgacaaagacCCCGTGCGTAATCAACTGGC	$pilT_{Vc}$(300)/$pilT_{Ec}$ F2
BBC3909	CTGCGCCACTTTATCTTCACGG	$pilT_{Ec}$(50)/$pilT_{Vc}$ R1
BBC3910	ccgtgaagataaagtggcgcagTTACCGCATGTTATTCAAACC	$pilT_{Ec}$(50)/$pilT_{Vc}$ F2
BBC3911	CTCCCGCAGTAGCTCTTCG	$pilT_{Ec}$(100)/$pilT_{Vc}$ R1
BBC3912	cgaagagctactgcgggagATCATGAATGATGCCCAGCGC	$pilT_{Ec}$(100)/$pilT_{Vc}$ F2
BBC3913	GCAGTGCGAAGGTAACAGCCG	$pilT_{Ec}$(150)/$pilT_{Vc}$ R1
BBC3914	cggctgttaccttcgcactgcCCGACGCTAGAGCAACTGG	$pilT_{Ec}$(150)/$pilT_{Vc}$ F2
BBC3915	CGCATCGGCATGTTGATTGAG	$pilT_{Ec}$(200)/$pilT_{Vc}$ R1
BBC3916	ctcaatcaacatgccgatgcgCATATTTTGACCATTGAAGATCCG	$pilT_{Ec}$(200)/$pilT_{Vc}$ F2
BBC3917	TCCGAGCAAAATCACATCAGG	$pilT_{Ec}$(250)/$pilT_{Vc}$ R1
BBC3918	cctgatgtgattttgctcggaGAGCTGCGTGACCAAGAAACG	$pilT_{Ec}$(250)/$pilT_{Vc}$ F2
BBC3919	ACGCACtGGGTCTTTTTCC	$pilT_{Ec}$(300)/$pilT_{Vc}$ R1
BBC3920	ggaaaaagacccagtgcgtTCCATGTTGTCGGAATCGC	$pilT_{Ec}$(300)/$pilT_{Vc}$ F2
DOG0565	AGATTGGTCTATTTTATGGCTCAAG	ΔpilB F1
DOG0570	gtcgacggatccccggaatCATTAGTTAATTGTTGCACCTGC	ΔpilB R1
DOG0571	gaagcagctccagcctacaCAGCGTGTGCTCTACTTTTAACTG	ΔpilB F2
DOG0572	AAAGAGTTGATGTTGAATCCCATC	ΔpilB R2
BBC1725	caatttcacacaggatcccgggAGGAGGTAGACAGATGGAGTTGAATCAATATC	Amplify $pilU_{Vc}$ for P_{BAD}-pilU F
BBC1726	tgtaggctggagctgcttcCTACTCCATATCAATCTTCACATTC	Amplify $pilU_{Vc}$ for P_{BAD}-pilU R
BBC2809	caatttcacacaggatcccgggAGGAGGTAATACGATGGACTTTAATGATTTACTC	Amplify $pilU_{Ab}$ for P_{BAD}-pilU F
BBC2810	tgtaggctggagctgcttcTTATTTGAGATTACGGCGTTGAC	Amplify $pilU_{Ab}$ for P_{BAD}-pilU R
JCP177	caatttcacacaggatcccgggAGGAGGTACGACGATGGAATTCGAGAAAC	Amplify $pilU_{Ps}$ for P_{BAD}-pilU F
JCP178	tgtaggctggagctgcttcGGCCTGTCAGCGGAAACTG	Amplify $pilU_{Ps}$ for P_{BAD}-pilU R
ABD334	AGTGCTCCGGACTCTTTGCTCTG	P_{BAD}-pilU @ lacZ F1
BBC1747	AACATCCGTTAGACAATGACC	P_{BAD}-pilU @ lacZ R2
BBC1952	caatttcacacaggatcccgggAGGAGGTTAACTAATGCTCACCAACCTTGTTG	Amplify $pilB_{Vc}$ for P_{BAD}-pilB F
BBC1953	tgtaggctggagctgcttcTTAAAAGTAGAGCACACGCTG	Amplify $pilB_{Vc}$ for P_{BAD}-pilB R
BBC3320	caatttcacacaggatcccgggaggaggtATTATAATGATATCAGAATCACAAGGCGG	Amplify $pilB_{Ab}$ for P_{BAD}-pilB F
BBC3310	tgtaggctggagctgcttcTTATTCACTGGTTACACGATTAATTTCC	Amplify $pilB_{Ab}$ for P_{BAD}-pilB R

BBC3327	caatttcacacaggatcccgggaggaggtAAATAAATGACATATCATTTTGAAATAGATA	Amplify tfpB$_{Ab}$ for P$_{BAD}$-tfpB F
BBC3328	tgtaggctggagctgcttcCTAAGTTAATGGGACCACCC	Amplify tfpB$_{Ab}$ for P$_{BAD}$-tfpB R
BBC3324	caatttcacacaggatcccgggaggaggtAACCTCATGAGCGAAAACATTTCCC	Amplify pilB$_{Ps}$ for P$_{BAD}$-pilB F
BBC3325	tgtaggctggagctgcttcTTAGTCCTTTGTCACCCGG	Amplify pilB$_{Ps}$ for P$_{BAD}$-pilB R
BBC2513	gaagcagctccagcctacaGATCCCCGGGTACCTAAGTAAC	Amplify BACTH pKT25-X to make T25-XX vectors F
BBC2512	tccacctgctccacctgcCTCTAGAGTCGACCCTGCAG	Amplify BACTH pKT25-X to make T25-XX vectors R
BBC2515	gaagcagctccagcctacaCCGAGCTCGAATTCATCGATATAAC	Amplify BACTH pUT18C to make T18-XX vectors F
BBC2514	tccacctgctccacctgcCTCTAGAGTCGACCTGCAG	Amplify BACTH pUT18C to make T18-XX vectors R
BBC2518	gcaggtggagcaggtggaAAAGCGACCCAAACCTTACC	BACTH TXX-PilC$_{Vc}$ F
BBC2524	tgtaggctggagctgcttcTTAACCCAATACACTCATTAAGTTAAAG	BACTH TXX-PilC$_{Vc}$ R
JCP027	gcaggtggagcaggtggaGATATCGCTGAGTTACTGGAATTTAG	BACTH TXX-PilT$_{Vc}$ F
BBC1724	tgtaggctggagctgcttcCTAAAATGCTTTTAAATCCAACTCG	BACTH TXX-PilT$_{Vc}$ R
BBC3194	gcaggtggagcaggtggaGATATAACAGAATTACTCGCGTT	BACTH TXX-PilT$_{Ab}$ F
BBC3195	tgtaggctggagctgcttcTTAGAACGACTCTGGAGATTTTG	BACTH TXX-PilT$_{Ab}$ R
BBC3921	gcaggtggagcaggtggaGATATCACAGAACTGCTGGCC	BACTH TXX-PilT$_{Ps}$ F
BBC3922	tgtaggctggagctgcttcTTAAAAATTTTCCGGCTGCTTGG	BACTH TXX-PilT$_{Ps}$ R
BBC3923	gcaggtggagcaggtggaAATATGGAAGAAATTGTGGCCC	BACTH TXX-PilT$_{Ec}$ F
BBC3924	tgtaggctggagctgcttcTCACAAACGCCCTTCCCCCAC	BACTH TXX-PilT$_{Ec}$ R
BBC2638	gcaggtggagcaggtggaCTCACCAACCTTGTTGCTATC	BACTH TXX-PilB$_{Vc}$ F
BBC2639	tgtaggctggagctgcttcTTAAAAGTAGAGCACACGCTG	BACTH TXX-PilB$_{Vc}$ R
BBC3182	gcaggtggagcaggtggaTCAGCATTTACAACACCACC	BACTH TXX-PilB$_{Ab}$ F
BBC3183	tgtaggctggagctgcttcTTATTCACTGGTTACACGATTAAT	BACTH TXX-PilB$_{Ab}$ R
JCP188	gcaggtggagcaggtggaACATATCATTTTGAAATAGATACACAATG	BACTH TXX-tfpB$_{Ab}$ F
BBC3328	tgtaggctggagctgcttcCTAAGTTAATGGGACCACCC	BACTH TXX-tfpB$_{Ab}$ R
BBC3890	gcaggtggagcaggtggaAGCGAAAACATTTCCCTTTCC	BACTH TXX-PilB$_{Ps}$ F

BBC3891	tgtaggctggagctgcttcTTAGTCCTTTGTCACCCGGTTG	BACTH TXX-PilB$_{Ps}$ R
BBC2434	gcaggtggagcaggtggaGAGTTGAATCAATATCTGGATGG	BACTH TXX-PilU$_{Vc}$ F
BBC1726	tgtaggctggagctgcttcCTACTCCATATCAATCTTCACATTC	BACTH TXX-PilU$_{Vc}$ R
BBC3197	gcaggtggagcaggtggaGACTTTAATGATTTACTCAATTTGATGG	BACTH TXX-PilU$_{Ab}$ F
BBC3198	tgtaggctggagctgcttcTTATTTGAGATTACGGCGTTGACC	BACTH TXX-PilU$_{Ab}$ R
BBC3981	gcaggtggagcaggtggaGAATTCGAGAAACTGTTGCG	BACTH TXX-PilU$_{Ps}$ F
JLC178	tgtaggctggagctgcttcGGCCTGTCAGCGGAAACTG	BACTH TXX-PilU$_{Ps}$ R
JCP318	GATTAACACACCCGCGaTtcGtAATTTGATTCGCGAAG	pilT$_{Ec}^{AVG\rightarrow AIR}$ F
JCP319	CTTCGCGAATCAAATTaCgaAtCGCGGGTGTGTTAATC	pilT$_{Ec}^{AVG\rightarrow AIR}$ R
JCP321	CTGGCGACGCCGGCAgTggGtAACTTGATCCGTGAAG	pilT$_{Vc}^{AIR\rightarrow AVG}$ F
JCP322	CTTCACGGATCAAGTTaCccAcTGCCGGCGTCGCCAG	pilT$_{Vc}^{AIR\rightarrow AVG}$ R
JCP264	tgtacttcttgcgcatcCAAACGCCCTTCCCCCAC	pilT$_{Ec}^{+15aa}$ R1
JCP259	ttggatttaaaagcatttTGAgaagcagctccagcctac	pilT$_{Ec}^{+15aa}$ F2
JCP262	gtaggctggagctgcttcCTACACCACACCACGCGCGATC	pilT$_{Vc}^{\Delta 15aa}$ R1
JCP263	gatcgcgcgtggtgtggtgTAGgaagcagctccagcctac	pilT$_{Vc}^{\Delta 15aa}$ F2