Retraction ATPase motors from three orthologous type IVa pilus systems support promiscuous retraction of the *Vibrio cholerae* competence pilus

Evan Couser[1], Jennifer L. Chlebek[1,*], and Ankur B. Dalia[1,*]

[1]Department of Biology, Indiana University, Bloomington, IN 47405.
*Correspondence to: ankdalia@indiana.edu or jennichlebek@gmail.com

Running title: T4aP retraction ATPase motors are promiscuous

ABSTRACT

Bacterial surface appendages called type IVa pili (T4aP) promote diverse activities including DNA uptake, twitching motility, and virulence. These activities rely on the ability of T4aP to dynamically extend and retract from the cell surface. Dynamic extension relies on a motor ATPase commonly called PilB. Most T4aP also rely on specific motor ATPases, commonly called PilT and PilU, to dynamically and forcefully retract. Here, we systematically assess whether motor ATPases from three orthologous T4aP can functionally complement *Vibrio cholerae* mutants that lack their endogenous motors. We found that the PilT and PilU retraction ATPases from the three T4aP systems tested are promiscuous and promote retraction of the *V. cholerae* competence T4aP despite a high degree of sequence divergence. In contrast, the orthologous extension ATPases from the same T4aP systems were not able to mediate extension of the *V. cholerae* competence T4aP despite exhibiting a similar degree of sequence divergence. Also, we show that one of the PilT orthologs characterized does not support PilU-dependent retraction and provide some data to indicate that the C-terminus of PilT is important for PilU-dependent retraction. Together, our data suggest that retraction ATPases may have maintained a high degree of promiscuity for promoting retraction of T4aP, while extension ATPases may have evolved to become specific for their cognate systems.

IMPORTANCE

One way that bacteria interact with their environments is via hair-like appendages called type IVa pili (T4aP). These appendages dynamically extend and retract from the cell surface via the action of distinct ATPase motors. T4aP are present in diverse bacterial species. Here, we demonstrate that retraction motors from three T4aP are promiscuous, and capable of promoting retraction of a heterologous T4aP system. By contrast, the extension ATPase motors from these same T4aP systems are specific and cannot promote extension of a heterologous T4aP. Thus, these results suggest that T4aP extension may be more tightly regulated compared to T4aP retraction.

INTRODUCTION

Type IVa pilus systems (T4aP) are encoded by diverse bacterial species. These systems promote the dynamic extension and retraction of filamentous appendages called pili from the bacterial surface. Through this dynamic activity, T4aP carry out a variety of functions including twitching motility, surface attachment, biofilm formation, virulence, and DNA uptake (1). Dynamic extension of all T4aP relies on the activity of a dedicated motor ATPase, which is

commonly called PilB (2). The gene that encodes this motor is generally found in close proximity
to other T4aP structural genes. Some systems also have an additional accessory extension
ATPase motor called TfpB (3). Dynamic retraction of T4aP canonically relies on a dedicated
retraction ATPase called PilT. Additionally, many T4aP systems contain an accessory retraction
ATPase called PilU. PilU is a PilT-dependent retraction ATPase that is required for forceful
retraction of the competence T4aP in *V. cholerae* (4) and is required for twitching motility in
other bacterial species (5). Unlike the extension motor, the genes encoding retraction motors
are generally not located near other T4aP structural genes. These hexameric extension and
retraction motors are thought to function via a mechanical interaction with the pilus platform,
which together take pilin subunits from the inner membrane and polymerize them into a filament
during pilus extension or depolymerize these subunits and return them to the inner membrane
during pilus retraction. Conformational changes in the ATPase motors due to ATP binding and
hydrolysis are thought to power these processes (6). A previous study demonstrated that the
PilT retraction motor from *Pseudomonas aeruginosa* was able to complement a *V. cholerae pilT*
mutant (7). However, a systematic analysis to assess motor promiscuity among T4aP is lacking.

Vibrio cholerae is a facultative pathogen that encodes a competence T4aP, which promotes
DNA uptake for natural transformation (8, 9). *Acinetobacter baylyi* and *Pseudomonas stutzeri*
encode functionally analogous competence T4aP that also promote horizontal gene transfer (3,
10-12). All of these T4aP systems encode canonical PilB, PilT, and PilU motor ATPases. For
this reason, *A. baylyi* and *P. stutzeri* were selected for this study. The T4aP of *A. baylyi* also
encodes an additional extension motor called TfpB that is required for initiating pilus extension
(3). Some strains of *Escherichia coli* also encode an ortholog of PilT (also referred to as *yggR*),
though these strains are not predicted to encode a PilU ortholog.

Here, we sought to formally test promiscuity among T4aP extension and retraction ATPases by
using the motors from the systems discussed above. We and others have demonstrated that the
competence pilus of *Vibrio cholerae* is a highly genetically tractable system for studying T4aP
dynamic activity (4, 7, 8, 13). Thus, we assessed promiscuity among T4aP motors by
determining if ectopic expression of heterologous motors could functionally complement *V.*
cholerae mutants that lack their endogenous motor ATPases. Our data suggests that while the
retraction ATPases PilT and PilU may be promiscuous in their ability to mediate retraction of
orthologous T4aP systems, extension ATPases display a high level of specificity for their
cognate T4aP. Furthermore, our data identify a PilT ortholog that supports ATPase-dependent
retraction, but not PilU-dependent retraction.

RESULTS

Retraction ATPase orthologs are promiscuous among T4aP systems, while extension ATPase
orthologs are not.
Because previous studies have shown that retraction ATPases from T4aP systems may be
promiscuous (7), we first wanted to determine if this characteristic is conserved among other
bacterial species. To test this, we ectopically expressed *pilT* from *A. baylyi* (*pilT$_{Ab}$*), *P. stutzeri*
(*pilT$_{Ps}$*), and *E. coli* (*pilT$_{Ec}$*) in *V. cholerae* strains where the native *pilT* or *pilTU* genes were
inactivated (i.e., Δ*pilT* and Δ*pilTU* backgrounds). As a control, we also ectopically expressed *V.*
cholerae pilT (*pilT$_{Vc}$*). Competence pili in *V. cholerae* promote DNA uptake during natural
transformation, a function that requires both pilus extension and retraction (8, 14). Thus, we
functionally assessed competence pilus dynamic activity by performing natural transformation
assays. Because PilU is an accessory retraction ATPase motor (4, 7), ectopic expression of
pilT$_{Vc}$ alone is able to rescue the transformation defect of the Δ*pilT* and Δ*pilTU* strains (**Fig. 1A**).
Furthermore, we observed that ectopic expression of *pilT$_{Ab}$*, *pilT$_{Ps}$*, and *pilT$_{Ec}$*, all fully

complemented natural transformation of Δ*pilT* and Δ*pilTU V. cholerae* (**Fig. 1A**). Western
blotting of functional N-terminally FLAG-tagged alleles of the PilT orthologs (4) confirmed that
their expression was equal to or greater than native PilT in *V. cholerae* (**Fig. S1**).These data
indicate that all of these retraction ATPases are capable of mediating retraction of the *V.*
cholerae competence T4aP, suggesting that the promiscuity of PilT retraction ATPases is
conserved across these species.

In addition to the dominant retraction ATPase PilT, many T4aP also encode an accessory
retraction ATPase called PilU that can promote pilus retraction in a PilT-dependent manner (4,
7). So next, we wanted to test whether PilU from *A. baylyi* and *P. stutzeri* could complement a
V. cholerae pilU mutant (the *E. coli* strain does not have a *pilU* ortholog). To test this, we
ectopically expressed *pilU*$_{Ab}$ and *pilU*$_{Ps}$ in a *V. cholerae* strain where native *pilT* contained a
Walker A mutation (*pilT*^{K136A} = *pilT*^{WA}) and the native *pilU* gene was inactivated (i.e., Δ*pilU*).
Mutation of the invariantly conserved lysine in the Walker A domain (K136A) inhibits the
ATPase activity of *pilT* (15), thus, efficient pilus retraction in this background (and
correspondingly a high frequency of natural transformation) is reliant on PilU ATPase activity (4,
7). As expected, ectopic expression of *pilU*$_{Vc}$ recovered the transformation defect of the *pilT*^{WA}
Δ*pilU* strain to parental levels (**Fig. 1B**). The expression of *pilU*$_{Ab}$ also showed almost full
recovery of transformation when compared to the parent strain (**Fig. 1B**). Interestingly, ectopic
expression of *pilU*$_{Ps}$ only partially complemented the transformation defect of this strain (**Fig.**
1B). Western blotting of functional N-terminally FLAG-tagged alleles of the PilU orthologs (4)
demonstrated that their expression was equal to or greater than native PilU in *V. cholerae* (**Fig.**
S1). This suggested that PilU$_{Ps}$ was either (1) generally a less functional PilU motor, or perhaps
(2) specifically less active at promoting pilus retraction in conjunction with PilT$_{Vc}$. To test this
further, we ectopically expressed *pilU*$_{Ps}$ in a background that also ectopically expressed a
Walker A mutant of its cognate PilT (*pilT*$_{Ps}$^{WA}). We found that in the presence of PilT$_{Ps}$^{WA}, PilU$_{Ps}$
was able to completely rescue natural transformation (**Fig. S2**). Together, this suggests that
while PilU is also promiscuous between these T4aP systems, it is perhaps less promiscuous
than PilT.

Because we found that retraction ATPases are, indeed, promiscuous between these T4aP, we
next wanted to test whether the extension ATPases from these same organisms were similarly
promiscuous. To test this, we ectopically expressed *pilB* orthologs from *A. baylyi* (*pilB*$_{Ab}$ and
tfpB$_{Ab}$), *P. stutzeri* (*pilB*$_{Ps}$), and *E. coli* (*pilB*$_{Ec}$; also referred to as *hofB*) in a *V. cholerae* Δ*pilB*
mutant and assessed natural transformation. While *pilB*$_{Vc}$ recovered natural transformation as
expected, ectopic expression of *pilB*$_{Ab}$, *tfpB*$_{Ab}$, *pilB*$_{Ps,}$ and *pilB*$_{Ec}$ failed to complement the Δ*pilB*
mutant (**Fig. 1C**). Importantly, Western blotting of N-terminally FLAG-tagged alleles of all PilB
orthologs (14) indicated that expression of each was equal to or greater than native PilB in *V.*
cholerae (**Fig. S1**). This suggests that extension ATPase orthologs from these species are not
able to mediate extension of the *V. cholerae* competence pilus. These data indicate that there
may be a high degree of specificity among extension ATPases, which is in stark contrast to the
observed promiscuity among the retraction ATPases studied here.

Extension and retraction ATPases likely interact with the pilus machinery platform protein PilC
to promote dynamic pilus activity (16). Thus, the observed promiscuity and/or specificity among
these motor ATPases may be attributed to their ability and/or inability to interact with PilC. To
test this, we assessed protein-protein interactions genetically using Bacterial Adenylate Cyclase
Two-Hybrid (BACTH) assays (17). First, looking at PilT, we found that each of the PilT orthologs
could self-interact, which is consistent with the hexameric conformation that likely all pilus
motors adopt (**Fig. 1D**). Also, consistent with the observation that all of the PilT orthologs tested
support natural transformation, we found that all PilT orthologs could interact with PilC$_{Vc}$ (**Fig.**
1D). Because PilU is a PilT-dependent retraction ATPase, it may need to interact with both PilT

and PilC to support pilus retraction. Indeed, all of the PilU orthologs could interact with PilT$_{Vc}$,
PilT$_{Vc}$^{WA}, and PilC$_{Vc}$, which is consistent with the ability of each PilU ortholog to support some
degree of natural transformation (**Fig. 1E**). For the PilB orthologs, we found that PilB$_{Vc}$ could
interact with PilC$_{Vc}$ as expected (**Fig. 1F**). Additionally, each of the PilB orthologs, apart from
PilB$_{Ec}$, could self-interact, which is consistent with the hexameric conformation that these
proteins likely adopt (**Fig. 1F**). Interestingly, PilB$_{Ab}$, TfpB$_{Ab}$, PilB$_{Ps}$, and PilB$_{Ec}$ varied in their
ability to interact with PilC$_{Vc}$. PilB$_{Ab}$, PilB$_{Ps}$, could not interact with PilC$_{Vc}$ and PilB$_{Ec}$ only
interacted with PilC$_{Vc}$ in one direction (**Fig. 1F**), which could suggest that these proteins fail to
support extension because they cannot interact properly with the platform. However, TfpB$_{Ab}$
could interact with PilC$_{Vc}$ (**Fig. 1F**), but still failed to support natural transformation (**Fig. 1C**).
These data indicate that while interactions between PilB and PilC may be required for extension
of the competence pilus, these interactions are not sufficient to promote extension.

The PilT ortholog from E. coli cannot support PilU-mediated retraction.
Above, our results indicate that PilT orthologs from distinct T4aP can promote retraction of the
V. cholerae competence pilus in the absence of native PilT$_{Vc}$ and PilU$_{Vc}$. PilT, however, can also
support PilU-dependent retraction, in a manner that is not dependent on PilT ATPase activity
(4). So next, we wanted to test if the PilT orthologs among the studied T4aP could support PilU-
dependent retraction in conjunction with PilU$_{Vc}$. To test this, Walker A mutants of each *pilT* gene
(*pilT$_{Vc}$^{WA}*, *pilT$_{Ab}$^{WA}*, *pilT$_{Ps}$^{WA}*, *pilT$_{Ec}$^{WA}*) were ectopically expressed in a Δ*pilT* background (i.e.,
where *pilU$_{Vc}$* is intact) and a Δ*pilTU* background. As expected, the ectopic expression of *pilT$_{Vc}$^{WA}*
fully restored the transformation efficiency of a Δ*pilT* background (since *pilU$_{Vc}$* is intact) (**Fig.
2A**). Importantly, transformation was not restored by ectopic expression of any Walker A *pilT*
alleles in a Δ*pilTU* background (**Fig. 2A**), which indicates that these ATPase defective alleles of
PilT only support PilU-dependent retraction and cannot promote retraction on their own. When
pilT$_{Ab}$^{WA} and *pilT$_{Ps}$^{WA}* were ectopically expressed in the Δ*pilT* background, transformation was
rescued to parental levels (**Fig. 2A**). These data suggest that PilU$_{Vc}$ is able to mediate retraction
of the *V. cholerae* competence pilus in a PilT$_{Ab}$- and PilT$_{Ps}$-dependent manner, which further
demonstrates the promiscuity among these retraction ATPases. Ectopic expression of the
Walker A mutant allele of *pilT$_{Ec}$*, however, did not restore natural transformation in the Δ*pilT*
background (**Fig. 2A**). This suggests that PilT$_{Ec}$^{WA} is incapable of promoting PilU-dependent
retraction in the presence of PilU$_{Vc}$.

The ability of PilU to interact with PilT is likely necessary for PilU-dependent retraction of the
competence pilus. As expected, PilT$_{Vc}$, PilT$_{Ab}$, PilT$_{Ps}$, and PilT$_{Ec}$ could self-interact (**Fig. 2B**) as
seen previously (**Fig. 1D**), and each allele could also interact with PilU$_{Vc}$ (**Fig. 2B**). Because
PilU$_{Vc}$-mediated retraction was possible when *pilT$_{Vc}$^{WA}*, *pilT$_{Ab}$^{WA}*, and *pilT$_{Ps}$^{WA}* were ectopically
expressed, we hypothesized that each of these PilT alleles would be able to self-interact and
also interact with PilU. PilT$_{Vc}$^{WA} and PilT$_{Ab}$^{WA} were able to self-interact (**Fig. 2B**), which suggests
that the Walker A mutation does not inhibit the ability of these proteins to self-assemble into
hexamers. This is consistent with our previous results which demonstrate that purified PilT$_{Vc}$^{WA}
can still form hexameric rings (4). Unexpectedly, PilT$_{Ps}$^{WA} and PilT$_{Ec}$^{WA} did not exhibit self-
interactions in these BACTH assays (**Fig. 2B**). We also observed that PilU$_{Vc}$ could interact with
PilT$_{Vc}$^{WA} and PilT$_{Ab}$^{WA}, however, it did not interact with PilT$_{Ec}$^{WA} or PilT$_{Ps}$^{WA} (**Fig. 2B**). The
absence of interactions between PilU$_{Vc}$ and PilT$_{Ec}$^{WA} (**Fig. 2B**) is consistent with the inability of
PilT$_{Ec}$^{WA} to promote PilU-dependent retraction (**Fig. 2A**). However, the absence of interactions
between PilU$_{Vc}$ and PilT$_{Ps}$^{WA} was particularly surprising since this PilT allele can support PilU-
dependent retraction when expressed in *V. cholerae* (**Fig. 2A**). Because PilT$_{Ps}$^{WA} supports PilU-
dependent retraction in *V. cholerae*, but does not exhibit self-interactions in BACTH assays, this
may suggest that self-assembly or hexamerization of PilT is not required for PilU-dependent
retraction. However, the inability of PilT$_{Ps}$^{WA} to interact with PilU$_{Vc}$ in these same BACTH assays

(**Fig. 2B**), despite the ability of this allele to support PilU-dependent retraction (**Fig. 2A**) makes
this difficult to conclude. The observed inability of $PilT_{Ec}^{WA}$ and $PilT_{Ps}^{WA}$ to self-interact and/or
interact with $PilU_{Vc}$ in BACTH assays could either be due to an absence of interactions between
these proteins, or could be due to a poor affinity interaction between these proteins that is below
the detection limit for the assay.

The molecular mechanism underlying PilU-mediated retraction in a PilT-dependent manner
remains unclear. The ability of PilT to self-assemble and form a hexamer may be a prerequisite
for PilU-mediated retraction. However, other characteristics of PilT may also be responsible for
supporting PilU-mediated retraction. Because $pilT_{Ec}^{WA}$ lacked self-interactions, we constructed
an alternative mutant that also disrupted the ATPase activity of $PilT_{Ec}$ by mutating an invariantly
conserved glutamate in the Walker B domain of $pilT_{Ec}$ ($pilT_{Ec}^{E201A} = pilT_{Ec}^{WB}$). Importantly,
$PilT_{Ec}^{WB}$ exhibited self-interactions in BACTH assays (**Fig. 2B**), suggesting that the Walker B
mutation allows for self-assembly and/or a higher affinity between subunits compared to the
Walker A mutant. When we ectopically expressed the $pilT_{Ec}^{WB}$ in a $\Delta pilT$ background (where
$pilU$ is intact), however, we observed that this mutant did not restore transformation (**Fig. 2A**),
which is similar to the $pilT_{Ec}^{WA}$ allele. This indicates that self-assembly of an ATPase defective
$PilT_{Ec}$ is not sufficient to promote PilU-dependent retraction. Together, these data suggest that
$PilT_{Ec}$ may have unique features that prevents it from promoting PilU-dependent retraction. We
hypothesized that analyzing the differences between $PilT_{Ec}$ and the other PilT alleles ($PilT_{Vc}$,
$PilT_{Ab}$, and $PilT_{Ps}$) may help define the features of PilT that allow for PilU-dependent retraction.

The C-terminus of $PilT_{Vc}$ contributes to PilU-mediated retraction.
In order to narrow down the region of $PilT_{Vc}$ that is necessary and/or sufficient to support PilU-
dependent retraction, we constructed defined chimeras of $pilT_{Vc}$ and $pilT_{Ec}$. $PilT_{Vc}$ and $PilT_{Ec}$ are
345 residues and 326 residues long, respectively. The chimeras were constructed in intervals of
50 amino acids where the residues of one protein were replaced with the corresponding amino
acids of the other. These chimeras were first ectopically expressed in a $\Delta pilTU$ background to
test their ability to mediate retraction of the competence pilus in a manner that is dependent on
their ATPase activity – a property that both $PilT_{Vc}$ and $PilT_{Ec}$ shared (**Fig. 1A**). Surprisingly, only
three of the chimeras were able to restore parental levels of transformation in the $\Delta pilTU$
background (**Fig. 3A**). Two of the functional chimeras consisted of portions of the N-terminus of
$PilT_{Vc}$ (100 or 200 amino acids) fused to the corresponding C-terminus of $PilT_{Ec}$. The remaining
functional chimera contained the first 300 amino acids of $PilT_{Ec}$ fused to the C-terminal 42 amino
acids of $PilT_{Vc}$. One potential explanation for this general lack of functionality is that the locations
of the junctions between the two proteins were not conducive to generating a functional ATPase
motor. PilT is composed of two domains, an N-terminal PAS-like domain and a C-terminal
RecA-like ATPase domain which are connected via a short linker (95-102 in $PilT_{Ec}$ and 98-105
in $PilT_{Vc}$). So, the chimeras at 100 residues should be within the linker that connects the two
domains of both proteins, and it was particularly surprising that the $PilT_{Ec}$(100) fusion was not
functional when the reciprocal $PilT_{Vc}$(100) fusion was functional. The three functional chimeras
of PilT obtained are the only alleles that we can be reasonably confident could potentially
support PilU-dependent retraction, thus, we only proceeded with characterizing these three
chimeras of PilT moving forward. To determine if these three chimeras could support PilU-
mediated retraction, we next introduced a Walker A mutation into each allele. None of the $pilT_{Vc}$
/ $pilT_{Ec}$ chimeras with the Walker A mutation were able to rescue PilU-dependent transformation
efficiency in a $\Delta pilT$ background (**Fig. 3A**). Separately, a Walker B mutation was introduced into
these chimeras. Again, none of the 3 chimeras were able to restore PilU-dependent
transformation in a $\Delta pilT$ background (**Fig. 3A**). We reasoned that since $PilT_{Vc}$ is capable of
PilU-mediated retraction, that the chimera that is predominantly composed of $PilT_{Vc}$ might help
define a portion of the protein that plays an important role in PilU-mediated retraction (i.e. that

the part of PilT$_{Vc}$ missing in this chimera plays an important role). PilT$_{Vc}$(200) had the largest
portion of PilT$_{Vc}$ and still made a functional retraction motor when ATPase activity was intact
(**Fig. 3A**). This chimera had 200 residues of the PilT$_{Vc}$ N-terminus fused to the C-terminal 129
residues of PilT$_{Ec}$. Walker A and Walker B mutants of PilT$_{Vc}$(200) did not support PilU-
dependent retraction (**Fig. 3A**). So, we reasoned that the C-terminal 145 residues of PilT$_{Vc}$
(which were replaced with PilT$_{Ec}$ in these chimeras) must be playing an important role in PilU-
dependent retraction. However, this does not exclude the possibility that the N-terminus of PilT$_{Vc}$
also plays an important role in promoting PilU-dependent retraction.

In order to further investigate potential components of PilT that are required for supporting PilU-
mediated retraction, we constructed an alignment of the 4 PilT orthologs that were studied here.
The alignment revealed that the middle section of PilT$_{Ec}$ was conserved relative to all of the
other PilT orthologs (**Fig. S3**). However, the N-terminus and the C-terminus of PilT$_{Ec}$ were
poorly conserved relative to the other PilT orthologs. Because our data suggests that the C-
terminus of PilT$_{Vc}$ may be important for supporting PilU-mediated retraction, that is where we
focused our investigation. In particular, we observed that *pilT$_{Ec}$* contains two substitutions in the
highly conserved AIRNLIRE domain of PilT (18). To determine if these substitutions were
responsible for the inability of PilT$_{Ec}$ to support PilU-mediated retraction (7), we mutated the two
residues in *pilT$_{Ec}$* to the residues present in the other PilT orthologs (PilT$_{Ec}$$^{AVG \rightarrow AIR}$). Conversely,
to determine if these changes in PilT$_{Ec}$ contributed to its inability to support PilU-mediated
retraction, the residues in *pilT$_{Vc}$* were mutated to the residues present in *pilT$_{Ec}$* (PilT$_{Vc}$$^{AIR \rightarrow AVG}$).
These mutations were introduced into Walker A (WA) or Walker B (WB) mutant alleles of PilT so
that we could assess PilU-mediated retraction. The introduction of these mutations into *pilT$_{Vc}$*
did not result in the inability of PilT$_{Vc}$ to facilitate PilU-mediated retraction (**Fig. 3B**). Moreover,
the restoration of the two AIRNLIRE substitutions in *pilT$_{Ec}$* did not rescue PilU-mediated
retraction (**Fig. 3B**). These data indicate that the substitutions in the AIRNLIRE domain of *pilT$_{Ec}$*
are not responsible for the inability of PilT$_{Ec}$ to support PilU-mediated retraction.

The alignment of the 4 PilT orthologs also revealed that 15 amino acids in the C-terminal
domain of PilT$_{Ec}$ were missing when compared to the other PilT orthologs that did support PilU-
mediated retraction (PilT$_{Vc}$, PilT$_{Ab}$, and PilT$_{Ps}$). We hypothesized that the deletion of the last 15
amino acids in PilT$_{Ec}$ could be responsible for the inability of this protein to support PilU-
mediated retraction. To test this, we deleted the last 15 amino acids from PilT$_{Vc}$ (*pilT$_{Vc}$$^{\Delta 15aa}$*)·
Additionally, we fused the last 15 amino acids present at the C-terminus of *pilT$_{Vc}$* to the end of
PilT$_{Ec}$ (*pilT$_{Ec}$$^{+15aa}$*). Again, these mutations were generated in Walker A and Walker B mutant
alleles of PilT so that we could assess PilU-mediated retraction. The *pilT$_{Vc}$$^{\Delta 15aa,\ WA}$* mutant
displayed no reduction in transformation (**Fig. 3B**), suggesting that PilT$_{Vc}$$^{\Delta 15aa,\ WA}$ is still able to
facilitate PilU-mediated retraction. Conversely, the *pilT$_{Ec}$$^{+15aa,\ WA}$* and *pilT$_{Ec}$$^{+15aa,\ WB}$* mutations did
not restore transformation to parental levels (**Fig. 3B**), which suggests that PilU$_{Vc}$ cannot
mediate retraction in the presence of these PilT$_{Ec}$ alleles. Together, these data suggest that the
last 15 amino acids present at the C-terminus of PilT$_{Vc}$ is not necessary or sufficient to promote
PilU-mediated retraction.

Our results have suggested that, individually, the substitutions present in the conserved
AIRNLIRE domain of PilT$_{Ec}$ and the deletion of the last 15 amino acids in the C-terminus are not
responsible for the inability of this PilT allele to support PilU-mediated retraction. However, it is
possible that the combination of these two factors diminishes PilU-mediated retraction. To test
this, we constructed *pilT$_{Vc}$ alleles* that contained both the AIRNLIRE domain substitutions and
the C-terminal deletion of 15 amino acids (*pilT$_{Vc}$$^{AIR \rightarrow AVG,\ \Delta 15aa}$*). Also, we generated a *pilT$_{Ec}$* allele
where we restored the AIRNLIRE domain and inserted the terminal 15 amino acids from *pilT$_{Vc}$*
onto the C-terminus (*pilT$_{Ec}$$^{AVG \rightarrow AIR,\ +15aa}$*). The combination of mutations in *pilT$_{Vc}$* resulted in a loss
of function, even in a strain where the ATPase activity of the allele was intact (**Fig. 3B**).

Additionally, the $pilT_{Ec}^{AVG \to AIR, +15aa, WA}$ and $pilT_{Ec}^{AVG \to AIR, +15aa, WB}$ alleles did not restore
transformation to parental levels in the $\Delta pilT$ background, suggesting that PilU was unable to
mediate retraction in these backgrounds. This suggests that the combination of the 15 amino
acid deletion and the substitutions in the AIRNLIRE domain are not solely responsible for the
inability of $PilT_{Ec}$ to support PilU-mediated retraction.

DISCUSSION

This work demonstrates that the retraction ATPases PilT and PilU are promiscuous among the
subset of T4aP systems studied here, and that their functionality extends beyond species
boundaries. This contrasts with the fidelity displayed by the extension ATPases from the same
T4aP systems. Additionally, our data suggests that the C-terminus of $PilT_{Vc}$ plays a role in
supporting PilU-mediated retraction. However, further analysis determined that the genetic
differences between $PilT_{Vc}$ and $PilT_{Ec}$ in the AIRNLIRE domain and the extended C-terminus are
neither necessary nor sufficient to explain this discrepancy in activity. Because of this, the exact
residues in the PilT C-terminus that contribute to PilU-mediated retraction remain elusive. Also,
it is likely that the N-terminus of $PilT_{Vc}$ also contributes to PilU-dependent retraction, however,
the paucity of functional $PilT_{Vc}$ / $PilT_{Ec}$ chimeras made this difficult to narrow down.

Because the orthologous retraction motor ATPases were overexpressed in this study, we
cannot comment on the relative efficiency of the ectopically expressed retraction motors
compared to native $PilT_{Vc}$ and $PilU_{Vc}$. But it is important to note that similar overexpression of
the extension motor orthologs from the same T4aP did not promote dynamic competence pilus
activity, which indicates that the overabundance of a motor ATPase is not necessarily sufficient
for motor functionality. One possible explanation for the discrepancy in the promiscuity /
specificity among motor ATPases could be due to differences in sequence divergence; where
retraction motor orthologs are more similar to one another compared to the extension motor
orthologs. When we performed this analysis, however, we found that the PilT orthologs ranged
from ~62-80% similar (with $PilT_{Ec}$ being the most divergent), the PilU orthologs ranged from
~70-74% similar, and the extension motor orthologs ranged from ~50-66% similar (**Fig. S4**).
These data indicate that overall sequence divergence among PilT and PilB orthologs is fairly
similar. Thus, suggesting that sequence divergence alone cannot account for the promiscuity /
specificity among these motor ATPases.

Why are retraction ATPases promiscuous and extension ATPases specific? One possibility is
that it is advantageous for T4aP systems to maintain promiscuous retraction ATPases. T4aP
are diverse and retraction ATPases are generally expressed at distinct loci from the other T4aP
components. Some T4aP systems are likely transferred between organisms through horizontal
gene transfer (19). The inability to retract T4aP is likely disadvantageous in many circumstances
since these surface appendages are highly immunogenic and can also be receptors for diverse
phages (20). Thus, it is tempting to speculate that one potential advantage to retraction ATPase
promiscuity is that this would allow cells that acquire a new T4aP the ability to retract pili from
the surface.

Some organisms, like *V. cholerae*, encode multiple T4P - competence pili, mannose sensitive
hemagglutinin (MSHA) pili, and toxin co-regulated pili (TCP) - and there are distinct
environmental conditions under which each pilus system is likely needed. Competence pili and
MSHA pili are likely critical for the survival of *V. cholerae* in its environmental reservoir, while
TCP are required for infection of its human host. In fact, it has previously been shown that the
inappropriate expression of MSHA pili during the context of an infection attenuates *V. cholerae*
(21). A high degree of specificity among the extension ATPase motors for these distinct T4P

may represent one mechanism by which cells can exhibit tight control over when these distinct
pili are made. In contrast, the retraction of these pili may not require the same level of regulation
because pilus extension is so tightly regulated. Indeed, in *V. cholerae,* PilT and PilU promote
retraction of both competence pili and MSHA pili (8, 22).

The molecular mechanism underlying the specificity of extension ATPase motors remains
unclear. One possibility is that the N-terminal domain of PilB provides specificity. It is thought
that the most N-terminal domain of PilB is responsible for interactions with PilM, which is likely
required for proper pilus extension (6). An alignment of the extension and retraction motor
orthologs showed that the extension motors contain an extended N-terminal domain (~184
residues), called N1D (6) that distinguishes them from the retraction ATPase motors (**Fig. S5**),
and we found that much of the sequence divergence among extension ATPases was attributed
to the N1D (**Fig. S4**). Thus, it is tempting to speculate that the N1D of extension ATPases plays
a critical role in defining the specificity of these motors for their cognate T4aP system.

While our results indicate that retraction motors are promiscuous and extension motors are
specific among the T4aP studied here, it remains unclear which of these phenotypes is the
exception vs the rule. To distinguish between these possibilities, it would be interesting to test
the degree of specificity / promiscuity among other T4aP components. Addressing this question
may continue to shed light on the evolutionary pressures that have helped shaped these nearly
ubiquitous molecular machines, and will be the focus of future research.

METHODS

Bacterial Strains and Growth Conditions
Vibrio cholerae strains were grown in LB Miller broth and LB Miller agar plates that contained
erythromycin (10 µg/mL), carbenicillin (20 µg/mL), and/or kanamycin (50 µg/mL) when
necessary. A list of all strains used in this study can be found in **Table S1**.

Construction of Mutant Strains and Constructs
All of the *V. cholerae* strains used in this study are derivatives of the El tor isolate E7946. The
motor ATPase orthologs from other species were amplified from genomic DNA of the following
strains: *Acinetobacter baylyi* ADP1 (23), *Pseudomonas stutzeri* JM300 (24), and *Escherichia*
coli MG1655 (25). All mutant constructs were made via splicing-by-overlap extension (SOE)
PCR, exactly as previously described (13, 26). A detailed list of all primers used to generate
mutant constructs can be found in **Table S2**. Mutant constructs were introduced onto the *V.*
cholerae genome by natural transformation and natural cotransformation exactly as previously
described (26). Strains were constructed by first inserting the ectopic expression construct of
the pilus component into the genome. This was followed by the deletion of the native pilus
component. All mutations and ectopic expression constructs were confirmed by PCR and/or
sequencing.

Natural Transformation Assays
In this study, we aimed to address the role of distinct T4aP components on competence pilus
function. Thus, we opted to bypass the natural mechanism of competence induction in *V.*
cholerae by ectopically expressing the master competence regulator TfoX (P_{tac}-*tfoX*) and
inactivating the quorum sensing negative regulator LuxO ($\Delta luxO$) as previously described (8).
Chitin-independent natural transformation assays were performed exactly as previously
described (8, 14). Briefly, strains were grown overnight in 3 mL of LB broth containing 100 µM
IPTG at 30 °C rolling. Next, cells were subcultured into 3 mL of LB supplemented with 20 mM
MgCl$_2$, 10 mM CaCl$_2$, 100 µM IPTG, and arabinose (when appropriate; 0.05-0.2%) and grown to

late-log phase at 30 $^{\circ}$C rolling. Next, ~10^8 cells were transferred to Instant Ocean medium (7
g/L; Aquarium Systems) containing 100 µM IPTG, arabinose (when appropriate; 0.05-0.2%),
and 200 ng of a transforming DNA (tDNA) PCR product. Reactions were then incubated
statically overnight at 30 $^{\circ}$C. The tDNA product used throughout this study targeted a "neutral
gene" (the frame-shifted transposase, VC1807) for inactivation (ΔVC1807::Erm^R) as previously
described (27). A negative control, containing no tDNA, was included for each strain in all
transformation assays. After the overnight incubation with tDNA, reactions were outgrown by
adding 1 mL of LB to each reaction and incubating at 37 $^{\circ}$C shaking for 2 hours. After the
outgrowth, reactions were plated for quantitative culture to distinguish between transformants
(plated on LB + erythromycin) and total viable counts (plated on plain LB). The transformation
frequency for each reaction was calculated by dividing the number of transformants by the total
viable counts.

Bacterial Adenylate Cyclase Two-Hybrid Assay
Using PCR, the desired motor ATPase genes were amplified and cloned into the BACTH
vectors pKT25 and pUT18C in order to produce N-terminal fusions between the motor ATPase
and the T25 and T18 fragments of adenylate cyclase, respectively. Miniprepped plasmids
(Qiagen) were then co-transformed into *E. coli* BTH101 cells. As a positive control, T25-zip and
T18-zip vectors were cotransformed into BTH101. And as a negative control, the pKT25 and
pUT18C empty vectors were cotransformed into BTH101. Transformations were plated onto LB
+ kanamycin (50 µg/mL) + carbenicillin (100 µg/mL) plates in order to select for transformants
that received both plasmids. These transformants were then selected and grown overnight at 30
$^{\circ}$C static in LB + kanamycin (50 µg/mL) + carbenicillin (100 µg/mL). Finally, 3 µL of the overnight
cultures were spotted onto LB agar plates supplemented with 500 µM IPTG, kanamycin (50
µg/mL), carbenicillin (100 µg/mL), and X-gal (40 µg/mL). These plates were allowed to incubate
statically at 30 $^{\circ}$C for ~48 hours prior to imaging on a HP Scanjet G4010 flatbed scanner.

Alignments
The sequences of the genes that encode for the desired proteins were downloaded from
GenBank. Sequences were aligned using T-Coffee (28) and the resulting alignments were run
through the Sequence Manipulation Suite: Color Align Conservation tool (29) to generate the
final figures.

Pairwise comparisons to assess similarity among ATPase motor orthologs
Pairwise comparisons between ATPase motor orthologs were performed via global alignment
using the Needleman-Wunsch algorithm (30). The percent similarities generated by these
pairwise comparisons are reported in **Fig. S4**.

Western Blotting

Strains were grown to late-log phase in LB supplemented with 20 mM MgCl$_2$, 10 mM CaCl$_2$, 100
µM IPTG, and arabinose (when appropriate) exactly as described above for transformation
assays. Cells were washed and concentrated to an OD$_{600}$ = 100 in 0.5X Instant Ocean medium,
and then lysed by adding Fastbreak lysis buffer (to 1X), lysozyme (80 µg/mL), and benzonase
(25U). Lysed samples were mixed 1:1 with 2X SDS sample buffer [220 mM Tris pH 6.8, 25%
glycerol, 1.8% SDS, 0.02% Bromophenol Blue, 5% β-mercaptoethanol] and 3 µL of each
sample was then separated on a 15% SDS PAGE gel. The proteins were transferred to a PVDF
membrane via electrophoresis, and the membrane was then incubated with the appropriate
primary antibody (α-FLAG polyclonal rabbit – for PilT and PilU; or α-FLAG M2 mouse
monoclonal – for PilB). A primary incubation with α-RpoA monoclonal mouse (BioLegend) was
also completed as a loading control. The blots were then washed and incubated with an
appropriate horseradish peroxidase (HRP) conjugated secondary antibody. And blots were

developed with Pierce™ ECL Western Blotting Substrate and imaged using a ProteinSimple
FluorChem R system.

ACKNOWLEDGEMENTS

We would like to thank David Kysela for helpful discussions. This work was supported by grant
R35GM128674 from the National Institutes of Health (to ABD).

References

437   1.   Craig L, Forest KT, Maier B. 2019. Type IV pili: dynamics, biophysics and functional
438   consequences. Nat Rev Microbiol 17:429-440.
439   2.   Jakovljevic V, Leonardy S, Hoppert M, Sogaard-Andersen L. 2008. PilB and PilT are
440   ATPases acting antagonistically in type IV pilus function in Myxococcus xanthus. J
441   Bacteriol 190:2411-21.
442   3.   Ellison CK, Dalia TN, Klancher CA, Shaevitz JW, Gitai Z, Dalia AB. 2021. Acinetobacter
443   baylyi regulates type IV pilus synthesis by employing two extension motors and a motor
444   protein inhibitor. Nat Commun 12:3744.
445   4.   Chlebek JL, Hughes HQ, Ratkiewicz AS, Rayyan R, Wang JC, Herrin BE, Dalia TN,
446   Biais N, Dalia AB. 2019. PilT and PilU are homohexameric ATPases that coordinate to
447   retract type IVa pili. PLoS Genet 15:e1008448.
448   5.   Whitchurch CB, Mattick JS. 1994. Characterization of a gene, pilU, required for twitching
449   motility but not phage sensitivity in Pseudomonas aeruginosa. Mol Microbiol 13:1079-91.
450   6.   McCallum M, Tammam S, Khan A, Burrows LL, Howell PL. 2017. The molecular
451   mechanism of the type IVa pilus motors. Nat Commun 8:15091.
452   7.   Adams DW, Pereira JM, Stoudmann C, Stutzmann S, Blokesch M. 2019. The type IV
453   pilus protein PilU functions as a PilT-dependent retraction ATPase. PLoS Genet
454   15:e1008393.
455   8.   Ellison CK, Dalia TN, Vidal Ceballos A, Wang JC, Biais N, Brun YV, Dalia AB. 2018.
456   Retraction of DNA-bound type IV competence pili initiates DNA uptake during natural
457   transformation in Vibrio cholerae. Nat Microbiol 3:773-780.
458   9.   Seitz P, Blokesch M. 2013. DNA-uptake machinery of naturally competent Vibrio
459   cholerae. Proc Natl Acad Sci U S A 110:17987-92.
460   10.   Leong CG, Bloomfield RA, Boyd CA, Dornbusch AJ, Lieber L, Liu F, Owen A, Slay E,
461   Lang KM, Lostroh CP. 2017. The role of core and accessory type IV pilus genes in
462   natural transformation and twitching motility in the bacterium Acinetobacter baylyi. PLoS
463   One 12:e0182139.
464   11.   Graupner S, Weger N, Sohni M, Wackernagel W. 2001. Requirement of novel
465   competence genes pilT and pilU of Pseudomonas stutzeri for natural transformation and
466   suppression of pilT deficiency by a hexahistidine tag on the type IV pilus protein PilAI. J
467   Bacteriol 183:4694-701.
468   12.   Graupner S, Frey V, Hashemi R, Lorenz MG, Brandes G, Wackernagel W. 2000. Type
469   IV pilus genes pilA and pilC of Pseudomonas stutzeri are required for natural genetic
470   transformation, and pilA can be replaced by corresponding genes from nontransformable
471   species. J Bacteriol 182:2184-90.
472   13.   Dalia TN, Chlebek JL, Dalia AB. 2020. A modular chromosomally integrated toolkit for
473   ectopic gene expression in Vibrio cholerae. Sci Rep 10:15398.
474   14.   Chlebek JL, Dalia TN, Biais N, Dalia AB. 2021. Fresh Extension of Vibrio cholerae
475   Competence Type IV Pili Predisposes Them for Motor-Independent Retraction. Appl
476   Environ Microbiol 87:e0047821.
477   15.   Chiang P, Sampaleanu LM, Ayers M, Pahuta M, Howell PL, Burrows LL. 2008.
478   Functional role of conserved residues in the characteristic secretion NTPase motifs of
479   the Pseudomonas aeruginosa type IV pilus motor proteins PilB, PilT and PilU.
480   Microbiology (Reading) 154:114-126.
481   16.   Bischof LF, Friedrich C, Harms A, Sogaard-Andersen L, van der Does C. 2016. The
482   Type IV Pilus Assembly ATPase PilB of Myxococcus xanthus Interacts with the Inner
483   Membrane Platform Protein PilC and the Nucleotide-binding Protein PilM. J Biol Chem
484   291:6946-57.

485   17.   Karimova G, Pidoux J, Ullmann A, Ladant D. 1998. A bacterial two-hybrid system based
486   on a reconstituted signal transduction pathway. Proc Natl Acad Sci U S A 95:5752-6.
487   18.   Aukema KG, Kron EM, Herdendorf TJ, Forest KT. 2005. Functional dissection of a
488   conserved motif within the pilus retraction protein PilT. J Bacteriol 187:611-8.
489   19.   Denise R, Abby SS, Rocha EPC. 2019. Diversification of the type IV filament superfamily
490   into machines for adhesion, protein secretion, DNA uptake, and motility. PLoS Biol
491   17:e3000390.
492   20.   Giltner CL, Nguyen Y, Burrows LL. 2012. Type IV pilin proteins: versatile molecular
493   modules. Microbiol Mol Biol Rev 76:740-72.
494   21.   Hsiao A, Liu Z, Joelsson A, Zhu J. 2006. Vibrio cholerae virulence regulator-coordinated
495   evasion of host immunity. Proc Natl Acad Sci U S A 103:14542-7.
496   22.   Floyd KA, Lee CK, Xian W, Nametalla M, Valentine A, Crair B, Zhu S, Hughes HQ,
497   Chlebek JL, Wu DC, Hwan Park J, Farhat AM, Lomba CJ, Ellison CK, Brun YV,
498   Campos-Gomez J, Dalia AB, Liu J, Biais N, Wong GCL, Yildiz FH. 2020. c-di-GMP
499   modulates type IV MSHA pilus retraction and surface attachment in Vibrio cholerae. Nat
500   Commun 11:1549.
501   23.   Barbe V, Vallenet D, Fonknechten N, Kreimeyer A, Oztas S, Labarre L, Cruveiller S,
502   Robert C, Duprat S, Wincker P, Ornston LN, Weissenbach J, Marliere P, Cohen GN,
503   Medigue C. 2004. Unique features revealed by the genome sequence of Acinetobacter
504   sp. ADP1, a versatile and naturally transformation competent bacterium. Nucleic Acids
505   Res 32:5766-79.
506   24.   Busquets A, Pena A, Gomila M, Bosch R, Nogales B, Garcia-Valdes E, Lalucat J,
507   Bennasar A. 2012. Genome sequence of Pseudomonas stutzeri strain JM300 (DSM
508   10701), a soil isolate and model organism for natural transformation. J Bacteriol
509   194:5477-8.
510   25.   Blattner FR, Plunkett G, 3rd, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J,
511   Glasner JD, Rode CK, Mayhew GF, Gregor J, Davis NW, Kirkpatrick HA, Goeden MA,
512   Rose DJ, Mau B, Shao Y. 1997. The complete genome sequence of Escherichia coli K-
513   12. Science 277:1453-62.
514   26.   Dalia AB. 2018. Natural Cotransformation and Multiplex Genome Editing by Natural
515   Transformation (MuGENT) of Vibrio cholerae. Methods Mol Biol 1839:53-64.
516   27.   Dalia AB, McDonough E, Camilli A. 2014. Multiplex genome editing by natural
517   transformation. Proc Natl Acad Sci U S A 111:8937-42.
518   28.   Notredame C, Higgins DG, Heringa J. 2000. T-Coffee: A novel method for fast and
519   accurate multiple sequence alignment. J Mol Biol 302:205-17.
520   29.   Stothard P. 2000. The Sequence Manipulation Suite: JavaScript Programs for Analyzing
521   and Formatting Protein and DNA Sequences. BioTechniques 28:1102-1104.
522   30.   Needleman SB, Wunsch CD. 1970. A general method applicable to the search for
523   similarities in the amino acid sequence of two proteins. J Mol Biol 48:443-53.

Fig. 1. *Retraction ATPase orthologs are promiscuous among T4aP systems, while extension ATPase orthologs are not.* Natural transformation assays were performed to test whether (**A**) PilT, (**B**) PilU, or (**C**) PilB/TfpB orthologs from 3 heterologous T4aP systems could complement *V. cholerae* mutants lacking their endogenous motor ATPases. The relevant genotypes of strains are indicated below each bar, which includes the status of the native *V. cholerae pilT/pilU/pilB* genes (top) as well as the ectopically expressed motor ATPase allele (induced via P_{BAD}) (bottom). The color of the bar corresponds to the species from which the ectopically expressed *motor ATPase* was derived (green bars = *Vc motors*, yellow bars = *Ab* motors, orange bars = *Ps* motors, and purple bars = *Ec* motors). Data are from at least 3 independent biological replicates and shown as the mean ± SD. The limit of detection for transformation frequency assays is ~10^{-7}. In **C**, Ab_1 denotes *pilB*$_{Ab}$ and Ab_2 denotes *tfpB*$_{Ab}$. In **C**, for all samples where the transformation frequency is ~10^{-7}, at least 1 replicate contained zero transformants and the limit of detection for that sample is displayed. BACTH assays to test whether the indicated (**D**) PilT, (**E**) PilU, or (**F**) PilB/TfpB alleles can interact with themselves ("self") or the indicated T4P components (PilC or PilT). For BACTH assays pilus components had an N-terminal fusion to the T25 or T18 fragment of adenylate cyclase as indicated. "WA" denotes a Walker A mutation in the tested allele. "+" and "−" denote positive and negative controls for the BACTH assay. "E.V." denotes an empty vector. Data are representative of at least 2 independent experiments.

pilT	+	Δ	Δ	WA	WA	Δ	Δ	Δ	Δ	Δ	Δ	Δ	Δ	Δ
pilU	+	+	Δ	+	Δ	+	+	Δ	+	Δ	+	Δ	+	Δ
P$_{BAD}$-pilT$_{XX}$^{WA}	-	-	-	-	-	-	Vc	-	Vc	Ab	Ab	Ps	Ps	Ec	-	Ec
P$_{BAD}$-pilT$_{XX}$^{WB}	-	-	-	-	-	-	Vc	-	-	-	-	-	-	Ec	-

Fig. 2. *The PilT homolog from E. coli cannot support PilU-mediated retraction*. (**A**) Natural transformation assays were performed to test whether retraction ATPase PilT mutants from 3 heterologous T4aP systems could complement *V. cholerae* mutants lacking their endogenous motor ATPases. The relevant genotypes of strains are indicated below each bar, which includes the status of the native *pilT/pilU* genes as well as the ectopically expressed motor ATPase allele (induced via P$_{BAD}$). The color of the bar corresponds to the species from which the ectopically expressed *pilT* was derived (green bars = *Vc pilT*, yellow bars = *Ab pilT*, orange bars = *Ps pilT*, and purple bars = *Ec pilT*). Data are from at least 3 independent biological replicates and shown as the mean ± SD. (**B**) BACTH assays to test whether the indicated PilT alleles can interact with themselves ("self") or PilU$_{Vc}$. For BACTH assays pilus components had an N-terminal fusion to the T25 or T18 fragment of adenylate cyclase as indicated. "^{WA}" denotes a mutation in the Walker A domain of the tested allele, while "^{WB}" denotes a mutation in the Walker B domain of the tested allele. "+" and "−" denote positive and negative controls for the BACTH assay. "E.V." denotes an empty vector. Data are representative of at least 2 independent experiments.

Fig. 3. *The C-terminus of PilTVc contributes to PilU-mediated retraction.* Natural transformation
assays were performed to test whether retraction ATPase chimeras between PilTVc and PilTEc can
complement *V. cholerae* mutants lacking their endogenous motor ATPases. PilT chimeras either have (**A**)
their ATPase activity intact to assess ATPase-dependent retraction, or their ATPase activity inactivated
via Walker A or Walker B mutations to assess PilU-mediated retraction. The relevant genotypes of strains
are indicated to the left of each bar, which includes the status of the native *pilT/pilU* genes as well as the
ectopically expressed motor ATPase chimera (induced via P_BAD). Diagrams to the left of bars denote the
composition of each chimera with the fragments of PilTVc denoted in green and PilTEc denoted in pink. For
strains that encode an ATPase active *pilT* allele, the bar is blue, while for strains that contain an ATPase
inactive *pilT* allele (via a WA or WB mutation as indicated) the bar is tan. The number in parentheses
indicates the number of N-terminal amino acids from the PilT allele indicated fused to its PilT ortholog. (**B**)
Natural transformation assays were performed to test whether two elements at the C-terminus of PilTVc
(the AIRNLIRE domain and a 15-residue domain at the C-terminus) contribute to PilU-mediated
retraction. The relevant genotypes of strains are indicated below each bar, which includes the status of

the native *pilT*/*pilU* genes as well as the ectopically expressed motor ATPase chimera (induced via P$_{BAD}$).
Four attributes of each ectopically expressed allele are indicated: the organism from which the PilT allele
is derived, the ATPase status of the allele ("+" denotes ATPase intact, "WA" denotes Walker A mutant,
and "WB" denotes Walker B mutant), the sequence of the AXXNLIRE domain (IR = AIRNLIRE, VG =
AVGNLIRE), and the presence / absence of the C-terminal 15 amino acids. If the attribute changes the
natural allele, it is displayed in red text. The color of the bar corresponds to the species from which the
ectopically expressed *pilT* was derived (green bars = *Vc pilT* and purple bars = *Ec pilT*). All data are from
at least 3 independent biological replicates and shown as the mean ± SD.

Fig. S1. *Expression of ectopic constructs is equal to or greater than native levels of expression.* Natural transformation assays were performed to confirm the functionality of the N-terminally FLAG-tagged alleles of **(A)** *pilT*, **(B)** *pilU*, and **(C)** *pilB*. The relevant genotypes of strains are indicated below each bar, which includes the status of the native *V. cholerae pilT/pilU/pilB* genes (top) as well as the ectopically expressed motor ATPase allele (induced via P_{BAD}) (bottom). The color of the bar corresponds to the species from which the ectopically expressed *motor ATPase* was derived (green bars = *Vc motors*, yellow bars = *Ab* motors, orange bars = *Ps* motors, and purple bars = *Ec* motors). Data are from at least 3 independent biological replicates and shown as the mean ± SD. The limit of detection for transformation frequency assays is ~10^{-7}. In **C**, Ab_1 denotes $pilB_{Ab}$ and Ab_2 denotes $tfpB_{Ab}$. In **C**, for all samples where the transformation frequency is ~10^{-7}, at least 1 replicate contained zero transformants and the limit of detection for that sample is displayed. Western blots to determine the levels of expression of the FLAG-tagged **(D)** *pilT*, **(E)** *pilU*, and **(F)** *pilB* alleles and RpoA as a loading control. The blots show that all ectopically expressed motors were expressed at or above the levels of the native *Vc* motor. Data is representative of at least two independent experiments.

Fig. S2. *PilU$_{Ps}$ is a fully functional retraction ATPase motor.* Natural transformation assays were performed to assess whether PilU$_{Ps}$ can support pilus retraction in conjunction with its cognate PilT$_{Ps}$^{WA} better than with the heterologous PilT$_{Vc}$^{WA}. The relevant genotypes of each strain are indicated below each bar, which includes the status of the native *pilT/pilU* genes as well as the ectopically expressed retraction motor ATPases. The color of the bar corresponds to the species from which the ectopically expressed PilT motor ATPase was derived (green bars = *Vc* motors, orange bar = *Ps* motors). Data are from at least 3 independent biological replicates and shown as the mean ± SD.

A

```
PilU Vc   MELNQYLDGMLTHKASDLYITVGAPILYRVDGELRAQGEALSVAD-VTALLHAMMDDARQAEFKQTREANFAVV-RD-SG  77
PilU Ab   MDFNDLLNLMAKHKASDLFITAGVEPSLKINGEITPVSKTKLDGDMIAQLIDSITTQKQRQEFIETHECNFAIINREKNE  80
PilU Ps   MEFEKLLRLMVEKGGSDLFITAGVPPSMKVNGKILPVNKTPMSPEMTRETVHGVMNEQQRREFTENHECNFAISARG-IG  79

PilU Vc   RFRVSAFFQRELPGAVIRRIETRIPTFEELKLPEVLQNLAIAKRGLVLVVGATGSGKSTTMAAMTGYRNQHRTGHILTVE  157
PilU Ab   RFRVSAFQQRDMPGMVIRRIETKIPSPDDLHLPPILKELAMSKRGIIIFSGATGTGKSTSLASMIQHRNQHSKGHIITIE  160
PilU Ps   RFRVSAFYQRNLAGMVLRRIETNIPTIEDLKLPDVLKKLALTKRGLVLFVGATGTGKSTSLAAMIGYRNKNSSGHIISIE  159

PilU Vc   DPIEFVHEHKRCIVTQREVGLDTESYEVALKNSLRQAPDMILIGEIRSRETMEYAMTFAETGHLCMATLHANNANQALER  237
PilU Ab   DPIEFIHEHAGCIITQREVGIDTDSFEIALKNTLRQAPDVILIGEIRSREVMDYALAFAETGHLVFATLHANNANQTLER  240
PilU Ps   DPIEFIHQHQNCIVTQREVGIDTDSFEVALKNTLRQAPDVILIGEVRTRETMDHAVAFAETGHLCLATLHANNANQALDR  239

PilU Vc   ILHLVPKEQREQFLLDLSLNLKGVIAQQLLRDKNGKGRHGVFEVLLNSPRIADLIRRGELHELKATMARSQEVGMQTFDQ  317
PilU Ab   IIHFFDADRHQQIFMDLSLNLKAVISQQLIPTHDGQSRRAAVEVLINTPLLSDYIRQGEIYKIKELMARSGELGMQTFDQ  320
PilU Ps   IINFFPADRQNQVWMDLSLNLKAIVAQQLVPTPDGKGRRAAVEVLINTPLAADLIRKGEVHELKGLMKRSTDLGMQTFDQ  319

PilU Vc   ALYQLVVDDKISEQDALHSADSANDLRLMLKTKRGDDYG--SGSL--QNVKIDME---------  368
PilU Ab   ALFDLYDHQHISYADALKHADSPNNLRLQIKLSEEANKP-LDNRERMSEMERKMTFDGQRRNLK  383
PilU Ps   ALYNLYVQGEITYEDALLHADSSNDLRLMIKLGSETDGEHLTSVS--QGLTLEMSDDDPGRSFR  381
```

B

```
PilT Vc   MDIAELLEFSVKHNASDLHLSAGVPPMVRIDGEVRKLGVPAFTHSDVHRLIFEIMNDAQRSEYEEKLEVDFSFELPNVGR  80
PilT Ab   MDITELLAFSVQNRASDLHLSAGIPPMIRVDGEVRRINLPAMDHSEVHNLIYDIMNDKQRRDYEEKLETDFSFEVPNLAR  80
PilT Ps   MDITELLAFSAKQGASDLHLSSGLPPMIRVDGDVRRINLPAMDHKQVHALIYDIMNDKQRKDFEEFLETDFSFEVPGVAR  80
PilT Ec   MNMEEIVALSVKHNVSDLHLCSAWPARWRIRGR---MEAAPFDTPDVEELLREWLDDDQRAILLENGQLDFAVSLAENQR  77

PilT Vc   FRVNAFHQARGCSAVFRTIPTVIPTLEQLDAPEIFSKIANYEKGLVLVTGPTGSGKSTTLAAMVNYVNAHHNKHILTIED  160
PilT Ab   FRVNVFNQNRGAGAVFRTIPSQVLTLEELGLGKIFRDICDYPRGLVLVTGPTGSGKSTTLAAMLDYINNHRYDHILTVED  160
PilT Ps   FRVNAFNQNRGAGAVFRTIPSKVLTMEDLGMGEVFRKITDVPRGLVLVTGPTGSGKSTTLAAMLDYLNNTKYHHILTIED  160
PilT Ec   LRGSAFAQRHGISLALDRLLPSHCPQLEQLGAPTVLPELLKSENGLILVTGATGSGKSTTLAAMVGYLNQHADAHILTLED  157

PilT Vc   PIEFVHSNNKCLINQREVHRDTHSFKNALRSALREDPDVILVGELRDQETISLALTAAETGHLVFGTLHTSSAAKTIDRI  240
PilT Ab   PIEFVHEPKKCLINQREVHRDTLGFNEALRSALREDPDIILVGEMRDLETIRLALTAAETGHLVFGTLHTTSAAKTIDRI  240
PilT Ps   PIEFVHESKKCLVNQREVHRDTLGFSEALRSALREDPDIILVGEMRDLETIRLALTAAETGHLVFGTLHTTSAAKTIDRV  240
PilT Ec   PVEYLYASQRCLIQQREIGLHCMTFASGLRAALREDPDVILLGELRDSETIRLALTAAETGHLVLATLHTRGAAQAVERL  237

PilT Vc   IDVFPGSDKDMVRSMLSESLRAVIAQKLLKRVGGGRVACHEIMLATP`AIRNLIRE`DKVAQMYSIQTGAAHGMQTMEQNA  320
PilT Ab   IDVFPAEEKDMVRTMLSESLQAVISQTLLKKEGGGRVAAHEIMIGTP`AIRNLIRE`GKVAQMYSSIQTGATHGMTTLDQSL  320
PilT Ps   VDVFPAEEKSMVRSMLSESLQAVISQTLLKKVGGGRVAAHEIMIGTP`AIRNLIRE`GKVAQMYSSIQTGGSLGMQTLDSCL  320
PilT Ec   VDSFPAQEKDPVRNQLAGSLRAVLSQKLEVDKQEGRVALFELLINTP`AVGNLIRE`GKTHQLPHVIQTGQQVGMITFQQSY  317
```

AIRNLIRE Domain

```
PilT Vc   KQLIARGVV`DAQEVQSKIELDLKAF`  345
PilT Ab   KQLLAKGII`NLPTARAVA-KSPESF`  344
PilT Ps   KGLLAKGLI`SRDSAKEKA-KQPENF`  344
PilT Ec   QHRVGEGRL`---------------`  326
```

C-term 15

Fig. S3. *Alignment of PilU and PilT sequences from the four T4aP studied.* Alignments of (**A**) PilU orthologs and (**B**) PilT orthologs. Residues highlighted in black are identical, while those highlighted in gray are similar. The AIRNLIRE and C-terminal 15 residue domains experimentally tested in this study are boxed in red. The GenBank Accession numbers for the protein sequences used are as follows: Vc, *Vibrio cholerae* PilT (AAF93635); Ab, *Acinetobacter baylyi* PilT (CAG67808); Ps, *Pseudomonas stutzeri* PilT (AFN76400); Ec, *Escherichia coli* PilT (NP_417425); Vc, *V. cholerae* PilU (AAF93636); Ab, *A. baylyi* PilU (CAG67807); Ps, *P. stutzeri* PilU (AFN76401).

A

	PilT$_{Vc}$	PilT$_{Ab}$	PilT$_{Ps}$	PilT$_{Ec}$
PilT$_{Vc}$	100%	83%	80%	62%
PilT$_{Ab}$	83%	100%	91%	60%
PilT$_{Ps}$	80%	91%	100%	59%
PilT$_{Ec}$	62%	60%	59%	100%

full length

B

	PilU$_{Vc}$	PilU$_{Ab}$	PilU$_{Ps}$
PilU$_{Vc}$	100%	70%	74%
PilU$_{Ab}$	70%	100%	78%
PilU$_{Ps}$	74%	78%	100%

full length

C

	PilB$_{Vc}$	PilB$_{Ab}$	TfpB$_{Ab}$	PilB$_{Ps}$	PilB$_{Ec}$
PilB$_{Vc}$	100%	64%	53%	66%	50%
PilB$_{Ab}$	64%	100%	55%	75%	48%
TfpB$_{Ab}$	53%	55%	100%	54%	45%
PilB$_{Ps}$	66%	75%	54%	100%	49%
PilB$_{Ec}$	50%	48%	45%	49%	100%

full length

	PilB$_{Vc}$	PilB$_{Ab}$	TfpB$_{Ab}$	PilB$_{Ps}$	PilB$_{Ec}$
PilB$_{Vc}$	100%	75%	64%	75%	64%
PilB$_{Ab}$	75%	100%	64%	86%	62%
TfpB$_{Ab}$	64%	64%	100%	61%	60%
PilB$_{Ps}$	75%	86%	61%	100%	63%
PilB$_{Ec}$	64%	62%	60%	63%	100%

N2D-CTD

	PilB$_{Vc}$	PilB$_{Ab}$	TfpB$_{Ab}$	PilB$_{Ps}$	PilB$_{Ec}$
PilB$_{Vc}$	100%	41%	32%	49%	20%
PilB$_{Ab}$	41%	100%	38%	52%	22%
TfpB$_{Ab}$	32%	38%	100%	41%	16%
PilB$_{Ps}$	49%	52%	41%	100%	21%
PilB$_{Ec}$	20%	22%	16%	21%	100%

N1D

Fig. S4. *Pairwise comparisons of the indicated motor ATPases.* Percent similarities are reported for pairwise comparisons of the indicated (**A**) PilT, (**B**) PilU, and (**C**) PilB/TfpB orthologs. The percent similarity is also denoted by the fill color of the box, where green represents high percent similarity and red represents lower percent similarity. PilB/TfpB orthologs were further analyzed for pairwise comparisons of the C-terminal domains that are shared between by extension and retraction motor ATPases (N2D and CTD) and the N-terminal domain (N1D) that is unique to extension motor ATPases (see **Fig. S5** for details on what portion of PilB represents the N1D vs N2D-CTD). The GenBank Accession numbers for the protein sequences used in these comparisons are as follows: Vc, *Vibrio cholerae* PilT (AAF93635); Ab, *Acinetobacter baylyi* PilT (CAG67808); Ps, *Pseudomonas stutzeri* PilT (AFN76400); Ec, *Escherichia coli* PilT (NP_417425); Vc, *V. cholerae* PilU (AAF93636); Ab, *A. baylyi* PilU (CAG67807); Ps, *P. stutzeri* PilU (AFN76401); Vc, *V. cholerae* PilB (AAF95567); Ab, *A. baylyi* PilB (CAG67312); Ab, *A. baylyi* TfpB (CAG69034); Ps, *P. stutzeri* PilB (AFN76867); Ec, *E. coli* PilB (AAC73218).

Fig. S5. *Alignment of the PilB/TfpB and PilT sequences from the T4aP studied.* Alignments
of PilB/TfpB orthologs and PilT orthologs. Residues highlighted in black are identical, while
those highlighted in gray are similar. The unique N-terminal domain in PilB orthologs is denoted
by the "N1D" bar above the sequence, while the two C-terminal domains shared by all motor

645   ATPases are denoted by "N2D-CTD". The GenBank Accession numbers for the protein
646   sequences used are as follows: Vc, *Vibrio cholerae* PilT (AAF93635); Ab, *Acinetobacter baylyi*
647   PilT (CAG67808); Ps, *Pseudomonas stutzeri* PilT (AFN76400); Ec, *Escherichia coli* PilT
648   (NP_417425); Vc, *V. cholerae* PilB (AAF95567); Ab, *A. baylyi* PilB (CAG67312); Ab, *A. baylyi*
649   TfpB (CAG69034); Ps, *P. stutzeri* PilB (AFN76867); Ec, *E. coli* PilB (AAC73218).

 Table S1. Strains used in this study

Strain name in manuscript	Genotype and antibiotic resistance	Figure	Strain # / Reference
Parent	E7946 SmR, ΔluxO::miniFRT, lacZ::lacIq, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX	Fig. 1, 2, 3, S1, S2	TND0905 (SAD2468) / (4)
ΔpilT	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔpilT::TmR	Fig. 1, 2, 3, S1	TND1035 (SAD2469) / (4)
ΔpilTU	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔpilTU::TmR	Fig. 1, 2, 3, S2	JLC227 (SAD2470) / (4)
P_{BAD}-pilT$_{Vc}$, ΔpilT	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Vc}$ CarbR, ΔpilT::TmR	Fig. 1, 3	TND1831 (SAD3199)
P_{BAD}-pilT$_{Vc}$, ΔpilTU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 1, 3	JLC1155 (SAD3222)
P_{BAD}-pilT$_{Ab}$, ΔpilT	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ab}$ CarbR, ΔpilT::TmR	Fig. 1	TND2148 (SAD3200)
P_{BAD}-pilT$_{Ab}$, ΔpilTU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ab}$ CarbR, ΔpilTU::TmR	Fig. 1	JLC1153 (SAD3223)
P_{BAD}-pilT$_{Ps}$, ΔpilT	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ps}$ CarbR, ΔpilT::TmR	Fig. 1	TND2125 (SAD3201)
P_{BAD}-pilT$_{Ps}$, ΔpilTU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ps}$ CarbR, ΔpilTU::TmR	Fig. 1	JLC1157 (SAD3224)
P_{BAD}-pilT$_{Ec}$, ΔpilT	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 1	JLC1473 (SAD3225)
P_{BAD}-pilT$_{Ec}$, ΔpilTU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-PilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 1	JLC1474 (SAD3226)
pilT^{WA}, ΔpilU	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, pilT^{K136A}, ΔpilU::TmR	Fig. 1, 2	JLC444 (SAD2472) / (4)
P_{BAD}-pilU$_{Vc}$, pilT^{WA}, ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}^{K136A}$, ΔlacZ:: P_{BAD}-PilU$_{Vc}$ CarbR, ΔpilU::TmR	Fig. 1	EHC062 (SAD3239)
P_{BAD}-pilU$_{Ab}$, pilT^{WA}, ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}^{K136A}$, ΔlacZ:: P_{BAD}-PilU$_{Ab}$ CarbR, ΔpilU::TmR	Fig. 1	EHC060 (SAD3240)
P_{BAD}-pilU$_{Ps}$, pilT^{WA}, ΔpilU	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}^{K136A}$, ΔlacZ:: P_{BAD}-PilU$_{Ps}$ CarbR, ΔpilU::TmR	Fig. 1	EHC061 (SAD3241)
Parent (P_{const}-tfoX)	E7946 SmR, ΔlacZ::lacIq, P_{const}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C	Fig. 1	TND1076 (SAD2513) / (4)
ΔpilB	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT	Fig. 1	TND2373 (SAD2787) / (13)

P_{BAD}-$pilB_{Vc}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P_{BAD}-pilB$_{Vc}$ CarbR	Fig. 1	TND2403 (SAD2789) / (13)
P_{BAD}-$pilB_{Ab}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P_{BAD}-pilB$_{Ab}$ CarbR	Fig. 1	TND2354 (SAD3202)
P_{BAD}-$tfpB_{Ab}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P_{BAD}-tfpB$_{Ab}$ CarbR	Fig. 1	TND2372 (SAD3203)
P_{BAD}-$pilB_{Ps}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P_{BAD}-pilB$_{Ps}$ CarbR	Fig. 1	TND2371 (SAD3204)
P_{BAD}-$pilB_{Ec}$, $\Delta pilB$	E7946 SmR, lacZ LPQEN:: KanR, P_{const}-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔlacZ:: P_{BAD}-pilB$_{Ec}$ CarbR	Fig. 1	EHC170 (SAD3283)
$pilT^{WA}$	E7946 SmR, ΔlacZ::lacIq, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT^{K136A}	Fig. 2	TND1077 (SAD2471) / (4)
P_{BAD}-$pilT_{Vc}^{WA}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Vc}^{K136A}$ CarbR, ΔpilT::TmR	Fig. 2, 3	JLC1609 (SAD3227)
P_{BAD}-$pilT_{Vc}^{WA}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Vc}^{K136A}$ CarbR, ΔpilTU::TmR	Fig. 2	EHC031 (SAD3242)
P_{BAD}-$pilT_{Vc}^{WB}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Vc}^{E204A}$ CarbR, ΔpilT::TmR	Fig. 2	EHC075 (SAD3243)
P_{BAD}-$pilT_{Ab}^{WA}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ab}^{K136A}$ CarbR, ΔpilT::TmR	Fig. 2	EHC015 (SAD3244)
P_{BAD}-$pilT_{Ab}^{WA}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ab}^{K136A}$ CarbR, ΔpilTU::TmR	Fig. 2	EHC019 (SAD3245)
P_{BAD}-$pilT_{Ps}^{WA}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ps}^{K136A}$ CarbR, ΔpilT::TmR	Fig. 2	EHC016 (SAD3246)
P_{BAD}-$pilT_{Ps}^{WA}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ps}^{K136A}$ CarbR, ΔpilTU::TmR	Fig. 2	EHC020 (SAD3247)
P_{BAD}-$pilT_{Ec}^{WA}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ec}^{K133A}$ CarbR, ΔpilT::TmR	Fig. 2	JLC1499 (SAD3228)
P_{BAD}-$pilT_{Ec}^{WA}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ec}^{K133A}$ CarbR, ΔpilTU::TmR	Fig. 2	JLC1498 (SAD3229)
P_{BAD}-$pilT_{Ec}^{WB}$, $\Delta pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ::P_{BAD}-pilT$_{Ec}^{E201A}$ CarbR, ΔpilT::TmR	Fig. 2, 3	EHC076 (SAD3248)
P_{BAD}-$pilT_{Vc}(50)$/$pilT_{Ec}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}$(50)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3069 (SAD3205)
P_{BAD}-$pilT_{Vc}(100)$/$pilT_{Ec}$, $\Delta pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-tfoX, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}$(100)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3070 (SAD3206)

P_{BAD}-*pilT*$_{Vc}$(150)/*pilT*$_{Ec}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}$(150)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3071 (SAD3207)
P_{BAD}-*pilT*$_{Vc}$(200)/*pilT*$_{Ec}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}$(200)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3073 (SAD3208)
P_{BAD}-*pilT*$_{Vc}$(300)/*pilT*$_{Ec}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}$(300)/pilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3072 (SAD3209)
P_{BAD}-*pilT*$_{Ec}$(50)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Ec}$(50)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3074 (SAD3210)
P_{BAD}-*pilT*$_{Ec}$(100)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Ec}$(100)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3075 (SAD3211)
P_{BAD}-*pilT*$_{Ec}$(150)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Ec}$(150)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3076 (SAD3212)
P_{BAD}-*pilT*$_{Ec}$(200)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Ec}$(200)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3077 (SAD3213)
P_{BAD}-*pilT*$_{Ec}$(250)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Ec}$(250)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3078 (SAD3214)
P_{BAD}-*pilT*$_{Ec}$(300)/*pilT*$_{Vc}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Ec}$(300)/pilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. 3	TND3079 (SAD3215)
P_{BAD}-*pilT*$_{Vc}^{WA}$(100)/*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}^{K136A}$(100)/pilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 3	TND3008 (SAD3216)
P_{BAD}-*pilT*$_{Vc}^{WA}$(200)/*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}^{K136A}$(200)/pilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 3	TND3010 (SAD3217)
P_{BAD}-*pilT*$_{Ec}^{WA}$(300)/*pilT*$_{Vc}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Ec}^{K133A}$(300)/pilT$_{Vc}$ CarbR, ΔpilT::TmR	Fig. 3	TND3018 (SAD3218)
P_{BAD}-*pilT*$_{Vc}^{WB}$(100)/*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}^{E204A}$(100)/pilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 3	EHC106 (SAD3249)
P_{BAD}-*pilT*$_{Vc}^{WB}$(200)/*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Vc}^{E204A}$(200)/pilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. 3	EHC108 (SAD3250)
P_{BAD}-*pilT*$_{Ec}^{WB}$(300)/*pilT*$_{Vc}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-3X Flag-pilT$_{Ec}^{E201A}$(300)/pilT$_{Vc}$ CarbR, ΔpilT::TmR	Fig. 3	EHC115 (SAD3251)
P_{BAD}-*pilT*$_{Vc}^{AIR \rightarrow AVG}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-pilT$_{Vc-}$$^{AIR \rightarrow AVG}$ CarbR, ΔpilTU::TmR	Fig. 3	JLC1622 (SAD3230)
P_{BAD}-*pilT*$_{Vc}^{WA, AIR \rightarrow AVG}$, Δ*pilT*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-pilT$_{Vc-}$$^{K136A, AIR \rightarrow AVG}$ CarbR, ΔpilT::TmR	Fig. 3	JLC1625 (SAD3231)
P_{BAD}-*pilT*$_{Vc}^{\Delta 15aa}$, Δ*pilTU*	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-*tfoX*, ΔlacZ:: P_{BAD}-pilT$_{Vc-}$$^{\Delta 15aa}$ CarbR, ΔpilTU::TmR	Fig. 3	TND2774 (SAD3219)

P_{BAD}-$pilT_{Vc}$^{WA, Δ15aa}, Δ$pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Vc}$-K136A, Δ15aa CarbR, ΔpilT::TmR	Fig. 3	JLC1611 (SAD3232)
P_{BAD}-$pilT_{Vc}$^{AIR→AVG, Δ15aa}, Δ$pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Vc}$-AIR→AVG, Δ15aa CarbR, ΔpilTU::TmR	Fig. 3	JLC1624 (SAD3233)
P_{BAD}-$pilT_{Vc}$^{WA, AIR→AVG, Δ15aa}, Δ$pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Vc}$-K136A, AIR→AVG, Δ15aa CarbR, ΔpilT::TmR	Fig. 3	JLC1627 (SAD3234)
P_{BAD}-$pilT_{Ec}$^{AVG→AIR}, Δ$pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-AVG→AIR CarbR, ΔpilTU::TmR	Fig. 3	JLC1614 (SAD3235)
P_{BAD}-$pilT_{Ec}$^{WA, AVG→AIR}, Δ$pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-K133A, AVG→AIR CarbR, ΔpilT::TmR	Fig. 3	JLC1615 (SAD3236)
P_{BAD}-$pilT_{Ec}$^{WB, AVG→AIR}, Δ$pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-E201A, AVG→AIR CarbR, ΔpilT::TmR	Fig. 3	EHC095 (SAD3252)
P_{BAD}-$pilT_{Ec}$^{+15aa}, Δ$pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-+15aa CarbR, ΔpilTU::TmR	Fig. 3	TND2730 (SAD3220)
P_{BAD}-$pilT_{Ec}$^{WA, +15aa}, Δ$pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-K133A, +15aa CarbR, ΔpilT::TmR	Fig. 3	TND2727 (SAD3221)
P_{BAD}-$pilT_{Ec}$^{WB, +15aa}, Δ$pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-E201A, +15aa CarbR, ΔpilT::TmR	Fig. 3	EHC093 (SAD3253)
P_{BAD}-$pilT_{Ec}$^{AVG→AIR, +15aa}, Δ$pilTU$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-AVG→AIR, +15aa CarbR, ΔpilTU::TmR	Fig. 3	JLC1620 (SAD3237)
P_{BAD}-$pilT_{Ec}$^{WA, AVG→AIR, +15aa}, Δ$pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-K133A, AVG→AIR, +15aa CarbR, ΔpilT::TmR	Fig. 3	JLC1617 (SAD3238)
P_{BAD}-$pilT_{Ec}$^{WB, AVG→AIR, +15aa}, Δ$pilT$	E7946 SmR, ΔluxO::miniFRT, pilA S56C, ΔVC1807::ZeoR, P_{tac}-$tfoX$, ΔlacZ:: P_{BAD}-pilT$_{Ec}$-E201A, AVG→AIR, +15aa CarbR, ΔpilT::TmR	Fig. 3	EHC097 (SAD3254)
P_{tac}-$pilT_{Vc}$^{WA}, P_{BAD}-$pilU_{Ps}$, Δ$pilTU$	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, ΔlacZ::P_{BAD}-PilU$_{Ps}$ CarbR, ΔVC0692::P_{tac}-PilT$_{Vc}$^{K136A} SpecR, ΔpilTU::TmR	Fig. S2	EHC099 (SAD3255)
P_{tac}-$pilT_{Ps}$^{WA}, P_{BAD}-$pilU_{Ps}$, Δ$pilTU$	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, ΔlacZ::P_{BAD}-PilU$_{Ps}$ CarbR, ΔVC0692::P_{tac}-PilT$_{Ps}$^{K136A} SpecR, ΔpilTU::TmR	Fig. S2	EHC100 (SAD3256)
P_{BAD}-3X FLAG-$pilT_{Vc}$, Δ$pilT$	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-3X FLAG-PilT$_{Vc}$ CarbR, ΔpilT::TmR	Fig. S1	EHC136 (SAD3285)
P_{BAD}-3X FLAG-$pilT_{Vc}$, Δ$pilTU$	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-3X FLAG-PilT$_{Vc}$ CarbR, ΔpilTU::TmR	Fig. S1	EHC137 (SAD3286)
P_{BAD}-3X FLAG-$pilT_{Ab}$, Δ$pilT$	E7946 SmR, P_{tac}-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P_{BAD}-3X FLAG-PilT$_{Ab}$ CarbR, ΔpilT::TmR	Fig. S1	EHC138 (SAD3287)

P$_{BAD}$-3X FLAG-*pilT*$_{Ab}$, Δ*pilTU*	E7946 SmR, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P$_{BAD}$-3X FLAG-PilT$_{Ab}$ CarbR, ΔpilTU::TmR	Fig. S1	EHC139 (SAD3288)
P$_{BAD}$-3X FLAG-*pilT*$_{Ps}$, Δ*pilT*	E7946 SmR, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P$_{BAD}$-3X FLAG-PilT$_{Ps}$ CarbR, ΔpilT::TmR	Fig. S1	EHC140 (SAD3289)
P$_{BAD}$-3X FLAG-*pilT*$_{Ps}$, Δ*pilTU*	E7946 SmR, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P$_{BAD}$-3X FLAG-PilT$_{Ps}$ CarbR, ΔpilTU::TmR	Fig. S1	EHC141 (SAD3290)
P$_{BAD}$-3X FLAG-*pilT*$_{Ec}$, Δ*pilT*	E7946 SmR, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P$_{BAD}$-3X FLAG-PilT$_{Ec}$ CarbR, ΔpilT::TmR	Fig. S1	TND2978 (SAD3291)
P$_{BAD}$-3X FLAG-*pilT*$_{Ec}$, Δ*pilTU*	E7946 SmR, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::ZeoR, pilA S56C, ΔlacZ::P$_{BAD}$-3X FLAG-PilT$_{Ec}$ CarbR, ΔpilTU::TmR	Fig. S1	TND2982 (SAD3292)
P$_{BAD}$-3X FLAG-*pilU*$_{Vc}$, *pilT*^{WA}, Δ*pilU*	E7946 SmR, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}$^{K136A}, ΔlacZ:: P$_{BAD}$-3X FLAG-PilU$_{Vc}$ CarbR, ΔpilU::TmR	Fig. S1	EHC161 (SAD3293)
P$_{BAD}$-3X FLAG-*pilU*$_{Ab}$, *pilT*^{WA}, Δ*pilU*	E7946 SmR, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}$^{K136A}, ΔlacZ:: P$_{BAD}$-3X FLAG-PilU$_{Ab}$ CarbR, ΔpilU::TmR	Fig. S1	EHC152 (SAD3294)
P$_{BAD}$-3X FLAG-*pilU*$_{Ps}$, *pilT*^{WA}, Δ*pilU*	E7946 SmR, P$_{tac}$-tfoX, ΔluxO::miniFRT, ΔVC1807::KanR, pilA S56C, pilT$_{Vc}$^{K136A}, ΔlacZ:: P$_{BAD}$-3X FLAG-PilU$_{Ps}$ CarbR, ΔpilU::TmR	Fig. S1	EHC145 (SAD3295)
P$_{BAD}$-3X FLAG-*pilB*$_{Vc}$, Δ*pilB*	E7946 SmR, lacZ LPQEN:: KanR, P$_{const}$-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P$_{BAD}$-3X FLAG-pilB$_{Vc}$ CarbR	Fig. S1	EHC166 (SAD3296)
P$_{BAD}$-3X FLAG-*pilB*$_{Ab}$, Δ*pilB*	E7946 SmR, lacZ LPQEN:: KanR, P$_{const}$-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P$_{BAD}$-3X FLAG-pilB$_{Ab}$ CarbR	Fig. S1	EHC167 (SAD3297)
P$_{BAD}$-3X FLAG-*tfpB*$_{Ab}$, Δ*pilB*	E7946 SmR, lacZ LPQEN:: KanR, P$_{const}$-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P$_{BAD}$-3X FLAG-tfpB$_{Ab}$ CarbR	Fig. S1	EHC168 (SAD3298)
P$_{BAD}$-3X FLAG-*pilB*$_{Ps}$, Δ*pilB*	E7946 SmR, lacZ LPQEN:: KanR, P$_{const}$-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔVCA0692:: P$_{BAD}$-3X FLAG-pilB$_{Ps}$ CarbR	Fig. S1	EHC169 (SAD3299)
P$_{BAD}$-3X FLAG-*pilB*$_{Ec}$, Δ*pilB*	E7946 SmR, lacZ LPQEN:: KanR, P$_{const}$-tfoX, ΔluxO::miniFRT, pilA S56C, ΔVC1807::CmR, ΔpilB::miniFRT, ΔlacZ:: P$_{BAD}$-3X FLAG-pilB$_{Ec}$ CarbR	Fig. S1	EHC172 (SAD3300)
T25-PilT$_{Vc}$	TG1, pkT25-PilT$_{Vc}$ KanR	Fig. 1, 2	SAD2248 / (4)
T25-PilT$_{Ab}$	TG1, pkT25-PilT$_{Ab}$ KanR	Fig. 1, 2	SAD2659
T25-PilT$_{Ps}$	TG1, pkT25-PilT$_{Ps}$ KanR	Fig. 1, 2	EHC043 (SAD3257)
T25-PilT$_{Ec}$	TG1, pkT25-PilT$_{Ec}$ KanR	Fig. 1, 2	EHC047 (SAD3258)
T18-PilT$_{Vc}$	TG1, pUT18C-PilT$_{Vc}$ CarbR	Fig. 1, 2	SAD2249 / (4)
T18-PilT$_{Ab}$	TG1, pUT18C-PilT$_{Ab}$ CarbR	Fig. 1, 2	SAD2660
T18-PilT$_{Ps}$	TG1, pUT18C-PilT$_{Ps}$ CarbR	Fig. 1, 2	EHC044

			(SAD3259)
T18-PilT$_{Ec}$	TG1, pUT18C-PilT$_{Ec}$ CarbR	Fig. 1, 2	EHC042 (SAD3260)
T25-PilU$_{Vc}$	TG1, pkT25-PilU$_{Vc}$ KanR	Fig. 1, 2	SAD2250 / (4)
T18-PilU$_{Vc}$	TG1, pUT18C-PilU$_{Vc}$ CarbR	Fig. 1, 2	SAD2251 / (4)
T25-PilC$_{Vc}$	TG1, pkT25-PilC$_{Vc}$ KanR	Fig. 1	SAD2260 / (4)
T18-PilC$_{Vc}$	TG1, pUT18C-PilC$_{Vc}$ CarbR	Fig. 1	SAD2261 / (4)
T25-PilU$_{Ab}$	TG1, pkT25-PilU$_{Ab}$ KanR	Fig. 1	EHC053 (SAD3261)
T25-PilU$_{Ps}$	TG1, pkT25-PilU$_{Ps}$ KanR	Fig. 1	EHC048 (SAD3262)
T18-PilU$_{Ab}$	TG1, pUT18C-PilU$_{Ab}$ CarbR	Fig. 1	SAD2664
T18-PilU$_{Ps}$	TG1, pUT18C-PilU$_{Ps}$ CarbR	Fig. 1	EHC064 (SAD3263)
T25-PilB$_{Vc}$	TG1, pkT25-PilB$_{Vc}$ KanR	Fig. 1	EHC052 (SAD3264) / (14)
T25-PilB$_{Ab}$	TG1, pkT25-PilB$_{Ab}$ KanR	Fig. 1	SAD2647 / (14)
T25-TfpB$_{Ab}$	TG1, pkT25-TfpB$_{Ab}$ KanR	Fig. 1	SAD2878
T25-PilB$_{Ps}$	TG1, pkT25-PilB$_{Ps}$ KanR	Fig. 1	SAD2959
T18-PilB$_{Vc}$	TG1, pUT18C-PilB$_{Vc}$ CarbR	Fig. 1	EHC050 (SAD3265) / (14)
T18-PilB$_{Ab}$	TG1, pUT18C-PilB$_{Ab}$ CarbR	Fig. 1	SAD2648
T18-TfpB$_{Ab}$	TG1, pUT18C-TfpB$_{Ab}$ CarbR	Fig. 1	SAD2879
T18-PilB$_{Ps}$	TG1, pUT18C-PilB$_{Ps}$ CarbR	Fig. 1	SAD2960
T25-PilT$_{Vc}^{WA}$	TG1, pkT25-PilT$_{Vc}^{K136A}$ KanR	Fig. 1, 2	SAD2285 / (4)
T25-PilT$_{Ab}^{WA}$	TG1, pkT25-PilT$_{Ab}^{K136A}$ KanR	Fig. 2	EHC054 (SAD3266)
T25-PilT$_{Ps}^{WA}$	TG1, pkT25-PilT$_{Ps}^{K136A}$ KanR	Fig. 2	EHC056 (SAD3267)
T25-PilT$_{Ec}^{WA}$	TG1, pkT25-PilT$_{Ec}^{K133A}$ KanR	Fig. 2	EHC057 (SAD3268)
T18-PilT$_{Vc}^{WA}$	TG1, pUT18C-PilT$_{Vc}^{K136A}$ CarbR	Fig. 1, 2	SAD2286 / (4)
T18-PilT$_{Ab}^{WA}$	TG1, pUT18C-PilT$_{Ab}^{K136A}$ CarbR	Fig. 2	EHC055 (SAD3269)
T18-PilT$_{Ps}^{WA}$	TG1, pUT18C-PilT$_{Ps}^{K136A}$ CarbR	Fig. 2	EHC059 (SAD3270)
T18-PilT$_{Ec}^{WA}$	TG1, pUT18C-PilT$_{Ec}^{K133A}$ CarbR	Fig. 2	EHC058 (SAD3271)
T25-PilT$_{Vc}^{WB}$	TG1, pkT25-PilT$_{Vc}^{E204A}$ KanR	Fig. 2	SAD2287
T25-PilT$_{Ec}^{WB}$	TG1, pkT25-PilT$_{Ec}^{E201A}$ KanR	Fig. 2	EHC078 (SAD3272)
T18-PilT$_{Vc}^{WB}$	TG1, pUT18C-PilT$_{Vc}^{E204A}$ CarbR	Fig. 2	SAD2288
T18-PilT$_{Ec}^{WB}$	TG1, pUT18C-PilT$_{Ec}^{E201A}$ CarbR	Fig. 2	EHC079 (SAD3273)
T25-PilB$_{Ec}$	TG1, pkT25-PilB$_{Ec}$ KanR	Fig. 1	EHC142

			(SAD3301)
T18-PilB_Ec	TG1, pUT18C-PilB_Ec CarbR	Fig. 1	EHC143 (SAD3302)
T25-Zip	TG1, pkT25-leucine zipper KanR	Fig. 1, 2	SAD2238 / (4)
T18-Zip	TG1, pUT18C-leucine zipper CarbR	Fig. 1, 2	SAD2241 / (4)
T25-vector	TG1, pkT25 vector KanR	Fig. 1, 2	SAD2236 / (4)
T18-vector	TG1, pUT18C vector CarbR	Fig. 1, 2	SAD2240 / (4)
Acinetobacter baylyi	ADP1	Purified gDNA	SAD631 / (23)
Pseudomonas stutzeri	JM300	Purified gDNA	SAD2178 / (24)
Escherichia coli	MG1655	Purified gDNA	SAD720 / (25)

653 **Table S2. Primers used in this study**

Primer name	Sequence (5' to 3')	Description
BBC374	TGGCAAAAAGCGAGAGAAGAAG	ΔluxO F1
BBC375	gtcgacggatccccggaatCATGAGGACATATTTTGTTTTCTGC	ΔluxO R1
BBC376	gaagcagctccagcctacaTAAGCGATGAGAGAATGGATCAAC	ΔluxO F2
BBC377	TCACACCCGAATTTCCATCATGC	ΔluxO R2
ABD344	GATTAGCAACGATTCTAGCGCAGGAG	ΔVC1807 F1
ABD340	gtcgacggatccccggaatACGTTTCATTAGTCACCTCTATTGT TAACTTGTTC	ΔVC1807 R1
ABD341	gaagcagctccagcctacaTAGTCGAAAATAAAAAAAGAGG CTCGCCTC	ΔVC1807 F2
ABD345	CTTGCTAACCGTTGGTGTTACCAGC	ΔVC1807 R2
DOG0400	ACTTCTGGCTGAAGGTCAATTTTC	Amplify ΔpilT/U mutations F1
DOG0403	TCACGTGTTCGGCCAAAATC	Amplify ΔpilT/U mutations R2
ABD332	GGCTGAACGTGGTTGTCGAAAATGAC	P_BAD-XX @ lacZ F1
ABD256	CCCAAATACGGCAACTTGGCG	P_BAD-XX @ lacZ R2
BBC832	GCTTTTTGCTACAACGACCG	P_tac-XX @ VCA0692 F1
BBC834	CACAATTTCTCGCTTAAAATGTCC	P_tac-XX @ VCA0692 R2
BBC1723	caatttcacacaggatcccgggAGGAGGTATTTAAATGGATATCGCTGAGTTAC	Amplify pilT_Vc for P_BAD-pilT or P_tac-pilT F
BBC1724	tgtaggctggagctgcttcCTAAAATGCTTTTAAATCCAACTCG	Amplify pilT_Vc for P_BAD-pilT or P_tac-pilT R
BBC2807	caatttcacacaggatcccgggAGGAGGTGAAAATATGGATATAACAGAATTACTCG	Amplify pilT_Ab for P_BAD-pilT F

BBC2808	tgtaggctggagctgcttcTTAGAACGACTCTGGAGATTTTG	Amplify $pilT_{Ab}$ for P_{BAD}-pilT R
BBC3143	caatttcacacaggatcccgggAGGAGGTAGTCCCATGGATATCACAGAAC	Amplify $pilT_{Ps}$ for P_{BAD}-pilT or P_{tac}-pilT F
BBC3144	tgtaggctggagctgcttcTTATCGGTTGTTAAAAATTTTCCGG	Amplify $pilT_{Ps}$ for P_{BAD}-pilT or P_{tac}-pilT R
BBC3140	caatttcacacaggatcccgggAGGAGGTCTTAGTATGAATATGGAAGAAATTGTGG	Amplify $pilT_{Ec}$ for P_{BAD}-pilT F
BBC3141	tgtaggctggagctgcttcCATGGCAACGTCACAAACG	Amplify $pilT_{Ec}$ for P_{BAD}-pilT R
HPQ0001	GGGCTCGGGTgctTCGACGACCTTGGCG	$pilT_{Vc}^{K136A}$ R1
HPQ0002	CGCCAAGGTCGTCGAagcACCCGAGCCC	$pilT_{Vc}^{K136A}$ F2
BBC2403	GCTGCAAGGGTGGTTGAagcACCGGAACCAGTGG	$pilT_{Ab}^{K136A}$ R1
BBC2404	CCACTGGTTCCGGTgctTCAACCACCCTTGCAGC	$pilT_{Ab}^{K136A}$ F2
JCP135	CGCCAGAGTGGTGGAcgcGCCCGAACCGGTCGG	$pilT_{Ps}^{K136A}$ R1
JCP134	CCGACCGGTTCGGGCgcgTCCACCACTCTGGCG	$pilT_{Ps}^{K136A}$ F2
JCP132	CGCCAGCGTGGTAGAcgcGCCACTCCCCGTCGC	$pilT_{Ec}^{K133A}$ R1
JCP131	GCGACGGGGAGTGGCgcgTCTACCACGCTGGCG	$pilT_{Ec}^{K133A}$ F2
BBC2508	GTTTCTTGGTCACGCAGagCACCGACGAGGATCAC	$pilT_{Vc}^{E204A}$ R1
BBC2509	GTGATCCTCGTCGGTGctCTGCGTGACCAAGAAAC	$pilT_{Vc}^{E204A}$ F2
BBC4065	CTCGCTGTCACGCAGagCTCCGAGCAAAATCACATCAGGATCTTCCCG	$pilT_{Ec}^{E201A}$ R1
BBC4066	GCTCGGAGctCTGCGTGACAGCGAGACAATCCGTCTGGCACTGAC	$pilT_{Ec}^{E201A}$ F2
BBC3897	CAAGCGATGTACATCAGAATGGG	$pilT_{Vc}$(50)/$pilT_{Ec}$ R1
BBC3898	cccattctgatgtacatcgcttgCTGCGGGAGTGGCTGGATG	$pilT_{Vc}$(50)/$pilT_{Ec}$ F2
BBC3899	GGGAATGGTACGAAACACCGCC	$pilT_{Vc}$(100)/$pilT_{Ec}$ R1
BBC3900	ggcggtgtttcgtaccattcccTCGCACTGCCCGCAGCTCGAAC	$pilT_{Vc}$(100)/$pilT_{Ec}$ F2
BBC3901	ATGAGCATTGACATAGTTCACC	$pilT_{Vc}$(150)/$pilT_{Ec}$ R1
BBC3902	ggtgaactatgtcaatgctcatGCCGATGCGCATATTCTGACG	$pilT_{Vc}$(150)/$pilT_{Ec}$ F2
BBC3903	GATCACATCTGGGTCTTCACGC	$pilT_{Vc}$(200)/$pilT_{Ec}$ R1
BBC3904	gcgtgaagacccagatgtgatcTTGCTCGGAGAGCTGCGTG	$pilT_{Vc}$(200)/$pilT_{Ec}$ F2
BBC3907	GTCTTTGTCGCTGCCCGGAAAAAC	$pilT_{Vc}$(300)/$pilT_{Ec}$ R1
BBC3908	gttttttccgggcagcgacaaagacCCCGTGCGTAATCAACTGGC	$pilT_{Vc}$(300)/$pilT_{Ec}$ F2
BBC3909	CTGCGCCACTTTATCTTCACGG	$pilT_{Ec}$(50)/$pilT_{Vc}$ R1
BBC3910	ccgtgaagataaagtggcgcagTTACCGCATGTTATTCAAACC	$pilT_{Ec}$(50)/$pilT_{Vc}$ F2
BBC3911	CTCCCGCAGTAGCTCTTCG	$pilT_{Ec}$(100)/$pilT_{Vc}$ R1
BBC3912	cgaagagctactgcgggagATCATGAATGATGCCCAGCGC	$pilT_{Ec}$(100)/$pilT_{Vc}$ F2
BBC3913	GCAGTGCGAAGGTAACAGCCG	$pilT_{Ec}$(150)/$pilT_{Vc}$ R1
BBC3914	cggctgttaccttcgcactgcCCGACGCTAGAGCAACTGG	$pilT_{Ec}$(150)/$pilT_{Vc}$ F2
BBC3915	CGCATCGGCATGTTGATTGAG	$pilT_{Ec}$(200)/$pilT_{Vc}$ R1
BBC3916	ctcaatcaacatgccgatgcgCATATTTTGACCATTGAAGATCCG	$pilT_{Ec}$(200)/$pilT_{Vc}$ F2
BBC3917	TCCGAGCAAAATCACATCAGG	$pilT_{Ec}$(250)/$pilT_{Vc}$ R1
BBC3918	cctgatgtgattttgctcggaGAGCTGCGTGACCAAGAAACG	$pilT_{Ec}$(250)/$pilT_{Vc}$ F2
BBC3919	ACGCACtGGGTCTTTTTCC	$pilT_{Ec}$(300)/$pilT_{Vc}$ R1
BBC3920	ggaaaaagacccagtgcgtTCCATGTTGTCGGAATCGC	$pilT_{Ec}$(300)/$pilT_{Vc}$ F2
DOG0565	AGATTGGTCTATTTTATGGCTCAAG	ΔpilB F1
DOG0570	gtcgacggatccccggaatCATTAGTTAATTGTTGCACCTGC	ΔpilB R1
DOG0571	gaagcagctccagcctacaCAGCGTGTGCTCTACTTTTAACTG	ΔpilB F2
DOG0572	AAAGAGTTGATGTTGAATCCCATC	ΔpilB R2
BBC1725	caatttcacacaggatcccgggAGGAGGTAGACAGATGGAGTTGAATCAATATC	Amplify $pilU_{Vc}$ for P_{BAD}-pilU F

BBC1726	tgtaggctggagctgcttcCTACTCCATATCAATCTTCACATTC	Amplify $pilU_{Vc}$ for P_{BAD}-pilU R
BBC2809	caatttcacacaggatcccgggAGGAGGTAATACGATGGACTTTAATGATTTACTC	Amplify $pilU_{Ab}$ for P_{BAD}-pilU F
BBC2810	tgtaggctggagctgcttcTTATTTGAGATTACGGCGTTGAC	Amplify $pilU_{Ab}$ for P_{BAD}-pilU R
JCP177	caatttcacacaggatcccgggAGGAGGTACGACGATGGAATTCGAGAAAC	Amplify $pilU_{Ps}$ for P_{BAD}-pilU F
JCP178	tgtaggctggagctgcttcGGCCTGTCAGCGGAAACTG	Amplify $pilU_{Ps}$ for P_{BAD}-pilU R
ABD334	AGTGCTCCGGACTCTTTGCTCTG	P_{BAD}-pilU @ lacZ F1
BBC1747	AACATCCGTTAGACAATGACC	P_{BAD}-pilU @ lacZ R2
BBC1952	caatttcacacaggatcccgggAGGAGGTTAACTAATGCTCACCAACCTTGTTG	Amplify $pilB_{Vc}$ for P_{BAD}-pilB F
BBC1953	tgtaggctggagctgcttcTTAAAAGTAGAGCACACGCTG	Amplify $pilB_{Vc}$ for P_{BAD}-pilB R
BBC3320	caatttcacacaggatcccgggaggaggtATTATAATGATATCAGAATCACAAGGCGG	Amplify $pilB_{Ab}$ for P_{BAD}-pilB F
BBC3310	tgtaggctggagctgcttcTTATTCACTGGTTACACGATTAATTTCC	Amplify $pilB_{Ab}$ for P_{BAD}-pilB R
BBC3327	caatttcacacaggatcccgggaggaggtAAATAAATGACATATCATTTTGAAATAGATA	Amplify $tfpB_{Ab}$ for P_{BAD}-tfpB F
BBC3328	tgtaggctggagctgcttcCTAAGTTAATGGGACCACCC	Amplify $tfpB_{Ab}$ for P_{BAD}-tfpB R
BBC3324	caatttcacacaggatcccgggaggaggtAACCTCATGAGCGAAAACATTTCCC	Amplify $pilB_{Ps}$ for P_{BAD}-pilB F
BBC3325	tgtaggctggagctgcttcTTAGTCCTTTGTCACCCGG	Amplify $pilB_{Ps}$ for P_{BAD}-pilB R
ECP0040	caatttcacacaggatcccgggAGGAGGTGCGGCAATGAATATTCCACAG	Amplify $pilB_{Ec}$ for P_{BAD}-$pilB_{Ec}$ F
ECP0041	tgtaggctggagctgcttcTTACTCGCCATGCGGCATC	Amplify $pilB_{Ec}$ for P_{BAD}-$pilB_{Ec}$ R
BBC2513	gaagcagctccagcctacaGATCCCCGGGTACCTAAGTAAC	Amplify BACTH pKT25-X to make T25-XX vectors F
BBC2512	tccacctgctccacctgcCTCTAGAGTCGACCCTGCAG	Amplify BACTH pKT25-X to make T25-XX vectors R
BBC2515	gaagcagctccagcctacaCCGAGCTCGAATTCATCGATATAAC	Amplify BACTH pUT18C to make T18-XX vectors F
BBC2514	tccacctgctccacctgcCTCTAGAGTCGACCTGCAG	Amplify BACTH pUT18C to make T18-XX vectors R
BBC2518	gcaggtggagcaggtggaAAAGCGACCCAAACCTTACC	BACTH TXX-$PilC_{Vc}$ F
BBC2524	tgtaggctggagctgcttcTTAACCCAATACACTCATTAAGTTAAAG	BACTH TXX-$PilC_{Vc}$ R
JCP027	gcaggtggagcaggtggaGATATCGCTGAGTTACTGGAATTTAG	BACTH TXX-$PilT_{Vc}$ F
BBC1724	tgtaggctggagctgcttcCTAAAATGCTTTTAAATCCAACTCG	BACTH TXX-$PilT_{Vc}$ R
BBC3194	gcaggtggagcaggtggaGATATAACAGAATTACTCGCGTT	BACTH TXX-$PilT_{Ab}$ F

BBC3195	tgtaggctggagctgcttcTTAGAACGACTCTGGAGATTTTG	BACTH TXX-PilT$_{Ab}$ R
BBC3921	gcaggtggagcaggtggaGATATCACAGAACTGCTGGCC	BACTH TXX-PilT$_{Ps}$ F
BBC3922	tgtaggctggagctgcttcTTAAAAATTTTCCGGCTGCTTGG	BACTH TXX-PilT$_{Ps}$ R
BBC3923	gcaggtggagcaggtggaAATATGGAAGAAATTGTGGCCC	BACTH TXX-PilT$_{Ec}$ F
BBC3924	tgtaggctggagctgcttcTCACAAACGCCCTTCCCCCAC	BACTH TXX-PilT$_{Ec}$ R
BBC2638	gcaggtggagcaggtggaCTCACCAACCTTGTTGCTATC	BACTH TXX-PilB$_{Vc}$ F
BBC2639	tgtaggctggagctgcttcTTAAAAGTAGAGCACACGCTG	BACTH TXX-PilB$_{Vc}$ R
BBC3182	gcaggtggagcaggtggaTCAGCATTTACAACACCACC	BACTH TXX-PilB$_{Ab}$ F
BBC3183	tgtaggctggagctgcttcTTATTCACTGGTTACACGATTAAT	BACTH TXX-PilB$_{Ab}$ R
JCP188	gcaggtggagcaggtggaACATATCATTTTGAAATAGATACACAATG	BACTH TXX-tfpB$_{Ab}$ F
BBC3328	tgtaggctggagctgcttcCTAAGTTAATGGGACCACCC	BACTH TXX-tfpB$_{Ab}$ R
BBC3890	gcaggtggagcaggtggaAGCGAAAACATTTCCCTTTCC	BACTH TXX-PilB$_{Ps}$ F
BBC3891	tgtaggctggagctgcttcTTAGTCCTTTGTCACCCGGTTG	BACTH TXX-PilB$_{Ps}$ R
ECP0043	gcaggtggagcaggtggaAATATTCCACAGCTCACTGC	BACTH TXX-PilB$_{Ec}$ F
ECP0044	tgtaggctggagctgcttcTTACTCGCCATGCGGCATC	BACTH TXX-PilB$_{Ec}$ R
BBC2434	gcaggtggagcaggtggaGAGTTGAATCAATATCTGGATGG	BACTH TXX-PilU$_{Vc}$ F
BBC1726	tgtaggctggagctgcttcCTACTCCATATCAATCTTCACATTC	BACTH TXX-PilU$_{Vc}$ R
BBC3197	gcaggtggagcaggtggaGACTTTAATGATTTACTCAATTTGATGG	BACTH TXX-PilU$_{Ab}$ F
BBC3198	tgtaggctggagctgcttcTTATTTGAGATTACGGCGTTGACC	BACTH TXX-PilU$_{Ab}$ R
BBC3981	gcaggtggagcaggtggaGAATTCGAGAAACTGTTGCG	BACTH TXX-PilU$_{Ps}$ F
JLC178	tgtaggctggagctgcttcGGCCTGTCAGCGGAAACTG	BACTH TXX-PilU$_{Ps}$ R
JCP318	GATTAACACACCCGCGaTtcGtAATTTGATTCGCGAAG	pilT$_{Ec}^{AVG \rightarrow AIR}$ F
JCP319	CTTCGCGAATCAAATTaCgaAtCGCGGGTGTGTTAATC	pilT$_{Ec}^{AVG \rightarrow AIR}$ R
JCP321	CTGGCGACGCCGGCAgTggGtAACTTGATCCGTGAAG	pilT$_{Vc}^{AIR \rightarrow AVG}$ F
JCP322	CTTCACGGATCAAGTTaCccAcTGCCGGCGTCGCCAG	pilT$_{Vc}^{AIR \rightarrow AVG}$ R
JCP264	tgtacttcttgcgcatcCAAACGCCCTTCCCCCAC	pilT$_{Ec}^{+15aa}$ R1
JCP259	ttggatttaaaagcatttTGAgaagcagctccagcctac	pilT$_{Ec}^{+15aa}$ F2
JCP262	gtaggctggagctgcttcCTACACCACACCACGCGCGATC	pilT$_{Vc}^{\Delta 15aa}$ R1
JCP263	gatcgcgcgtggtgtggtgTAGgaagcagctccagcctac	pilT$_{Vc}^{\Delta 15aa}$ F2
ECP0004	tccaccacttccacctgcCATGGGACTACCTCCTcccgggatcctg	3X FLAG-pilT$_{Ps}$ R1
ECP0005	gcaggtggagcaggtggaGATATCACAGAACTGCTGGCC	3X FLAG-pilT$_{Ps}$ F2
ECP0010	tccaccacttccacctgcCATATTTTCACCTCCTcccgggatcctg	3X FLAG-pilT$_{Ab}$ R1
ECP0011	gcaggtggagcaggtggaGATATAACAGAATTACTCGCGTTTTC	3X FLAG-pilT$_{Ab}$ F2

ECP0016	tccaccacttccacctgcCATCTGTCTACCTCCTcccgggatcctg	3X FLAG-pilU$_{Vc}$ R1
ECP0017	gcaggtggagcaggtggaGAGTTGAATCAATATCTGGATGGC	3X FLAG-pilU$_{Vc}$ F2
ECP0019	tccaccacttccacctgcCATTTAAATACCTCCTCCCGGGATCCTG	3X FLAG-pilT$_{Vc}$ R1
ECP0020	gcaggtggagcaggtggaGATATCGCTGAGTTACTGGAATTTAG	3X FLAG-pilT$_{Vc}$ F2
ECP0022	tccaccacttccacctgcCATCGTATTACCTCCTcccgggatcct	3X FLAG-pilU$_{Ab}$ R1
ECP0023	gcaggtggagcaggtggaGACTTTAATGATTTACTCAATTTGATGGC	3X FLAG-pilU$_{Ab}$ F2
ECP0025	tccaccacttccacctgcCATCGTCGTACCTCCTcccgggatcct	3X FLAG-pilU$_{Ps}$ R1
ECP0026	gcaggtggagcaggtggaGAATTCGAGAAACTGTTGCGC	3X FLAG-pilU$_{Ps}$ F2
ECP0028	tccaccacttccacctgcCATTAGTTAACCTCCTcccgggatcct	3X FLAG-pilB$_{Vc}$ R1
ECP0029	gcaggtggagcaggtggaCTCACCAACCTTGTTGCTATC	3X FLAG-pilB$_{Vc}$ F2
ECP0031	tccaccacttccacctgcCATTAGACCacctcctcccgggatcct	3X FLAG-pilB$_{Ab}$ R1
ECP0032	gcaggtggagcaggtggaTCAGCATTTACAACACCACCTAAGTTTTC	3X FLAG-pilB$_{Ab}$ F2
ECP0034	tccaccacttccacctgcCATTTATTTacctcctcccgggatcct	3X FLAG-tfpB$_{Ab}$ R1
ECP0035	gcaggtggagcaggtggaACATATCATTTTGAAATAGATACACAATGG	3X FLAG-tfpB$_{Ab}$ F2
ECP0037	tccaccacttccacctgcCATGAGGTTacctcctcccgggatcct	3X FLAG-pilB$_{Ps}$ R1
ECP0038	gcaggtggagcaggtggaAGCGAAAACATTTCCCTTTCC	3X FLAG-pilB$_{Ps}$ F2
ECP0045	tccaccacttccacctgcCATTGCCGCacctcctcccgggatcct	3X FLAG-pilB$_{Ec}$ R1
ECP0046	gcaggtggagcaggtggaAATATTCCACAGCTCACTGCCCTG	3X FLAG-pilB$_{Ec}$ F2