Multiple paths lead to salt tolerance - pre-adaptation vs dynamic responses from two closely related extremophytes

Kieu-Nga Tran, Guannan Wang, Dong-Ha Oh, John C. Larkin, Aaron P. Smith, Maheshi Dassanayake*

Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA
*Address correspondence to: maheshid@lsu.edu

Keywords: Extremophyte, *Schrenkiella parvula*, *Eutrema salsugineum*, omics, convergent response, pre-adaptation, stress-preparedness, dynamic response

Abstract

Salinity stress is an ongoing problem for global crop production. *Schrenkiella parvula* and *Eutrema salsugineum* are salt-tolerant extremophytes closely related to *Arabidopsis thaliana*. We investigated multi-omics salt stress responses of the two extremophytes in comparison to *A. thaliana*. Our results reveal that *S. parvula* limits Na accumulation while *E. salsugineum* shows high tissue tolerance to excess Na. Despite this difference, both extremophytes maintained their nutrient balance, while *A. thaliana* failed to sustain its nutrient content. The root metabolite profiles of the two extremophytes, distinct at control conditions, converged upon prolonged salt stress. This convergence was achieved by a dynamic response in *S. parvula* roots increasing its amino acids and sugars to the constitutively high basal levels observed in *E. salsugineum*. The metabolomic adjustments were strongly supported by the transcriptomic responses in the extremophytes. The predominant transcriptomic signals in all three species were associated with salt stress. However, root architecture modulation mediated by negative regulators of auxin and ABA signaling supported minimally affected root growth unique to each extremophyte during salt treatments. Overall, *E. salsugineum* exhibited more preadapted responses at the metabolome level while *S. parvula* showed predominant pre-adaptation at the transcriptome level to salt stress. Our work shows that while salt

tolerance in these two species shares common features, they substantially differ in pathways

leading to convergent adaptive traits.

Introduction

Plants differ greatly in their tolerance to salt stress and there is a metabolic cost for

adaptation to salt stress, reflected by differences in growth and yield when grown in high saline

soils (Santiago-Rosario et al., 2021; Flowers et al., 2015). Only ~2 % of angiosperms are adapted

to grow in high saline environments while the remaining 98 %, including most crop plants, are

highly sensitive to salt stress (Flowers and Colmer, 2008; Hajiboland et al., 2018). Previous

studies have largely used salt-sensitive model plants or crops to understand genetic

mechanisms underlying salt tolerance. Insight gained from examining the molecular processes

in salt tolerant extremophytes contrasted to a salt-sensitive model offers a promising prospect

to discover tolerance mechanisms that are absent in the salt-sensitive plants and proven to be

evolutionary successful against salt stress.

Schrenkiella parvula and *Eutrema salsugineum* (Brassicaceae) are two extremophyte

models with foundational genomic resources made available. They are also closely related to

Arabidopsis thaliana making it feasible to readily deduce orthologous relationships among the

three models (Dassanayake et al., 2011; Wu et al., 2012). These extremophytes, currently used

as leading models to investigate genetic mechanisms underlying salt stress adaptation

(Kazachkova et al., 2018), show salt resilient growth even at salinities reaching seawater

strengths (Inan et al., 2004; Orsini et al., 2010; Kazachkova et al., 2018). While *S. parvula* is

found near salt lakes in the Irano-Turanian region (Hajiboland et al., 2018; Tug et al., 2019), *E.*

salsugineum has a wider distribution from coastal to inland saline fields in the northern

temperate to sub-arctic regions including the United States, Canada, Russia, and China (Lee et

al., 2016). Despite multiple studies highlighting some of the metabolomic and transcriptomic

responses of the extremophytes to salt stress (Gong et al., 2005; Kazachkova et al., 2013; Oh et

al., 2014; Lee et al., 2016; Prerostova et al., 2017), molecular phenotypes determining how both

extremophytes have convergently achieved salt adapted growth unlike their salt stress-

sensitive relative, *A. thaliana* have not been explored.

Excess salt exerts cellular stress as osmotic, oxidative, ionic, and water-deficit stresses

(Pantha and Dassanayake, 2020; Zhao et al., 2020; van Zelm et al., 2020). Multiple genetic

mechanisms mediated by ABA as well as non-ABA dependent pathways have been shown to

modulate salt stress responses in *A. thaliana*, all major crops, and selected halophytes

(Takahashi and Shinozaki, 2019; Takahashi et al., 2020; Zhao et al., 2020). These studies

collectively support the view that salt stress-adapted plants will show a highly coordinated

response to survive salt stress that requires synergistic coordination between root and shoot

responses. Despite the complexity of adapting to salt stress at the whole plant level, multiple

angiosperm lineages have evolved this complex trait repeatedly (Flowers et al., 2010; Bennett

et al., 2013). Therefore, it is one of the main complex traits studied for convergent evolution in

plants. Even if a large body of work exist to identify many of the individual cellular, molecular,

or physiological responses in selected tissues that contribute to salt stress adaptations, we still

have a large gap in understanding how a common set of orthologs from a closely related group

of plants selectively modulate salt stress adaptations at a cellular level (Isayenkov and

Maathuis, 2019) and which pathways are synergistically used to achieve salt stress adaptation

facilitating stress resilient growth in extremophytes.

In this study, we used a combined transcriptomic, ionomic, and metabolomic

experimental design to investigate the coordinated cellular responses of *S. parvula*, *E.*

salsugineum, and *A. thaliana* to salt treatments in root and shoot tissues. We examined the

coordination of multiple genetic pathways used by the three models at different salt stress

intensities. We identified preadapted responses mostly at the metabolome level for *E.*

salsugineum while *S. parvula* showed more stress-preparedness at the transcriptome level.

Additionally, both extremophyte showed molecular phenotypes suggesting induction of

complementary cellular processes that used core pathways present in all plants, but with

modifications to those in ways that optimized balance between stress tolerance and growth.

Results

Previous studies using either *S. parvula* or *E. salsugineum* in a comparative study with *A.*

thaliana have successfully used 150 mM NaCl to elicit salt stress responses in the extremophyte

while avoiding induction of immediate tissue necrosis in *A. thaliana* in the short-term

(Kazachkova et al., 2018). We grew *S. parvula, E. salsugineum*, and *A. thaliana* hydroponically

for 28 days, added NaCl to the growth medium (Figure S1A). *Arabidopsis thaliana* did not show

severe stress symptoms until four days of exposure to salt. Therefore, in our comparative

system with three species, we decided to capture the short-term effects of salt at 0, 3, 24, and

72 hr durations with 150 mM NaCl to include timepoints that would precede the onset of stress

phenotypes. Additionally, 250 mM NaCl was used to further deduce salt stress responses

induced in the extremophytes. We used the initial three timepoints to detect immediate

responses of the ionome and the transcriptome to salt and used 0 hr and the latter two

timepoints to detect the subsequent metabolome level changes (Figure S1B).

High tissue tolerance to Na accumulation vs limiting Na accumulation in tissues

All three species accumulated salt as the duration of the treatment or the concentration

of NaCl increased. *Arabidopsis thaliana* accumulated Na in shoots much more than in roots but

did not limit Na accumulation in roots compared to *S. parvula* (Figure 1A). *Arabidopsis thaliana*

shoots showed a 19-fold increase of Na compared to control within 24 hr. Tissue accumulation

of Na was remarkably low in both roots and shoots of *S. parvula* compared to the other two

species. Interestingly, *E. salsugineum* allowed high Na accumulation in both roots and shoots

compared to *S. parvula* and at levels similar to those observed for *A. thaliana* roots under salt

stress (Figure 1A). This suggests that *S. parvula* limited total Na accumulation in tissues while *E.*

salsugineum showed high tissue tolerance to Na accumulation, a trend maintained for 250 mM

NaCl treatments in extremophytes (Figure 1A). High Na levels interfere with K and other

nutrient uptake in plants (Munns and Tester, 2008). We examined whether Na accumulation

triggered nutrient imbalance in any of the test species, by quantifying the abundance of 13

plant nutrients during salt stress treatments. The K level significantly dropped within 3 hr in *A.*

thaliana roots but did not change in the extremophytes regardless of the duration and intensity

of the Na^+ treatments (Figure 1B). In shoots, we did not observe any significant changes in K

levels by salt stress treatments in any of the three species. When we examined the elemental

profiles that include all quantified nutrients, we found that shoots exhibited minimal changes in

all three species in response to salt stress (Figure 1C). Intriguingly, *A. thaliana* roots showed

significant depletion of 6 out of 13 nutrients under salt treatments whereas both

extremophytes showed minimal nutrient disturbances (Figure 1C and Table S1). We then

examined nutrient compositions in the control samples (basal levels) to identify if any nutrient

was preferentially enriched in any of the species (Figure 1D and Table S1). Notably, Ca and Mg,

which are known to aid in selectivity of Non-Selective Cation Channels (NSCCs) for K over Na in

plants (Shabala et al., 2006), and B were found at higher basal levels in *S. parvula* roots

compared to other species. Similarly, Cu and Co were high in *E. salsugineum* whereas Fe, P, K,

and Mo were high in *A. thaliana*.

Primary metabolite pools decreased from high basal levels in *E. salsugineum* but increased

from low basal levels in *S. parvula* upon salt treatments

Salt stress requires metabolic adjustments in plants (Pantha and Dassanayake, 2020).

Therefore, we examined whether the extremophytes showed similar metabolic adjustment

strategies that were distinct from the salt-sensitive *A. thaliana* when treated with salt. We

quantified a total of 716 metabolites using GC-MS for each species (see Methods), of which 182

with known specific structures were referred to as "known metabolites'' while the remaining

were collectively referred to as "unknown metabolites'' in this study (Table S2). The overall

quantified metabolite pool in roots showed a relatively strong correlation in metabolite

abundance between selected pairwise comparisons resulting in two distinct trends (Figure 2A).

First, the highest pairwise correlation was detected between the control groups of *S. parvula*

and *A. thaliana* which decreased with time as the duration of salt treatment increased (Figure

2A, upper panel). Second, *S. parvula* and *E. salsugineum* profiles were moderately correlated at

control conditions, but as both species were treated with salt, the correlation became stronger

indicating that the majority of metabolites in the extremophytes adjusted to similar levels in

the roots in response to salt treatment. This trend was observed at a shorter duration of salt

treatment at 24 hr when the NaCl concentration was higher at 250 mM (Fig. 2A, lower panel).

In contrast to roots, the shoot metabolomes of *S. parvula* and *E. salsugineum* remained

divergent following salt treatment (Figure S2A).

We also found distinct patterns in the direction of salt responses among species. Under

salt treatments, metabolite abundances in both *A. thaliana* and *E. salsugineum* roots largely

decreased while *S. parvula* increased metabolite abundances (Figure 2B). This trend for *S.*

parvula was not upheld in shoots, where all three species predominantly decreased in

metabolite abundances (Figure S2B). We next examined if the dynamically changing

metabolites were categorically associated with sugars and amino acids often known for their

roles as organic osmolytes (Slama et al., 2015). Figure 2C shows percent abundance among the

three species at control levels in roots, for all sugars, amino acids, and their immediate

derivatives (see Methods) quantified in our study that showed significant difference at basal

level abundance. Even before the salt treatment, *E. salsugineum* had accumulated much higher

levels for most of these metabolites than the other species. For instance, the percent

abundance of sucrose in *E. salsugineum* accounted for 98.6 % of the combined sucrose

abundance in the roots of all three species (Figure 2C) and this was 155- and 131-fold higher

than that in *A. thaliana*, and *S. parvula*, respectively (Table S2). Similarly, glucose, raffinose, and

fructose sugars and proline were much higher in *E. salsugineum* compared to the other species

(Figures 2C and Table S2).

We then tested the possibility of one extremophyte maintaining a higher basal

abundance for key metabolites associated with osmoregulation while the other extremophyte

actively induced those under salt stress, eventually converging to a metabolic status distinct

from that of *A. thaliana* under salt stress, as we observed in Figure 2A. We clustered the

profiles of all 146 known metabolites that changed abundance significantly differently at basal

level or in at least one stress condition from roots of all three species (Figure 2D). Two readily

identifiable clusters emerged (boxes outlined in Figure 2D), composed of metabolites that

showed lower abundances in *A. thaliana* and *S. parvula* at the control condition compared to *E.*

salsugineum, but increased in abundance exclusively in *S. parvula* upon prolonged salt

treatment, to the high level constitutively found in *E. salsugineum*. These clusters were

enriched in metabolites known for their role as osmoprotectants or antioxidants (Slama et al.,

2015), including sucrose, fructose, glucose, GABA, proline, and dehydroascorbic acid (full list in

Table S3). These results indicate a basal level metabolic "preparedness" in *E. salsugineum* roots

compared to active induction of many osmoprotectants and antioxidants in *S. parvula* when

responding to salt treatments. On the other hand, *A. thaliana* lacks neither a preadapted- nor

dynamic-strategy found in the two extremophytes.

***A. thaliana* showed stronger transcriptomic responses compared to the extremophytes**

during salt stress

Previous studies have reported that nearly 20 % of the *A. thaliana* transcriptome

responds to salt stress (Gong et al., 2005; Oh et al., 2014). We tested if the transcriptomic salt

responses from *A. thaliana* aligned more with either extremophyte or if the extremophytes

showed a largely overlapping response that were distinguished from *A. thaliana*. The overall

root and shoot transcriptome profiles were grouped into species-tissue clusters (Figures 3A,

S3). This dominant species-level distinctions in transcript profiles were further demonstrated

with pairwise correlations of entire transcriptome profiles where none of the comparisons

showed correlations (Figure S3) unlike the similarities detected in metabolite profiles (Figure 2A

and Figure S2A). Among differently regulated genes (DEGs), we observed more transcripts

induced, than suppressed, by salt treatments in all three species, and the majority of the DEGs

were identified in 1-to-1 ortholog groups (OGs) (see Methods) (Figure 3B, Table S4). *A. thaliana*

showed the largest number of DEGs in response to salt stress in both roots and shoots. The *A.*

thaliana shoot transcriptome contained more than twice the number of DEGs than in the root

transcriptome (Figure 3B), despite fewer changes in the nutrient profile were observed in the

shoot than in the root (Figure 1C). Notably, the shoot transcriptome of *A. thaliana* included

orders of magnitude more salt-responsive DEGs than those of the extremophytes.

We next examined if transcriptomic changes in cellular homeostasis functions during

salt stress observed for *A. thaliana* entailed processes already maintained or induced in the

extremophytes. This search was done at two levels. First, we clustered expression of all 1-to-1

orthologs at their basal level (from control samples), filtered the clusters to show expression

patterns that differed at least in one species compared to the other two, and identified

enriched cellular processes in each cluster (Table 1). Responses associated with abiotic stress

were among the top highly representative processes in each of these functional clusters.

Second, we investigated the changes in transcriptomic profiles during salt treatments, by

identifying functional clusters enriched among all OGs that included at least one DEG in a

species, for roots and shoots separately (Figure 3C and Table S5 and S6). Response to stress

formed the largest functional cluster including the most OGs in both roots and shoots (RC1 and

SC1 in Figure 3C), which were further sub-clustered to highlight various salt responsive

pathways mediated by auxin and ABA regulation and oxidative stress responses (Figure S4).

Interestingly, amino acid metabolism (RC2) and sugar metabolism (RC3) formed the next two

largest root clusters followed by transmembrane transport (RC4) and ion transport (RC5)

(Figure 3C). In shoots, the second and third largest functional clusters (SC2 and SC3 in Figure 3C)

suggested cellular processes involved in plant development and growth/cell cycle, in which the

majority of the *A. thaliana* orthologs were notably salt-repressed. While the *A. thaliana*

orthologs were the most regulated among these OGs, extremophytes responded in similar

functional processes associated with abiotic stresses when treated with 250 mM NaCl (Figure

S5).

We have identified ortholog clusters showing salt-responsive co-expression across time

points within tissues per species. We present the two most dominant co-expression clusters in

roots and shoots (Figure 3D and Table S7). In roots, the largest co-expression cluster showed *A.*

thaliana orthologs substantially salt-induced compared to a much smaller magnitude of

induction seen in the extremophyte orthologs (Figure 3D- left panel). In shoots, the largest co-

expression cluster included *A. thaliana* orthologs suppressed in response to salt while

expression of extremophyte orthologs were stable (Figure 3D- right panel).

The predominant transcriptomic signature in roots supports divergent auxin-dependent root

growth during salt stress

In all three species, response to stress was the most representative function in roots

enriched among ortholog sets showing either different basal-level expression (Table 1) or salt-

responses (Figure 3C) among species. The most dominant sub-functional clusters/processes

was response to hormones in all ortholog sets especially in roots (Table 1, Figure S4A, and

Figure S5 A and C). Hormonal regulation plays an important role in salt stress responses (Yu et

al., 2020). Furthermore, our comparative ortholog expression profiling suggested that

regulation of hormone signal transduction was distinct among the target species (Figures 4A, S6

and Table S8). We found that genes mediating auxin response via auxin/indole acetic acid

(Aux/IAA) repressors (Gallei et al., 2020) and the type 2C protein phosphatases (PP2Cs) that

negatively regulate abscisic acid (ABA) signaling (Hauser et al., 2011) showed contrasting salt-

responsive expression among the three species, while those of other regulatory phases in auxin

and ABA signaling together with other hormone signaling pathways largely conserved (Figures

4A, S6, and Table S8). Notably, none of the *AUX/IAAs* and *PP2Cs* showed any differential

expression in response to salt in *S. parvula* while 15 out of total 24 *AUX/IAAs* and *PP2Cs*

orthologs in the other two species were significantly induced when treated with salt (Figure

4A). When the basal expression was compared, 11 out of 17 *E. salsugineum AUX/IAAs* orthologs

and 5 out of 7 *S. parvula PP2Cs* orthologs showed higher levels compared to their respective

orthologs in the other species (Figure 4A, the bottom panel, and Table S8).

Auxin and ABA suppress root elongation and lateral root initiation during salt stress in *A.*

thaliana (Ding and De Smet, 2013; Ding et al., 2015). Therefore, we first curated genes involved

in primary root development (GO:0080022) and lateral root development (GO:0048527) and

checked for functionally verified phenotypes associated with them that describe root growth in

A. thaliana. Then, we assigned a binary category of promoted or suppressed root growth to the

orthologs based on their basal expression levels in our target species and the genetically

verified functions of *A. thaliana* ortholog (see a full list of references used for the selected

genotype-phenotype associations in Table S9). We identified 35 single-copy orthologs that

showed a distinct expression level in one species compared to the other two species and

assigned their binary category on their effects on root growth, given their presumed effect on

root phenotypes (Figures 4B). We further mapped them to a lateral root development gene

network (modified from Banda et al., 2019; De Rybel et al., 2010 and Figure S7). Among the

total of 35, we found 27 orthologs in *S. parvula* to suggest increased primary root elongation or

suppressed lateral root initiation compared to the other species, while 23 orthologs in *E.*

salsugineum suggested slower primary and lateral root growth (Figure 4B and S7).

Contrastingly, *A. thaliana* showed 25 orthologs that supported its fast primary root growth and

increased lateral root number. Notably, the majority (71%) of the 35 orthologs were annotated

as auxin-responsive genes (Figure 4B).

In line with the observed difference in expression of orthologs associated with root

development (Figure 4B), *S. parvula* and *A. thaliana* seedlings have comparable primary root

lengths at control conditions, longer than that in *E. salsugineum* (Figure 4C and D). Moreover, *S.*

parvula showed uncompromised primary root growth compared to *E. salsugineum* and *A.*

thaliana when treated with salt for an extended time (Figures 4C, D, S8). *S. parvula* and *E.*

salsugineum seedlings also had fewer lateral roots compared to *A. thaliana* at control

conditions (Figure 4E). However, lateral root growth, assessed using total lateral root number

and density during a week-long salt treatment, indicated that *S. parvula* not only sustained

uninterrupted root growth but also induced lateral root growth upon salt treatment, in contrast

to the responses observed for *A. thaliana* and *E. salsugineum* (Figure 4E, F, and S8).

We aimed to further identify orthologs in *S. parvula* that may support its uninterrupted

primary and lateral root initiation during salt stress that distinguishes it from the other two

species. Therefore, we checked co-expression clusters that included differentially expressed

ortholog groups (OGs) in roots and found two clusters where *S. parvula* showed a different

pattern compared to the other two species (Figure 4G). The first cluster showed salt-induced

expression uniquely observed for *S. parvula.* This accounted for 11% (294 ortholog groups) of

differentially expressed OGs (DEOGs) in roots (Table S10). The second cluster with 113 OGs (5%

of all root DEOGs) groups showed induction of genes in response to salt stress in *A. thaliana*

and *E. salsugineum* while the orthologs in *S. parvula* did not alter their expression (Figure 4G).

Interestingly, within these two clusters, nearly 50% of orthologs did not have a specific GO

annotation indicating the level of functional obscurity associated with those genes that

uniquely respond in *S. parvula* to salt stress (Table S10).

Salt-responsive transcriptome-metabolome coordination supports cellular protection in

extremophytes

In response to salt treatments there were a total of 7852 orthologs identified as

differentially expressed genes (DEGs) and a total of 634 metabolites identified as differently

abundant metabolites (DAMs) for the three species in all salt treatments (Figures 2 and 3, Table

S2 and 4). The difference in % change in response to salt between DAMs and DEGs was the

lowest in *A. thaliana*, while in extremophytes metabolic responses were relatively larger

compared to the overall transcriptomic response during salt treatments (Figure 5A and S9A).

This indicates that there is more transcriptome level preparedness to respond to salt in the

extremophytes than in *A. thaliana*.

The second and third largest functional clusters over-represented among DEOGs in roots

were amino acid and sugar metabolism (550 orthologs) in roots (Figure 3C, RC2 and RC3).

Additionally, our overall metabolite response indicated a high correlation of metabolite levels

between the two extremophytes *S. parvula* and *E. salsugineum* during salt treatments, and the

DAMs shared between the extremophytes were enriched in amino acids and sugars (Figure 2).

Collectively, these findings led us to examine whether DEGs associated with amino acid and

sugar metabolism support a salt-induced metabolic response associated with amino acid and

sugars in the extremophytes. In line with the global trend (Figure 5A), we observed fewer DEGs

associated with amino acid and sugar metabolism in both extremophyte roots than in *A.*

thaliana roots, but more amino acid and sugar metabolites were found as DAMs in

extremophytes in roots (Figure 5B).

Given the established role of amino acids and sugars as organic osmoprotectants during

salt stress (Slama et al., 2015), we focused on the most highly responsive DAMs that are also

osmoprotectants and the key genes that are involved in the metabolism of those metabolite

(Figures 5 C and D, and S9A). Proline was one of the most significant DAMs found in roots of

both extremophytes that not only had a higher basal level than in *A. thaliana*, but also its

abundance further increased with salt treatments concordantly with the transcriptional

induction of *pyrroline-5-carboxylate synthetase* (*P5CS1*) and suppression of *Pro-dehydrogenase*

(*ProDH1*) in extremophytes (Figure 5C and S9B)*.* Furthermore, shikimic acid derived amino

acids, tyrosine, and phenylalanine (precursors of phenylpropanoids) showed higher basal levels

or significant induction in roots of both extremophytes compared to *A. thaliana* (Figure 5C).

Interestingly, genes coding for the enzymes directly involved in the conversion of

phosphoenolpyruvate to shikimic acid such as DAHPS1, DAHPS2, DHQS, DHQ, SK1, ESPS, and CS

were not detected as DEGs in any of the three species (Figure S9C). However, *MYB15,* a master

regulator of shikimic acid biosynthesis pathway (Chen et al., 2006), was highly induced in all

three species (Figure 5C). *Chorismate mutases 1* (*CM1*), coding for the enzyme involved in the

first committed step of tyrosine and phenylalanine biosynthesis was salt-induced in *E.*

salsugineum, while it was constitutively expressed at a high level in *S. parvula* (Figure 5C).

E. salsugineum maintained a higher basal abundance than the other two species for

multiple sugars in roots used as organic osmolytes in plants, while these metabolites increased

their abundances in response to salt in *S. parvula* (Figure 5D). The genes involved in the

biosynthesis of these metabolites such as *sucrose synthase 1* (*SUS1*), *cell wall invertase 1*

(*cwIN1*), *galactinol synthase 2* (*GolS2*), and *raffinose synthase* were mostly induced under salt

or constitutively expressed with high basal levels in the extremophytes (Figure 5D). This

concordant alignment of DAMs and DEGs or maintenance of high constitutive abundance of

metabolites with high basal expression of genes highlighted for osmoprotectants in the

extremophytes was similar to coordination observed between known antioxidants and their

associated genes such as the dehydroascorbic acid pathway (Figure S9D).

Coordination between nutrient balance and gene expression associated with ion transport

Two other dominant functional clusters in roots, including 156 OGs, assessed using salt

stress responsive ortholog groups identified ion transport and membrane transport as the main

processes, (Figure 3C, RC4 and RC5). We also observed extremophyte ionomic profiles that

maintained nutrient balance unlike that of *A. thaliana* during salt treatments (Figure 1C).

Hence, we investigated all 148 *A. thaliana* genes encoding transporters based on Araport 11

annotation (Cheng et al., 2017) in the following four categories: (1) aquaporins (*PIPs*, *NIPs*, *TIPs*,

and *SIPs*), (2) cation transporters (*CAXs*, *NHXs*, *KEAs*, *CHXs*, *KUPs*, *HKT1*, and *TRH1*), (3) non-

selective cation channels (*CNGCs* and *GLRs*), and (4) K channels (*SKOR, GORK, AKTs* and *KCOs*).

We found 109 corresponding one-to-one ortholog groups (OGs), among which 64 OGs were

differentially expressed at basal level in the three species (Table S11). We focused on the OGs

encoding transporters and channels that were expressed at significantly different basal levels in

both extremophytes compared to *A. thaliana* (Figure 6A). Genes encoding transporters known

to exclude Na from the cell (*SOS1*), transporters involved in Ca uptake and transport (*CNGC1*,

CNGC12, *CAX1*, and *CAX5*), K channel (*KAT1*), and endosomal K transporter (*KEA5*) that aid in

pH and ion homeostasis (Zhu et al., 2018; Sustr et al., 2019) showed higher basal expression in

the extremophytes than the *A. thaliana* orthologs, while genes involved in K efflux/Na influx to

the cell or water transport such as *TIP1;1*, *PIP2;2*, *SIP1;2*, *SIP2;1*, *KUP5/6/10/11*, and *NHX3*

(Figure 6A). *NHX1/2*, the main transporters involved in Na sequestration in the vacuole, showed

higher basal level expression in *E. salsugineum* compared to the other two species (Table S11).

We then examined how these transporter classes changed in expression in the three

species when treated with salt (Figure 6B and Table S12). Interestingly, the majority of the

genes across the four categories were either induced or constant in expression during salt

treatments in *S. parvula*, whereas *A. thaliana* and *E. salsugineum* had a mixed response (Figure

6B). Notably, *E. salsugineum* mostly repressed the expression of genes encoding non-selective

cation channels (e.g. *GLRs*) in response to all salt treatments (Figure 6B and Table S12). Other

observable trends included the induction of *CHX17* (functions in K uptake) identified as the only

induced transporter in all three species in our selected set, a 50-fold induction of *CHX2*

(involved in Na transport to vacuole) in *A. thaliana* (Table S12), and *GORK* as the only K channel

up-regulated in *E. salsugineum* under salt treatment while maintaining high basal level

expression in *S. parvula*.

We next expanded the focus on the transporter genes to include copy number variation

among species and expression partitioned to paralogs present in each species (Figure 6C). We

searched for transporter gene orthologs with increased copy numbers in the extremophytes:

SpNIP6;1/2 in *S. parvula*, *EsCNGC4;1/2* in *E. salsugineum*, *SpKUP9;1/2* and *EsKUP9;1/2* in both

extremophytes (Figure 6C, top panels, and Table S13). All extremophyte paralogs either showed

differential expression in response to salt or high constitutive expression compared to *A.*

thaliana (Figure 6C, bottom panels).

Finally, we examined the expression profiles of all non-Na/K transporters involved in

nutrient uptake that were included in an ortholog group in Figure 3C functional clusters RC4

and RC5 (Figure 6D and Table S13). Notably, *S. parvula* induced orthologs encoding a large and

diverse set of transporters involved in nutrient uptake upon salt treatments, while *A. thaliana*

and *E. salsugineum* showed salt-induction of a limited group associated with Fe, Zn, and Cu

uptake (e.g. *FER1/2/3*, *MTPB1*, *ZF14*, *ZIP2/11*,) (Figure 6D).

Discussion

The direct flow of Na^+ ions into plant tissue is unavoidable in saline soils. Plants show

adaptations at varying degrees to limit accumulation of Na^+ or mitigate cellular toxicity caused

by Na^+ in tissues when exposed to increasing levels of salt (Flowers and Colmer, 2008; Pantha

and Dassanayake, 2020; Santiago-Rosario et al., 2021). In the tolerance spectrum for salt, *S.*

parvula and *E. salsugineum* are considered halophytes whereas *A. thaliana* is not (Kazachkova

et al., 2018). We show that both extremophytes accumulate less Na at the whole plant level

than *A. thaliana* during salt treatments, with *S. parvula* accumulating the lowest levels (Figure

1A), consistent with previous reports (Volkov et al., 2004; Ghars et al., 2008; Oh et al., 2014).

Figure 7 summarizes the most notable ionomic, metabolomic, transcriptomic, and phenotypic

adjustments that distinguish each model plant from the other two when responding to salt

treatments.

Upholding nutrient balance is a shared outcome in the extremophytes when responding to

increasing external salt levels

When compared to *A. thaliana*, which loses its nutrient balance under salt stress as the

Na content increases (Figure 1A-C)*,* the two extremophytes present two distinct paths for

regulating Na levels in roots coincident to how each plant achieves nutrient balance as a shared

outcome. *S. parvula* regulates Na accumulation to maintain it at levels observed for control

conditions, while preventing the loss or maintaining the uptake of K and Ca during salt stress. In

contrast, *E. salsugineum* does not show such a restriction to Na accumulation in roots but

reaches the same outcome as *S. parvula* for both 150 and 250 mM NaCl treatments at 24 hr

preventing a significant drop in its K and Ca contents.

Regulation of Ca levels coordinated with K is a critical requirement for ion homeostasis

and excess Na in plant tissues when exposed to salt stress challenges this balance (Flowers and

Colmer, 2008; Pantha and Dassanayake, 2020). Ca activates the efflux of Na via SOS2-SOS3

signaling pathway which in turn activates the plasma membrane localized Na-exporter, *SOS1*

(Halfter et al., 2000). Additionally, the plasma membrane localized non-selective cation

channels (NSCCs) are reportedly a main entry point of Na into root cells and are regulated by Ca

(Demidchik and Tester, 2002; Han et al., 2014). Genes encoding for NSCCs are highly

suppressed in *E. salsugineum* consistent with previous findings during salt stress (Volkov and

Amtmann, 2006) (Figure 6B and 7). However, counterintuitively, these *NSCCs* were induced or

unaltered in *S. parvula* during salt treatments (Figure 6B), while *S. parvula* maintains a relatively

smaller Na content than *E. salsugineum* when treated with salt (Figure 1A). The high basal level

of Ca in *S. parvula* roots may alleviate Na-induced cellular toxicity by limiting the selectivity of

NSCCs to Na (Han et al., 2014) (Figure 1C). This suggests that Ca may play a key role to achieve

ion homeostasis in *S. parvula* during salt stress at a much higher level than reported for *A.*

thaliana (Choi et al., 2014).

Halophytes are known for their ability to maintain high K/Na ratios in roots when

exposed to salt even if the species-dependent ratios can have large differences between

halophytes compared to more salt-sensitive species (Flowers and Colmer, 2008; Cuin et al.,

2008; Sun et al., 2009; Cheng et al., 2015). Under salt stress, the negative effect of Na-induced

membrane depolarization at the root epidermis could be reversed by the activation of K

outward rectifying channel, *GORK* by releasing some of the cytosolic K as shown in *A. thaliana*

root hairs (Ivashikina et al., 2001; Shabala and Cuin, 2008). *GORK* expression was induced in *E.*

salsugineum when exposed to 250 mM NaCl stress and maintained at high basal level in *S.*

parvula while it was low at basal level and unaltered in *A. thaliana* roots under salt stress (Table

S4 and 12). Therefore, it may help to prevent the membrane from further depolarization in

extremophytes at high salinities. However, how the extremophytes regulate their K

transporters to allow K uptake and simultaneously prevent excessive leakage of K during salt

stress is unclear. The extremophytes provide an alternative co-regulation of Na and K

transporters different from *A. thaliana*. One such modified pathway in the extremophytes

compared to *A. thaliana* from our comparative analysis point to a possible synergistic activity

between *GORK* and *KAT1* (K channels) synchronized with Na exclusion mediated by *SOS1* to

regulate cytosolic K levels, pH, and ion homeostasis in the two extremophytes when exposed to

salt. Coordinated expression of *GORK* and *KAT1* is a prevalent pathway mostly reported for

guard cells in *A. thaliana* and *KAT1* expression is thought to be highly reduced in roots (Ache et

al., 2000; Philippar et al., 2004). In contrast, *KAT1* is highly expressed even at basal expression

levels in both extremophytes (Figure 6A). Notably, these K channels are regulated by ABA and

auxin and both extremophytes seem to differently regulate these two hormones in conjunction

with overall root growth modulation in response to salt differently compared to that in *A.*

thaliana roots (Figure 4, S7, and 7). The intracellular control of K transport appears to be

uniquely regulated in the extremophytes compared to *A. thaliana.* For example, *KEA5* primarily

expressed in the trans-Golgi network (Zhu et al., 2018; Zhang et al., 2020) is expressed at much

higher levels in both extremophytes than in *A. thaliana* (Figures 6A and 7)*.* Additionally, *KUP9*

which has undergone tandem duplication in both extremophytes, has one paralog in each

extremophyte that show much higher basal expression as well as induction following salt

treatments, in contrast to lower constitutive expression of the *A. thaliana* ortholog (Figure 6).

KUP9 in *A. thaliana* mediates auxin efflux from endoplasmic reticulum (Zhang et al., 2020). It

would be interesting to find out if the *KUP9* duplication in the extremophytes have led to

divergent regulation in auxin homeostasis. Future research exploring this aspect may lead to a

missing link in how root growth is modulated under salt stress in salt adapted species.

High tissue tolerance to Na is illustrated by high Na accumulation in *E. salsugineum*

roots without losing its nutrient balance during salt treatments (Figure 1). This is thought to be

primarily facilitated by vacuolar Na-K/H transporters, *NHX1* and *NHX2* (Bassil et al., 2011). *NHX1*

and *NHX2* are co-induced with *V-ATPase* and this coordinated transcription concurrent to

increasing salt treatments is observed as expected in *E. salsugineum*, but not in the other two

models (Table S11 and Figure 7). *S. parvula* may have reduced the need for vacuolar

sequestration of Na differently from *E. salsugineum*. If it is achieved by regulating Na entry to

the cell via CNGCs or aquaporins as proposed by previous studies conducted on *A. thaliana*

(Demidchik et al., 2002; Byrt et al., 2017), transcript level coordination alone is insufficient to

deduce the orthologous functions in *S. parvula* as both groups are categorically highly induced

under salt treatments instead of being suppressed to reduce Na entry (Figure 6B).

Metabolic preadaptation or a dynamic response equally allow adaptation to salt stress in

extremophytes

Sugars, amino acids, or their associated derivatives are used in all plants for
osmoregulation during salt stress (Slama et al., 2015). The most striking metabolic feature
among the three species in our study is the extraordinary level of sugars and amino acids in *E.*
salsugineum even in control conditions compared to the other two species of which many of
those metabolites can serve as osmoprotectants or antioxidants (Figure 2C and 7). The
metabolite profiles we observed are consistent with earlier studies that examined metabolites
in *E. salsugineum* (Gong et al., 2005; Kazachkova et al., 2013; MacLeod et al., 2015; Lee et al.,
2016; Eshel et al., 2017; Shamustakimova et al., 2017; Yin et al., 2018; Pinheiro et al., 2019).
Previous work had suggested that *E. salsugineum* leaves and seedlings are metabolically
preadapted to salt stress (Gong et al., 2005; Kazachkova et al., 2013; Lee et al., 2016). Our
results reinforce this view and extend these observations to roots (Figure 2C and 7).
Complementary to a dynamic metabolic response, *S. parvula* in comparison to the other two
species exhibits a higher transcriptome level preadaptation to salt stress (concordant with its
fewer DEGs in both roots and shoots; Figures 3B and 5A). Interestingly, during salt treatments,
S. parvula which maintained much lower levels of sugars and amino acids comparable to levels
seen in *A. thaliana* at control conditions, raised its metabolite pools of amino acids and sugars
to the high basal level present in *E. salsugineum* (Figures 2 and 7).

Proline is among the most studied metabolites during salt stress in *E. salsugineum* and
generally in extremophytes due to its role as an osmoprotectant and an antioxidant (Taji et al.,
2004; Kant et al., 2008; Flowers and Colmer, 2008; Bartels and Dinakar, 2013). While the key
proline biosynthesis gene, *P5CS1* was induced in all three species under salt treatments, *Pro-*
dehydrogenase (*ProDH1*), encoding an enzyme that degrades proline, was suppressed only in
the extremophytes, coincident to a significant boost in proline exclusively in the extremophytes
(Figure 5C). The expression of *ProDH1* is known to be suppressed by sucrose (Funck et al.,
2010). Our results may suggest a novel regulatory alteration involving high sucrose levels
concomitant to the induction of *sucrose synthase* (*SUS1*) to maintain high proline levels in the
extremophytes via sucrose mediated suppression of *ProDH1* (Figure 5D). *A. thaliana SUS1* is

known to be regulated by osmotic stress independent of ABA and is also less expressed in roots

compared to other *SUS* members (Déjardin et al., 1999; Bieniawska et al., 2007).

An alternative view to why sugars and amino acids are maintained in *E. salsugineum* at

high levels as a preadapted feature to abiotic stress tolerance is proposed based on those

metabolites serving as a carbon or nitrogen source to facilitate growth when photosynthesis

and nitrogen acquisition may decrease with abiotic stress (Grime and Hunt, 1975; Pinheiro et

al., 2019; Yu et al., 2020). *E. salsugineum* roots grow much slower than both *A. thaliana* and *S.*

parvula primary roots under control conditions (Figure S8) (Orsini et al., 2010). A high

proportional allocation in roots for sugars such as sucrose and raffinose together with amino

acids known for nitrogen storage, especially proline is a common trait associated with slow

growing plants (Poorter, 1989; Atkinson et al., 2012). This idea aligns better with *E. salsugineum*

viewed as a slow growing annual compared to the other two models. Some of these

metabolites that are high at basal levels then have the added benefits for serving as

osmoprotectants or antioxidants when the plants experience high salinity levels.

Accumulation of sugars is reported as a metabolic trait in response to drought in *A.*

thaliana leaves (Sperdouli and Moustakas, 2012). Contradictory to viewing the accumulation of

sugars as a growth promoting adaptation or an adaptation to stress tolerance, it has been

proposed as an outcome of photosynthates passively accumulating in the absence of active

growth under stressed conditions (Martínez-Vilalta et al., 2016; Granda and Camarero, 2017).

Schrenkiella parvula dynamically accumulates these metabolites during salt treatments without

an indication of compromised growth to match the levels maintained in *E. salsugineum* (Figures

4 and 5 and Tran et al., 2021)*. The source of increased sugars in *S. parvula* is expected to result

from starch hydrolysis mediated by ABA upon salt stress (Thalmann et al., 2016). Therefore, at

least in the extremophytes, the shared outcome of high sugars and amino acids seems to be an

active metabolic state that could be considered as an adaptive feature to cope with salt stress

more than a passive outcome of excess photosynthesis under interrupted growth during stress.

Root transcriptional networks contain multiple divergent points between the extremophytes

with convergent outcomes for stress-resilient growth

Schrenkiella parvula and *Eutrema salsugineum* have lower lateral root densities

compared to *A. thaliana* (Figure 4F). Auxin is among the main hormones that regulate root

architecture including lateral root development (Wang et al., 2009). Lateral root initiation is

regulated by binding of auxin to its receptor TIR1, resulting in degradation of auxin signaling

repressors, the Aux/IAAs (Lakehal et al., 2019). The *TIR* orthologs in both extremophytes are

suppressed in control conditions coincident to successive suppression of multiple *Aux/IAAs* and

other downstream transcription factors in control or salt treated conditions, a pattern

collectively suggesting limited lateral root initiation (Banda et al., 2019) compared to *A.*

thaliana (Figure 4B and S7). Both extremophytes either having a slower root growth rate (in *E.*

salsugineum) or higher primary root growth rate but with lower lateral root density (in *S.*

parvula), which may reduce the total surface contact area with salt by reducing lateral root

density. Therefore, the transcriptional network mediated by auxin and other hormones in the

extremophytes suggests multiple regulatory points for root growth modulation that could

result in different root architecture (Figure S7). The overall transcriptional response promoting

root growth is unique to each extremophyte, but supports uninterrupted nutrient acquisition,

water transport, and Na$^+$ sequestration to synchronize both developmental and metabolic

coordination during salt stress differently from *A. thaliana* (Figure 4 and 6).

Molecular phenotypes for transcriptional preadaptation associated with multiple abiotic

stresses have been reported for *S. parvula* and *E. salsugineum* separately to highlight genes

uniformly expressed before and after stress treatments compared to differential expression of

orthologs in a stress sensitive sister species (Gong et al., 2005; Oh et al., 2014; Lee et al., 2016;

Simopoulos et al., 2020; Wang et al., 2021; Pantha et al., 2021). Our work indicated that stress-

associated functional gene clusters that exhibit transcriptional preadaptation in one

extremophyte is in many instances contrasted by a dynamic induced response by the orthologs

from the other extremophyte that ultimately match the expression level found in the

preadapted species (e.g. Figure 4G). This highlights the diversity of independent and divergent

regulation of orthologs even between closely related extremophytes in response to the same

treatments given in the same growth conditions but demonstrating different adaptive

strategies.

Overall, our study highlights the multiple different combinatorial expression modules

that can be examined for stress optimized growth using two extremophytes adapted to grow in

high salinities. The hormone-mediated root growth and especially the selective release of

repression in auxin and ABA signaling pathways that may lead to different *S. parvula* and *E.*

salsugineum root growth strategies absent in *A. thaliana* needs to be further investigated to

identify additional regulatory dependencies using target studies in the extremophyte models.

Similarly, comparisons from multiple extremophytes can serve as training data to identify

compatible regulatory pathways that can coexist but are currently absent in crops. Such

pathways need to be investigated for their functionality to metabolic cost to evaluate whether

constitutive expression (as pre-adaptive traits) or induced expression (as dynamic responses)

offer an optimum strategy depending on a stress condition being constant or intermittent in

certain environments. The scarcity of halophytes despite their recurrent evolution implies a

high metabolic cost and complex regulation required for salt stress adaptation for coordinated

growth from cellular to whole plant level (Flowers and Colmer, 2015). Extremophytes will be a

direct resource when modeling stress resilient growth based on which combination of core

stress-response pathways are critical to deliver growth and survival during environmental stress

(Zandalinas et al., 2021). For sustainable global food security, our crops need to be diversified,

less dependent on fresh water and high nutrient soils while being adapted to varying levels of

marginal soils with different salinities (Bailey-Serres et al., 2019; Pareek et al., 2020). Therefore,

basic research examining genetic regulation underlying stress optimized growth will be a

prerequisite when selecting new crops in light of a climate crisis.

Materials and methods

Plant growth and treatments

Schrenkiella parvula (ecotype Lake Tuz, Turkey; Arabidopsis Biological Resource

Center/ABRC germplasm CS22663)*, Eutrema salsugineum* (ecotype Shandong, China; ABRC

germplasm CS22504)*,* and *Arabidopsis thaliana* (ecotype Col-0) seeds were surface-sterilized

and stratified at 4 °C for 7 days (for *A. thaliana* and *S. parvula*) or 14 days (for *E. salsugineum*).

Stratified seeds were germinated and grown in a hydroponic system as described by Conn et al.,

(2013) for transcriptomic, ionomic, and metabolomic experiments (Figure S1B). The plants were

grown in aerated 1/5x strength Hoagland's solution in a growth cabinet set to 22-23 °C,

photosynthetic photon flux density at 80-120 mM m^{-2}s^{-1}, and a 12 hr light / 12 hr dark cycle.

Fresh Hoagland's solution was replaced every two weeks. Four-week-old plants were randomly

placed in 1/5x strength Hoagland's solution with and without NaCl and incubated for indicated

duration in each experiment.

Root growth analysis

Five-day old seedlings were transferred to 1/4x MS plates supplied with 150 mM NaCl.

Root growth was recorded every two days for one week. Control and treated plates were

scanned and analyzed using ImageJ (Schneider et al., 2012) to quantify primary root length and

number of lateral roots. Three biological replicates were used with 7 seedlings per replicate

from each species.

Elemental analysis

Elemental quantification was conducted for Na, K, Ca, P, S, Mg, Fe, B, Zn, Mn, Mo, Cu,

Ni, and Co using Inductively Coupled Plasma - Mass Spectrometry (ICP-MS, Elan 6000 DRC-

PerkinElmer SCIEX) at the US Department of Agricultural Research Service at Donald Danforth

Plant Science Center. We used dried root and shoot samples from 3-4 biological replicates from

each control and treated condition harvested at 3, and 24 hr (Figure S1) processed as described

in Baxter et al., (2014). Changes in element contents were calculated as the log$_2$ fold change of

the element level in treated samples over that in control samples and visualized with the

pheatmap package in R. Significant differences between treatments within species were

determined by one-way ANOVA followed by Tukey post-hoc test using agricolae package in R

with an adjusted p-value cutoff of 0.05. Basal level of each element per species was calculated

as a % contribution from each species that added to 100% for each element, using the following

formula: Xi/ (Ati+Spi+Esi)*100 where Xi represents the abundance of element i in *A. thaliana*, *S.*

parvula and *E. salsugineum*, respectively. Only elements that showed significant differences in

abundance among the three species were visualized on ternary diagrams using ggtern R.

Metabolite analysis

Untargeted high throughput metabolite profiling was conducted using gas

chromatography-mass spectrometry (GC-MS) service at the West Coast Metabolomics Center,

University of California Davis. Root and shoot samples in 4 biological replicates were harvested

at 24, and 72 hr (Figure S1B), flash frozen in liquid N_2, and processed as described in Fiehn

(2017) and quantified as described in Pantha et al., (2021).

Pearson correlation coefficients per each test condition were calculated between

species using normalized metabolite abundances. Significant differences in metabolite

abundances across treatments within species (differently abundant metabolites, DAMs) was

determined by one-way ANOVA followed by Tukey post-hoc test using an agricolae package in R

with an adjusted *p*-value cutoff of 0.05. Structurally annotated (known) metabolites were

further categorized into functional groups according to the refmet database

(https://www.metabolomicsworkbench.org/databases/refmet/index.php) and Kyoto

Encyclopedia of Genes and Genomes (KEGG). Metabolites were clustered into amino acids,

sugars, nucleic acids, and other organic acids. Derivatives or precursors of those were included

in the same metabolite category if those individual metabolites were found to be within three

steps of the main metabolite category identified in a KEGG pathway (Table S2). Basal level of

each metabolite per species was calculated as a % contribution from each species that added to

100% for each metabolite similar to the elemental basal level calculation described earlier.

Metabolites that showed significant differences in abundance at basal level or under salt

treatments at least in one species were used for K-mean clustering (k= 9) (Mannor et al., 2011).

The metabolite profiles across samples were done using hierarchical clustering function in

pheatmap package.

Orthologous group identification

Ortholog groups were identified using *S. parvula* gene models version 2.2

(https://phytozome-next.jgi.doe.gov/); *A. thaliana* gene models version 10 (TAIR10)

(https://www.arabidopsis.org/download/), and *E. salsugineum* gene models (Wu et al., 2012)

which were updated based on reference-guided transcripte assembly using hisat2 (version

2.0.1)-stringtie (version 1.2.1) pipeline (Pertea et al., 2015, 2016) and RNA-seq reads from this

study. The updated gene models for *Eutrema salsugineum* and *Schrenkiella parvula* are

available at www.lsugenomics.org. The ortholog gene pairs between species were identified by

reciprocal blastp with default parameters and an e-value cutoff of 1E-5. The blastp results were

filtered using a custom python script to select the best High-scoring Segment Pair (HSPs) to

ensure only unique query-subject pairs were retained. Pairs with an alignment coverage smaller

than 50% of the query or the subject were removed. The ortholog-pair list included only non-

redundant pairs with the highest bit score optimized for highest % coverage, and % identity.

Ortholog pairs were additionally filtered to exclude any orthologs that had a more than ± 30 %

length difference in the coding sequence between the two sequences. A total of 16,591 one-to-

one ortholog groups were identified among the three species and used for all downstream

analyses (Table S3).

Gene expression profiling and analysis

Total RNA was extracted from control and treated samples at 3 and 24 hr with three

biological replicates (Figure S1) using Qiagen RNeasy Plant Mini kit, with an on-column DNase

treatment. mRNA enriched samples were converted to libraries using True-Seq stranded

RNAseq Sample Prep kit (Illumina, San Diego, CA, USA),multiplexed, and sequenced on a

HiSeq4000 (Illumina) platform at the Roy K. Carver Biotechnology Center, University of Illinois

at Urbana-Champaign. A minimum of >15 million 50-nucleotide single-end reads per sample

were sequenced for a total of 138 RNAseq samples.

After quality checks, the reads were mapped to reference transcript model sequences of

gene models in each species (described earlier) using Bowtie (Langmead and Slazberg, 2013)

with -m 1 --best -n 1 -l 50. A custom python script was used to count uniquely mapped reads for

each gene model. Differentially expressed genes (DEGs) across treatments within each species

were identified using DESeq2 (Love et al., 2014). Genes with an adjusted p-value ≤ 0.01 were

further filtered using the following criteria: 1) $|\log_2$ fold change$| \geq 1$ or 2) $|\log_2$ fold change$| \geq$

0.5 if normalized mean expression across samples ≥ 100. Genes that passed these filters were

considered as DEGs. Reads per kilobase per million reads (RPKM) were calculated for each gene

from the raw read counts for all samples. These RPKM values were $\log_2$-transformed and

median-normalized when used in PCA-UMAP (McInnes et al., 2018) and for co-expression

cluster determinations. Co-expression clusters were done using fuzzy K-mean clustering with a

membership cutoff ≥ 0.4 and log2 fold change clustering. $\log_2$ fold changes were computed for

all ortholog groups (OGs) from three species, used to calculate Pearson correlation coefficients

when comparing transcriptome profiles between species/samples, and were visualized with

ggplot2 in R. Ortholog groups where at least one of the genes in the group was identified as a

DEG in any species under any stress condition was considered to be a differentially expressed

ortholog group (DEOG).

To compare the basal expression levels of orthologs among species, we generated a raw

count matrix based on reads uniquely mapped to coding sequences (CDS) for the 16,591

orthologs from control samples. Orthologs that are differently expressed between any two of

the three species were identified using DESeq2 at an adjusted p-value ≤ 0.001 in a pairwise

manner. We next assigned the expression of each gene within any given OG as High (H),

Medium (M), and Low (L) based on their relative expression level resulting in 12 clusters. The

six largest clusters in both roots and shoots highlighted the expression difference in one species

compared to the other two species. OGs were assigned to these clusters represented by HLH,

LHL, HHL, LLH, LHH, and HLL (expression profiles). Each expression profile was then subjected to

functional gene enrichment analysis.

BiNGO (Maere et al., 2005) was used to identify Gene Ontology (GO) terms enriched in

selected DEGs and DEOGs from shoots and roots. To reduce the redundancy between enriched

GO terms, we further grouped these GO terms into GO clusters using GOMCL (Wang et al.,

2020) with settings for -Ct 0.5 -I 1.5 -Sig 0.05 -hm -nw -hgt -d -gosize 3500 -gotype BP. Sub-

clustering of selected clusters was performed using GOMCL-sub with the same parameters.

OrthNets for selected genes were generated using the CLfinder-OrthNet pipeline with the

reciprocal blastp results as input using default OrthNet settings (Oh and Dassanayake, 2019),

and visualized in Cytoscape.

Curated gene sets for primary metabolism, hormone signaling, root development, and transporter functions

DEOGs and OGs that showed significant differences at basal level/control condition or under treatments were mapped to the KEGG pathways associated with primary metabolism, hormone regulation, or root development. Log$_2$ fold changes were computed for all selected OGs and visualized with pheatmap in R. Genes involved in root development were identified from Gene Ontology annotations, GO:0048527 for lateral root development and GO:0080022 for primary root development. Genes coding for transport functions associated with K/Na transport were mined from Araport 11. To assess the correlation between changes at the transcriptomic level and the metabolic level, we calculated the percentage of DEGs and differently abundant metabolites (DAMs) in each species at early (3 hr for transcriptome and 24 hr for metabolome) and late (24 hr for transcriptome and 72 hr for metabolome) response to salt. Percent DEGs was calculated by dividing the number of DEGs by the total number of expressed genes (RPKM ≥ 1) in each species. Similarly, the percent DAMs was calculated by dividing the number of DAMs by the total number of quantified metabolites within a species. We extracted DEGs and DAMs related to amino acid and sugar metabolism based on GO (GO:0006520 for amino acid metabolism and GO:0005975 for carbohydrate metabolism). The count of DEGs and DAMs were separately normalized to the total number of genes and metabolites in each of the two categories.

Data availability

All Illumina sequence data are deposited at National Center for Biotechnology Information BioProject PRJNA63667. Mapped RNAseq data can be browsed using genome browsers created for *Schrenkiella parvula* and *Eutrema salsugineum* at www.lsugenomics.org.

Acknowledgement

This work was supported by the US National Science Foundation IOS-EDGE 1923589/2019610, NSF-MCB-1616827, US Department of Energy BER-DE-SC0020358, Next-Generation BioGreen21 Program of Republic of Korea (PJ01317301) awards. Graduate students K.T., G.W. were supported by an Economic Development Assistantship from LSU. We thank the

LSU High Performance Computing facility for providing computational resources;

undergraduate students Jordan Vivien, Christine Tran, Ashley Doan, and Jason Garcia at LSU for

assisting with plant growth and phenotyping; Drs. Alvaro Hernandez and John Cheeseman at

University of Illinois for providing assistance with Illumina sequencing.

Author contribution: K.T. prepared plant samples and conducted data analyses; K.T., G.W., D-

H.O., J.L., A.S., and M.D. contributed to data interpretation. K.T. and M.D. wrote the manuscript

with input from all co-authors who revised and approved the final manuscript. M.D.

conceptualized and supervised the overall project.

Main figure captions

Figure 1: Sodium accumulation and its effect on nutrient balance in the extremophytes

compared to *A. thaliana*. [A] Na^+ and [B] K^+ content in roots and shoots. Data are mean ± SD (n ≥

4, at least 4 plants per replicate). Asterisks indicate significant differences ($p ≤ 0.05$) between

the treated samples and their respective control samples, determined by Student's t-test. Open

circles indicate the biological replicates. [C] Macro- and micro-nutrient fold changes between

treated and its respective control sample. [D] Percent abundance of nutrients that were

significantly different at basal levels among the three species. The three axes of the ternary

plots are marked with *A. thaliana, S. parvula*, and *E. salsugineum*. The gridlines in species

designated colors point to the relevant % abundance of the element in each axis. Significant

difference in elemental abundance was determined by one-way ANOVA with post-hoc Tukey's

test at p ≤ 0.05.

Figure 2: Overall metabolic preparedness and adjustments in the roots of the extremophytes

compared to *A. thaliana*. [A] Pairwise correlation of 716 quantified metabolites(182 known, and

534 unknown) across all conditions and species. Correlation calculated using Pearson

correlation coefficients. [B] Total number of metabolites that significantly changed in

abundance in each species compared to its respective control sample. Known metabolite

numbers given in parenthesis. [C] Basal level metabolite abundances in all three species for

sugars, amino acids, and their derivatives. [D] Hierarchical clustering of known metabolites

shown in [C]. Blue box indicates clusters of interest. Treatment concentrations are 150 and 250

mM NaCl; Treatment durations are 24 and 72 hr. Significant differences were determined by

one-way ANOVA with post-hoc Tukey's test at $p \leq 0.05$ (n = 4, at least 4 plants per replicate).

Figure 3: Transcriptomic overview *S. parvula, E. salsugineum,* and *A. thaliana* in response to

salt. [A] Overall transcriptome clustering using one-to-one ortholog groups (OGs) in shoots and

roots from all replicates in all conditions. [B] Number of differently expressed genes (DEGs) in

response to salt treatments. [C] Functionally enriched processes clustered among OGs that

included a DEG from at least one species in roots and shoots. Functional annotations were

based on GO annotations assigned to OGs. The bar graphs assigned to each cluster represent

percent allocation of induced and suppressed DEGs with the number of genes that did not

significantly change in each species given in the white space between induced (red) and

suppressed (blue) bars. [D] Dominant gene expression clusters of orthologs across time points

and species from roots and shoots that included an *A. thaliana* ortholog that was either

induced or suppressed in response to salt. Differentially expressed genes identified at *p*-adj ≤

0.01. N= 3 (at least 4 plants per replicate).

Figure 4: Root growth responses in line with gene expression mediated by auxin and ABA in

Arabidopsis thaliana, Schrenkiella parvula, and *Eutrema salsugineum* when treated with salt.

[A] Expression responses of auxin/indole acetic acid (Aux/IAA) repressors and type 2C protein

phosphatases (PP2Cs) that regulate auxin and ABA signaling. [B] Curated genes from lateral root

development (GO:0048527) and primary root development (GO:0080022) used to infer root

growth phenotypes based on functional genetic studies in *A. thaliana*. Orthologs in the three

species are indicated as high (yellow), low (blue), and indistinguishable levels between species

(no-color) assigned to relative basal expression levels and their effect inferred with arrows (up –

promote growth and down – suppress growth). Genes associated with Auxin, ABA, and other

hormones were labeled in pink, gold, and black respectively. [C] Root growth assessment of 12-

day-old seedlings in response to 150 mM NaCl treated for 7 days. Quantification of [D] primary

root growth, [E] number of lateral roots, and [F] lateral root density. Data given as mean ± SD (n

= 3, at least 7 plants per replicate). Open circles indicate the measurements from each plant.

Asterisks indicate significant differences ($p \leq 0.05$) between treated samples and their

respective control samples, determined by one-way ANOVA with post-hoc Tukey's test. [G]

Gene co-expression clusters of differentially expressed orthologs that showed similar

expression pattern in *A. thaliana* and *E. salsugineum* roots compared to a different pattern

observed for *S. parvula* under salt stress.

Figure 5: Coordination between differently expressed genes and differently abundant

metabolites during responses to salt in *Arabidopsis thaliana, Schrenkiella parvula,* and *Eutrema*

salsugineum. [A] The difference in percent change in total differentially expressed genes (DEGs)

and total differently abundant metabolites (DAMs) in response to salt treatments. [B] % DEGs

and % DAMs involved in the metabolism of amino acids, sugars, and their immediate

derivatives (GO:0006520 and GO:0005975). [C and D] Selected pathways in amino acids and

sugar metabolism. Line graphs represent normalized Log2 relative metabolite abundance.

Boxplots represent normalized expression values. Center line shows median; box indicates

interquartile range (IQR); notch is for 1.58 × IQR/sqrt(n); whiskers show 1.5 × IQR. Asterisks

indicate DEGs and DAMs assigned using 3-4 biological replicates. Early (E) and late (L) responses

for transcripts refer to 3 and 24 hr. E and L responses for metabolites refer to 24 and 72 hr.

Metabolites are shown in the backbone of the pathway while genes encoding for key

enzymes/transcription factors are placed beside arrows. Black labels in pathway maps indicate

quantified metabolites and genes while grey labels indicate metabolites not quantified.

Figure 6: Basal and differential expression associated with salt and nutrient transport in

Arabidopsis thaliana, Schrenkiella parvula, and *Eutrema salsugineum*. [A] Basal level expression

given as fold changes between species. The dashed lines indicate 2-fold differences to highlight

genes either highly suppressed or highly expressed in the extremophytes in roots at control

conditions. [B] Aggregated expression profiles of transporters/channels associated with Na^+/K^+

transport activity during salt treatments. [C] Selected cation transporters that showed higher

gene copy number in extremophytes in ortholog groups and their expression during salt

treatments. Nodes of each ortholog network is assigned with species colors connected by edges

based on the relationship properties of the homologs. Edges show relationships for co-linear

reciprocal orthologs (cl-rc), co-linear unidirectional orthologs (cl-uni), transposed -unidirectional

duplicates (tr-uni), or tandem duplicated-unidirectional paralogs (td-uni). Asterisks indicate

differentially expressed genes identified at *p*-adj ≤ 0.05 and ** *p*-adj ≤ 0.01. N= 3 (at least 4

plants per replicate). [D] Genes associated with nutrient transport identified in Figure 3C.

Figure 7: Summary of ionomic, metabolomic, transcriptomic, and phenotypic adjustments and

preadaptation trends to salt stress in *Arabidopsis thaliana*, *Schrenkiella parvula* and *Eutrema*

salsugineum roots. Top 9 metabolites that increased abundance in *S. parvula* were Fru,

fructose; Asn, asparagine, Raf, raffinose, DHA, dehydroascorbic acid; Pro, proline; Shkm,

shikimic acid; Suc, sucrose, GABA, gamma-aminobutyric acid; Gln, glutamine.

Orthologs in the three species are indicated as high (yellow), low (blue) assigned to relative

basal expression levels. Blue and red indicate the induction and suppression gene expression,

respectively. KEA5, K^+ efflux antiporter 5; KUP9, K^+ uptake permease 9; KAT1, K^+ channel in

Arabidopsis thaliana 1; GLRs, glutamate-like receptors; NHX1/2, Na^+/H^+ exchanger1/2; V-

ATPases, vacuoles-ATPases; SOS1, salt overly sensitive 1; GORK, gated outwardly-rectifying K^+

channel.

Main table caption

Table 1: Table 1: Enriched cellular processes of ortholog groups at basal expression level

clustered in shoots and roots of *S. parvula* (purple), *E. salsugineum*, (red) and *A. thaliana*

(green).

Supplement figure and table captions

Figure S1: Figure S1: Effect of salt stress on phenotype of Schrenkiella parvula (Sp), Eutrema

salsugineum (Es) and Arabidopsis thaliana (At). [A] Four-week-old hydroponic grown plants

were treated with indicated salt concentrations. [B] Experimental design and sample scheme.

There were at least 4 replicates used for ionomic and metabolomic profiling, and 3 replicates

for transcriptomic profiling with at least 4 plants per replicate.

Figure S2: Overall shoot metabolite responses in the extremophytes compared to *A. thaliana*.

[A] Pairwise correlation of 725 quantified metabolites across all conditions and species.

Correlation calculated using Pearson correlation coefficients. [B] Total number of metabolites

that significantly changed in abundance in each species compared to its respective control

sample. Known metabolite numbers given in parenthesis. [C] Basal level metabolite abundances

in all three species for sugars, amino acids, and their derivatives. Significant differences were

determined by one-way ANOVA with post-hoc Tukey's test at $p \leq 0.05$ (n = 4, at least 4 plants

per replicate).

Figure S3: Transcriptomic profiles of *S. parvula*, *E. salsugineum,* and *A. thaliana* were diverged

in response at basal level and under salt stress. [A] roots, [B] shoots. Correlation was calculated

using log2 fold change of treated over control samples from differentially expressed ortholog

pairs which included DEGs from at least one condition in one species using Pearson correlation

coefficients. *S. parvula* (purple), *E. salsugineum* (red) and *A. thaliana* (green)*.

Figure S4: Sub-clustering of dominant cluster - Response to stresses. Overlap and similarity

between sub-clusters determined using GO-MCL-sub and cumulative number of ortholog

groups (OGs) in [A and B] roots and [C and D] shoots sub-clusters.

Figure S5: Species dependent responses to salt treatments in *Schrenkiella parvula* and *Eutrema*

salsugineum. Differently expressed genes (DEGs) in *S. parvula* [A] root and [B] shoot and *E.*

salsugineum [C] root and [D] shoot in response to 150 and 250 mM NaCl stress at 3 and 24 hr of

treatment. Yellow highlighted data points in the Upset plots for each panel show DEGs that

were uniquely expressed at 250 mM salt treatments. Functionally enriched processes for these

selected DEGs are shown in the horizontal bar graphs given as insets outlined in yellow for each

panel. Differentially expressed genes identified at p-adj ≤ 0.01. N = 3 (at least 4 plants per

replicate).

Figure S6: Hormonal signaling pathways in response to salt treatment in *Arabidopsis thaliana,*

Schrenkiella parvula, and *Eutrema salsugineum*. Expression responses of genes associated with

hormonal signaling pathways obtained from KEGGs. Note that only genes that were

significantly different at either basal level condition or under salt treatments were shown.

Orthologs in the three species are indicated as high (yellow), low (blue), and indistinguishable

levels between species (no-color) assigned to relative basal expression levels.

Figure S7: Lateral root development gene network adopted from Banda et al., 2019 and Rybel

et al., 2010. The expression levels of orthologs were annotated as high (H), medium (M), or low

(L) relative to each other at basal levels in each ortholog group. Specific paths in the network in

which *S. parvula* and *E. salsugineum* consistently regulated to inhibit lateral root formation

were highlighted in ribbons across the network in species-specific colors.

Figure S8: Root growth of *E. salsugineum*, *S. parvula* and *A. thaliana* before and after salt

treatments. *S. parvula* (Sp) promoted primary root growth and reduced lateral root formation

while *E. salsugineum* (Es) showed slower primary root growth with reduced lateral root number

compared to *A. thaliana* (At) roots that showed severe root growth inhibition under salt stress.

Five-day-old seedlings were treated for 11 days on plates supplemented with 150 mM NaCl.

Figure S9: Metabolite abundance and expression of associated genes encoding the enzymes in

shikimic acid and ascorbic acid pathways. [A] percent change in total differentially expressed

genes (DEGs) and total differently abundant metabolites (DAMs) in response to salt treatments.

[B] pathway involved in proline metabolism. [C] pathway involved in the conversion from

phosphoenolpyruvate to chorismate. [D] pathway involved in ascorbic acid metabolism. Line

graphs represent normalized Log2 relative metabolite abundance. Boxplots represent

normalized expression values. Center line shows median; box indicates interquartile range

(IQR); notch is for $1.58 \times IQR/sqrt(n)$; whiskers show $1.5 \times IQR$. Asterisks indicate DEGs

(Differently expressed genes) and DAMs (Differently abundant metabolites). Early (E) and late

(L) responses for transcripts refer to 3 and 24 hr. E and L responses for metabolites refer to 24

and 72 hr. Metabolites are shown in the backbone of the pathway while genes encoding for key

enzymes/transcription factors are placed beside arrows. Black labels in pathway maps indicate

quantified metabolites and genes while grey labels indicate metabolites not shown.

References

Ache, P., Becker, D., Ivashikina, N., Dietrich, P., Roelfsema, M.R.G., and Hedrich, R. (2000).

GORK, a delayed outward rectifier expressed in guard cells of Arabidopsis thaliana , is a K +

-selective, K + -sensing ion channel. FEBS Lett. **486**: 93–98.

Atkinson, R.R.L., Burrell, M.M., Osborne, C.P., Rose, K.E., and Rees, M. (2012). A non-targeted

metabolomics approach to quantifying differences in root storage between fast- and slow-

growing plants. New Phytol. **196**: 200–211.

Bailey-Serres, J., Parker, J.E., Ainsworth, E.A., Oldroyd, G.E.D., and Schroeder, J.I. (2019).

Genetic strategies for improving crop yields. Nature **575**: 109–118.

Banda, J., Bellande, K., von Wangenheim, D., Goh, T., Guyomarc'h, S., Laplaze, L., and

Bennett, M.J. (2019). Lateral Root Formation in Arabidopsis: A Well-Ordered LRexit.

Trends Plant Sci. **24**: 826–839.

Bartels, D. and Dinakar, C. (2013). Balancing salinity stress responses in halophytes and non-

halophytes: a comparison between Thellungiella and Arabidopsis thaliana. Funct. Plant

Biol. **40**: 819.

Bassil, E., Tajima, H., Liang, Y.-C.Y.-C., Ohto, M.M. -a., Ushijima, K., Nakano, R., Esumi, T.,

Coku, A., Belmonte, M., and Blumwald, E. (2011). The Arabidopsis Na+/H+ Antiporters

NHX1 and NHX2 Control Vacuolar pH and K+ Homeostasis to Regulate Growth, Flower

Development, and Reproduction. Plant Cell **23**: 3482–3497.

Bennett, T.H., Flowers, T.J., and Bromham, L. (2013). Repeated evolution of salt-tolerance in

grasses. Biol. Lett. **9**: 20130029.

Bieniawska, Z., Paul Barratt, D.H., Garlick, A.P., Thole, V., Kruger, N.J., Martin, C., Zrenner, R.,

and Smith, A.M. (2007). Analysis of the sucrose synthase gene family in Arabidopsis. Plant

J. **49**: 810–828.

Byrt, C.S., Zhao, M., Kourghi, M., Bose, J., Henderson, S.W., Qiu, J., Gilliham, M., Schultz, C.,

Schwarz, M., Ramesh, S.A., Yool, A., and Tyerman, S. (2017). Non-selective cation channel

activity of aquaporin AtPIP2;1 regulated by Ca 2+ and pH. Plant. Cell Environ. **40**: 802–815.

Chen, Y., Zhang, X., Wu, W., Chen, Z., Gu, H., and Qu, L.-J. (2006). Overexpression of the

Wounding-Responsive Gene AtMYB15 Activates the Shikimate Pathway in Arabidopsis. J.

Integr. Plant Biol. **48**: 1084–1095.

Cheng, C.-Y.Y., Krishnakumar, V., Chan, A.P., Thibaud-Nissen, F., Schobel, S., and Town, C.D.

(2017). Araport11: a complete reannotation of the Arabidopsis thaliana reference genome.

Plant J. **89**: 789–804.

Cheng, D., Wu, G., and Zheng, Y. (2015). Positive correlation between potassium uptake and

salt tolerance in wheat. Photosynthetica **53**: 447–454.

Choi, W.-G., Toyota, M., Kim, S.-H., Hilleary, R., and Gilroy, S. (2014). Salt stress-induced Ca2+

waves are associated with rapid, long-distance root-to-shoot signaling in plants. Proc. Natl.

Acad. Sci. **111**: 6497–6502.

Conn, S.J., Hocking, B., Dayod, M., Xu, B., Athman, A., Henderson, S., Aukett, L., Conn, V.,

Shearer, M.K., Fuentes, S., Tyerman, S.D., and Gilliham, M. (2013). Protocol: Optimising

hydroponic growth systems for nutritional and physiological analysis of Arabidopsis

thaliana and other plants. Plant Methods **9**: 1–11.

Cuin, T.A., Betts, S.A., Chalmandrier, R., and Shabala, S. (2008). A root's ability to retain K+

correlates with salt tolerance in wheat. J. Exp. Bot. **59**: 2697–2706.

Dassanayake, M., Oh, D.-H., Haas, J.S., Hernandez, A., Hong, H., Ali, S., Yun, D.-J., Bressan,

R.A., Zhu, J.-K., Bohnert, H.J., and Cheeseman, J.M. (2011). The genome of the

extremophile crucifer Thellungiella parvula. Nat. Genet. **43**: 913–918.

Déjardin, A., Sokolov, L.N., and Kleczkowski, L.A. (1999). Sugar/osmoticum levels modulate

differential abscisic acid-independent expression of two stress-responsive sucrose

synthase genes in Arabidopsis. Biochem. J. **344**: 503–509.

Demidchik, V., Davenport, R.J., and Tester, M. (2002). Nonselective cation channels in plants.

Annu. Rev. Plant Biol. **53**: 67–107.

Demidchik, V. and Tester, M. (2002). Sodium Fluxes through Nonselective Cation Channels in

the Plasma Membrane of Protoplasts from Arabidopsis Roots. Plant Physiol. **128**: 379–387.

Ding, Z. and De Smet, I. (2013). Localised ABA signalling mediates root growth plasticity. Trends

Plant Sci. **18**: 533–535.

Ding, Z.J., Yan, J.Y., Li, C.X., Li, G.X., Wu, Y.R., and Zheng, S.J. (2015). Transcription factor

WRKY46 modulates the development of Arabidopsis lateral roots in osmotic/salt stress

conditions via regulation of ABA signaling and auxin homeostasis. Plant J. **84**: 56–69.

Eshel, G., Shaked, R., Kazachkova, Y., Khan, A., Eppel, A., Cisneros, A., Acuna, T., Gutterman,

Y., Tel-Zur, N., Rachmilevitch, S., Fait, A., and Barak, S. (2017). Anastatica hierochuntica,

an Arabidopsis Desert Relative, Is Tolerant to Multiple Abiotic Stresses and Exhibits

Species-Specific and Common Stress Tolerance Strategies with Its Halophytic Relative,

Eutrema (Thellungiella) salsugineum. Front. Plant Sci. **7**: 1–18.

Fiehn, O. (2017). Metabolomics by Gas Chromatography-Mass Spectrometry: the combination

of targeted and untargeted profiling.

Flowers, T.J. and Colmer, T.D. (2015). Plant salt tolerance: Adaptations in halophytes. Ann. Bot.

115: 327–331.

Flowers, T.J. and Colmer, T.D. (2008). Salinity tolerance in halophytes. New Phytol. **179**: 945–

963.

Flowers, T.J., Galal, H.K., and Bromham, L. (2010). Evolution of halophytes: multiple origins of

salt tolerance in land plants. Funct. Plant Biol. **37**: 604.

Flowers, T.J., Munns, R., and Colmer, T.D. (2015). Sodium chloride toxicity and the cellular

basis of salt tolerance in halophytes. Ann. Bot. **115**: 419–431.

Funck, D., Eckard, S., and Müller, G. (2010). Non-redundant functions of two proline

dehydrogenase isoforms in Arabidopsis. BMC Plant Biol. **10**: 70.

Gallei, M., Luschnig, C., and Friml, J. (2020). Auxin signalling in growth: Schrödinger's cat out of

the bag. Curr. Opin. Plant Biol. **53**: 43–49.

Ghars, M.A., Parre, E., Debez, A., Bordenave, M., Richard, L., Leport, L., Bouchereau, A.,

Savouré, A., and Abdelly, C. (2008). Comparative salt tolerance analysis between

Arabidopsis thaliana and Thellungiella halophila, with special emphasis on K+/Na+

selectivity and proline accumulation. J. Plant Physiol. **165**: 588–599.

Gong, Q., Li, P., Ma, S., Indu Rupassara, S., and Bohnert, H.J. (2005). Salinity stress adaptation

competence in the extremophile Thellungiella halophila in comparison with its relative

Arabidopsis thaliana. Plant J. **44**: 826–839.

Granda, E. and Camarero, J.J. (2017). Drought reduces growth and stimulates sugar

accumulation: new evidence of environmentally driven non-structural carbohydrate use.

Tree Physiol. **37**: 997–1000.

Grime, J. and Hunt, R. (1975). Relative Growth-Rate : Its Range and Adaptive Significance in a

Local Flora. J. Ecol. **63**: 393–422.

Hajiboland, R., Bahrami-Rad, S., Akhani, H., and Poschenrieder, C. (2018). Salt tolerance

mechanisms in three Irano-Turanian Brassicaceae halophytes relatives of Arabidopsis

thaliana. J. Plant Res. **131**: 1029–1046.

Halfter, U., Ishitani, M., and Zhu, J.-K. (2000). The Arabidopsis SOS2 protein kinase physically

interacts with and is activated by the calcium-binding protein SOS3. Proc. Natl. Acad. Sci.

97: 3735–3740.

Han, S., Wang, C., Wang, W., and Jiang, J. (2014). Mitogen-activated protein kinase 6 controls

root growth in Arabidopsis by modulating Ca2+-based Na+ flux in root cell under salt

stress. J. Plant Physiol. **171**: 26–34.

Hauser, F., Waadt, R., and Schroeder, J.I. (2011). Evolution of Abscisic Acid Synthesis and

Signaling Mechanisms. Curr. Biol. **21**: R346–R355.

Inan, G. et al. (2004). Salt Cress. A Halophyte and Cryophyte Arabidopsis Relative Model System

and Its Applicability to Molecular Genetic Analyses of Growth and Development of

Extremophiles. Plant Physiol. **135**: 1718–1737.

Isayenkov, S. V. and Maathuis, F.J.M. (2019). Plant Salinity Stress: Many Unanswered

Questions Remain. Front. Plant Sci. **10**: 80.

Ivashikina, N., Becker, D., Ache, P., Meyerhoff, O., Felle, H.H., and Hedrich, R. (2001). K+

channel profile and electrical properties of Arabidopsis root hairs. FEBS Lett. **508**: 463–469.

Kant, S., Bi, Y.-M., Weretilnyk, E., Barak, S., and Rothstein, S.J. (2008). The Arabidopsis

Halophytic Relative Thellungiella halophila Tolerates Nitrogen-Limiting Conditions by

Maintaining Growth, Nitrogen Uptake, and Assimilation. Plant Physiol. **147**: 1168–1180.

Kazachkova, Y., Batushansky, A., Cisneros, A., Tel-Zur, N., Fait, A., and Barak, S. (2013).

Growth Platform-Dependent and -Independent Phenotypic and Metabolic Responses of

Arabidopsis and Its Halophytic Relative, Eutrema salsugineum, to Salt Stress. Plant Physiol.

162: 1583–1598.

Kazachkova, Y., Eshel, G., Pantha, P., Cheeseman, J.M., Dassanayake, M., and Barak, S. (2018).

Halophytism: What Have We Learnt From Arabidopsis thaliana Relative Model Systems?

Plant Physiol. **178**: 972–988.

Lakehal, A. et al. (2019). A Molecular Framework for the Control of Adventitious Rooting by

TIR1/AFB2-Aux/IAA-Dependent Auxin Signaling in Arabidopsis. Mol. Plant **12**: 1499–1514.

Langmead, B. and Slazberg, S.L. (2013). Fast gapped-read alignmnet with Bowtie 2. Nat.

Methods **9**: 357–359.

Lee, Y.P., Funk, C., Erban, A., Kopka, J., Köhl, K.I., Zuther, E., and Hincha, D.K. (2016). Salt

stress responses in a geographically diverse collection of Eutrema/Thellungiella spp.

accessions. Funct. Plant Biol. **43**: 590–606.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. **15**: 1–34.

MacLeod, M.J.R., Dedrick, J., Ashton, C., Sung, W.W.L., Champigny, M.J., and Weretilnyk, E.A.

(2015). Exposure of two Eutrema salsugineum (Thellungiella salsuginea) accessions to

water deficits reveals different coping strategies in response to drought. Physiol. Plant.

155: 267–280.

Maere, S., Heymans, K., and Kuiper, M. (2005). BiNGO: A Cytoscape plugin to assess

overrepresentation of Gene Ontology categories in Biological Networks. Bioinformatics **21**:

3448–3449.

Mannor, S., Jin, X., Han, J., Jin, X., Han, J., Jin, X., Han, J., and Zhang, X. (2011). K-Means

Clustering. In Encyclopedia of Machine Learning (Springer US: Boston, MA), pp. 563–564.

Martínez-Vilalta, J., Sala, A., Asensio, D., Galiano, L., Hoch, G., Palacio, S., Piper, F.I., and

Lloret, F. (2016). Dynamics of non-structural carbohydrates in terrestrial plants: a global

synthesis. Ecol. Monogr. **86**: 495–516.

McInnes, L., Healy, J., Saul, N., and Großberger, L. (2018). UMAP: Uniform Manifold

Approximation and Projection. J. Open Source Softw. **3**: 861.

Oh, D.-H., Hong, H., Lee, S.Y., Yun, D.-J., Bohnert, H.J., and Dassanayake, M. (2014). Genome

Structures and Transcriptomes Signify Niche Adaptation for the Multiple-Ion-Tolerant

Extremophyte Schrenkiella parvula. Plant Physiol. **164**: 2123–2138.

Oh, D.H. and Dassanayake, M. (2019). Landscape of gene transposition-duplication within the

Brassicaceae family. DNA Res.

Orsini, F. et al. (2010). A comparative study of salt tolerance parameters in 11 wild relatives of

Arabidopsis thaliana. J. Exp. Bot. **61**: 3787–3798.

Pantha, P. and Dassanayake, M. (2020). Living with Salt. Innov. **1**: 100050.

Pantha, P., Oh, D.-H., Longstreth, D., and Dassanayake, M. (2021). Living with high potassium:

an asset or a hindrance. bioRxiv: 2021.07.01.450778.

Pareek, A., Dhankher, O.P., and Foyer, C.H. (2020). Mitigating the impact of climate change on

plant productivity and ecosystem sustainability. J. Exp. Bot. **71**: 451–456.

Pertea, M., Kim, D., Pertea, G.M., Leek, J.T., and Salzberg, S.L. (2016). Transcript-level

expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nat.

Protoc. **11**: 1650–1667.

Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.-C., Mendell, J.T., and Salzberg, S.L.

(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.

Nat. Biotechnol. **33**: 290–295.

Philippar, K., Ivashikina, N., Ache, P., Christian, M., Lüthen, H., Palme, K., and Hedrich, R.

(2004). Auxin activates KAT1 and KAT2 , two K + -channel genes expressed in seedlings of

Arabidopsis thaliana. Plant J. **37**: 815–827.

Pinheiro, C., Dickinson, E., Marriott, A., Ribeiro, I.C., Pintó-Marijuan, M., António, C., Zarrouk,

O., Chaves, M.M., Dodd, I.C., Munné-Bosch, S., Thomas-Oates, J., and Wilson, J. (2019).

Distinctive phytohormonal and metabolic profiles of Arabidopsis thaliana and Eutrema

salsugineum under similar soil drying. Planta **249**: 1417–1433.

Poorter, H. (1989). Interspecific variation in relative growth rate: on ecological causes and

physiological consequences. Causes consequences Var. growth rate Product. High. plants

24: 45–68.

Prerostova, S., Dobrev, P.I., Gaudinova, A., Hosek, P., Soudek, P., Knirsch, V., and Vankova, R.

(2017). Hormonal dynamics during salt stress responses of salt-sensitive Arabidopsis

thaliana and salt-tolerant Thellungiella salsuginea. Plant Sci. **264**: 188–198.

De Rybel, B. et al. (2010). A Novel Aux/IAA28 Signaling Cascade Activates GATA23-Dependent

Specification of Lateral Root Founder Cell Identity. Curr. Biol. **20**: 1697–1706.

Santiago-Rosario, L.Y., Harms, K.E., Elderd, B.D., Hart, P.B., and Dassanayake, M. (2021). No

escape: The influence of substrate sodium on plant growth and tissue sodium responses.

Ecol. Evol.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of

image analysis. Nat. Methods.

Shabala, S. and Cuin, T.A. (2008). Potassium transport and plant salt tolerance. Physiol. Plant.

133: 651–669.

Shabala, S., Demidchik, V., Shabala, L., Cuin, T.A., Smith, S.J., Miller, A.J., Davies, J.M., and

1074 **Newman, I.A.** (2006). Extracellular Ca2+ ameliorates NaCl-induced K+ loss from

1075 Arabidopsis root and leaf cells by controlling plasma membrane K +-permeable channels.

1076 Plant Physiol. **141**: 1653–1665.

1077 **Shamustakimova, A.O., Leonova, T.G., Taranov, V.V., de Boer, A.H., and Babakov, A.V.** (2017).

1078 Cold stress increases salt tolerance of the extremophytes Eutrema salsugineum

1079 (Thellungiella salsuginea) and Eutrema (Thellungiella) botschantzevii. J. Plant Physiol. **208**:

1080 128–138.

1081 **Simopoulos, C.M.A., MacLeod, M.J.R., Irani, S., Sung, W.W.L., Champigny, M.J., Summers,**

1082 **P.S., Golding, G.B., and Weretilnyk, E.A.** (2020). Coding and long non-coding RNAs provide

1083 evidence of distinct transcriptional reprogramming for two ecotypes of the extremophile

1084 plant Eutrema salsugineum undergoing water deficit stress. BMC Genomics **21**: 396.

1085 **Slama, I., Abdelly, C., Bouchereau, A., Flowers, T., and Savouré, A.** (2015). Diversity,

1086 distribution and roles of osmoprotective compounds accumulated in halophytes under

1087 abiotic stress. Ann. Bot. **115**: 433–447.

1088 **Sperdouli, I. and Moustakas, M.** (2012). Interaction of proline, sugars, and anthocyanins during

1089 photosynthetic acclimation of Arabidopsis thaliana to drought stress. J. Plant Physiol. **169**:

1090 577–585.

1091 **Sun, J. et al.** (2009). Calcium mediates root K+/Na+ homeostasis in poplar species differing in

1092 salt tolerance. Tree Physiol. **29**: 1175–1186.

1093 **Taji, T., Seki, M., Satou, M., Sakurai, T., Kobayashi, M., Ishiyama, K., Narusaka, Y., Narusaka,**

1094 **M., Zhu, J.K., and Shinozaki, K.** (2004). Comparative genomics in salt tolerance between

1095 arabidopsis and arabidopsis-related halophyte salt cress using arabidopsis microarray.

Plant Physiol. **135**: 1697–1709.

Takahashi, F. and Shinozaki, K. (2019). Long-distance signaling in plant stress response. Curr.

Opin. Plant Biol. **47**: 106–111.

Takahashi, Y., Zhang, J., Hsu, P.-K., Ceciliato, P.H.O., Zhang, L., Dubeaux, G., Munemasa, S.,

Ge, C., Zhao, Y., Hauser, F., and Schroeder, J.I. (2020). MAP3Kinase-dependent SnRK2-

kinase activation is required for abscisic acid signal transduction and rapid osmotic stress

response. Nat. Commun. **11**: 12.

Thalmann, M., Pazmino, D., Seung, D., Horrer, D., Nigro, A., Meier, T., Kölling, K., Pfeifhofer,

H.W., Zeeman, S.C., and Santelia, D. (2016). Regulation of leaf starch degradation by

abscisic acid is important for osmotic stress tolerance in plants. Plant Cell **28**: 1860–1878.

Tran, K.-N. et al. (2021). Balancing growth amidst salinity stress – lifestyle perspectives from

the extremophyte model Schrenkiella parvula. bioRxiv: 6.

Tug, G.N., Yaprak, A.E., and Vural, M. (2019). The Floristical, Ecological, and Syntaxonomical

Characteristics of Salt Marshes and Salt Steppes in Turkey. In Sabkha Ecosystems (Springer:

Cham), pp. 413–446.

Volkov, V. and Amtmann, A. (2006). Thellungiella halophila, a salt-tolerant relative of

Arabidopsis thaliana, has specific root ion-channel features supporting K+/Na+

homeostasis under salinity stress. Plant J. **48**: 342–353.

Volkov, V., Wang, B., Dominy, P.J., Fricke, W., and Amtmann, A. (2004). Thellungiella

halophila, a salt-tolerant relative of Arabidopsis thaliana, possesses effective mechanisms

to discriminate between potassium and sodium. Plant, Cell Environ. **27**: 1–14.

Wang, G., DiTusa, S.F., Oh, D.H., Herrmann, A.D., Mendoza-Cozatl, D.G., O'Neill, M.A., Smith,

1118 **A.P., and Dassanayake, M.** (2021). Cross species multi-omics reveals cell wall

1119 sequestration and elevated global transcript abundance as mechanisms of boron tolerance

1120 in plants. New Phytol. **230**: 1985–2000.

1121 **Wang, G., Oh, D.H., and Dassanayake, M.** (2020). GOMCL: A toolkit to cluster, evaluate, and

1122 extract non-redundant associations of Gene Ontology-based functions. BMC

1123 Bioinformatics **21**: 1–9.

1124 **Wang, Y., Li, K., and Li, X.** (2009). Auxin redistribution modulates plastic development of root

1125 system architecture under salt stress in Arabidopsis thaliana. J. Plant Physiol. **166**: 1637–

1126 1645.

1127 **Wu, H.-J. et al.** (2012). Insights into salt tolerance from the genome of Thellungiella salsuginea.

1128 Proc. Natl. Acad. Sci. **109**: 12219–12224.

1129 **Yin, J., Gosney, M.J., Dilkes, B.P., and Mickelbart, M. V.** (2018). Dark period transcriptomic and

1130 metabolic profiling of two diverse Eutrema salsugineum accessions. Plant Direct **2**: e00032.

1131 **Yu, Z., Duan, X., Luo, L., Dai, S., Ding, Z., and Xia, G.** (2020). How Plant Hormones Mediate Salt

1132 Stress Responses. Trends Plant Sci. **25**: 1117–1130.

1133 **Zandalinas, S.I., Fritschi, F.B., and Mittler, R.** (2021). Global Warming , Climate Change , and

1134 Environmental Pollution : Recipe for a Multifactorial Stress Combination Disaster. Trends

1135 Plant Sci. **26**: 588–599.

1136 **van Zelm, E., Zhang, Y., and Testerink, C.** (2020). Salt Tolerance Mechanisms of Plants. Annu.

1137 Rev. Plant Biol. **71**: 403–433.

1138 **Zhang, M., Huang, P., Ji, Y., Wang, S., Wang, S., Li, Z., Guo, Y., Ding, Z., Wu, W., and Wang, Y.**

1139 (2020). <scp>KUP</scp> 9 maintains root meristem activity by regulating K^+ and auxin

1140   homeostasis in response to low K. EMBO Rep. **21**: e50164.

1141   **Zhao, C., Zhang, H., Song, C., Zhu, J., and Shabala, S.** (2020). Mechanisms of Plant Responses

1142   and Adaptation to Soil Salinity. Innov. **1**: 100017.

1143   **Zhu, X. et al.** (2018). K + Efflux Antiporters 4, 5, and 6 Mediate pH and K + Homeostasis in

1144   Endomembrane Compartments. Plant Physiol. **178**: 1657–1678.

1145

Figure S1: Effect of salt stress on phenotype of *Schrenkiella parvula* (Sp), *Eutrema salsugineum* (Es) and *Arabidopsis thaliana* (At). [A] Four-week-old hydroponic grown plants were treated with indicated salt concentrations. [B] Experimental design and sample scheme. There were at least 4 replicates used for ionomic and metabolomic profiling, and 3 replicates for transcriptomic profiling with at least 4 plants per replicate.

Figure 1: Sodium accumulation and its effect on nutrient balance in the extremophytes compared to *A. thaliana*. [A] Na^+ and [B] K^+ content in roots and shoots. Data are mean ± SD (n ≥ 4, at least 4 plants per replicate). Asterisks indicate significant differences ($p ≤ 0.05$) between the treated samples and their respective control samples, determined by Student's t-test. Open circles indicate the biological replicates. [C] Macro- and micro-nutrient fold changes between treated and its respective control sample. [D] Percent abundance of nutrients that were significantly different at basal levels among the three species. The three axes of the ternary plots are marked with *A. thaliana, S. parvula*, and *E. salsugineum*. The gridlines in species designated colors point to the relevant % abundance of the element in each axis. Significant difference in elemental abundance was determined by one-way ANOVA with post-hoc Tukey's test at $p ≤ 0.05$.

Figure 2: Overall metabolic preparedness and adjustments in the roots of the extremophytes compared to *A. thaliana*. [A] Pairwise correlation of 716 quantified metabolites (182 known, and 534 unknown) across all conditions and species. Correlation calculated using Pearson correlation coefficients. [B] Total number of metabolites that significantly changed in abundance in each species compared to its respective control sample. Known metabolite numbers given in parenthesis. [C] Basal level metabolite abundances in all three species for sugars, amino acids, and their derivatives. [D] Hierarchical clustering of known metabolites shown in [C]. Blue box indicates clusters of interest. Treatment concentrations are 150 and 250 mM NaCl; Treatment durations are 24 and 72 hr. Significant differences were determined by one-way ANOVA with post-hoc Tukey's test at $p \leq 0.05$ (n = 4, at least 4 plants per replicate).

Figure S2: Overall shoot metabolite responses in the extremophytes compared to *A. thaliana*. [A] Pairwise correlation of 725 quantified metabolites across all conditions and species. Correlation calculated using Pearson correlation coefficients. [B] Total number of metabolites that significantly changed in abundance in each species compared to its respective control sample. Known metabolite numbers given in parenthesis. [C] Basal level metabolite abundances in all three species for sugars, amino acids, and their derivatives. Significant differences were determined by one-way ANOVA with post-hoc Tukey's test at $p \leq 0.05$ (n = 4, at least 4 plants per replicate).

Figure 3: Transcriptomic overview *S. parvula, E. salsugineum,* and *A. thaliana* in response to salt. [A] Overall transcriptome clustering using one-to-one ortholog groups (OGs) in shoots and roots from all replicates in all conditions. [B] Number of differently expressed genes (DEGs) in response to salt treatments. [C] Functionally enriched processes clustered among OGs that included a DEG from at least one species in roots and shoots. Functional annotations were based on GO annotations assigned to OGs. The bar graphs assigned to each cluster represent percent allocation of induced and suppressed DEGs with the number of genes that did not significantly change in each species given in the white space between induced (red) and suppressed (blue) bars. [D] Dominant gene expression clusters of orthologs across time points and species from roots and shoots that included an *A. thaliana* ortholog that was either induced or suppressed in response to salt. Differentially expressed genes identified at *p*-adj ≤ 0.01. N= 3 (at least 4 plants per replicate).

Table 1: Enriched cellular processes of ortholog groups at basal expression level clustered in shoots and roots of *S. parvula* (purple), *E. salsugineum*, (red) and *A. thaliana* (green).

Patterns	Clstr	# GO	# genes	Cumulative genes %	Functional category
2159 OGs (Sh:1215 - Rt:1351)	1	101	703	51.4%	Hormone and abiotic/biotic stimuli signaling
	2	58	458	70.2%	Ion and vesicular transport
	3	65	417	77.6%	Organ development
	4	26	337	82.5%	Transcriptional control of organ development
2042 OGs (Sh:1141 - Rt:1241)	1	74	708	63.8%	Hormone and abiotic stimuli regulation
	2	83	497	88.1%	RNA and amino acid processes
	3	53	437	99.4%	Organelle biogenesis
	4	6	23	99.8%	Phospholipid metabolism
1898 OGs (Sh:1134 - Rt:1084)	1	160	776	67.4%	Hormone and abiotic stimuli regulation
	2	100	520	88.9%	RNA and amino acid processes
	3	29	376	99%	Ion and vesicular transport
	4	4	112	99.7%	Homeostasis processes
1879 OGs (Sh:1098 - Rt:1050)	1	32	606	52.8%	Organ development
	2	58	519	74.0%	Hormone and abiotic/biotic stimuli signaling
	3	21	324	80.6%	Transcriptional regulation of development
	4	11	291	88.7%	Lipid metabolism
2086 OGs (Sh:1199 - Rt:1239)	1	44	671	65.8%	RNA metabolism and gene expression
	2	11	331	82.9%	Shoot and flower development
	3	19	268	91.5%	Cell cycle regulation
	4	4	159	95.5%	Abiotic stimuli responses
2225 OGs (Sh:1321 - Rt:1263)	1	106	775	55.3%	Hormone and abiotic/biotic stimuli signaling
	2	92	572	72.6%	Organ development and reproduction
	3	74	398	84.4%	Amino acid & lipid metabolism
	4	43	348	92.1%	Ion and vesicular transport

Normalized expression

Figure S3: Transcriptomic profiles of *S. parvula*, *E. salsugineum,* and *A. thaliana* were diverged in response at basal level and under salt stress. [A] roots, [B] shoots. Correlation was calculated using log2 fold change of treated over control samples from differentially expressed ortholog pairs which included DEGs from at least one condition in one species using Pearson correlation coefficients. *S. parvula* (purple), *E. salsugineum* (red) and *A. thaliana* (green).

Figure S4: Sub-clustering of dominant cluster - Response to stresses. Overlap and similarity between sub-clusters determined using GO-MCL-sub and cumulative number of ortholog groups (OGs) in [A and B] roots and [C and D] shoots sub-clusters.

Figure S5: Species dependent responses to salt treatments in *Schrenkiella parvula* and *Eutrema salsugineum*. Differently expressed genes (DEGs) in *S. parvula* [A] root and [B] shoot and *E. salsugineum* [C] root and [D] shoot in response to 150 and 250 mM NaCl stress at 3 and 24 hr of treatment. Yellow highlighted data points in the Upset plots for each panel show DEGs that were uniquely expressed at 250 mM salt treatments. Functionally enriched processes for these selected DEGs are shown in the horizontal bar graphs given as insets outlined in yellow for each panel. Differentially expressed genes identified at *p*-adj ≤ 0.01. N = 3 (at least 4 plants per replicate).

Figure 4: Root growth responses in line with gene expression mediated by auxin and ABA in *Arabidopsis thaliana, Schrenkiella parvula,* and *Eutrema salsugineum* when treated with salt. [A] Expression responses of auxin/indole acetic acid (Aux/IAA) repressors and type 2C protein phosphatases (PP2Cs) that regulate auxin and ABA signaling. [B] Curated genes from lateral root development (GO:0048527) and primary root development (GO:0080022) used to infer root growth phenotypes based on functional genetic studies in *A. thaliana*. Orthologs in the three species are indicated as high (yellow), low (blue), and indistinguishable levels between species (no-color) assigned to relative basal expression levels and their effect inferred with arrows (up – promote growth and down – suppress growth). Genes associated with Auxin, ABA, and other hormones were labeled in pink, gold, and black respectively. [C] Root growth assessment of 12-day-old seedlings in response to 150 mM NaCl treated for 7 days. Quantification of [D] primary root growth, [E] number of lateral roots, and [F] lateral root density. Data given as mean ± SD (n = 3, at least 7 plants per replicate). Open circles indicate the measurements from each plant. Asterisks indicate significant differences ($p ≤ 0.05$) between treated samples and their respective control samples, determined by one-way ANOVA with post-hoc Tukey's test. [G] Gene co-expression clusters of differentially expressed orthologs that showed similar expression pattern in *A. thaliana* and *E. salsugineum* roots compared to a different pattern observed for *S. parvula* under salt stress.

Figure S6: Hormonal signaling pathways in response to salt treatment in *Arabidopsis thaliana, Schrenkiella parvula,* and *Eutrema salsugineum*. Expression responses of genes associated with hormonal signaling pathways obtained from KEGGs. Note that only genes that were significantly different at either basal level condition or under salt treatments were shown. Orthologs in the three species are indicated as high (yellow), low (blue), and indistinguishable levels between species (no-color) assigned to relative basal expression levels

Figure S7: Lateral root development gene network adopted from Banda et al., 2019 and Rybel et al., 2010. The expression levels of orthologs were annotated as high (H), medium (M), or low (L) relative to each other at basal levels in each ortholog group. Specific paths in the network in which *S. parvula* and *E. salsugineum* consistently regulated to inhibit lateral root formation were highlighted in ribbons across the network in species-specific colors.

Figure S8: Root growth of *E. salsugineum*, *S. parvula* and *A. thaliana* before and after salt treatments. *S. parvula* (Sp) promoted primary root growth and reduced lateral root formation while *E. salsugineum* (Es) showed slower primary root growth with reduced lateral root number compared to *A. thaliana* (At) roots that showed severe root growth inhibition under salt stress. Five-day-old seedlings were treated for 11 days on plates supplemented with 150 mM NaCl.

Figure 5: Coordination between differently expressed genes and differently abundant metabolites during responses to salt in *Arabidopsis thaliana, Schrenkiella parvula,* and *Eutrema salsugineum*. [A] The difference in percent change in total differentially expressed genes (DEGs) and total differently abundant metabolites (DAMs) in response to salt treatments. [B] % DEGs and % DAMs involved in the metabolism of amino acids, sugars, and their immediate derivatives (GO:0006520 and GO:0005975). [C and D] Selected pathways in amino acids and sugar metabolism. Line graphs represent normalized Log2 relative metabolite abundance. Boxplots represent normalized expression values. Center line shows median; box indicates interquartile range (IQR); notch is for 1.58 × IQR/sqrt(n); whiskers show 1.5 × IQR. Asterisks indicate DEGs and DAMs assigned using 3-4 biological replicates. Early (E) and late (L) responses for transcripts refer to 3 and 24 hr. E and L responses for metabolites refer to 24 and 72 hr. Metabolites are shown in the backbone of the pathway while genes encoding for key enzymes/transcription factors are placed beside arrows. Black labels in pathway maps indicate quantified metabolites and genes while grey labels indicate metabolites not quantified.

Figure S9: Metabolite abundance and expression of associated genes encoding the enzymes in shikimic acid and ascorbic acid pathways. [A] percent change in total differentially expressed genes (DEGs) and total differently abundant metabolites (DAMs) in response to salt treatments. [B] pathway involved in proline metabolism. [C] pathway involved in the conversion from phosphoenolpyruvate to chorismate. [D] pathway involved in ascorbic acid metabolism. Line graphs represent normalized Log2 relative metabolite abundance. Boxplots represent normalized expression values. Center line shows median; box indicates interquartile range (IQR); notch is for 1.58 × IQR/sqrt(n); whiskers show 1.5 × IQR. Asterisks indicate DEGs (Differently expressed genes) and DAMs (Differently abundant metabolites). Early (E) and late (L) responses for transcripts refer to 3 and 24 hr. E and L responses for metabolites refer to 24 and 72 hr. Metabolites are shown in the backbone of the pathway while genes encoding for key enzymes/transcription factors are placed beside arrows. Black labels in pathway maps indicate quantified metabolites and genes while grey labels indicate metabolites not shown.

Figure 6: Basal and differential expression associated with salt and nutrient transport in *Arabidopsis thaliana, Schrenkiella parvula,* and *Eutrema salsugineum*. [A] Basal level expression given as fold changes between species. The dashed lines indicate 2-fold differences to highlight genes either highly suppressed or highly expressed in the extremophytes in roots at control conditions. [B] Aggregated expression profiles of transporters/channels associated with Na⁺/K⁺ transport activity during salt treatments. [C] Selected cation transporters that showed higher gene copy number in extremophytes in ortholog groups and their expression during salt treatments. Nodes of each ortholog network is assigned with species colors connected by edges based on the relationship properties of the homologs. Edges show relationships for co-linear reciprocal orthologs (cl-rc), co-linear unidirectional orthologs (cl-uni), transposed -unidirectional duplicates (tr-uni), or tandem duplicated-unidirectional paralogs (td-uni). Asterisks indicate differentially expressed genes identified at *p-adj ≤ 0.05 and ** p-adj ≤ 0.01. N= 3 (at least 4 plants per replicate). [D] Genes associated with nutrient transport identified in Figure 3C.

Figure 7: Summary of ionomic, metabolomic, transcriptomic, and phenotypic adjustments and preadaptation trends to salt stress in *Arabidopsis thaliana*, *Schrenkiella parvula* and *Eutrema salsugineum* roots. Top 9 metabolites that increased abundance in *S. parvula* were Fru, fructose; Asn, asparagine, Raf, raffinose, DHA, dehydroascorbic acid; Pro, proline; Shkm, shikimic acid; Suc, sucrose, GABA, gamma-aminobutyric acid; Gln, glutamine. Orthologs in the three species are indicated as high (yellow), low (blue) assigned to relative basal expression levels. Blue and red indicate the induction and suppression gene expression, respectively. KEA5, K$^+$ efflux antiporter 5; KUP9, K$^+$ uptake permease 9; KAT1, K$^+$ channel in *Arabidopsis thaliana* 1; GLRs, glutamate-like receptors; NHX1/2, Na$^+$/H$^+$ exchanger1/2; V-ATPases, vacuoles-ATPases; SOS1, salt overly sensitive 1; GORK, gated outwardly-rectifying K$^+$ channel.