Clarifying intercellular signalling in yeast: *Saccharomyces cerevisiae* does

not undergo a quorum sensing-dependent switch to filamentous growth

Michela Winters[a], Violetta Aru[b], Kate Howell[a,#], and Nils Arneborg[b]

[a] School of Agriculture and Food, Faculty of Veterinary and Agricultural Science,

University of Melbourne, Parkville 3010, Australia

[b] Department of Food Science, University of Copenhagen, Frederiksberg 1958,

Denmark

Running Head: Unipolar budding of *Saccharomyces cerevisiae*

#Address correspondence to Kate Howell, khowell@unimelb.edu.au

Word counts

Abstract: 235

Importance: 150

Text: 5244

 ABSTRACT

Saccharomyces cerevisiae can alter its morphology to a filamentous form

associated with unipolar budding in response to environmental stressors. Induction of

filamentous growth is suggested under nitrogen deficiency in response to alcoholic

signalling molecules through a quorum sensing mechanism. To investigate this claim,

we analysed the budding pattern of *S. cerevisiae* cells over time under low nitrogen

while concurrently measuring cell density and extracellular metabolite concentration. We

found that the proportion of cells displaying unipolar budding increased between local

cell densities of 4.8×10^6 and 5.3×10^7 cells/ml. However, the observed increase in

unipolar budding could not be reproduced when cells were prepared at the critical cell

density and in conditioned media. Removing the nutrient restriction by growth under

high nitrogen conditions also resulted in an increase in unipolar budding between local

cell densities of 5.2×10^6 and 8.2×10^7 cells/ml, but there were differences in metabolite

concentration compared to the low nitrogen conditions. This suggests that neither cell

density, metabolite concentration, nor nitrogen deficiency were necessary or sufficient to

increase the proportion of unipolar budding cells. It is therefore unlikely that quorum

sensing is the mechanism controlling the switch to filamentous growth in *S. cerevisiae*.

Only a high concentration of the putative signalling molecule, 2-phenylethanol resulted

in an increase in unipolar budding, but this concentration was not physiologically

relevant. We suggest that the compound 2-phenylethanol acts through a toxicity

mechanism, rather than quorum sensing, to induce filamentous growth.

IMPORTANCE

Investigating dimorphism in the model organism *Saccharomyces cerevisiae* has been instrumental in understanding the signalling pathways that control hyphal growth and virulence in human pathogenic fungi. Quorum sensing was proposed to signal morphogenesis in *S. cerevisiae* populations. This mechanism requires the switch to filamentous growth to occur at a critical quorum sensing molecule concentration corresponding to a critical cell density. However, evidence for this mechanism is sparse and limited by the use of non-physiologically relevant concentrations of signalling metabolites. Our study designed a methodology to address this gap and may be applied to further studies of dimorphism in other types of yeasts. A significant implication of our findings is that morphogenesis does not occur in *S. cerevisiae* via a quorum sensing mechanism, and this important definition needs to be corrected. Mechanistic studies to understand dimorphism in yeasts, by considering metabolite concentrations, will further shed light onto this important cellular behaviour.

KEYWORDS

Saccharomyces cerevisiae, filamentous growth, budding pattern, signalling metabolites, cell density

INTRODUCTION

The yeast *Saccharomyces cerevisiae* has the ability to exist in different multicellular forms depending on its environment. In liquid environments, the yeast cells can be sessile or planktonic and may form flocs and flors, and in solid and semi-solid substrates, colonies, biofilms, filaments and mats are observed (1). Filamentous growth of *S. cerevisiae* commonly occurs under nutrient starvation and has been hypothesised as a mechanism to allow the cells to grow into their immediate environment and forage for available nutrients (2, 3). This morphogenesis is described as pseudohyphal or invasive depending on whether the cells exist as diploids or haploids and can be observed both at a colony and cellular level (2, 4-9). When growing on agar plates, colony morphology and invasion reveal filamentous growth, and these macro-observations are determined by budding pattern and cell shape that indicate filamentous growth at the cellular level (2, 4-6, 10, 11).

Budding is the method by which *S. cerevisiae* cells divide during their mitotic cell cycle. The budding pattern is determined by the orientation of new buds with respect to the mother-daughter junction and can be differentiated as axial, bipolar or unipolar (Fig. 1) (2, 5, 12-15). Whether the cell is a diploid or haploid will determine which budding pattern is exhibited when cells grow planktonically (12, 13). Haploid cells bud in an axial manner whereby the new bud consistently emerges adjacent to or overlapping the pole of the previous mother-daughter junction. Diploid cells, however, bud according to a bipolar pattern, where buds can emerge either adjacent to the mother-daughter junction or at the opposite pole. Usually, the first bud site of a new daughter cell has a strong opposite pole bias, while second and subsequent buds have no bias towards a pole (2,

12, 14, 15). Both haploids and diploids transition to a unipolar budding pattern when

cells switch to filamentous growth. This pattern occurs when the daughter buds

consistently from the pole opposite the junction to its mother (2, 5, 7-10). Cell division is

thus polarized by the serial reiteration of unipolar budding away from the mass of cells

in the colony into chains of interconnected cells (2).

S. cerevisiae cells undergoing filamentous growth have a higher proportion of

unipolar budding compared to non-filamentous cells. Previous research established that

when two bud sites are observed, non-filamentous cells show 70-71% unipolar budding,

while filamentous cells show 90-100% unipolar budding (2, 5). For cells with three bud

sites, this proportion of unipolar budding was observed to be 26% for non-filamentous

cells and 97% for filamentous cells (5). Furthermore, under filamentous growth inducing

conditions (nitrogen deficiency), unipolar budding was found to increase by an average

of 35% (16). Observation of a higher proportion of unipolar budding cells thus signifies

that cells are on track to grow as filaments.

Filamentous growth is induced in *S. cerevisiae* cells in response to different

environmental cues and stressors, including nutrient deficiency and signalling

molecules. Filamentous cells are considered more able to scavenge nutrients when

exposed to a variety of extracellular stressors (17). Nutrient starvation, including lack of

nitrogen (2, 9, 18-20), can trigger the morphological change to filamentous growth. An

additional adaptation to nitrogen starvation is the secretion of autoregulatory aromatic

alcohols that stimulate filamentous growth through a quorum sensing mechanism (18,

21-23).

Quorum sensing is a form of intercellular signalling where interactions between

organisms are mediated by intercellular signalling molecules (24). Quorum sensing

involves a cell-density dependent regulation of a behaviour that occurs after a critical

signal concentration of quorum sensing molecules is achieved (25). Quorum sensing

molecules are excreted and accumulate in the extracellular environment proportional to

cell density. At the critical cell density a corresponding threshold concentration of

quorum sensing molecules is reached and a synchronised response is triggered in the

community of cells (26-31). A prior study observed a quorum sensing-controlled

mechanism for the switch to filamentous growth in *S. cerevisiae,* connecting nutrient

sensing with aromatic alcohol secretion. Nitrogen and cell density were demonstrated to

regulate the production of three auto-induced aromatic alcohols via a positive feedback

loop. These included tyrosol (produced in the range 1-8 µM), 2-phenylethanol (in the

range of 1-8 µM) and tryptophol (in the range of 1-2 µM). 2-phenylethanol and

tryptophol, but not tyrosol, were then found to stimulate filamentous growth when added

exogenously at ≥20 µM under nitrogen limiting conditions (18). Further studies have

linked exogenous addition of different alcohols to induction of morphogenesis under

both high (32) and low nitrogen (9) conditions. These include 1-butanol (9), isoamylol (9,

32), 1-propanol (9, 32) and n-hexanol (32). However, all these studies induced

morphogenesis by using exogenous alcohol concentrations much higher than those

found to be produced when the cell is growing in defined or undefined media (25).

Intercellular signalling interactions need to be defined by specific criteria to be

accurately identified and understood. Previous definitions of quorum sensing have

tended to be very broad so that any morphological behaviours in yeast may have been

inaccurately categorised as being a result of quorum sensing. In our previous work, we

set out a series of criteria for what constitutes an intercellular signalling molecule and

quorum sensing molecule to promote clarity in classifying microbial interactions (25).

When these defined criteria were applied to studies of quorum sensing-controlled

filamentous growth in *S. cerevisiae* we found that two criteria were consistently not met

in prior research (25). Firstly, there is a lack of evidence that the switch in cellular

behaviour occurs at a critical quorum sensing molecule concentration corresponding to

a critical cell density. Secondly, no studies used physiologically relevant concentrations

of intercellular signalling molecules to trigger the switch to filamentous growth.

In the current study, we aimed to address these two missing criteria on the role of

quorum sensing in the switch to filamentous growth in *S. cerevisiae*. We designed a

new methodology which uses budding pattern as a proxy for filamentous growth. This

involved observing an individual cell, growing over time, to determine the orientation of

buds from new daughter cells. By using an oCelloScope™ to capture time-lapse images

of budding yeast growing in liquid media, we observed a switch to a filamentous

budding pattern. Since our method simultaneously quantified the budding pattern, cell

density and metabolite concentration under physiological conditions, we successfully

identified the critical cell density and metabolite concentration at which cells switched to

filamentous growth. An untargeted metabolome approach through proton (^{1}H) NMR

meant both previously reported and putative quorum sensing molecules could be

monitored. Next, we individually assessed the role of cell density, metabolites, and

nitrogen concentration in triggering this switch to filamentous growth. Additionally, we

investigated the role of exogenous addition of 2-phenylethanol, a putative quorum

sensing molecule, in morphogenesis. Together, our findings suggest that *S. cerevisiae*

does not undergo a quorum sensing-dependent switch to filamentous growth, and that

2-phenylethanol acts through a toxicity mechanism rather than as a quorum sensing

molecule. Finally, the novel methodology developed here opens the way for further

studies of dimorphism in other strains and species of yeasts.

RESULTS

157 **RESULTS**

S. cerevisiae Σ1278b displays a switch in budding pattern under low nitrogen.

S. cerevisiae Σ1278b displays a switch in budding pattern under low nitrogen.

We designed a time series experiment such that *S. cerevisiae* strains could grow

in the same media to obtain an accurate link between our three factors of interest:

budding pattern, cell density and metabolite concentration. The cropped images

obtained from the oCelloScope™ were analyzed manually and the bud site sequence of

new daughters were recorded followed by a classification of their corresponding

budding pattern (Fig. 1). A representative example of this assignment process can be

seen in Figure 2.

The growth of *S. cerevisiae* strains S288c and Σ1278b were observed under low

nitrogen conditions over 30 hours (Fig. 3). Both S288c and Σ1278b strains grew

exponentially up to 20 hours after which their growth slightly declined (Fig. 3A).

A predominantly axial budding pattern was observed for the negative control

strain S288c for both the two and three bud sites, with no significant changes in budding

pattern observed over time (Fig. 3B). Strain Σ1278b budded in a predominantly bipolar

or unipolar pattern. Additionally, the proportion of unipolar budding was observed to

increase over time, with a corresponding decrease in bipolar budding, for both two and

three bud sites (Fig. 3D and E). An increase in the proportion of unipolar budding from

72% to 91% (Fig. 3D) was observed in cells with two bud sites. For cells with three bud

sites, it increased from 20% to 36% (Fig. 3E). Due to the growth limitation, most

probably caused by nitrogen starvation between 20 and 30 hours (Table S1), the cells

were only able to achieve two bud sites and therefore budding pattern data for three bud

sites were not available during this period (Fig. 3E). The significant (P≤ 0.05) increase in

filamentous unipolar budding occurred between the local cell densities of $4.8x10^6$ and

$5.3x10^7$ cells/ml within 10 and 20 hours of growth. These values signify the critical cell

density for the switch to filamentous growth.

Physiological concentrations of metabolites present at the critical cell density

and change in budding pattern.

To correlate the change in concentration of metabolites over time to the changes

in budding pattern and cell density observed, media from the time series experiments

were collected, measured by ^{1}H NMR spectroscopy, and compounds identified and

quantified with SigMa (33). In total, 30 compounds were identified in the media with

concentrations quantified for each time period (Table 1). Tryptophol was not identified in

our samples though we specially looked for it during the NMR identification stage. 2-

phenylethanol and tyrosol were present but did not show a significant increase in

concentration until 30 hours with very low concentrations prior to that (Table 1).

Metabolites potentially involved in the budding pattern switch were chosen from

the identified compound list as those that were not present as nutrients in the raw media

and had concentrations that increased during growth. The metabolite concentrations

that corresponded to the observed switch in budding pattern were 1-propanol (1.4 – 3.7

µM), 2-hydroxyisovalerate (0.7 – 1.3 µM), 2-oxoglutarate (4.4 – 9.3 µM), 2-

phenylethanol (0.7 – 1.5 µM), acetaldehyde (2.3 – 19.2 µM), acetate (17 - 160 µM),

acetoin (6.7 – 8.0 µM), ethanol (369 – 5403 µM), fumarate (0.5 – 1.1 µM), isoamylol (0.6

– 1.0 µM), isopropanol (0.4 – 2.1 µM), lactate (29 – 224 µM), malate (2.9 – 6.8 µM),

pyruvate (5.0 – 30.6 µM), succinate (2.5 – 12.0 µM), and tyrosol (0.3 – 0.8 µM). These

values represent the physiological metabolite concentration present at the critical cell

density at which cells switch to filamentous growth.

Critical cell density does not recapitulate the switch in budding pattern.

To establish whether the increase in unipolar budding was dependent on cell

density alone, we inoculated cells at the critical cell density, recorded at 10 hours, into

low nitrogen media (Fig. 4). This cell density was chosen as it was at the beginning of

the time period where we observed the significant increase in unipolar budding. At the

critical cell density, 71% of cells with two bud sites had a unipolar pattern (Fig. 4A). This

was analogous to 72% unipolar budding observed between 0 to 10 hours in the time

series experiment (Fig. 3D). This parallel was also seen for cells with three bud sites at

the critical cell density where we measured 23% unipolar budding (Fig. 4B) compared to

20% between 0 to 10 hours in the time series results (Fig. 3E). Since the critical cell

density did not replicate the increase in unipolar budding, these findings demonstrate

that cell density alone is not responsible for the behaviour.

Conditioned media and cells at the critical cell density do not recapitulate the

switch in budding pattern.

To test whether the combination of physiological metabolites and the critical cell

density was a trigger for the increase in unipolar budding, we determined the budding

pattern of cells growing in conditioned media from 20 hours of growth at the critical cell

density in low nitrogen media (Fig. 4). Growth of cells with two bud sites in conditioned

media had 71% unipolar budding (Fig. 4A) and cells with three bud sites had 18% (Fig.

4B). The proportion of unipolar budding observed between the critical cell density and

conditioned media conditions were not significantly different; both had 71% for two bud

sites (Fig. 4A), and 20% and 18% respectively for three bud sites (Fig. 4B). In both

cases the values paralleled the proportion of unipolar budding observed between 0 to

10 hours in the time series experiment; 72% for two bud sites (Fig. 3D) and 20% for

three bud sites (Fig. 3E). Thus, the combination of critical cell density and physiological

metabolite concentrations were also not accountable for the observed increase in

unipolar budding.

Growth under high nitrogen conditions replicates the switch budding pattern but

with different produced metabolite concentrations.

To determine whether budding pattern behaviour over time was impacted by

nitrogen conditions, we carried out the time series experiment in high nitrogen media.

Evidence for the differences in the nitrogen concentration between the two conditions is

provided in Table S1.

Similar to growth in low nitrogen, S288c and Σ1278b cells grew exponentially

from 0 to 20 hours of the experiment, followed by a slight decrease between 20 and 30

hours (Fig. 5A). However, there was a less pronounced decrease in growth rate for the

Σ1278b compared to the S288c under high nitrogen conditions compared to low

nitrogen conditions.

Budding pattern behaviour was not significantly different between the two

nitrogen conditions. Just as in low nitrogen, S288c showed a predominantly axial

budding pattern with no changes over time (Fig. 5B and C) and Σ1278b showed a

consistently bipolar or unipolar budding pattern (Fig. 5D and E). Again, we saw a

significant increase to a unipolar budding pattern accompanied by a significant decrease

in bipolar budding between 10 and 20 hours for cells with both two and three bud sites

(Fig. 5D and E). For cells with two bud sites we saw an increase in unipolar budding

from 67% to 84% (Fig. 5D). For cells with three bud sites, it increased from 23% to 43%

(Fig. 5E). The increase in unipolar budding was associated with local cell densities of

5.2×10^6 and 8.2×10^7 cells/ml, respectively, which signifies the critical cell density under

high nitrogen. These values are of the same order of magnitude as the low nitrogen

critical cell density (4.8×10^6 and 5.3×10^7 cells/ml). Between 20 and 30 hours under high

nitrogen conditions we were not able to observe many cells with three bud sites (Fig.

5E) due to the cell density becoming so high that single buds were no longer visible in

the images. For this reason, we did not see a statistically significant increase in the

budding pattern between 0 to 10 hours and 20 to 30 hours.

Media from the time series experiment under high nitrogen conditions were also

analysed by ^{1}H NMR to obtain compound concentrations over time. A list of the 30

compounds identified in our high nitrogen samples and their concentrations can be seen

in Table 2. Metabolites potentially involved in the budding pattern switch were again

chosen from the identified compound list as those that were not present as nutrients in

the raw media and had concentrations that increased during growth. Here the

metabolite concentrations which corresponded to the observed switch in budding

pattern were 1-propanol (1.4 – 3.4 µM), 2-hydroxyisovalerate (0.4 – 0.5 µM), 2-

oxoglutarate (1.5 – 2.0 µM), 2-phenylethanol (0.4 – 0.6 µM), acetaldehyde (2.0 – 18.0

µM), acetate (27 - 186 µM), acetoin (1.0 – 5.6 µM), ethanol (472 – 4972 µM), fumarate

(0.3 – 0.4 µM), isoamylol (0.7 – 0.9 µM), isopropanol (0.3 – 2.9 µM), lactate (33-231

µM), malate (2.1- 2.2 µM), pyruvate (11.5 – 38.8 µM), succinate (0.9 – 1.6 µM), and

tyrosol (0.3 – 0.6 µM).

Comparing these values to the concentration of metabolites produced under low

nitrogen conditions at 20 hours of growth, the alcohols 2-phenylethanol and tyrosol were

60% and 63% lower under higher nitrogen respectively, while 1-propanol, isoamylol,

and isopropanol concentrations were equivalent. The concentrations of some acids

were also lower under high nitrogen conditions, 2-hydroxyisovalerate (↓62%), 2-

oxogluatare (↓79%), fumarate (↓73%), malate (↓68%) and succinate (↓87%). However,

acetate and pyruvate were 16% and 27% higher under high nitrogen conditions,

respectively. Thus, while the same budding pattern behaviour was observed under

different nitrogen conditions, there were significant differences in the metabolome.

Collectively, these findings concur with our findings mentioned above that physiological

metabolite concentrations are not responsible for the observed increase in unipolar

budding.

Proposed signalling molecule, 2-phenylethanol, can induce unipolar budding at

high concentrations.

Exogenous 2-phenylethanol has been reported to induce cellular signalling in *S.*

cerevisiae (18, 21), but at much higher concentrations than those observed in growth

media. To investigate the effects of these non-physiological concentrations, we added to

low nitrogen media a concentration of 2-phenylethanol 13 times higher than those

produced by the cells in our growth experiments. This higher concentration was used in

previous studies to stimulate filamentous growth (18). Here we observed 91% unipolar

budding for two bud sites (Fig. 4A) and 43% for three bud sites (Fig. 4B). These

proportions of unipolar budding corresponded to the higher percentage of unipolar

budding observed in both time series experiments between 10 to 20 hours; 84-91% for

two bud sites (Fig. 3D and 5D) and 37-43% for three bud sites (Fig. 3E and 5E). These

results indicate that a high non-physiological concentration of 2-phenylethanol can

recapitulate the increase in unipolar budding, while physiological concentrations do not.

DISCUSSION

The transition to filamentous growth has been proposed as a cell density

dependent, chemically mediated switch in *S. cerevisiae.* This conclusion was drawn

from studies which examined the growth of yeasts, metabolites, and exogenous

compounds, but work has not conclusively shown the dependency of cellular density,

induction, and accumulation of a chemical signal (25).

The objective of this study was to address the research gaps in the transition to

filamentous growth in *S. cerevisiae.* Specifically, we focused on the switch to

filamentous growth at a critical quorum sensing molecule concentration corresponding

to a critical cell density, and that the intercellular signalling molecule concentrations

used to induce the response were physiologically relevant (25). To achieve this, we

designed a new methodology to correlate the filamentous growth of *S. cerevisiae* to

both cell density and metabolite concentration. We combined chemical cell

immobilisation in liquid media, automated time-lapse scanning microscopy, cell counting

and ^{1}H NMR. Filamentous yeast cells show a significantly higher proportion of unipolar

budding overall compared to non-filamentous cells (2, 5). Thus, by using changes in

budding pattern as a proxy for filamentous growth, this study was able to identify the

critical cell density at which the cells switched morphology and to correlate this to the

physiological concentrations of metabolites present.

Regardless of the nitrogen condition, both *S. cerevisiae* strains were in their

exponential growth phase during the first 20 hours of the experiments after which their

growth rate slightly declined (Fig. 3A and 5A). This trend was more pronounced for the

Σ1278b strain under nitrogen deficiency (Fig. 3A). This general decline in growth is likely

due to the accumulation of toxic metabolites that occurred toward the end of the

experiment and nutrient starvation.

The use of the oCelloScope™ for imaging enabled us to clearly observe the

budding pattern of single yeast cells over time. Previous work has applied this imaging

technique to bacteriological studies on real-time detection of bacterial growth and

antimicrobial susceptibility (34), and to quantify filamentous bacteria (35). The

methodology of monitoring yeast budding patterns presented in this paper is unique

since previous studies have used photomicrographs of developing pseudohyphae in

colonies growing on agar (2) or calcofluor staining of bud scars (5, 12). However, growth

on agar plates does not allow the monitoring of metabolite production, budding pattern,

and cell density simultaneously over time (2, 9, 18), and calcofluor staining does not

permit reliable assignment of the birth end of cells (5, 11).

The highest proportion of unipolar budding measured here for cells with two bud

sites was 85-91% (Fig. 3D and 5D). This finding is in accordance with previous studies

that observed >85% unipolar buds for filamentous cells (2, 5). Conversely, in cells with

three bud sites we found a maximum of 36-43% unipolar budding (Fig. 3E and 5E).

Even though we saw a significant increase over time, this proportion was relatively low

compared to the expected 97% from previous findings (5). A possible explanation for

this might be that prior studies have not always reported on the position of third bud

sites (2) and/or have tended to concentrate on cells at the periphery of the colony (2, 5).

Our experiment analysed the whole population of cells which could explain this lower

proportion. Additionally, our cells were in the exponential phase of growth and growing

in liquid culture, while previous studies used cells in the stationary phase (18) or

investigated the budding pattern of colonies growing on agar (2, 5).

S. cerevisiae strain S288c has a mutation in a gene required for filamentous

growth (FLO8)(36) and therefore acted as our negative control for filamentation.

Additionally, it is a haploid and thus expected to bud in a consistently axial manner (12,

13). Our findings are consistent with this since we observed S288c to bud axially at all

time points under both high and low nitrogen conditions. Additionally, as anticipated for

a diploid able to display filamentous growth, we observed strain Σ1278b to bud only in a

bipolar or unipolar way. Furthermore, Σ1278b was observed to increase in unipolar

budding, which suggests a transition to filamentous growth (2, 12-15). Our study is the

first to identify the cell density at which the cells switch to a filamentous budding pattern.

We define this cell density as the critical cell density, and when converted to local cell

densities, we find it to be in the range of 4.8×10^6 and 5.3×10^7 cells/ml under low nitrogen

conditions. Local cell densities are a more accurate representation of the cell density

experienced by the cells in our experiments, as all cells were immobilised to the bottom

of the wells.

We sought to obtain the physiological metabolite concentrations at which the

cells switch to filamentous growth. For our growth phases, these were the metabolite

concentrations present between 10 and 20 hours of incubation. We were able to

correlate this physiological concentration of metabolites present with the critical cell

density and identified a total of 29 compounds in our media (Table 1). This is

comparable to other metabolome yeast studies using [1]H NMR which identified 18-39

total compounds (37-41). The actual number of metabolites produced by *S. cerevisiae* is

likely to be in the range of 600-1000 (42, 43). Our observed metabolome may be

somewhat limited by the sensitivity of NMR spectroscopy. It should be noted, however,

that the highest number of identified compounds achieved using currently available

techniques, such as GC-TOF-MS, ranges from 80 (44) to 110 (45), so much more

research needs to be done before having any overview of the full yeast metabolome

and the physiological concentrations present when cells switch to filamentous growth.

Experiments to probe the mechanisms of the observed change in budding

pattern were carried out with the Σ1278b strain under low nitrogen conditions. These

conditions were chosen for comparison with previous studies that have used nitrogen

deficiency to induce filamentous growth (2, 9, 18). Here, to investigate the effects of cell

density and metabolites independently, an overnight culture growing in YPD broth was

directly diluted to an inoculation concentration that corresponded to the critical cell

density (Fig. 7A). In contrast to the time series experiment (Fig. 6A), there was no

previous cell growth and metabolization for 10 hours in the media. Collectively, results

from these experiments suggest that neither the critical cell density alone, nor a

combination of critical cell density and physiological metabolites, could recapitulate the

increase in unipolar budding pattern (Fig. 4A and B). The difference in the growth

environment and metabolism between the different experiments could explain why we

saw no change in budding pattern in either of these conditions. The observed change in

budding pattern seems therefore to be linked to the specific growth environment found

in the time series experiment. Additionally, when comparing metabolite concentrations

under nitrogen rich and deficient conditions we see there are significant differences

(Table 1 and 2). This provides extra support to the conclusion that the measured

metabolites are not triggering the increase in unipolar budding.

Some small molecule alcohols have been associated with induction of

filamentous growth through a quorum sensing mechanism whereby their production is

autoinduced under low nitrogen. The identified alcohols are tyrosol, 2-phenylethanol and

tryptophol (18, 21-23). We were unable to demonstrate a correlation between these

alcohols and the change in budding pattern. Both tyrosol and 2-phenylethanol did not

show a significant increase in concentration until 30 hours under low nitrogen conditions

and had very low concentrations prior to that (Table 1). Since the increase in unipolar

budding occurred between 10 and 20 hours (Fig. 3D and E) we concluded that tyrosol

and 2-phenylethanol could not induce the change in budding pattern. Furthermore,

tryptophol was not identified in our samples, though we specifically looked for it during

the NMR metabolite identification stage. Tryptophol could have been present but at very

low concentrations and non detectable to the ^{1}H NMR. Avbelj et al. (23) found that

tryptophol was only detectable at a conc of less than 5 µM when cells reached approx.

1.5×10^7 cells/ml. Since our cells ended their growth at a lower cell density than this, it is

likely that tryptophol was present but at a non-detectable concentration. Finally, all three

alcohols have relatively low volatility (2-phenylethanol $bp_{760} = 220°C$ (46), tyrosol $bp_{760} =$

$325°C$ (47) and tryptophol $bp_{760} = 360°C$ (48)). Thus, the measured concentrations were

not likely to be inaccurate due to loss during the experiment. The accumulated evidence

in this paper indicates that these alcohols are not responsible for triggering the increase

in unipolar budding. The discrepancy with previous studies may be attributed to

differences in experimental design and execution. For example, our experiment was

carried out in liquid media for 30 hours instead of colonies growing on agar plates for 3 -

5 days (9, 18, 21).

Nitrogen deficiency has commonly been linked to inducing filamentous growth in

S. cerevisiae (2, 18). Contrary to expectations, under both nitrogen rich and deficient

conditions we see a significant increase in unipolar budding between 10 to 20 hours of

growth (Fig. 3B-E, 5B-E) occurring at local cell densities of a similar order of magnitude

(between 5.2×10^6 and 8.2×10^7 cells/ml under high nitrogen and 4.8×10^6 and 5.3×10^7

cells/ml under low nitrogen). Therefore, nutrient deficiency alone appears not to trigger

the switch to unipolar budding and perhaps the switch relies on other factors.

The results presented in this study have important implications in terms of

satisfying the necessary criteria to be classified as a quorum sensing mechanism;

specifically the criteria that filamentous growth is triggered at a critical cell density and is

reproducible at physiological signal molecule concentrations (25). Therefore, our

findings that the critical cell density and physiological metabolite concentration are not

responsible for the shift to a filamentous budding pattern leads to the conclusion that

this phenomenon is not occurring through a quorum sensing mechanism but is being

controlled in an alternative way. While the change in budding pattern observed in the

time series experiment appears to be cell density dependent, it does not however align

with an authentic quorum sensing mechanism based on the previously defined criteria

(25). It is only when the cells could increase in cell density in the same media over time

that this shift to unipolar budding was observed. These findings raise intriguing

questions regarding the mechanism of action controlling this transition to filamentous

growth and highlight the need for further studies into this important issue.

An alternative mechanism could be one involving cell-to-cell-contact that was

found to be a mechanism of growth arrest in yeast (49, 50). Cellular communication

mediated by direct cell-to-cell contact has also been found in bacteria. For example,

direct cell-to-cell contact results in growth inhibition in *Escherichia coli* (51), coordinates

cellular motility in Mycobacterium (52), and mediates auto aggregation and

adhesiveness in *Lactobacillus acidophilus* cell surface (53). Thus, there may be a signal

present on the cell surface, only present there because of metabolism, that signals the

change in budding pattern. Further investigation is therefore required to determine the

mechanism by which the cells increase in unipolar budding since there appears to be

more transpiring than a quorum sensing mechanism.

As mentioned above, 2-phenylethanol has previously been linked to inducing

filamentous growth under low nitrogen conditions (18, 21). In these previous

experiments, the compound was added at concentrations much higher than those

produced by the cells growing in defined media (18, 21, 22). In our experiments, a high

non-physiological concentration of 2-phenylethanol, i.e. 20 µM compared to the

maximum of 4.3 µM measured in our low nitrogen media, induced an increase in

unipolar budding whereas physiological levels of both 2-phenylethanol and other

metabolites did not. This finding confirms the association between 2-phenylethanol and

filamentous growth. However, an implication of the non-physiological concentration

used is that 2-phenylethanol-induced filamentous growth cannot be occurring through a

quorum sensing mechanism as previously suggested (18, 23). Table 3 compares our 2-

phenylethanol findings with the criteria defined by Winters et al. (25) and finds that it

does not satisfy two key points. Therefore, we hypothesise that 2-phenyethanol induces

filamentous growth through a toxicity mechanism due to its high concentration rather

than through an intercellular signalling mechanism.

A recent study by Lenhart et al. (54) also provides evidence contrary to previous

assumptions of quorum sensing mechanisms in *S. cerevisiae.* They found that the

induction of morphogenesis by high concentrations of 2-phenylethanol and tryptophol

(100 µM) is not observed in most environmental isolates and that the filamentous

induction response in Σ1278b was small compared to the extent of that reported by

Chen and Fink (18). This supports our conclusion that quorum sensing controlled

filamentous growth in yeast is not straightforward and that the mechanism of this

biologically critical phenomenon requires more examination.

In conclusion, the methodology presented in this paper uses budding pattern as a

proxy for filamentous growth. This study monitors the budding pattern of yeast cells over

time while simultaneously collecting data on metabolite concentration and cell density.

This allowed us to probe the question of whether filamentous growth is induced though

a quorum sensing mechanism based on more specific criteria. We identified the critical

cell density and physiological metabolite concentration present during the shift to a

filamentous budding pattern in *S. cerevisiae*. However, neither cell density, metabolite

concentration nor nitrogen condition triggered the increase in unipolar budding. An

implication of these findings is that the criteria for filamentous growth to occur via an

intercellular signalling and quorum sensing mechanism were not fulfilled and that the

change in budding pattern does not therefore occur through these mechanisms. Our

results also indicated that 2-phenylethanol induces filamentous growth through another

mechanism, proposed as toxicity, and not through a quorum sensing mechanism as

previously accepted.

The contribution of this study has been to provide clarity of this transition in the

model and industrially useful species, *S. cerevisiae*, highlighting the need for further

investigation to determine the mechanism controlling the shift to filamentous growth

under physiological conditions. It thus challenges the assumptions around quorum

sensing in *S. cerevisiae* widely accepted in literature. The mechanism by which the

switch to filamentous growth occurs in *S. cerevisiae* needs to be elucidated through

further research without resorting to categorisation into quorum sensing mechanisms.

MATERIALS AND METHODS

Yeast Strains and Media.

The *Saccharomyces cerevisiae* strains S288c and Σ1278b were used as a

negative and positive control for filamentation respectively (Fig. 6A). The strains were

maintained using accepted protocols, detailed below.

Yeast Extract Peptone Dextrose (YPD) broth was made by combining 1 g yeast

extract (BD Bacto™, Herlev, Denmark), 2 g bacteriological peptone and 2 g glucose per

100 ml of distilled water. The solution was stirred with heating until dissolved. 2 % YPD

agar plates were made in the same way but with addition of 2g/100ml bacteriological

agar to the YPD broth before autoclaving. Synthetic Low Ammonium Dextrose (SLAD)

broth was made by combining 0.67 g Yeast Nitrogen Base without amino acids and

ammonium sulphate (BD Difco™, Herlev, Denmark), 2 g glucose and 50 µl of

ammonium sulphate solution (1M) per 100 ml of distilled water. Synthetic Ammonium

Dextrose (SAD) liquid broth was made by combining 0.67 g Yeast Nitrogen Base

without amino acids and ammonium sulphate, 2 g glucose and 3.7 ml of ammonium

sulphate solution (1M) per 100 ml of distilled water. All media were autoclaved and

cooled to room temperature and sterile filtered (Q-Max 25 mm 0.22 µm CA) before use.

Concanavalin A (biotin conjugate, Type IV) at 1mg/ml MilliQ water (Con A) was used for

chemical immobilization of cells. Where not indicated otherwise, all chemicals were

purchased from Sigma–Aldrich (Søborg, Denmark).

Image Acquisition.

Images of growing cells in liquid media were obtained using an oCelloScope™

(BioSense Solutions ApS, Farum, Denmark). The oCelloScope™ is a digital time-lapse

microscopy technology that generates very clear images of growing microorganisms.

This is achieved by scanning through a sample to produce a series of images using a

tilted imaging plane. Combining the tilted images give the best focus images and

enables time-lapse images of single cells (34, 35).

Time series determination of budding pattern under low and high nitrogen conditions.

S288c and Σ1278b strains of *S. cerevisiae* (Fig. 6A) were used in the time series experiment which linked budding pattern with cell density and physiological metabolite concentrations. Figure 6B represents a schematic of the overall time series experimental process used in this research. The experiment was carried out in both high (SAD) and low nitrogen (SLAD) liquid media with the same starting cell density of 10^3 cells/ml (Fig. 6C). The overnight culture was made by inoculating 5 ml of YPD broth in a 15 ml centrifuge tube with a single yeast colony grown on a 2% YPD agar plate. This was incubated overnight at 25°C with shaking. After overnight incubation, the cells were counted using a hemocytometer (Neubauer Improved counting chamber, Assistent), washed twice with the relevant media, and diluted to obtain a cell density of 1×10^3 cells/ml in the relevant media. Three 5 ml stock cultures were made in 15 ml centrifuge tubes with lids to produce biological replicates A, B and C for each condition (Fig. 6C). During the experiment these stock cultures grew at 25°C without shaking. Wells of the 12-well microtiter plate (VWR 734-2324 T C-treated 12-well plates) were prepared just before inoculation and imaging by coating the bottom of each well with 15 μl of Con A and allowing the wells to dry. 1 ml of each stock culture was pipetted into wells of the microtiter plate. After inoculation with the cell suspension the plate was placed in the oCelloScope™ and allowed to rest for 30 minutes before starting image acquisition. This was to ensure that the cells were all immobilized on the bottom of the well before imaging to aid focusing. Scanning was set to take place every 10 minutes for 10 hours with one scan area per well at 25°C. Two scan areas per well were used for the final 20-to-30-hour time point to ensure enough single cells were captured at the higher cell

density. After 10 hours of image acquisition, the plate was removed, and the media from

each well was collected in 1.5 ml Eppendorf tubes. The media was spun down in a

centrifuge at 4°C for 10 min at 4400 g (Centrifuge 5920 R, Eppendorf Nordic, Hørsholm,

Denmark) and the supernatant collected and stored at -80° for ^{1}H NMR analysis. Direct

transfer of 1 ml of each stock solution occurred again at 10 and 20 hours to a freshly

Con A coated microtiter plate, as previously described. Images at 10 and 20 hours were

acquired as previously described. Thus, this process resulted in a total of 30 hours of

imaging to give three image acquisitions time periods of 10 hours each. Cell density of

the stock cultures was also measured at 10, 20 and 30 hours with a hemocytometer.

Cell densities were averages of the three biological replicates A, B and C.

Investigating the factors responsible for change in budding pattern

Figure 7A represents a schematic of the overall process for the experiments used

to investigate the effect of various factors of interest on the budding pattern. This

included investigating the effect of critical cell density, conditioned media and high 2-

phenylethanol concentrations (Fig. 7B). All experiments were carried out using a cell

density which corresponded to the critical cell density identified under low nitrogen

conditions in the time series experiment (2×10^4 cells/ml).

Effect of critical cell density alone on budding pattern.

To test the effect of critical cell density on the budding pattern, we analyzed the

effect of using cells at the critical cell density in low nitrogen broth. The critical cell

density condition used SLAD media inoculated with the critical cell density of 2×10^4

cells/ml (Fig. 7B). 5 ml of YPD broth in a 15 ml centrifuge tube was inoculated with a

single yeast colony grown on a 2% YPD agar plate and incubated overnight at 25°C

with shaking. After overnight incubation, the cells were counted using a hemocytometer.

These cells were then washed twice with SLAD media. A cell density of 2×10^4 cells/ml

(critical cell density) was obtained to make three 1 ml solutions (Fig. 7A). This produced

three biological replicates, A, B and C. The microtiter plate was prepared as in the time

series experiment. The three 1 ml solutions were pipetted into the relevant wells before

analysis. Image acquisition in the oCelloScope™ and image analysis to obtain the

budding pattern was carried out as in the time series budding pattern assay.

Combined effect of critical cell density and metabolites on budding pattern.

To test the effect of critical cell density and metabolites on budding pattern, we

inoculated the critical cell density of 2×10^4 cells/ml into conditioned media grown for 20

hours (Fig. 7B). Conditioned media was produced by growing 10^3 cells/ml of freshly

washed cells in SLAD broth for 20 hours followed by centrifugation and supernatant

removal. This conditioned media supernatant was used immediately in the experiment

to wash and inoculate cells, as described in the previous section.

Effect of non-physiological 2-phenylethanol on budding pattern

To investigate the effect of non-physiological concentrations of 2-phenylethanol

on the budding pattern we used SLAD broth containing 20 µM of 2-phenylethanol. This

concentration was previously found to induce filamentation by Chen and Fink (18) under

low nitrogen. Cells were inoculated into the spiked broth at the critical cell density of

2×10^4 cells/ml (Fig. 7B). Cell preparation, inoculation and imaging was carried out as

described in the previous section.

Image analysis to determine budding pattern.

Manual image analysis was carried out on the images acquired from the

oCelloScope™ to determine the budding patterns of individual cells. Firstly, the scan

areas were cropped using a custom-made code in MATLAB (version R202a, The

MathWorks Inc., Natick, MA, USA) such that thirty images were cropped from each scan

area from the three biological replicate wells. Samples of cropped images were

randomly taken from the biological replicates A, B and C. Each cropped image was then

analyzed by eye to determine the bud site sequence of new daughter cells over each

period of image acquisition to obtain a bud site sequence for >200 cells and >300 bud

sites for each condition (Fig. 1). This was done by observing new daughter cells and

recording the positions of the first two or three bud sites (dependent on the number of

times the cell budded within the time frame) to assign each cell with a bud site

sequence. An example of this process on a representative cell can be found in Figure 2

(for full time-lapse see Movie S1). These were then assigned their corresponding

budding pattern classification (axial, bipolar or unipolar) according to Figure 1. From the

budding pattern assignments the percentage value of each budding pattern was

determined for the different experimental conditions.

Cell Density Data Analysis

As mentioned above, the cell density in the 1 ml (V_{sample}) of stock solution was

determined before imaging by counting with a hemocytometer. This measured cell

density value (measured CD) was converted to a local cell density (local CD) that the

cells would experience based on being immobilized on the bottom of the well.

Assuming a yeast cell size of 10 µm, we calculated the volume that the yeasts

would occupy at the bottom of the well. Since the well had a bottom surface area of 3.85

cm^2, the new volume occupied by the yeast at the bottom of the well would be 3.85x10^{-3}

cm^3 (V_{bottom}) if the whole bottom surface was covered with yeast. This value was used to

convert the cell-density per ml value to a local cell density according to the following

formula:

$$Local\ CD = \frac{Measured\ CD \times V_{sample}}{V_{bottom}}$$

NMR Chemicals and Buffer Preparation

Analytical grade sodium phosphate monobasic (NaH2PO3, ≥99.0%), sodium

phosphate dibasic dihydrate (Na_2HPO_3, 2 H_2O, ≥98.0%), sodium azide (NaN_3, ≥99.5%),

deuterium oxide (D_2O, 99.9 atom % D), and 3-trimethyl-silyl-[2,2,3,3-2H4] propionic acid

sodium salt (TSP-d4, 98 atom % D) were purchased from Sigma–Aldrich (Søborg,

Denmark). Water used throughout the study was purified using a Millipore lab water

system (Merck KGaA, Darmstadt, Germany) equipped with a 0.22 mm filter membrane.

NMR Sample Preparation

An aliquot of 700 µL of sample was transferred into a 2 mL Eppendorf tube

containing 300 µL of phosphate buffer (0.15 M, 20% D2O). The mixture was vigorously

vortexed for 30 s. A volume of 600 µL was transferred into 5 mm O.D. NMR SampleJet

tubes (Bruker Biospin, Ettlingen, Germany).

NMR Measurements

Proton ([1]H) NMR spectra were recorded on a Bruker Avance III 600 operating at

a proton's Larmor frequency of 600.13 MHz. The spectrometer was equipped with a 5

mm broadband inverse (BBI) probe and an automated sample changer (SampleJet™,

Bruker Biospin, Ettlingen, Germany). The SampleJet was equipped with a refrigerated

sample storage station with cooling racks (278 K) and heating/drying station (298 K).

Data acquisition and processing were carried out in the TopSpin software (version 3.5,

Bruker, Rheinstetten, Germany). Automation of the overall measurement procedure was

controlled by iconNMR™ (Bruker Biospin, Rheinstetten, Germany). NMR spectra were

measured at 298 K using the standard pulse sequence for pre-saturation of the water

signal (zgcppr pulse program, Bruker nomenclature), a 90° pulse, a sweep width of

9615.385 Hz (16 ppm), and an acquisition time of 3 s. The relaxation delay (d1) was set

to 4 s. The receiver gain (RG) value was determined experimentally for each sample.

NMR data was collected into 32 K data points after 512 scans. Each free induction

decay was apodized by a Lorentzian line-broadening of 0.3 Hz. Automatic phase and

baseline correction were performed in TopSpin.

Processing of the NMR data and metabolites quantification

NMR spectra were imported into the SigMa software (33) where the ^{1}H NMR

signals were referenced to the TSP signal at 0.00 ppm. Signals alignment and

quantification (relative) were performed in SigMa by icoshift and multivariate curve

resolution, respectively (55, 56). The yeast metabolome database (http://www.ymdb.ca/)

and the Chenomx NMR suite (Chenomx Inc. Edmonton, Canada) were used for

metabolite identification.

Metabolite concentrations (absolute) were calculated using the following

equation:

648
$$C(X) = \frac{\text{Area}(X)}{\text{Area}(\text{Ref})} * \frac{H(\text{Ref}) * C(\text{Ref})}{H(X)}$$

Where C(X) is the concentration of the metabolite X, Ref is the reference

compound (TSP), and H indicates the number of protons giving rise to the signal from

metabolite X and the reference compound (TSP). Metabolite concentrations are

averages of the three biological replicates A, B and C.

Due to the short recycle delay (D1) employed for the ^{1}H NMR experiments

(D1=4) - too short for an accurate absolute quantification of TSP (D1≥20) - metabolite

concentrations (absolute) were calculated in two-steps: firstly, the average glucose

concentration in the raw media (known) was used to retrieve an estimate of the TSP

concentration (absolute) in the very same samples. Secondly, the average absolute

concentration of TSP was used to calculate the metabolite concentrations (absolute) in

the remaining samples. Therefore, metabolites concentrations reported in the present

study are to be considered as estimates of absolute concentrations.

Statistical Analysis

Statistical analyses were performed using Minitab® 19.2020.1 (64-bit). To

determine statistically significant differences between conditions of budding pattern

proportions, a 2x2 contingency table using Fisher's Exact Test at a significance level of

p=0.05 was applied. For metabolite concentrations, a minimum of two biological

replicates were averaged to obtain the mean, and standard deviations calculated to

determine the standard error of the mean. For cell densities, the three biological

replicates were averaged to obtain the mean value, and the standard deviations

calculated.

ACKNOWLEDGEMENTS

This research was supported by the Faculty of Veterinary and Agricultural Science at the University of Melbourne and an Australian Government Research Training Program (RTP) Scholarship.

We are grateful to Nadia Devargue, Chu Chu Huang and Sebastian Bech-Terkilsen from the University of Copenhagen for assistance in the laboratory, and Franciscus Winfried J van der Berg from the University of Copenhagen for allowing us use of an oCelloScope™.

We declare that we have no conflict of interest.

Experiment designed by MW, KH and NA; experiment and analysis carried out by MW; NMR data collected by VA; paper written by MW. All authors have read, critically revised, and approved the final version of the manuscript.

REFERENCES

1. Brückner S, Mösch H-U. 2012. Choosing the right lifestyle: adhesion and development in *Saccharomyces cerevisiae*. FEMS Microbiol Rev 36:25-58.

2. Gimeno CJ, Ljungdahl PO, Styles CA, Fink GR. 1992. Unipolar cell divisions in the yeast *S. cerevisiae* lead to filamentous growth: Regulation by starvation and RAS. Cell 68:1077-1090.

3. Cullen PJ, Sprague GF. 2012. The regulation of filamentous growth in yeast. Genetics 190:23-49.

4.   Tronnolone H, Gardner JM, Sundstrom JF, Jiranek V, Oliver SG, Binder BJ.

2017. Quantifying the dominant growth mechanisms of dimorphic yeast using a

lattice-based model. J R Soc Interface 14:20170314.

5.   Kron SJ, Styles CA, Fink GR. 1994. Symmetric cell division in pseudohyphae of

the yeast *Saccharomyces cerevisiae*. Mol Biol Cell 5:1003-1022.

6.   Roberts RL, Fink GR. 1994. Elements of a single MAP kinase cascade in

Saccharomyces cerevisiae mediate two developmental programs in the same

cell type: mating and invasive growth. Genes Dev 8:2974-2985.

7.   Dickinson J. 2008. Filament formation in *Saccharomyces cerevisiae* — a review.

Folia Microbiol 53:3-14.

8.   Lengeler KB, Davidson RC, D'souza C, Harashima T, Shen W-C, Wang P, Pan

X, Waugh M, Heitman J. 2000. Signal transduction cascades regulating fungal

development and virulence. Microbiol Mol Biol Rev 64:746-785.

9.   Lorenz MC, Cutler NS, Heitman J. 2000. Characterization of alcohol-induced

filamentous growth in *Saccharomyces cerevisiae*. Mol Biol Cell 11:183-199.

10.   Cullen PJ, Sprague GF. 2000. Glucose depletion causes haploid invasive growth

in yeast. Proc Natl Acad Sci U S A 97:13619-13624.

11.   Cullen PJ, Sprague Jr GF. 2002. The roles of bud-site-selection proteins during

haploid invasive growth in yeast. Mol Biol Cell 13:2990-3004.

12.   Chant J, Pringle JR. 1995. Patterns of bud-site selection in the yeast

Saccharomyces cerevisiae. J Cell Biol 129:751-765.

13.   Freifelder D. 1960. Bud position in *Saccharomyces cerevisiae*. Journal of

bacteriology 80:567.

14.   Gimeno CJ, Fink GR. 1992. The logic of cell division in the life cycle of yeast.

Science 257:626.

15.   Madden K, Costigan C, Snyder M. 1992. Cell polarity and morphogenesis in

Saccharomyces cerevisiae. Trends Cell Biol 2:22-29.

16.   Cali BM, Doyle TC, Botstein D, Fink GR. 1998. Multiple functions for actin during

filamentous growth of *Saccharomyces cerevisiae*. Mol Biol Cell 9:1873-1889.

17.   Karunanithi S, Joshi J, Chavel C, Birkaya B, Grell L, Cullen PJ. 2012. Regulation

of mat responses by a differentiation MAPK pathway in *Saccharomyces*

cerevisiae. PLoS One 7:e32294.

18.   Chen H, Fink GR. 2006. Feedback control of morphogenesis in fungi by aromatic

alcohols. Genes Dev 20:1150-1161.

19.   Gancedo JM. 2001. Control of pseudohyphae formation in *Saccharomyces*

cerevisiae. FEMS Microbiol Rev 25:107-123.

20.   Lambrechts MG, Bauer FF, Marmur J, Pretorius IS. 1996. Muc1, a mucin-like

protein that is regulated by Mss10, is critical for pseudohyphal differentiation in

yeast. Proc Natl Acad Sci U S A 93:8419-8424.

21.   González B, Vázquez J, Cullen PJ, Mas A, Beltran G, Torija M-J. 2018. Aromatic

Amino Acid-Derived Compounds Induce Morphological Changes and Modulate

the Cell Growth of Wine Yeast Species. Front Microbiol 9:1-16.

22.   González B, Vázquez J, Morcillo-Parra MÁ, Mas A, Torija MJ, Beltran G. 2018.

The production of aromatic alcohols in non-*Saccharomyces* wine yeast is

modulated by nutrient availability. Food Microbiol 74:64-74.

23.   Avbelj M, Zupan J, Kranjc L, Raspor P. 2015. Quorum-Sensing Kinetics in

Saccharomyces cerevisiae: A Symphony of ARO Genes and Aromatic Alcohols.

J Agric Food Chem 63:8544-8550.

24.   Monds RD, O'Toole GA. 2008. Metabolites as intercellular signals for regulation

of community-level traits, p 105-129. *In* Winans S, Bassler B (ed), Chemical

Communication Among Bacteria. American Society of Microbiology, Washington,

DC.

25.   Winters M, Arneborg N, Appels R, Howell K. 2019. Can community-based

signalling behaviour in *Saccharomyces cerevisiae* be called quorum sensing? A

critical review of the literature. FEMS Yeast Res 19.

26.   Miller M, Bassler B. 2001. Quorum Sensing in Bacteria. Annu Rev Microbiol

55:165-199.

27.   Henke JM, Bassler BL. 2004. Bacterial social engagements. Trends Cell Biol

14:648-656.

28.   Fuqua C, Winans SC, Greenberg EP. 1996. Census and Consensus in Bacterial

Ecosystems: The LuxR-LuxI Family of Quorum-Sensing Transcriptional

Regulators. Annu Rev Microbiol 50:727-751.

29.   Kleerebezem M, Quadri LEN, Kuipers OP, De Vos WM. 1997. Quorum sensing

by peptide pheromones and two-component signal-transduction systems in

Gram-positive bacteria. Mol Microbiol 24:895-904.

30.   West SA, Winzer K, Gardner A, Diggle SP. 2012. Quorum sensing and the

confusion about diffusion. Trends Microbiol 20:586-594.

31.   Wuster A, Babu MM. 2008. Chemical Molecules that Regulate Transcription and

Facilitate Cell-to-Cell Communication, p 1-8. *In* Begley TP (ed), Wiley

Encyclopedia of Chemical Biology. John Wiley & Sons, New Jersey.

32.   Dickinson J. 1996. Fusel alcohols induce hyphal-like extensions and

pseudohyphal formation in yeasts. Microbiology 142:1391-1397.

33.   Khakimov B, Mobaraki N, Trimigno A, Aru V, Engelsen SB. 2020. Signature

Mapping (SigMa): An efficient approach for processing complex human urine 1H

NMR metabolomics data. Anal Chim Acta 1108:142-151.

34.   Fredborg M, Andersen KR, Jørgensen E, Droce A, Olesen T, Jensen BB,

Rosenvinge FS, Sondergaard TE. 2013. Real-Time Optical Antimicrobial

Susceptibility Testing. J Clin Microbiol 51:2047-2503.

35.   Fredborg M, Rosenvinge FS, Spillum E, Kroghsbo S, Wang M, Sondergaard TE.

2015. Automated image analysis for quantification of filamentous bacteria. BMC

Microbiol 15:255.

36.   Liu H, Styles CA, Fink GR. 1996. *Saccharomyces cerevisiae* S288C Has a

Mutation in FL08, a Gene Required for Filamentous Growth. Genetics 144:967-

978.

37.   Puig-Castellví F, Alfonso I, Piña B, Tauler R. 2015. A quantitative 1 H NMR

approach for evaluating the metabolic response of *Saccharomyces cerevisiae* to

mild heat stress. Metabolomics 11:1612-1625.

38.   Nilsson M, Duarte IF, Almeida C, Delgadillo I, Goodfellow BJ, Gil AM, Morris GA.

2004. High-resolution NMR and diffusion-ordered spectroscopy of port wine. J

Agric Food Chem 52:3736-3743.

39.   Peng C, Viana T, Petersen MA, Larsen FH, Arneborg N. 2018. Metabolic

footprint analysis of metabolites that discriminate single and mixed yeast cultures

at two key time-points during mixed culture alcoholic fermentations.

Metabolomics 14:93.

40.   Kumar Babele P. 2019. Zinc oxide nanoparticles impose metabolic toxicity by de-

regulating proteome and metabolome in Saccharomyces cerevisiae. Toxicol Rep

6:64-73.

41.   Lourenço AB, Roque FC, Teixeira MC, Ascenso JR, Sá-Correia I. 2013.

Quantitative 1H-NMR-Metabolomics Reveals Extensive Metabolic

Reprogramming and the Effect of the Aquaglyceroporin FPS1 in Ethanol-

Stressed Yeast Cells. PLoS One 8:e55439.

42.   Raamsdonk LM, Teusink B, Broadhurst D, Zhang N, Hayes A, Walsh MC,

Berden JA, Brindle KM, Kell DB, Rowland JJ. 2001. A functional genomics

strategy that uses metabolome data to reveal the phenotype of silent mutations.

Nat Biotechnol 19:45-50.

43.   Mo ML, Palsson BØ, Herrgård MJ. 2009. Connecting extracellular metabolomic

measurements to intracellular flux states in yeast. BMC Syst Biol 3:37.

44.   Ding M-Z, Zhou X, Yuan Y-J. 2010. Metabolome profiling reveals adaptive

evolution of *Saccharomyces cerevisiae* during repeated vacuum fermentations.

Metabolomics 6:42-55.

45.   Kim S, Lee DY, Wohlgemuth G, Park HS, Fiehn O, Kim KH. 2013. Evaluation

and optimization of metabolome sample preparation methods for *Saccharomyces*

cerevisiae. Anal Chem 85:2169-2176.

46.   Hua D, Lin S, Li Y, Chen H, Zhang Z, Du Y, Zhang X, Xu P. 2010. Enhanced 2-

phenylethanol production from L-phenylalanine via in situ product adsorption.

Biocatal Biotransformation 28:259-266.

47.   Napolitano A, De Lucia M, Panzella L, d'Ischia M. 2010. The Chemistry of

Tyrosol and Hydroxytyrosol: Implications for Oxidative Stress, p 1225-1232. *In*

Preedy VR, Watson RR (ed), Olives and Olive Oil in Health and Disease

Prevention. Academic Press, San Diego.

48.   Lide DR. 1995. CRC Handbook of Chemistry and Physics, 75th ed. CRC Press,

Boca Raton.

49.   Nissen P, Arneborg N. 2003. Characterization of early deaths of non-

Saccharomyces yeasts in mixed cultures with Saccharomyces cerevisiae. Arch

Microbiol 180:257-263.

50.   Nissen P, Nielsen D, Arneborg N. 2003. Viable *Saccharomyces cerevisiae* cells

at high concentrations cause early growth arrest of non-Saccharomyces yeasts in

mixed cultures by a cell–cell contact-mediated mechanism. Yeast 20:331-341.

51.   Aoki SK, Pamma R, Hernday AD, Bickham JE, Braaten BA, Low DA. 2005.

Contact-dependent inhibition of growth in *Escherichia coli*. Science 309:1245-

1248.

52.   Kaiser D. 2004. Signaling in myxobacteria. Annu Rev Microbiol 58:75-98.

53.   Kos B, Šušković J, Vuković S, Šimpraga M, Frece J, Matošić S. 2003. Adhesion

and aggregation ability of probiotic strain *Lactobacillus acidophilus* M92. J Appl

Microbiol 94:981-987.

54.   Lenhart BA, Meeks B, Murphy HA. 2019. Variation in filamentous growth and

response to quorum-sensing compounds in environmental isolates of

Saccharomyces cerevisiae. G3: Genes, Genomes, Genetics 9:1533-1544.

55.   Savorani F, Tomasi G, Engelsen SB. 2010. icoshift: A versatile tool for the rapid

alignment of 1D NMR spectra. J Magn Reson Open 202:190-202.

56.   De Juan A, Jaumot J, Tauler R. 2014. Multivariate Curve Resolution (MCR).

Solving the mixture analysis problem. Anal Methods 6:4964-4976.

TABLES & FIGURES

FIG 1 Schematic diagram of how the budding pattern is obtained from the bud site sequence of a new daughter cell. Bud site sequence is assigned for new daughters with both two and three bud sites and categorised to its corresponding budding pattern as shown. For cells growing in non-filamentous form, haploids will adopt an axial pattern (red). This pattern is defined as new buds emerging consistently adjacent to the junction to the mother. Diploids will adopt a bipolar pattern (orange) whereby buds may emerge either adjacent or opposite the junction to the mother. When filamentous growth occurs, both cell types transition to a unipolar budding pattern (blue). This pattern is defined as new buds emerging consistently from the end opposite the junction to the mother. In this way, budding pattern can be used as a proxy for tracking filamentous growth in liquid media (2, 5, 12-15).

863
FIG 2 Representative cropped time-lapse images of a new daughter cell being assigned
a budding pattern by using its bud site sequence. Time-lapse images were obtained
with the oCelloScope™. For the 1st daughter (green), the first bud (orange) emerges
opposite the birth end (O), followed by a second bud (orange) also opposite its birth end
(OO) and finally the third bud (orange) emerges adjacent to the birth end (OOA). Thus,
this daughter would have a bud site sequence of OOA which categorizes it as bipolar
budding pattern according to Fig 1.

901
FIG 3 *S. cerevisiae* strain Σ1278b shows an increase in the filamentous unipolar
budding pattern under low nitrogen conditions between local cell densities of 4.8x106
and 5.3x107 cells/ml. (A) Change in local cell density under low nitrogen conditions
measured by cell counting with a hemocytometer. Means and standard deviation for
three biological replicates are shown. (B) Budding pattern for S288cs cells with two bud
sites. (C) Budding pattern for S288cs cells with three bud sites. (D) Budding pattern for
Σ1278b cells with two bud sites. (E) Budding pattern for Σ1278b cells with three bud
sites. The percentage of total cells exhibiting each budding pattern are shown. Budding
pattern of cells during growth were determined by manual image analysis of time-lapse
images obtained using the oCelloScope™. Different symbols indicate a P value of ≤0.05
as determined by Fishers Exact test.

TABLE 1: Heatmap of the physiological concentrations of compounds detected under
low nitrogen conditions during growth of *S. cerevisiae* strain Σ1278b.[a]

Low Concentration Metabolites (µM)				
Metabolite Name	0 hrs	10 hrs	20 hrs	30 hrs
1-Propanol	0.8 ± 0.1	1.4 ± 0.3	3.7 ± 0.3	6.7 ± 0.5
2-Hydroxyisovalerate	0.6 ± 0.2	0.7 ± 0.2	1.3 ± 0.8	2.9 ± 0.2
2-Oxoglutarate	3.4 ± 1.2	4.4 ± 0.2	9.3 ± 2.5	61.7 ± 1.3
2-Phenylethanol	0.5 ± 0.1	0.7 ± 0.2	1.5 ± 0.8	4.3 ± 0.4
3-Methylxanthine	5.9 ± 0.8	3.9 ± 0.1	4.2 ± 0.4	3.7 ± 0.2
Acetaldehyde	0.4 ± 0.1	2.3 ± 0.3	19.2 ± 0.6	30.0 ± 3.8
Acetoin	0.4 ± 0.1	6.7 ± 4.7	8.0 ± 2.1	20.1 ± 1.3
Choline	2.3 ± 0.2	2.9 ± 0.9	2.0 ± 0.8	1.6 ± 0.4
Diacetyl	5.0 ± 0.4	5.4 ± 0.4	4.8 ± 0.8	5.0 ± 0.1
Fumarate	0.6 ± 0.1	0.5 ± 0.2	1.1 ± 0.5	14.6 ± 1.4
Hydroxyacetone	3.6 ± 1.6	4.2 ± 1.7	2.4 ± 0.2	3.3 ± 0.1
Isoamylol	0.6 ± 0.2	0.6 ± 0.4	1.0 ± 0.7	11.1 ± 1.1
Isopropanol	0.3 ± 0.1	0.4 ± 0.1	2.1 ± 0.2	4.8 ± 0.7
Malate	2.4 ± 0.6	2.9 ± 0.4	6.8 ± 3.7	10.2 ± 0.7
Methionine	0.2 ± 0.1	0.2 ± 0.1	0.6 ± 0.1	1.0 ± 0.1
Myoinositol	47.3 ± 1.0	43.6 ± 7.3	33.1 ± 13.0	41.4 ± 0.7
Niacin	9.5 ± 0.2	7.8 ± 1.6	7.6 ± 2.9	9.6 ± 0.2
Pantothenate	4.3 ± 0.2	4.8 ± 0.4	5.3 ± 0.6	6.0 ± 0.1
Pyridoxine	7.8 ± 0.2	8.3 ± 0.3	7.8 ± 0.3	7.4 ± 0.1
Pyruvate	4.9 ± 0.9	5.0 ± 0.1	30.6 ± 1.5	52.8 ± 1.9
Succinate	1.8 ± 0.2	2.5 ± 0.1	12.0 ± 7.3	45.0 ± 0.2
Trehalose	91.3 ± 0.5	88.1 ± 8.3	66.9 ± 26.6	85.9 ± 1.3
Tyrosol	0.2 ± 0.1	0.3 ± 0.1	0.8 ± 0.2	4.9 ± 0.2
Xanthine	0.9 ± 0.1	1.3 ± 0.3	0.6 ± 0.1	6.6 ± 0.1

0 µM

92 µM

High Concentration Metabolites (µM)				
Metabolite Name	0 hrs	10 hrs	20 hrs	30 hrs
Acetate	9 ± 1	17 ± 1	160 ± 6	411 ± 12
Ethanol	189 ± 40	369 ± 8	5403 ± 160	14334 ± 1023
Formate	221 ± 2	238 ± 2	231 ± 9	193 ± 4
Glucose (x10³)	107 ± 1	103 ± 1	115 ± 5	99 ± 1
Lactate	8 ± 2	29 ± 1	224 ± 8	343 ± 38

0 µM

115 mM

[a]Metabolites identified through [1]H-NMR and quantified using the SigMa programme (Khakimov et al. 2020). Values are expressed as mean ± SE for two or three biological replicates.

917
FIG 4 High-concentration 2-phenylethanol recapitulates the increase in unipolar
budding, while cells at the critical cell density and growing in conditioned media do not.
Experiments were carried out using *S. cerevisiae* strain Σ1278b under low nitrogen
condition at the critical cell density to investigate the effect of cell density alone (CCD),
the combination of cell density and signalling metabolites (CM), and non-physiological
2-phenylethanol (2PE) on the budding pattern. (A) Budding pattern for Σ1278b cells with
two bud sites. (B) Budding pattern for Σ1278b cells with three bud sites. The percentage
of total cells exhibiting each budding pattern are shown. Different symbols indicate a P
value of ≤0.05 as determined by Fishers Exact test.

951
FIG 5 *S. cerevisiae* strains growing under high nitrogen conditions display the same
budding pattern behaviour as low nitrogen conditions, with an increase in unipolar
budding occurring between local cell densities of 5.2×10^6 and 8.2×10^7 cells/ml. (A)
Change in local cell density under high nitrogen conditions measured by cell counting
with a hemocytometer. Means and standard deviation for three biological replicates are
shown. (B) Budding pattern for S288cs cells with two bud sites. (C) Budding pattern for
S288cs cells with three bud sites. (D) Budding pattern for Σ1278b cells with two bud
sites. (E) Budding pattern for Σ1278b cells with three bud sites. The percentage of total
cells exhibiting each budding pattern are shown. Budding pattern of cells during growth
were determined by manual image analysis of time-lapse images obtained using the
oCelloScope™. Different symbols indicate a P value of ≤0.05 as determined by Fishers
Exact test.

TABLE 2: Heatmap of the physiological concentrations of compounds detected under
high nitrogen conditions during growth of *S. cerevisiae* strain Σ1278b.[a]

Low Concentration Metabolites (µM)			
Metabolite Name	**0 hrs**	**10 hrs**	**20 hrs**	**30 hrs**
1-Propanol	1.3 ± 0.1	1.4 ± 0.1	3.4 ± 0.4	21.4 ± 10.5
2-Hydroxyisovalerate	0.3 ± 0.1	0.4 ± 0.1	0.5 ± 0.2	1.8 ± 0.3
2-Oxoglutarate	0.7 ± 0.2	1.5 ± 0.1	2.0 ± 0.1	13.6 ± 3.2
2-Phenylethanol	0.4 ± 0.1	0.4 ± 0.1	0.6 ± 0.1	0.9 ± 0.4
3-Methylxanthine	6.9 ± 0.3	7.8 ± 0.1	8.3 ± 0.4	6.7 ± 1.1
Acetaldehyde	1.5 ± 0.4	2.0 ± 0.6	18.0 ± 5.3	18.8 ± 8.8
Acetoin	1.2 ± 0.1	1.0 ± 0.1	5.6 ± 1.9	25.3 ± 15.4
Choline	2.2 ± 0.6	3.9 ± 1.0	3.7 ± 1.0	2.1 ± 0.8
Diacetyl	4.5 ± 0.1	4.5 ± 0.1	4.5 ± 0.2	4.2 ± 0.1
Fumarate	0.2 ± 0.1	0.4 ± 0.1	0.3 ± 0.1	4.8 ± 0.6
Hydroxyacetone	8.1 ± 0.1	7.9 ± 0.2	8.1 ± 0.3	8.4 ± 0.2
Isoamylol	0.8 ± 0.1	0.7 ± 0.1	0.9 ± 0.1	13.0 ± 6.6
Isopropanol	0.3 ± 0.1	0.3 ± 0.1	2.9 ± 0.8	2.5 ± 0.8
Malate	3.0 ± 0.3	2.1 ± 0.5	2.2 ± 0.2	6.2 ± 1.1
Methionine	0.2 ± 0.1	0.3 ± 0.1	0.5 ± 0.1	6.8 ± 3.8
Myoinositol	51.3 ± 0.9	50.3 ± 0.7	48.2 ± 1.6	43.7 ± 4.2
Niacin	9.4 ± 0.3	9.4 ± 0.3	11.6 ± 0.9	6.5 ± 1.7
Pantothenate	4.9 ± 0.1	5.3 ± 0.3	5.3 ± 0.2	5.5 ± 0.2
Pyridoxine	8.9 ± 0.2	9.0 ± 0.3	9.6 ± 0.2	10.5 ± 1.1
Pyruvate	8.5 ± 0.8	11.5 ± 0.5	38.8 ± 6.3	92.2 ± 48.2
Succinate	1.4 ± 0.2	0.9 ± 0.1	1.6 ± 0.1	18.5 ± 10.9
Trehalose	95.3 ± 1.5	94.4 ± 2.3	92.0 ± 1.1	64.1 ± 18.6
Tyrosol	0.4 ± 0.1	0.6 ± 0.1	0.3 ± 0.1	0.4 ± 0.1
Xanthine	1.1 ± 0.2	1.3 ± 0.1	1.2 ± 0.4	0.9 ± 0.2

0 µM

96 µM

High Concentration Metabolites (µM)			
Metabolite Name	**0 hrs**	**10 hrs**	**20 hrs**	**30 hrs**
Acetate	9 ± 1	17 ± 1	160 ± 6	411 ± 12
Ethanol	189 ± 40	369 ± 8	5403 ± 160	14334 ± 1023
Formate	221 ± 2	238 ± 2	231 ± 9	193 ± 4
Glucose (x10^3)	107 ± 1	103 ± 1	115 ± 5	99 ± 1
Lactate	8 ± 2	29 ± 1	224 ± 8	343 ± 38

0 µM

115 mM

[a]Metabolites identified through ^{1}H-NMR and quantified using the SigMa programme (Khakimov et al. 2020). Values are expressed as mean ± SE for two or three biological replicates.

968 **TABLE 3** 2-phenylethanol does not act as an intercellular signaling and quorum sensing
969 molecule to induce the change in budding pattern based on our experimental results.[a]

	Criteria	2-phenylethanol
Intercellular Signalling Molecule	Extracellular & Identified	Y
	Specific signalling mechanism	Y
	Reproducible response	N[b]
	Adaptive at community level	Y
Quorum Sensing Molecule	Proportional to cell density	Y
	Critical threshold trigger	N[c]

971
972 [a] Criteria from (23).
973 [b] Response only observed when 2-phenylethanol used at much higher concentrations
974 (20 µM) than found to be physiologically produced (0.7 – 1.5 µM)
975 [c] Since the response is only induced at high levels it may be a starvation or stress
976 response and not a QS mechanism

(A)

Strain Name	Source	Budding Pattern	Filamentation	Reference
S288c	Lab strain	Axial	-	ATCC® 26108™
Σ1278b	NCYC 1391	Bipolar	+	(Grenson, Hou & Crabeel 1970)

(B)

(C)

Media	Strain	Starting Cell Density (cells/ml)
Low Nitrogen (SLAD)	S288c	
	Σ1278b	
High Nitrogen (SAD)	S288c	1×10^3
	Σ1278b	

1000
FIG 6 The budding patterns of *S. cerevisiae* cells growing in low and high nitrogen liquid
media were tracked during growth and correlated to cell density and metabolite
concentration. (A) *S. cerevisiae* strains used in this study. (B) Schematic of the time
series experiment that utilised the oCelloScope™ for image acquisition and ¹H NMR for
metabolite identification and quantification. (C) Experimental conditions used in the time
series experiment.
1007

(A)

(B)

Label	Description	Strains	Media	Starting Cell Density (cells/ml)
CCD	Critical Cell Density		SLAD	
CM	20 hours CM	Σ1278b	20 hours SLAD conditioned media	2×10^4
2-PE	Non-physiological 2-phenylethanol		SLAD + 20 µM 2-phenylethanol	

1008
1009   **FIG 7** The factors responsible for change in budding pattern were investigated under
1010   low nitrogen conditions using strain Σ1278b. (A) Schematic diagram of experimental set-
1011   up. (B) Experimental conditions used included the critical cell density (CCD), 20 hours
1012   conditioned media (CM), and media with a high concentration of 2-phenylethanol (2PE).
1013   All conditions inoculated cells at the critical cell density 2×10^4 cells/ml.
1014

1015

1016

1017

1018

1019

1020

SUPPLEMENTAL MATERIAL

MOVIE S1 Full representative cropped image time-lapse with bud site sequence assignment and budding pattern categorisation from Figure 2.

TABLE S1 Results from Nitrogen Concentration Assay which measured the nitrogen concentration from the media and the supernatant from the three biological replicates A, B and C at the end of the time series experiment i.e. after 30 hours of growth.

Nitrogen Condition	High Nitrogen			Low Nitrogen		
Condition	Media	Σ1278b	S288c	Media	Σ1278b	S288c
Nitrogen Conc (mM)	82.9	80.0	85.0	1.06	0.04	0.03

1040　**TABLE S2** [1]H-NMR metabolite shifts and intervals with J-coupling. Shifts in bold were
1041　used for quantification.
1042

Metabolite Name	Shift (ppm)	Multiplicity	J-coupling (Hz)	Notes
1-propanol	0.88	t	7.5	
	1.53	m	7.1	
	3.55	t	(6.6)	
2-hydroxyisovalerate	0.83	d	6.86	
	0.95	d	6.94	
	2.01	m	(13.81,6.88)	
	3.84	d	(3.75)	
2-oxoglutarate	2.44	t	6.83	
	3.01	t	6.82	
2-phenylethanol	**2.85**	t	6.7	
	3.83	t	-	
	7.30	m	-	
	7.37	t	7.5	
3-methylxanthine	3.51	s		
	8.02	s		
Acetaldehyde	2.25	d	3.0	
	9.68	q	3.0	
Acetate	**1.92**	s		
Acetoin	1.38	d	7.1	
	2.22	s		
	4.41	q	7.1	
Arabinose	3.51	dd	(9.7,7.8)	NQ
	3.68	m		
	3.83	dd	(9.7, 3.5)	
	3.90	m		
	3.95	m		
	4.05	m		
	4.50	d	7.7	
	5.25	d	(3.57)	
Choline	**3.18**	s		
	3.50	dd	(5.8,4.1)	
	4.05	ddd		
Diacetyl	**2.31**	s		
Ethanol	**1.18**	t	7.0	
	3.66	q	7.0	
Formate	**8.45**	s		

Fructose	3.58 3.69 3.82 3.90 4.00 4.03 4.12	m m m dd m dd m	(9.9,3.4) 12.8, 1.0	NQ
Fumarate	**6.51**	s		
α-glucose	3.81 3.82 3.38 3.70 3.53 5.24	m m m m m d	 3.8	Media nutrient
β-glucose	3.89 3.72 3.38 3.45 3.23 **4.63**	m m m m dd d	 (7.8, 9.2) 8.0	Media nutrient
Hydroxyacetone	**2.16** 3.27 4.26	s s d		
Isoamylol	0.86 **1.44** 1.66 3.5 4.3	d q m t s	6.9 7.0	
Isopropanol	**1.16** 4.01	d m	6.0	
Lactic acid	**1.31** 4.10	d q	7.0 7.0	
Malic acid	2.34 **2.66** 4.29	dd dd dd	(15.37,10.2) 15.4, 2.9 (10.23,2.9)	
Myo-inositiol	3.26 3.52 3.61 **4.06**	t dd t t	(9.3) 9.7 2.8	Media nutrient
Niacin	7.53 8.26 8.61 **8.94**	t d d s	(7.94, 4.90) (4.98, 1.67)	Media nutrient

Pantothenate	0.88 **0.92** 2.40 3.37 3.4 3.48 3.51 3.98	s s t s m s s s		Media nutrient
Pyridoxine	2.47 4.73 7.69	s s s		Media nutrient
Pyruvate	2.38	s		
Succinate	2.41	s		
Taurine	3.25 3.42	t t	6.1 (6.1)	NQ
Trehalose	3.42 3.64 3.75 3.82 5.11	t dd m m d	(9.47) (9.93,3.8) 3.8	
Tyrosol	2.77 3.77 6.86 7.17	t t d d	6.7 - 8.4 8.5	

1043

FIG S1 Representative ¹H-NMR spectra of samples. The major resonances used for
quantification have been assigned: 1. TSP; 2. Pantothenate; 3. 2-hydroxyisovalerate; 4.
Isopropanol; 5. Ethanol; 6. Lactate; 7. Isoamylol; 8. 1-Propanol; 9. Acetate; 10.
Hydroxyacetone; 11. Acetoin; 12. Diacetyl; 13. Pyruvate; 14. Succinate; 15. Pyridoxine;
16. Malate; 17. 2-Phenylethanol; 18. 2-oxoglutarate; 19. Choline; 20. Myoinositol; 21. β-
Glucose; 22. Trehalose; 23. α-Glucose; 24. Fumarate; 25. Tyrosol; 26. Xanthine; 27. 3-
Methylxanthine; 28. Formate; 29. Niacin; 30. Acetaldehyde.

1051
1052
1053

1054 **FIG S2** Representative ^{1}H-NMR spectra showing the results of the spiking experiment
1055 conducted by adding (A) 10 µL 2-phenylethanol and (B) 5 µL tyrosol to the samples.