(–)-Epicatechin alters reactive oxygen and nitrogen species production

independent of mitochondrial respiration in human vascular endothelial

cells

Daniel G. Sadler[1]; Jonathan Barlow[2]; Richard Draijer[3]; Helen Jones[1]; Dick H. J. Thijssen[1];

Claire E. Stewart[1].

[1]School of Sport and Exercise Science, Liverpool John Moores University, Liverpool, United

Kingdom

[2]Mitochondrial Profiling Centre, School of Sport, Exercise and Rehabilitation Sciences,

University of Birmingham, Birmingham, United Kingdom

[3]Unilever Foods Innovation Centre, Bronland 14, 6708 WH Wageningen, The Netherlands

Corresponding Author: Claire E. Stewart

Address Correspondence: Research Institute for Sport and Exercise Sciences, Liverpool

John Moores University, Byrom Street, L3 3AF, Liverpool, UK

Telephone number: +44 (0)151 904 6234

E-mail address: C.E.Stewart@ljmu.ac.uk

Author email addresses:

Daniel G. Sadler: Dan_sadler94@hotmail.co.uk

Jonathan Barlow: J.P.Barlow@bham.ac.uk

Richard Draijer: Richard.Draijer@unilever.com

Helen Jones: H.Jones1@ljmu.ac.uk

Dick H. J. Thijssen: D.Thijssen@ljmu.ac.uk

Running head: (–)-Epicatechin alters reactive oxygen and nitrogen species production

Abstract

Introduction: Vascular endothelial dysfunction is characterised by lowered nitric oxide (NO)

bioavailability, which may be explained by increased production of reactive oxygen species

(ROS), mitochondrial dysfunction and altered cell signalling. (-)-Epicatechin (EPI) has proven

effective in the context of vascular endothelial dysfunction, but the underlying mechanisms

associated with EPI's effects remain unclear.

Objective(s): Our aim was to investigate whether EPI impacts reactive oxygen and nitrogen

species (RONS) production and mitochondrial function of human vascular endothelial cells

(HUVECs). We hypothesised that EPI would attenuate ROS production, increase NO

bioavailability, and enhance indices of mitochondrial function.

Methods: HUVECs were treated with EPI (0-20 µM) for up to 48 h. Mitochondrial and cellular

ROS were measured in the absence and presence of antimycin A (AA), an inhibitor of the

mitochondrial electron transport protein complex III, favouring ROS production. Genes

associated with mitochondrial remodelling and the antioxidant response were quantified by

RT-qPCR. Mitochondrial bioenergetics were assessed by respirometry and signalling

responses determined by western blotting.

Results: Mitochondrial superoxide production without AA was increased 32% and decreased

53% after 5 and 10 µM EPI treatment vs. CTRL ($P<0.001$). With AA, only 10 µM EPI

increased mitochondrial superoxide production vs. CTRL (25%, $P<0.001$). NO bioavailability

was increased by 45% with 10 µM EPI vs. CTRL ($P=0.010$). However, EPI did not impact

mitochondrial respiration. NRF2 mRNA expression was increased 1.5- and 1.6-fold with 5 and

10 µM EPI over 48 h vs. CTRL ($P=0.015$ and $P=0.001$, respectively). Finally, EPI transiently

enhanced ERK1/2 phosphorylation (2.9 and 3.2-fold over 15 min and 1 h vs. 0 h, respectively;

$P=0.035$ and $P=0.011$).

Conclusion(s): EPI dose dependently alters RONS production of HUVECs but does not impact

mitochondrial respiration. The induction of NRF2 mRNA expression with EPI might relate to

enhanced ERK1/2 signalling, rather than RONS production. In humans, EPI may improve

vascular endothelial dysfunction via alteration of RONS and activation of cell signalling.

Introduction

Globally, cardiovascular disease (CVD) is the leading cause of morbidity and mortality (WHO, 2018). One major risk factor for CVD is vascular endothelial dysfunction, which is typified by impaired vasodilation and diminished blood flow (Heitzer et al., 2001; Suwaidi et al., 2000). Several factors contribute to vascular endothelial dysfunction, including reduced nitric oxide (NO) bioavailability and elevated oxidative stress (Ungvari et al., 2018). A decline in NO bioavailability has been attributed to lower endothelial nitric oxide synthase (eNOS) content and activity, in part due to lower phosphorylation of eNOS at Ser1177 - at least in the aged vascular endothelium (Gliemann et al., 2018; Musicki et al., 2005). Elevated oxidative stress might be explained increased production of ROS (Cernadas et al., 1998; Chou et al., 1998; Donato et al., 2007, 2009), that are cytosolic and mitochondrial in origin (Csiszar et al., 2002, 2007; Donato et al., 2007; Hamilton et al., 2001; Jablonski et al., 2007; Sun et al., 2004; Van Der Loo et al., 2000).

Vascular endothelial health is also impacted by mitochondrial function. Indeed, ageing is associated with reduced mitochondrial content in endothelial cells of conduit arteries, feed arteries and capillaries (Burns et al., 1979; S. H. Park et al., 2018; S.-Y. Park et al., 2018; Ungvari et al., 2008), which could be due to blunted transcriptional responses (S.-Y. Park et al., 2018; Ungvari et al., 2008). This potential reduction in mitochondrial biogenesis with ageing may result from diminished NO bioavailability (Gouill et al., 2007; Miller et al., 2013) and/or lowered AMP-activated protein kinase (AMPK) signalling (Lesniewski et al., 2012), both of which are known to activate peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1α). A link between mitochondria and vascular endothelial dysfunction has also been made from observations that human skeletal muscle feed arteries exhibit impaired respiratory capacity and lower coupling efficiency in middle- (55 years) and older-age (70 years) compared to young adults (S. H. Park et al., 2018; S.-Y. Park et al., 2020). Furthermore, mitochondrial-targeted antioxidants like Mitoquinone (MitoQ) restore vascular endothelial dysfunction in aged mice and patients with peripheral artery disease, which likely results from reductions in levels of mitochondrial superoxide (Gioscia-Ryan et al., 2014; S.-Y. Park et al., 2020). Together, evidence points towards mitochondria as promising targets for interventions aimed at combatting vascular endothelial dysfunction.

(-)-Epicatechin (EPI) belongs to a subclass of flavonoids known as the flavanols. Not only
is EPI highly bioavailable – reaching up to 10 µM in circulation in humans (Hollands et al.,
2013) - but EPI is also associated with several health benefits (Arts et al., 2001; Hertog et al.,
1993), including improved vascular endothelial function (Galleano et al., 2013; Karim et al.,
2000; Schroeter et al., 2006) and increased NO bioavailability (Schroeter et al., 2006).
Additional mechanisms thought to underly the therapeutic effects of EPI include: increased
eNOS phosphorylation (Ramirez-Sanchez et al., 2011, 2012, 2018), enhanced content or
activity of enzymatic antioxidant proteins (Moreno-Ulloa, Nogueira, et al., 2015; Ramirez-
Sanchez et al., 2014) and augmented mitochondrial biogenesis (Lee et al., 2015; Ramirez-
Sanchez et al., 2018; Taub et al., 2012, 2016). Whilst the potential of EPI to enhance markers
of mitochondrial biogenesis and antioxidant capacity seems promising, it is unclear whether
this effect translates to enhanced respiratory function or altered ROS production of vascular
endothelial cells. In-fact, one recent study reported no impact of EPI on mitochondrial
respiration (Keller et al., 2020), but the authors demonstrated that EPI may lower mitochondrial
ROS production of HUVECs in the presence of high glucose. The challenge remains to resolve
whether EPI modulates ROS production and mitochondrial function of vascular endothelial
cells.
112
113   To this end, our aim was to investigate whether EPI modulates reactive oxygen and nitrogen
114   species (RONS) production and mitochondrial function of human vascular endothelial cells.
115   We hypothesised that EPI would attenuate ROS production, augment NO bioavailability, and
116   enhance indices of mitochondrial function.
117
118

Materials and Methods

Cell culture and treatment

Human umbilical endothelial vein endothelial cells (HUVECs; Thermo Fisher Scientific,
Waltham, MA, USA) at passages 3-7 were used in this study. HUVECs were not passaged
more than 7 times because of changes in cell phenotype that can occur with multiple population
doublings that ultimately lead to senescence (Chang et al., 2005; Cheung, 2007; Grillari et al.,
2000). Following the plating of cells onto pre-gelatinised well-plates (0.2% gelatin) in
complete endothelial cell growth medium (EGM; Cell Applications Inc, San Diego, CA, USA),
~80% confluent HUVECs were washed twice with Dulbecco's phosphate-buffered saline (D-

PBS) and switched to pre-warmed (37°C) EGM in the absence (vehicle [H_2O], 'CTRL') or

presence of EPI (0.5-20 µM) over 24 h and 48 h. Human umbilical endothelial vein endothelial

cells (HUVECs; Thermo Fisher Scientific, Waltham, MA, USA) at passages 3-7 were used in

this study. HUVECs were not passaged more than 8 times because of changes in cell phenotype

that can occur with multiple population doublings that ultimately lead to senescence (Chang et

al., 2005; Cheung, 2007; Grillari et al., 2000). Following the plating of cells onto pre-

gelatinised well-plates (0.2% gelatin) in complete endothelial cell growth medium (EGM; Cell

Applications Inc, San Diego, CA, USA), ~80% confluent HUVECs were washed twice with

Dulbecco's phosphate-buffered saline (D-PBS) and switched to pre-warmed (37°C) EGM in

the absence (vehicle [H_2O], 'CTRL') or presence of EPI (0-20 µM) over 24 h and 48 h.

Cell viability

The fluorescent CyQUANT® Proliferation Assay kit was used to determine cell viability.

HUVECs were grown to 60-70% confluency in EGM in gelatinised 96-well plates. Cells were

subsequently dosed for 24 h in EGM +/- EPI at 0-20 µM. After 24 h, wells were aspirated,

washed twice with D-PBS, and then frozen immediately at -80°C. On the day of the assay,

plates were thawed at room temperature and CyQUANT® GR dye/cell-lysis buffer was added

to each well according to manufacturer instructions. Plates were gently mixed on an orbital

shaker (80 rpm) for 5 minutes protected from light. Sample fluorescence was measured using

a CLARIOStar plate reader (BMG Labtech, Ortenberg, Germany) with 485/520 Ex/Em.

Mitochondrial ROS production

Mitochondrial superoxide was detected in HUVECs using MitoSOX (Thermo Fisher

Scientific, Waltham, USA), a hydroethidine probe which is targeted to mitochondria by a

conjugated triphenyl-phosphonium moiety (Zielonka et al., 2017). HUVECs were seeded at 3

$\times 10^4$ cells/mL in gelatinised 12-well microplates and at ~80% confluence dosed +/- EPI (0-10

µM) for 24 h. Next, cells were washed in Krebs-Ringer buffer (KRH; 135 mM NaCl, 3.6 mM

KCl, 10 mM HEPES (pH 7.4), 0.5 mM $MgCl_2$, 1.5 mM $CaCl_2$, 0.5 mM NaH_2PO_4, 2 mM

glutamine and 5 mM D(+)-Glucose) prior to incubation at 37ºC for 30 minutes, with or without

15 µM antimycin A (AA) to stimulate mitochondrial superoxide production. Next, AA-

containing KRH was removed and MitoSOX was loaded into cells in fresh pre-warmed KRH

to a final concentration of 2.5 µM. Plates were immediately transferred to a plate reader

(ClarioStar, BMG Labtech) and fluorescence was monitored continuously at 30-sec intervals

over 30 min at excitation/emission of 510/580 nm. Rates of mitochondrial superoxide

production were determined from the slope of the resultant progress curve over the 30-minute

recording. Finally, plates were immediately fixed in 1% (v/v) acetic acid in methanol for the

determination of cell density by the Sulforhodamine B (SRB) assay, which was used to

normalise obtained fluorescence values. The plate reader's focal height and gain were

optimised and fixed between different experiments.

Cellular ROS

Cellular ROS were detected using the CellROX® Deep Red reagent by spectrophotometry.

HUVECs were seeded at 3×10^4 cells/mL into gelatinised 12-well microplates and at ~80%

confluence dosed +/- EPI (0-10 µM) for 24 h. After treatment, HUVECs were washed in KRH

with or without 15 µM AA and incubated at 37ºC for 30 minutes, prior to KRH removal and

CellROX loading using fresh, pre-warmed KRH buffer, to a final concentration of 2.5 µM.

Following 30 minutes CellROX incubation protected from light, cells were washed 2 × with

D-PBS and immediately transferred to a plate reader (ClarioStar, BMG Labtech), where

fluorescent CellROX oxidation products were excited at 640 nm and light emission detected at

665 nm. The plate reader's focal height and gain were optimised and fixed between

experiments. Upon completion of the reading, plates were immediately fixed for the

determination of cell density by as discussed above.

Nitric oxide bioavailability

To assess intracellular NO bioavailability, HUVECs were plated and cultured as previously

described. At ~80% confluency, cells were treated with 0, 5 or 10 µM EPI for 24 h. After

treatment, HUVECs were washed 2 × with D-PBS and loaded with DAF-FM™ diacetate (4-

amino-5-methylamino- 2′,7′-difluorofluorescein diacetate; Molecular Probes, Invitrogen) to a

final concentration of 1 µM in KRH buffer and incubated at 37°C for 45 minutes protected

from light. Following dye loading, cells were washed 2 × with D-PBS and immediately

trypsinised prior to pelleting and resuspension in D-PBS. Sample fluorescence was

subsequently detected at 495/515 Ex/Em by flow cytometry (BD Accuri C6, BD Biosciences,

Wokingham, UK). Data were recorded from 5,000 events and median values reported.

RT-qPCR – Gene expression quantification

HUVECs were grown to ~70% confluency in EGM in gelatinised 12-well plates and lysed

in 125 µL TRIzol. Total RNA was then extracted using the phenol-chloroform method. RNA

concentrations were determined by spectrophotometry (NanoDrop™ 2000, Thermo Fisher

Scientific, Waltham, USA). Specific primers used in each PCR are outlined in supplementary

Table 1. After preparation, reaction tubes were transferred to a Rotor-Gene Q PCR thermal

cycler for product amplification using a one-step protocol (QuantiFast SYBR® Green RT-PCR

Kit, Qiagen, UK). The amplification protocol was as follows: reverse transcription (10 minutes

at 50°C), transcriptase inactivation and initial denaturation (95°C for 5 min) followed by 40 ×

amplification cycles consisting of: 95°C for 10 s (denaturation) and 60°C for 30 s (annealing

and extension); followed by melt curve detection. Critical threshold (C_T) values were derived

from setting a threshold of 0.09 for all genes. To quantify gene expression, C_T values were used

to quantify relative gene expression using the comparative Delta Delta C_T ($2^{-\Delta\Delta CT}$) equation

(Livak & Schmittgen, 2001), whereby the expression of the gene of interest was determined

relative to the internal reference gene (RPL13a) in the treated sample, compared with the

untreated zero-hour control.

Mitochondrial respiration

HUVECs (passages 4-6) were seeded in XFe 24 well plates (Agilent, Santa Clara, CA,

USA) at 30,000 cells per well in 200 µL EGM for 48 h. After 48 h, HUVECs were washed

twice with D-PBS and replaced with fresh EGM containing 0, 5 and 10 µM EPI for 24 h. On

the day of the assay, HUVECs were washed with pre-warmed XF Dulbecco's Modified Eagle

Medium pH 7.4 (DMEM; Agilent, Santa Clara, CA, USA), supplemented with 5.5 mM

glucose, 1 mM sodium pyruvate and 2 mM L-Glutamine, and brought to a final well volume

of 500 µL. The cells were incubated in this medium for 45 minutes at 37°C in a non-CO_2

incubator and then transferred to a Seahorse XFe24 extracellular flux analyser (Agilent, Santa

Clara, CA, USA) maintained at 37°C. After an initial 15-minute calibration, oxygen

consumption rate (OCR) was measured by a 3-4 loop cycle consisting of a 1-min mix, 2-min

wait and 3-min measure to record cellular respiration. After measuring basal respiration, 2 mM

oligomycin was added to selectively inhibit the mitochondrial ATP synthase. Subsequently, 3

µM BAM15 was added to uncouple OCR to determine maximal respiration, and finally a

mixture of 2 µM rotenone and 2 µM AA was added to inhibit complex I and III of the electron

transfer system, respectively, to determine non-mitochondrial respiration. Rates of oxygen

consumption were corrected for non-mitochondrial respiration and expressed relative to the

cell number of the appropriate well, determined by the CyQUANT® assay. The raw values of
extracellular acidification rate (ECAR) and OCR were divided into component rates to
calculate the relative contribution of glycolytic (ATP_{glyc}) and oxidative ATP-producing
reactions (ATP_{ox}) to total ATP production, as previously described (Mookerjee & Brand,
2015). Three independent experiments were performed that contained at least two technical
replicates.

SDS-PAGE and immunoblotting

Total protein and phosphoprotein levels were detected by western blot. Following
treatment, HUVECs were lysed and scraped in ice-cold 1x precipitation assay buffer
containing: 25 mM Tris-HCl pH 7.6, 150 mM NaCl, 1% NP-40, 1% sodium deoxycholate and
0.1% SDS, supplemented with 1x Protease Inhibitor Cocktail Set V (Merck Life Science, UK).
Cell lysates were centrifuged for 15 minutes at $18,000 \times g$ (4°C) and the supernatant was stored
at -80°C before analyses for total protein by the Pierce BCA™ assay. Samples were
subsequently resuspended in 4x Laemmli buffer (Bio-Rad laboratories, Hertfordshire, UK)
containing reducing agent (1x working concentration: 31.5 mM Tris-HCl [pH 6.8], 10%
glycerol, 1% SDS, 0.005% Bromophenol Blue and 355 mM 2-mercaptoethanol) and were
loaded and electrophoresed on 10% SDS-stain-free polyacrylamide gels (supplementary Figure
1). Semi-dry transfer of proteins to a nitrocellulose membrane was performed using the Trans-
Blot® Turbo™ Transfer System. Following blocking for 1-hour in Tris-buffered saline Tween-
20 (TBS-T) containing 5% non-fat dried milk (NFDM), membranes were incubated overnight
with rabbit anti-phosphorylated or total antibodies: AMPKα, pThr172-AMPK, p44/42 MAPK,
pThr202/Tyr204-p44/42 MAPK, eNOS and pSer1177-eNOS, at a dilution of 1:500-1:4000 in
5% bovine serum albumin (BSA) made up in TBS-T (see Table 1). After overnight incubation,
the membranes were washed 3 times in TBS-T for 5 minutes and incubated for 1 hour in HRP-
conjugated anti-rabbit antibodies (Cell Signaling Technology, London, UK) at dilution of
1:5000-1:10,000, following optimisation. Proteins were visualised by enhanced
chemiluminescence (Thermo Fisher Scientific inc, Waltham, USA) and quantified by
densitometry (ChemiDoc™ MP imaging system, Bio-Rad Laboratories, Inc. CA, USA).
Analysed western blot images are presented in supplementary figure 2.

Table 1. List of antibodies and dilutions used.

Antibody	Primary Ab Dilution	Secondary Ab Dilution
AMPKα	1:1000	1:5000
pThr172-AMPK	1:1000	1:10,000
p44/42 MAPK	1:2000	1:10,000
pThr202/Tyr204-p44/42 MAPK	1:2000	1:10,000
eNOS	1:500	1:5000
pSer1177-eNOS	1:500	1:5000

All antibodies were purchased from Cell Signaling Technology.

Statistical analysis

One-way ANOVAs were employed to detect effects of EPI treatment. Two-way ANOVAs

were employed to detect main effects (e.g., dose, time or antimycin A) and potential significant

interactions between two main independent factors. Multiple comparisons between

experimental conditions were adjusted for multiple tests, using Dunnett's or Sidak's where

appropriate. All data are presented as mean±SEM and significance accepted when $P<0.05$.

Results

EPI does not cause vascular endothelial cell toxicity

After 24 h EPI treatment (0.5-20 µM dose responses), there was a significant main effect

of dose on cell proliferation (P=0.018; Figure 1). However, multiple comparisons revealed no

significant difference between doses of EPI versus CTRL. Given that EPI did not cause cell

toxicity, and prior knowledge of physiologic EPI concentrations *in vivo* (up to 10 µM),

subsequent experiments were performed with doses of 5 and 10 µM.

Figure 1. EPI does not cause vascular endothelial cell toxicity. HUVECs were treated with 0-20 μM EPI for 24 h. Data are means±SEM, representative of 3 independent repeats with 3 replicates of each condition. Statistical significance was tested for by one-way ANOVA and Dunnett's test for multiple comparisons. $^{\alpha}$ Significant main effect of dose ($P<0.05$).

EPI dose-dependently modulates mitochondrial RONS production

Next, we assessed whether EPI, in the absence or presence of the complex III inhibitor, AA, impacted mitochondrial superoxide emission. There was a significant main effect of dose and AA on rates of MitoSOX oxidation ($P<0.001$), and a significant dose × AA interaction ($P<0.001$; Figure 2A). Post-hoc comparisons revealed that, in the absence of AA (-AA), 5 μM EPI significantly increased and 10 μM EPI decreased rates of MitoSOX oxidation compared to CTRL, respectively (CTRL: $8.1\times10^{-5}\pm0.2\times10^{-5}$; 5 μM EPI: $10.7\times10^{-5}\pm0.2\times10^{-5}$; 10 μM EPI: $3.8\times10^{-5}\pm0.2\times10^{-5}$ RFU/sec^{-1}/cell^{-1}; $P<0.001$). The associated raw traces of MitoSOX oxidation are displayed in Figure 2B. In the presence of AA (+AA), 5 μM EPI did not affect MitoSOX oxidation versus CTRL (5 μM EPI: $35.4\times10^{-5}\pm0.4\times10^{-5}$ vs. CTRL: $35.4\times10^{-5}\pm0.5\times10^{-5}$ RFU/sec^{-1}/cell^{-1}; Figure 2A). Whereas 10 μM EPI significantly increased rates of MitoSOX oxidation compared to CTRL (10 μM EPI: $44.4\times10^{-5}\pm0.6\times10^{-5}$ vs. CTRL: $35.4\times10^{-5}\pm0.5\times10^{-5}$ RFU/sec^{-1}/cell^{-1}; $P<0.001$). In contrast to mitochondrial ROS, cellular ROS production (not mitochondrial-specific) was not altered by EPI (Figure 2C).

After revealing that rates of mitochondrial ROS production were dose-dependently altered by EPI, we assessed whether EPI modified intracellular NO levels. There was a significant main effect of EPI dose on NO levels ($P=0.003$; Figure 2D). Whilst 5 μM EPI did not impact NO levels, 10 μM EPI significantly increased NO levels compared to CTRL conditions (10 μM EPI: $4.38\times10^{5}\pm0.43\times10^{5}$ vs. CTRL: $3.02\times10^{5}\pm0.18\times10^{5}$ AU; $P=0.010$).

Figure 2. EPI dose-dependently impacts RONS production in vascular endothelial cells. A) MitoSOX oxidation rates in HUVECs after 24 h EPI treatment and 30 minutes incubation with or without antimycin A. B) Mean unnormalized MitoSOX oxidation rates measured in 30 second intervals over 30 minutes. C) CellROX oxidation in HUVECs after 24 h EPI treatment. D) DAF-FM oxidation in HUVECs after 24 h EPI treatment measured by flow cytometry. Data are means±SEM of three independent repeats with two replicates per treatment. Statistical significance was tested for by a two-way ANOVA, with dose and antimycin A as factors: [α] Significant main effect of dose; [δ] Significant main effect of AA ($P<0.05$). *$P<0.05$ and ****$P<0.0001$.

EPI dose-dependently impacts the expression of genes associated with energy metabolism in vascular endothelial cells

Next, experiments were performed to resolve whether EPI alters the expression of genes linked with mitochondrial function and the antioxidant response. Firstly, the expression of genes associated with mitochondrial function and remodelling were quantified. In the presence of EPI, there was a significant main effect of dose ($P=0.018$) and time ($P=0.002$) on dynamin-

related protein 1 (DRP1) expression (Figure 3A), but no significant dose × time interaction. At 48 h, 10 μM EPI increased DRP1 expression 2.2-fold compared to CTRL ($P=0.010$). There was a significant main effect of dose on mitofusin-2 (MFN2) mRNA expression in cells treated with EPI ($P=0.035$; Figure 3B). With 10 μM EPI, MFN2 expression increased 1.6-fold versus CTRL ($P=0.024$). Parkin, PGC-1α, and transcription Factor A (TFAM) expression were not changed by EPI treatment (supplementary Figure 3).

Genes associated with the antioxidant response were also quantified. There was no effect of dose or time on catalase or eNOS expression. There was a significant main effect of dose and time on NADPH oxidase 4 (NOX4) expression in EPI treated cells, respectively ($P=0.015$ and $P=0.006$). A significant main effect of dose was found on nuclear factor-erythroid factor 2-related factor 2 (NRF2) expression in the presence of EPI ($P<0.001$; Figure 3C). NRF2 mRNA abundance was increased 1.5-fold and 1.6-fold with 5 and 10 μM EPI over 48 h when compared to CTRL ($P=0.015$ and $P=0.001$, respectively). There was a significant effect of time on superoxide dismutase 2 (SOD2) expression in EPI treated cells only ($P=0.024$). Multiple comparisons revealed that SOD2 expression was increased 2.1-fold in the presence of 10 μM EPI versus CTRL conditions ($P=0.040$; Figure 3D).

Figure 3. Gene expression responses following acute EPI treatment. HUVECs were treated with 0, 5 and 10 μM EPI over 48 h and lysed for analysis of gene expression. A) DRP1, B) MFN2, C) NRF2 and D) SOD2. Data are means±SEM from 3 independent experiments. Statistical significance was determined by a two-way ANOVA, with dose and time as factors. Multiple comparisons were performed by Dunnett's test to determine differences in gene expression between conditions. [α] Main effect of dose; [β] main effect of time ($P<0.05$); *$P<0.05$, **$P<0.01$.

EPI does not alter mitochondrial bioenergetics of vascular endothelial cells

Having described that EPI influences RONS production and alters the expression of

genes linked with mitochondrial function, we tested whether EPI impacted vascular endothelial

cell bioenergetics. There was no significant main effect of EPI on rates of basal respiration,

maximal respiration, ADP phosphorylation, proton leak, spare respiratory capacity (%) or

coupling efficiency, regardless of dose (Figure 4A & 4B & 4D). There was no effect of EPI on

the relative contribution of ATP_{glyc} or ATP_{ox} to total ATP production (Figure 4C).

Figure 4. EPI does not directly affect mitochondrial bioenergetics. A) Mitochondrial bioenergetics of HUVECs following 24 h EPI treatment (0, 5 and 10 µM). B) Coupling efficiency of oxidative phosphorylation. C) Relative contribution of glycolytic (grey) and oxidative (black) ATP production to total ATP production rates. D) Representative trace of oxygen consumption rates during the mitochondrial stress test. Data from 3 independent experiments are normalised to cell number (1×10^3) and presented as mean±SEM. Max, maximal respiratory capacity; PL, proton leak; ADP_{phos}, ADP phosphorylation; SRC, spare respiratory capacity.

EPI rapidly and transiently promotes ERK1/2 phosphorylation, independent of AMPK

348   To further probe how EPI might alter vascular endothelial transcription, we assessed

349   cell signalling responses. There was a significant main effect of treatment and time on AMPKα

350   phosphorylation, and a significant interaction was observed ($P<0.001$; Figure 5A). Multiple

comparisons revealed a significant increase in phosphorylation of AMPKα at Thr172 at 1 h versus 0 h (1 h: 2.24±0.22 vs. 0 h: 1.16±0.19 AU) under CTRL conditions ($P=0.006$), whereas there was no significant change at 1 h versus 0 h with EPI (1 h: 0.47±0.18 vs. 0 h: 1.16±0.19 AU; $P=0.157$). However, there was a significant reduction in AMPKα phosphorylation in the presence of EPI vs. CTRL at 15 min (EPI: 0.35±0.09 vs. CTRL: 1.86±0.29 AU; $P<0.001$) and 1 h (EPI: 0.47±0.18 vs. CTRL: 2.24±0.22 AU; $P<0.001$). From 3 hours, AMPKα phosphorylation was suppressed under both CTRL (3 h: 0.54±0.08 and 24 h: 0.39±0.12 AU vs. 0 h: 1.16±0.19 AU; $P=0.046$ and $P=0.026$, respectively) and EPI conditions compared to 0 h CTRL (3 h: 0.37±0.08 and 24 h: 0.46±0.08 AU vs. 0 h: 1.16±0.19 AU; $P=0.021$ and $P=0.049$, respectively).

Whilst ERK1/2 is not involved in the AMPK/eNOS pathway, ERK1/2 signalling may mediate the effects of EPI upon vascular endothelial cell adaptation. There was no main effect of treatment ($P=0.141$), but a significant main effect of time on ERK1/2 phosphorylation ($P=0.039$; Figure 5B). There was also a significant treatment × time interaction ($P=0.003$). Under CTRL conditions, despite a 2.7-fold increase in ERK1/2 phosphorylation at 3 h, ERK1/2 phosphorylation did not reach significance vs. 0 h until 24 hours post treatment (0 h: 0.38±0.10 vs. 24 h: 1.15±0.28 AU; $P=0.022$). By contrast, EPI significantly increased ERK1/2 phosphorylation at 15 minutes vs. 0 h (15 min: 1.09±0.24 vs. 0 h: 0.38±0.10 AU; $P=0.035$), which was retained at 1 hour (1 h: 1.22±0.07 vs. 0 h: 0.38±0.10 AU; $P=0.011$), before returning to baseline levels, suggesting a change in the temporal pattern of ERK1/2 activation because of EPI treatment. Indeed at 1-hour, EPI treatment resulted in a 4.3-fold increase in ERK1/2 phosphorylation vs. the 1-hour untreated CTRL ($P=0.007$).

Figure 5. EPI rapidly and transiently activates ERK1/2 signalling whilst blunting AMPK phosphorylation. A) AMPKα phosphorylation at Thr172 in HUVECs in the absence (-; clear bars) or presence (+; green bars) of EPI. B) ERK1/2 phosphorylation at Thr202/Tyr204. C) eNOS phosphorylation at Ser1177. D) Representative images of n=3 independent experiments. Cell lysates were analysed by SDS-PAGE and western blotting with indicated antibodies. Data are expressed as means±SEM; *$P<0.05$ and ***$P<0.001$. [α] Significant main effect of treatment; [β] significant main effect of time ($P<0.05$).

375   To help establish whether the increase in NO brought about by EPI was associated with

376 eNOS signalling, phosphorylation of eNOS at Ser1177 was assessed. There was no main effect

377 of treatment or time on eNOS phosphorylation (Figure 5C), and no treatment × time interaction

378 ($P=0.100$). At 3 h, eNOS phosphorylation was ~60% higher under CTRL versus EPI conditions

379 (CTRL: 0.38±0.10 vs EPI: 1.15±0.28 AU, $P=0.038$).

381 **Discussion**

382  Resolving EPI's mode of action using vascular endothelial cells as a model will help to

383 establish its potential efficacy in mitigating vascular endothelial dysfunction. We tested the

384 hypothesis that EPI would attenuate ROS production, augment NO bioavailability and enhance

385 mitochondrial function of human vascular endothelial cells in culture. We demonstrated that

386 physiologic EPI concentrations, dose-dependently, modulated RONS emission but did not

387 directly impact mitochondrial respiration. Moreover, the influence of EPI on RONS emission

388 was associated with the induction of increased NRF2 mRNA, which appears downstream of

389 rapid and transient activation of ERK1/2 signalling. Taken together our findings expand our

390 knowledge of EPI's mechanisms of action *in vitro* and support further research on EPI as a

391 potential instigator of cell signalling and NRF2 activation *in vivo*.

EPI dose-dependently modulates RONS production

We have demonstrated that EPI dose-dependently altered mitochondrial ROS production in the absence of additional cell stress. Increased rates of mitochondrial ROS production, reported in the presence of 5 µM EPI supports previous observations that 10 days EPI supplementation increased superoxide production in mitochondria isolated from mouse heart tissue (Panneerselvam et al., 2013). However, EPI has been demonstrated to increase the abundance and/or activity of key antioxidant proteins like SOD2 and catalase (Bettaieb et al., 2014; Calabró et al., 2016), and even to blunt hydrogen peroxide production from isolated brain and heart mitochondria (Lagoa et al., 2011). In a similar way, we found that higher (10 µM) EPI concentrations attenuated the rate of mitochondrial superoxide production, which may have been facilitated by increased SOD2 mRNA expression. Clearly and importantly, the divergent effects of EPI on mitochondrial ROS production suggest EPI's biological effects are highly dose dependent. Future studies should investigate whether EPI alters the vascular endothelial cell redox state and its potency relative to known mitochondrial antioxidants. Another noteworthy observation of this study was that EPI did not rescue AA-induced increases in mitochondrial ROS. Likewise, one recent study reported that 1 µM EPI did not lower the production of mitochondrial superoxide in HUVECs after acute AA treatment (Keller et al., 2020), suggesting limited efficacy for EPI in overcoming conditions associated with elevated mitochondrial ROS production in the vascular endothelium of humans.

The potent stimulation of NO production by EPI is well documented in studies using cell, human and rodent models (Loke et al., 2008; Moreno-Ulloa, Mendez-Luna, et al., 2015; Ramirez-Sanchez et al., 2011, 2018; Schroeter et al., 2006), although not all studies have reported such effects (Dower et al., 2015). Here, we demonstrated that EPI increased NO levels of vascular endothelial cells, suggesting that EPI may be a promising strategy to combat vascular endothelial dysfunction, although further *in vivo* studies are required. Previous studies have attributed elevated NO bioavailability in the presence of EPI to increased phosphorylation of eNOS at Ser177 (Carnevale et al., 2014; Ramirez-Sanchez et al., 2010, 2012, 2018; Ramírez-Sánchez et al., 2016). Given that we and others demonstrated unaltered eNOS phosphorylation in the presence of EPI (Keller et al., 2020), it is plausible that arginase inhibition is the potential mechanism by which EPI increases NO production in HUVECs (Schnorr et al., 2008). However, this may not be the case in arterial endothelial cells, where EPI has repeatedly been

425 shown to enhance eNOS Ser1177 phosphorylation (Ramirez-Sanchez et al., 2012, 2018;
426 Ramírez-Sánchez et al., 2016).

EPI augments the expression of genes linked with the antioxidant response and mitochondrial dynamics

To better understand the mechanisms underlying EPI's effects, we measured the
transcription of genes associated with energy metabolism. Interestingly, EPI increased the
expression of genes DRP1 and MFN2, respectively involved in mitochondrial fission and
fusion. Despite these effects, EPI did not impact mitochondrial respiration, at least over 24
hours. Although mitochondrial dynamics can influence respiratory function (Chen et al., 2010;
Glancy et al., 2020), the lack of functional change in mitochondrial respiration could reflect a
discord between cellular mRNA responses and changes in protein abundance and/or function.
Regardless of dose, EPI significantly enhanced NRF2 mRNA expression. This observation
implies that the induction of NRF2 mRNA by EPI is independent of RONS production and
might be explained by alternate factors that control NRF2 activity, like phosphorylation status
(Robledinos-Antón et al., 2019). Although post-translational modifications of NRF2 were not
assessed in this study, flavonoids have been shown to promote NRF2 phosphorylation and
nuclear translocation (Lan et al., 2017; Shi et al., 2018), which would likely increase the
transcription of genes related to the antioxidant response, including NRF2 itself.

EPI does not impact mitochondrial respiration

One important finding was that EPI does not impact mitochondrial respiration of HUVECs
in culture. Although mitochondria have been proposed as potential molecular targets of EPI
(Duluc et al., 2012), previous investigations into EPI's effects on mitochondrial respiration
have produced equivocal results. Some studies have demonstrated increased state 3 respiration
in rat beta cells following EPI (0.1-2.5 µM) supplementation (Kener et al., 2018; Rowley et al.,
2017), whilst others have reported inhibited or similar state 3 respiration rates with EPI,
depending on the substrates provided (Kopustinskiene et al., 2015). One recent study using
HUVECs as a model reported that 0.1 and 1 µM EPI treatment over 2 hours had negligible
impact on mitochondrial respiration (Keller et al., 2020). Taken together, our data suggest that
the therapeutic potential of EPI is not related to changes in mitochondrial respiration in vascular

endothelial cells, pointing to alternate potential mechanisms of action like cell signalling
activation (Fraga et al., 2018).

EPI rapidly and transiently activates ERK1/2 signalling

The serine/threonine protein kinase AMPK is an important regulator of mitochondrial
adaptation (Herzig & Shaw, 2018). However, in our studies AMPK phosphorylation was
suppressed in the presence of EPI for up to 1 h and was without further impact for up to 24 h,
compared with controls. Supporting these findings, 2 h EPI treatment (1 μM) did not affect the
phosphorylation of AMPK in HUVECs (Keller et al., 2020). Although EPI is capable of
augmenting AMPK activity in liver and muscle tissue, and several cell types (Murase et al.,
2009; Papadimitriou et al., 2014; Si et al., 2011), it seems that AMPK activation is not
responsible for the therapeutic actions of EPI in HUVECs in culture. Together with the
negligible impact of EPI on mitochondrial respiration, the data suggest that EPI does not
directly alter vascular endothelial cell energy metabolism, *in vitro*.

Importantly and of novelty, we reported that ERK1/2 phosphorylation at Thr202/Tyr204
was rapidly and transiently increased by EPI in HUVECs (Figure 5). This finding supports
recent observations of increased ERK1/2 signalling after 0.1 μM EPI treatment in bovine
coronary artery endothelial cells, that may be associated with phosphorylation of CaMKII
(Moreno-Ulloa, Mendez-Luna, et al., 2015). How EPI promotes ERK1/2 phosphorylation in
vascular endothelial cells remains to be described, but current evidence suggests EPI activates
ERK1/2 by binding to the G-protein coupled estrogen receptor (GPER) on the cell membrane
(Moreno-Ulloa, Mendez-Luna, et al., 2015; Yang & Chan, 2018). Given the induction of NRF2
mRNA expression found in this study, it would be useful to determine if acute activation of
ERK1/2 signalling by EPI is a prerequisite for the induction of NRF2 activity.

Limitations

We did not use metabolites of EPI that appear in circulation after ingestion of EPI-
containing foods or supplements *in vivo* (Ottaviani et al., 2016). Therefore, the present findings
may have limited translational potential. In our study we harnessed HUVECs to model the
vascular endothelial cell, and given that these cells are venous in nature, their physiology may
not well reflect the arterial vasculature or microcirculation where EPI potentially exerts its
beneficial effects. Further, not all assays were performed in the presence of additional cell

stress. Thus, caution should be taken when translating our findings to human populations with

disease. Finally, post-translational modifications of NRF2, which are critical for regulating

NRF2's activity, were not measured in this study.

Conclusion

We demonstrate that physiologic EPI concentrations do not impact mitochondrial

respiration but do modulate reactive oxygen and nitrogen species production and the signalling

and transcriptional activities of vascular endothelial cells *in vitro*. EPI's dose-dependent

alteration of reactive oxygen and nitrogen species production occurred in parallel with

enhanced and transient ERK1/2 signalling, and the induction of NRF2 mRNA (Figure 6). The

fact that EPI enhanced NRF2 mRNA expression regardless of dose, implies that alterations in

reactive oxygen and nitrogen species production alone were not solely responsible. Further

research will help clarify the precise way in which EPI promotes ERK1/2 signalling and NRF2

activity, and its relevance to vascular endothelial health *in vivo*.

Figure 6. Schematic of the potential mechanisms by which EPI exerts its biological effects in vascular
endothelial cells. Solid arrows/lines signify EPI's mode of action demonstrated in this study. Dashed
arrows represent potential activity of EPI not examined in this study but reported previously.

Availability of data and material

The datasets generated and analysed during this study are available from the corresponding
author upon request.

Conflict of interest

Daniel G. Sadler, Jonathan Barlow, Helen Jones, Dick H. J. Thijssen and Claire E. Stewart had
no conflict of interest associated with this manuscript. Richard Draijer is employed by
Unilever.

Funding

The study was supported by funding received from the Biotechnology and Biological Sciences
Research Council (BBSRC) and Unilever.

Authors' contributions

DGS and CES conceived the study and designed experiments. DGS and JB designed the
respiration experiments using the Seahorse XFe96 Analyzer. DGS collected and
analysed the data, DGS, JB, RD, HJ, DHJT and CES interpreted the data. DGS and CES wrote
the manuscript and all authors revised it critically. All authors provided final approval of the
version to be published and agree to be accountable for all aspects of the work in ensuring that
questions related to the accuracy or integrity of any part of the work are appropriately
investigated and resolved. All people designated as authors qualify for authorship, and all those
who qualify for authorship are listed. CES is the guarantor for the work and/or conduct of the
study, had full access to all the data in the study and takes responsibility for the integrity of
data and the accuracy of the data analysis, and controlled the decision to publish.

Acknowledgements

The authors acknowledge use of the Mitochondrial Profiling Centre, a core resource supported
by the University of Birmingham. This study was supported by the BBSRC funded Industrial
CASE (iCASE) award (BB/P504385/1).

539
540 **References**
541

Arts, I. C. W., Jacobs, D. R., Harnack, L. J., Gross, M., & Folsom, A. R. (2001). Dietary

catechins in relation to coronary heart disease death among postmenopausal women.

Epidemiology. https://doi.org/10.1097/00001648-200111000-00015

Bettaieb, A., Vazquez Prieto, M. A., Rodriguez Lanzi, C., Miatello, R. M., Haj, F. G., Fraga,

C. G., & Oteiza, P. I. (2014). (−)-Epicatechin mitigates high-fructose-associated

insulin resistance by modulating redox signaling and endoplasmic reticulum stress.

Free Radical Biology and Medicine, *72*, 247–256.

https://doi.org/10.1016/j.freeradbiomed.2014.04.011

Burns, E. M., Kruckeberg, T. W., Comerford, L. E., & Buschman, M. T. (1979). Thinning of

capillary walls and declining numbers of endothelial mitochondria in the cerebral

cortex of the aging primate, Macaca nemestrina. *Journals of Gerontology*.

https://doi.org/10.1093/geronj/34.5.642

Calabró, V., Piotrkowski, B., Fischerman, L., Vazquez Prieto, M. A., Galleano, M., & Fraga,

C. G. (2016). Modifications in nitric oxide and superoxide anion metabolism induced

by fructose overload in rat heart are prevented by (−)-epicatechin. *Food & Function*,

7(4), 1876–1883. https://doi.org/10.1039/C6FO00048G

Carnevale, R., Loffredo, L., Nocella, C., Bartimoccia, S., Bucci, T., De Falco, E., Peruzzi,

M., Chimenti, I., Biondi-Zoccai, G., Pignatelli, P., Violi, F., & Frati, G. (2014).

Epicatechin and catechin modulate endothelial activation induced by platelets of

patients with peripheral artery disease. *Oxidative Medicine and Cellular Longevity*.

https://doi.org/10.1155/2014/691015

Cernadas, M. R., De Miguel, L. S., García-Durán, M., González-Fernández, F., Millás, I.,

Montón, M., Rodrigo, J., Rico, L., Fernández, P., De Frutos, T., Rodríguez-Feo, J. A.,

Guerra, J., Caramelo, C., Casado, S., & López-Farré, A. (1998). Expression of

constitutive and inducible nitric oxide synthases in the vascular wall of young and

aging rats. *Circulation Research*. https://doi.org/10.1161/01.RES.83.3.279

Chang, M. W. F., Grillari, J., Mayrhofer, C., Fortschegger, K., Allmaier, G., Marzban, G.,

Katinger, H., & Voglauer, R. (2005). Comparison of early passage, senescent and

hTERT immortalized endothelial cells. *Experimental Cell Research*.

https://doi.org/10.1016/j.yexcr.2005.05.002

Chen, H., Vermulst, M., Wang, Y. E., Chomyn, A., Prolla, T. A., McCaffery, J. M., & Chan,

D. C. (2010). Mitochondrial fusion is required for mtDNA stability in skeletal muscle

and tolerance of mtDNA mutations. *Cell*, *141*(2), 280–289. PubMed.

https://doi.org/10.1016/j.cell.2010.02.026

Cheung, A. L. (2007). Isolation and Culture of Human Umbilical Vein Endothelial Cells

(HUVEC). In *Current Protocols in Microbiology*.

https://doi.org/10.1002/9780471729259.mca04bs4

Chou, T. C., Yen, M. H., Li, C. Y., & Ding, Y. A. (1998). Alterations of nitric oxide synthase

expression with aging and hypertension in rats. *Hypertension*.

https://doi.org/10.1161/01.HYP.31.2.643

Csiszar, A., Labinskyy, N., Orosz, Z., Xiangmin, Z., Buffenstein, R., & Ungvari, Z. (2007).

Vascular aging in the longest-living rodent, the naked mole rat. *American Journal of*

Physiology - Heart and Circulatory Physiology.

https://doi.org/10.1152/ajpheart.01287.2006

Csiszar, A., Ungvari, Z., Edwards, J. G., Kaminski, P., Wolin, M. S., Koller, A., & Kaley, G.

(2002). Aging-induced phenotypic changes and oxidative stress impair coronary

arteriolar function. *Circulation Research*.

https://doi.org/10.1161/01.RES.0000020401.61826.EA

Donato, A. J., Eskurza, I., Silver, A. E., Levy, A. S., Pierce, G. L., Gates, P. E., & Seals, D.

R. (2007). Direct evidence of endothelial oxidative stress with aging in humans:

Relation to impaired endothelium-dependent dilation and upregulation of nuclear

factor-κB. *Circulation Research*.

https://doi.org/10.1161/01.RES.0000269183.13937.e8

Donato, A. J., Gano, L. B., Eskurza, I., Silver, A. E., Gates, P. E., Jablonski, K., & Seals, D.

R. (2009). Vascular endothelial dysfunction with aging: Endothelin-1 and endothelial

nitric oxide synthase. *American Journal of Physiology - Heart and Circulatory*

Physiology. https://doi.org/10.1152/ajpheart.00689.2008

Dower, J. I., Geleijnse, J. M., Gijsbers, L., Zock, P. L., Kromhout, D., & Hollman, P. C. H.

(2015). Effects of the pure flavonoids epicatechin and quercetin on vascular function

and cardiometabolic health: A randomized, double-blind, placebo-controlled,

crossover trial. *American Journal of Clinical Nutrition*.

https://doi.org/10.3945/ajcn.114.098590

Duluc, L., Soleti, R., Clere, N., Andriantsitohaina, R., & Simard, G. (2012). Mitochondria As

Potential Targets of Flavonoids: Focus on Adipocytes and Endothelial Cells. *Current*

Medicinal Chemistry. https://doi.org/10.2174/092986712803251467

Fraga, C. G., Oteiza, P. I., & Galleano, M. (2018). Plant bioactives and redox signaling: (–)-

Epicatechin as a paradigm. *Molecular Aspects of Medicine*.

https://doi.org/10.1016/j.mam.2018.01.007

Galleano, M., Bernatova, I., Puzserova, A., Balis, P., Sestakova, N., Pechanova, O., & Fraga,

C. G. (2013). Epicatechin reduces blood pressure and improves vasorelaxation in

spontaneously hypertensive rats by NO-mediated mechanism. *IUBMB Life*.

https://doi.org/10.1002/iub.1185

Gioscia-Ryan, R. A., LaRocca, T. J., Sindler, A. L., Zigler, M. C., Murphy, M. P., & Seals,

D. R. (2014). Mitochondria-targeted antioxidant (MitoQ) ameliorates age-related

arterial endothelial dysfunction in mice. *Journal of Physiology*.

https://doi.org/10.1113/jphysiol.2013.268680

Glancy, B., Kim, Y., Katti, P., & Willingham, T. B. (2020). The Functional Impact of

Mitochondrial Structure Across Subcellular Scales. *Frontiers in Physiology*, *11*, 1462.

https://doi.org/10.3389/fphys.2020.541040

Gliemann, L., Rytter, N., Piil, P., Nilton, J., Lind, T., Nyberg, M., Cocks, M., & Hellsten, Y.

(2018). The Endothelial Mechanotransduction Protein Platelet Endothelial Cell

Adhesion Molecule-1 Is Influenced by Aging and Exercise Training in Human

Skeletal Muscle. *Frontiers in Physiology*, *9*, 1807.

https://doi.org/10.3389/fphys.2018.01807

Gouill, E. Le, Jimenez, M., Binnert, C., Jayet, P.-Y., Thalmann, S., Nicod, P., Scherrer, U., &

Vollenweider, P. (2007). Endothelial Nitric Oxide Synthase (eNOS) Knockout Mice

Have Defective Mitochondrial β-Oxidation. *Diabetes*, *56*(11), 2690 LP – 2696.

https://doi.org/10.2337/db06-1228

Grillari, J., Hohenwarter, O., Grabherr, R. M., & Katinger, H. (2000). Subtractive

hybridization of mRNA from early passage and senescent endothelial cells.

Experimental Gerontology. https://doi.org/10.1016/S0531-5565(00)00080-2

Hamilton, C. A., Brosnan, M. J., McIntyre, M., Graham, D., & Dominiczak, A. F. (2001).

Superoxide excess in hypertension and aging a common cause of endothelial

dysfunction. *Hypertension*. https://doi.org/10.1161/01.HYP.37.2.529

Heitzer, T., Schlinzig, T., Krohn, K., Meinertz, T., & Münzel, T. (2001). Endothelial

Dysfunction, Oxidative Stress, and Risk of Cardiovascular Events in Patients With

Coronary Artery Disease. *Circulation*, *104*(22), 2673–2678.

https://doi.org/10.1161/hc4601.099485

Hertog, M. G. L., Feskens, E. J. M., Kromhout, D., Hertog, M. G. L., Hollman, P. C. H.,

Hertog, M. G. L., & Katan, M. B. (1993). Dietary antioxidant flavonoids and risk of

coronary heart disease: The Zutphen Elderly Study. *The Lancet*.

https://doi.org/10.1016/0140-6736(93)92876-U

Herzig, S., & Shaw, R. J. (2018). AMPK: guardian of metabolism and mitochondrial

homeostasis. *Nature Reviews. Molecular Cell Biology*, *19*(2), 121–135. PubMed.

https://doi.org/10.1038/nrm.2017.95

Hollands, W. J., Hart, D. J., Dainty, J. R., Hasselwander, O., Tiihonen, K., Wood, R., &

Kroon, P. A. (2013). Bioavailability of epicatechin and effects on nitric oxide

metabolites of an apple flavanol-rich extract supplemented beverage compared to a

whole apple puree: A randomized, placebo-controlled, crossover trial. *Molecular*

Nutrition & Food Research, *57*(7), 1209–1217.

https://doi.org/10.1002/mnfr.201200663

Jablonski, K. L., Seals, D. R., Eskurza, I., Monahan, K. D., & Donato, A. J. (2007). High-

dose ascorbic acid infusion abolishes chronic vasoconstriction and restores resting leg

blood flow in healthy older men. *Journal of Applied Physiology*.

https://doi.org/10.1152/japplphysiol.00533.2007

Karim, M., McCormick, K., & Kappagoda, C. T. (2000). Effects of Cocoa Extracts on

Endothelium-Dependent Relaxation. *The Journal of Nutrition*.

https://doi.org/10.1093/jn/130.8.2105s

Keller, A., Hull, S. E., Elajaili, H., Johnston, A., Knaub, L. A., Chun, J. H., Walker, L.,

Nozik-Grayck, E., & Reusch, J. E. B. (2020). (–)-Epicatechin Modulates

Mitochondrial Redox in Vascular Cell Models of Oxidative Stress. *Oxidative*

Medicine and Cellular Longevity, *2020*, 6392629.

https://doi.org/10.1155/2020/6392629

Kener, K. B., Munk, D. J., Hancock, C. R., & Tessem, J. S. (2018). High-resolution

respirometry to measure mitochondrial function of intact beta cells in the presence of

natural compounds. *Journal of Visualized Experiments*. https://doi.org/10.3791/57053

Kopustinskiene, D. M., Savickas, A., Vetchý, D., Masteikova, R., Kasauskas, A., &

Bernatoniene, J. (2015). Direct effects of (-)-Epicatechin and procyanidin B2 on the

respiration of rat heart mitochondria. *BioMed Research International*.

https://doi.org/10.1155/2015/232836

Lagoa, R., Graziani, I., Lopez-Sanchez, C., Garcia-Martinez, V., & Gutierrez-Merino, C.

(2011). Complex I and cytochrome c are molecular targets of flavonoids that inhibit

hydrogen peroxide production by mitochondria. *Biochimica et Biophysica Acta -*

Bioenergetics. https://doi.org/10.1016/j.bbabio.2011.09.022

Lan, X., Han, X., Li, Q., & Wang, J. (2017). (-)-Epicatechin, a Natural Flavonoid Compound,

Protects Astrocytes Against Hemoglobin Toxicity via Nrf2 and AP-1 Signaling

Pathways. *Molecular Neurobiology*, *54*(10), 7898–7907. PubMed.

https://doi.org/10.1007/s12035-016-0271-y

Lee, I., Hüttemann, M., Kruger, A., Bollig-Fischer, A., & Malek, M. H. (2015). (-)-

Epicatechin combined with 8 weeks of treadmill exercise is associated with increased

angiogenic and mitochondrial signaling in mice. *Frontiers in Pharmacology*.

https://doi.org/10.3389/fphar.2015.00043

Lesniewski, L. A., Zigler, M. C., Durrant, J. R., Donato, A. J., & Seals, D. R. (2012).

Sustained activation of AMPK ameliorates age-associated vascular endothelial

dysfunction via a nitric oxide-independent mechanism. *Mechanisms of Ageing and*

Development. https://doi.org/10.1016/j.mad.2012.03.011

688 Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using

689   real-time quantitative PCR and the 2-ΔΔCT method. *Methods*.

690   https://doi.org/10.1006/meth.2001.1262

691 Loke, W. M., Hodgson, J. M., Proudfoot, J. M., McKinley, A. J., Puddey, I. B., & Croft, K.

692   D. (2008). Pure dietary flavonoids quercetin and (-)-epicatechin augment nitric oxide

693   products and reduce endothelin-1 acutely in healthy men. *American Journal of*

694   *Clinical Nutrition*. https://doi.org/10.1093/ajcn/88.4.1018

695 Miller, M. W., Knaub, L. A., Olivera-Fragoso, L. F., Keller, A. C., Balasubramaniam, V.,

696   Watson, P. A., & Reusch, J. E. B. (2013). Nitric oxide regulates vascular adaptive

697   mitochondrial dynamics. *American Journal of Physiology - Heart and Circulatory*

698   *Physiology*. https://doi.org/10.1152/ajpheart.00987.2012

699 Mookerjee, S. A., & Brand, M. D. (2015). Measurement and analysis of extracellular acid

700   production to determine glycolytic rate. *Journal of Visualized Experiments*.

701   https://doi.org/10.3791/53464

702 Moreno-Ulloa, A., Mendez-Luna, D., Beltran-Partida, E., Castillo, C., Guevara, G., Ramirez-

703   Sanchez, I., Correa-Basurto, J., Ceballos, G., & Villarreal, F. (2015). The effects of (-

704   )-epicatechin on endothelial cells involve the G protein-coupled estrogen receptor

705   (GPER). *Pharmacological Research*. https://doi.org/10.1016/j.phrs.2015.08.014

706 Moreno-Ulloa, A., Nogueira, L., Rodriguez, A., Barboza, J., Hogan, M. C., Ceballos, G.,

707   Villarreal, F., & Ramirez-Sanchez, I. (2015). Recovery of Indicators of Mitochondrial

708   Biogenesis, Oxidative Stress, and Aging With (-)-Epicatechin in Senile Mice.

709   *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*.

710   https://doi.org/10.1093/gerona/glu131

Murase, T., Misawa, K., Haramizu, S., & Hase, T. (2009). Catechin-induced activation of the LKB1/AMP-activated protein kinase pathway. *Biochemical Pharmacology*. https://doi.org/10.1016/j.bcp.2009.03.021

Musicki, B., Kramer, M. F., Becker, R. E., & Burnett, A. L. (2005). Age-Related Changes in Phosphorylation of Endothelial Nitric Oxide Synthase in the Rat Penis. *The Journal of Sexual Medicine*, *2*(3), 347–357. https://doi.org/10.1111/j.1743-6109.2005.20349.x

Ottaviani, J. I., Borges, G., Momma, T. Y., Spencer, J. P. E., Keen, C. L., Crozier, A., & Schroeter, H. (2016). The metabolome of [2-(14)C](-)-epicatechin in humans: Implications for the assessment of efficacy, safety, and mechanisms of action of polyphenolic bioactives. *Scientific Reports*, *6*, 29034. https://doi.org/10.1038/srep29034

Panneerselvam, M., Ali, S. S., Finley, J. C., Kellerhals, S. E., Migita, M. Y., Head, B. P., Patel, P. M., Roth, D. M., & Patel, H. H. (2013). Epicatechin regulation of mitochondrial structure and function is opioid receptor dependent. *Molecular Nutrition & Food Research*, *57*(6), 1007–1014. https://doi.org/10.1002/mnfr.201300026

Papadimitriou, A., Peixoto, E. B. M. I., Silva, K. C., Lopes de Faria, J. M., & Lopes de Faria, J. B. (2014). Increase in AMPK brought about by cocoa is renoprotective in experimental diabetes mellitus by reducing NOX4/TGFβ-1 signaling. *The Journal of Nutritional Biochemistry*, *25*(7), 773–784. https://doi.org/10.1016/j.jnutbio.2014.03.010

Park, S. H., Kwon, O. S., Park, S.-Y., Weavil, J. C., Andtbacka, R. H. I., Hyngstrom, J. R., Reese, V., & Richardson, R. S. (2018). Vascular mitochondrial respiratory function: The impact of advancing age. *American Journal of Physiology. Heart and Circulatory*

Physiology, *315*(6), H1660–H1669. PubMed.

https://doi.org/10.1152/ajpheart.00324.2018

Park, S.-Y., Kwon, O. S., Andtbacka, R. H. I., Hyngstrom, J. R., Reese, V., Murphy, M. P., &

Richardson, R. S. (2018). Age-related endothelial dysfunction in human skeletal

muscle feed arteries: The role of free radicals derived from mitochondria in the

vasculature. *Acta Physiologica*, *222*(1), e12893. https://doi.org/10.1111/apha.12893

Park, S.-Y., Pekas, E. J., Headid, R. J., Son, W.-M., Wooden, T. K., Song, J., Layec, G.,

Yadav, S. K., Mishra, P. K., & Pipinos, I. I. (2020). Acute mitochondrial antioxidant

intake improves endothelial function, antioxidant enzyme activity, and exercise

tolerance in patients with peripheral artery disease. *American Journal of Physiology-*

Heart and Circulatory Physiology, *319*(2), H456–H467.

https://doi.org/10.1152/ajpheart.00235.2020

Ramirez-Sanchez, I., Aguilar, H., Ceballos, G., & Villarreal, F. (2012). (-)-Epicatechin-

induced calcium independent eNOS activation: Roles of HSP90 and AKT. *Molecular*

and Cellular Biochemistry. https://doi.org/10.1007/s11010-012-1405-9

Ramirez-Sanchez, I., De Los Santos, S., Gonzalez-Basurto, S., Canto, P., Mendoza-Lorenzo,

P., Palma-Flores, C., Ceballos-Reyes, G., Villarreal, F., Zentella-Dehesa, A., & Coral-

Vazquez, R. (2014). (-)-Epicatechin improves mitochondrial-related protein levels

and ameliorates oxidative stress in dystrophic δ-sarcoglycan null mouse striated

muscle. *FEBS Journal*. https://doi.org/10.1111/febs.13098

Ramirez-Sanchez, I., Mansour, C., Navarrete-Yañez, V., Ayala-Hernandez, M., Guevara, G.,

Castillo, C., Loredo, M., Bustamante, M., Ceballos, G., & Villarreal, F. J. (2018). (−)-

Epicatechin induced reversal of endothelial cell aging and improved vascular

function: Underlying mechanisms. *Food and Function*.

https://doi.org/10.1039/c8fo00483h

Ramirez-Sanchez, I., Maya, L., Ceballos, G., & Villarreal, F. (2010). (-)-Epicatechin

activation of endothelial cell endothelial nitric oxide synthase, nitric oxide, and

related signaling pathways. *Hypertension*, *55(6)*, 1398–1405.

https://doi.org/10.1161/HYPERTENSIONAHA.109.147892

Ramirez-Sanchez, I., Maya, L., Ceballos, G., & Villarreal, F. (2011). (-)-Epicatechin induces

calcium and translocation independent eNOS activation in arterial endothelial cells.

American Journal of Physiology - Cell Physiology.

https://doi.org/10.1152/ajpcell.00406.2010

Ramírez-Sánchez, I., Rodríguez, A., Moreno-Ulloa, A., Ceballos, G., & Villarreal, F. (2016).

(-)-Epicatechin-induced recovery of mitochondria from simulated diabetes: Potential

role of endothelial nitric oxide synthase. *Diabetes and Vascular Disease Research*.

https://doi.org/10.1177/1479164115620982

Robledinos-Antón, N., Fernández-Ginés, R., Manda, G., & Cuadrado, A. (2019). Activators

and Inhibitors of NRF2: A Review of Their Potential for Clinical Development.

Oxidative Medicine and Cellular Longevity, *2019*, 9372182.

https://doi.org/10.1155/2019/9372182

Rowley, T. J., Bitner, B. F., Ray, J. D., Lathen, D. R., Smithson, A. T., Dallon, B. W.,

Plowman, C. J., Bikman, B. T., Hansen, J. M., Dorenkott, M. R., Goodrich, K. M.,

Ye, L., O'Keefe, S. F., Neilson, A. P., & Tessem, J. S. (2017). Monomeric cocoa

catechins enhance β-cell function by increasing mitochondrial respiration. *Journal of*

Nutritional Biochemistry. https://doi.org/10.1016/j.jnutbio.2017.07.015

Schnorr, O., Brossette, T., Momma, T. Y., Kleinbongard, P., Keen, C. L., Schroeter, H., &

Sies, H. (2008). Cocoa flavanols lower vascular arginase activity in human

endothelial cells in vitro and in erythrocytes in vivo. *Archives of Biochemistry and*

Biophysics. https://doi.org/10.1016/j.abb.2008.02.040

Schroeter, H., Heiss, C., Balzer, J., Kleinbongard, P., Keen, C. L., Hollenberg, N. K., Sies,

H., Kwik-Uribe, C., Schmitz, H. H., & Kelm, M. (2006). (-)-Epicatechin mediates

beneficial effects of flavanol-rich cocoa on vascular function in humans. *Proceedings*

of the National Academy of Sciences of the United States of America, *103(4)*, 1024–

1029. https://doi.org/10.1073/pnas.0510168103

Shi, L., Hao, Z., Zhang, S., Wei, M., Lu, B., Wang, Z., & Ji, L. (2018). Baicalein and baicalin

alleviate acetaminophen-induced liver injury by activating Nrf2 antioxidative

pathway: The involvement of ERK1/2 and PKC. *Biochemical Pharmacology*, *150*, 9–

23. https://doi.org/10.1016/j.bcp.2018.01.026

Si, H., Fu, Z., Babu, P. V. A., Zhen, W., LeRoith, T., Meaney, M. P., Voelker, K. A., Jia, Z.,

Grange, R. W., & Liu, D. (2011). Dietary epicatechin promotes survival of obese

diabetic mice and Drosophila melanogaster. *Journal of Nutrition*.

https://doi.org/10.3945/jn.110.134270

Sun, D., Huang, A., Yan, E. H., Wu, Z., Yan, C., Kaminski, P. M., Oury, T. D., Wolin, M. S.,

& Kaley, G. (2004). Reduced release of nitric oxide to shear stress in mesenteric

arteries of aged rats. *American Journal of Physiology - Heart and Circulatory*

Physiology. https://doi.org/10.1152/ajpheart.00854.2003

Suwaidi, J. A., Hamasaki, S., Higano, S. T., Nishimura, R. A., Holmes, D. R., & Lerman, A.

(2000). Long-Term Follow-Up of Patients With Mild Coronary Artery Disease and

Endothelial Dysfunction. *Circulation*, *101*(9), 948–954.

https://doi.org/10.1161/01.CIR.101.9.948

Taub, P. R., Ramirez-Sanchez, I., Ciaraldi, T. P., Perkins, G., Murphy, A. N., Naviaux, R.,

Hogan, M., Maisel, A. S., Henry, R. R., Ceballos, G., & Villarreal, F. (2012).

Alterations in Skeletal Muscle Indicators of Mitochondrial Structure and Biogenesis

in Patients with Type 2 Diabetes and Heart Failure: Effects of Epicatechin Rich

Cocoa. *Clinical and Translational Science*. https://doi.org/10.1111/j.1752-

8062.2011.00357.x

Taub, P. R., Ramirez-Sanchez, I., Patel, M., Higginbotham, E., Moreno-Ulloa, A., Román-

Pintos, L. M., Phillips, P., Perkins, G., Ceballos, G., & Villarreal, F. (2016).

Beneficial effects of dark chocolate on exercise capacity in sedentary subjects:

Underlying mechanisms. A double blind, randomized, placebo controlled trial. *Food*

and Function. https://doi.org/10.1039/c6fo00611f

Ungvari, Z., Labinskyy, N., Gupte, S., Chander, P. N., Edwards, J. G., & Csiszar, A. (2008).

Dysregulation of mitochondrial biogenesis in vascular endothelial and smooth muscle

cells of aged rats. *American Journal of Physiology - Heart and Circulatory*

Physiology. https://doi.org/10.1152/ajpheart.00012.2008

Van Der Loo, B., Labugger, R., Skepper, J. N., Bachschmid, M., Kilo, J., Powell, J. M.,

Palacios-Callender, M., Erusalimsky, J. D., Quaschning, T., Malinski, T., Gygi, D.,

Ullrich, V., & Lüscher, T. F. (2000). Enhanced peroxynitrite formation is associated

with vascular aging. *Journal of Experimental Medicine*.

https://doi.org/10.1084/jem.192.12.1731

Yang, K., & Chan, C. B. (2018). Epicatechin potentiation of glucose-stimulated insulin

secretion in INS-1 cells is not dependent on its antioxidant activity. *Acta*

Pharmacologica Sinica. https://doi.org/10.1038/aps.2017.174

Zielonka, J., Joseph, J., Sikora, A., Hardy, M., Ouari, O., Vasquez-Vivar, J., Cheng, G.,

Lopez, M., & Kalyanaraman, B. (2017). Mitochondria-Targeted

Triphenylphosphonium-Based Compounds: Syntheses, Mechanisms of Action, and

Therapeutic and Diagnostic Applications. *Chemical Reviews*, *117*(15), 10043–10120.

PubMed. https://doi.org/10.1021/acs.chemrev.7b00042

834
835