A MademoiseLLE domain binding platform
links the key RNA transporter to endosomes

Senthil-Kumar Devan[1, $], Stephan Schott-Verdugo[2, 3, $], Kira Müntjes[1], Lilli Bismar[1], Jens Reiners[4], Eymen Hachani[5], Lutz Schmitt[5], Astrid Höppner[4], Sander HJ Smits[4, 5, *], Holger Gohlke[2, 3, *], Michael Feldbrügge[1, *]

$ shared first authorship

* shared corresponding authorship

[1] Institute of Microbiology, Heinrich Heine University Düsseldorf, Cluster of Excellence on Plant Sciences, 40204 Düsseldorf, Germany

[2] John von Neumann Institute for Computing (NIC), Jülich Supercomputing Centre (JSC), Institute of Biological Information Processing (IBI-7: Structural Bioinformatics), and Institute of Bio- and Geosciences (IBG-4: Bioinformatics), Forschungszentrum Jülich GmbH, Wilhelm-Johnen-Str., 52425 Jülich, Germany

[3] Institute for Pharmaceutical and Medicinal Chemistry, Heinrich Heine University Düsseldorf, Universitätsstraße 1, 40225 Düsseldorf, Germany

[4] Center for Structural Studies, Heinrich Heine University Düsseldorf, 40204 Düsseldorf, Germany

[5] Institute of Biochemistry I, Heinrich Heine University Düsseldorf, 40204 Düsseldorf, Germany

Running title: A novel MLLE platform

Key words: Endosome / PAM2 / Poly(A)-binding protein / RNA transport / *Ustilago maydis*

[*] **Corresponding authors:**

Dr. Sander HJ Smits

Institute of Biochemistry,

Heinrich Heine University Düsseldorf, 40204 Düsseldorf, Germany

Phone: +49 (211) 81-12647

sander.smits@hhu.de

Dr. Holger Gohlke

Institute for Pharmaceutical and Medicinal Chemistry,

Heinrich Heine University Düsseldorf, 40204 Düsseldorf, Germany

Phone: +49 (211) 81-13662

&

John von Neumann Institute for Computing (NIC), Jülich Supercomputing Centre (JSC), Institute of Biological Information Processing (IBI-7: Structural Bioinformatics), and Institute of Bio- and Geosciences (IBG-4: Bioinformatics), Forschungszentrum Jülich GmbH, 52425 Jülich, Germany

Phone: +49 (2461) 61-85550

gohlke@uni-duesseldorf.de & h.gohlke@fz-juelich.de

Dr. Michael Feldbrügge

Institute of Microbiology,

Bioeconomy Science Center, Cluster of Excellence on Plant Sciences

Heinrich Heine University Düsseldorf, 40204 Düsseldorf, Germany

Phone: +49 (211) 81-15475

feldbrue@hhu.de

Abstract

Spatiotemporal expression is mostly achieved by transport and translation of mRNAs at defined

subcellular sites. An emerging mechanism mediating mRNA trafficking is microtubule-

dependent co-transport of mRNAs on shuttling endosomes. Although progress has been made

in identifying various components of the endosomal mRNA transport machinery, a mechanistic

understanding of how these RNA-binding proteins are connected to endosomes is still lacking.

Here, we demonstrate that a flexible MademoiseLLE (MLLE) domain platform within Rrm4

of *Ustilago maydis* is crucial for endosomal attachment. Our structure/function analysis

uncovered three MLLE domains at the C-terminus of Rrm4 with a functionally defined

hierarchy. MLLE3 recognizes two PAM2-like sequences of the adaptor protein Upa1 and is

essential for endosomal shuttling of Rrm4. MLLE1 and MLLE2 are most likely accessory

domains that exhibit a variant binding mode for interaction with currently unknown partners.

Thus, endosomal attachment of the mRNA transporter is orchestrated by a sophisticated MLLE

domain binding platform.

Introduction

mRNA localisation and local translation are essential for spatiotemporal control of protein
expression. An important mechanism to achieve localised translation is the active transport of
mRNAs along the cytoskeleton (Das *et al.*, 2021; Fernandopulle *et al.*, 2021; Mofatteh &
Bullock, 2017). Mainly, long-distance transport of mRNA is mediated by motor-dependent
movement along microtubules. Important carriers are transport endosomes that move
messenger ribonucleoprotein complexes (mRNPs), consisting of RNA-binding proteins and
cargo mRNAs on their cytoplasmic surface (Mofatteh & Bullock, 2017; Müntjes *et al.*, 2021;
Niessing *et al.*, 2018). This process is evolutionarily conserved in fungi, plants, and animals
(Baumann *et al.*, 2012; Cioni *et al.*, 2019; Tian *et al.*, 2020a; Liao *et al.*, 2019; Müntjes *et al.*.,
2021; Quentin *et al.*, 2021; Schuhmacher *et al.*, 2021). In endosperm cells of developing rice
seeds, cargo mRNAs are transported to cortical ER by the action of the two RNA recognition
motif (RRM)-containing proteins RBP-P and RBP-L. These form a quaternary complex with
membrane trafficking factor NSF (*N*-ethylmaleimide-sensitive factor) and small GTPase Rab5a
on the endosomal surface (Tian *et al.*, 2020b). In neurons, mRNA transport has been linked to
early and late endosomes as well as lysosomal vesicles. Especially, local translation of mRNAs
encoding mitochondrial proteins on the surface of late endosomes is needed for mitochondrial
function. Importantly, this trafficking process has been associated with the neuronal Charcot
Marie-Tooth disease (Cioni *et al.*, 2019). Annexin 11, a factor implicated in amyotrophic lateral
sclerosis (ALS), was found as an mRNP linker on motile lysosomal vesicles (Liao *et al.*, 2019).
Recently, the five-membered FERRY complex was identified connecting mRNAs encoding
mitochondrial proteins to neuronal endosomes by interaction with the active form of Rab5
(Quentin *et al.*, 2021; Schuhmacher *et al.*, 2021).

Among the best-studied examples for membrane-coupled mRNA transport is the
endosomal mRNA transport in the corn pathogen *Ustilago maydis* (Béthune *et al.*, 2019; Haag
et al., 2015; Müntjes *et al.*, 2021). Extensive peripheral movement of mRNAs is needed for
efficient unipolar growth of infectious hyphae. These hyphae grow highly polarised by
expanding at the growing tip and inserting regularly spaced septa at the basal pole. Loss of
mRNA distribution causes aberrant bipolar growth (Baumann *et al.*, 2012; Becht *et al.*, 2006;
Pohlmann *et al.*, 2015). Key vehicles of cargo mRNAs are Rab5a-positive endosomes that
shuttle along microtubules by the concerted action of plus-end directed kinesin-3 and minus-
end directed cytoplasmic dynein (Baumann *et al.*, 2012). Important cargo mRNAs are, for
example, all four septin mRNAs. Their local translation during transport is needed to form
heteromeric septin complexes on the surface of transport endosomes. Endosomes deliver these

complexes to the hyphal tip forming a defined gradient of septin filaments at the growing pole

(Baumann *et al.*, 2014; Olgeiser *et al.*, 2019; Zander *et al.*, 2016).

Rrm4 is the key RNA-binding protein of the transport process that recognises defined sets

of cargo mRNAs via its three N-terminal RRMs (Fig. 1A; Olgeiser *et al.*, 2019). Rrm4 and

bound cargo mRNAs are linked to endosomes by Upa1, containing a FYVE zinc finger for

interaction with PI$_3$P lipids (phosphatidylinositol 3-phosphate; Fig 1A; Pohlmann *et al.*, 2015;

Stenmark *et al.*, 2002). The adaptor protein contains a PAM2 motif (poly [A]-binding protein

interacting motif 2; Albrecht & Lengauer, 2004; Jinek *et al.*, 2010; Kozlov *et al.*, 2004) and

two PAM2 like (PAM2L) sequences. These motifs are crucial for interaction with

MademoiseLLE (MLLE) domains of the poly(A)-binding protein Pab1 and Rrm4, respectively

(Fig. 1A; Pohlmann *et al.*, 2015).

The MLLE domain was first identified as a conserved domain at the C-terminus of the

human cytoplasmic poly(A)-binding protein 1 (PABP1C; Mangus *et al.*, 1998; Mangus *et al.*,

2003). Solution and crystal structures of PABC domains from PABP1C and ubiquitin ligase

UBR5 showed that they are structurally conserved (Deo *et al.*, 2001; Kozlov *et al.*, 2001). The

domain is about 70 amino acids in length and consists of five bundled α-helices. Interaction

with the PAM2 binding motif (consensus sequence xxLNxxAxEFxP) is characterised by the

central α-helix 3 with the sequence KITG<u>MLLE</u> and mediated by two adjacent hydrophobic

pockets (Xie *et al.*, 2014), with the binding of the Phe residue of the PAM2 motif being the

major determinant for this interaction (Kozlov & Gehring, 2010). Besides human PABPC1,

there are currently only two additional proteins with MLLE domains described: the ubiquitin

ligase UBR5 functioning, for example, during microRNA-mediated gene silencing (Su *et al.*,

2011) and Rrm4-type RNA-binding proteins from fungi (Fig. 1B; Müller *et al.*, 2019).

Mutations in the C-terminal MLLE domain of Rrm4 result in the loss of Rrm4 motility,

suggesting that the link to endosomes is disrupted (Becht *et al.*, 2006). Consistently, the C-

terminus of Rrm4 recognizes the PAM2L sequence of the adaptor protein Upa1 (Pohlmann *et*

al., 2015), suggesting that the interaction of MLLE domains with PAM2L sequences is

responsible for its endosome association. In this study, we combine structural biology with

fungal genetics to demonstrate that the C-terminal half of Rrm4 has three divergent MLLE

domains with a flexible arrangement that contribute differentially to the endosomal attachment.

Results

Iterative structural modeling predicts three MLLE domains at the C-terminus of Rrm4

To generate structural models of the MLLE domains present in Rrm4, we focused on the C-
terminal part of the protein (residues 421 to 792). This excluded the three N-terminal RRMs
but included the previously predicted two C-terminal MLLE domains (Fig. 1A-B; Müller *et al.*,
2019). Subjecting this Rrm4 sequence region to iterative comparative modeling with TopModel
(Fig. 1C; Mulnaes *et al.*, 2020a) revealed, as expected, the previously identified two regions
with homology for MLLE domains located at residues 571-629 and 712-791 (denoted MLLE2
and MLLE3; Fig. 1C; Müller *et al.*, 2019). Unexpectedly, using the TopModel workflow with
its efficient template selection capabilities (Mulnaes *et al.*, 2020a), we identified an additional
de novo predicted MLLE domain located at residues 451-529 (denoted MLLE1; Fig. 1B-C; Fig.
EV1A). Although the sequence identity between templates and their respective Rrm4 sequence
stretches is only 17 to 32% (Fig. 1B, Fig. EV1A), the generated MLLE domain models have a
high predicted local structural quality, as assessed by TopScore (Fig. 1C; Mulnaes & Gohlke,
2018). The generated models were also verified by the current deep neural network modeling
approaches AlphaFold2 and RoseTTAFold (Fig. EV1B; Baek *et al.*, 2021; Jumper *et al.*, 2021),
further indicating that the C-terminal half of Rrm4 has three MLLE domains instead of the
previously identified two. All of these MLLE domains might be relevant for the interaction
with Upa1.

X-ray analysis of the second MLLE domain confirms the predicted structural models

To verify the structural models further, we expressed and purified an N-terminally truncated
version of the Rrm4 carrying the three MLLE domains in *Escherichia coli* (Fig. EV2A; version
H-Rrm4-NT4 carrying an N-terminal hexa-histidine tag; Pohlmann *et al.*, 2015; Materials and
methods). Size exclusion chromatography combined with multiangle light scattering (MALS)
indicates that the protein is homogenous and does not form aggregates (Fig. EV2B-C). We thus
set out to crystallize the protein for X-ray diffraction analysis (see Material and Methods).
Testing 2016 different conditions, crystals were only obtained in individual cases after at least
7 days of incubation. A complete dataset was collected from a single crystal diffracting to 2.6 Å
resolution and a $P4_32_12$ symmetry. Data and refinement statistics are given in Appendix Table
S2. Surprisingly, the unit cell dimensions were small and, with a Matthews coefficient assuming
50% solvent content, only 128 amino acids would fit into the asymmetric unit of the crystal.
Hence, the unit cell has an insufficient size to cover H-Rrm4-NT4, which contains 380 amino

acids. Using the predicted models of MLLE1-3 as templates for molecular replacement, only

MLLE2 gave a clear solution, showing after refinement that two copies of MLLE2 (residues

567-630) were present in the asymmetric unit. For comparison, previously, two copies of the

MLLE domain in the asymmetric unit were reported in crystals of MLLE of UBR5 (Munoz-

Escobar *et al.*, 2015). The structural data indicates that the protein was truncated from both

termini during crystallization, resulting in a shortened version of the H-Rrm4-NT4 protein that

formed stable crystals (see Material and methods). Both MLLE2[Rrm4] copies adopt the same

overall fold (Fig. 2A) as seen by the RMSD of 0.29 Å over 59 C-alpha atoms. The MLLE2[Rrm4]

crystal structure displays high similarity with the MLLE domain of the ubiquitin ligase UBR5

(MLLE[UBR5]; Munoz-Escobar *et al.*, 2015; PDB code 3NTW, RMSD of 0.97 Å over 56 amino

acids) and the MLLE domain of PABPC1 (Kozlov *et al.*, 2010; PDB code 3KUS, RMSD of

1.34 Å over 61 amino acids). The MLLE2[Rrm4] domain consists of four helices (designated α2 -

5; Fig. 2A) that are arranged as a right-handed superhelix similar to MLLE[UBR5]. In comparison

to the MLLE domain of PABPC1, the first short helix is absent in both MLLE2[Rrm4] and

MLLE[UBR5] structures.

When comparing the obtained crystal structure with the MLLE2 [Rrm4] model generated by

TopModel, the average RMSD is 0.62 Å over the backbone atoms, close to the uncertainty of

the atomic coordinates of the experimental structure (Fig. 2B). Importantly, this confirms our

structural model of MLLE2[Rrm4] and strongly suggests that the modeled MLLE1[Rrm4] and

MLLE3[Rrm4] domains should be of equally high quality.

We compared the models obtained of MLLE1-3[Rrm4] with the known structure of the human

PABPC1 focusing on the well described PAM2 peptide binding pocket. This revealed that

MLLE3[Rrm4] maintains a characteristic Gly residue at position 736 that binds the conserved Phe

residue of the PAM2 motifs, a major binding determinant in PABPC1 and UBR5 (Kozlov &

Gehring, 2010; Munoz-Escobar *et al.*, 2015; Fig. 2C). However, the binding interfaces of

MLLE1[Rrm4] and MLLE2[Rrm4] are altered compared to the 'canonical' binding site in PABPC1

and UBR5 (Fig. 2C). Instead of Gly, MLLE1[Rrm4] and MLLE2[Rrm4] have a Ser and Arg in the

corresponding positions 471 and 573. The notion that MLLE1[Rrm4] and MLLE2[Rrm4] may differ

from canonical MLLE domains is also supported by the lower sequence identity of MLLE1[Rrm4]

and MLLE2[Rrm4] when compared to previously characterized MLLE domains (Fig. EV1A). In

summary, structural modeling revealed the presence of three MLLE domains at the C-terminus

of Rrm4. Furthermore, the structure of the MLLE2[Rrm4] domain was successfully verified by X-

ray crystallographic analysis. MLLE1[Rrm4] and MLLE2[Rrm4] are divergent in the key region of

PAM2 binding, suggesting that these domains might employ a different binding mode or show

a different binding specificity.

The MLLE domains of Rrm4 form a binding platform with flexible arrangement of the

individual domains

To study the relative arrangement of all three MLLE^{Rrm4} domains and the orientation to the N-

terminal RRMs, we performed Small-Angle X-ray Scattering (SAXS) experiments. We

expressed and purified H-Rrm4-NT4 as well as full-length protein with N-terminal GST fusion

(glutathione S-transferase; G-Rrm4) from *E. coli* (see Materials and methods). Primary data

analysis of the scattering curves (Tria *et al.*, 2015) revealed that both proteins are monomeric

and highly flexible in solution (Appendix Table S3; Fig. 2D-E; Fig. EV2D-E). To visualise the

different conformations, we performed an Ensemble Optimization Method (EOM) analysis for

both the G-Rrm4 and H-Rrm4-NT4 proteins (Fig 2D-E). We used our MLLE models and a

GST model (PDB entry: 1ua5) together with the protein sequence for G-Rrm4 as input, yielding

a distribution of different conformations of the protein in solution (representative models in Fig.

EV2E). One model of G-Rrm4, representing 25% of the population, revealed that the C-

terminal part containing MLLE1-3^{Rrm4} adopts an elongated and mainly unfolded but open

conformation (Fig. 2D; Fig. EV2D-E). The N-terminal part, containing RRM domains of the

GST fusion protein, adopts a more globular structure, indicating less flexibility within this

region (Fig. 2D; Fig. EV2D-E). Studying only the C-terminal part of Rrm4 revealed that the

most prominent model of this analysis (75 % of the population) has a nearly identical

conformation as the one selected for G-Rrm4 (Fig. 2D-E, Fig. EV2E). It suggests that the C-

terminal part of Rrm4-NT4 adopts a very similar orientation when expressed by itself. From

this analysis, one can deduce that the MLLE1-3^{Rrm4} domains form a C-terminal binding

platform with a flexible arrangement for multiple contact sites for binding partners. The RRM

domains and thus the RNA interaction site are clearly spatially separated from the protein

interaction platform.

The third MLLE is essential for interaction with PAM2 like sequences of Upa1

To evaluate the interaction capacity of MLLE1-3^{Rrm4}, we performed *in vitro* binding studies.

We expressed different deletion versions of Rrm4 as N-terminal GST fusions in *E. coli*. As a

control, we expressed an N-terminal GST fusion of the MLLE domain of Pab1 (Fig. 3A;

Materials and methods). To check the physical interaction with PAM2 and PAM2L sequences

of Upa1, we expressed 18 amino acid fragments (Fig. 3A, Fig. EV3B) as N-terminal hexa-

histidine-SUMO (HS) fusion proteins (see Materials and methods). In GST pull-down

experiments using GST fusion proteins as bait, G-Pab1-MLLE interacts with HS-PAM2 but

not with the HS-PAM2L motifs of Upa1 (Fig. EV3B, lane 2). Conversely, G-Rrm4-NT4

recognizes the two HS-PAM2L motifs of Upa1 but not the HS-PAM2 motif (Fig. EV3B, lane 3;

Pohlmann *et al.*, 2015). Interestingly, the interaction with both PAM2L motifs was lost when

MLLE3[Rrm4] was deleted (G-Rrm4-NT4-M3Δ; Fig. EV3B, lane 6), while constructs with

deletion of MLLE1[Rrm4] or MLLE2[Rrm4], or MLLE1[Rrm4] and MLLE2[Rrm4], still interacted with the

HS-PAM2L motifs of Upa1 (Fig. EV3B, lane 4,5 and 7).

To obtain quantitative data on the protein/peptide interactions, we performed isothermal

titration calorimetry (ITC) experiments with purified proteins (Fig. EV3C) using synthetic

peptides with a length of 18 amino acids (PAM2[Upa1], PAM2L1[Upa1], and PAM2L2[Upa1]; Fig. 3A).

From the curves, the binding constant K_D and the binding stoichiometry was calculated, which

in all cases indicated a 1:1 ratio between the G-Rrm4-NT4 protein and the binding partner.

Testing G-Pab1-MLLE with the peptides revealed a K_D of 14.6 μM for PAM2[Upa1] (Fig.

EV3D), which is within the range of observed K_D of 0.2 to 40 μM for known MLLE/PAM2

interactions like the MLLE domain of PABPC1 with various PAM2 sequences (Mattijssen *et*

al., 2021). Testing G-Pab1-MLLE with PAM2L1[Upa1] and PAM2L2[Upa1] peptides, no indication

for binding was observed. PAM2L sequences are rich in acidic residues and exhibit a different

FxxP spacing than the canonical FXP sequence of PAM2 sequences in the core region (Fig.

3A, EV3D; see Discussion). The observed binding behaviour indicates a clear binding

specificity differentiating PAM2 and PAM2L peptides. This is in line with the previously

published GST pull-down experiments (Pohlmann *et al.*, 2015).

In comparison, testing G-Rrm4-NT4 with the peptides revealed a K_D of 14.9 μM for

PAM2L1[Upa1] and 5.1 μM for PAM2L2[Upa1] and no binding to PAM2[Upa1] (Fig. 3B). This suggests

a similar affinity when compared to the interactions of MLLE[Pab1] with PAM2 and demonstrates

the high sequence specificity of the MLLE domains to their respective PAM2L sequences (see

Discussion).

Analysing G-Rrm4-NT4-M3Δ with a deletion of MLLE3[Rrm4] revealed that binding to

PAM2L1[Upa1] and PAM2L2[Upa1] was no longer detectable (Fig. 3C). This is in line with

observations from the GST pull-down experiments (Fig. EV3B). This suggests that MLLE3[Rrm4]

is essential for binding. Testing G-Rrm4-NT4 versions carrying deletions in MLLE1[Rrm4] and

MLLE2[Rrm4] revealed no difference in binding affinity (Fig. EV3E-F). Even testing G-Rrm4-

NT4 with both MLLE domains deleted revealed a binding affinity in the same range as the

wildtype version containing all three MLLEs (Fig. 3D). We conclude that (i) MLLE3[Rrm4] is

vital for recognizing PAM2L sequences with a higher affinity to PAM2L2 and (ii) neither

MLLE1[Rrm4] nor MLLE2[Rrm4] contribute to the binding of PAM2L or PAM2 motifs. This is

consistent with our structural analysis revealing a differences in the binding site for these MLLE

domains (see Discussion).

The third MLLE domain of Rrm4 is essential for its function

To address how the different MLLE domains contribute to the biological function of Rrm4, we

generated *U. maydis* strains carrying deletions in the respective domains of Rrm4 (Fig. 4A). As

genetic background, we used laboratory strain AB33, expressing the heteromeric master

transcription factor of hyphal growth (bE/bW) under control of the nitrate inducible promoter

P*nar1*. Thereby, polar hyphal growth can be elicited efficiently and in a highly reproducible

fashion by changing the nitrogen source (Fig. 4B, top; Brachmann *et al.*, 2001). To investigate

dynamic endosomal transport, we used strains expressing functional C-terminal fusion Upa1-

Gfp and Rrm4-Kat (fusion with enhanced version of the green fluorescent protein, Clontech,

and monomeric version of red fluorescent protein mKate2, respectively; Müntjes *et al.*, 2020;

Pohlmann *et al.*, 2015; see Materials and methods).

The resulting hyphae grew with a defined axis of polarity, i.e., they expanded at the hyphal

tip and inserted basal septa leading to the formation of empty sections (Fig. 4B-C). Loss of

Rrm4 (*rrm4Δ* strain) caused the formation of hyphae growing at both ends, characteristic for

aberrant bipolar growth (Fig. 4B-C; Pohlmann *et al.*, 2015). Testing Rrm4-Kat versions

carrying deletions of MLLE1[Rrm4] or MLLE2[Rrm4] did not cause alterations in unipolar growth

(Fig. 4B-C). Furthermore, endosomal shuttling and co-localisation were indistinguishable from

wildtype (Fig. 4D-E). Also, the number of Rrm4-Kat-positive endosomes (number of signals /

10 μm, Fig. EV4A), velocity, and processivity (Fig. EV4B-C) were comparable to wildtype.

Hence, the first two MLLE domains were dispensable for polar growth and endosomal shuttling

under optimal growth conditions. Since the deletion of the first two MLLEs did not substantially

alter the function of Rrm4, we can infer that the deletion neither affects the overall structure of

the protein nor interfere with other domains like the RNA-binding domain of the protein. This

supports the conclusions of our biochemical experiments (see above).

Importantly, testing strains expressing Rrm4-Kat with deletion of the third MLLE domain

revealed a loss-of-function phenotype similar to *rrm4Δ* strains. The amount of bipolar hyphae

was comparable to *rrm4Δ* strains (Fig. 4B-C; Becht *et al.*, 2006). We observed drastic alteration

in shuttling, and Rrm4 aggregates did not co-localise with motile Upa1-positive signals (Fig.

4D-E). While the Rrm4 signals were static (Fig. 4D-E; Fig. EV4A-C) the amount of motile

Upa1-Gfp positive endosomes, their velocity, and their processivity were not affected (Fig.

EV4A-C, summarised in Fig. EV4D). This is consistent with previous results showing that the

third MLLE domain is important for the movement of Rrm4 and that endosomal shuttling of

Upa1 is not affected if Rrm4 is missing (Baumann *et al.*, 2012; Becht *et al.*, 2006). To conclude,

MLLE3^{Rrm4} is an essential domain for Rrm4 attachment to endosomes in contrast to MLLE1

and MLLE2^{Rrm4}.

The second MLLE domain plays accessory roles in endosomal Rrm4 attachment

To investigate the biological role of MLLE1^{Rrm4} and MLLE2^{Rrm4} in more detail, we generated

strains expressing Rrm4-M12Δ-Kat, lacking MLLE1^{Rrm4} and MLLE2^{Rrm4} domains, and tested

the influence on hyphal growth. Unipolar growth was not disturbed (Fig. 5A-B). To challenge

endosomal attachment of Rrm4, we expressed Upa1-Gfp versions carrying mutations in

PAM2L motif 1 or 2 as well as in both motifs; these motifs are important for Rrm4 interaction

(Fig. 4A; Pohlmann *et al.*, 2015). Strains expressing Rrm4-M12Δ-Kat in combination with

mutated PAM2L1 or PAM2L2 of Upa1 showed unipolar growth comparable to wildtype (Fig.

5A-B), indicating that MLLE1^{Rrm4} and MLLE2^{Rrm4} are dispensable for unipolar growth even

when the endosomal attachment was weakened by expressing Upa1 versions with mutated

PAM2L motifs (Fig. EV4D). When studying Upa1 mutated in both PAM2L motifs, we

observed an aberrant bipolar growth phenotype comparable to the *upa1Δ* strain (Fig. 5A-B).

This was expected, since the interaction of Rrm4 to endosomes is mediated by both PAM2L

motifs (Pohlmann *et al.*, 2015). Analysing Rrm4-M12Δ-Kat in this genetic background

revealed no additive phenotype (Fig. 5A-B). This reinforces that the interactions of PAM2L

motifs of Upa1 are the major determinants for endosomal attachment of Rrm4.

Next, we investigated endosomal shuttling. In strains expressing Rrm4-M12Δ-Kat missing

MLLE1^{Rrm4} and MLLE2^{Rrm4} endosomal shuttling was not disturbed (Fig. 5C). The amount of

motile Rrm4-M12Δ-Kat positive signals, their velocity, and their processivity were not affected

(Fig. EV5A). Like above, we challenged endosomal attachment of Rrm4 by expressing Upa1

versions with mutations in the PAM2L motifs. As expected, simultaneous mutation of both

PAM2L motifs of Upa1 resulted in a reduction in the number of Rrm4-Kat positive shuttling

endosomes (Fig. 5C, EV5E; Pohlmann *et al.*, 2015). In about 80% of hyphae, Rrm4-Kat found

to be aberrantly attached to microtubules (MT) when both PAM2L motifs are mutated (Fig.

5C-D). This is a characteristic feature for altered endosomal attachment of Rrm4 due to the

static accumulation of Rrm4-Kat, which can be also seen in *upa1Δ* strains (Fig. EV5C;

Jankowski *et al.*, 2019).

To analyse the influence of individual PAM2L motifs we determined the amount of hyphae

with aberrant microtubule staining in strains co-expressing Rrm4-Kat and an Upa1-Gfp version

with mutations of PAM2L sequence 1 or 2 (Fig. 5D). Mutations in PAM2L1 and PAM2L2

causes 8% and 19% of hyphae with aberrant MT staining, respectively. This indicates that the

interaction of PAM2L2 is more important for correct endosomal attachment of Rrm4. This is

consistent with our biochemical results demonstrating that MLLE3^{Rrm4} binds stronger to

PAM2L2 of Upa1 than to PAM2L1 (Fig. 3D).

Next, we investigated the association of Rrm4-M12Δ-Kat in strains expressing Upa1-

PAM2L1. Thus, the endosomal attachment was solely dependent on the interaction of

MLLE3^{Rrm4} with the PAM2L2 sequence of Upa1. We did observe 6% of hyphae with aberrant

MT staining (Fig. 5D; Fig. EV4D). This was comparable to strains expressing Rrm4-Kat

suggesting no clear difference (Fig. 5D). However, testing Upa1 with its PAM2L2 mutated,

leaving only PAM2L1 for interaction with Rrm4, we observed a clear increase in hyphae with

aberrant MT staining when comparing strains co-expressing Rrm4-M12Δ-Kat versus Rrm4-

Kat (52% versus 19%; Fig. 5D-E). Hence, the region covering MLLE1^{Rrm4} and MLLE2^{Rrm4} was

important for Rrm4 attachment. Finally, we tested individual deletions in MLLE1^{Rrm4} and

MLLE2^{Rrm4} in combination with mutated PAM2L2 in Upa1 to dissect the role of the different

MLLE domains. In the strains expressing Rrm4-Kat or Rrm4-M1Δ-Kat, the amount of hyphae

with aberrant MT staining was comparable (18% versus 11%, respectively; Fig. 5D). However,

strains expressing Rrm4-M2Δ-Kat exhibited an increased amount of hyphae with aberrant MT

staining that was comparable to Rrm4-M12Δ-kat (51% versus 52% respectively; Fig. 5D; Fig.

EV4D). To conclude, for MLLE1^{Rrm4} we were unable to assign a clear function yet. However,

MLLE2^{Rrm4} plays an accessory role in the endosomal attachment of Rrm4. In essence, the C-

terminus of Rrm4 contains three MLLE domains, with MLLE2^{Rrm4} fulfilling an accessory role

and MLLE3 having an essential function during the attachment of mRNPs to endosomes.

Discussion

Combining structural biology and biophysical techniques with fungal genetics and cell biology,

we addressed how mRNPs can be mechanistically linked to endosomes in the model fungus *U.*

maydis. Previously, it was found that the C-terminal MLLE domain of Rrm4 is needed for

shuttling (Becht *et al.*, 2006) and that the C-terminus of Rrm4 interacts with two PAM2L motifs

of Upa1 (Pohlmann *et al.*, 2015). Now, we demonstrate that the C-terminus of Rrm4 contains

not only two MLLE domains, but a sophisticated binding platform consisting of three MLLE

domains that collaborate for optimal functionality in linking the key RNA transporter to

endosomes. We disclose a strict hierarchy with main and accessory domain. The accessory

MLLE2 domain shows variations in the critical region of the predicted PAM2 binding pocket,

suggesting a novel mode of interaction with currently unknown interaction partners. To the best

of our knowledge, Rrm4 represents the first protein containing multiple MLLE domains to form

a binding platform. This interaction unit is essential for correct endosomal attachment and,

hence, mRNP trafficking.

The MLLE / PAM2 connection

The founding member of the MLLE domain family is present at the C-terminus of the poly(A)-

binding protein PABPC1. This domain interacts with PAM2 motifs of numerous interaction

partners such as GW182, eRF3, and the RNA-binding protein LARP4 functioning in

microRNA biology, translational termination, and posttranscriptional control, respectively

(Jinek *et al.*, 2010; Kozlov & Gehring, 2010; Yang *et al.*, 2011). Structural analysis revealed a

common mode of binding, where the Leu and particularly the Phe of the PAM2 consensus motif

xx<u>L</u>NxxAxE<u>F</u>xP (Fig. EV3A) are interacting with helix 2 and 3 as well as helix 3 and 5 of

MLLE domain, respectively (Kozlov & Gehring, 2010; Xie *et al.*, 2014). Indeed, the interaction

of MLLE with a hydrophobic amino acid is highly conserved, which in most cases is Phe with

a known exception in the variant PAM2w motif of LARP4 and LARP4A, where Trp is found

(Fig. EV3A; Grimm *et al.*, 2020; Yang *et al.*, 2011; Xie *et al.*, 2014).

Studying the MLLE domain-containing protein Rrm4, we discover that the protein has

three MLLE domains in its C-terminal half. MLLE3^{Rrm4} binds PAM2L motifs of Upa1 with a

K_D of 5 and 15 µM for PAM2L2^{Upa1} and -1, respectively. The binding affinities are in the same

range as described for other MLLE/PAM2 interactions: for example, the binding of

MLLE^{PABPC1} with PAM2^{LARP1}, PAM2$^{Tob2-125}$, PAM2^{LARP4} exhibit a K_D of 3.8, 16 and 22 µM,

respectively (Mattijssen *et al.*, 2021). Importantly, our biophysical assessment confirms the

exquisite binding specificity of MLLE^{Rrm4} that recognizes PAM2L1^{Upa1} and -2 but not

PAM2^{Upa1} version. PAM2L sequences contain a stretch of acidic amino acids in the N-terminal

half, and the spacing of FxxP in the core sequence is altered (Fig. EV3A). These variations

might account for the differential binding mode. Visual inspection of the potential PAM2L

binding region in the predicted model revealed that MLLE3^{Rrm4} contains a Gly at position 736

to sustain the binding of an aromatic residue of PAM2L as described for other MLLE domains

(see above). However, we were unable to uncover the structural basis for the sequence

specificity. Towards this end, future structural studies are required to provide detailed

information on the interaction of MLLE3[Rrm4] with PAM2L sequences.

Differential PAM2 binding has also been described for the MLLE[UBR5]. This MLLE domain

interacts with PAM2[PAIP] with an affinity of 3.4 µM (Lim *et al.*, 2006), whereas it binds a

PAM2L sequence (Fig. EV3A) in its own HECT domain with lower affinity (K_D of 50 µM).

The latter interaction has been implicated in the regulation of the HECT ligase activity (Munoz-

Escobar *et al.*, 2015). Interestingly, the PAM2L sequence within the HECT domain of UBR5

is highly similar to the PAM2L1 and -2 of Rrm4: (i) the sequences contain an acidic stretch N-

terminal to the conserved Phe (Fig. EV3A), (ii) the distance between Phe and Pro is two instead

of one amino acid, and (iii) the PAM2L sequence contains an additional bulky Tyr in close

vicinity to the Phe residue. Remarkably, MLLE[PABC1] does not recognise the PAM2L sequence

of UBR5 (Munoz-Escobar *et al.*, 2015). In essence, although the strong sequence specificity of

MLLE[Rrm4] and MLLE[Pab1] from *U. maydis* is, to the best of our knowledge, so far unique, we

hypothesise that differential PAM2 and PAM2L interactions are evolutionarily conserved and

might be more widespread than currently anticipated.

We observed also clear binding specificity for MLLE[Pab1] from *U. maydis* that interacts with

PAM2[Upa1] but not the PAM2L sequences from Upa1 (Fig. 6). MLLE[Pab1] binds with comparable

affinity to the PAM2[Upa1] (K_D of about 14 µM, Fig EV3D). Previously, we showed that mutations

in PAM2[Upa1] strongly decreased MLLE[Pab1] binding but did not interfere with the endosomal

shuttling of Pab1 (Pohlmann *et al.*, 2015). Thus, there might be other members of the endosomal

mRNPs interacting with Pab1 and stabilising its endosomal association. In fact, the dimerising

scaffold protein Upa2 of endosomal mRNA transport contains four PAM2 motifs offering eight

potential PAM2 motifs for interaction with Pab1 (Fig. 6). However, mutating all four PAM2

motifs did not interfere with endosomal mRNA transport, although interaction with MLLE[Pab1]

was lost (Jankowski *et al.*, 2019), confirming a potential redundancy. Consistently, mutations

in PAM2 of human LARP4B did not interfere with the function of stress granule recruitment,

also suggesting in this case the presence of additional factors (Grimm *et al.*, 2020).

Studying the other two MLLE domains of Rrm4 revealed that both lack the canonical Gly

for interactions with PAM2 or PAM2L sequences. MLLE1[Rrm4] and MLLE2[Rrm4] have Ser471

and Arg573 instead, respectively. Consistently, MLLE1[Rrm4] and MLLE2[Rrm4] do not bind PAM2

or PAM2L sequences. Thus, although the general fold of the MLLE domain is probably

conserved in MLLE1[Rrm4] and MLLE2[Rrm4], these domains most likely exhibit a different binding

specificity to their potential protein partner. Our detailed *in vivo* analysis revealed that MLLE2

carries out an accessory function for the correct attachment of Rrm4 during endosomal

shuttling. In case of MLLE1 we did not identify a clear function so far. However, we believe
that all three MLLE domains are functional important. This is supported by the fact that the
presence of a MLLE binding platform with three MLLE domains is evolutionarily conserved.
Even Rrm4 versions of the distantly related fungus *Rhizophagus irregularis* contains three
MLLE domains (Mucoromycota, determined by AlphaFold; Müller *et al.*, 2019).

Studying the spatial arrangement of the three MLLE domains revealed that they form a
highly flexible binding platform. This would allow for the simultaneous interaction of several
binding partners and potential rearrangements like an induced fit after binding. Such a scenario
might be particularly important during the loading and unloading of mRNPs to endosomes.
Noteworthy, the N-terminal RNA-binding domain consisting of three RRMs is clearly
separated from the MLLE domains for endosomal attachment. This is comparable with the
arrangement of RRM and MLLE domains in human PABPC1: the four N-terminal RRM
domains interact with the poly(A) tail of mRNAs, and the MLLE domain is exposed by a
flexible spacer region for protein/protein interactions (Schäfer *et al.*, 2019). Within the spacer
region, additional interactions with the RRM2 of PABPC1 were found, suggesting a function
in multimerization of the protein on the poly(A) tail of mRNAs (Sawazaki *et al.*, 2018).

Conclusion

Endosomal mRNA transport is evolutionarily highly conserved. Besides hyphal growth in
fungi, it is important for endosperm development in plants as well as neuronal functions in
animals and humans (Béthune *et al.*, 2019; Fernandopulle *et al.*, 2021; Müntjes *et al.*, 2021;
Tian *et al.*, 2020a). Malfunction causes defects in polar growth in fungi and has been implicated
in neuronal diseases such as Charcot-Marie-Tooth type 2B neuropathy or amyotrophic lateral
sclerosis in humans (Cioni *et al.*, 2019; Liao *et al.*, 2019).
A key question is how mRNPs are linked to endosomes. In plants, two RRM-type RNA-binding
proteins form a complex with cargo mRNAs and the endosomal component *N*-ethylmaleimide-
sensitive factor NSF as well as Rab5a (Tian *et al.*, 2018; Tian *et al.*, 2020a; Tian *et al.*, 2020b).
Comparably, the FERRY complex (Five-subunit Endosomal Rab5 and RNA/ribosome
intermediarY) interacts with the activated GTP-bound form of Rab5 during endosomal mRNA
transport in neurons (Schuhmacher *et al.*, 2021; Quentin *et al.*, 2021). Further examples are the
membrane-associated protein Annexin 11 that functions in linking large RNA granules to
lysosomal vesicles during mRNA transport in axons and dendrites (Liao *et al.*, 2019). Thus, a
number of components and interactions are known, however detailed structural insights are
scarce. Here, we demonstrate that in hyphae, endosomal attachment of Rrm4 is mediated by an

410 MLLE binding platform with a non-canonical accessory domain joining an essential MLLE

411 domain for perfect interaction to Upa1 on the endosomal surface (Fig. 6). This binary

412 interaction in the core of the transport mRNPs is supported by numerous interactions of

413 additional protein partners such as Upa2 and Pab1 that assist in the attachment of components

414 to the endosomal surfaces (Fig. 6). In essence, studying endosomal mRNP transport in fungal

415 model systems might guide future research endeavours in plant and neuronal systems.

Materials and methods

Structural modeling of C-terminal MLLE domains of Rrm4

To obtain structural models of the C-terminal region of Rrm4, an iterative homology modeling
approach was used with the TopModel workflow (Mulnaes *et al.*, 2020a). Initially, the complete
C-terminal region (421 to 792) was submitted as input in TopModel and identified templates
for MLLE3^{Rrm4} (665 – 792 AA; Fig. 1C, Fig. EV1A). Then, the rest of the C-terminal part
comprising amino acids 421 to 664 was resubmitted as input and identified other templates as
a new starting point for the MLLE2^{Rrm4} (571-629). Likewise, the remaining C-terminal
sequence comprising amino acids 421 to 549 was resubmitted as input, which identified other
templates as a new starting point for the MLLE1^{Rrm4} (446-530). In total, this led to the
identification of three MLLE domains, for which structural models were generated using
default TopModel parameters. The quality of the structural models was assessed with TopScore
(Mulnaes & Gohlke, 2018).

Plasmids, strains, and growth conditions

For molecular cloning of plasmids, *Escherichia coli* Top10 cells (Thermofisher C404010) and
for recombinant protein expression *E .coli Lobstr* cells (Kerafast EC1002) were used
respectively. Sequence encoding H-Rrm4-NT4 was subcloned into pET22 vector (Merck
69744) with an N-terminal hexa-histidine tag for crystallization studies. Sequence encoding
MLLE variants were inserted into pGEX-2T vector (Merck GE28-9546-53) containing GST
sequence in N-terminus for pulldown and ITC experiments. Sequence encoding PAM2 variants
were inserted into Champion pET-Sumo vector (Thermofisher K30001). pRarepLys plasmid
was co-transformed in *E. coli Lobstr strain* to supplement the rare codons for efficient
recombinant protein production. *E. coli* transformation, cultivation, and plasmid isolation were
conducted using standard techniques. All *U. maydis* strains are derivatives of AB33, in which
hyphal growth can be induced by switching the nitrogen source in the medium (Brachmann *et*
al., 2001). *U. maydis* yeast cells were incubated in complete medium (CM) supplemented with
1% glucose, whereas hyphal growth was induced by changing to nitrate minimal medium (NM)
supplemented with 1% glucose, both at 28°C (Brachmann *et al.*, 2001). Detailed growth
conditions and general cloning strategies for *U. maydis* are described elsewhere (Baumann *et*
al., 2012; Brachmann *et al.*, 2004; Terfrüchte *et al.*, 2014). All plasmids were verified by
sequencing. Strains were generated by transforming progenitor strains with linearized plasmids.
Successful integration of constructs was verified by diagnostic PCR and by Southern blot

analysis (Brachmann *et al.*, 2004). For ectopic integration, plasmids were linearized with *Ssp*I

and targeted to the *ip*[S] locus (Loubradou *et al.*, 2001). A detailed description of all plasmids,

strains, and oligonucleotides is given in Appendix Tables S3–S7. Sequences are available upon

request.

Recombinant protein expression and purification

E. coli cells from freshly transformed plates were inoculated in 20 ml expression media. In

order to produce high density expression cultures with tight regulation of induction and

expression in shake flasks we designed a complex media inspired from the principle of Studier's

autoinduction media (Studier, 2005). We use adequate amount of glucose to prevent the

unintended induction and leaky expression of target protein as well as phosphate buffer to

prevent acidity as a result of glucose metabolism from the excessive glucose in the media. In

addition, the medium contained glycerol, nitrogen, sulphur, and magnesium for promoting high

density growth. Unlike the studier's autoinduction media our media lack lactose therefore

expression can be induced with IPTG and expressed at required temperature (Studier, 2005; 1.6

% Trypton, 1% Yeast extract, 50 mM Na_2HPO_4, 50 mM KH_2PO_4, 25 mM $[NH_4]_2SO_4$, 0.5%

Glycerol, 0.5% Glucose, 2 mM $MgSO_4$) with ampicillin (100 mg/ml) and chloramphenicol (34

mg/ml) or kanamycin (200 mg/ml) and chloramphenicol (34 mg/ml) and grown overnight (16

hours) at 37 °C, 200 rpm. Note, high concentration of kanamycin was used to prevent the

unintended resistance promoted by high phosphate concentration (Studier, 2005). Supernatant

from the overnight culture was removed by centrifugation at 4 °C, 5000 g for 2 minutes. Cells

were resuspended in fresh media with a starting OD_{600} of 0.1 and grown at 37 °C, 200 rpm for

about 2 hours 30 minutes until the $OD_{600} = 1$. Protein expression was induced at 28 °C, 200

rpm, for 4 hours by addition of 1 mM IPTG, and harvested by centrifugation at 4°C, 6,000 g

for 5 minutes. Protein purification was performed as per the previous report (Abts *et al.*, 2013).

Hexa-histidine tagged protein was purified using Nickle-based affinity chromatography

(HisTrap HP GE healthcare) on Akta prime FPLC system. Cells were thawed on ice and

resuspended in buffer A (20 mM HEPES pH 8.0, 200 mM NaCl, 1 mM EDTA, 10 mM

Imidazole pH 8.0; 1 mM PMSF, 0.5 mg/ml Lysozyme, 0.5 mg/ml DNase, 1mM β

mercaptoethanol [β-ME]). Subsequently, cells were lysed by sonication on ice and centrifuged

at 4 °C 18,000 g for 30 minutes. Resulting supernatant was loaded on to a pre-equilibrated

column with buffer B (20 mM HEPES pH 8.0, 200 mM NaCl,10 mM Imidazole), washed with

buffer C (20 mM HEPES pH 8.0, 200 mM NaCl, 50 mM Imidazole, 1 mM β-ME), eluted with

buffer D (20 mM HEPES pH 8.0, 200 mM NaCl, 300 mM Imidazole, 1mM β-ME) and further

purified by Size Exclusion Chromatography (HiLoad 26/600 Superdex 200, GE healthcare),

pre-equilibrated with buffer E (20 mM HEPES pH 8.0, 200 mM NaCl, 1 mM β-ME). For

crystallization studies, H-Rrm4-NT4 was purified as above except that the buffers were

prepared with high salt (500 mM NaCl) and without β-ME.

GST-tagged protein was purified using Glutathione-based affinity chromatography (GSTrap

FF GE healthcare). Cells were thawed on ice and resuspend in Buffer F (20 mM HEPES pH

8.0, 200 mM NaCl, 1 mM EDTA, pH 8.0; 1 mM PMSF, 0.1 mg/ml Lysozyme, 1 mM β-ME).

Subsequently, cells were lysed by sonication on ice and centrifuged at 4 °C, 18,000 g for 30

minutes. The resulting supernatant was loaded onto a pre-equilibrated column with buffer E (20

mM HEPES pH 8.0, 200 mM NaCl, 1 mM β-ME) and washed with the same buffer, eluted

with buffer H (20 mM HEPES pH 8.0, 200 mM NaCl, 10 mM reduced glutathione, 1 mM β-

ME), and further purified by size exclusion chromatography (HiLoad 16/600 Superdex 200 GE

healthcare), pre-equilibrated with buffer E. Protein purity was analysed on SDS PAGE. All the

purified proteins were concentrated, centrifuged at 4°C, 100,000 g for 30 minutes, quantified

by Nanodrop A280, aliquoted, and stored at -80°C. Peptides for ITC experiments were custom-

synthesised and purchased from Genscript, USA (see Fig. 3A for peptide sequence).

GST pull-down experiments

Pull-down assays were performed as per the previous report (Jankowski *et al.*, 2019). GST-

MLLE variants and HS-PAM2 variants were expressed in *E. coli* and harvested as described

above. Cell pellets from 50 ml *E. coli* expression culture were resuspended in 10 ml buffer F

(20 mM HEPES pH 8.0, 200 mM NaCl, 1 mM EDTA; 0.5% Nonidet P-40, 1 mM PMSF, 0.1

mg/ml Lysozyme). Cells were lysed by sonication on ice and centrifuged at 4 °C, 16,000 g for

10 minutes. 1 mL of the resulting supernatant was incubated for 1 hour at 4°C on constant

agitation of 1,000 rpm with 100 µL glutathione sepharose (GS) resin (GE healthcare), pre-

equilibrated in buffer F. The GS resin was washed three times with 1 ml of buffer G (20 mM

HEPES pH 8.0, 200 mM NaCl, 1 mM EDTA, 0.5 % Nonidet P-40). Subsequently, supernatant

of HS-PAM2 variants was added to the GST-MLLE variant bound resins and incubated for 1

hour at 4 °C on agitation. The resins were washed as aforementioned and resuspended in 100

µL of 2x Laemmli loading buffer and boiled for 10 minutes at 95 °C and analysed by western

blotting.

SDS-PAGE and immunoblotting

All SDS–PAGE and Western blotting experiments were performed as per the previous report.

(Jankowski *et al.*, 2019). For Western blotting, protein samples were resolved by 12 % SDS–

PAGE and transferred to a PVDF membrane (GE Healthcare) by semi-dry blotting. Western

blot analysis was performed with anti-His (Sigma H1029). Anti-mouse IgG HRP conjugate

(Promega W4021) was used as a secondary antibody. The activity was detected using the

AceGlow blotting detection system (VWR Peqlab, Erlangen, Germany).

Multiangle laser light scattering (MALS)

MALS was performed as per the previous report (Weiler *et al.*, 2021). Superdex 200 Increase

10/300 GL column (GE Healthcare) was pre-equilibrated overnight at a flow rate of

0.1 ml/minute with buffer E (20 mM HEPES pH 8.0, 200 mM NaCl, 1 mM β-ME). For each

analysis, 200 µl of protein sample at the concentrations of 2.0 mg/ml was loaded onto the

column at the flow rate of 0.6 ml/minute using 1260 binary pump (Agilent Technologies).

Scattered light was measured with miniDAWN TREOS II light scatterer, (Wyatt Technologies),

and the refractive index was measured with Optilab T-rEX refractometer, (Wyatt

Technologies). Data analysis was performed with the software ASTRA 7.3.2.21 (Wyatt

Technologies; Slotboom *et al.*, 2008).

Crystallization of H-Rrm4 NT4

Initial crystallization conditions were searched using MRC 3 96-well sitting drop plates and

various commercially available crystallization screens at 12 °C. 0.1 µl homogeneous protein

solution (10 mg/ml in 20 mM Hepes pH 8.0, 500 mM NaCl) was mixed with 0.1 µl reservoir

solution and equilibrated against 40 µl of the reservoir. After one week, initial rod-shaped

crystals were found, which were then further optimized by slightly varying the precipitant

concentrations. Optimisation was also performed in sitting drop plates (24-well) at 12 °C but

by mixing 1 µl protein solution with 1 µl of the reservoir solution, equilibrated against 300 µl

reservoir solution. Best diffracting crystals were grown within 7 days in 0.1 M Hepes pH 7.5,

20% (w/v) PEG 10000 (Qiagen PEG I, D5). Crystal-containing drops were overlaid with 2 µl

mineral oil before harvesting the crystal and immediately flash-frozen in liquid nitrogen.

Data collection, processing, and structure refinement

A complete data set of the H-Rrm4-NT4 were collected at beamline ID23EH1 (ESRF, France)

at 100 K and wavelength 0.98 Å up to 2.6 Å resolution. All data were processed using the

automated pipeline at the EMBL HAMBURG and reprocessed afterwards using XDS (Kabsch,

2014). The above obtained model for MLLE2 by TopModel was successfully used to phase the

2.6 Å data set of Rrm4 MLLE using the program Phaser from the program suite Phenix

(Afonine *et al.*, 2012). The structure was then refined in iterative cycles of manual building and

refinement in coot (Emsley & Cowtan, 2004) followed by software-based refinements using

the program suite Phenix (Afonine *et al.*, 2012.) All residues were in the preferred and

additionally allowed regions of the Ramachandran plot (Appendix Table S2). The data

collection and refinement statistics are listed in Appendix Table S2. The comparison of the

structure and models were performed using the superpose tool of PHENIX calculating the

corresponding RMSD. The images of the models were prepared using PyMOL. The structure

was deposited at the worldwide protein data bank under the accession code 7PZE.

Small angle X-ray scattering

We collected all SAXS data on beamline BM29 at the ESRF Grenoble (Franke *et al.*, 2017

Pernot *et al.*, 2013; Pernot *et al.*, 2010). The beamline was equipped with a PILATUS 2M

detector (Dectris) with a fixed sample to distance of 2.827 m. To prevent concentration-

dependent oligomerization, we performed the measurements with concentrations of 0.6 mg/ml

at a temperature of 10 °C in buffer E. We collected one frame each second and scaled the data

to absolute intensity against water. All used programs for data processing were part of the

ATSAS Software package (Version 3.0.3;Manalastas-Cantos *et al.*, 2021). The primary data

reduction was performed with the program Primus (Konarev *et al.*, 2003). With Primus and the

included Guinier approximation (Franke & Svergun, 2009), we determined the forward

scattering $I(0)$ and the radius of gyration (R_g). The pair-distribution function $p(r)$ was calculated

with Gnom (Svergun, 1992) and was used to estimate the maximum particle dimension (D_{max}).

Due to the high flexibility of the proteins we performed an Ensemble Optimization Method

(EOM; Tria *et al.*, 2015; default parameters, 10,000 models in initial ensemble, native-like

models, constant subtraction allowed) with the predicted MLLE domains from TopModel

(Mulnaes *et al.*, 2021; Mulnaes *et al.*, 2020b) for H-Rrm4-NT4 and G-Rrm4 with an additional

GST (PDB1ua5). We uploaded the data to the Small Angle Scattering Biological Data Bank

(SASBDB; Kikhney *et al.*, 2020; Valentini *et al.*, 2015) with the accession codes SASDMS5(G-

Rrm4) and SASDMT5 (H-Rrm4-NT4).

Isothermal titration calorimetry

All ITC experiments were performed as per the previous report (Abts *et al.*, 2013). All the protein samples used in ITC were centrifuged at 451,000 g for 30 minutes and quantified by Nanodrop (A280) before use. The concentration of GST-MLLE variants was adjusted to 30 µM and PAM2 peptide variants was adjusted to 300 µM using buffer G (20 mM HEPES pH 8.0, 200 mM NaCl, 1 mM 2 ME) at 25°C. Using an MicroCal iTC200 titration calorimeter (Malvern Panalytical technologies), a PAM2 peptide variant with a volume of 40 µL was titrated to the different GST-MLLE variants. All experiments were repeated at least twice. ITC measurements were performed at 25 °C with 40 injections (1 µL each). Only the first injection had a volume of 0.5 µL and was discarded from the isotherm. The other technical parameters were reference power, 5 µcal s^{-1}; stirring speed, 1000 rpm; spacing time, 120 s, and a filter period, 5 s. The resulting isotherm was fitted with a one-site binding model using MicroCal Origin for ITC software (MicroCal LLC).

Microscopy, image processing and image analysis

Laser-based epifluorescence-microscopy was performed on a Zeiss Axio Observer.Z1 as previously described (Jankowski *et al.*, 2019). Co-localisation studies of dynamic processes were carried out with a two-channel imager (DV2, Photometrics, Tucson, AZ, USA; Baumann *et al.*, 2016). To quantify uni- and bipolar hyphal growth, cells were grown in 30 ml cultures to an OD$_{600}$ of 0.5, and hyphal growth was induced. After 6 hours, more than 100 hyphae were analysed per strains towards their growth behaviour (n = 3). Cells were assessed for unipolar and bipolar growth as well as the formation of a basal septum. To analyse the signal number, velocity, and travelled distance of fluorescently labelled proteins, movies with an exposure time of 150 ms and 150 frames were recorded. More than 25 hyphae were analysed per strain (n = 3). All movies and images were possessed and analysed using the Metamorph software (Version 7.7.0.0, Molecular Devices, Seattle, IL, USA). For the generation of kymographs, 20 µm of hyphal cell starting at the hyphal tip were used. For statistical analysis of signal number, velocity, and travelled distance, processive signals with a travelled distance of more than 5 µm were conducted and counted manually. Data points represent means from three independent experiments (*n* = 3) with mean of means (red line) and SEM. For all statistical evaluations, two-tailed Student's *t*-tests were used.

Acknowledgements

We thank Drs. Florian Altegoer and Dierk Niessing as well as laboratory members for critical reading of the manuscript. We acknowledge Drs. Georg Groth, Alexander Minges, and Mohan Gopalswamy for support with ITC experiments. We gratefully acknowledge the computational support provided by the "Center for Information and Media Technology" (ZIM) at the Heinrich Heine University Düsseldorf and the computing time provided by the John von Neumann Institute for Computing (NIC) on the supercomputer JUWELS at Jülich Supercomputing Centre (JSC) (user IDs: HKF7, VSK33). The synchrotron MX data were collected at ESRF ID23-EH1. We thank Sylvain Engilberge for the assistance in using the beamline. The final SAXS experiments were performed on beamline BM29 at the European Synchrotron Radiation Facility (ESRF), Grenoble, France. We are grateful to Dihia Moussaoui at the ESRF for providing assistance during data collection. The work was funded by grants from the Deutsche Forschungsgemeinschaft under Germany's Excellence Strategy EXC-2048/1 - Project ID 39068111 to MF; Project-ID 267205415 – SFB 1208 to MF (project A09), HG (project A03), and LS (project A01). The Center for Structural Studies is funded by the Deutsche Forschungsgemeinschaft (DFG Grant number 417919780; INST 208/740-1 FUGG; INST 208/761-1 FUGG).

Author contributions

SKD, SSV, LS, SHJS, HG, and MF designed this study and analysed the data. SKD and SSV contributed equally to the structural biology of the MLLE binding platform; SKD performed the wet lab experiments and SSV was responsible for modelling. KM and SKD performed the cell biological experiments. LB coordinated strain generation and experimental design. JR, EH, AH contributed to SAXS, MALS, and X ray analysis, respectively. SKD, SSV, LS, SHJS, HG, and MF drafted and revised the manuscript with input from all co-authors. SKD, SSV, SHJS, HG and MF directed the project.

Conflict of interest

The authors declare that they have no competing interests.

References

Abts A, Montalban-Lopez M, Kuipers OP, Smits SH, Schmitt L (2013) NisC binds the FxLx motif of the nisin leader peptide. *Biochemistry* 52: 5387-5395

Afonine PV, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, Terwilliger TC, Urzhumtsev A, Zwart PH, Adams PD (2012) Towards automated crystallographic structure refinement with phenix.refine. *Acta Crystallogr D Biol Crystallogr* 68: 352-367

Albrecht M, Lengauer T (2004) Survey on the PABC recognition motif PAM2. *Biochem Biophys Res Commun* 316: 129-138

Baek M, DiMaio F, Anishchenko I, Dauparas J, Ovchinnikov S, Lee GR, Wang J, Cong Q, Kinch LN, Schaeffer RD *et al.* (2021) Accurate prediction of protein structures and interactions using a three-track neural network. *Science* 373: 871-876

Baumann S, König J, Koepke J, Feldbrügge M (2014) Endosomal transport of septin mRNA and protein indicates local translation on endosomes and is required for correct septin filamentation. *EMBO Rep* 15: 94-102

Baumann S, Pohlmann T, Jungbluth M, Brachmann A, Feldbrügge M (2012) Kinesin-3 and dynein mediate microtubule-dependent co-transport of mRNPs and endosomes. *J Cell Sci* 125: 2740-2752

Baumann S, Zander S, Weidtkamp-Peters S, Feldbrügge M (2016) Live cell imaging of septin dynamics in *Ustilago maydis*. In: *Methods in Cell Biol: Septins*, Gladfelter A.S. (ed.) pp. 143-159. Elsevier Inc.:

Becht P, König J, Feldbrügge M (2006) The RNA-binding protein Rrm4 is essential for polarity in *Ustilago maydis* and shuttles along microtubules. *J Cell Sci* 119: 4964-4973

Béthune J, Jansen RP, Feldbrügge M, Zarnack K (2019) Membrane-associated RNA-binding proteins orchestrate organelle-coupled translation *Trends in cell biology* 29: 178-188

Brachmann A, König J, Julius C, Feldbrügge M (2004) A reverse genetic approach for generating gene replacement mutants in *Ustilago maydis*. *Mol Gen Genom* 272: 216-226

Brachmann A, Weinzierl G, Kämper J, Kahmann R (2001) Identification of genes in the bW/bE regulatory cascade in *Ustilago maydis*. *Mol Microbiol* 42: 1047-1063

Cioni JM, Lin JQ, Holtermann AV, Koppers M, Jakobs MAH, Azizi A, Turner-Bridger B, Shigeoka T, Franze K, Harris WA *et al.* (2019) Late endosomes act as mRNA translation platforms and sustain mitochondria in axons. *Cell* 176: 56-72 e15

Das S, Vera M, Gandin V, Singer RH, Tutucci E (2021) Intracellular mRNA transport and localized translation. *Nat Rev Mol Cell Biol* 22: 483-504

Deo RC, Sonenberg N, Burley SK (2001) X-ray structure of the human hyperplastic discs
protein: an ortholog of the C-terminal domain of poly(A)-binding protein. *Proc Natl Acad*
Sci U S A 98: 4414-4419

Emsley P, Cowtan K (2004) Coot: model-building tools for molecular graphics. *Acta*
Crystallogr D Biol Crystallogr 60: 2126-2132

Fernandopulle MS, Lippincott-Schwartz J, Ward ME (2021) RNA transport and local
translation in neurodevelopmental and neurodegenerative disease. *Nature neuroscience*

Franke D, Petoukhov MV, Konarev PV, Panjkovich A, Tuukkanen A, Mertens HDT, Kikhney
AG, Hajizadeh NR, Franklin JM, Jeffries CM *et al.* (2017) ATSAS 2.8: a comprehensive
data analysis suite for small-angle scattering from macromolecular solutions. *J Appl*
Crystallogr 50: 1212-1225

Franke D, Svergun DI (2009) DAMMIF, a program for rapid ab-initio shape determination in
small-angle scattering. *J Appl Crystallogr* 42: 342-346

Grimm C, Pelz JP, Schneider C, Schaffler K, Fischer U (2020) Crystal structure of a variant
PAM2 motif of LARP4B bound to the MLLE domain of PABPC1. *Biomol* 10

Haag C, Steuten B, Feldbrügge M (2015) Membrane-coupled mRNA trafficking in fungi. *Annu*
Rev Microbiol 69: 265-281

Jankowski S, Pohlmann T, Baumann S, Müntjes KM, Devan SK, Zander S, Feldbrügge M
(2019) The multi PAM2 protein Upa2 functions as novel core component of endosomal
mRNA transport. *EMBO Rep* 24: e47381

Jinek M, Fabian MR, Coyle SM, Sonenberg N, Doudna JA (2010) Structural insights into the
human GW182-PABC interaction in microRNA-mediated deadenylation. *Nat Struct Mol*
Biol 17: 238-240

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
R, Zidek A, Potapenko A *et al.* (2021) Highly accurate protein structure prediction with
AlphaFold. *Nature* 596: 583-589

Kabsch W (2014) Processing of X-ray snapshots from crystals in random orientations. *Acta*
Crystallogr D Biol Crystallogr 70: 2204-2216

Kikhney AG, Borges CR, Molodenskiy DS, Jeffries CM, Svergun DI (2020) SASBDB:
Towards an automatically curated and validated repository for biological scattering data.
Protein Sci 29: 66-75

Konarev PV, Volkov VV, Sokolova AV, Koch MHJ, Svergun DI (2003) PRIMUS: a Windows
PC-based system for small-angle scattering data analysis. *J Appl Crystallogr* 36: 1277-
1282

Kozlov G, De Crescenzo G, Lim NS, Siddiqui N, Fantus D, Kahvejian A, Trempe JF, Elias D,
Ekiel I, Sonenberg N *et al.* (2004) Structural basis of ligand recognition by PABC, a highly
specific peptide-binding domain found in poly(A)-binding protein and a HECT ubiquitin
ligase. *EMBO J* 23: 272-281

Kozlov G, Gehring K (2010) Molecular basis of eRF3 recognition by the MLLE domain of
poly(A)-binding protein. *PLoS One* 5: e10169

Kozlov G, Menade M, Rosenauer A, Nguyen L, Gehring K (2010) Molecular determinants of
PAM2 recognition by the MLLE domain of poly(A)-binding protein. *J Mol Biol* 397: 397-
407

Kozlov G, Trempe JF, Khaleghpour K, Kahvejian A, Ekiel I, Gehring K (2001) Structure and
function of the C-terminal PABC domain of human poly(A)-binding protein. *Proc Natl*
Acad Sci U S A 98: 4409-4413

Liao YC, Fernandopulle MS, Wang G, Choi H, Hao L, Drerup CM, Patel R, Qamar S, Nixon-
Abell J, Shen Y *et al.* (2019) RNA granules hitchhike on lysosomes for long-distance
transport, using annexin A11 as a molecular tether. *Cell* 179: 147-164 e120

Lim NS, Kozlov G, Chang TC, Groover O, Siddiqui N, Volpon L, De Crescenzo G, Shyu AB,
Gehring K (2006) Comparative peptide binding studies of the PABC domains from the
ubiquitin-protein isopeptide ligase HYD and poly(A)-binding protein. Implications for
HYD function. *The Journal of biological chemistry* 281: 14376-14382

Loubradou G, Brachmann A, Feldbrügge M, Kahmann R (2001) A homologue of the
transcriptional repressor Ssn6p antagonizes cAMP signalling in *Ustilago maydis. Mol*
Microbiol 40: 719-730

Manalastas-Cantos K, Konarev PV, Hajizadeh NR, Kikhney AG, Petoukhov MV, Molodenskiy
DS, Panjkovich A, Mertens HDT, Gruzinov A, Borges C *et al.* (2021) ATSAS 3.0:
expanded functionality and new tools for small-angle scattering data analysis. *J Appl*
Crystallogr 54

Mangus DA, Amrani N, Jacobson A (1998) Pbp1p, a factor interacting with *Saccharomyces*
cerevisiae poly(A)-binding protein, regulates polyadenylation. *Molecular and cellular*
biology 18: 7383-7396

Mangus DA, Evans MC, Jacobson A (2003) Poly(A)-binding proteins: multifunctional
scaffolds for the post-transcriptional control of gene expression. *Genome Biol* 4: 223

Mattijssen S, Kozlov G, Fonseca BD, Gehring K, Maraia RJ (2021) LARP1 and LARP4: up
close with PABP for mRNA 3' poly(A) protection and stabilization. *RNA Biol* 18: 259-274

Mofatteh M, Bullock SL (2017) SnapShot: Subcellular mRNA localization. *Cell* 169: 178.e171

Müller J, Pohlmann T, Feldbrügge M (2019) Core components of endosomal mRNA transport
are evolutionarily conserved in fungi. *Fungal Genet Biol* 126: 12-16

Mulnaes D, Gohlke H (2018) TopScore: using deep neural networks and large diverse data sets
for accurate protein model quality assessment. *J Chem Theory Comput* 14: 6117-6126

Mulnaes D, Koenig F, Gohlke H (2021) TopSuite Web Server: A Meta-Suite for Deep-
Learning-Based Protein Structure and Quality Prediction. *Journal of chemical information*
and modeling 61: 548-553

Mulnaes D, Porta N, Clemens R, Apanasenko I, Reiners J, Gremer L, Neudecker P, Smits SHJ,
Gohlke H (2020a) TopModel: template-based protein structure prediction at low sequence
identity using top-down consensus and deep neural networks. *J Chem Theory Comput* 16:
1953-1967

Mulnaes D, Porta N, Clemens R, Apanasenko I, Reiners J, Gremer L, Neudecker P, Smits SHJ,
Gohlke H (2020b) TopModel: Template-Based Protein Structure Prediction at Low
Sequence Identity Using Top-Down Consensus and Deep Neural Networks. *J Chem*
Theory Comput 16: 1953-1967

Munoz-Escobar J, Matta-Camacho E, Kozlov G, Gehring K (2015) The MLLE domain of the
ubiquitin ligase UBR5 binds to its catalytic domain to regulate substrate binding. *The*
Journal of biological chemistry 290: 22841-22850

Müntjes K, Devan SK, Reichert AS, Feldbrügge M (2021) Linking transport and translation of
mRNAs with endosomes and mitochondria. *EMBO reports* accepted for publication

Müntjes K, Philipp M, Husemann L, Heucken N, Weidtkamp-Peters S, Schipper K, Zurbriggen
MD, Feldbrügge M (2020) Establishing polycistronic expression in the model
microorganism *Ustilago maydis*. *Front Microbiol* 11: 1384

Niessing D, Jansen RP, Pohlmann T, Feldbrügge M (2018) mRNA transport in fungal top
models. *Wiley interdisciplinary reviews RNA* 9: e1453

Olgeiser L, Haag C, Boerner S, Ule J, Busch A, Koepke J, König J, Feldbrügge M, Zarnack K
(2019) The key protein of endosomal mRNP transport Rrm4 binds translational landmark
sites of cargo mRNAs. *EMBO Rep* 20: e46588

Pernot P, Round A, Barrett R, De Maria Antolinos A, Gobbo A, Gordon E, Huet J, Kieffer J,
Lentini M, Mattenet M *et al.* (2013) Upgraded ESRF BM29 beamline for SAXS on
macromolecules in solution. *Journal of synchrotron radiation* 20: 660-664

Pernot P, Theveneau P, Giraud T, Fernandes RN, Nurizzo D, Spruce D, Surr J, McSweeney S,
Round A, Felisaz F *et al.* (2010) New beamline dedicated to solution scattering from

biological macromolecules at the ESRF. *Journal of Physics: Conference Series* 247:
012009

Pohlmann T, Baumann S, Haag C, Albrecht M, Feldbrügge M (2015) A FYVE zinc finger
domain protein specifically links mRNA transport to endosome trafficking. *eLife* 4: e06041

Quentin D, Schuhmacher JS, Klink BU, Lauer J, Shaikh TR, Huis in t´t Veld PJ, Welp LM,
Urlaub H, Zerial M, Raunser S (2021) Structure of the human FERRY Rab5 effector
complex. *bioRiv* 10.1101/2021.06.21.449265

Sawazaki R, Imai S, Yokogawa M, Hosoda N, Hoshino SI, Mio M, Mio K, Shimada I, Osawa
M (2018) Characterization of the multimeric structure of poly(A)-binding protein on a
poly(A) tail. *Sci Rep* 8: 1455

Schäfer IB, Yamashita M, Schuller JM, Schussler S, Reichelt P, Strauss M, Conti E (2019)
Molecular bsis for poly(A) RNP architecture and recognition by the Pan2-Pan3
deadenylase. *Cell* 177: 1619-1631 e1621

Schuhmacher JS, Dieck ST, Christoforidis S, Landerer C, Hersemann L, Seifert S, Giner A,
Toth-Petroczy A, Kalaidzidis Y, Schumann EM *et al.* (2021) The novel Rab5 effector
FERRY links early endosomes with the translation machinery. *BioRiv*
10.1101/2021.06.20.449167

Slotboom DJ, Duurkens RH, Olieman K, Erkens GB (2008) Static light scattering to
characterize membrane proteins in detergent solution. *Methods* 46: 73-82

Stenmark H, Aasland R, Driscoll PC (2002) The phosphatidylinositol 3-phosphate-binding
FYVE finger. *FEBS Lett* 513: 77-84

Studier FW (2005) Protein production by auto-induction in high density shaking cultures.
Protein Expr Purif 41: 207-234

Su H, Meng S, Lu Y, Trombly MI, Chen J, Lin C, Turk A, Wang X (2011) Mammalian
hyperplastic discs homolog EDD regulates miRNA-mediated gene silencing. *Mol Cell* 43:
97-109

Svergun DI (1992) Determination of the regularization parameter in indirect-transform methods
using perceptual criteria. . *J Appl Crystallogr* 25: 495-503

Terfrüchte M, Joehnk B, Fajardo-Somera R, Braus G, Riquelme M, Schipper K, Feldbrügge M
(2014) Establishing a versatile Golden Gate cloning system for genetic engineering in
fungi. *Fungal Genet Biol* 62: 1-10

Tian L, Chou HL, Fukuda M, Kumamaru T, Okita TW (2020a) mRNA localization in plant
cells. *Plant Physiol* 182: 97-109

Tian L, Chou HL, Zhang L, Hwang SK, Starkenburg SR, Doroshenk KA, Kumamaru T, Okita
TW (2018) RNA-binding protein RBP-P is required for glutelin and prolamine mRNA
localization in rice endosperm cells. *Plant Cell* 30: 2529-2552

Tian L, Doroshenk KA, Zhang L, Fukuda M, Washida H, Kumamaru T, Okita T (2020b)
Zipcode RNA-binding proteins and membrane trafficking proteins cooperate to transport
glutelin mRNAs in rice endosperm. *Plant Cell* 32: 2566-2581

Tria G, Mertens HD, Kachala M, Svergun DI (2015) Advanced ensemble modelling of flexible
macromolecules using X-ray solution scattering. *IUCrJ* 2: 207-217

Valentini E, Kikhney AG, Previtali G, Jeffries CM, Svergun DI (2015) SASBDB, a repository
for biological small-angle scattering data. *Nucleic Acids Res* 43: D357-363

Weiler AJ, Spitz O, Gudzuhn M, Schott-Verdugo SN, Kamel M, Thiele B, Streit RW, Kedrov
A, L. S, Gohlke H *et al.* (2021) Novel intracellular phospholipase B from *Pseudomonas*
aeruginosa with activity towards endogenous phospholipids affects biofilm assembly.
bioRiv doi.org/10.1101/2021.06.15.448513

Xie J, Kozlov G, Gehring K (2014) The "tale" of poly(A) binding protein: the MLLE domain
and PAM2-containing proteins. *Biochim Biophys Acta* 1839: 1062-1068

Yang R, Gaidamakov SA, Xie J, Lee J, Martino L, Kozlov G, Crawford AK, Russo AN, Conte
MR, Gehring K *et al.* (2011) La-related protein 4 binds poly(A), interacts with the poly(A)-
binding protein MLLE domain via a variant PAM2w motif, and can promote mRNA
stability. *Mol Cell Biol* 31: 542-556

Zander S, Baumann S, Weidtkamp-Peters S, Feldbrügge M (2016) Endosomal assembly and
transport of heteromeric septin complexes promote septin cytoskeleton formation. *J Cell*
Sci 129: 2778-2792

Figures

Figure 1. The C-terminal half of Rrm4 contains three MLLE domains.

(**A**) Schematic representation of protein variants drawn to scale (bar, 200 amino acids, number
of amino acids indicated next to protein bars) using the following colouring: dark green, RNA
recognition motif (RRM); orange, MLLE^{Rrm4} domains; dark blue, MLLE^{Pab1}; light blue PAM2;

light orange PAM2L sequence (PL1 – 2) Ankyrin repeats (5xANK), FYVE domain, and RING
domain of Upa1 are given in dark grey. (**B**) Sequence alignment of previously determined
MLLE domains showing the degree of similarity to the three Rrm4-MLLE domains and the
positions (Hs - *Homo sapiens*, Ta - *Triticum aestivum*, La - *Leishmania major*, Sc -
Saccharomyces cerevisiae, Tc - *Trypanosoma cruzi*, Rn - *Rattus norvegicus*, Um - *Ustilago*
maydis, PABPC1, Pab1 – poly [A]-binding protein, UBR5 - E3 ubiquitin-protein ligase).
Accession number and sequence coverage are listed in the Appendix Table S1. Multiple
sequence alignment was performed by ClustalW. (**C**) Identification and modeling of C-terminal
MLLE domains of Rrm4. The iterative process is depicted graphically. The best identified
template for each run, and the region of that template that aligns with Rrm4, are displayed (see
also Fig. EV1A for the templates used for the final models). The structural models obtained are
shown for the span of the first identified template and are coloured according to their per-
residue TopScore, where the scale from 0 to 1 indicates a low to high local structural error.
844

Figure 2. Rrm4 contains a C-terminal tripartite MLLE binding platform

(**A**) Crystal structure of the MLLE2 domain is highlighted in orange. The four helices are
indicated by α2-5 according to the 5 helix nomenclature found in MLLE domains (Xie *et al.*,
2014). Note that the first short helix α1 is missing. Arg573 and Glu591 are highlighted in
sticks. These sides chains would interfere with binding of the canonical Phe of PAM2 type
motifs. (**B**) Structural alignment of the MLLE2 model generated by TopModel and the X-ray
crystal structure of this domain (grey or orange, respectively). The all-atom RMSD is 0.69 Å,
resulting mostly from different rotamers of solvent-exposed sidechains. (**C**) Comparison of
peptide binding sites after structural alignment of the models of Rrm4 MLLE domains (orange
shades) and the canonical MLLE domain of HsPABPC1 (blue; PDB ID 3KUS) and manually
placing the PAM2 motif of PAIP2 (lilac). In the interaction of MLLE^{PABPC1} with PAM2 of
PAIP2, Phe118 of PAM2 is the major determinant for binding and present in all the PAM2
motifs except LARP4a and b (Kozlov & Gehring, 2010; Xie *et al.*, 2014; Fig. EV3A). Of the
identified Rrm4 MLLE domains, only MLLE3 retains all sidechains that favour the binding of
this characteristic Phe; particularly, Gly736 should allow the Phe to bind into a pocket. MLLE1
and MLLE2 have Ser471 and Arg573 instead of G in this position, suggesting that Phe binding
would be sterically hindered in these interfaces. (**D**) *left panel* Experimental data curve for GST-
Rrm4 is shown in black dots with grey error bars, the EOM fit as a red line (χ^2 =1.289). The
intensity is displayed as a function of momentum transfers. *right panel*. Selected model of the
EOM analysis from GST-Rrm4 with a R_g of 8.75 nm, a D_{max} of 23.99 nm with a volume fraction
of~0.25. (**E**) *left panel* Experimental data curve for H-Rrm4NT4 is shown in black dots with
grey error bars, the EOM fit as red line (χ^2 =1.262). The intensity is displayed as a function of
momentum transfers. *right panel* Selected model of the EOM analysis from H-Rrm4NT4 with
a R_g of 5.10 nm, a D_{max} of 16.43 nm, and a volume fraction of~0.75. The MLLE subdomains
are shown in cartoon representation (MLLE1 in light orange, MLLE2 in orange, MLLE 3 in
dark orange, and the GST in dark grey) and the missing amino acids as grey spheres (all other
models and the SAXS data are available in Figure EV2E).
873
874

Figure 3. MLLE3^{Rrm4} is crucial for PAM2L1^{Upa1} and PAM2L2^{Upa1} binding.

(**A**) Schematic representation of protein variants (Molecular weight in kilo Dalton indicated)
using the following colouring: dark green, RNA recognition motif (RRM); orange, MLLE^{Rrm4}
domains; dark blue, MLLE^{Pab1}; light blue PAM2; light orange PAM2L sequence (PL1 – 2).
Ankyrin repeats (5xANK), FYVE domain, and RING domain of Upa1 are given in dark grey.
GST and SUMO tags are labelled. Amino acids of the FxP and FxxP of PAM2 and PAM2L
sequences are printed in grey fond. (**B-D**) Representative isothermal titration calorimetry (ITC)
binding curves of MLLE domains. Experiments were performed using GST- or Histidine-
tagged MLLE variants and synthetic PAM2 peptide variants. K_D values of two independent
measurements are given (values corresponding to the indicated data are given in bold).
886

Figure 4. MLLE3 is key for endosomal mRNA transport.

(**A**) Schematic representation of Rrm4 and Upa1 variants drawn not to scale (number of amino acids indicated next to protein bars) using the following colouring: dark green, RNA recognition motif (RRM); orange, MLLE domains; red, mKate2, blue, PAM2, light orange PAM2 like sequence (PL1 – 2) and light green, Gfp. Ankyrin repeats (5xANK), FYVE domain and RING domain of Upa1 are given in dark grey. (**B**) Growth of AB33 derivatives in their hyphal form (6 h.p.i.; size bar 10 μm). Growth direction is marked by arrows. (**C**) Quantification of hyphal growth of AB33 derivatives shown in B (6 h.p.i.): unipolarity, bipolarity and basal septum formation were quantified (error bars, SEM.; n = 3 independent experiments, > 100 hyphae were counted per strain; for statistical evaluation, the percentage of uni- and bipolarity was investigated by using unpaired two-tailed Student's t-test (α<0.05). (**D**) Kymographs of AB33 hyphae derivatives (6 h.p.i.; inverted fluorescence images) expressing pairs of red and green fluorescent proteins as indicated. Fluorescence signals were detected simultaneously using

901   dual-view technology (arrow length on the left and bottom indicates time and distance,
902   respectively). Processive co-localising signals are marked by red arrowheads. (**E**) Percentage
903   of processive signals exhibiting co-localisation for strains shown in D (data points represent
904   means from n = 3 independent experiments, with mean of means, red line and SEM; unpaired
905   two-tailed Student's t-test (α<0.05); for each experiment, 10 hyphae per strains were analysed).
906
907

Figure 5. MLLE2 plays an accessory role in endosomal attachment of Rrm4

(**A**) Growth of AB33 derivatives in their hyphal form (6 h.p.i.; size bar 10 μm). Growth direction is marked by arrows. (**B**). Quantification of hyphal growth of AB33 derivatives shown in A (6 h.p.i.): unipolarity, bipolarity and basal septum formation were quantified (error bars,

SEM; n = 3 independent experiments, > 100 hyphae were analysed per strain; For statistical
analysis, the percentage of uni- and bipolarity was investigated by using unpaired two-tailed
Student's t-test (α<0.05). (**C**) Micrographs (inverted fluorescence image; size bar, 10 µm) and
corresponding kymographs of AB33 hyphae derivatives (6 h.p.i.) co-expressing various Upa1-
Gfp and Rrm4-Kat versions as indicated. Movement of Rrm4-Kat versions is shown (arrow
length on the left and bottom indicates time and distance, respectively). Bidirectional movement
is visible as diagonal lines (red arrowheads). Aberrant microtubule staining is indicated by a
yellow arrowhead. (**D**) Percentage of hyphae (6 h.p.i.) exhibiting aberrant microtubule
association as indicated in panel C and EV5C. Set of strains that were analysed simultaneously
are shown in the same yellow shading (error bars, SEM; for statistical evaluation, the percentage
of hyphae with abnormal microtubule staining was compared by using unpaired two-tailed
Student's t-test (α<0.05); n = 3 independent experiments, > 25 hyphae were analysed per
strain).
926
927

Figure 6. Schematic model of endosomal attachment of mRNPs via MLLE domains

Cargo mRNAs (green) are bound by the N-terminal RRM domains of Rrm4 (green). The C terminal MLLE domains (orange) form a binding platform: MLLE3^{Rrm4} interacts with PAM2L1^{Upa1} and PAM2L2^{Upa1}(orange), MLLE1 and -2^{Rrm4} might interact with a currently unknown factor to support endosomal binding. The four RRMs of Pab1 (green) interact with the poly(A) tail, and the MLLE^{Pab1} (blue) interacts with PAM2 of Upa1 and with the four PAM2 motifs of Upa2 (dark blue), a dimerising scaffold protein. Upa1 is attached to endosomes via its FYVE domain, and the C-terminal GWW motif of Upa2 is crucial for its endosomal binding.

938 **Expanded View Figures**

A

Domain	Templates[a]	Rrm4 span	Template span	Identity (Similarity)[b]
MLLE1	6H7A:A	451-529	481-560	22.1 (38.4)
	3PTH:A	451-528	544-618	18.4 (34.5)
	1I2T:A	464-528	1009-1069	25.8 (40.9)
MLLE2	1I2T:A	571-629	1009-1069	23.8 (44.4)
	3KTP:A	550-631	554-620	17.6 (35.3)
	3KUR:A	550-627	554-616	18.5 (37.0)
MLLE3	2X04:A	712-791	540-616	30.6 (44.7)
	3KUR:A	716-791	544-616	32.9 (45.6)
	3KTP:A	716-791	544-616	32.9 (45.6)

a. PDB ID and chain identifier. b. in percentage

B

Figure EV1. The presence of three MLLEs is verified by additional modeling predictions.

(**A**) Compilation of MLLE sequences used for modeling with the highest similarity of MLLE1-
3[Rrm4]. (**B**) Structural models obtained with TopModel overlaid to Rrm4 full-length models
obtained with the recently available tools as indicated. Natural alignments between
corresponding MLLE domains have an RMSD < 2Å, mutually confirming the quality of the
independently modeled structures. The differences in the relative domain arrangements in both
full-length models and the disordered regions in between the domains suggest a high mobility
within Rrm4.

Figure EV2. The three MLLEs of Rrm4 are located in a flexible C-terminal region.

(**A**) Schematic representation of protein variants drawn to scale (molecular weight in kilo Dalton indicated next to protein bar) using the following coloring: dark green, RNA recognition motif (RRM); orange, MLLE[Rrm4] domains; (**B**) SDS PAGE analysis of purified G-Rrm4, H-Rrm4-NT4 used in crystallography and SAXS measurement. (**C**) MALS-SEC analysis of H-Rrm4-NT4. Graph shows the elution profile. Dotted line in red indicate the apparent molecular

weight as observed in the light scattering. (**D**) R_g distribution calculated by EOM pool is shown
in grey bars and the selected models in blue bars *left* GST_Rrm4 *right* H-Rrm4NT4 (**E**) *Left*
Selected models of the EOM analysis for GST-Rrm4. The MLLE subdomains and the GST are
shown in cartoon representation (MLLE1 in light orange, MLLE2 in orange, MLLE 3 in dark
orange) and the missing amino acids as grey spheres. **I:** The model has a R_g of 8.94 nm, a D_{max}
of 29.22 nm with a volume fraction of~0.38. **II:** The model has a R_g of 8.75 nm, a D_{max} of 23.99
with a volume fraction of~0.25. **III:** The model has a R_g of 7.74 nm, a D_{max} of 25.90 with a
volume fraction of~0.12. **IV:** The model has a R_g of 8.33 nm, a D_{max} of 28.79 with a volume
fraction of~0.12. **V:** The model has a R_g of 9.14 nm, a D_{max} of 33.73 with a volume fraction
of~0.12. *Right* Selected models of the EOM analysis for H-Rrm4NT4. The MLLE subdomains
are shown in cartoon representation (MLLE1 in light orange, MLLE2 in orange, MLLE3 in
dark orange) and the missing amino acids as grey spheres. **I:** The model has a R_g of 5.12 nm,
a D_{max} of 15.56 with a volume fraction of~0.17. **II:** The model has a R_g of 5.90 nm, a D_{max} of
18.73 nm with a volume fraction of~0.08. **III:** The model has a R_g of 5.10 nm, a D_{max} of 16.43
nm with a volume fraction of~0.75.
972

Figure EV3. MLLE1 ^{Rrm4} , -2^{Rrm4} are not essential for PAM2L1^{Upa1} and -L2 ^{Upa1} binding.

(**A**) Comparison of PAM2 sequences found in Upa1 (UniprotKB ID A0A0D1E015) with those of human proteins, such as Usp10 (Q14694), GW182 Q9HCJ0), Mkrn1 (Q9UHC7), Paip1 (Q9H074), Paip2 (Q9BPZ3), Atx2 (-Q99700), NFX (Q12986), eRF3 (P15170), PAN3 (Q58A45), LARP4 (Q71RC2), LARP4b (Q92615), Tob (P50616), HECT (O95071), Asp and Glu are indicated in red stressing the highly negative charges in PAM2L sequences. (**B**)

Western blot analysis of GST co-purification experiments with components expressed in *E.*
coli: N-terminal Hexa-Histidine-SUMO-tagged PAM2 variants were pulled down by N-
terminal GST fused MLLE variants of Rrm4 and Pab1. Experiment was performed with the
soluble fraction of *E. coli* cell lysate to demonstrate specific binding. Results were analysed
with αHis immunoblotting (**C**) SDS PAGE analysis of purified GST-MLLE variants used in
ITC experiments (**D-G**) Representative isothermal titration calorimetry (ITC) binding curves
of MLLE domains. Experiments were performed using GST or hexa-histidine-tagged MLLE
variants and synthetic PAM2 peptide variants. K_D values of two independent measurements are
given (values corresponding to the indicated data are given in bold). (**H**) Summary of ITC
results shown in the Fig3 and Fig EV3. K_D values are given in μM.
992

Figure EV4. Deletion of MLLE3^{Rrm4} abolishes endosomal movement of Rrm4

(**A-C**) Quantification of processive Rrm4-Kat (top) and Upa1-Gfp signals (bottom; (**A**)),
velocity of fluorescent Rrm4-Kat (top) and Upa1-Gfp signals (bottom; (**B**)) and the travelled
distance of processive Rrm4-Kat (top) and Upa1-Gfp signals (bottom; (**C**); per 10 µm of hyphal
length; only particles with a processive movement of > 5 µm were conducted; data points
representing mean from n =3 independent experiments, with mean of means, red line and SEM;
unpaired two-tailed Student's t-test ($\alpha < 0.05$), for each experiment at least 25 hyphae were
analysed per strain). (**D**) Summary of the *in vivo* analysis shown in Figure 4,5 and EV5A-C.

Figure EV5. Deletion of MLLE1^{Rrm4} and -2 cause defects in microtubule association.

(**A**) Growth of AB33 derivatives in their hyphal form (6 h.p.i.; size bar 10 μm). Growth
direction is marked by arrows. (**B**) Quantification of hyphal growth of AB33 derivatives shown
in panel A (6 h.p.i.): unipolarity, bipolarity and basal septum formation were quantified (error
bars, SEM.; n = 3 independent experiments, > 100 hyphae were analysed per strain; For
statistical evaluation, the percentage of uni- and bipolarity was investigated by using unpaired
two-tailed Student's t-test ($\alpha<0.05$). (**C**) Micrograph and Kymograph of AB33 hyphae
derivates (6 h.p.i.) expressing red and green fluorescent proteins as indicated. Fluorescence
signals were detected simultaneously using dual-view technology (arrow length on the left and
bottom indicates time and distance, respectively). Processive co-localising signals are marked
by red arrowheads. Aberrant microtubule staining is indicated by a yellow arrowhead. (**D-E**)
Quantification of processive Rrm4-Kat signals (left), velocity of fluorescent Rrm4-Kat (middle)
and the travelled distance of processive Rrm4-Kat signals (right) related to figure 5 C and
EV5C, respectively (per 10 μm of hyphal length; only particles with a processive movement
of > 5 μm were conducted; data points representing mean from n =3 independent experiments,
with mean of means, red line and SEM; unpaired two-tailed Student's t-test ($\alpha<0.05$), for each
experiment at least 25 hyphae were analysed per strain).