Effects of perfluoroalkyl and polyfluoroalkyl substances on soil

structure and function

*Baile Xu[1,2] *, Gaowen Yang[1,2], Anika Lehmann[1,2], Sebastian Riedel[3], Matthias C. Rillig[1,2]*

[1]Institute of Biology, Freie Universität Berlin, D-14195 Berlin, Germany

[2]Berlin-Brandenburg Institute of Advanced Biodiversity Research, D-14195 Berlin, Germany

[3]Institute of Chemistry and Biochemistry, Freie Universität Berlin, D-14195 Berlin, Germany

*Corresponding author. E-mail: bxu@zedat.fu-berlin.de

Abstract Art

Abstract

Soils are impacted at a global scale by several anthropogenic factors, including chemical

pollutants. Among those, perfluoroalkyl and polyfluoroalkyl substances (PFAS) are of

concern due to their high environmental persistence, and as they might affect soil health and

functions. However, data on impacts of PFASs on soil structure and microbially-driven

processes are currently lacking. This study explored the effects of perfluorooctanesulfonic

acid (PFOS), perfluorooctanoic acid (PFOA) and perfluorobutanesulfonic acid (PFBS) at

environmental-relevant nominal concentrations (1 ~ 1000 ng g^{-1}) on soil functions, using a 6-

week microcosm experiment. We measured soil respiration, litter decomposition, enzyme

and microbial activities, soil aggregates, and bacterial abundance. PFAS (even at 1 ng g^{-1}

for PFBS) significantly increased litter decomposition, associated with positive effects on

bacterial abundance, and β-glucosidase activities. This effect increased with PFAS

concentrations. Soil respiration was significantly inhibited by PFAS in the 3rd week, while this

effect was more variable in week 6. Water-stable aggregates were negatively affected by

PFOS and PFOA, possibly related to microbial shifts. The general microbial activities and β-

D-cellobiosidase and phosphatase activities were barely affected by PFAS treatments. Our

work highlights the potential effects of PFAS on soil health, and we argue that this substance

class could be a factor of environmental change of potentially broad relevance in terrestrial

ecosystem functioning.

Keywords: Litter decomposition; Soil respiration; Water-stable aggregates; Soil bacteria;

Perfluorobutanesulfonate (PFBS); Perfluorooctanesulfonic acid (PFOS); Perfluorooctanoic

acid (PFOA); Enzyme activities

Synopsis: PFAS are likely to affect soil health.

Introduction

Human activity is progressively and fundamentally affecting the Earth's surface, including soils, which operate at the interface between biosphere, hydrosphere, atmosphere, and lithosphere, and are suffering from physical, chemical, and biological stressors related to anthropogenic activities [1,2]. One significant category of these influences is chemical pollution, which is widely recognized as a global change factor [1,3]. Perfluoroalkyl and polyfluoroalkyl substances (PFAS) are a highly diverse family of chemicals of concern [4]. These compounds contain the perfluoroalkyl moiety (C_nF_{2n+1}–) [5], and the relatively high-energy carbon-fluorine (C-F) bonds make them extremely resistant to breakdown, and subsequently persistent in the environment. Therefore, they are also called "forever chemicals" in public discourse [6]. Certain PFAS, perfluorooctanesulfonic acid (PFOS) and perfluorooctanoic acid (PFOA) are on the list and on the waiting list under the Stockholm Convention [7], respectively.

Soil is a major sink for persistent organic chemicals in the environment. There are many pathways for PFAS entering the soil environment. Typically, fluoride factory emission, sludge application, the degradation of aqueous film-forming foam, and landfills contribute direct sources, and atmospheric deposition and runoff constitute non-point sources [8,9]. PFAS have been widely detected in soils with a broad range of concentrations. Generally, PFAS concentrations in non-hotspot soil are lower than 300 ng g^{-1}, while in hotspots of PFAS-contaminated soil, they can be as high as several or even tens of µg g^{-1}, mostly dominated by PFOA and PFOS [10]. Jin et al. [11] also reported that perfluorobutanesulfonic acid (PFBS) was a prominent type of PFAS in the soil around a fluoride-factory park.

Existing evidence has demonstrated that PFAS can exert some impacts on soil functions, in particular on soil enzymes, and microbial activities and communities, and these influences were closely related to properties of soil and PFASs. According to the number of carbon atoms, perfluoroalkyl carboxylic acids with 7 or more and perfluoroalkyl sulfonate with 6 or more carbons are categorized as long-chain PFAS, otherwise, short-chain PFAS [5]. Short-chain PFBS might activate sucrase and urease activities, while long-chain PFOS might

reduce these activities in soil [12]. Cai et al. [13] showed that PFAS with sulfonic groups and

longer chains had higher toxicity to soil microbial activities, and that soil with higher organic

matter content and higher pH (neutral) exhibited lower impact by PFAS. Differences in

sorption affinity of PFAS to soil are likely to regulate the impact of PFAS on soil microbes[8].

Changes in structure and function of microbial communities by PFAS were shown by

previous research [12,14–16], and these shifts were suggested to affect soil processes and

ecosystem functions. However, the implications on process rates in soil, for example, litter

decomposition and soil aggregation, were not addressed.

Given the persistent nature of PFAS it is important to explore if these chemicals can affect

soil process rates and properties. Here, we investigate PFAS effects on soil processes,

including soil respiration, litter decomposition, soil aggregation, enzyme and microbial

activities, as well as bacterial abundance. We discuss the environmental implications of our

results and suggest that PFAS be considered as a global change factor of importance in

terrestrial ecosystems.

Materials and Methods

Test soil and PFAS

The test soil was Albic Luvisol collected at the agricultural field station of Freie Universität

Berlin in December 2020. The soil has a sandy loam texture (73.6% sand, 18.8% silty and

7.6% clay), with 1.87% total C, 0.12% total N and a soil pH of 7.1 [17,18]. Fresh soil samples

were thoroughly mixed, passed through a 2-mm sieve, and then stored at 4°C.

Three PFAS, namely perfluorooctanesulfonic acid (PFOS), perfluorooctanoic acid (PFOA)

and perfluorobutanesulfonic acid (PFBS) were selected in this study due to their wide

occurrence in the soil environment [19]. Values of their Log K_{ow} are 5.26, 4.59 and 2.73,

respectively [20], and their chemical structures and other physicochemical properties are listed

in Supporting Information (SI) Table S1.

Experimental setup

PFAS standards were dissolved in sterilized deionized water to prepare the stock solution

with the concentration of 100 mg L^{-1}. A five-gram portion of previously sterilized soil samples

(121°C, 20 min, twice) was supplemented with appropriate doses of PFAS in solution (we

used this sterilized 'loading soil' to avoid any exaggerated effects on soil communities [21]),

and thoroughly mixed this soil with 25 g of soil by manual stirring for 2 mins. A total of 30 g

soil was placed in 50-mL mini-bioreactor tubes (Corning Inc., Corning, USA) with vented lids

to establish experimental microcosms. Each tube was watered to 70% soil water holding

capacity (WHC) with deionized water. The nominal concentrations of PFOS and PFOA in

soil were 1, 10, 100 and 1000 ng g^{-1}, and that of PFBS were 0.5, 5, 50 and 500 ng g^{-1},

corresponding to their environmentally-relevant levels [10]. The analytical method and actual

concentrations of PFAS in soil are reported in SI Text S1, and Table S2, respectively. For

the convenience of comparison of three PFAS, we also used the nominal concentration

below. Tubes were placed in a randomized fashion inside a dark temperature-controlled

incubator at 20 °C for 6 weeks, and each tube was watered weekly to maintain soil moisture.

This experiment ran with 10 replications of blank control (without any PFAS added, but

handled exactly the same way) and 8 replications of each treatment, for a total of 106

microcosms.

Proxies for soil health and function

We measured well-established proxies for soil health and functions, including soil respiration,

litter decomposition, enzyme activities, soil aggregates, and soil bacterial abundance. Soil

respiration was measured with an infrared gas analyzer (LI-6400XT, LI-COR Inc., Bad

Homburg, Germany), and litter decomposition was determined by the mass loss of tea bags

[18]. Soil enzymes activities were measured, including four enzymes concerning C (β-

glucosidase and β-D-1,4-cellobiosidase), N (β-1,4-N-acetyl-glucosaminidase), and P

(phosphatase) cycling, and fluorescein diacetate hydrolase (FDA) representing general soil

microbial activity. Water-stable aggregates, as the basic unit of soil structure, were qualified

using a wet-sieving apparatus (Eijkelkamp, Giesbeek, Netherlands) with an established

method [22,23] . Soil DNA was extracted with DNeasy PowerSoil Pro Kit (QIAGEN GmbH,

Germany) following the technical protocol, and we amplified using the universal primers

515F (5'-GTGCCAGCMGCCGCGGTAA-3') and 806R (5'-GGACTACHVGGGTWTCTAAT-3')

with quantitative polymerase chain reactions (qPCR) in a CFX 96 Real-Time System (Bio-

Rad Lab., Hercules, USA). For more information on measurement procedures, qPCR

conditions and quality control, see SI Text S2.

Statistical analysis

All statistical analyses and data visualization were performed in R [24]. The effects of PFAS

treatment (four concentrations per PFAS type) on soil functions were tested with a two-step

method. Firstly, we calculated the 95% confidence interval (CI) of unpaired mean differences

(treatment minus control) using the R package "dabestr" [25]. This approach focuses on the

effect size and its precision, and can avoid the pitfalls of significance testing. Secondly, one-

way analysis of variance (ANOVA) followed by Dunnett's test in the R package "multcomp"

was implemented to compare each treatment with the control [26]. Model residuals were

checked for heteroscedasticity and normal distribution. Spearman correlations among actual

concentrations of PFAS and soil structure and function were performed with the package

"corrplot" [27]. Adjusted p values by a single-step method are reported in the SI Table S3. All

data used for analyses and plotting are available online [28], and plots were generated with the

package "ggplot2" [29].

Results and Discussion

PFAS increased litter decomposition associated with bacterial abundance

Positive effects of three PFASs on litter decomposition were observed (Figure 1A), and

PFBS, particularly, at all tested concentrations significantly increased the decomposition rate

($p < 0.05$, Table S3). Regardless of tested type, increasing concentrations of PFAS

significantly enhanced litter decomposition (r = 0.15, $p = 0.0221$, Figure S1 and S2B). In

terms of PFAS type, PFOA and PFBS resulted in significantly positive effects on litter

decomposition ($p < 0.01$, Table S3), and PFBS exerted the most remarkable positive effect

(Figure S1C).

The treatment of PFASs significantly increased soil bacterial abundance (F = 1.868, $p =$

0.0485), although the multiple comparison tests did not reveal a significant difference (Table

S3). PFBS seems to have the most obvious effect among three PFAS, showing a positive

effect with concentrations higher than 0.5 ng g^{-1} (Figure 1B). Independently of concentration,

PFAS treatments had a positive effect on soil bacterial abundance in terms of copy number

per gram of dry soil (Figure S3C).

Microorganisms, including bacteria and fungi, play an essential role in the biological

decomposition of organic matter in soil, a process in which bacteria generally predominate in

neutral or alkaline soils, while fungi are more important in acidic soil [30,31]. Correspondingly,

we found that in our pH-neutral soil, with various PFAS treatments, the number of bacteria

increased compared to the control (Figure 1B), and there was a trend for a correlation

between bacteria abundance and litter decomposition rate (F=2.705, $p = 0.103$). The

increased bacterial populations probably contributed to the increased decomposition rate of

organic matter.

Litter decomposition and the ensuing nutrient release, governing carbon and nutrient cycling,

is a key process in terrestrial ecosystems. Our results showed that PFAS had a positive

effect on litter decomposition, and particularly the PFBS treatment, even at 1 ng g^{-1} in soil,

resulting in a significant enhancement of litter decomposition, likely associated with

increased soil bacterial abundance.

Figure 1. Effects of per- and polyfluoroalkyl substances (PFAS) on litter decomposition (A)

and bacterial abundance (B) in soil. In the first row of each panel, raw data are presented as

both scatter points and the corresponding mean and 95% confidence intervals (CIs) (N = 8

for each treatment, and N = 10 for blank control). In the second row, estimation plots present

the unpaired mean difference between each treatment and the shared control. Circles in

grey and triangles (arrow head up) in black represent neutral and positive effects,

respectively. PFOS, perfluorooctanesulfonic acid; PFOA, perfluorooctanoic acid; PFBS,

perfluorobutanesulfonic acid. Summary of effects of PFAS concentration or type refers to SI

Figure S2 and S3, respectively, and outcomes of ANOVA followed by Dunnett's test is

presented in SI Table S3.

Soil respiration is inhibited by PFAS

We observed that the tested PFAS produced significantly negative effects on soil respiration

in week 3 (Figure 2A and S4), while more variable effects were present in week 6 (Figure

S5). In week 3, PFOA and PFBS at all tested concentrations exerted negative effects on soil

respiration, while effects of PFOS were dependent on its concentration ($F = 5.461$, $p <$

0.001). Irrespective of concentration, our tested PFAS had negative effects on soil

respiration in week 3 (Figure S4C). Compared with soil respiration in week 3, less

pronounced effects were observed in week 6. Similar to effects in week 3, PFBS had the

most apparent effect on respiration in week 6 (Figure S5C). However, this negative effect

was significant only at its highest concentration (500 ng g^{-1}, $p = 0.019$, Table S3).

Waning effects of organic compounds on soil respiration during the incubation period have

been previously reported [32,33]. For example, the fungicides tebuconazole and carbendazim

significantly suppressed soil respiration during the first 30 days, while this effect was no

longer present on the 90th day [34]. Microorganisms would react to the exogenous PFAS

during the first stage, expressed as an inhibitory effect, and gradually recover from this

inhibition during the incubation. Another possible explanation is that as litter decomposition

progressed, easily available carbon (e.g., sugars) was progressively utilized by microbes

and subsequently transferred to CO_2, which modulated the suppressed production of CO_2.

Figure 2. Effects of per- and polyfluoroalkyl substances (PFAS) on soil respiration at the 3rd week (A) and water-stable aggregates (B). In the first row of each panel, raw data are presented as both scatter points and the corresponding mean and 95% confidence intervals (CIs) (N = 8 for each treatment, and N = 10 for blank control). In the second row, estimation plots present the unpaired mean difference between each treatment and the shared control. Circles in grey and triangles (arrow head down) in black represent neutral and negative effects, respectively. PFOS, perfluorooctanesulfonic acid; PFOA, perfluorooctanoic acid; PFBS, perfluorobutanesulfonic acid. Soil respiration at the 6th week refers to SI Figure S5. Summary of effects of PFAS concentration or type refers to SI Figure S4 and S6, and outcomes of ANOVA followed by Dunnett's test is presented in SI Table S3.

Limited effects on soil enzyme and microbial activities

Four enzymes were not significantly affected by individual PFAS treatments (Figures S7–

S10), nor the general microbial activities (Figure S11, Table S3), but regardless of

concentration, PFBS significantly increased β-glucosidase activity (p = 0.029, Figure S9C).

Measuring enzyme activities provides evidence on how soil biochemical processes might be

affected. β-glucosidase is responsible for catalyzing the hydrolysis of cellobiose (a product of

cellulose breakdown) to glucose [35]. We observed a positive trend on β-glucosidase by

PFASs, particularly PFBS, which probably contributed to the litter decomposition by PFASs.

The only significant effect on β-glucosidase corresponded with the most marked effect on

litter decomposition by PFBS. Additionally, there was a significant correlation between

decomposition rate and β-glucosidase activity (r = 0.24, p = 0.004) (Figure S1).

Previous studies reported that soil dehydrogenase (proxy for total microbial activity), urease

and sucrase activities were only insignificantly impacted by PFOA and PFOS with

concentrations lower than 10 µg g^{-1} [12,36]. Cai et al.[13] also reported that microbial activity was

barely affected by PFAS at 100 µg g^{-1} in selected soils. Changes in enzyme activities are

highly dynamic processes [12,37], and thus insignificant effects observed at harvest do not

necessarily indicate that there were no remarkable changes during the incubation. In

addition, there was no significant relationship between microbial activities (FDA) with other

parameters (Figure S1). Overall, general microbial activities were affected only to a very

limited degree.

Different effect sizes caused by three PFAS

The three PFAS examined here appeared to exert similar impact on soil microbes and

functions, but with different effect sizes, which is likely related to their bioavailability and

bioaccumulation [8]. Of the three PFASs, PFBS even at lower concentrations seemed to have

the most remarkable impact on soil respiration, litter decomposition and soil bacterial

abundance, while PFOS had a larger effect size on water-stable aggregates.

Sorption affinities of PFASs followed the order PFOS > PFOA > PFBS on soils with various

soil textures and organic carbon contents, showing the same order of their hydrophobicity [20].

With a low sorption affinity to soil particles, PFBS likely had an increased likelihood to

interact with soil microbes, subsequently causing an impact. However, it is not a simple

effect of hydrophobicity, because, for example, the higher hydrophobicity might result in

higher bioaccumulation and hence exert higher toxicity in soil microorganisms [12,13]. This

might explain the more apparent effect by PFOS on some processes via soil microbes.

Environmental implications and future perspectives

The present results highlight the potential of PFAS to induce changes in soil properties and

functions. Our study comprehensively analyzed PFAS impacts on soil functions, and thus we

hope that our result inspires further studies that consider the impact of PFAS on soil

ecosystem functions.

Within environmentally relevant concentrations, three PFAS, especially the short-chain

PFBS, had a positive effect on litter decomposition in neutral soil. This effect indicates that

the PFAS present in soils now might already affect ecosystem processes. The elevated

decomposition might increase the release of carbon as CO_2, CH_4 and dissolved organic

carbon, affecting carbon sinks in soil [38]. The precise mechanisms underpinning litter

decomposition effects caused by PFAS need to be explored in further studies, with changes

in microbial community composition likely playing a main role. In addition, effects are likely

influenced by soil and PFAS properties.

Soil aggregation is an essential feature of soil structure, principally driven by soil biota and

their interactions [39]. Our finding that certain PFAS negatively affected water-stable

aggregates could indicate far-reaching consequences for soil health, given the many

influences of soil structure on virtually all soil processes. Thus, future studies might explore

these effects on the soil aggregation process in greater depth, including the formation, size

distribution of soil aggregates and their intrinsic connections with soil biota.

We introduce the possibility of PFAS as persistent chemicals being a potential environmental

change factor. The effects of various PFAS on soil functions should now be addressed in the

context of global patterns of contamination.

Acknowledgements

BX thanks the China Scholarship Council and Deutscher Akademischer Austauschdienst

(CSC-DAAD) for a postdoctoral scholarship. MCR acknowledges support from an ERC

Advanced Grant (694368). We thank Daniel Lammel, Yun Liang, Tingting Zhao, and Lili

Rong for their help with experimental measurements. We thank Rosolino Ingraffia for

providing soil samples.

References

(1) Zhu, Y. G.; Penuelas, J. Changes in the Environmental Microbiome in the

Anthropocene. *Glob. Chang. Biol.* **2020**, *26* (6), 3175–3177.

https://doi.org/10.1111/gcb.15086.

(2) Morgado, R. G.; Loureiro, S.; González-Alcaraz, M. N. Changes in Soil Ecosystem

Structure and Functions Due to Soil Contamination. *Soil Pollut. From Monit. to*

Remediat. **2018**, 59–87. https://doi.org/10.1016/B978-0-12-849873-6.00003-0.

(3) Rillig, M. C.; Ryo, M.; Lehmann, A. Classifying Human Influences on Terrestrial

Ecosystems. *Glob. Chang. Biol.* **2021**, gcb.15577. https://doi.org/10.1111/gcb.15577.

(4) Lim, X. Tainted Water: The Scientists Tracing Thousands of Fluorinated Chemicals in

Our Environment. *Nature* **2019**, *566* (7742), 26–29. https://doi.org/10.1038/d41586-

019-00441-1.

(5) Buck, R. C.; Franklin, J.; Berger, U.; Conder, J. M.; Cousins, I. T.; Voogt, P. De;

Jensen, A. A.; Kannan, K.; Mabury, S. A.; van Leeuwen, S. P. J. Perfluoroalkyl and

Polyfluoroalkyl Substances in the Environment: Terminology, Classification, and

Origins. *Integr. Environ. Assess. Manag.* **2011**, *7* (4), 513–541.

https://doi.org/10.1002/ieam.258.

(6) Beans, C. News Feature: How "Forever Chemicals" Might Impair the Immune System.

Proc. Natl. Acad. Sci. **2021**, *118* (15). https://doi.org/10.1073/PNAS.2105018118.

(7) UNEP. Stockholm Convention on Persistent Organic Pollutants (POPS) - Texts and

Annexes. *United Nations Environ. Program.* **2019**.

(8) Cai, Y.; Wang, Q.; Zhou, B.; Yuan, R.; Wang, F.; Chen, Z.; Chen, H. A Review of

Responses of Terrestrial Organisms to Perfluorinated Compounds. *Sci. Total Environ.*

2021, *793*. https://doi.org/10.1016/j.scitotenv.2021.148565.

(9)   Ma, D.; Zhong, H.; Lv, J.; Wang, Y.; Jiang, G. Levels, Distributions, and Sources of

Legacy and Novel per- and Perfluoroalkyl Substances (PFAS) in the Topsoil of Tianjin,

China. *J. Environ. Sci.* **2022**, *112*, 71–81. https://doi.org/10.1016/J.JES.2021.04.029.

(10)   Brusseau, M. L.; Anderson, R. H.; Guo, B. PFAS Concentrations in Soils: Background

Levels versus Contaminated Sites. *Sci. Total Environ.* **2020**, *740*, 140017.

https://doi.org/10.1016/j.scitotenv.2020.140017.

(11)   Jin, H.; Zhang, Y.; Zhu, L.; Martin, J. W. Isomer Profiles of Perfluoroalkyl Substances

in Water and Soil Surrounding a Chinese Fluorochemical Manufacturing Park. *Environ.*

Sci. Technol. **2015**, *49* (8), 4946–4954. https://doi.org/10.1021/acs.est.5b00212.

(12)   Qiao, W.; Xie, Z.; Zhang, Y.; Liu, X.; Xie, S.; Huang, J.; Yu, L. Perfluoroalkyl

Substances (PFASs) Influence the Structure and Function of Soil Bacterial

Community: Greenhouse Experiment. *Sci. Total Environ.* **2018**, *642*, 1118–1126.

https://doi.org/10.1016/j.scitotenv.2018.06.113.

(13)   Cai, Y.; Chen, H.; Yuan, R.; Wang, F.; Chen, Z.; Zhou, B. Toxicity of Perfluorinated

Compounds to Soil Microbial Activity: Effect of Carbon Chain Length, Functional

Group and Soil Properties. *Sci. Total Environ.* **2019**, *690*, 1162–1169.

https://doi.org/10.1016/j.scitotenv.2019.06.440.

(14)   Xu, R.; Tao, W.; Lin, H.; Huang, D.; Su, P.; Gao, P.; Sun, X.; Yang, Z.; Sun, W.

Effects of Perfluorooctanoic Acid (PFOA) and Perfluorooctane Sulfonic Acid (PFOS)

on Soil Microbial Community. *Microb. Ecol.* **2021**. https://doi.org/10.1007/s00248-021-

01808-6.

(15)   Cai, Y.; Chen, H.; Yuan, R.; Wang, F.; Chen, Z.; Zhou, B. Metagenomic Analysis of

Soil Microbial Community under PFOA and PFOS Stress. *Environ. Res.* **2020**, *188*

(May), 109838. https://doi.org/10.1016/j.envres.2020.109838.

(16)   Chen, H.; Wang, Q.; Cai, Y.; Yuan, R.; Wang, F.; Zhou, B.; Chen, Z. Effect of per Fl

Uorooctanoic Acid on Microbial Activity in Wheat Soil under Different Fertilization

Conditions. *Environ. Pollut.* **2020**, *264*, 114784.

https://doi.org/10.1016/j.envpol.2020.114784.

(17)   Rillig, M. C.; Mardatin, N. F.; Leifheit, E. F.; Antunes, P. M. Mycelium of Arbuscular

Mycorrhizal Fungi Increases Soil Water Repellency and Is Sufficient to Maintain

Water-Stable Soil Aggregates. *Soil Biol. Biochem.* **2010**, *42* (7), 1189–1191.

https://doi.org/10.1016/j.soilbio.2010.03.027.

(18)   Lehmann, A.; Leifheit, E. F.; Feng, L.; Bergmann, J.; Wulf, A.; Rillig, M. C.

Microplastic Fiber and Drought Effects on Plants and Soil Are Only Slightly Modified

by Arbuscular Mycorrhizal Fungi. *Soil Ecol. Lett.* **2020**.

https://doi.org/10.1007/s42832-020-0060-4.

(19)   Ahmed, M. B.; Johir, M. A. H.; McLaughlan, R.; Nguyen, L. N.; Xu, B.; Nghiem, L. D.

Per- and Polyfluoroalkyl Substances in Soil and Sediments: Occurrence, Fate,

Remediation and Future Outlook. *Sci. Total Environ.* **2020**, *748*, 141251.

https://doi.org/10.1016/j.scitotenv.2020.141251.

(20)   Milinovic, J.; Lacorte, S.; Vidal, M.; Rigol, A. Sorption Behaviour of Perfluoroalkyl

Substances in Soils. *Sci. Total Environ.* **2015**, *511*, 63–71.

https://doi.org/10.1016/j.scitotenv.2014.12.017.

(21)   Rillig, M. C.; Ryo, M.; Lehmann, A.; Aguilar-Trigueros, C. A.; Buchert, S.; Wulf, A.;

Iwasaki, A.; Roy, J.; Yang, G. The Role of Multiple Global Change Factors in Driving

Soil Functions and Microbial Biodiversity. *Science (80-.).* **2019**, *366* (6467), 886–890.

https://doi.org/10.1126/science.aay2832.

(22)   Kemper, W. D.; Rosenau, R. C. Aggregate Stability and Size Dlstributlon. *Methods*

soil Anal. Part 1 Phys. Mineral. Methods **1986**.

(23) Liang, Y.; Lehmann, A.; Ballhausen, M.-B.; Muller, L.; Rillig, M. C. Increasing

Temperature and Microplastic Fibers Jointly Influence Soil Aggregation by Saprobic

Fungi. *Front. Microbiol.* **2019**, *10* (September), 1–10.

https://doi.org/10.3389/fmicb.2019.02018.

(24) R Core Team. R: A Language and Environment for Statistical Computing. Vienna

2020.

(25) Ho, J.; Tumkaya, T.; Aryal, S.; Choi, H.; Claridge-Chang, A. Moving beyond P Values:

Data Analysis with Estimation Graphics. *Nat. Methods* **2019**, *16* (7), 565–566.

https://doi.org/10.1038/s41592-019-0470-3.

(26) Hothorn, T.; Bretz, F.; Westfall, P. Simultaneous Inference in General Parametric

Models. *Biometrical Journal*. John Wiley & Sons, Ltd June 1, 2008, pp 346–363.

https://doi.org/10.1002/bimj.200810425.

(27) Wei, T.; Simko, V. R Package "Corrplot": Visualization of a Correlation Matrix. 2017.

(28) Xu, B.; Yang, G.; Lehmann, A.; Hasenstab-Riedel, S.; Rillig, M. C. PFAS Effects on

Soil Structure and Function. **2021**. https://doi.org/10.6084/m9.figshare.16967401.v1.

(29) Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*; Springer-Verlag: New York,

2016.

(30) Valentín, L.; Nousiainen, A.; Mikkonen, A. Introduction to Organic Contaminants in

Soil: Concepts and Risks; 2013; pp 1–29. https://doi.org/10.1007/698_2012_208.

(31) Rousk, J.; Brookes, P. C.; Bååth, E. Investigating the Mechanisms for the Opposing

PH Relationships of Fungal and Bacterial Growth in Soil. *Soil Biol. Biochem.* **2010**, *42*

(6), 926–934. https://doi.org/10.1016/j.soilbio.2010.02.009.

(32) Zhang, W.; Wang, J.; Wang, J.; Zhu, L.; Lv, N.; Wang, R.; Ahmad, Z. New Insights

into Dose- and Time-Dependent Response of Five Typical PPCPs on Soil Microbial

Respiration. *Bull. Environ. Contam. Toxicol.* **2019**, *103* (1), 193–198.

https://doi.org/10.1007/s00128-019-02655-5.

(33) Xiong, D.; Li, Y.; Xiong, Y.; Li, X.; Xiao, Y.; Qin, Z.; Xiao, Y. Influence of Boscalid on

the Activities of Soil Enzymes and Soil Respiration. *Eur. J. Soil Biol.* **2014**, *61*, 1–5.

https://doi.org/10.1016/j.ejsobi.2013.12.006.

(34) Wang, C.; Wang, F.; Zhang, Q.; Liang, W. Individual and Combined Effects of

Tebuconazole and Carbendazim on Soil Microbial Activity. *Eur. J. Soil Biol.* **2016**, *72*,

6–13. https://doi.org/10.1016/j.ejsobi.2015.12.005.

(35) German, D. P.; Weintraub, M. N.; Grandy, A. S.; Lauber, C. L.; Rinkes, Z. L.; Allison,

S. D. Optimization of Hydrolytic and Oxidative Enzyme Methods for Ecosystem

Studies. *Soil Biol. Biochem.* **2011**, *43* (7), 1387–1397.

https://doi.org/10.1016/j.soilbio.2011.03.017.

(36) He, W.; Megharaj, M.; Naidu, R. Toxicity of Perfluorooctanoic Acid towards

Earthworm and Enzymatic Activities in Soil. *Environ. Monit. Assess.* **2016**, *188* (7).

https://doi.org/10.1007/s10661-016-5416-y.

(37) Zhao, T.; Lozano, Y. M.; Rillig, M. C. Microplastics Increase Soil PH and Decrease

Microbial Activities as a Function of Microplastic Shape, Polymer Type, and Exposure

Time. *Front. Environ. Sci.* **2021**, *9*, 1. https://doi.org/10.3389/fenvs.2021.675803.

(38) Dieleman, C. M.; Lindo, Z.; McLaughlin, J. W.; Craig, A. E.; Branfireun, B. A. Climate

Change Effects on Peatland Decomposition and Porewater Dissolved Organic Carbon

Biogeochemistry. *Biogeochemistry* **2016**, *128* (3), 385–396.

https://doi.org/10.1007/s10533-016-0214-8.

(39) Lehmann, A.; Zheng, W.; Rillig, M. C. Soil Biota Contributions to Soil Aggregation. *Nat.*

Ecol. Evol. **2017**, *1* (12), 1828–1835. https://doi.org/10.1038/s41559-017-0344-y.

389