Effects of perfluoroalkyl and polyfluoroalkyl substances on soil

structure and function

Baile Xu* , Gaowen Yang, Anika Lehmann, Matthias C. Rillig

Institute of Biology, Free University of Berlin, D-14195 Berlin, Germany

Berlin-Brandenburg Institute of Advanced Biodiversity Research, D-14195 Berlin, Germany

* Corresponding author. E-mail: bxu@zedat.fu-berlin.de

Abstract Art

Abstract

Soils are impacted at a global scale by several anthropogenic factors, including chemical

pollutants. Among those, perfluoroalkyl and polyfluoroalkyl substances (PFAS) are of

concern due to their high environmental persistence, and as they might affect soil health and

functions. However, data on impacts of PFASs on soil structure and microbially-driven

processes are currently lacking. This study explored the effects of perfluorooctanesulfonic

acid (PFOS), perfluorooctanoic acid (PFOA) and perfluorobutanesulfonic acid (PFBS) at

environmental-relevant nominal concentrations ($1 \sim 1000$ ng g^{-1}) on soil functions, using a 6-

week microcosm experiment. We measured soil respiration, litter decomposition, enzyme

and microbial activities, soil aggregates, and bacterial abundance. PFAS (even at 1 ng g^{-1}

for PFBS) significantly increased the litter decomposition, associated with positive effects on

bacterial abundance, and β-glucosidase activities. This effect increased with PFAS

concentrations. Soil respiration was significantly inhibited by PFAS in the 3rd week, while this

effect was more variable in week 6. Water-stable aggregates were negatively affected by

PFOS and PFOA, possibly related to microbial shifts. The general microbial activities and β-

D-cellobiosidase and phosphatase activities were barely affected by PFAS treatments. Our

work highlights the potential effects of PFAS on soil health, and we argue that this substance

class could be a factor of environmental change of potentially broad relevance in terrestrial

ecosystem functioning.

Keywords: Litter decomposition; Soil respiration; Water-stable aggregates; Soil bacteria;

Perfluorobutanesulfonate (PFBS); Perfluorooctanesulfonic acid (PFOS); Perfluorooctanoic

acid (PFOA); Enzyme activities

Synopsis: PFAS are likely to affect soil health.

Introduction

Human activity is progressively and fundamentally affecting the Earth's surface, including soils, which operate at the interface between biosphere, hydrosphere, atmosphere, and lithosphere, and are suffering from physical, chemical, and biological stressors related to anthropogenic activities [1,2]. One significant category of these influences is chemical pollution, which is widely recognized as a global change factor [1,3]. Perfluoroalkyl and polyfluoroalkyl substances (PFAS) are a highly diverse family of chemicals of concern. These compounds contain at least one perfluoroalkyl moiety ($C_nF_{2n+1}-$) [4], and the high-energy carbon-fluorine (C-F) bonds make them extremely resistant to break down, and subsequently persistent in the environment. Therefore, they are also called "forever chemicals" in public discourse [5]. Certain PFAS, perfluorooctanesulfonic acid (PFOS) and perfluorooctanoic acid (PFOA) are on the list and on the waiting list under the Stockholm Convention [6], respectively.

Soil is known as a major sink for persistent organic chemicals in the environment. There are many pathways for PFAS entering the soil environment. Typically, fluoride factory emission, sludge application, the degradation of aqueous film-forming foam, and landfills contribute direct sources, and atmospheric deposition and runoff constitute non-point sources [7,8]. PFAS have been widely detected in soils with a broad range of concentrations. Generally, PFAS concentrations in non-hotspot soil are lower than 300 ng g^{-1}, while in hotspots of PFAS-contaminated soil, they can be as high as several or even tens of μg g^{-1}, mostly dominated by PFOA and PFOS [9]. Jin et al. [10] also reported that perfluorobutanesulfonic acid (PFBS) was a prominent type of PFAS in the soil around a fluoride-factory park.

Existing evidence has demonstrated that PFAS can exert some impacts on soil functions, in particular on soil enzymes, and microbial activities and communities, and these influences were closely related to properties of soil and PFASs. According to the number of carbon atoms, perfluoroalkyl carboxylic acids with 7 or more and perfluoroalkyl sulfonate with 6 or more carbons are categorized as long-chain PFAS, otherwise, short-chain PFAS [4]. Short-chain PFBS might activate sucrase and urease activities, while long-chain PFOS might

reduce these activities in soil [11]. Cai et al. [12] showed that PFAS with sulfonic groups and

longer chains had higher toxicity to soil microbial activities, and that soil with higher organic

matter content and higher pH (neutral) exhibited lower impact by PFAS. Differences in

sorption affinity of PFAS to soil are likely to regulate the impact of PFASs on soil microbes[7].

Changes in structure and function of microbial communities by PFAS were shown by

previous research [11,13–15], and these shifts were suggested to affect soil processes and

ecosystem functions. However, the implications on process rates in soil, for example, litter

decomposition and soil aggregation, were not addressed.

Given the persistent nature of PFAS it is important to explore if these chemicals can affect

soil process rates and properties. Here, we investigate the PFAS effects on soil processes,

including soil respiration, litter decomposition, soil aggregation, enzyme and microbial

activities, as well as bacterial abundance. We discuss the environmental implications of our

results and highlight the possibility of PFAS to be considered as a global change factor of

importance in terrestrial ecosystems.

Materials and Methods

Test soil and PFAS

The test soil was Albic Luvisol collected at the agricultural field station of Freie Universität

Berlin in December 2020. The soil has a sandy loam texture (73.6% sand, 18.8% silty and

7.6% clay), with 1.87% total C, 0.12% total N and a soil pH of 7.1 [16,17]. Fresh soil samples

were thoroughly mixed, passed through a 2-mm sieve, and then stored at 4°C.

Three PFAS, namely PFOS, PFOA, and PFBS were selected in this study due to their wide

occurrence in the soil environment [18]. Values of their Log K_{ow} are 5.26, 4.59 and 2.73,

respectively [19], and their chemical structures and other physicochemical properties are listed

in Supporting Information (SI) Table S1.

Experimental setup

PFAS standards were dissolved in sterilized deionized water to prepare the stock solution

with the concentration of 100 mg L^{-1}. A five-gram portion of previously sterilized soil samples

(121°C, 20 min, twice) was supplemented with appropriate doses of PFAS in solution (we

used this sterilized 'loading soil' to avoid any exaggerated effects on soil communities [20]),

and thoroughly mixed this soil with 25 g of soil by manual stirring for ~2 mins. A total of 30 g

soil was placed in 50-mL mini-bioreactor tubes (Corning Inc., Corning, USA) with vented lids

to establish experimental microcosms. Each tube was watered to 70% soil water holding

capacity (WHC) with deionized water. The nominal concentrations of PFAS in soil were 1, 10,

100 and 1000 ng g^{-1}, corresponding to the environmental-relevant levels [9]. Although the

actual concentrations might differ from the nominal concentrations of PFAS in soil, this

difference would be minor and not impact the understanding of soil response to gradients of

PFAS concentrations. Tubes were randomly placed in a dark temperature-controlled

incubator at 20°C for 6 weeks, and each tube was watered weekly to maintain soil moisture.

This experiment ran with 10 replications of blank control (without any PFAS added, but

handled exactly the same way) and 8 replications of each treatment, for a total of 106

microcosms.

Proxies for soil health and function

We measured well-established proxies for soil health and functions, including soil respiration,

litter decomposition, enzyme activities, soil aggregates, and soil bacterial abundance. Soil

respiration was measured with an infrared gas analyzer (LI-6400XT, LI-COR Inc., Bad

Homburg, Germany), and litter decomposition was determined by the mass loss of tea bags

[17]. Soil enzymes activities were measured, including four enzymes concerning C (β-

glucosidase and β-D-1,4-cellobiosidase), N (β-1,4-N-acetyl-glucosaminidase), and P

(phosphatase) cycling, and fluorescein diacetate hydrolase (FDA) representing general soil

microbial activity. Water-stable aggregates, as the basic unit of soil structure, were qualified

using a wet-sieving apparatus (Eijkelkamp, Giesbeek, Netherlands) with the reported

method [21,22]. Soil DNA was extracted with DNeasy PowerSoil Pro Kit (QIAGEN GmbH,

Germany) following the technical protocol, and we amplified using the universal primers

515F (5'-GTGCCAGCMGCCGCGGTAA-3') and 806R (5'-GGACTACHVGGGTWTCTAAT-3')

with quantitative polymerase chain reactions (qPCR) in a CFX 96 Real-Time System (Bio-

Rad Lab., Hercules, USA). For more information on measurement procedures, qPCR

conditions and quality control, see SI Text 1.

Statistical analysis

All statistical analyses and data visualization were performed in R [23]. The effects of PFAS

treatment (four concentrations per PFAS type) on soil functions were tested with a two-step

method. Firstly, we calculated the 95% confidence interval (CI) of unpaired mean differences

(treatment minus control) using the R package "dabestr" [24]. This approach focuses on the

effect size and its precision, and can avoid the pitfalls of significance testing. Secondly, one-

way analysis of variance (ANOVA) followed by Dunnett's test in the R package "multcomp"

was implemented to compare each treatment with the control [25]. Spearman correlations

among PFAS concentrations and soil structure and function were performed with the

package "corrplot" [26]. Adjusted p values by a single-step method are reported in the SI Table

S2. All plots were generated with the package "ggplot2" [27].

Results and Discussion

PFAS increased litter decomposition associated with bacterial abundance

Positive effects of three PFASs on litter decomposition were observed (Figure 1A), and

PFBS, particularly, at all tested concentrations significantly increased the decomposition rate

($p < 0.05$, Table S2). Regardless of tested type, PFAS at each concentration positively

affected the decomposition, and increasing concentrations significantly enhanced this effect

($p = 0.0222$, Figure S1 and S2). In terms of PFAS type, PFOA and PFBS resulted in

significantly positive effects on litter decomposition ($p < 0.01$, Table S2), and PFBS exerted

the most remarkable positive effect (Figure S1C).

Figure 1. Effects of per- and polyfluoroalkyl substances (PFASs) on litter decomposition (A) and bacterial abundance (B) in soil. In the first row of each panel, raw data are presented as both scatter points and the corresponding mean and 95% confidence intervals (CIs) (N = 8 for each treatment, and N = 10 for blank control). In the second row, estimation plots present the unpaired mean difference between each treatment and the shared control. Circles in grey and triangles (arrow head up) in black represent neutral and positive effects, respectively. PFOS, perfluorooctanesulfonic acid; PFOA, perfluorooctanoic acid; PFBS, perfluorobutanesulfonic acid. Summary of effects of PFAS concentration or type refers to SI Figure S2 and S3, respectively, and outcomes of ANOVA followed by Dunnett's test is presented in SI Table S2.

The treatment of PFASs significantly increased soil bacterial abundance (F = 1.868, p =

0.0485), although the multiple comparison tests did not reveal a significant difference (Table

S2). PFBS seems to have the most obvious effect among three PFAS, showing a positive

effect with concentrations higher than 1 ng g^{-1} (Figure 1B). Independently of its concentration

or type, PFAS treatment had a positive effect on soil bacterial abundance in terms of copy

number per gram of dry soil (Figure S3B and S3C).

It is widely accepted that microorganisms, including bacteria and fungi, play an essential role

in the biological decomposition of organic matter in soil, in which bacteria govern in neutral

or alkaline soils, while fungi do in acidic soil [28,29]. Correspondingly, we found that in our

neutral soil, with various PFAS treatments, the number of bacteria rose compared to the

control (Figure 1B), and there was a trend for a correlation between bacteria abundance and

litter decomposition rate (F=2.705, p = 0.103). The increased bacterial populations probably

contributed to the increased decomposition rate of organic matter.

Litter decomposition and the ensuing nutrient release, governing carbon and nutrient cycling,

is a key process in terrestrial ecosystems. Our results showed that PFAS had a positive

effect on litter decomposition, and particularly the PFBS treatment, even at 1 ng g^{-1} in soil,

resulting in a significant enhancement of litter decomposition, likely associated with

increased soil bacterial abundance.

Soil respiration is inhibited by PFAS

We observed that PFAS produced significantly negative effects on soil respiration in week 3

(Figure 2A and S4), while more variable effects were present in week 6 (Figure S5). In week

3, PFOA and PFBS at all tested concentrations exerted negative effects on soil respiration,

while effects of PFOS were dependent on its concentration (F = 5.461, p < 0.001).

Irrespective of concentration or type, PFAS had negative effects on soil respiration in week 3

(Figure S4B and S4C). Compared with soil respiration in week 3, less pronounced effects

were observed in week 6. Similar to effects in week 3, PFBS had the most apparent effect

on respiration in week 6 (Figure S5C). However, this negative effect was significant only at

its highest concentration (1000 ng g^{-1}, p = 0.019, Table S2).

Figure 2. Effects of per- and polyfluoroalkyl substances (PFASs) on soil respiration at the 3rd

week (A) and water-stable aggregates (B). In the first row of each panel, raw data are

presented as both scatter points and the corresponding mean and 95% confidence intervals

(CIs) (N = 8 for each treatment, and N = 10 for blank control). In the second row, estimation

plots present the unpaired mean difference between each treatment and the shared control.

Circles in grey and triangles (arrow head down) in black represent neutral and negative

effects, respectively. PFOS, perfluorooctanesulfonic acid; PFOA, perfluorooctanoic acid;

PFBS, perfluorobutanesulfonic acid. Soil respiration at the 6th week refer to SI Figure S5.

Comprehensive comparisons irrespective PFAS concentration or type refer to SI Figure S4

and S6, and outcomes of ANOVA followed by Dunnett's test is presented in SI Table S2.

Waning effects by organic compounds on soil respiration during the incubation period have

been previously reported [31,32]. For example, the fungicides tebuconazole and carbendazim

significantly suppressed soil respiration during the first 30 days, while this effect was no

longer present on the 90[th] day [33]. Microorganisms would react to the exogenous PFAS

during the first stage, expressed as the inhibitory effect, and gradually recover from this

inhibition during the incubation. Another possible explanation is that as litter decomposition

progressed, easily available carbon (e.g., sugars) was progressively utilized by microbes

and subsequently transferred to CO_2, which modulated the suppressed production of CO_2.

Long-chain PFOS suppressed water-stable aggregates

Although effects of PFOS and PFOA treatments on soil aggregate stability at the single

concentration were insignificant (Figure 2B, Table S2), PFOS treatment significantly

decreased water-stable aggregates irrespective of concentrations ($p = 0.044$, Figure S6C).

Fungi are likely to play more important roles in the formation of macroaggregate (250~2000

μm), whilst bacteria contribute more to microaggregate stability (53~250 μm) [34]. Although

fungal abundance was not measured, based on discussion above, we speculate that long-

chain PFOA and PFOS might suppress fungi and hence negatively affect the stability of soil

aggregates.

Limited effects on soil enzyme and microbial activities

Four enzymes were not significantly affected by PFAS treatments (Figures S7–S10), nor the

general microbial activities (Figure S11, Table S2), although there were positive trends on β-

glucosidase and β-glucosaminidase activities (Figures S9 and S10).

Measuring enzyme activities provides evidence on soil biochemical processes. β-

glucosidase is responsible for catalyzing the hydrolysis of cellobiose (the product of cellulose)

to glucose [37]. We observed a positive trend on β-glucosidase by PFASs, particularly PFBS,

which partly contributed to the litter decomposition by PFASs. The only positive effect on β-

glucosidase corresponded with the most marked effect on litter decomposition by PFBS.

Additionally, there was a significant correlation between decomposition rate and β-

glucosidase activity (R^2 = 0.24, p = 0.004) (Figure S1).

Previous studies reported that soil dehydrogenase (proxy for total microbial activity), urease

and sucrase activities were only insignificantly impacted by PFOA and PFOS with

concentrations lower than 10 µg g^{-1} [11,38]. Cai et al.[12] also reported that microbial activity was

barely affected by PFAS at 100 µg g^{-1} in selected soils. Changes in enzyme activities are

highly dynamic processes [11,39], and thus insignificant effects observed at harvest do not

necessarily indicate that there were no remarkable changes during the incubation. In

addition, there was no significant relationship between microbial activities (FDA) with other

parameters (Figure S1). Overall, general microbial activities were affected only to a very

limited degree.

Different effect sizes caused by three PFAS

The three PFAS examined here appeared to exert similar impact on soil microbes and

functions, but with different effect sizes, which is likely related to their bioavailability and

bioaccumulation [7]. Of the three PFASs, PFBS seemed to have the most remarkable impact

on soil respiration, litter decomposition and soil bacterial abundance, while PFOS had a

larger effect size on water-stable aggregates.

Sorption affinities of PFASs followed the order PFOS > PFOA > PFBS on soils with various

soil textures and organic carbon contents, showing the same order of their hydrophobicity [19].

With a low sorption affinity to soil particles, PFBS likely had an increased likelihood to

interact with soil microbes, subsequently causing an impact. However, it is not a simple

effect of hydrophobicity, because, for example, the higher hydrophobicity might result in

higher bioaccumulation and hence exert higher toxicity in soil microorganisms [11,12]. This

might explain the more apparent effect by PFOS on some processes via soil microbes.

Environmental implications and future perspectives

The present results highlight the potential of PFAS to induce changes in soil properties and

functions. Our study comprehensively analyzed PFAS impacts on soil functions, and thus we

hope that our result inspires further studies that consider PFAS impact on soil ecosystem

functions.

Within environmentally relevant concentrations, three PFASs, especially the short-chain

PFBS, had a positive effect on litter decomposition ratio in neutral soil. This effect indicates

that the presence of PFAS might already affect ecosystem processes. The elevated

decomposition might increase the release of carbon as CO_2, CH_4 and dissolved organic

carbon, affecting carbon sinks in soil [40]. The precise mechanisms underpinning litter

decomposition effects caused by PFAS need to be explored in further studies, with changes

in microbial community composition likely playing a main role. In addition, effects are likely

closely associated with soil and PFAS properties.

Soil aggregation is an essential feature of soil structure, principally driven by soil biota and

their interactions [41]. Our finding that certain PFAS negatively affected water-stable

aggregates could indicate far-reaching consequences for soil health, given the many

influences of soil structure on virtually all soil processes. Thus, future studies might explore

these effects on the soil aggregation process in greater depth, including the formation, size

distribution of soil aggregates and their intrinsic connections with soil biota.

We introduce the possibility of PFAS as persistent chemicals being a potential environmental

change factor. The effects of various PFAS on soil functions should now be addressed in the

context of global patterns of contamination.

Acknowledgements

BX thanks the China Scholarship Council and Deutscher Akademischer Austauschdienst

(CSC-DAAD) for a postdoc scholarship. MCR acknowledges support from an ERC

Advanced Grant (694368). We thank Daniel Lammel, Yun Liang, Tingting Zhao, and Lili

Rong for their helps with experimental measurements. We thank Rosolino Ingraffia for

providing soil samples.

References

(1)   Zhu, Y. G.; Penuelas, J. Changes in the Environmental Microbiome in the

272   Anthropocene. *Glob. Chang. Biol.* **2020**, *26* (6), 3175–3177.

273   https://doi.org/10.1111/gcb.15086.

274   (2)   Morgado, R. G.; Loureiro, S.; González-Alcaraz, M. N. Changes in Soil Ecosystem

275   Structure and Functions Due to Soil Contamination. *Soil Pollut. From Monit. to*

276   *Remediat.* **2018**, 59–87. https://doi.org/10.1016/B978-0-12-849873-6.00003-0.

277   (3)   Rillig, M. C.; Ryo, M.; Lehmann, A. Classifying Human Influences on Terrestrial

278   Ecosystems. *Glob. Chang. Biol.* **2021**, gcb.15577. https://doi.org/10.1111/gcb.15577.

279   (4)   Buck, R. C.; Franklin, J.; Berger, U.; Conder, J. M.; Cousins, I. T.; Voogt, P. De;

280   Jensen, A. A.; Kannan, K.; Mabury, S. A.; van Leeuwen, S. P. J. Perfluoroalkyl and

281   Polyfluoroalkyl Substances in the Environment: Terminology, Classification, and

282   Origins. *Integr. Environ. Assess. Manag.* **2011**, *7* (4), 513–541.

283   https://doi.org/10.1002/ieam.258.

284   (5)   Beans, C. News Feature: How "Forever Chemicals" Might Impair the Immune System.

285   *Proc. Natl. Acad. Sci.* **2021**, *118* (15). https://doi.org/10.1073/PNAS.2105018118.

286   (6)   UNEP. Stockholm Convention on Persistent Organic Pollutants (POPS) - Texts and

287   Annexes. *United Nations Environ. Program.* **2019**.

288   (7)   Cai, Y.; Wang, Q.; Zhou, B.; Yuan, R.; Wang, F.; Chen, Z.; Chen, H. A Review of

289   Responses of Terrestrial Organisms to Perfluorinated Compounds. *Sci. Total Environ.*

290   **2021**, *793*. https://doi.org/10.1016/j.scitotenv.2021.148565.

291   (8)   Ma, D.; Zhong, H.; Lv, J.; Wang, Y.; Jiang, G. Levels, Distributions, and Sources of

292   Legacy and Novel per- and Perfluoroalkyl Substances (PFAS) in the Topsoil of Tianjin,

293   China. *J. Environ. Sci.* **2022**, *112*, 71–81. https://doi.org/10.1016/J.JES.2021.04.029.

(9)   Brusseau, M. L.; Anderson, R. H.; Guo, B. PFAS Concentrations in Soils: Background

Levels versus Contaminated Sites. *Sci. Total Environ.* **2020**, *740*, 140017.

https://doi.org/10.1016/j.scitotenv.2020.140017.

(10) Jin, H.; Zhang, Y.; Zhu, L.; Martin, J. W. Isomer Profiles of Perfluoroalkyl Substances

in Water and Soil Surrounding a Chinese Fluorochemical Manufacturing Park. *Environ.*

Sci. Technol. **2015**, *49* (8), 4946–4954. https://doi.org/10.1021/acs.est.5b00212.

(11) Qiao, W.; Xie, Z.; Zhang, Y.; Liu, X.; Xie, S.; Huang, J.; Yu, L. Perfluoroalkyl

Substances (PFASs) Influence the Structure and Function of Soil Bacterial

Community: Greenhouse Experiment. *Sci. Total Environ.* **2018**, *642*, 1118–1126.

https://doi.org/10.1016/j.scitotenv.2018.06.113.

(12) Cai, Y.; Chen, H.; Yuan, R.; Wang, F.; Chen, Z.; Zhou, B. Toxicity of Perfluorinated

Compounds to Soil Microbial Activity: Effect of Carbon Chain Length, Functional

Group and Soil Properties. *Sci. Total Environ.* **2019**, *690*, 1162–1169.

https://doi.org/10.1016/j.scitotenv.2019.06.440.

(13) Xu, R.; Tao, W.; Lin, H.; Huang, D.; Su, P.; Gao, P.; Sun, X.; Yang, Z.; Sun, W.

Effects of Perfluorooctanoic Acid (PFOA) and Perfluorooctane Sulfonic Acid (PFOS)

on Soil Microbial Community. *Microb. Ecol.* **2021**. https://doi.org/10.1007/s00248-021-

01808-6.

(14) Cai, Y.; Chen, H.; Yuan, R.; Wang, F.; Chen, Z.; Zhou, B. Metagenomic Analysis of

Soil Microbial Community under PFOA and PFOS Stress. *Environ. Res.* **2020**, *188*

(May), 109838. https://doi.org/10.1016/j.envres.2020.109838.

(15) Chen, H.; Wang, Q.; Cai, Y.; Yuan, R.; Wang, F.; Zhou, B.; Chen, Z. Effect of per Fl

Uorooctanoic Acid on Microbial Activity in Wheat Soil under Different Fertilization

Conditions. *Environ. Pollut.* **2020**, *264*, 114784.

https://doi.org/10.1016/j.envpol.2020.114784.

(16) Rillig, M. C.; Mardatin, N. F.; Leifheit, E. F.; Antunes, P. M. Mycelium of Arbuscular

Mycorrhizal Fungi Increases Soil Water Repellency and Is Sufficient to Maintain

Water-Stable Soil Aggregates. *Soil Biol. Biochem.* **2010**, *42* (7), 1189–1191.

https://doi.org/10.1016/j.soilbio.2010.03.027.

(17) Lehmann, A.; Leifheit, E. F.; Feng, L.; Bergmann, J.; Wulf, A.; Rillig, M. C.

Microplastic Fiber and Drought Effects on Plants and Soil Are Only Slightly Modified

by Arbuscular Mycorrhizal Fungi. *Soil Ecol. Lett.* **2020**.

https://doi.org/10.1007/s42832-020-0060-4.

(18) Ahmed, M. B.; Johir, M. A. H.; McLaughlan, R.; Nguyen, L. N.; Xu, B.; Nghiem, L. D.

Per- and Polyfluoroalkyl Substances in Soil and Sediments: Occurrence, Fate,

Remediation and Future Outlook. *Sci. Total Environ.* **2020**, *748*, 141251.

https://doi.org/10.1016/j.scitotenv.2020.141251.

(19) Milinovic, J.; Lacorte, S.; Vidal, M.; Rigol, A. Sorption Behaviour of Perfluoroalkyl

Substances in Soils. *Sci. Total Environ.* **2015**, *511*, 63–71.

https://doi.org/10.1016/j.scitotenv.2014.12.017.

(20) Rillig, M. C.; Ryo, M.; Lehmann, A.; Aguilar-Trigueros, C. A.; Buchert, S.; Wulf, A.;

Iwasaki, A.; Roy, J.; Yang, G. The Role of Multiple Global Change Factors in Driving

Soil Functions and Microbial Biodiversity. *Science (80-.).* **2019**, *366* (6467), 886–890.

https://doi.org/10.1126/science.aay2832.

(21) Kemper, W. D.; Rosenau, R. C. Aggregate Stability and Size DIstributIon. *Methods*

soil Anal. Part 1 Phys. Mineral. Methods **1986**.

(22) Liang, Y.; Lehmann, A.; Ballhausen, M.-B.; Muller, L.; Rillig, M. C. Increasing

Temperature and Microplastic Fibers Jointly Influence Soil Aggregation by Saprobic

Fungi. *Front. Microbiol.* **2019**, *10* (September), 1–10.

https://doi.org/10.3389/fmicb.2019.02018.

(23) R Core Team. R: A Language and Environment for Statistical Computing. Vienna

2020.

(24)   Ho, J.; Tumkaya, T.; Aryal, S.; Choi, H.; Claridge-Chang, A. Moving beyond P Values:

Data Analysis with Estimation Graphics. *Nat. Methods* **2019**, *16* (7), 565–566.

https://doi.org/10.1038/s41592-019-0470-3.

(25)   Hothorn, T.; Bretz, F.; Westfall, P. Simultaneous Inference in General Parametric

Models. *Biometrical Journal*. John Wiley & Sons, Ltd June 1, 2008, pp 346–363.

https://doi.org/10.1002/bimj.200810425.

(26)   Wei, T.; Simko, V. R Package "Corrplot": Visualization of a Correlation Matrix. 2017.

(27)   Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*; Springer-Verlag: New York,

2016.

(28)   Valentín, L.; Nousiainen, A.; Mikkonen, A. Introduction to Organic Contaminants in

Soil: Concepts and Risks; 2013; pp 1–29. https://doi.org/10.1007/698_2012_208.

(29)   Rousk, J.; Brookes, P. C.; Bååth, E. Investigating the Mechanisms for the Opposing

PH Relationships of Fungal and Bacterial Growth in Soil. *Soil Biol. Biochem.* **2010**, *42*

(6), 926–934. https://doi.org/10.1016/j.soilbio.2010.02.009.

(30)   Munier-Lamy, C.; Borde, O. Effect of a Triazole Fungicide on the Cellulose

Decomposition by the Soil Microflora. *Chemosphere* **2000**, *41* (7), 1029–1035.

https://doi.org/10.1016/S0045-6535(99)00512-3.

(31)   Zhang, W.; Wang, J.; Wang, J.; Zhu, L.; Lv, N.; Wang, R.; Ahmad, Z. New Insights

into Dose- and Time-Dependent Response of Five Typical PPCPs on Soil Microbial

Respiration. *Bull. Environ. Contam. Toxicol.* **2019**, *103* (1), 193–198.

https://doi.org/10.1007/s00128-019-02655-5.

(32)   Xiong, D.; Li, Y.; Xiong, Y.; Li, X.; Xiao, Y.; Qin, Z.; Xiao, Y. Influence of Boscalid on

the Activities of Soil Enzymes and Soil Respiration. *Eur. J. Soil Biol.* **2014**, *61*, 1–5.

https://doi.org/10.1016/j.ejsobi.2013.12.006.

(33) Wang, C.; Wang, F.; Zhang, Q.; Liang, W. Individual and Combined Effects of

Tebuconazole and Carbendazim on Soil Microbial Activity. *Eur. J. Soil Biol.* **2016**, *72*,

6–13. https://doi.org/10.1016/j.ejsobi.2015.12.005.

(34) Lynch, J. M.; Bragg, E. Microorganisms and Soil Aggregate Stabililty. In *Advances in*

soil science. Vol. 2; 1985; pp 133–171. https://doi.org/10.1007/978-1-4612-5088-3_3.

(35) Leifheit, E. F.; Verbruggen, E.; Rillig, M. C. Arbuscular Mycorrhizal Fungi Reduce

Decomposition of Woody Plant Litter While Increasing Soil Aggregation. *Soil Biol.*

Biochem. **2015**, *81*, 323–328. https://doi.org/10.1016/j.soilbio.2014.12.003.

(36) Piccolo, A.; Pietramellara, G.; Mbagwu, J. S. C. Use of Humic Substances as Soil

Conditioners to Increase Aggregate Stability. *Geoderma* **1997**, *75* (3–4), 267–277.

https://doi.org/10.1016/S0016-7061(96)00092-4.

(37) German, D. P.; Weintraub, M. N.; Grandy, A. S.; Lauber, C. L.; Rinkes, Z. L.; Allison,

S. D. Optimization of Hydrolytic and Oxidative Enzyme Methods for Ecosystem

Studies. *Soil Biol. Biochem.* **2011**, *43* (7), 1387–1397.

https://doi.org/10.1016/j.soilbio.2011.03.017.

(38) He, W.; Megharaj, M.; Naidu, R. Toxicity of Perfluorooctanoic Acid towards

Earthworm and Enzymatic Activities in Soil. *Environ. Monit. Assess.* **2016**, *188* (7).

https://doi.org/10.1007/s10661-016-5416-y.

(39) Zhao, T.; Lozano, Y. M.; Rillig, M. C. Microplastics Increase Soil PH and Decrease

Microbial Activities as a Function of Microplastic Shape, Polymer Type, and Exposure

Time. *Front. Environ. Sci.* **2021**, *9*, 1. https://doi.org/10.3389/fenvs.2021.675803.

(40) Dieleman, C. M.; Lindo, Z.; McLaughlin, J. W.; Craig, A. E.; Branfireun, B. A. Climate

Change Effects on Peatland Decomposition and Porewater Dissolved Organic Carbon

Biogeochemistry. *Biogeochemistry* **2016**, *128* (3), 385–396.

https://doi.org/10.1007/s10533-016-0214-8.

(41)   Lehmann, A.; Zheng, W.; Rillig, M. C. Soil Biota Contributions to Soil Aggregation. *Nat.*

396   *Ecol. Evol.* **2017**, *1* (12), 1828–1835. https://doi.org/10.1038/s41559-017-0344-y.

397   (42)   Kurz-Besson, C.; Coûteaux, M. M.; Thiéry, J. M.; Berg, B.; Remacle, J. A Comparison

398   of Litterbag and Direct Observation Methods of Scots Pine Needle Decomposition

399   Measurement. *Soil Biol. Biochem.* **2005**, *37* (12), 2315–2318.

400   https://doi.org/10.1016/J.SOILBIO.2005.03.022.

401   (43)   Xie, Y. A Meta-Analysis of Critique of Litterbag Method Used in Examining

402   Decomposition of Leaf Litters. *J. Soils Sediments 2020 204* **2020**, *20* (4), 1881–1886.

403   https://doi.org/10.1007/S11368-020-02572-9.

404   (44)   Jackson, C. R.; Tyler, H. L.; Millar, J. J. Determination of Microbial Extracellular

405   Enzyme Activity in Waters, Soils, and Sediments Using High Throughput Microplate

406   Assays. *J. Vis. Exp.* **2013**, No. 80, 50399. https://doi.org/10.3791/50399.