TITLE PAGE

Title

Functional and historical drivers of leaf shape evolution in Palms (Arecaceae)

Running title:

Drivers of leaf shape in evolution palms

Abstract

Aim: Leaves display a remarkable variety of shapes, each with potential ecological advantages

in specific climates. While relations between leaf shape and either climate or height has been

relatively well studied in eudicots, the macroecological drivers of shape remain poorly known in

monocots. Here, we investigated associations between climate and plant height with the

evolution of leaf shape in a clade with high species and morphological diversity.

Location: Global.

Time period: Cretaceous to contemporary.

Major taxa studied: Palms (Arecaceae).

Methods: We apply a Bayesian phylogenetic mixed model to test for associations between

climate and leaf shape (all entire-leaved, pinnate-dissected, palmate-dissected, and

costapalmate). We further reconstruct the ancestral leaf shape using multistate speciation and

extinction models and compare the frequency of shapes with global temperatures through time.

Results: We find that plant height associates with dissected leaves and that annual precipitation

associates with pinnate shapes. The ancestral leaf shape is unclear but early diversification was

dominated by pinnate-dissected palms, which has remained the most species-rich form of leaves

throughout palm history.

Main conclusions: Palms that are tall and live in humid regions are more likely to have pinnate

leaves. Through geological time scales, temperature did not play an obvious role in determining

leaf shapes. This study contributes to our understanding of how the diversity of leaf shapes is

linked to biological and climatic factors.

Keywords:

Arecaceae, annual precipitation, dissection, leaf shape, macroevolution, MuSSE, palms, plant

height.

1. Introduction

Leaves are the engines of most life on land. They capture atmospheric carbon dioxide and convert it into accessible nutrients that sustain ecosystem functioning and health. The basic functions they carry out – photosynthesis, transpiration, and respiration – depend on temperature and water availability. Because they carry out such critical functions, leaves are likely under strong natural selection, resulting in morphological adaptations to environmental conditions (Nicotra et al., 2011). Throughout angiosperm evolution, variation not only in length and width but also in blade dissection, has evolved numerous times independently (Nicotra et al., 2011). Nonetheless, the drivers of leaf shape evolution and its adaptive potential are poorly known, with research being limited to a handful of species or to model eudicots (Nicotra et al., 2011; Chitwood & Sinha, 2016; Leigh et al., 2017), hindering generalisations at a larger taxonomic scale (Conklin et al., 2019). The evolution of leaf shape results from trade-offs between physiological and allometric constraints, phylogenetic history, and natural selection (Givnish, 1987; Tsukaya, 2006; Nicotra et al., 2011). Therefore, understanding the extent to which leaf shapes have evolved in response to climate, allometry, or natural selection, and how they have changed over time and across geographies, may shed further light on the evolution of life forms and biological adaptations.

General expectations regarding leaf function can be drawn from examining their adaptations in different environments. Traits like leaf dissection, length and effective width (diameter of the largest circle drawn within a leaf margin; Leigh et al., 2017) vary across climates, which affects plant temperature regulation and interacts with water availability (Nicotra et al., 2011; Peppe et al., 2011). In environments with extremely hot temperatures, plants tend to have small or dissected leaves to avoid reaching damaging temperatures (Leigh et al., 2017; Wright et al., 2017). Deeply dissected leaves effectively function as a collection of small leaf units, with faster heat loss through convection than entire-leaved ones of the same area (Givnish, 1979; Gurevitch & Schuepp, 1990), and are less likely to reach damaging temperatures when exposed to extreme heat. Thus, shapes that reduce damage at high temperatures are expected to be beneficial in hot and dry environments (Nicotra et al., 2007; 2008; Leigh et al., 2017). Under drought conditions where water for transpiration is limited, stomata close and compounding high temperatures threaten leaf function. In such cases, smaller or deeply dissected shapes reduce

overheating and optimise leaf safety (Nardini & Luglio, 2014). Species with large and wide

leaves, optimised for high gas exchange, could benefit from dissection if water transportation to

blade areas farthest from the rachis is more effective (Givnish, 1979). In contrast, cool climatic

conditions may favour entire-leaved over dissected, where leaf heating is not extreme and

photosynthetic surfaces can be wide.

Leaf shape and size may also be optimised to maximise light capture. This is particularly

relevant for monopodial plants (non-branching plants with a single growth axis) or understory

species that depend on leaf size, angle, or number to increase the total photosynthetic area

(Givnish, 1988; Valladares et al., 2002; Renninger & Phillips, 2016). However, leaf size is

limited by allometry (the proportional change in size of different parts of an organism). Corner's

rule states that "the larger and sturdy an axis is, the larger and more complicated are its

appendages" (Corner, 1949; Tomlinson, 2006). Dissecting large leaves enables independent

movement of leaflets and reduces wind drag (Niklas, 1992; Vogel, 2009). Similarly, the "rapid

growth hypothesis" links dissection and allometry (Givnish 1984; Niinemets, 1998), and states

that dissection (or compound leaves) enables fast growth in high light by investing in longer

rachises, which is energetically cheaper than wood density and branching (Malhado et al., 2010).

To maximize light capture in the absence of branching, elongated petioles or rachises can be

used as alternative strategies (Givnish, 1988), facilitating optimal trade-offs between leaflets'

angle and self-shade (Valladares et al., 2002). When petiole lenghts are similar across species,

elongated rachises could be effective for outstretching leaves or leaflets to radii farther from the

plant stem. In addition, elongating light structures like the rachis could solve limitations imposed

by supporting heavier structures. In the absence of branching, alternative strategies for

maximising light capture could be elongated petioles or rachises (Givnish, 1988); particularly

rachises for outstretching leaves or leaflets to radii farther from the plant stem and facilitating

optimal trade-offs between leaflets' angle and self-shade (Valladares et al., 2002). Moreover,

long petioles for better light capture, elongating light rachises (and thus the leaf blade) can solve

limitations imposed by supporting long and heavy petioles.

Here, we use palms (Arecaceae) to understand the macroevolutionary drivers of leaf

shape variation. Palms are an ideal group to address this question given their wide distribution

and morphological diversity. They are tropical and subtropical, with 80% of the species

distributed within a 15-30°C mean annual temperature range (Dransfield et al., 2008), and

exhibit wide leaf shape variation (pinnate, palmate, and costapalmate, all of which can be entire-

leaved or dissected; Dransfield et al., 2008; Horn et al., 2009). Palms are primarily monopodial

with non-deciduous leaves (Tomlinson, 2006; Dransfield et al., 2008), allowing us to control for

the effect of branching strategies over leaf-variable relationships. Understanding the evolution of

leaf shape through past and current environmental conditions provides a context for predicting

plant responses to changing climates (Chitwood & Sinha., 2016).

In this study, we aim at disentangling the contributions of climatic (extrinsic) and

allometric (intrinsic) factors on palm leaf shape by testing four hypotheses: **H1:** Climate, and in

particular temperature, contributes more to leaf shape than allometry. Dissected species (pinnate-

and palmate-leaved) should be found at higher temperatures (or lower precipitation and higher

aridity) than entire-leaved species regardless of height. **H2:** Allometry, likely plant height,

contributes more to leaf shape. We test H1–H2 by comparing i) extant dissected and non-

dissected shaped species and ii) all shapes in a pairwise manner. **H3:** Either plant's height in

relation to forest canopy height or elongated rachises contribute to leaf shape by trading light

capture maximisation and leaf support, particularly when comparing pinnate and palmate

species. Longer rachises (in pinnate species) translate into long blades and more surface for light

capture in cases for which petioles and stems are short. We test H3 by comparing all shapes in a

pairwise manner. **H4:** If climate is a strong contributor to leaf shape evolution through time, this

should be reflected in a temporal congruence between leaf shape shifts and major global climatic

changes since the Late Cretaceous, when palms are estimated to have originated (Couvreur et al.,

2011; Baker & Couvreur, 2013). We test these hypotheses by estimating variable effects and

estimating ancestral character states.

2. Methods

We conducted Generalised Linear Mixed Models (GLMMs) and ancestral trait estimation

analyses at the species level using a time-calibrated Maximum Clade Credibility tree generated

from the tree distribution in Faurby et al. (2016) and updated by Hill et al. (2021). The

phylogeny included 2550 species for which we annotated leaf shapes, recovered geographic

records from the Global Biodiversity Information Facility (GBIF), and estimated species

medians for the climatic and allometric variables. We standardised the taxonomy across all data

sources using Kew's World Checklist of Selected Plant Families (WCSP) for Arecaceae

(Govaerts et al., 2020) and removed records that could not be unambiguously assigned to

accepted species.

Leaf shape in palms

Leaf shape variation in palms can be described by three features: size, plication and

dissection. Since plication is phylogenetically conserved, we focused on dissection and shape,

but see supplementary material for a brief description of size and plication. Leaves are either

dissected or entire-leaved. Dissected leaves can be pinnate, palmate, or costapalmate depending

on rachis length and the presence of a costa (here treated as the equivalent of rachis in pinnate

leaves but it is an extension of the leaf axis; Dransfield et al., 2008). Polymorphism, where

intraspecific variation in leaf shape occurs, only involves either pinnate-entire or pinnate-

dissected shapes. Genomic analyses (e.g. Loiseau et al., 2019) confirm most intraspecific

variation as true polymorphisms within populations, rather than separate taxa grouped under one

species name.

We classified species by dissection (dissected vs. entire) and shape (pinnate, palmate, and

costapalmate) based on Genera Palmarum II (Dransfield et al., 2008), additional information on

the PalmWeb (https://palmweb.org/, last consulted January 2023), and herbarium specimens

(access through GBIF). We merged the costapalmate and palmate shapes because the only

difference between them is the presence of a costa. We removed the bipinnate category from the

GLMM analyses but merged them to pinnate for ancestral state estimation because bipinnate

includes only 14 out of 2550 species. Of the species analysed, the majority are pinnate (66.18 %;

Table S1, Supplementary Material), followed by palmate+costapalmate (21.26 %) and entire

(5.12%). Only 75 (6.76 %) species are polymorphic and while they were included in the

ancestral state estimation, they were excluded from the GLMMs because they cannot be assigned

to a unique shape category (Table S2).

Palm allometry data

We used Palm_Traits v.1 (Kissling et al., 2019) to calculate plant height by adding the

variables 'MaxStemHeight_m', 'Max_Petiole_length_m', and 'Max_Blade_Length_m', with

'MaxStemHeight_m' set to zero for acaulescent species. We estimated an index that measures

plant height controlled by canopy height (height over canopy) as a proxy to wind-drag exposure

and the Understorey/Canopy variable in Kissling et al. (2019). The index differentiates between

e.g. two tall species, one occurring in a high canopy and another in a low canopy forest. Values

of height over canopy are higher than 2.70e+10 for tall species in low canopies, close to one for

species with the same height as the canopy, and smaller than 2.65e-08 for short species under

high canopies (based on the 0.25 and 0.75 quantiles). To calculate it, we divided the palm height

species value by the height extracted at the coordinate point from the Global Vegetation Height

layer (Simmard, 2011). Calculations and code are available at

doi.org/10.6084/m9.figshare.21230453. We annotated 61-100% of the species in the phylogeny.

Species without annotations had no data in Palm_Traits v.1, were climbing, or had both climbing

and no-climbing habits (479 species, Table S3); climbing species were removed from the

GLMM analyses because their life strategies differ from other palms and their stem height is not

comparable.

Climatic data

We downloaded 994,084 raw occurrences from GBIF (last consulted in April 2022) and

excluded fossils and records without coordinates. We kept observations, living and preserved

specimens, and material samples. We used the R package (R Core Team, 2018)

'CoordinateCleaner' v2.0-20 (Zizka et al., 2019) to remove duplicate coordinates per species and

records nearby science institutions, bodies of water, and city/country centroids, using a buffer of

5000 m and 10000 m for centroids and cites respectively. We used a Python script

(doi.org/10.6084/m9.figshare.21230453) and the World Geographical Scheme for Recording

Plant Distributions maps (TDWG; Brummitt, 2001) to remove records falling outside their native

botanical countries according to the WCSP. We obtained 124,703 clean records representing

61.56% of the 2550 species in the phylogeny (Table S4), which we used to extract all Chelsa

v2.1 bioclim variables (Karger et al., 2017; Karger at al., 2018) with a python script

(doi.org/10.6084/m9.figshare.21230453). Finally, we estimated the species medians for every

variable from which we could extract the information (Table S4).

We determined correlations between variables and chose those with a Spearman's

coefficient $-0.7 < \square < 0.7$ (Fig. S1). We additionally estimated the variance inflation factor for

all variables using python's 'statsmodels' v0.13.5 (Seabold and Perktold, 2010) and kept those

with values below two. We kept five climatic and three allometric variables: mean annual

temperature (°C*100), temperature seasonality (standard deviation °C*100), annual precipitation

(kg m-2); precipitation seasonality (standard deviation kg m-2), aridity index (mean annual

precipitation/mean annual potential evapotranspiration; kg m-2/time unit), plant height (m),

rachis length (m), and height over canopy. Temperature seasonality and mean annual

precipitation, and plant height and height over canopy were correlated (Spearman's correlation

coefficient = -0.8 and 0.73 respectively; Fig. S1); however, we assessed them in separate models

to determine if the effect of e.g. temperature is larger than e.g. precipitation on leaf shape. All

variables used were $\log_{10}$-transformed and standardised to have zero mean and a standard

deviation of one (Table S4).

Generalised linear mixed models

We fit a series of GLMMs for the first three hypotheses using logistic regressions and

pairwise comparisons; entire (0) vs. dissected (1), pinnate (0) vs. palmate (1), entire (0) vs.

palmate (1), and entire (0) vs. pinnate (1); these were implemented in the R package

'MCMCglmmRAM' v2.24 (Hadfield, 2015). The GLMMs assess shape rather than shape

evolution, thus, we treat pinnate-entire and palmate-entire as entire-leaved, pinnate-dissected as

pinnate, and palmate-dissected as palmate. For computational constraints, we ran every model on

42 (Adams, 1979) phylogenies randomly selected from the distribution, each as a random effect.

For every model, we ran four independent chains of 10,000,000 iterations, a thinning of 5,000,

and 8000 burn-in. Analyses were run on a HPE Cray OS computer through the Swedish National

Infrastructure for Computing. We estimated effective sampling sizes (ESS) and evaluated chain

convergence with a python script (doi.org/10.6084/m9.figshare.21230453). Chains converged

and ESS scored higher than 200. We evaluated the significance of the effects for every predictor

based on whether the 2.5 and 97.5% quartiles of the estimated density overlapped zero.

Ancestral state estimation

To reconstruct the evolution of leaf shapes we used the Multi-State Speciation and

Extinction (MuSSE) model from the R package 'diversitree' v0.9-16 (FitzJohn, 2012). We

adjusted the leaf shape dataset to avoid over parametrization of our models: 1) excluded plication

due to high phylogenetic clustering (Fig. 1; but see Couturier et al., 2011); 2) used a

"polymorphic" category for species exhibiting more than one shape; 3) excluded all species with

missing data for leaf shape. The final dataset included 2543 species. Ancestral state estimation

concerns shape and dissection evolution, thus we designed three different model structures – two

binary state structure excluding polymorphic, pinnate vs. palmate, entire vs. dissected, and a

five-state structure (pinnate-entire, pinnate-dissected, palmate-entire, palmate-dissected and

polymorphic).

For the two binary structures we fitted all possible parameter constraints using maximum

likelihood (ML) parameter estimations and compared them from AIC scores. For the five-state

structure, we refined the basic model structure by assuming constraints on transition rates. We

did not allow simultaneous change of shape and dissection (*e.g.*, from palmate-entire to pinnate-

dissected). As polymorphism only includes pinnate leaves, we assumed direct transitions

between palmate and polymorphic states to be impossible and constrained the corresponding

transition parameters to zero. Given the large number of parameters remaining (22), we selected

the best-fitting model using a backward model selection procedure using ML estimation on the

maximum clade credibility tree only. Once we obtained the best-fitting model for the two binary

and the five-state structures, we used it to perform an MCMC analysis to estimate the posterior

distribution of parameter estimates. We ran the MCMC for 3,000,000 generations, sampling

every 3,000 generations and a burnin of 10%. We used the ML parameter estimation as starting

points for the MCMC. In addition, we performed ML estimations of the best-fitting model (1) for

100 trees randomly sampled from the posterior distribution to account for phylogenetic

uncertainty and (2) for a range of starting parameters. More details about model selection and

parameter constraints are available in the Supplementary Material.

For all model structures, we used the best-fitting MuSSE model to estimate ancestral

states, and the relative frequency of leaf shape through time by sampling the relative

probabilities of shape estimated for each node 100 times. For each iteration, at each branch

where a state transition occurred, we sampled a random timing from a uniform distribution for

the event along that branch. Finally, for each iteration, we counted the number of transitions

within a 5 million-year sliding window and calculated the rate of transitions through time by

dividing the number of transitions by the sum of branch lengths within each time interval. We

repeated this procedure for 100 trees randomly sampled from the posterior distribution. Finally,

we compared the relative shape frequency through time with the global temperature change scale

computed for an ice-free ocean (Zachos et al. 2001; Condamine et al., 2020).

3. **Results**

Leaf shape and variable distributions

Pinnate and palmate species tend to be larger than entire-leaved species with median

plant heights of 7.39 m and 7 m, respectively, while entire-leaved species had a median height of

3.8 m (Table S5), leaf and petiole lengths included in all cases. Dissected shapes together had a

median height twice that of non-dissected species. Pinnate-leaved species had a median rachis

length of 2.9 m and a median height over canopy of 4e+10, while palmate-leaved species had a

median rachis length of 2.4 m and a median height over canopy of 7e+10. Entire-leaved species

had a median height over canopy of 1.3e+10. Pinnate and palmate species tend to be equal or

taller than the canopy, whereas entire-leaved species were rarely tall enough to approach the

canopy and do not occur in open habitats. alms are widely distributed across climatic gradients

except for habitats where soil temperatures fall below -2□ for a long period of time. Pinnate and

palmate species are distributed farther from the equator with wider climatic seasonality, while

the distribution of entire-leaved species is concentrated in tropical areas near the equator where

annual temperature and precipitation are more constant. Pinnate and palmate species have wider

median distributions for most climatic variables than entire-leaved species, except for narrower

annual mean temperature and temperature seasonality ranges in palmates. The median of the

distribution for mean annual temperatures is highest for palmate species, then entire-leaved

species, and lowest for pinnate species (Table S5, Supplementary Material).

Generalised Linear Mixed Models

We found that plant height was positively associated with dissected leaves, particularly

pinnate shapes. The association is significant when comparing entire-leaved vs. pinnate-leaved

species (P; proportion of samples above/below zero=0.99; Fig. 2, Table S8) and entire-leaved vs.

dissected species (P=0.99). The pattern was consistent across models regardless of which

climatic variables were included. For the model including annual precipitation instead of annual

temperature seasonality, annual precipitation showed a positive and almost significant

association with pinnate species when compared to entire-leaved species (P=0.952), but not

when comparing dissected and entire-leaved shapes (P=0.89). Nevertheless, the effect of plant

height is an order of magnitude larger than the effect of precipitation (1.58E+05 and 6.68E+04

respectively, Table S8, Supplementary Material). For models comparing pinnate and entire-

leaved shapes in which height over canopy was included instead of plant height (these variables

are correlated and were not included in the same model), only height over canopy was almost

significantly and positively associated with pinnate shapes (P=0.953). Height over canopy had a

similar effect as plant height on leaf shape for pinnnate vs. entire-leaved models (1.02E+05 and

1.58E+05). However, the significance of height over canopy' effect was lost if the model did not

account for annual precipitation. No climatic variables were consistently and significantly

associated with leaf shape (0.35<P< 0.94) and all variables were not significant in models

involving palmate shapes (Fig. 2, Table S8).

When disentangling the effects of rachis length, height over canopy, and climate, we

found a significant positive association between rachis length and pinnate shapes when compared

to entire-leaved species; plants with longer rachises are more likely to be pinnate. In addition to

rachis length, annual precipitation had a significantly positive association with pinnate species

and height over canopy was almost significant (P=0.99 and P=0.968, respectively). Here, the

effect of rachis length was one order of magnitude larger than precipitation and height over

canopy, which had similar effects. The models resulted in no association between rachis length

and shape when comparing pinnate and palmate species (0.34< P<0.5; Fig. 2, Table S8).

Ancestral state estimation

The best-fitting model for dissection vs. entire-leaved state estimation had six parameters

(Fig. S3). The speciation rate (lineage/unit of time, $l^{-1}my^{-1}$) for entire-leaved lineages was more

than twice higher as that of dissected lineages and the extinction rate was simultaneously much

lower for entire-leaved lineages, but transition rates were highly asymmetrical favouring

transitions from entire-leaved to dissected than the reverse (Fig. S8). For pinnate vs. palmate model structure, we found no difference in speciation or extinction rates between shapes. Transitions from pinnate to palmate leaves were three times higher than the reverse transition, but both were very low (Fig. S9). The best-fitting model for the five-state model had 12 parameters (Fig. S5). Most notably, the speciation rate of polymorphic species was more than twice higher as that of the others. The extinction rate for the polymorphic and palmate-dissected lineages was much lower than that of pinnate-dissected and palmate-entire lineages. The resulting net diversification was highest for polymorphic lineages, followed by pinnate-entire leaves. Both palmate-entire and pinnate-dissected net diversification rates were close to zero. Transitions were highest from polymorphic leaves to pinnate-dissected, twice higher than transitions from pinnate-entire to polymorphic leaves and near 10 times higher than transitions from polymorphic to pinnate-entire leaves. All other types of transition rates were very low (Fig. S10). For all three model structures, we obtained identical parameter estimations from varying starting parameters, a posterior distribution of trees or MCMC analyses.

Across a posterior distribution of 100 trees, the two-state structures estimated the root to be pinnate and entire-leaved (Fig. S6-7). According to the five-state structure, the shape at the root was most likely polymorphic or pinnate-entire (Fig. 3). However, we found the results to poorly represent the pattern of early diversification of shapes. The relative frequency of shapes through time, calculated from the ancestral state estimations, shows instead that pinnate-dissected leaves dominated through the history of palms, representing between 50 to 80% of all lineages. Despite appearing as likely leaf shape at the root, entire-leaved lineages diversified only during the second half of the Cenozoic. According to our estimations, palmate-dissected palms appeared about 75-85 My ago and represented about 20% of palm lineages throughout history. Polymorphic lineages were poorly represented during the middle Cenozoic, despite being estimated as a probable shape for the root of the tree and increasing in frequency during the last 30 My (Fig. 4).

4. Discussion

In this study, we aimed at disentangling the contributions of plant allometry and climate on leaf shape by testing whether **H1)** temperature contributes more than allometry to leaf shape;

or **H2)** allometry, as plant height, contributes to shape, or **H3)** rachis length or height over

canopy contribute to shape; and whether **H4)** there is temporal congruence between shape

frequency and major global temperature changes. This suggest that taller plants are more likely

to have dissected leaves and, regarding climate, only annual precipitation contributes and makes

species more likely to have pinnate leaves (Fig. 2). We also found that longer rachises make

pinnate leaves more likely. In congruence with our results, we observed a lack of correspondence

between leaf shape frequency and temperature fluctuations through time. Our models of

ancestral trait estimation revealed that the ancestor of all palms likely had entire-leaved and

pinnate or polymorphic leaves (Fig. 3-4c).

Leaf dissection and climate

Contrary to H1, neither mean annual temperature nor temperature seasonality had an

effect on leaf shape (Fig. 2, Table S8). We expected that plants exposed to extremely high

temperatures would have dissected leaves as a strategy to avoid overheating their leaves faster

(Nicotra et al., 2007; 2008; Leigh et al., 2017). Similarly, we expected that a high aridity index –

a proxy for a lack of water in the environment– would favour pinnate leaves over other shapes.

Our results showed no significant associations between climate and leaf shape for most of the

models and only the effect of annual precipitation was significant when comparing pinnate and

entire-leaved species, thus rejecting **H1**. Species that experience high annual precipitation are

more likely to be pinnate than entire-leaved when plant height is also present in the model. Since

plant height and annual precipitation are not correlated, our reuslts suggests that even though the

effect of plant height is greater, annual precipitation and plant height both seem to explain

independent variation.

Moreover, the association between climate and palmate shapes is not significant (Fig. 2);

three non-mutually exclusive hypotheses could explain the general lack of associations, of which

the first two are more likely. First, the discrete shape categories in our models might not fully

capture interspecific variation in dissection depth in palmate species and could be insufficient for

unveiling the climate-palmate associations. Second, leaf traits unrelated to shape could be key

adaptations to different environments (Horn et al., 2009). For example, the loss of non-lignified

fibre bundles in the leaf mesophyll mostly observed in coryphoid palms or the presence of

bridge-like veins connecting the adaxial and abaxial layers of the leaf are traits thought to be
advantageous in dry environments (Kenzo et al., 2007; Horn et al., 2009). Third, in the case of
palmate and costapalmate species, the correlation between climate and shape is difficult to
disentangle from a phylogenetic effect due to the strong geographic structuring of related
lineages, particularly in the Coryphoideae subfamily to which the majority of palmate and
costapalmate species belong.
378
Leaf shape and plant allometry
We expected plant height to influence leaf shape via allometry so that large plants would
have large and more likely dissected leaves to avoid mechanical damage (Corner, 1949;
Chazdon, 1991). We found that plant height has a positive effect on dissection and taller plants
are more likely to have pinnate leaves (Fig. 2, Table S8). Our results were consistent when
comparing dissected –pinnate and palmate together– and pinnate species against entire-leaved
species, supporting **H2**. Only when comparing palmate- vs. entire-leaved species, all effect
estimates were not significant (Fig. 2). This suggests that the association between plant height
and dissection is primarily driven by the number of pinnate species (66.18% vs. 21.26% species
with palmate leaves) and that other factors are involved when palmate species are considered (as
discussed in the section above).
390
Following the "rapid growth hypothesis" (Givnish, 1984; Niinemets, 1998), we expected
plant height to relate to dissection. Malhado et al. (2010) tested this hypothesis in Amazonian
species (except palms and polymorphic eudicots), and found an association between dissection
and both low wood density and rapid diameter growth. They argue that dissection is adaptive
under favourable light conditions because producing compound or dissected leaves is
physiologically less expensive since enlongater rachises are cheaper than supporting structures
(petioles in the case of palms) and thus promotes rapid vertical growth (Malhado et al., 2010).
We found a positive association between height over canopy and pinnate shapes when these are
compared to entire-leaved shapes (Fig. 2, Table S8), although marginally significant. The
significance of this association, however, should be interpreted carefully as height over canopy
values are derived from, and correlated with, plant height. The lack of association between
height over canopy when comparing pinnate and palmate-leaved species is explained by the

large proportion of species occupying similar spaces in the canopy where overall light
availability conditions are similar. Malhado et al. (2010) excluded palms from their analyses and
testing whether rapid growth or light capture optimisation better explains dissection is not
possible with our data; a deep look at species-level measurements of growth rates and light
capture efficiency could shed a light on this question. Nonetheless, our results suggest that
allometry is more important than light capture in palms and further exploration of this
association is worthwhile in future studies.

Finally, we expected light capture optimisation to drive rachis length and contribute to
shape when comparing pinnate vs. palmate species. Pinnate leaves differ from palmate leaves in
that pinnae attach to a long axis and not a centroid. On one hand, the elongated rachis could be a
strategy for maximising light capture for the pinnate-leaved species living under the canopy.
Pinnae distributed along a longer axis can reach farther from the plant's standing point and
reduce self-shading in closed habitats. On the other hand, most species with palmate and
costapalmate leaves in which the leaflets are arranged radially, are as tall or taller than the
cannopy they inhabit (Fig. S2) where light availability is high. However, we found no
association between rachis length and shape for pinnate vs. palmate species, thus rejecting **H3**
(Fig. 2, Table S8). The lack of association between height over canopy and shape when
comparing pinnate vs. palmate-leaved species is consistent with these results, meaning that
neither the plant's position with respect to the canopy nor the rachis length explain leaf shape.
The association between rachis length and shape is only significant when comparing pinnate- vs.
entire-leaved species, which can be explained by Corner's rule: large plants have large leaves
which necessarily have longer rachises.

Leaf shape evolution
We found that dissected lineages appeared frequently during the history of palms and
from entire-leaved lineages (Fig. 3, S6-7). Additionally, palmate (palmate+costapalmate)
lineages appeared at least twice from pinnate lineages, a result consistent with Horn et al. (2009).
Using 178 taxa with pinnate or palmate species only, their study concluded that shape is
homoplasious and changes between states are frequent. Overall, leaf shape is particularly labile

among pinnate-leaved palms and evolutionary changes are frequent between pinnate-entire and
pinnate-dissected, involving polymorphism as well.

Based on our ancestral state estimation, the origin of palm leaves is not entirely resolved.
We found clear support for an ancestral pinnate leaf, but conflicting results for dissection. On the
one hand, entire-leaved states are repeatedly inferred at the root across our models. However,
pinnate-entire leaves are almost entirely absent from early diversification of palms, instead
dominated by pinnate-dissected leaves (Fig. 4a,b). Palmate and costapalmate leaf shapes
appeared between 75-85 Mya, which corresponds to the stem of Coryphoideae. The appearance
of palmate shapes roughly coincides with the age of the costapalmate fossil *Sabalites*
carolinensis (Santonian, 86.3-83.6 Mya; Berry, 1914), commonly cited as the earliest palm leaf
fossil. However, uncertainty as to the stratigraphic assessment of the formation from which *S.*
carolinensis was reported calls for the reassessment of the fossil's age to be Campanian (83.6-
72.1 Mya; Greenwood et al., 2022). If that is the case, the oldest costapalmate (here merged with
palmate) fossil would postdate the older Coniacian–early Campanian "pinnate-entire"
Phoenicites imperialis (Dawson) emend. Greenwood et Conran comb. nov. (89.8 to
approximately 80 Mya; Greenwood et al., 2022) and be contemporaneous to the mid-Campanian
"undivided" *Plicatophyllum* (81-76 Mya; Crabtree 1987; Greenwood et al., 2022). Macrofossils
remain rare and provide at best a minimum age for any taxon or morphological character; the
absence of older fossils is likely explained by the incomplete nature of the fossil record and its
identification. Nevertheless, results on root state estimations should be interpreted with caution
since validating them would require a comprehensive taxonomic sampling beyond Arecaceae
and our analyses do not consider potential correlations between leaf shape and other
morphological characters.

One interesting outcome of our five-state model is the role played by polymorphic
lineages. Polymorphism – pinnate-dissected and pinnate-entire – appears as a transitional state
between these shapes forming an evolutionary bridge between lineages with non-polymorphic
leaf shapes (Fig. 3). Maximum Likelihood estimates of transition parameters also suggest a
strong directionality; the highest transition rates being from polymorphic toward pinnate shapes
and the second highest was transitions from entire-leaved shapes towards polymorphism (Fig. 4a
and S3). As a result, polymorphic lineages are maintained in our ancestral state estimation

throughout time despite that only 3% of extant species are polymorphic. Dissected and entire-

leaved of pinnate and palmate shapes do not have significantly different speciation rates but the

speciation rate for polymorphic lineages is more than twice higher, which also translated into a

higher net diversification rate. Thus, according to our best model, the high number of extant

pinnate-dissected lineages (1363 species) does not result from its high diversification rates. It

could instead be a result of either a high speciation rate of polymorphic lineages with frequent

independent transitions to pinnate-dissected leaves or an early success during diversification that

facilitated the accumulation of pinnate-dissected lineages throughout the Cenozoic. In either

case, the prevalence of pinnate-dissected leaves is an ancient and deeply rooted pattern in the

group's history.

Leaf shape through time

We found no clear relationship between temperature variations and changes in the

relative frequency of leaf shapes, thus we reject **H4** (Fig. 4c). The late Cretaceous and early

Eocene were periods of high global temperatures, with ever-wet tropical areas distributed

towards high latitudes. Palms thrived during these periods forming the Palmae Province, and

becoming ecologically dominant in South America, Africa and India (Pan et al., 2006). During

the Eocene, global temperatures decreased at a relatively fast pace (Zachos et al., 2001) and kept

declining towards the present, except during the Mid-Miocene climatic optimum. But there is

also evidence of aridification across continents over the past 40-50 My alongside temperature

variation, leading to the contraction of tropical-like forests towards the equatorial latitudes. The

pollen record has revealed patterns of palm extinctions including groups such as Mauritiinae and

Eugeissoneae, typical rainforest groups and pinnate-dissected (Bacon et al. 2022, Lim et al.

2021). From our results, this period corresponds to the diversification of entire-leaved lineages,

increasing in relative frequency at the expense of pinnate-dissected palms mainly (Figure 4a, b).

Global temperature change did not translate into a clear continuous shift in relative leaf shape

frequency, but increasing aridity since the Eocene may have promoted the diversification of

entire-leaved palms, considering the positive association between precipitation and pinnate-

dissected shapes we found in modern palms.

Conclusion

We explore the drivers of leaf shape evolution in palms by disentangling the associations between shape, climate, and allometric variables related to plant height, and by reconstructing the evolution of shape throughout their evolutionary history. We highlight the importance of considering biotic (intrinsic) and abiotic (extrinsic) factors when studying the evolution of traits. Regarding palms, tall plants are more likely to be dissected and this tendency is mostly driven by pinnate species but not palmate species. High annual precipitation increases the likelihood of having pinnate leaves when controlling for plant height and temperature was not significantly associated with shape. Palms are important representative taxa of tropical forests with more than 2,500 species distributed globally. Exploring how their leaf shape diversity emerged contributes to our understanding of shape and its adaptive potential to future climate conditions.

REFERENCES

Adams, D. (1979). The Hitchhiker's Guide to the Galaxy. London, UK: Pan Books.

Bacon, C.D., Silvestro, D., Hoorn, C., Bogotá-Ángel, G., Antonelli, A. & Chazot, N. (2022). The origin of modern patterns of continental diversity in Mauritiinae palms: the Neotropical museum and the Afrotropical graveyard. *Biology Letters* **18**(11): 20220214.

Baker, W.J. & Couvreur, T.L. (2013). Global biogeography and diversification of palms sheds light on the evolution of tropical lineages. I. Historical biogeography. *Journal of Biogeography* **40**(2): 274-285.

Berry, E. W. (1914). The upper Cretaceous and Eocene floras of South Carolina and Georgia (No. 84). US: US Government Printing Office.

Brummitt, R. K. (2001). World Geographical Scheme for Recording Plant Distributions, Edition 2. *Biodiversity Information Standards (TDWG)*, http://www.tdwg.org/standards/109

Chazdon, R. L. (1991). Plant size and form in the understory palm genus *Geonoma*: are species variations on a theme? *American Journal of Botany* **78**(5): 680-694.

Chitwood, D. H., & Sinha, N. R. (2016). Evolutionary and environmental forces sculpting leaf development. *Current Biology* **26**(7): R297-R306.

Crabtree, D.R., (1987). The early Campanian flora of the Two Medicine Formation, northcentral Montana. University of Montana. USA.

Condamine, F. L., Silvestro, D., Koppelhus, E. B., & Antonelli, A. (2020). The rise of angiosperms pushed conifers to decline during global cooling. *Proceedings of the National Academy of Sciences* **117**(46): 28867-28875.

Conklin, P. A., Strable, J., Li, S., & Scanlon, M. J. (2019). On the mechanisms of development in monocot and eudicot leaves. *New Phytologist* **221**(2): 706-724.

Corner, E. J. H. (1949). The durian theory or the origin of the modern tree. *Annals of Botany* **13**(52): 367-414.

Couturier, E., Du Pont, S. C., & Douady, S. (2011). The filling law: a general framework for leaf folding and its consequences on leaf shape diversity. *Journal of Theoretical Biology* **289:** 47-64.

Couvreur, T.L., Forest, F. & Baker, W.J. (2011). Origin and global diversification patterns of
tropical rain forests: inferences from a complete genus-level phylogeny of palms. *BMC*
biology, **9***:* 1-12.
Dransfield, J., Uhl, N.W., Asmussen, C.B., Baker, W.J., Harley, M.M. & Lewis, C.E. (2008)
Genera Palmarum - the evolution and classification of palms. Royal Botanic Gardens,
Kew, Richmond, 732 pp.
Faurby, S., Eiserhardt, W. L., Baker, W. J., & Svenning, J. C. (2016). An all-evidence
species-level supertree for the palms (Arecaceae). *Molecular Phylogenetics and*
Evolution **100**: 57-69.
FitzJohn, R. G. (2012). Diversitree: comparative phylogenetic analyses of diversification in R.
Methods in Ecology and Evolution **3**(6): 1084-1092.
the Global Biodiversity Information Facility. GBIF Occurrence Download
https://doi.org/10.15468/dl.w8djyx retrieved on 26 April 2022.
Givnish, T. J. (1979). On the adaptive significance of leaf form. Topics. In: O.T. Solbrig, G.B.
Jain, G.B. Johnson and P.H. Raven eds. *Plant Population Biology.* New York, US:
Columbia University Press, 375-407.
Givnish, T. J. (1984). Leaf and canopy adaptations in tropical forests. In: E. Medina, H.A.
Mooney and C. Vásquez-Yánes eds. *Physiological ecology of plants in the wet tropics.*
Tasks for vegetation Science, vol 12. Dordrecht, Netherlands: Springer, 51–84.
Givnish, T. J. (1987). Comparative studies of leaf form: assessing the relative roles of selective
pressures and phylogenetic constraints. *New phytologist* **106**: 131-160.
Givnish, T.J., (1988). Adaptation to sun and shade: a whole-plant perspective. *Functional Plant*
Biology 15(2): 63-92.
Greenwood, D.R., Conran, J.G. & West, C.K., (2022). Palm fronds from western Canada are
the northernmost palms from the Late Cretaceous of North America and may include the
oldest Arecaceae. *Review of Palaeobotany and Palynology* **301**: 104641.
Govaerts, R., Dransfield, J., Zona, S., Hodel, D. R., & Henderson, A. (2020). *World Checklist*
of Arecaceae. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet;
http://wcsp.science.kew.org/ Retrieved 25 February 2020
Gurevitch, J., & Schuepp, P. H. (1990). Boundary layer properties of highly dissected leaves:
an investigation using an electrochemical fluid tunnel. *Plant, Cell & Environment* **13**(8):
783-792.
Hadfield, J. D. (2015). Increasing the efficiency of MCMC for hierarchical phylogenetic models
of categorical traits using reduced mixed models. *Methods in Ecology and Evolution*
6(6): 706-714.
Hill, A., Torres Jimenez, M. F. , Chazot, N., Cássia-Silva, C., Faurby, S., & Bacon, C. D.
(2021). Fruit colour and range size interact to influence diversification. *bioRxiv*.
10.1101/2021.10.26.465838.
Horn, J. W., Fisher, J. B., Tomlinson, P. B., Lewis, C. E., & Laubengayer, K. (2009).
Evolution of lamina anatomy in the palm family (Arecaceae). *American Journal of*
Botany **96**(8): 1462-1486.
Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W.,
Zimmermann, N.E., Linder, P., & Kessler, M. (2017). Climatologies at high resolution
for the Earth land surface areas. *Scientific Data* 4-170122.
Karger D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W.,
Zimmermann, N.E, Linder, H.P., & Kessler, M. (2018). Data from: Climatologies at

high resolution for the earth's land surface areas. EnviDat.
https://doi.org/10.16904/envidat.228.v2.1

Kenzo, T., Ichie, T., Watanabe, Y., & Hiromi, T. (2007). Ecological distribution of homobaric
and heterobaric leaves in tree species of Malaysian lowland tropical rainforest. *American
Journal of Botany* **94**(5): 764-775.

**Kissling, W. D., Balslev, H., Baker, W. J., Dransfield, J., Göldel, B., Lim, J. Y., Onstein, R.
E., & Svenning, J. C. (2019).** PalmTraits 1.0, a species-level functional trait database of
palms worldwide. *Scientific Data* **6**(1): 1-13.

**Kissling, W. D., Walls, R., Bowser, A., Jones, M. O., Kattge, J., Agosti, D., Amengual, J.,
Basset, A., Van Bodegom, P. M., Cornelissen, J. H., & Denny, E. G. (2018).** Towards
global data products of Essential Biodiversity Variables on species traits. *Nature Ecology
& Evolution* **2**(10): 1531-1540.

Leigh, A., Sevanto, S., Close, J. D., & Nicotra, A. B. (2017). The influence of leaf size and
shape on leaf thermal dynamics: does theory hold up under natural conditions? *Plant,
Cell & Environment* **40**(2): 237-248.

**Lim, J.Y., Huang, H., Farnsworth, A., Lunt, D.J., Baker, W.J., Morley, R.J., Kissling, W.D.
& Hoorn, C. (2022).** The Cenozoic history of palms: Global diversification,
biogeography and the decline of megathermal forests. *Global Ecology and Biogeography*
31(3), pp.425-439.

**Loiseau, O., Olivares, I., Paris, M., de La Harpe, M., Weigand, A., Koubínová, D., Rolland,
J., Bacon, C. D., Balslev, H., Borchsenius, F., & Cano, A. (2019).** Targeted capture of
hundreds of nuclear genes unravels phylogenetic relationships of the diverse Neotropical
palm tribe Geonomateae. *Frontiers in Plant Science* **10**:864.

**Malhado, A. C., Whittaker, R. J., Malhi, Y., Ladle, R. J., tTer Steege, H., Phillips, O.,
Aragão, L. E. O. C., Baker, T. R., Arroyo, L., Almeida, S., & Higuchi, N. (2010).** Are
compound leaves an adaptation to seasonal drought or to rapid growth? Evidence from
the Amazon rain forest. *Global Ecology and Biogeography* **19**(6): 852-862.

Nardini, A., & Luglio, J. (2014). Leaf hydraulic capacity and drought vulnerability: possible
trade☐offs and correlations with climate across three major biomes. *Functional Ecology*
28(4): 810-818.

Nicotra, A. B., Cosgrove, M. J., Cowling, A., Schlichting, C. D., & Jones, C. S. (2008). Leaf
shape linked to photosynthetic rates and temperature optima in South African
Pelargonium species. *Oecologia* **154**(4): 625-635.

Nicotra, A. B., Hermes, J. P., Jones, C. S., & Schlichting, C. D. (2007). Geographic variation
and plasticity to water and nutrients in *Pelargonium australe*. *New Phytologist* **176**(1):
136-149.

**Nicotra, A. B., Leigh, A., Boyce, C. K., Jones, C. S., Niklas, K. J., Royer, D. L., & Tsukaya,
H. (2011).** The evolution and functional significance of leaf shape in the angiosperms.
Functional Plant Biology **38**(7): 535-552.

Niinemets, Ü. (1998). Are compound-leaved woody species inherently shade-intolerant? An
analysis of species ecological requirements and foliar support costs. *Plant Ecology*
134(1): 1-11.

Niklas, K. J. (1992). Petiole mechanics, light interception by lamina, and "economy in design".
Oecologia **90**(4): 518-526.

Pan, A. D., Jacobs, B. F., Dransfield, J., & Baker, W. J. (2006). The fossil history of palms
(Arecaceae) in Africa and new records from the Late Oligocene (28–27 Mya) of north-
western Ethiopia. *Botanical Journal of the Linnean Society* **151**(1): 69-81.
Peppe, D. J., Royer, D. L., Cariglino, B., Oliver, S. Y., Newman, S., Leight, E., Enikolopov,
G., Fernandez☐Burgos, M., Herrera, F., Adams, J. M., & Correa, E. (2011).
Sensitivity of leaf size and shape to climate: global patterns and paleoclimatic
applications. *New phytologist* **190**(3): 724-739.
R Core Team. (2018). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
Renninger, H. J., & Phillips, N. G. (2016). Palm physiology and distribution in response to
global environmental change. In: Goldstein G., Santiago L. eds. *Tropical tree physiology*.
Cham, Switzerland. Springer, 67-101
Seabold, S. & Perktold, J., (2010). Statsmodels: Econometric and statistical modelling with
python. In Proceedings of the 9th Python in Science Conference **57**(61): 10-25080.
Simard, M., Pinto, N., Fisher, J.B. & Baccini, A., (2011). Mapping forest canopy height
globally with spaceborne lidar. Journal of Geophysical Research: Biogeosciences,
116(G4).
Tomlinson, P. B. (2006). The uniqueness of palms. *Botanical Journal of the Linnean Society*
151(1): 5-14.
Tsukaya, H. (2006). Mechanism of leaf-shape determination. *Annual Review of Plant Biology*
57: 477-496.
Valladares, F., Skillman, J.B. & Pearcy, R.W., (2002). Convergence in light capture
efficiencies among tropical forest understory plants with contrasting crown architectures:
a case of morphological compensation. *American Journal of Botany*, **89**(8): 1275-1284.
Vogel, S. (2009). Leaves in the lowest and highest winds: temperature, force and shape. *New*
Phytologist **183**(1): 13-26.
Wright, I. J., Dong, N., Maire, V., Prentice, I. C., Westoby, M., Díaz, S., Gallagher, R. V.,
Jacobs, B. F., Kooyman, R., Law, E. A., & Leishman, M. R. (2017). Global climatic
drivers of leaf size. *Science* **357**(6354): 917-921.
Zachos, J., Pagani, M., Sloan, L., Thomas, E., & Billups, K. (2001). Trends, rhythms, and
aberrations in global climate 65 Ma to present. *Science* **292**(5517): 686-693.
Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D., Farooq,
H., Herdean, A., Ariza, M., Scharn, R., & Svantesson, S. (2019). CoordinateCleaner:
Standardized cleaning of occurrence records from biological collection databases.
Methods in Ecology and Evolution **10**(5): 744-751.

671
672
673

Data Availability Statement

All data generated during this study and the coding scripts used to carry out analyses are
available as Supplementary Material in the following repository:
doi.org/10.6084/m9.figshare.21230453. Chelsa Bioclimate data are available at https://chelsa-
climate.org/downloads/ and can be downloaded using the wget command. PalmTraits v1.0
containing morphological traits for all palm species is available from Kissling et al. (2019),
Scientific Data. The Global Vegetation Height raster from Simard et al. (2011; Biogeosciences)
is available at https://www.sciencebase.gov/catalog/item/50f59de2e4b0114312ab0280. Scripts
for data curation are available at https://github.com/mftorres/palm_leaf

    Figures

Figure 1. Leaf shape distribution at a global scale and across the Palm phylogeny. Left: The maps show the approximate distribution of shapes based on cleaned GBIF records (see Methods). Leaf silhouettes, originally by Marion Ruff Sheahan, were modified from Genera Palmarum (Dransfield et al., 2008). **Right:** The calibrated maximum clade credibility tree of 2550 palm species (Faurby et al., 2016; Hill et al., 2021). Dark and light greys are used to distinguish between subfamilies. The outermost ring shows the distribution of leaf shapes across the phylogeny and follows the same colour scheme in the maps. Species with no leaf shape colours are climbing and were removed from the Generalised Linear Mixed Models. **a)** Palmate + Costapalmate, **b)** Pinnate, **c)** Polymorphic, **d)** Entire, **e)** Bipinnate. **Cer.=** Ceroxyloideae. **N.**=Nypa.

Figure 2. Estimate distributions from the Generalised Linear Mixed Models (GLMMs).
Leaf shapes show the pair compared in the models (logistic regression; left=0, right=1). Each row shows the results from the models concerning hypotheses H1-3. Palmate = (costapalmate+palmate). We tested hypotheses H1-2 for each shape separately and for entire vs. dissected (palmate + costapalmate + pinnate), H1-2a, H1-2b, H1-2c, and H1-2d correspond to the models including either plant height or height over canopy, or temperature seasonality or precipitation. Hypothesis H3 was tested for each shape separately and H3a and H3b correspond to the models including either temperature seasonality or precipitation. Leaf silhouettes, originally by Marion Ruff Sheahan, were modified from Genera Palmarum (Dransfield et al., 2008).

Figure 3. Ancestral state estimation of leaf shape across palms (Arecaceae) using the
calibrated maximum clade credibility tree (Faurby et al., 2016; Hill, et al., 2021). The pie
charts show the probability of each state at the given node. States are colour-coded as follows:
Blue: Palmate (costapalmate+palmate); Red: Pinnate; Green: Entire; Yellow: Polymorphic
(where individuals of the same species have either entire or pinnate leaves). Leaf silhouettes,
originally by Marion Ruff Sheahan, were modified from Genera Palmarum (Dransfield et al.,
2008). **N.**=Nypa, **Cal.**=Calamoideae, **Cor.**=Coryphoideae, **Cer.**=Ceroxyloideae,
Are.=Arecoideae.

Figure 4. Palm leaf shape through time. (a). Top: Relative proportion of lineages with either entire-leaved or dissected shapes compared to global temperature variation through time. **Bottom:** Relative proportion of lineages with either pinnate or palmate (palmate+costapalmate) shapes compared to global temperature variation through time. **(b).** Relative proportion of lineages with (from top to bottom) palmate-dissected leaves, pinnate-dissected leaves, pinnate-entire leaves, or are polymorphic (intraspecific variation between pinnate-entire and pinnate-dissected). The proportion of palmate-entire lineages is close to zero and not plotted. In **(a)** and **(b)**, the white lines show the average proportion and the coloured lines show the relative proportion sampled from the 100 ancestral state estimations. **(c).** Relative probability of the ancestral state at the root of Arecaceae across 100 random trees sampled from the posterior distribution of the MCMC based on the five-state model. The horizontal bars at the bottom show the width of the posterior distribution and triangles show the default starting parameter. Leaf

795 silhouettes, originally by Marion Ruff Sheahan, were modified from Genera Palmarum
796 (Dransfield et al., 2008).
797
798 **Supplementary Material:**
799
Palmleaf_suppmat_20230417.zip
– **palm_leaf_shape_20221201** contains the leaf shape annotations
– **palms_alltraits_curated_20220620 c**ontains the median variables annotated by
species
– **SuppMat**
– Figure S1 Correlation coefficients across all variables.
– Figure S2 Species median distributions by leaf shape.
– Figure S3 Paramenters of the best MuSSE model for entire- and dissected-leaved
states.
– Figure S4 Paramenters of the best MuSSE model for pinnate- and palmate-leaved
states.
– Figure S5 Paramenters of the best MuSSE model for the five-state model.
– Figure S6 Entire- and dissected-leaved ancestral state estimation across palms.
– Figure S7 Pinnate- and palmate-leaved ancestral state estimation across palms.
– Figure S8 Posterior distribution of parameters for the MuSSE model for entire-
and dissected-leaved states.
– Figure S9 Posterior distribution of parameters for the MuSSE model for pinnate-
and palmate-leaved states.
– Figure S10 Five-trait ancestral state estimation across palms.
– Figure S11 State probability at the root based on the best-fitting model for entire-
and dissected-leaved states.
– Figure S12 State probability at the root based on the best-fitting model for
pinnate- and palmate-leaved states.
– Figure S13 Map of species distribution centroids for all palms
– Figure S14 Map of global distribution of plant heights for all palm species by leaf
shape.
– **Supplementary publication tables**
– Table S1 Number and percentage of species by leaf shape as used in the
Generalised Linear Mixed Models.
– Table S2 Palms exhibiting within-species leaf shape variation.
– Table S3 Species not considered in the analyses. Most species in the list are
climbing or have climbing and no-climbing habits.
– Table S4 Selected predictive variables used to fit the Bayesian phylogenetic
Generalised Linear Mixed Models.

- Table S5 Median (Minimum - Maximum) and range (delta) of species medians by shape and variable for species used in the Generalised Linear Mixed Models.
- Table S6 Median (Minimum - Maximum) and range (delta) of species medians by shape and variable for species used in the five-state ancestral state estimation.
- Table S7 Median (Minimum - Maximum) and range (delta) of species medians by shape and dissection for species used in the two-state ancestral state estimations.
- Table S8 Median, effective sample sizes, and 95% confidence intervals for the estimates in all Generalised Linear Mixed Models.
- Table S9 Entire-leaved versus dissected leaves. Medians and standard deviation of parameter estimates for alternative ML runs on the best fitting model.
- Table S10 Pinnate versus palmate shapes. Medians and standard deviation of parameter estimates for alternative ML runs on the best fitting model.
- **SupplementaryMaterial_20230105 contains supplementary methods and figure legends**