Transcriptomic profiling of sex-specific olfactory neurons reveals subset-specific receptor

expression in *Caenorhabditis elegans*

Author Names and Affiliations:

Douglas K. Reilly[1,7,†], Erich M. Schwarz[2,†], Caroline S. Muirhead[1], Annalise N. Robidoux[1,3],

Anusha Narayan[4,5], Meenakshi K. Doma[4,6], Paul W. Sternberg[4], Jagan Srinivasan[1,8,*]

[1]Department of Biology and Biotechnology, Worcester Polytechnic Institute, Worcester, MA

01605, USA

[2]Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853, USA

[3]Department of Chemistry and Biochemistry, Worcester Polytechnic Institute, Worcester, MA

01605, USA

[4]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena,

CA 91125, USA

[5]Present address: Merck and Co, 2000 Galloping Hill Road, Kenilworth, NJ 07033, USA

[6]Present address: Individualized Cell and Gene Therapies, South San Francisco, Genentech, CA

94080, USA

[7]Present address: Biology Department, Tufts University, Medford, MA 02155, USA

[8]Bioinformatics and Computational Biology Program, Worcester Polytechnic Institute, Worcester,

MA 01605, USA

*Corresponding author: Jagan Srinivasan, *jsrinivasan@wpi.edu*

†These authors contributed equally to this work

Key words: *C. elegans*, Pheromone, Sex-specific, G Protein-Coupled Receptors, Transcriptomes,

Single-Cell

Summary

The nematode *Caenorhabditis elegans* utilizes chemosensation to navigate an ever-

changing environment for its survival. A class of secreted small-molecule pheromones, termed

ascarosides, play an important role in olfactory perception by affecting biological functions

ranging from development to behavior. The ascaroside ascr#8 mediates sex-specific behaviors,

driving avoidance in hermaphrodites and attraction in males. Males sense ascr#8 via the ciliated

male-specific cephalic sensory (CEM) neurons, which exhibit radial symmetry along dorsal-

ventral and left-right axes. Calcium imaging studies suggest a complex neural coding mechanism

that translates stochastic physiological responses in these neurons to reliable behavioral outputs.

To test the hypothesis that neurophysiological complexity arises from differential expression of

genes, we performed cell-specific transcriptomic profiling; this revealed between 18 and 62 genes

with at least two-fold higher expression in a specific CEM neuron type versus both other CEM

neurons and adult males. These included two G protein coupled receptor (GPCR) genes, *srw-97*

and *dmsr-12*, that were specifically expressed in non-overlapping subsets of CEM neurons and

whose expression was confirmed by GFP reporter analysis. Single CRISPR-Cas9 knockouts of

either *srw-97* or *dmsr-12* resulted in partial defects, while a double knockout of both *srw-97* and

dmsr-12 completely abolished the attractive response to ascr#8. Together, our results suggest that

the evolutionarily distinct GPCRs SRW-97 and DMSR-12 act non-redundantly in discrete

olfactory neurons to facilitate male-specific sensation of ascr#8.

Introduction

The ability of an organism to find a mate is critical to the survival of a species. Many

species utilize small molecule pheromones to signal mate location (PUNGALIYA *et al.* 2009;

NARAYAN *et al.* 2016), sexual maturity (APRISON AND RUVINSKY 2015; APRISON AND RUVINSKY

2017), and receptivity (HOUCK *et al.* 2007; JANG *et al.* 2017), which are sensed and processed via

the nervous system, driving behavioral and developmental responses.

The nematode *Caenorhabditis elegans* communicates with conspecifics almost exclusively

using pheromones called ascarosides (LUDEWIG *et al.* 2019; MCGRATH AND RUVINSKY 2019), a

structurally conserved, modular class of small molecule pheromones (VON REUSS AND SCHROEDER

2015; ZHANG *et al.* 2017) composed of a core ascarylose sugar and a fatty-acid derived side chain

(BUTCHER *et al.* 2007; VON REUSS *et al.* 2012; LUDEWIG *et al.* 2019). Ascarosides signal a host of

environmental and developmental information, including the sexual maturity and location of

potential mates (NARAYAN *et al.* 2016; APRISON AND RUVINSKY 2017), and this information is

often conserved across nematodes (CHOE *et al.* 2012; RAGSDALE *et al.* 2013; DONG *et al.* 2016;

DONG *et al.* 2018; REILLY *et al.* 2019).

While most male-specific neurons are located in the tail, the major contributor to male-

specific chemosensory-driven behaviors are the CEM neurons, located in the amphid. Previous

studies have shown these four radially symmetric neurons to be involved in sensing ascaroside #3

and #8 (ascr#3 and ascr#8), two pheromones that serve to attract males to mates (PUNGALIYA *et*

al. 2009; NARAYAN *et al.* 2016; REILLY *et al.* 2017). Ascr#3 also plays a role in dauer formation,

functioning alongside ascr#2 and ascr#4 as the "dauer pheromone" (BUTCHER *et al.* 2008). Ascr#8

is unique in that it contains a *p*-aminobenzoic acid moiety on its terminus (PUNGALIYA *et al.* 2009;

ARTYUKHIN *et al.* 2018). In animals lacking CEM the attractive response to ascr#8 is abolished,

while further ablation of the ASK is required for complete loss of the ascr#3 behavioral response

(NARAYAN *et al.* 2016). Ablating three of the four CEM neurons, while leaving one CEM intact,

revealed that the four CEM neurons cooperate to drive a tuned attractive response to an

intermediate (1 µM) concentration of both ascr#3 and ascr#8 (NARAYAN *et al.* 2016).

Calcium imaging and electrophysiological experiments demonstrated that CEM neurons

show variable responses to stimuli not only between different animals, but within a single animal

(NARAYAN *et al.* 2016; REILLY *et al.* 2017). While neurons such as AWC exhibit variable calcium

responses between different animals, such responses are consistent between the left and right AWC

neurons of a single animal (COCHELLA *et al.* 2014). In contrast, the responses of individual CEM

neurons are stochastic within single animals, and yet four CEM neurons within one animal

consistently generate proper behavioral responses (NARAYAN *et al.* 2016). To understand this

pattern of stochastic neuronal activity yielding consistent behavioral outputs, it is imperative to

uncover genes encoding components of the CEM response.

A recent study in *C. elegans* hermaphrodites generated transcriptomic landscapes of 118

neuronal classes from 302 neurons in order to link functional and anatomical properties of

individual neurons with their molecular identities (TAYLOR *et al.* 2021). Discrete neuronal classes

were successfully identified via their combinations of expressed neuropeptides and neuropeptide

receptor genes. However, a similar feat has yet to be performed on male *C. elegans*, and the

transcriptomic profiles of individual male neurons remain enigmatic.

In a more focused transcriptomic approach, gene expression profiles of extracellular

vesicle-releasing neurons (EVNs) were identified (WANG *et al.* 2015; KALETSKY *et al.* 2016).

While CEM neurons are a subset of EVN neurons, and CEM transcriptomic data are embedded in

these data, there were thousands of EVNs in each replicate, diluting CEM-specific expression

values.

To begin understanding how CEM neurons achieve stochastic yet reliable physiological

responses to ascr#8, we performed single-cell RNA-seq (SCHWARZ *et al.* 2012) on CEM neurons,

and uncovered a small number of highly enriched genes encoding G protein-coupled receptors

(GPCRs). Given that all ascaroside receptors identified to date have been GPCRs (KIM *et al.* 2009;

MCGRATH *et al.* 2011; PARK *et al.* 2012; GREENE *et al.* 2016a; GREENE *et al.* 2016b; CHUTE *et al.*

2019), we tested whether any of these enriched genes contribute to male *C. elegans* sensation of

ascr#8. We identified two distantly related GPCR genes that contribute to the ascr#8 behavioral

response, *srw-97* and *dmsr-12*. Loss of each receptor resulted in partially deficient behavioral

responses to ascr#8. Following generation of a double *srw-97;dmsr-12* mutant, we found that loss

of both receptors results in a complete loss of behavioral response to ascr#8. Phylogenetic analysis

further indicates that both receptors are homologous to closely related receptors present throughout

the *Caenorhabditis* genus, and robust ascr#8 responses may be a trait recently evolved in

C. elegans.

Results

The transcriptomic landscape of CEM neurons is variable

Individual CEM neurons were isolated from *C. elegans* expressing an integrated GFP

labeling male-specific EVNs (p*pkd-2*::GFP [**Figure 2A**]), as previously described (GOODMAN *et*

al. 1998; NARAYAN *et al.* 2011; NARAYAN *et al.* 2016). Cells were separated by anatomical identity

(i.e., CEM dorsal left [DL], dorsal right [DR], ventral left [VL], and ventral right [VR]); single-

cell CEM cDNA libraries were constructed and used for RNA-seq (SCHWARZ *et al.* 2012).

Enriched genes in each CEM neuron were identified by comparing RNA-seq profiles of

distinct CEM types to one another and to whole adult *C. elegans* males (**Table S1, S5**). We

observed 267 genes with at least two-fold higher expression in CEM neurons than in adult males.

Of these, we observed 164 genes that were specific to individual CEM types, defining specificity

as being at least two-fold higher expression in one type than in any other type: 62 genes specific

to CEM DL, 40 to CEM DR, 18 to CEM VL, and 44 to CEM VR. Uniquely mapped reads ranged

from 5.2 to 9.2 million per CEM type, matching the trend for alignment rates of each CEM neuron,

which ranged from 18.0% to 47.2%, (**Table S2**) with an average of 1,426 genes showing robust

RNA expression in each neuron (**Table S3**).

Five GPCR-encoding genes expressed in CEMs (*seb-3, srr-7, srw-97, dmsr-12, srd-32*)

were selected for study; of these, *srw-97* and *dmsr-12* showed specific expression in CEM VR and

CEM DL. Four of these GPCR genes were uncharacterized; one (*seb-3*) has previously been shown

to play roles in locomotion, stress response, and ethanol tolerance (JEE *et al.* 2013). *dmsr-12* is

related to *daf-37* (ROBERTSON AND THOMAS 2006), a previously identified ascaroside receptor

gene (PARK *et al.* 2012), although it is more closely related to the neuropeptide receptor gene *dmsr-*

1, and more distantly to *srw-97* (ROBERTSON AND THOMAS 2006). *srd-32* belongs to a divergent

branch of the SRD phylogeny (ROBERTSON AND THOMAS 2006). *srr-7* belongs to the one of the

smallest families of *C. elegans* chemoreceptor genes (ROBERTSON AND THOMAS 2006).

seb-3 and *srw-97* exhibited similar enrichment profiles across the CEM neurons, with 3.0-

and 3.3-fold higher expression in CEM VR than in any other CEM neuron (**Table 1; Figure 1D;**

Table S5). *dmsr-12* showed 25-fold higher expression in CEM DL versus any other CEM neuron

(**Table 1; Figure 1A; Table S5**), while *srd-32* was only 1.8-fold higher in CEM DL (**Table 1;**

Figure 1A; Table S5). *srr-7* showed 4.2-fold higher expression in CEM VL versus any other CEM

neuron (**Table 1; Figure 1C; Table S5**), which correlated with previous transcriptomic analyses

that found *srr-7* to be enriched in *C. elegans* EVNs (WANG *et al.* 2015).

CEM-specific receptor expression patterns

To confirm our single-cell RNA-seq results, we generated transgenic GFP fusions for the

five receptor genes (BOULIN *et al.* 2006). Roughly 3 kb of promoter region upstream of the start

codon was included in these constructs, along with the majority of the coding sequence (**Table**

S6); this would have automatically included any large 5'-ward introns that might contain cis-

regulatory elements of these genes (FUXMAN BASS *et al.* 2014). The GFP coding sequence was

cloned from the Fire Kit vector pPD95.75 (BOULIN *et al.* 2006). *pha-1; lite-1; him-5* animals were

injected with reporter constructs and the co-injection marker pBX (*pha-1*(+)). We isolated GFP$^+$

strains and imaged GFP$^+$ males for expression at 63x magnification (**Figure 2; Fig. S1**). An

integrated p*pkd-2*::GFP line was used as a CEM-specific control (**Figure 2A**).

The previously characterized GPCR gene, *seb-3*, displayed a non-CEM-specific expression

pattern matching that previously described (**Figure 2B**). The other four receptor genes showed

transgenic expression patterns similar to their RNA-seq data (**Figure 2C-E**). SRR-7 was observed

in CEM VR, as well as another neuron that is likely to be the mechanoreceptor CEP VR (SAWIN

et al. 2000) (**Figure 2F**). Both *seb-3* and *srr-7* were excluded from further analyses, as we aimed

to identify CEM-specific regulators of the ascr#8 response.

Except for SRD-32, all GFP-tagged receptors showed subcellular localization patterns that

included the sensory cilia (**Figure 2, white bars**). Our failure to see this for SRD-32 may be an

artifact of our p*srd-32*::GFP construct design, because only 52% of the *srd-32* coding sequence

was included in our transgene (**Table S6**). We also observed non-GPCR genes to be specifically

expressed in single CEM neurons, such as *trf-1*, which encodes a TNF receptor homolog (TENOR

AND ABALLAY 2008) expressed in CEM VR (**Table 1**); *trf-1* has an EVN-specific promoter (WANG

et al. 2015), providing an alternative to the *pkd-2* and *klp-6* promoters typically used to drive

transgenes in EVNs (**Fig. S1**) (PEDEN AND BARR 2005; BAE *et al.* 2006).

RNAi-mediated knockdown of CEM Receptors

To test whether these CEM-enriched genes encode receptors that are required in ascr#8

sensation, we fed dsRNA to a strain that is hypersensitive to neuronal RNA interference,

nre-1; lin-15B (SCHMITZ *et al.* 2007; POOLE *et al.* 2011).

To confirm that our system for male neuronal RNAi could affect behavioral phenotypes,

we first fed animals *osm-3* and *osm-9* dsRNA clones from the Ahringer Library (FRASER *et al.*

2000; KAMATH *et al.* 2003). Using a Spot Retention Assay (NARAYAN *et al.* 2016), we assayed

animals for their behavioral dwell time in ascr#8 (**Figure S2A**). Animals fed *osm-3* dsRNA

showed significantly defective responses to ascr#8, as observed for loss-of-function alleles

(**Figure S2B**). In contrast, animals fed *osm-9* showed only a slight decrease in their times spent

within ascr#8 (**Figure S2A**). Because we could successfully abolish ascr#8 attraction through male

neuronal RNAi of *osm-3*, we were confident that RNAi would be effective at functionally verifying

CEM-enriched genes encoding components of the ascr#8 response.

We fed animals dsRNA clones targeting three CEM-enriched receptor genes: *srd-32*, *dmsr-*

12, and *srw-97*. A *srd-32* clone was unavailable in the Ahringer library (FRASER *et al.* 2000;

KAMATH *et al.* 2003), but a present in the Vidal library (RUAL *et al.* 2004). Each library uses the

same backbone vector, allowing the same control to be used for both.

RNAi of *srd-32* caused no defect in ascr#8 responses (**Figure S2C**). Both *dmsr-12* or *srw-*

97 dsRNA caused partial defects of ascr#8 dwell time (**Figure S2C**). Neither knockdown

statistically lowered the dwell time in ascr#8 (**Figure S2D**), though they did abolish the statistically

significant increase in time spent in ascr#8 over vehicle controls (**Figure S2C**).

CRISPR-generated null mutants of candidate receptors

For phenotypic analysis of null mutants, we used our Single Worm Attraction Assay

(SWAA), as previously described (REILLY *et al.* 2021). Wild-type (*him-5*) animals were strongly

attracted to ascr#8 in the SWAA, replicating previous observations (**Figure 3A, B; Fig. S3**)

(REILLY *et al.* 2021).

srw-97(knu456) males displayed partially defective ascr#8 attraction: their ascr#8 dwell

time was no different than that of vehicle, nor was it different that *him-5* animals (**Figure 3A, B;**

Fig. S3). We were able to restore normal ascr#8 attraction in *srw-97(knu456)* males through

transgenesis with a translational fusion construct (p*srw-97*::*srw-97*::GFP) (**Fig. S4**). Expression of

the rescue transgene matched that of our initial fusion, with GFP visible in both ventral CEM

neurons (**Figure 3C; Fig. S5**).

Unlike *srw-97(knu456)*, *dmsr-12(tm8706)* males exhibited completely defective responses

to ascr#8 (**Figure 3A, B; Fig. S3**). Similarly, the expression profile of the p*dmsr-12*::*dmsr-*

12::GFP rescue construct (**Fig. S4**) matched its earlier reporter expression (**Figure 3D; Fig. S5**),

and completely restored attraction to ascr#8 (**Figure 3A, B; Fig. S3**). Quantification of GFP

expression confirmed SRW-97 to be enriched within the CEM_VL neuron, and to a lesser extent

CEM_VR and CEM_DL neurons, while DMSR-12 was observed within CEM_DL and CEM_DR

neurons (**Table S7**).

Given that *dmsr-12* is expressed in dorsal CEM neurons, and *srw-97* is expressed in ventral

CEM neurons (**Figure 3C, D**), we speculated that single mutants of either receptor partially retain

the ability to sense and respond to ascr#8, with the remaining CEM-specific receptor gene

conferring some residual response to ascr#8. We thus generated a *dmsr-12; srw-97* double mutant

strain and assayed animals for their ability to respond to ascr#8. These double mutants were not

attracted to ascr#8 compared to the vehicle control (**Figure 3A; Fig. S3**), though they were not

significantly defective in their attraction compared to wild-type animals either (**Figure 3B**).

However, this is likely a result of lower vehicle dwell times in the double mutants. This defect was

specific to ascr#8; double mutants showed no defect in their responses to ascr#3 (**Figure 3E, F;**

Fig. S6).

Together, these data suggest that at least two GPCR receptors are expressed in discrete

pairs of CEM neurons and act in parallel during sensation of ascr#8.

Phylogenetic analyses of ascr#8 receptors reveal likely gene duplication events

The ability to attract mates through pheromones is often essential for a species' survival.

We have recently observed that different *Caenorhabditis* species show variable levels of attraction

to ascr#8 (REILLY *et al.* 2019). To better understand variability in levels of attraction, we analyzed

the evolution of both *srw-97* and *dmsr-12*.

We identified orthologs of *srw-97* from *C. elegans* (*Cel-srw-97*) in other *Caenorhabditis*

proteomes via OrthoFinder (**Figure 4A**). A closely related *C. elegans* paralog, *Cel-srw-98*,

underwent a species-specific expansion within *C. inopinata*. The coding DNA sequences (CDS

DNAs) of both *Cel-srw-97* and *Cel-srw-98* are 66.2% identical to one another, while their amino

acid sequences are only 57.1% identical (MADEIRA *et al.* 2019). *Cel-srw-98* shows enriched

expression in CEM VR that is similar to (though weaker than) that of *Cel-srw-97* (**Table 1; Table**

S5).

C. tropicalis, like *C. elegans* and *C. briggsae*, is a hermaphroditic *Caenorhabditis* species

(NOBLE *et al.* 2021) which encodes a reduced number of ascr#8 receptor gene paralogs, with no

Cel-dmsr-12 paralogs, and only one *Cel-srw-97* paralog (**Figure 4**). We previously observed that

C. tropicalis fails to be attracted to ascr#8, and it is possible that this loss of receptor genes is one

reason for that failure (REILLY *et al.* 2019).

In contrast, the other candidate ascr#8 receptor, *dmsr-12*, does not appear in our

phylogenetic analysis to have undergone any species-specific expansion (**Figure 4B**). Rather, there

are generally fewer paralogs per non-*C. elegans* species.

Discussion

Pheromones are important for mating in many animals. In the nematode *C. elegans*, males

are attracted to hermaphrodites as possible mates by the small molecule pheromone ascaroside #8

(PUNGALIYA *et al.* 2009; NARAYAN *et al.* 2016). While the ascaroside class of small molecule

pheromones utilized by nematodes is rapidly being elucidated (with over 230 known ascaroside

structures so far; *https://smid-db.org*) (ARTYUKHIN *et al.* 2018), the neuronal receptor proteins that

mediate pheromone signals remain largely unknown. For only a select few ascarosides have

sensory components been identified at the cellular (GREENE *et al.* 2016a; GREENE *et al.* 2016b;

CHUTE *et al.* 2019), receptor (KIM *et al.* 2009; MCGRATH *et al.* 2011; PARK *et al.* 2012), or signal

transduction levels (ZWAAL *et al.* 1997).

Here, we identify two novel G protein-coupled receptors as active, required components in

sensory and behavioral responses to the mating pheromone ascr#8. Transcripts for the two GPCRs,

dmsr-12 and *srw-97*, are enriched in single CEM neurons (**Table 1**), and they express in non-

overlapping subsets of the male-specific chemosensory neurons (**Figure 1, 2**). There are, however,

other receptors present in these same neurons that contribute to the navigation of a vast and ever-

changing array of environmental cues, such as the widely expressed ethanol sensor *seb-3* (**Figure**

2B) (JEE *et al.* 2013).

One mechanism for sensory flexibility in CEM neurons may be heterodimerization of

receptor proteins. Previous work identified two receptors for ascr#2 in the ASK neuron: DAF-37

and DAF-38 (PARK *et al.* 2012). While both are required for proper ascr#2-induced dauer

formation, only DAF-38 is involved in the sensation of other ascarosides (PARK *et al.* 2012). Such

heterodimers may also exist for pheromone receptors in CEM neurons.

Another mechanism for sensory flexibility in CEM neurons is to have diverse receptor

proteins with distinct specificities co-expressed within a single neuron. Multiple ascarosides are

sensed by the male-specific CEM neurons, including ascr#8 and ascr#3 (NARAYAN *et al.* 2016).

The two receptors identified here may function as ascr#8-specific receptors, as *dmsr-12;srw-97*

double mutant animals do not exhibit defective responses to ascr#3 (**Figure 3E, F**). The related

receptors *dmsr-10*, *dmsr-13*, *dmsr-16*, and *srw-98* are enriched in ventral CEM neurons, and may

heterodimerize with *srw-97*, to result in fine-tuned sensation of ascr#8. Similarly, there may be

other receptors may be enriched in the dorsal CEM neurons to heterodimerize with *dmsr-12*. The

role of DAF-37 as an ascr#2-specific receptor further supports our hypothesis, as *dmsr-12* is related

to DAF-37 (ROBERTSON AND THOMAS 2006). The ventral CEM receptor SRW-97 falls within the

same large family of GPCRs as well (KRISHNAN *et al.* 2014), while the promiscuous DAF-38 does

not (ROBERTSON AND THOMAS 2006).

The initial goal of our single-cell RNA-seq analysis of male-specific chemosensory CEM

neurons was to identify GPCRs encoding pheromone receptors. However, further analysis of other

genes with CEM-enriched expression should also uncover specific and novel promoter profiles,

which will enable optogenetic manipulation and calcium imaging of defined individual CEM

neurons (NARAYAN *et al.* 2016). The development of advanced transgenic reagents will also permit

chemical biology of CEM neurons. We have recently developed an active ascr#8 bioaffinity probe

to employ in the targeted elucidation of ascr#8 receptors (ZHANG *et al.* 2019). Use of this probe

will lead to further elucidation of the identity of acsr#8 receptors, by confirming either SRW-97

or DMSR-12 as receptors or identifying their heterodimeric partners. The combination of these

technologies should clarify how heterogeneous CEM neurons achieve homogeneous sensory

responses.

Methods

Single cell isolation and RT-PCR

Individual ciliated male-specific cephalic sensory (CEM) neurons were isolated from adult

males of the *C. elegans* strain CU607 (smIs23 [p*pkd-2*::GFP+ pBX]; *him-5*(e1490)), as previously

described (NARAYAN *et al.* 2016). This strain expresses an integrated GFP transgene that labels

male-specific extracellular vesicle-releasing neurons (EVNs, p*pkd-2*::GFP [Figure 2A]). Cells

were separated by anatomical identity (i.e., CEM dorsal left [CEM_DL], CEM dorsal right

[CEM_DR], CEM ventral left [CEM_VL], and CEM ventral right [CEM_VR]). Microdissection

and single-cell RT-PCR of individual CEM_DL, CEM_DR, CEM_VL, and CEM_DR neurons

was performed essentially as described (SCHWARZ *et al.* 2012). For all four neuronal types, single-

end 50-nucleotide (50-nt) to 100-nt RNA-seq was performed on an Illumina HiSeq 2000. To

identify so-called housekeeping genes and genes primarily active outside the nervous system, we

compared our results from CEM_DL, CEM_DR, CEM_VL, and CEM_DR to equivalent

reanalyses of published single-end 38-nt RNA-seq data from mixed-stage whole *C. elegans*

hermaphrodite larvae (SCHWARZ *et al.* 2012), and of published single-end 36-nt RNA-seq data

from adult whole *C. elegans* males and females (THOMAS *et al.* 2012).

Transcriptional analysis

Reads from RT-PCR-amplified CEM neurons and larvae were quality-filtered as follows:

neuronal reads that failed Chastity filtering were discarded (Chastity filtering had not been

available for the larval reads); raw 38-nt larval reads were trimmed 1 nt to 37 nt; all reads were

trimmed to remove any indeterminate ("N") residues or residues with a quality score of less than

3; and larval reads that had been trimmed below 37 nt were deleted, as were neuronal reads that

had been trimmed below 50 nt. For uniformity of analysis, all RNA-seq reads of 51-100 nt in

length were trimmed to 50 nt before further RNA-seq analysis. This left a total ranging from

55,902,297 (CEM_VL) to 73,412,777 (CEM_VR) filtered reads for analysis of each neuronal type,

versus 23,369,056 filtered reads for whole larvae (**Table S2**). Previously published RNA-seq reads

(THOMAS *et al.* 2012) from three biological replicates of adult males (accession numbers

SRR580386, SRR580387, and SRR580388) and adult females (accession numbers SRR580383,

SRR580384, and SRR580385) were downloaded from the European Nucleotide Archive

(*ftp://ftp.sra.ebi.ac.uk*) and used without further trimming or filtering.

We used RSEM version 1.2.17 (LI AND DEWEY 2011) with bowtie2 version 2.2.3

(LANGMEAD AND SALZBERG 2012) and SAMTools version 1.0 (LI *et al.* 2009) to map filtered reads

to a *C. elegans* gene index and generate read counts and gene expression levels in transcripts per

million (TPM). To create the *C. elegans* gene index, we ran RSEM's *rsem-prepare-reference* with

the arguments "*--bowtie2 --transcript-to-gene-map*' upon a collection of coding DNA sequences

(CDSes) from both protein-coding and non-protein-coding *C. elegans* genes in WormBase release

WS245 (HOWE *et al.* 2016). The CDS sequences were obtained from *ftp://ftp.sanger.ac.uk/pub2/*

wormbase/releases/WS245/species/c_elegans/PRJNA13758/c_elegans.

PRJNA13758.WS245.mRNA_transcripts.fa.gz and *ftp://ftp.sanger.ac.uk/pub2/wormbase/*

*releases/WS245/species/c_elegans/PRJNA13758/c_elegans.PRJNA13758.WS245.ncRNA_*

transcripts.fa.gz. For each RNA-seq data set of interest, we computed mapped reads and

expression levels per gene by running RSEM''s *rsem-calculate-expression* with the arguments

"*--bowtie2 -p 8 --no-bam-output --calc-pme --calc-ci --ci-credibility-level*

0.99 --fragment-length-mean 200 --fragment-length-sd 20 --estimate-rspd --ci-memory 30000'.

These arguments, in particular '*--estimate-rspd*', were aimed at dealing with single-end data from

3'-biased RT-PCR reactions; the argument '*--phred64-quals*' was used for the larval reads, while

'*--phred33-quals*' was used for all other reads (neuronal, adult males, and adult females). We

computed posterior mean estimates (PMEs) both for read counts and for gene expression levels,

and rounded PMEs of read counts down to the nearest lesser integer. We also computed 99%

credibility intervals (Cis) for expression data, so that we could use the minimum value in the 99%

CI for TPM as a robust minimum estimate of a gene's expression (minTPM).

We observed the following overall alignment rates of reads to the WS245 *C. elegans* gene

index: 47.16% for the CEM_DL read set, 27.37% for the CEM_DR read set, 40.20% for the

CEM_VL read set, 17.99% for the CEM_VR read set, and 76.41% for the larval read set (**Table**

S2). A similar discrepancy between lower alignment rates for hand-dissected linker cell RNA-seq

reads versus higher alignment rates for whole larval RNA-seq reads was previously observed, and

found to be due to a much higher rate of human contaminant RNA sequences in the hand-dissected

linker cells (SCHWARZ *et al.* 2012). For previously published adult male and female reads, we

observed mapping rates of 88.20% to 96.11%; this may reflect greater difficulty of getting

mappable reads from our RT-PCR-amplified cDNAs than from unamplified cDNA syntheses.

We defined detectable expression for a gene in a given RNA-seq data set by that gene

having an expression level of 0.1 TPM; we defined robust expression by that gene having a

minimum estimated expression level (termed minTPM) of at least 0.1 TPM in a credibility interval

of 99% (i.e., $\geq$ 0.1 minTPM). The numbers of genes being scored as expressed in a given neuronal

type above background levels, for various data sets, are given in **Table S3**. Other results from

RSEM analysis are given in **Table S5**.

We annotated *C. elegans* genes and the encoded gene products in several ways (**Table S5**).

For the products of protein-coding genes, we predicted classical signal and transmembrane

sequences with Phobius 1.01 (KÄLL *et al.* 2004), regions of low sequence complexity with pseg

(SEG for proteins, from *ftp://ftp.ncbi.nlm.nih.gov/pub/seg/pseg*) (WOOTTON 1994), and coiled-coil

domains with ncoils (from *http://www.russell.embl-heidelberg.de/coils/coils.tar.gz*) (LUPAS

1996). Protein domains from Pfam 31.0 (FINN *et al.* 2016) were detected with HMMER 3.1b2/

hmmsearch (EDDY 2009), using the argument '--*cut_ga*' to invoke model-specific reporting

thresholds. Memberships of genes in orthology groups from eggNOG 3.0 (POWELL *et al.* 2012)

were extracted from WormBase WS245 with the TableMaker function of ACEDB 4.3.39. Genes

with likely housekeeping status (based on ubiquitous expression in both larvae and linker cells)

were as identified in our previous work (SCHWARZ *et al.* 2012). Genes were predicted to encode

GPCRs on the basis of their encoding a product containing one or more of the following Pfam-A

protein domains: 7tm_1 [PF00001.16], 7tm_2 [PF00002.19], 7tm_3 [PF00003.17], 7tm_7

[PF08395.7], 7TM_GPCR_Srab [PF10292.4], 7TM_GPCR_Sra [PF02117.11],

7TM_GPCR_Srbc [PF10316.4], 7TM_GPCR_Srb [PF02175.11], 7TM_GPCR_Srd [PF10317.4],

7TM_GPCR_Srh [PF10318.4], 7TM_GPCR_Sri [PF10327.4], 7TM_GPCR_Srj [PF10319.4],

7TM_GPCR_Srsx [PF10320.4], 7TM_GPCR_Srt [PF10321.4], 7TM_GPCR_Sru [PF10322.4],

7TM_GPCR_Srv [PF10323.4], 7TM_GPCR_Srw [PF10324.4], 7TM_GPCR_Srx [PF10328.4],

7TM_GPCR_Srz [PF10325.4], 7TM_GPCR_Str [PF10326.4], ABA_GPCR [PF12430.3], Sre

[PF03125.13], and Srg [PF02118.16]. By this criterion, we identified 1,615 genes encoding

GPCRs in the WS245 version of the *C. elegans* genome; this resembles a previous estimate of

~1,470 *C. elegans* genes encoding chemoreceptors and other GPCRs, identified through extensive

computational and manual analysis (HOBERT 2013). The memberships of genes in orthology

groups from eggNOG 3.0 (POWELL *et al.* 2012) were extracted directly from WormBase WS245

with the TableMaker function of ACEDB 4.3.39. Genes with likely housekeeping status (based on

ubiquitous expression in both larvae and linker cells) were as identified in our previous work

(SCHWARZ *et al.* 2012). Gene Ontology (GO) annotations for *C. elegans* genes were extracted from

WormBase-computed annotations in *ftp://ftp.wormbase.org/pub/wormbase/releases/WS245/*

ONTOLOGY/gene_association.WS245.wb.c_elegans; human-readable text descriptions for GO

term IDs were extracted from *term.txt* in the Gene Ontology archive *http://archive.geneontology.*

org/full/2014-07-01/go_201407-termdb-tables.tar.gz.

GFP reporter construction

Reporter fusion constructs were generated using previously described techniques (BOULIN

et al. 2006). Approximately 2-3 kb of upstream promoter region of each gene was included in

construct generation, as well as a portion of the coding sequence (**Table S6**). This was then fused

to GFP (from the Fire Vector Kit plasmid, pPD95.75), via PCR fusion (BOULIN *et al.* 2006).

Primers were designed using Primer 3 and ordered from IDT (Integrated DNA Technologies).

Primer sequences available in **Table S9**. Successful fusion was confirmed via gel electrophoresis

prior to injection.

Reporter fusion constructs were injected into the gonads of *pha-1(e2123ts); lite-1(ce314);*

him-5(e1490) animals, along with a co-injection marker of pBX(*pha-1*(+)). In this manner, positive

array animals will propagate normally at 20 °C. Strains were confirmed via GFP expression, with

multiple array lines being generated per injection (**Table S8**). Injections were performed by InVivo

Biosystems, with strain isolation being performed in house.

Imaging

Animals were imaged for GFP expression using previously described techniques. GFP+

young adult male animals were mounted on a 1% agarose pad and immobilized with sodium azide.

Animals were then imaged on a spinning disk confocal microscope at 63x magnification. Z-stack

imaging was performed, generating 3D reconstructions of the heads of the imaged animals.

Central/optimal z-plane images were used to generate the images used to verify expression (**Figure**

2; Fig. S1).

RNAi Feeding

VH624 (*rhIs13* [*unc-119*::GFP + *dpy-20*(+)]; *nre-1(hd20)*; *lin-15B(hd126)*) animals

(SCHMITZ *et al.* 2007; POOLE *et al.* 2011) were crossed with *him-5(e1490)* animals to integrate

male production into a strain hypersensitive to neuronal RNAi-knockdown, generating JSR44

(*nre-1(hd20)*; *lin-15B(hd126); him-5(e1490)*). During the cross, insertion of the *him-5(e1490)*

allele displaced the integrated array *rhIs13*, suggesting location of the array on Chromosome V.

Presence of *lin-15B(hd126)* in JSR44 was confirmed via sequencing. The non-annotated

nre-1(hd20) is linked with *lin-15B*, being retained alongside *lin-15B* (SCHMITZ *et al.* 2007; POOLE

et al. 2011).

RNAi clones were grown overnight in cultures of LB containing 50 µg/mL ampicillin.

Cultures were then diluted to an OD_{600} of 1.0 before plating on NGM agar plates containing 50

µg/mL ampicillin and 1 mM IPTG (Isopropyl-β-D-thiogalactoside) to select for RNAi clones and

induce expression. Lawns were allowed to grow at room temperature for 8-16 hours, before JSR44

young adult hermaphrodites were placed on the plates and left to propagate at 16°C. Young adult

males of the F1 progeny were then selected for behavioral testing suing the *Spot Retention Assay*.

Empty vector controls (VC-1 clone) were run alongside every targeted knockdown experiment.

Spot Retention Assay

Following previously described methods, young adult males were isolated from

hermaphrodites 5-16 hours prior to testing (PUNGALIYA *et al.* 2009; NARAYAN *et al.* 2016). In

short, at the time of the assay, 0.6 µL of either vehicle control (-) or ascaroside #8 (+) was added

to the NGM plates covered in a thin lawn of OP50 *E. coli*. Ten males were then divided between

two pre-marked spots on the agar, equidistant from the cues. The plate was then recorded for 20

minutes. The time spent of each visit in either vehicle or ascaroside #8 (if greater than 10 seconds)

was scored and averaged. Plates in which the average was greater than two standard deviations

removed from the population average were removed from the final analysis as outliers. To compare

between strains or conditions, the vehicle was subtracted from the ascaroside dwell time for each

plate. The average of these differences was then compared statistically.

Single Worm Behavioral Assay

Following previously described methods (PUNGALIYA *et al.* 2009; NARAYAN *et al.* 2016;

REILLY *et al.* 2021), animals were isolated and prepared in an identical manner to the Spot

Retention Assay. The two outside rings of wells in a 48-well tissue culture plate were filled with

200 µL of NGM agar, which was then seeded with 65 µL of OP50 *E. coli*. The plates and lawns

were then dried at 37 °C for 4 hours. Alternating wells were then prepared as spatial controls ("S",

nothing done), vehicle controls ("V", 0.85 µL of dH$_2$O was placed in the center of the well), or

ascaroside well ("A", 0.85 µL of ascaroside was placed in the center of the well). This was

performed over four quadrants. Animals were scored for their visits and duration to the center of

the well and/or the cue.

The average duration of each worm's visits was calculated, and these values were again

averaged together to generate a Mean Dwell Time in seconds for each plate. When comparing

across strains or conditions, the Spatial controls were then compared for statistical difference. If

none was observed, the Log(fold-change) A/V was then calculated by taking the log of the

ascaroside mean dwell time divided by the vehicle mean dwell time for each plate. The number of

times each worm visited the center was averaged to generate the Visit Counts.

The Percent Attraction values were calculated by first determining the "attractive" cut off

as two standard deviations above the vehicle average. Any visit longer than this was deemed

"attractive" and scored as a "1"; non-attractive visits were scored as "0". The percent attraction

was then calculated for each worm was the percent of visits scored a "1". The average was then

calculated across the plate to determine Percent Attraction.

CRISPR Design and Strain Generation

A novel null mutation was generated for *srw-97* by InVivo Biosystems. The *srw-*

97(knu456) allele was generated in a *him-5(e1490)* strain using two sgRNAs

(TTTAGTAGAGCAGAAATTAA and TACAGCTTTAACTTTCAAC) to generate a 1,620-nt

deletion which removed the start codon and left only the terminal exon intact. The *knu456*

knockout allele was generated by donor homology using the pNU1361odn oligo:

(TTTTCTTGTATTTCCAAAAATTGTAAAAACCTTTATGAAAGTTAAAGCTGTAAGGAT

TTTCAGACATTTA). Following generation of a homozygous deletion by InVivo Biosystems,

the line was then backcrossed twice into *him-5(e1490)*.

The *dmsr-12(tm8706)* allele, provided by the National BioResource Group (NRBP)

contains a 118-nucleotide deletion that was generated by Dr. Mitani of the NRBP. The allele was

crossed into a *him-5(e1490)* background prior to testing. The deletion spans intron 2 and exon 3

of the coding sequence. Whether this results in a correctly spliced gene remains unknown, although

the expected remaining coding sequence remains in frame.

Phylogenetic Analyses

For phylogenetic analysis of selected CEM genes, we downloaded proteomes for

C. elegans and related *Caenorhabditis* nematodes from WormBase (release WS275), the Blaxter

Caenorhabditis database (release 1), or our unpublished work, as listed in **Table S4**. From each

proteome, we extracted the longest predicted isoform for each gene with *get_largest_isoforms.pl*

(*https://github.com/SchwarzEM/ems_perl/blob/master/fasta/get_largest_isoforms.pl*).   We

observed that the predicted isoform for *dmsr-12* in the WormBase WS275 release of *C. elegans*'

proteome was shorter than past versions of *dmsr-12*, and that the WS275 isoform omitted exons

that our transgenic expression data (based on older gene models for *dmsr-12*) indicated were likely

to be real. We therefore manually replaced the WS275 version of *dmsr-12* with an older version

(extracted from the *C. elegans* proteome in the WS250 release of WormBase). We then computed

orthology groups for *C. elegans* and its related species with OrthoFinder version 2.3.11 (EMMS

AND KELLY 2015; EMMS AND KELLY 2019), using the arguments '*-a 1 -S diamond -og*'. We

identified which orthology groups contained the *C. elegans* genes *srr-7*, *srw-97*, and *dmsr-12*, and

extracted their sequences from a concatenation of all 11 proteomes via *extract_fasta_subset.pl*

(*https://github.com/SchwarzEM/ems_perl/blob/master/fasta/extract_fasta_subset.pl*).   For   each

orthogroup's member sequences, we aligned the sequences with MAFFT version 7.455 (KATOH

AND STANDLEY 2013) and filtered the alignments twice with trimAl version 1.4.rev15 (CAPELLA-

GUTIÉRREZ *et al.* 2009), using first the argument '*-automated1*' and then the arguments

'*-resoverlap 0.50 -seqoverlap 50*'. From the filtered alignments, we computed maximum-

likelihood protein phylogenies with IQ-TREE version 2.0-rc1 (NGUYEN *et al.* 2015;

KALYAANAMOORTHY *et al.* 2017), using the arguments '*-m MFP -b 100 --tbe*'. In particular, we

used transfer bootstrap expectation ('*--tbe*') which provides more reliable confidence values than

classic bootstrapping (LEMOINE *et al.* 2018). We visualized the resulting phylogenies with FigTree

version 1.4.4 (*http://tree.bio.ed.ac.uk/software/figtree*).

Statistical Analyses

Prior to any statistical analyses, outliers were identified and removed. Outliers were

defined as any data points greater than two standard deviations removed from the average. All data

were then tested for normality using a Shapiro-Wilk Normality Test. This test was chosen over the

more conventional D'Agostino-Pearson Normality Test as many data sets were below 10 in

number (due to the statistical power offered by the Single Worm Behavioral Assay (REILLY *et al.*

2021).

The Spot Retention Assay data was analyzed using paired *t*-tests or Wilcoxon Matched-

Pairs Signed Rank tests to compare vehicle control and ascaroside dwell times within strains

following tests for normality (**Figure S2**). When comparing the values of multiple conditions or

strains, the data was first normalized to account for vehicle dwell time variation between plates

using a base-2 exponentiation, as described previously, to transform all data points into non-zero

values. This allows for the calculation of the fold-change as the log(base2) of the ascaroside dwell

time divided by the vehicle dwell time. These normalized values were then compared using a

Mann-Whitney test or a One-Way ANOVA followed by a Dunnett's multiple comparisons test

(**Figure S2**).

The Single Worm Behavioral Assay was first analyzed within each strain by performing

either a Repeated-Measures ANOVA followed by a Dunnett's multiple comparisons test or

Friedman test followed by Dunn's correction, comparing both Spatial control and Ascaroside

dwell times to that of the Vehicle Control. The log(fold-change) of raw ascaroside and vehicle

dwell time values were calculated and compared using either a One-Way ANOVA followed by a

Bonferroni's multiple corrections test or Friedman Test followed by Dunn's correction (**Figure 3**).

The ascr#3 log(fold-change) values were compared using a Student's *t*-test (**Figure 3F**). Visit

Counts were compared in the same manner as the Mean Dwell Time data, while the Percent

Attraction data was analyzed using paired *t*-tests or Wilcoxon Matched-Pairs Signed Rank tests to

compare the attractive values of the vehicle and ascaroside (**Figures S3, S4, S5**).

Data availability

RNA-seq reads have been deposited in the NCBI Sequence Read Archive (SRA) under the

BioProject accession number PRJNA781271.

Acknowledgements

We thank the *Caenorhabditis* Genetics Center, which is funded by the NIH Office of

Research Infrastructure Programs (P40 OD01044), as well as the National BioResource Project,

Ding Xue (University of Colorado, Boulder), L. Rene Garcia (Texas A&M University), and

Douglas Portman (University of Rochester Medical Center) for providing strains. We also thank

Igor Antoshechkin, Caltech Genomics facility for sequencing and InVivo Biosystems for

generating transgenic and CRISPR knockout animals. The synthetic ascr#8 utilized in this study

was provided by Frank Schroeder (Cornell University). The research reported in this publication

was supported by NIH R01 DC016058 (J.S.), R01 GM084389 (P.W.S.), the Howard Hughes

Medical Institute (P.W.S.), Moore Foundation Grant No. 4551 (E.M.S.), and Cornell startup

funding (E.M.S.). We thank Titus Brown and the Michigan State University High-Performance

Computing Center (supported by U.S. Department of Agriculture grant 2010-65205-20361 and

NIFA–National Science Foundation (NSF) grant IOS-0923812) for computational support;

additional computing was enabled by start-up and research allocations from NSF XSEDE (TG-

MCB180039 and TG-MCB190010).

Figure and Table Legends

Gene	CEM DL	CEM DR	CEM VL	CEM VR	CEM/others	CEM/males
srw-97	0.02	0.03	0.02	**0.10**	3.33	2.50
dmsr-12	**1.26**	0.03	0.02	0.05	25.2	18.0
seb-3	0.01	0.02	0.01	**0.06**	3.00	0.02
srd-32	**0.07**	0.02	0.02	0.04	1.75	0.17
srr-7	0.03	0.03	**0.17**	0.04	4.25	0.68
pkd-2	**1.36**	0.09	0.42	1.15	1.18	0.15
trf-1	0.23	0.04	0.01	**78.02**	339.22	8.72
srw-98	0.02	0.03	0.02	**0.05**	1.67	0.20
dmsr-13	0.02	0.03	0.02	**0.05**	1.67	0.45
dmsr-10	0.02	0.04	0.03	**0.07**	1.75	1.40
dmsr-11	0.01	0.02	0.02	**0.04**	2.00	0.33
dmsr-16	0.02	0.04	0.03	**0.06**	1.50	0.67

Table 1. Expression levels for genes of interest in CEM neurons. For each gene, observed

RNA-seq expression levels in each CEM neuron type are given in transcripts per million (TPM);

CEM types with the highest expression are in boldface for each gene. Two measures of gene

enrichment and specificity are also shown: first, the ratio of a gene's expression for the highest-

expressing CEM neuron type divided by that gene's expression in the highest-expressing of the

other three CEM neuron types ('CEM/others'; 'CEM_any/CEM_non_any.max' in **Table S5**); and,

the ratio of that gene's highest-expressing CEM neuron type divided by that gene's expression in

the highest-expressing of three replicates of adult *C. elegans* males ('CEM/males';

'CEM_all.max_TPM/male.max_TPM' in **Table S5**). The first ratio gives the skew for a gene's

expression towards a single CEM type; the second ratio gives the degree to which that gene's

expression cannot be explained merely by its being a male-specific gene, perhaps expressed

throughout the male body. Genes listed here are of interest either because we tested them for

transgenic GFP expression or biological function, because they have previously known CEM-

specificity, or because they are evolutionary related to *srw-97* and *dmsr-12*.

Figure 1. Transcriptomic landscapes of the CEM neurons. (A-D) Expression plots of genes

expressed in individual CEM DL **(A)**, CEM DR **(B)**, CEM VL **(C)**, and CEM VR **(D)** neurons.

For each gene, its X-axis position displays the ratio of its expression in an individual CEM neuron

divided by its highest expression in three replicates of adult *C. elegans* males; its Y-axis position

displays expression in an individual CEM neuron type. Expression levels are measured in

transcripts per million (TPM). Genes of interest are denoted by colored symbols, defined in the

legend.

Figure 2. Expression profiles of CEM-enriched GPCR genes of interest. Transcriptional GFP

fusions of GPCR genes enriched in the CEM neurons. **(A)** A previously published CEM reporter,

p*pkd-2*::GFP. **(B)** p*seb*-3::GFP, matching previously published expression, with no discernable

enrichment in the CEM. **(C)** p*dmsr-12*::GFP is strongly expressed in the CEM DL neuron, as well

as the CEM DR soma. **(D)** p*srd-32*::GFP is weakly expressed in the somas of CEM DR and DL.

(E) p*srw-97*::GFP is expressed in both CEM VR and VL, with localization in the cilia as well as

the soma. **(F)** p*srr-7*::GFP is expressed in CEM VR, both the soma and cilia, as well the cilial of

a neighboring neuron, presumably CEP VR. Dorsal/ventral axes and anterior/posterior directions

shown in (B). Scale bars denote distal, cilia region.

Figure 3. The GPCRs, SRW-97 and DMSR-12, act in non-overlapping CEM neurons to aid

in driving the ascr#8 response. (A) Raw dwell times in the centers of spatial control (S), vehicle-

only control (V), and ascaroside (A) wells, and **(B)** Log(fold-change) of A versus V for *srw-97*

knockout and transgenic rescue (WorEx57), *dmsr-12* and transgenic rescue (WorEx54), as well as

srw-97;*dmsr-12* double mutant animals. **(C)** Expression of WorEx61 (p*srw-97*::*srw-97*::GFP) in

CEM VR and CEM VL. **(D)** Expression of WorEx60 (p*dmsr-12*::*dmsr-12*::GFP) in CEM DL and

CEM DR. IL2_LV and IL2_L somas denoted. Cellular identities confirmed through co-expression

of p*klp-6*::dtTomato (**Fig. S5**). Anterior/Posterior and Right/Left axes denoted. **(E)** Raw dwell

times and **(F)** Log(fold-change) of *him-5* and *dmsr-12;srw-97* animals in response to the

structurally related attracted pheromone, ascr#3. **(G)** Structural comparison of ascr#8 (left) and

ascr#3 (right). Error bars denote SEM. $n \geq 5$. **(A, E)** RM-ANOVA followed by Dunnett's post-hoc

test or Friedman test followed by Dunn's (within strain), **(B)** ANOVA followed by Bonferroni's

post-hoc test (between strain log(fold-change) values). **(F)** Student's t-test between strain log(fold-

change) values * $p < 0.05$, ** $p < 0.01$.

Figure 4. Phylogenetic analysis of ascr#8-receptor candidate paralogs across the

***Caenorhabditis* genus. (A)** The phylogeny of *Cel_srw-97* reveals a *C. inopinata*-specific

amplification of *Cel_srw-98*. **(B)** The phylogeny of *Cel_dmsr-12* shows conserved numbers of

orthologs across the genus. Genes for *C. elegans* are denoted in purple, *C. inopinata* in pink,

C. nigoni in light blue, *C. briggsae* in dark blue, *C. japonica* in red, and *C. tropicalis* in dark green.

Distance reference bars (0.1) depict substitutions per site.

References

Aprison, E. Z., and I. Ruvinsky, 2015 Sex Pheromones of *C. elegans* Males Prime the Female
Reproductive System and Ameliorate the Effects of Heat Stress. PLoS Genet 11:
e1005729.
Aprison, E. Z., and I. Ruvinsky, 2017 Counteracting Ascarosides Act through Distinct Neurons
to Determine the Sexual Identity of *C. elegans* Pheromones. Curr Biol 27: 2589-
2599.e2583.
Artyukhin, A. B., Y. K. Zhang, A. E. Akagi, O. Panda, P. W. Sternberg *et al.*, 2018 Metabolomic
"Dark Matter" Dependent on Peroxisomal beta-Oxidation in *Caenorhabditis elegans*. J
Am Chem Soc 140: 2841-2852.
Bae, Y. K., H. Qin, K. M. Knobel, J. Hu, J. L. Rosenbaum *et al.*, 2006 General and cell-type
specific mechanisms target TRPP2/PKD-2 to cilia. Development 133: 3859-3870.
Boulin, T., J. F. Etchberger and O. Hobert, 2006 Reporter gene fusions. WormBook: 1-23.
Butcher, R. A., M. Fujita, F. C. Schroeder and J. Clardy, 2007 Small-molecule pheromones that
control dauer development in *Caenorhabditis elegans*. Nat Chem Biol 3: 420-422.
Butcher, R. A., J. R. Ragains, E. Kim and J. Clardy, 2008 A potent dauer pheromone component
in Caenorhabditis elegans that acts synergistically with other components. Proc Natl
Acad Sci U S A 105: 14288-14292.
Capella-Gutiérrez, S., J. M. Silla-Martínez and T. Gabaldón, 2009 trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics (Oxford,
England) 25: 1972-1973.
Choe, A., S. H. von Reuss, D. Kogan, R. B. Gasser, E. G. Platzer *et al.*, 2012 Ascaroside
Signaling is Widely Conserved Among Nematodes. Current Biology 22: 772-780.
Chute, C. D., E. M. DiLoreto, Y. K. Zhang, D. K. Reilly, D. Rayes *et al.*, 2019 Co-option of
neurotransmitter signaling for inter-organismal communication in *C. elegans*. Nat
Commun 10: 3186.
Cochella, L., B. Tursun, Y. W. Hsieh, S. Galindo, R. J. Johnston *et al.*, 2014 Two distinct types
of neuronal asymmetries are controlled by the *Caenorhabditis elegans* zinc finger
transcription factor die-1. Genes Dev 28: 34-43.
Dong, C., F. Dolke and S. H. von Reuss, 2016 Selective MS screening reveals a sex pheromone
in Caenorhabditis briggsae and species-specificity in indole ascaroside signalling. Org
Biomol Chem 14: 7217-7225.
Dong, C., D. K. Reilly, C. Bergame, F. Dolke, J. Srinivasan *et al.*, 2018 Comparative Ascaroside
Profiling of *Caenorhabditis* Exometabolomes Reveals Species-Specific (ω) and (ω - 2)-
Hydroxylation Downstream of Peroxisomal beta-Oxidation. J Org Chem.
Eddy, S. R., 2009 A new generation of homology search tools based on probabilistic inference.
Genome informatics. International Conference on Genome Informatics 23: 205-211.
Emms, D. M., and S. Kelly, 2015 OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome Biology 16:
157.
Emms, D. M., and S. Kelly, 2019 OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biology 20: 238.
Finn, R. D., P. Coggill, R. Y. Eberhardt, S. R. Eddy, J. Mistry *et al.*, 2016 The Pfam protein
families database: towards a more sustainable future. Nucleic acids research 44: D279-
D285.

Fraser, A. G., R. S. Kamath, P. Zipperlen, M. Martinez-Campos, M. Sohrmann *et al.*, 2000
Functional genomic analysis of *C. elegans* chromosome I by systematic RNA
interference. Nature 408**:** 325-330.
Fuxman Bass, J. I., A. M. Tamburino, A. Mori, N. Beittel, M. T. Weirauch *et al.*, 2014
Transcription factor binding to Caenorhabditis elegans first introns reveals lack of
redundancy with gene promoters. Nucleic Acids Res 42**:** 153-162.
Goodman, M. B., D. H. Hall, L. Avery and S. R. Lockery, 1998 Active currents regulate
sensitivity and dynamic range in *C. elegans* neurons. Neuron 20**:** 763-772.
Greene, J. S., M. Brown, M. Dobosiewicz, I. G. Ishida, E. Z. Macosko *et al.*, 2016b Balancing
selection shapes density-dependent foraging behaviour. Nature 539**:** 254-258.
Greene, J. S., M. Dobosiewicz, R. A. Butcher, P. T. McGrath and C. I. Bargmann, 2016a
Regulatory changes in two chemoreceptor genes contribute to a *Caenorhabditis elegans*
QTL for foraging behavior. Elife 5.
Hobert, O., 2013 The neuronal genome of *Caenorhabditis elegans*. WormBook**:** 1-106.
Houck, L., C. Palmer, R. Watts, S. Arnold, P. Feldhoff *et al.*, 2007 A new vertebrate courtship
pheromone, PMF, affects female receptivity in a terrestrial salamander. Animal
Behaviour 73**:** 315-320.
Howe, K. L., B. J. Bolt, S. Cain, J. Chan, W. J. Chen *et al.*, 2016 WormBase 2016: expanding to
enable helminth genomic research. Nucleic acids research 44**:** D774-D780.
Jang, Y. H., H. S. Chae and Y. J. Kim, 2017 Female-specific myoinhibitory peptide neurons
regulate mating receptivity in *Drosophila melanogaster*. Nat Commun 8**:** 1630.
Jee, C., J. Lee, J. P. Lim, D. Parry, R. O. Messing *et al.*, 2013 SEB-3, a CRF receptor-like
GPCR, regulates locomotor activity states, stress responses, and ethanol tolerance in *C.*
elegans. Genes, brain, and behavior 12**:** 10.1111/j.1601-1183X.2012.00829.x.
Kaletsky, R., V. Lakhina, R. Arey, A. Williams, J. Landis *et al.*, 2016 The C. elegans adult
neuronal IIS/FOXO transcriptome reveals adult phenotype regulators. Nature 529**:** 92-96.
Käll, L., A. Krogh and E. L. L. Sonnhammer, 2004 A combined transmembrane topology and
signal peptide prediction method. Journal of molecular biology 338**:** 1027-1036.
Kalyaanamoorthy, S., B. Q. Minh, T. K. F. Wong, A. von Haeseler and L. S. Jermiin, 2017
ModelFinder: fast model selection for accurate phylogenetic estimates. Nature methods
14**:** 587-589.
Kamath, R. S., A. G. Fraser, Y. Dong, G. Poulin, R. Durbin *et al.*, 2003 Systematic functional
analysis of the *Caenorhabditis elegans* genome using RNAi. Nature 421**:** 231-237.
Katoh, K., and D. M. Standley, 2013 MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Molecular biology and evolution 30**:** 772-
780.
Kim, K., K. Sato, M. Shibuya, D. M. Zeiger, R. A. Butcher *et al.*, 2009 Two Chemoreceptors
Mediate Developmental Effects of Dauer Pheromone in *C. elegans*. Science 326**:** 994-
998.
Krishnan, A., M. S. Almén, R. Fredriksson and H. B. Schiöth, 2014 Insights into the Origin of
Nematode Chemosensory GPCRs: Putative Orthologs of the Srw Family Are Found
across Several Phyla of Protostomes. PLoS ONE 9**:** e93048.
Langmead, B., and S. L. Salzberg, 2012 Fast gapped-read alignment with Bowtie 2. Nature
methods 9**:** 357-359.

Lemoine, F., J. B. Domelevo Entfellner, E. Wilkinson, D. Correia, M. Dávila Felipe *et al.*, 2018
Renewing Felsenstein's phylogenetic bootstrap in the era of big data. Nature 556**:** 452-
456.
Li, B., and C. N. Dewey, 2011 RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. BMC bioinformatics 12**:** 323-323.
Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence
Alignment/Map format and SAMtools. Bioinformatics (Oxford, England) 25**:** 2078-2079.
Ludewig, A. H., A. B. Artyukhin, E. Z. Aprison, P. R. Rodrigues, D. C. Pulido *et al.*, 2019 An
excreted small molecule promotes *C. elegans* reproductive development and aging. Nat
Chem Biol 15**:** 838-845.
Lupas, A., 1996 Prediction and analysis of coiled-coil structures. Methods in enzymology 266**:**
513-525.
Madeira, F., Y. M. Park, J. Lee, N. Buso, T. Gur *et al.*, 2019 The EMBL-EBI search and
sequence analysis tools APIs in 2019. Nucleic acids research 47**:** W636-W641.
McGrath, P. T., and I. Ruvinsky, 2019 A primer on pheromone signaling in *Caenorhabditis*
elegans for systems biologists. Curr Opin Syst Biol 13**:** 23-30.
McGrath, P. T., Y. Xu, M. Ailion, J. L. Garrison, R. A. Butcher *et al.*, 2011 Parallel evolution of
domesticated *Caenorhabditis* species targets pheromone receptor genes. Nature 477**:** 321-
325.
Narayan, A., G. Laurent and P. W. Sternberg, 2011 Transfer characteristics of a thermosensory
synapse in *Caenorhabditis elegans*. Proceedings of the National Academy of Sciences of
the United States of America 108**:** 9667-9672.
Narayan, A., V. Venkatachalam, O. Durak, D. K. Reilly, N. Bose *et al.*, 2016 Contrasting
responses within a single neuron class enable sex-specific attraction in *Caenorhabditis*
elegans. Proc Natl Acad Sci U S A 113**:** E1392-1401.
Nguyen, L.-T., H. A. Schmidt, A. von Haeseler and B. Q. Minh, 2015 IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies.
Molecular biology and evolution 32**:** 268-274.
Noble, L. M., J. Yuen, L. Stevens, N. Moya, R. Persaud *et al.*, 2021 Selfing is the safest sex for
Caenorhabditis tropicalis. Elife 10.
Park, D., I. O'Doherty, R. K. Somvanshi, A. Bethke, F. C. Schroeder *et al.*, 2012 Interaction of
structure-specific and promiscuous G-protein-coupled receptors mediates small-molecule
signaling in *Caenorhabditis elegans*. PNAS 109**:** 9917-9922.
Peden, E. M., and M. M. Barr, 2005 The KLP-6 kinesin is required for male mating behaviors
and polycystin localization in *Caenorhabditis elegans*. Curr Biol 15.
Poole, R. J., E. Bashllari, L. Cochella, E. B. Flowers and O. Hobert, 2011 A Genome-Wide
RNAi Screen for Factors Involved in Neuronal Specification in *Caenorhabditis elegans*.
PLoS Genet 7**:** e1002109.
Powell, S., D. Szklarczyk, K. Trachana, A. Roth, M. Kuhn *et al.*, 2012 eggNOG v3.0:
orthologous groups covering 1133 organisms at 41 different taxonomic ranges. Nucleic
acids research 40**:** D284-D289.
Pungaliya, C., J. Srinivasan, B. W. Fox, R. U. Malik, A. H. Ludewig *et al.*, 2009 A shortcut to
identifying small molecule signals that regulate behavior and development in
Caenorhabditis elegans. Proceedings of the National Academy of Sciences of the United
States of America 106**:** 7708-7713.

Ragsdale, E. J., M. R. Muller, C. Rodelsperger and R. J. Sommer, 2013 A developmental switch
coupled to the evolution of plasticity acts through a sulfatase. Cell 155: 922-933.
Reilly, D. K., D. E. Lawler, D. R. Albrecht and J. Srinivasan, 2017 Using an Adapted
Microfluidic Olfactory Chip for the Imaging of Neuronal Activity in Response to
Pheromones in Male *C. Elegans* Head Neurons. Journal of Visualized Experiments:
e56026.
Reilly, D. K., E. J. McGlame, E. Vandewyer, A. N. Robidoux, C. S. Muirhead *et al.*, 2021
Distinct neuropeptide-receptor modules regulate a sex-specific behavioral response to a
pheromone. Communications Biology 4: 1018.
Reilly, D. K., L. J. Randle and J. Srinivasan, 2019 Evolution of hermaphroditism decreases
efficacy of Ascaroside#8-mediated mate attraction in *Caenorhabditis* nematodes.
microPublication Biology.
Robertson, H. M., and J. H. Thomas, 2006 The putative chemoreceptor families of *C. elegans*.
WormBook: 1-12.
Rual, J. F., J. Ceron, J. Koreth, T. Hao, A. S. Nicot *et al.*, 2004 Toward improving
Caenorhabditis elegans phenome mapping with an ORFeome-based RNAi library.
Genome Res 14: 2162-2168.
Sawin, E. R., R. Ranganathan and H. R. Horvitz, 2000 C. elegans locomotory rate is modulated
by the environment through a dopaminergic pathway and by experience through a
serotonergic pathway. Neuron 26: 619-631.
Schmitz, C., P. Kinge and H. Hutter, 2007 Axon guidance genes identified in a large-scale RNAi
screen using the RNAi-hypersensitive *Caenorhabditis elegans* strain *nre-1(hd20) lin-*
15b(hd126). Proceedings of the National Academy of Sciences 104: 834-839.
Schwarz, E. M., M. Kato and P. W. Sternberg, 2012 Functional transcriptomics of a migrating
cell in Caenorhabditis elegans. Proceedings of the National Academy of Sciences of the
United States of America 109: 16246-16251.
Taylor, S. R., G. Santpere, A. Weinreb, A. Barrett, M. B. Reilly *et al.*, 2021 Molecular
topography of an entire nervous system. Cell 184: 4329-4347.e4323.
Tenor, J. L., and A. Aballay, 2008 A conserved Toll-like receptor is required for Caenorhabditis
elegans innate immunity. EMBO Rep 9: 103-109.
Thomas, C. G., R. Li, H. E. Smith, G. C. Woodruff, B. Oliver *et al.*, 2012 Simplification and
desexualization of gene expression in self-fertile nematodes. Curr Biol 22: 2167-2172.
von Reuss, S. H., N. Bose, J. Srinivasan, J. J. Yim, J. C. Judkins *et al.*, 2012 Comparative
Metabolomics Reveals Biogenesis of Ascarosides, a Modular Library of Small-Molecule
Signals in *C. elegans*. Journal of the American Chemical Society 134: 1817-1824.
von Reuss, S. H., and F. C. Schroeder, 2015 Combinatorial chemistry in nematodes: modular
assembly of primary metabolism-derived building blocks. Natural product reports 32:
994-1006.
Wang, J., R. Kaletsky, M. Silva, A. Williams, L. A. Haas *et al.*, 2015 Cell-Specific
Transcriptional Profiling of Ciliated Sensory Neurons Reveals Regulators of Behavior
and Extracellular Vesicle Biogenesis. Curr Biol 25: 3232-3238.
Wootton, J. C., 1994 Non-globular domains in protein sequences: automated segmentation using
complexity measures. Computers & chemistry 18: 269-285.
Zhang, Y. K., D. K. Reilly, J. Yu, J. Srinivasan and F. C. Schroeder, 2019 Photoaffinity probes
for nematode pheromone receptor identification. Journal of Organic & Biomolecular
Chemistry: 10.1039/C1039OB02099C.

Zhang, Y. K., M. A. Sanchez-Ayala, P. W. Sternberg, J. Srinivasan and F. C. Schroeder, 2017
Improved Synthesis for Modular Ascarosides Uncovers Biological Activity. Org Lett 19**:**
2837-2840.
Zwaal, R. R., J. E. Mendel, P. W. Sternberg and R. H. Plasterk, 1997 Two neuronal G proteins
are involved in chemosensation of the *Caenorhabditis* elegans Dauer-inducing
pheromone. Genetics 145**:** 715-727.

Figure - 1

Figure - 2

Figure - 3

Figure - 4