Transcriptomic profiling of sex-specific olfactory neurons reveals subset-specific receptor

expression in *C. elegans*

Author Names and Affiliations:

Douglas K. Reilly[1,7,†], Erich M. Schwarz[2,†], Caroline S. Muirhead[1], Annalise N. Robidoux[1,3], Igor

Antoshechkin[4], Anusha Narayan[4,5], Meenakshi K. Doma[4,6], Paul W. Sternberg[4], Jagan

Srinivasan[1,8,*]

[1] Department of Biology and Biotechnology, Worcester Polytechnic Institute, Worcester, MA

01605, USA

[2] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853, USA

[3] Department of Chemistry and Biochemistry, Worcester Polytechnic Institute, Worcester, MA

01605, USA

[4] Division of Biology and Biological Engineering, California Institute of Technology, Pasadena,

CA 91125

[5] Present address: Strategic Marketing, Evidation Health, San Mateo, CA 94402, USA

[6] Present address: Genomics Lead, Individualized Cell and Gene Therapies, South San Francisco,

Genentech, CA 94080, USA

[7] Present address: Biology Department, Tufts University, Medford, MA 02155, USA

[8] Program of Bioinformatics and Computational Biology, Worcester Polytechnic Institute,

Worcester, MA 01605, USA

* Corresponding author

† These authors contributed equally to this work

Short Title: Male pheromone G protein-coupled receptors in *C. elegans*

Key words: *C. elegans*, Pheromone, Sex-specific, G Protein-Coupled Receptors, Transcriptomes,

Single-Cell

Corresponding author:

Jagan Srinivasan

Life Sciences and Bioengineering Center, 4th Floor

60 Prescott Street, Worcester, MA, 01605

(508) 831-6564

jsrinivasan@wpi.edu

SUMMARY

The nematode *Caenorhabditis elegans* utilizes chemosensation to navigate an ever-changing

environment for its survival. A class of secreted small-molecule pheromones, termed ascarosides,

play an important role in olfactory perception by affecting a host of biological function ranging

from development to behavior. The ascaroside ascr#8 mediates sex-specific behaviors, driving

avoidance in hermaphrodites and attraction in males. Males sense ascr#8 via the ciliated male-

specific cephalic sensory (CEM) neurons, which exhibit radial symmetry along dorsal-ventral and

left-right axes. Calcium imaging studies suggest a complex neural coding mechanism that

translates stochastic physiological responses in these neurons to reliable behavioral outputs. To

test the hypothesis that the neurophysiological complexity arises from differential expression of

genes within subsets of these neurons, we performed cell-specific transcriptomic profiling of these

sensory neurons. Expression profiling revealed between 20 and 639 genes enriched at least two-

fold per CEM neuron and identified multiple G protein coupled receptor (GPCR) candidates

enriched in non-overlapping subsets of CEM neurons. GFP reporter analysis confirmed that RNA

expression of two of the GPCR genes, *srw-97* and *dmsr-12*, is enriched in specific subsets of the

CEM neurons. Single CRISPR-Cas9 knockouts of either *srw-97* or *dmsr-12* resulted in partial

defects, while a double knockout of both *srw-97* and *dmsr-12* completely abolished the attractive

response to ascr#8, suggesting that each receptor acts in a non-redundant manner in discrete

olfactory neurons. Together, our results suggest that the evolutionarily distinct GPCRs SRW-97

and DMSR-12 act to facilitate male-specific sensation of ascr#8 through discrete subsets of CEM

neurons.

INTRODUCTION

The ability of an organism to find a mate is critical to the survival of a species. Many species utilize

small molecule pheromones to signal mate location (PUNGALIYA *et al.* 2009; NARAYAN *et al.*

2016), sexual maturity (APRISON AND RUVINSKY 2015; APRISON AND RUVINSKY 2017), and

receptivity (HOUCK *et al.* 2007; JANG *et al.* 2017). These signals are sensed and processed through

the nervous system to generate proper behavioral and developmental responses. The nematode

Caenorhabditis elegans communicates with conspecifics almost exclusively using pheromones

called ascarosides (LUDEWIG *et al.* 2019; MCGRATH AND RUVINSKY 2019).

Ascarosides are a large, structurally conserved class of small molecule pheromones (VON

REUSS AND SCHROEDER 2015; ZHANG *et al.* 2017). Built in a modular fashion, they are composed

of a core ascarylose sugar and a fatty-acid derived side chain (BUTCHER *et al.* 2007; VON REUSS *et*

al. 2012; LUDEWIG *et al.* 2019). Ascarosides signal a host of environmental and developmental

information, including the sexual maturity and location of potential mates (NARAYAN *et al.* 2016;

APRISON AND RUVINSKY 2017). This communication system is conserved across nematodes, with

different species using different combinations of ascarosides or structural derivatives (CHOE *et al.*

2012; RAGSDALE *et al.* 2013; DONG *et al.* 2016; DONG *et al.* 2018; REILLY *et al.* 2019). While the

effects of some ascarosides are dependent on physiological state (BUTCHER *et al.* 2009; BUTCHER

2017; CHUTE *et al.* 2019), others are able to elicit sex-specific responses (PUNGALIYA *et al.* 2009;

NARAYAN *et al.* 2016; FAGAN *et al.* 2018).

Most neurons in the *C. elegans* nervous system are bilaterally symmetrical (WHITE *et al.*

1986), though radially symmetric classes of sensory neurons exist, such as the inner labial (IL1

and IL2) neurons (HRUS *et al.* 2007; WANG *et al.* 2014; WANG *et al.* 2015) which are required for

normal dauer development, nictation behavior, and foraging (LEE *et al.* 2012; SCHROEDER *et al.*

2013). *C. elegans* also has several sex-specific neurons, such as the HSN neuron (hermaphrodite-

specific neuron) involved in egg laying (APIRSON AND RUVINSKY 2019a; APIRSON AND RUVINSKY

2019b), and the MCM neurons in the male that contribute to neuronal plasticity and learning

(SAMMUT *et al*. 2015). However, while most male-specific neurons are located in the tail, the major

contributor to male-specific behaviors are the chemosensory CEM neurons, located in the amphid.

These four radially symmetric neurons have been found, through laser ablation,

electrophysiology, and calcium imaging studies to be involved in sensing the ascaroside

pheromones ascr#8 and ascr#3 (PUNGALIYA *et al*. 2009; NARAYAN *et al*. 2016; REILLY *et al*. 2017).

Ascr#3 plays a role in dauer formation, functioning alongside ascr#2 and ascr#4 as the "dauer

pheromone" (BUTCHER *et al*. 2008). Ascr#8, meanwhile, is unique in that it contains a *p*-

aminobenzoic acid moiety on its terminus (PUNGALIYA *et al*. 2009; ARTYUKHIN *et al*. 2018). We

have previously shown by both laser and genetic ablation experiments that ascr#8 is sensed

primarily by male-specific CEM neurons located in the head region of the male nervous system

(NARAYAN *et al*. 2016). In animals lacking CEM (through either genetic or laser ablation), the

response to ascr#8 is abolished, and further ablation of the ASK results in complete loss of the

ascr#3 response (NARAYAN *et al*. 2016). Through studies ablating three of the four CEM neurons,

leaving one CEM neuron intact, it was found that four CEM neurons cooperate to drive a tuned

response to an intermediate, 1 µM concentration of both ascr#3 and ascr#8 (NARAYAN *et al*. 2016).

Although CEM neurons are indeed the primary pathway by which male *C. elegans* sense

and response to ascr#8, this response is modulated by additional neural circuitry via neuropeptides.

The FMRFamide-like peptide gene *flp-3* encodes ten discrete peptides, some of which act through

two neuropeptide receptors both to repress *C. elegans*' normal avoidance of ascr#8, and to

simultaneously make *C. elegans* positively attracted to ascr#8 (REILLY *et al.* 2021). However, it

remains unknown how CEM neurons themselves sense and drive responses to ascr#8.

Calcium imaging and electrophysiological experiments demonstrated that CEM neurons

show variable responses to stimuli not only between different animals, but within a single animal

(NARAYAN *et al.* 2016; REILLY *et al.* 2017). While other neurons such as the stochastically

asymmetric AWC chemosensory neurons show variable calcium responses between different

animals, these responses are consistent between the left and right AWC neurons of a single animal

(COCHELLA *et al.* 2014). In contrast, we have observed that responses of individual CEM neurons

are stochastic within single animals, and yet four CEM neurons within one animal are somehow

consistently able to generate proper behavioral responses to both ascr#3 and ascr#8 (NARAYAN *et*

al. 2016). To understand this remarkable pattern of stochastic neuronal activity yielding consistent

neuronal outputs, it is imperative to uncover genes encoding components of the CEM response.

A recent study in *C. elegans* hermaphrodites generated transcriptomic landscapes of 118

neuronal classes from 302 neurons in order to link functional and anatomical properties of

individual neurons with their molecular identities (TAYLOR *et al.* 2021). Discrete neuronal classes

were successfully identified via their combinations of expressed neuropeptides and neuropeptide

receptor genes. However, a similar feat is yet to be performed on male *C. elegans*, and the

transcriptomic profiles of male neurons remain enigmatic.

In a more focused transcriptomic approach, gene expression profiles of extracellular

vesicle-releasing neurons (EVNs) as a whole were determined by expressing GFP under the EVN-

specific *klp-6* promoter, mechanically and proteolytically dissociating male worms expressing this

GFP, selecting the target GFP+ neurons with FACS, and performing RNA-seq on the collected

GFP+ neurons (WANG *et al.* 2015; KALETSKY *et al.* 2016). This gave thousands of EVNs in each

biological replicate and offered increased statistical power in analyzing transcriptional landscapes.

CEM neurons are a subset of EVN neurons, so CEM transcriptomic data are embedded in these

data, but the expression values of CEM and EVN are not equivalent.

To begin understanding how CEM neurons achieve stochastic yet reliable physiological

responses to ascr#8, we performed single-cell RNA-seq (SCHWARZ *et al.* 2012) on male-specific

CEM neurons. We uncovered a small number of genes encoding G protein-coupled receptors

(GPCRs) highly enriched in single CEM neurons. Given that all ascaroside receptors identified to

date have been GPCRs (KIM *et al.* 2009; MCGRATH *et al.* 2011; PARK *et al.* 2012; GREENE *et al.*

2016a; GREENE *et al.* 2016b; CHUTE *et al.* 2019), we tested whether any of these enriched genes

contribute to male *C. elegans* sensation of ascr#8. Through neuronal RNAi knockdown and

CRISPR null mutagenesis, we identified two distantly related GPCR genes that contribute to the

ascr#8 behavioral response, *srw-97* and *dmsr-12*; expression of these genes are enriched in ventral

and dorsal CEMs, respectively. When expression of each receptor was knocked down through

RNAi, behavioral responses to ascr#8 were partially deficient. Using CRISPR and genetic crosses

to generate a double *srw-97 dmsr-12* mutant, we found that loss of both receptors results in a

complete loss of behavioral response to ascr#8. Phylogenetic analysis further indicates that both

receptors are homologous to closely related receptors across the *Caenorhabditis* genus, and robust

ascr#8 responses may be a trait recently evolved in *C. elegans*.

141   **RESULTS**

The transcriptomic landscape of CEM neurons is variable

Individual CEM neurons were isolated from *C. elegans* expressing an integrated GFP

labeling extracellular vesicle-releasing neurons (EVNs, p*pkd-2*::GFP [**Figure 2A**]), as previously

described (GOODMAN *et al.* 1998; NARAYAN *et al.* 2011; NARAYAN *et al.* 2016). Cells were

separated by anatomical identity (i.e., CEM dorsal left [DL], dorsal right [DR], ventral left [VL],

and ventral right [VR]), and cDNA libraries were constructed.

Enriched genes in each CEM neuron were identified by comparing RNA-seq profiles of

distinct CEM types; genes expressed at least two times more strongly in one CEM type than others

were listed and checked for biological functions, as annotated by their WormBase Gene Ontology

(GO) terms (HARRIS *et al.* 2020; GENE ONTOLOGY 2021). A variety of enriched gene sets were

defined for different CEM neurons, ranging from 20 genes enriched in CEM VL to over 600 in

CEM VR (**Table S1, S6, S7**). Dorsal CEM neurons were more consistent, expressing 98 and 105

enriched genes in the right and left neurons, respectively (**Table S1, S8, S9**). Uniquely mapped

reads ranged from 1.9 to 3.8 million per CEM type, matching the trend for alignment rates of each

CEM neuron, which ranged from 19.73% to 48.01%, (**Table S2**) with an average of 1.410 genes

showing robust RNA expression in each neuron (**Table S3**).

Five genes encoding G protein-coupled receptors (*seb-3, srr-7, srw-97, dmsr-12, srd-32*)

were expressed at levels four times higher in one CEM neuron than in other CEM neurons. Of

these, four were uncharacterized; one (*seb-3*) has previously been shown to play roles in

locomotion, stress response, and ethanol tolerance (JEE *et al.* 2013). *dmsr-12* is related to *daf-37*

(ROBERTSON AND THOMAS 2006), a previously identified ascaroside receptor gene (PARK *et al.*

2012), although it is more closely related to the neuropeptide receptor gene *dmsr-1*, and more

distantly to *srw-97* (ROBERTSON AND THOMAS 2006). *srd-32* belongs to a divergent branch of the

SRD phylogeny, which is itself a divergent family of the STR superfamily (ROBERTSON AND

THOMAS 2006). *srr-7* belongs to the one of the smallest families of *C. elegans* chemoreceptor

genes, outside of the single opsin family member, *sro-1*, and the *srm* family encoding five

chemoreceptor genes (ROBERTSON AND THOMAS 2006)).

seb-3 and *srw-97* exhibited similar enrichment profiles across the CEM neurons, with ~3.5-

fold enrichment in CEM VR (**Table 1; Figure 1D; Table S7**). *dmsr-12* showed almost five-fold

stronger expression in CEM DL versus other CEM neurons (**Table 1; Figure 1A; Table S8**), while

srd-32 was only 1.62-fold enriched (**Table 1; Figure 1A; Table S8**). *srr-7* showed only two-fold

enrichment in CEM VL (**Table 1; Figure 1C; Table S6**), which correlated with previous

transcriptomic analyses that found *srr-7* to be enriched in *C. elegans* EVNs (WANG *et al.* 2015).

CEM-specific receptor expression patterns

To confirm our single-cell RNA-seq results, we generated transgenic GFP fusions for the

five receptor genes (BOULIN *et al.* 2006). Roughly 3 kb of promoter region upstream of the start

codon was included in these constructs, along with the majority of the coding sequence (**Table**

S10); this would have automatically included any large 5'-ward introns that might contain cis-

regulatory elements of these genes (FUXMAN BASS *et al.* 2014). The GFP coding sequence was

cloned from the Fire Kit vector pPD95.75 (BOULIN *et al.* 2006). *pha-1; lite-1; him-5* animals were

injected with reporter constructs and the co-injection marker pBX (*pha-1*(+)).We isolated GFP$^+$

strains and imaged and GFP$^+$ males for expression at 63x magnification (**Figure 2; Fig. S1**). An

integrated p*pkd-2*::GFP line was used as a CEM-specific control (**Figure 2A**).

The previously characterized GPCR gene, *seb-3*, displayed a non-CEM-specific expression

pattern matching that previously described (**Figure 2B**) (JEE *et al.* 2013), and was excluded from

further analyses. The other four receptor genes showed transgenic expression patterns similar to

their RNA-seq data: GFP-tagged DMSR-12 was heavily enriched in CEM DL, but also exhibited

CEM DR expression, and was observed in both soma and cilia (**Figure 2C**). GFP-tagged SRD-32

was found faintly in both dorsal CEM neurons but exhibited no ciliary localization (**Figure 2D**).

GFP-tagged SRW-97 was found in both ventral CEM soma and cilia, with slightly higher

expression in CEM VL (**Figure 2E**). GFP-tagged SRR-7 was found in CEM VR, as well as another

neuron that is likely to be CEP VR (**Figure 2F**). As with *seb-3*, *srr-7* was excluded from further

analyses, as we aimed to identify CEM-specific regulators of the ascr#8 response.

Except for SRD-32, all GFP-tagged receptors showed subcellular localization patterns that

included the sensory cilia (**Figure 2, white bars**). Our failure to see this for SRD-32 may be an

artifact of our p*srd-32*::GFP construct design, because only 52% of the *srd-32* coding sequence

was included in our transgene (**Table S10**).

We also observed non-GPCR genes to be heavily enriched in single CEM neurons, such as

trf-1, which encodes a TNF receptor homolog (TENOR AND ABALLAY 2008); *trf-1* has an EVN-

specific promoter (WANG *et al.* 2015), which provides an alternative to the *pkd-2* and *klp-6*

promoters typically used to drive transgenes in EVNs (**Fig. S1**) (PEDEN AND BARR 2005; BAE *et*

al. 2006). This fits previous observations of *trf-1*::GFP expression in CEM, HOB, and RnB

neurons (WANG *et al.* 2015).

RNAi-mediated knockdown of CEM Receptors

To test whether these CEM-enriched genes encoded receptors that are required in ascr#8

sensation, we used RNAi in a strain that is hypersensitive to neuronal RNA interference,

nre-1; lin-15B (SCHMITZ *et al.* 2007; POOLE *et al.* 2011). We crossed *nre-1; lin-15B* from this

strain into a *him-5* background so that we could assay ascr#8 responses in males. Young adults

were then reared on NGM agar plates containing 1 mM IPTG to induce expression of dsRNA

within the food source which would cause receptor-specific RNAi. Young adult males of the

subsequent generation were then assayed for their response to ascr#8 after being grown on either

a control vector (pL4440) or targeted dsRNA vector.

To confirm that our system for male neuronal RNAi could affect behavioral phenotypes,

we first fed animals M02B7.3 (α-*osm-3*) and B0212.5 (α-*osm-9*) RNAi clones from the Ahringer

Library (FRASER *et al.* 2000; KAMATH *et al.* 2003). Using our single-worm Spot Retention Assay

(NARAYAN *et al.* 2016), we assayed animals subjected to RNAi against *osm-3* or *osm-9* for their

behavioral dwell time in ascr#8 (**Figure 3A**). Animals fed *osm-3* dsRNA showed significantly

defective responses to ascr#8, like that observed loss-of-function alleles (**Figure 3B**). In contrast,

animals fed *osm-9* showed only a slight decrease in their times spent within ascr#8 (**Figure 3A**).

Because we could successfully abolish ascr#8 attraction through male neuronal RNAi of *osm-3*,

we could be confident that RNAi would be broadly effective at functionally verifying CEM-

enriched genes encoding components of the ascr#8 response.

We thus fed animals dsRNA clones targeting three CEM-enriched receptor genes:

T18H12.5 (α-*srd-32*), H34P18.1 (α-*dmsr-12*), and ZC204.15 (α-*srw-97*). The α-*srd-32* clone was

the only one not available in the Ahringer library (FRASER *et al.* 2000; KAMATH *et al.* 2003), but

a clone from the Vidal library was available (RUAL *et al.* 2004). These clones were in the same

backbone vector, allowing for the same control vector to be utilized across studies.

RNAi of *srd-32* caused no defect in ascr#8 responses (**Figure 3C**). Either *dmsr-12* or *srw-*

97 dsRNA caused partial defects of ascr#8 dwell time (**Figure 3C**). Neither knockdown

statistically lowered the dwell time in ascr#8 (**Figure 3D**), though they did abolish the statistically

significant increase in time spent in ascr#8 over vehicle controls (**Figure 3C**).

CRIPSR-generated null mutants of candidate receptors

We used CRISPR to generate a null allele of *srw-97(knu456)* via InVivo Biosystems and

backcrossed it into a *him-5* background for analysis of male attraction to ascr#8. For phenotypic

analysis of null mutants, we used our Single Worm Attraction Assay (SWAA) (REILLY *et al.*

2021). In this assay, animals are placed in a well of a 48-well tissue culture plate containing NGM

coated in *E. coli* OP50, and either nothing (S), a drop of the vehicle control (V), or the attractive

ascaroside (A). The location of animals is recorded over 15 minutes, and the time spent in the

center of the well, or cue, is compared (REILLY *et al.* 2021).

Wild-type (*him-5*) animals were strongly attracted to ascr#8 in the SWAA, replicating our

previous observations (**Figure 4A, B; Fig. S2**) (REILLY *et al.* 2021). *srw-97(knu456)* males

displayed partially defective ascr#8 attraction: their ascr#8 dwell time was no different than that

of vehicle (**Figure 4B; Fig. S2**), while their increase over the vehicle was no different than that of

him-5 (**Figure 4A; Fig. S2**).

We were able to restore normal ascr#8 attraction to *srw-97(knu456)* males through

transgenesis with a translational fusion construct (p*srw-97*::*srw-97*::GFP) that employed the same

promoter as our transcriptional fusion (**Figure 2E**). Expression of the rescue transgene matched

that of our initial fusion, with GFP visible in both ventral CEM neurons (**Figure 4C**). Single-worm

assays revealed that the *srw-97* rescue construct completely rescued the partial defect in the ascr#8

response (**Figure 4A, B; Fig. S2**).

We obtained a *dmsr-12(tm8706)* null allele from the National BioResource Program in

Japan and crossed it into a *him-5* background as well. Like *srw-97(knu456)*, *dmsr-12(tm8706)*

males exhibited partially defective responses to ascr#8 (**Figure 5A, B; Fig. S3**). Similarly, the

expression profile of the p*dmsr-12*::*dmsr-12*::GFP rescue construct matched its earlier reporter

expression (**Figure 2C, 5C**). This rescue construct also completely restored normal attraction to

ascr#8 (**Figure 5A, B; Fig. S3**).

Given that *dmsr-12* is expressed in dorsal CEM neurons, and *srw-97* is expressed in ventral

CEM neurons (**Figure 2C, E**), we speculated that single mutants of either *dmsr-12* or *srw-97*

partially retained the ability to sense and respond to ascr#8, with the opposing wild-type CEM-

specific receptor gene conferring some residual response to ascr#8. We thus generated a *dmsr-*

12; srw-97 double mutant strain and assayed animals for their ability to respond to ascr#8. These

double mutants completely failed to be attracted to or retained by ascr#8 activity (**Figure 5; Fig.**

S3). This defect, though strong, was specific to ascr#8 sensation; double mutants showed no defect

in their responses to ascr#3 (**Figure 5D, E; Fig. S4**).

Together, these data suggest that at least two GPCR receptors are expressed in opposite

pairs of CEM neurons and act in parallel during sensation of ascr#8.

Phylogenetic analyses of ascr#8 receptors reveal likely gene duplication events

The ability to attract mates through pheromones is often essential for a species' survival.

However, *C. elegans* is a self-fertile hermaphrodite: male mating is useful in creating genetic

diversity but is not absolutely required for species propagation. We have recently observed that

different *Caenorhabditis* species show quite different responses to ascr#8 (REILLY *et al.* 2019).

To understand this pattern better, we analyzed the evolution of both *srw-97* and *dmsr-12*.

We identified orthologs of *srw-97* from *C. elegans* (*Cel-srw-97*) in other *Caenorhabditis*

proteoms via OrthoFinder (**Figure 6A**). A closely related *C. elegans* paralog, *Cel-srw-98*,

underwent a species-specific expansion within *C. inopinata*. However, other species, such as

C. briggsae and *C. nigoni*, have closely related paralogs to *Cel-srw-97*. The coding DNA

sequences (CDS DNAs) of both *Cel-srw-97* and *Cel-srw-98* are 66.2% identical to one another,

while their amino acid sequences are only 57.1% identical (MADEIRA *et al.* 2019). In our single-

cell CEM RNA-seq data, *Cel-srw-98* shows enrichment similar to that of to *Cel-srw-97*; it is most

strongly expressed in CEM VR, although below the two-fold enrichment cutoff that we chose in

selecting candidates for functional analysis (**Table 1**).

C. tropicalis, like *C. elegans* and *C. briggsae*, is a hermaphroditic *Caenorhabditis* species

for which male mating is optional (NOBLE *et al.* 2021). *C. tropicalis* encodes a reduced number of

ascr#8 receptor gene paralogs, with no *Cel-dmsr-12* paralogs, and only one *Cel-srw-97* paralog

(**Figure 6**). We previously observed that *C. tropicalis* fails to be attracted to ascr#8, and it is

possible that this loss of receptor genes is one reason for that failure (REILLY *et al.* 2019). Given

the conserved activity of cis-regulatory elements between different *Caenorhabditis* species (WANG

et al. 2004; GORDON *et al.* 2015), it would be interesting to express a *Cel-srw-97* construct in *C.*

tropicalis to see if it can confer attraction to and retention by ascr#8.

In contrast, the other candidate ascr#8 receptor, *dmsr-12*, does not appear in our

phylogenetic analysis to have undergone any species-specific expansion (**Figure 6B**). In fact,

outside of *C. elegans*, there are generally fewer paralogs per species. The most closely related *C.*

elegans gene in our phylogeny, *Cel-dmsr-13*, does not show CEM enrichment in our single-neuron

RNA-seq data, although *Cel-dmsr-10*, *Cel-dmsr-11*, and *Cel-dmsr-16* are somewhat enriched

within CEM VR (**Table S7**). Notably, CEM VR is the same neuron in which *srw-97* is enriched.

301   **DISCUSSION**

Pheromones are important for mating in many animals. In the nematode *C. elegans*, males are

attracted to hermaphrodites as possible mates by the small molecule pheromone ascaroside #8

(PUNGALIYA *et al.* 2009; NARAYAN *et al.* 2016). While the ascaroside class of small molecule

pheromones utilized by nematodes is rapidly being elucidated (with over 230 known ascaroside

structures so far; *https://smid-db.org*) (ARTYUKHIN *et al.* 2018), the neuronal receptor proteins that

mediate pheromone signals remain largely unknown. For only a select few ascarosides have

sensory components been identified at the cellular (GREENE *et al.* 2016a; GREENE *et al.* 2016b;

CHUTE *et al.* 2019), receptor (KIM *et al.* 2009; MCGRATH *et al.* 2011; PARK *et al.* 2012), or signal

transduction levels (ZWAAL *et al.* 1997).

311   Here, we identify two novel G protein-coupled receptors as active, required components in

the sensation and behavioral response to the mating pheromone, ascr#8. Transcripts for the two

GPCRs, *dmsr-12* and *srw-97*, are enriched in single CEM neurons (**Table 1**), and they express in

non-overlapping subsets of the male-specific chemosensory neurons (**Figure 1, 2**). There are,

however, other receptors present in these same neurons that contribute to the navigation of a vast

and ever-changing array of environmental cues, such as the widely expressed ethanol sensor, *seb-*

3 (**Figure 2B**) (JEE *et al.* 2013).

318   One mechanism for sensory flexibility in CEM neurons may be heterodimerization of

319   receptor proteins. Previous work identified two receptors for ascr#2 in the ASK neuron: DAF-37

320   and DAF-38 (PARK *et al.* 2012). While both are required for proper ascr#2-induced dauer

321   formation, only DAF-38 is involved in the sensation of other ascarosides (PARK *et al.* 2012). Such

322   heterodimers may also exist for pheromone receptors in CEM neurons.

323   Another mechanism for sensory flexibility in CEM neurons is to have diverse receptor

proteins with distinct specificities co-expressed within a single neuron. Multiple ascarosides are

sensed by the male-specific CEM neurons, including ascr#8 and ascr#3 (NARAYAN *et al.* 2016).

The two receptors identified here may function as ascr#8-specific receptors, as *dmsr-12;srw-97*

double mutant animals do not exhibit defective responses to ascr#3 (**Figure 5D, E**). The related

receptors *dmsr-10*, *dmsr-13*, *dmsr-16*, and *srw-98* are enriched in ventral CEM neurons, and may

heterodimerize with *srw-97*, to result in fine-tuned sensation of ascr#8. Similarly, there may be

other receptors may be enriched in the dorsal CEM neurons to heterodimerize with *dmsr-12*. The

role of DAF-37 as an ascr#2-specific receptor further supports our hypothesis, as *dmsr-12* is related

to DAF-37 (ROBERTSON AND THOMAS 2006). The ventral CEM-receptor, *srw-97* falls within the

same large family of GPCRs as well (KRISHNAN *et al.* 2014), while the promiscuous DAF-38 does

not (ROBERTSON AND THOMAS 2006).

The initial goal of our single-cell RNA-seq analysis of male-specific chemosensory CEM

neurons was to identify GPCRs encoding pheromone receptors. However, further analysis of other

genes with CEM-enriched expression should also uncover specific and novel promoter profiles,

which will enable optogenetic manipulation and calcium imaging of defined individual CEM

neurons (NARAYAN *et al.* 2016). The development of advanced transgenic reagents will also permit

chemical biology of CEM neurons. We have recently developed an active ascr#8 bioaffinity probe

to employ in the targeted elucidation of ascr#8 receptors (ZHANG *et al.* 2019). Use of this probe

will lead to further elucidation of the identity of acsr#8 receptors, by confirming either SRW-97

or DMSR-12 as receptors or identifying their heterodimeric partners. The combination of these

technologies should clarify how heterogeneous CEM neurons achieve homogeneous sensory

responses.

346   **METHODS**

Single cell isolation and RT-PCR

Microdissection and single-cell RT-PCR of individual CEM_DL, CEM_DR, CEM_VL,

and CEM_DR neurons was performed essentially as described (SCHWARZ *et al.* 2012). For all four

neuronal types, single-end 50-nucleotide (nt) RNA-seq was performed on an Illumina HiSeq 2000.

To identify so-called housekeeping genes and genes primarily active outside the nervous system,

we compared our results from CEM_DL, CEM_DR, CEM_VL, and CEM_DR to equivalent

analyses of published single-end 38-nt RNA-seq data from mixed-stage whole *C. elegans*

hermaphrodite larvae (SCHWARZ *et al.* 2012).

Transcriptional analysis

Reads were quality-filtered as follows: neuronal reads that failed Chastity filtering were

discarded (Chastity filtering had not been available for the larval reads); raw 38-nt larval reads

were trimmed 1 nt to 37 nt; all reads were trimmed to remove any indeterminate ("N") residues or

residues with a quality score of less than 3; and larval reads that had been trimmed below 37 nt

were deleted, as were neuronal reads that had been trimmed below 50 nt. This left a total ranging

from 21,554,964 (CEM_VL) to 24,546,096 (CEM_DL) filtered reads for analysis of each neuronal

type, versus 23,369,056 filtered reads for whole larvae (**Table S2**).

We used RSEM version 1.2.17 (LI AND DEWEY 2011) with bowtie2 version 2.2.3

(LANGMEAD AND SALZBERG 2012) and SAMTools version 1.0 (LI *et al.* 2009) to map filtered reads

to a *C. elegans* gene index and generate read counts and gene expression levels in transcripts per

million (TPM). To create the *C. elegans* gene index, we ran RSEM's *rsem-prepare-reference* with

the arguments '*--bowtie2 --transcript-to-gene-map*' upon a collection of coding DNA sequences

(CDSes) from both protein-coding and non-protein-coding *C. elegans* genes in WormBase release

WS245 (Howe *et al.* 2016). The CDS sequences were obtained from *ftp://ftp.sanger.ac.uk/pub2/*

wormbase/releases/WS245/species/c_elegans/PRJNA13758/c_elegans.

PRJNA13758.WS245.mRNA_transcripts.fa.gz and *ftp://ftp.sanger.ac.uk/pub2/wormbase/*

*releases/WS245/species/c_elegans/PRJNA13758/c_elegans.PRJNA13758.WS245.ncRNA_*

transcripts.fa.gz. For each RNA-seq data set of interest, we computed mapped reads and

expression levels per gene by running RSEM's *rsem-calculate-expression* with the arguments

'*--bowtie2 -p 8 --no-bam-output --calc-pme --calc-ci --ci-credibility-level*

0.99 --fragment-length-mean 200 --fragment-length-sd 20 --estimate-rspd --ci-memory 30000'.

These arguments, in particular '*--estimate-rspd*', were aimed at dealing with single-end data from

3'-biased RT-PCR reactions; the arguments '*--phred33-quals*' and '*--phred64-quals*' were also used

for the neuronal and larval reads, respectively. We computed posterior mean estimates (PMEs)

both for read counts and for gene expression levels, and rounded PMEs of read counts down to the

nearest lesser integer. We also computed 99% credibility intervals (CIs) for expression data, so

that we could use the minimum value in the 99% CI for TPM as a robust minimum estimate of a

gene's expression (minTPM).

We observed the following overall alignment rates of the reads to the WS245 *C. elegans*

gene index: 48.01% for the CEM_DL read set, 28.19% for the CEM_DR read set, 40.75% for the

CEM_VL read set, 19.73% for the CEM_VR read set, and 76.41% for the larval read set (**Table**

S2). A similar discrepancy between lower alignment rates for hand-dissected linker cell RNA-seq

reads versus higher alignment rates for whole larval RNA-seq reads was previously observed, and

found to be due to a much higher rate of human contaminant RNA sequences in the hand-dissected

linker cells (Schwarz *et al.* 2012). We defined detectable expression for a gene in a given

RNA-seq data set by that gene having an expression level of 0.1 TPM; we defined robust

expression by that gene having a minimum estimated expression level (termed minTPM) of at least

0.1 TPM in a credibility interval of 99% (i.e., $\geq$ 0.1 minTPM). The numbers of genes being scored

as expressed in a given neuronal type above background levels, for various data sets, are given in

Table S3. Other results from RSEM analysis are given in **Table S3**.

We annotated *C. elegans* genes and the encoded gene products in several ways (**Table S5**).

For the products of protein-coding genes, we predicted classical signal and transmembrane

sequences with Phobius 1.01 (KÄLL *et al.* 2004), regions of low sequence complexity with pseg

(SEG for proteins, from *ftp://ftp.ncbi.nlm.nih.gov/pub/seg/pseg*) (WOOTTON 1994), and coiled-coil

domains with ncoils (from *http://www.russell.embl-heidelberg.de/coils/coils.tar.gz*) (LUPAS

1996). PFAM 27.0 protein domains from PFAM (FINN *et al.* 2016) were detected with HMMER

3.0/hmmsearch (EDDY 2009) at a threshold of E $\leq 10^{-5}$. The memberships of genes in orthology

groups from eggNOG 3.0 (POWELL *et al.* 2012) were extracted from WormBase WS245 with the

TableMaker function of ACEDB 4.3.39. Genes with likely housekeeping status (based on

ubiquitous expression in both larvae and linker cells) were as identified in our previous work

(SCHWARZ *et al.* 2012). Genes were predicted to encode GPCRs on the basis of their encoding a

product containing one or more of the following Pfam-A protein domains: 7tm_1 [PF00001.16],

7tm_2 [PF00002.19], 7tm_3 [PF00003.17], 7tm_7 [PF08395.7], 7TM_GPCR_Srab [PF10292.4],

7TM_GPCR_Sra   [PF02117.11],   7TM_GPCR_Srbc   [PF10316.4],   7TM_GPCR_Srb

[PF02175.11], 7TM_GPCR_Srd [PF10317.4], 7TM_GPCR_Srh [PF10318.4], 7TM_GPCR_Sri

[PF10327.4], 7TM_GPCR_Srj [PF10319.4], 7TM_GPCR_Srsx [PF10320.4], 7TM_GPCR_Srt

[PF10321.4], 7TM_GPCR_Sru [PF10322.4], 7TM_GPCR_Srv [PF10323.4], 7TM_GPCR_Srw

[PF10324.4], 7TM_GPCR_Srx [PF10328.4], 7TM_GPCR_Srz [PF10325.4], 7TM_GPCR_Str

[PF10326.4], ABA_GPCR [PF12430.3], Sre [PF03125.13], and Srg [PF02118.16]. By this

criterion, we identified 1,615 genes encoding GPCRs in the WS245 version of the *C. elegans*

genome; this resembles a previous estimate of ~1,470 *C. elegans* genes encoding chemoreceptors

and other GPCRs, identified through extensive computational and manual analysis (HOBERT

2013). The memberships of genes in orthology groups from eggNOG 3.0 (POWELL *et al.* 2012)

were extracted directly from WormBase WS245 with the TableMaker function of ACEDB 4.3.39.

Genes with likely housekeeping status (based on ubiquitous expression in both larvae and linker

cells) were as identified in our previous work (SCHWARZ *et al.* 2012). Gene Ontology (GO)

annotations for *C. elegans* genes were extracted from WormBase-computed annotations in *ftp://*

ftp.wormbase.org/pub/wormbase/releases/WS245/ONTOLOGY/

gene_association.WS245.wb.c_elegans; human-readable text descriptions for GO term IDs were

extracted from *term.txt* in the Gene Ontology archive *http://archive.geneontology.org/full/2014-*

07-01/go_201407-termdb-tables.tar.gz.

GFP reporter construction

Reporter fusion constructs were generated using previously described techniques (BOULIN

et al. 2006). Approximately 2-3 kb of upstream promoter region of each gene was included in

construct generation, as well as a portion of the coding sequence (**Table S10**). This was then fused

to GFP (from the Fire Vector Kit plasmid, pPD95.75), via PCR fusion (BOULIN *et al.* 2006).

Primers were designed using Primer 3 and ordered from IDT (Integrated DNA Technologies).

Primer sequences available in **Table S12**. Successful fusion was confirmed via gel electrophoresis

prior to injection.

Reporter fusion constructs were injected into the gonads of *pha-1(e2123ts); lite-1(ce314);*

him-5(e1490) animals, along with a co-injection marker of pBX(*pha-1*(+)). In this manner, positive

array animals will propagate normally at 20 °C. Strains were confirmed via GFP expression, with

multiple array lines being generated per injection (**Table S11**). Injections were performed by

InVivo Biosystems, with strain isolation being performed in house.

Imaging

Animals were imaged for GFP expression using previously described techniques. GFP+

young adult male animals were mounted on a 1% agarose pad and immobilized with sodium azide.

Animals were then imaged on a spinning disk confocal microscope at 63x magnification. Z-stack

imaging was performed, generating 3D reconstructions of the heads of the imaged animals.

Central/optimal z-plane images were used to generate the images used to verify expression (**Figure**

2; Fig. S1).

RNAi Feeding

VH624 (*rhIs13* [*unc-119*::GFP + *dpy-20*(+)]; *nre-1(hd20)*; *lin-15B(hd126)*) animals

(SCHMITZ *et al.* 2007; POOLE *et al.* 2011) were crossed with *him-5(e1490)* animals to integrate

male production into a strain hypersensitive to neuronal RNAi-knockdown, generating JSR44

(*nre-1(hd20)*; *lin-15B(hd126); him-5(e1490)*). During the cross, insertion of the *him-5(e1490)*

allele displaced the integrated array *rhIs13*, suggesting location of the array on Chromosome V.

Presence of *lin-15B(hd126)* in JSR44 was confirmed via sequencing. The non-annotated

nre-1(hd20) is linked with *lin-15B*, being retained alongside *lin-15B* (SCHMITZ *et al.* 2007; POOLE

et al. 2011).

RNAi clones were grown overnight in cultures of LB containing 50 µg/mL ampicillin.

Cultures were then diluted to an OD_{600} of 1.0 before plating on NGM agar plates containing 50

µg/mL ampicillin and 1 mM IPTG (Isopropyl-β-D-thiogalactoside) to select for RNAi clones and

induce expression. Lawns were allowed to grow at room temperature for 8-16 hours, before JSR44

young adult hermaphrodites were placed on the plates and left to propagate at 16°C. Young adult

males of the F1 progeny were then selected for behavioral testing suing the *Spot Retention Assay*.

Empty vector controls (VC-1 clone) were run alongside every targeted knockdown experiment.

Spot Retention Assay

Following previously described methods, young adult males were isolated from

hermaphrodites 5-16 hours prior to testing (PUNGALIYA *et al.* 2009; NARAYAN *et al.* 2016). In

short, at the time of the assay, 0.6 µL of either vehicle control (-) or ascaroside #8 (+) was added

to the NGM plates covered in a thin lawn of OP50 *E. coli*. Ten males were then divided between

two pre-marked spots on the agar, equidistant from the cues. The plate was then recorded for 20

minutes. The time spent of each visit in either vehicle or ascaroside #8 (if greater than 10 seconds)

was scored, and averaged. Plates in which the average was greater than two standard deviations

removed from the population average were removed from the final analysis as outliers.   To

compare between strains or conditions, the vehicle was subtracted from the ascaroside dwell time

for each plate. The average of these differences was then compared statistically.

Single Worm Behavioral Assay

Following previously described methods (PUNGALIYA *et al.* 2009; NARAYAN *et al.* 2016;

REILLY *et al.* 2021), animals were isolated and prepared in an identical manner to the Spot

Retention Assay. The two outside rings of wells in a 48-well tissue culture plate were filled with

200 µL of NGM agar, which was then seeded with 65 µL of OP50 *E. coli*. The plates and lawns

were then dried at 37 °C for 4 hours. Alternating wells were then prepared as spatial controls

(nothing done), vehicle controls (0.85 µL of dH$_2$O was placed in the center of the well), or

ascaroside well (0.85 µL of ascaroside was placed in the center of the well). This was performed

over four quadrants. Animals were scored for their visits and duration to the center of the well

and/or the cue.

The average duration of each worm's visits was calculated, and these values were again

averaged together to generate a Mean Dwell Time in seconds for each plate. When comparing

across strains or conditions, the Spatial controls were then compared for statistical difference. If

none was observed, the Log(fold-change) A/V was then calculated by taking the log of the

ascaroside mean dwell time divided by the vehicle mean dwell time for each plate. The amount of

times each worm visited the center was averaged to generate the Visit Counts.

The Percent Attraction values were calculated by first determining the "attractive" cut off

as two standard deviations above the vehicle average. Any visit longer than this was deemed

"attractive" and scored as a "1"; non-attractive visits were scored as "0". The percent attraction

was then calculated for each worm was the percent of visits scored a "1". The average was then

calculated across the plate to determine Percent Attraction.

CRISPR Design and Strain Generation

A novel null mutation was generated for *srw-97* by InVivo Biosystems. The *srw-*

97(knu456) allele was generated in a *him-5(e1490)* strain using two sgRNAs

(TTTAGTAGAGCAGAAATTAA and TACAGCTTTAACTTTCAAC) to generate a 1,620-nt

deletion which removed the start codon and left only the terminal exon intact. The *knu456*

knockout allele was generated by donor homology using the pNU1361odn oligo:

(TTTTCTTGTATTTCCAAAAATTGTAAAAACCTTTATGAAAGTTAAAGCTGTAAGGAT

TTTCAGACATTTA). Following generation of a homozygous deletion by InVivo Biosystems,

the line was then backcrossed twice into *him-5(e1490)*.

The *dmsr-12(tm8706)* allele, provided by the National BioResource Group (NRBP)

contains a 118-nucleotide deletion that was generated by Dr. Mitani of the NRBP. The allele was

crossed into a *him-5(e1490)* background prior to testing. The deletion spans intron 2 and exon 3

of the coding sequence. Whether this results in a correctly spliced gene remains unknown, although

the expected remaining coding sequence remains in frame.

Phylogenetic Analyses

For phylogenetic analysis of selected CEM genes, we downloaded proteomes for

C. elegans and related *Caenorhabditis* nematodes from WormBase (release WS275), the Blaxter

Caenorhabditis database (release 1), or our unpublished work, as listed in **Table S4**. From each

proteome, we extracted the longest predicted isoform for each gene with *get_largest_isoforms.pl*

(*https://github.com/SchwarzEM/ems_perl/blob/master/fasta/get_largest_isoforms.pl*).   We

observed that the predicted isoform for *dmsr-12* in the WormBase WS275 release of *C. elegans*'

proteome was shorter than past versions of *dmsr-12*, and that the WS275 isoform omitted exons

that our transgenic expression data (based on older gene models for *dmsr-12*) indicated were likely

to be real. We therefore manually replaced the WS275 version of *dmsr-12* with an older version

(extracted from the *C. elegans* proteome in the WS250 release of WormBase). We then computed

orthology groups for *C. elegans* and its related species with OrthoFinder version 2.3.11 (EMMS

AND KELLY 2015; EMMS AND KELLY 2019), using the arguments '-*a 1 -S diamond -og*'. We

identified which orthology groups contained the *C. elegans* genes *srr-7*, *srw-97*, and *dmsr-12*, and

extracted their sequences from a concatenation of all 11 proteomes via *extract_fasta_subset.pl*

(*https://github.com/SchwarzEM/ems_perl/blob/master/fasta/extract_fasta_subset.pl*).   For   each

orthogroup's member sequences, we aligned the sequences with MAFFT version 7.455 (KATOH

AND STANDLEY 2013) and filtered the alignments twice with trimAl version 1.4.rev15 (CAPELLA-

GUTIÉRREZ *et al.* 2009), using first the argument '-*automated1*' and then the arguments '-*resoverlap*

0.50 -seqoverlap 50'. From the filtered alignments, we computed maximum-likelihood protein

phylogenies with IQ-TREE version 2.0-rc1 (NGUYEN *et al.* 2015; KALYAANAMOORTHY *et al.*

2017), using the arguments '-*m MFP -b 100 --tbe*'. In particular, we used transfer bootstrap

expectation ('--*tbe*') which provides more reliable confidence values than classic bootstrapping

(LEMOINE *et al.* 2018). We visualized the resulting phylogenies with FigTree version 1.4.4 (*http:/*

/tree.bio.ed.ac.uk/software/figtree).

Statistical Analyses

544   Prior to any statistical analyses, outliers were identified and removed. Outliers were

545   defined as any data points greater than two standard deviations removed from the average. All data

546   were then tested for normality using a Shapiro-Wilk Normality Test. This test was chosen over the

547   more conventional D'Agostino-Pearson Normality Test as many data sets were below 10 in

548   number (due to the statistical power offered by the Single Worm Behavioral Assay (REILLY *et al.*

549   2021).

550   The Spot Retention Assay data was analyzed using paired *t*-tests or Wilcoxon Matched-

Pairs Signed Rank tests to compare vehicle control and ascaroside dwell times within strains

following tests for normality (**Figure 3**). When comparing the values of multiple conditions or

strains, the data was first normalized to account for vehicle dwell time variation between plates

using a base-2 exponentiation, as described previously, to transform all data points into non-zero

values. This allows for the calculation of the fold-change as the log(base2) of the ascaroside dwell

time divided by the vehicle dwell time. These normalized values were then compared using a

Mann-Whitney test or a One-Way ANOVA followed by a Dunnett's multiple comparisons test

(**Figure 3**).

The Single Worm Behavioral Assay was first analyzed within each strain by performing

either a Repeated-Measures ANOVA followed by a Dunnett's multiple comparisons test or

Friedman test followed by Dunn's correction, comparing both Spatial control and Ascaroside

dwell times to that of the Vehicle Control. The log(fold-change) of raw ascaroside and vehicle

dwell time values were calculated and compared using either a One-Way ANOVA followed by a

Bonferroni's multiple corrections test or Friedman Test followed by Dunn's correction (**Figure 4,**

5). The ascr#3 log(fold-change) values were compared using a Student's *t*-test (**Figure 5E**). Visit

Counts were compared in the same manner as the Mean Dwell Time data, while the Percent

Attraction data was analyzed using paired *t*-tests or Wilcoxon Matched-Pairs Signed Rank tests to

compare the attractive values of the vehicle and ascaroside (**Figures S2, S3, S4**).

Data availability

RNA-seq reads will be submitted to the Sequence Read Archive (SRA; *https://*

www.ncbi.nlm.nih.gov/sra).

ACKNOWLEDGEMENTS

We thank the *Caenorhabditis* Genetics Center, which is funded by the NIH Office of

Research Infrastructure Programs (P40 OD01044), as well as the National BioResource

Project, L. Rene Garcia (Texas A&M University), and Douglas Portman (University of

Rochester Medical Center) for providing strains. We also thank InVivo Biosystems for

generating transgenic and CRISPR knockout animals. The synthetic ascr#8 utilized in this

study was provided by Frank Schroeder (Cornell University). The research reported in this

publication was supported by NIH R01 DC016058 (J.S.), R01 GM084389 (P.W.S.), the Howard

Hughes Medical Institute (P.W.S.), Moore Foundation Grant No. 4551 (E.M.S.), and Cornell

startup funding (E.M.S.). We thank Titus Brown and the Michigan State University High-

Performance Computing Center (supported by U.S. Department of Agriculture grant 2010-65205-

20361 and NIFA–National Science Foundation (NSF) grant IOS-0923812) for computational

support; additional computing was enabled by start-up and research allocations from NSF XSEDE

(TG-MCB180039 and TG-MCB190010).

FIGURE AND TABLE LEGENDS

Gene	CEM DL	CEM DR	CEM VL	CEM VR	Max CEM
pkd-2	1.30	0.09	0.25	0.77	143
seb-3	0.15	0.31	0.23	**3.25**	6.50
dmsr-12	**4.96**	0.06	0.04	0.11	4.96
srd-32	**1.62**	0.38	0.23	0.62	4.33
srw-97	0.14	0.27	0.18	**3.67**	5.50
srr-7	0.16	0.26	**1.90**	0.53	6.33
trf-1	0.00	0.00	0.00	226.28	226.28

Table 1. Enrichment levels of candidate GPCR genes in CEM neurons. Enrichment levels

displayed as normalized count within a single CEM cell type over the normalized count in the

remaining three CEM neurons. Max CEM is the enrichment in all four CEM neurons over the

remainder of the animal. The CEM with highest enrichment of each GPCR gene are denoted by

bolded-underlined values.

Figure 1. Transcriptomic landscapes of the CEM neurons. (A-D) TPKM plots of individual

CEM DL **(A)**, CEM DR **(B)**, CEM VL **(C)**, and CEM VR **(D)** neurons. X-axes display enrichment

in individual CEM neurons compared to whole larvae, while the Y-axes display total transcript

counts (TPM). Genes of interest are denoted by colored symbols, defined in the legend.

Figure 2. Expression profiles of CEM-enriched GPCR genes of interest. Transcriptional GFP

fusions of GPCR genes enriched in the CEM neurons. **(A)** A previously published CEM reporter,

p*pkd-2*::GFP. **(B)** p*seb-3*::GFP, matching previously published expression, with no discernable

enrichment in the CEM. **(C)** p*dmsr-12*::GFP is strongly expressed in the CEM DL neuron, as well

as the CEM DR soma. **(D)** p*srd-32*::GFP is weakly expressed in the somas of CEM DR and DL.

(E) p*srw-97*::GFP is expressed in both CEM VR and VL, with localization in the cilia as well as

the soma. **(F)** p*srr-7*::GFP is expressed in CEM VR, both the soma and cilia, as well the cilial of

a neighboring neuron, presumably CEP VR. Dorsal/ventral axes and anterior/posterior directions

shown in **(B)**. Scale bars denote distal, cilia region.

Figure 3. Knockdown of CEM-enriched GPCR candidates results in aberrant behavioral

response to ascr#8. (A) Raw dwell times and **(B)** Log2(fold-change of ascr#8/vehicle) of RNAi

knockdown of the kinesin motor, *osm-3*, and the TRPV channel, *osm-9* confirm the ability of RNAi

feeding to affect pheromone-driven behaviors in a targeted manner. $n \geq 5$. **(C)** Raw dwell times

and **(D)** Log2(fold-change) of genetic null *osm-3* mutants reveal an inability to respond to ascr#8.

$n \geq 17$. **(E)** Raw dwell time and **(F)** Log2(fold-change) of GPCR of interest (*srd-32*, *dmsr*-12, and

srw-97) knockdown reveals *dmsr-12* and *srw-97* as contributors to the ascr#8 behavioral response.

$n \geq 4$. Error bars denote SEM. **(A, C, E)** Paired *t*-tests and Wilcoxon tests of vehicle control versus

ascr#8 dwell time, **(B)** ANOVA, followed by Bonferroni post-hoc tests comparing Log2(fold-

change) of RNAi knockdowns to VC-1 control and to each other, **(D)** Mann-Whitney test

comparing *him-5* and *osm-3* Log2(fold-change) in response to ascr#8, **(F)** ANOVA, followed by

Dunnett's post-hoc test comparing Log2(fold-change) of RNAi knockdowns to VC-1 control. * *p*

< 0.05, ** *p* < 0.01, **** *p* < 0.0001.

Figure 4. The GPCR, SRW-97, is involved in the ascr#8 behavioral response. (A) Raw dwell

times and **(B)** Log(fold-change) of *srw-97* knockout and transgenic rescue (WorEx57) animals.

(C) Expression of WorEx57 (p*srw-97*::*srw-97*::GFP) in CEM VR. Anterior/Posterior and

Right/Left axes denoted. Error bars denote SEM. $n \geq 5$. **(A)** ANOVA followed by Dunnett's post-

hoc test or Friedman test followed by Dunn's (within strain), **(B)** Friedman test, followed by

Dunn's post-hoc test (between strain log(fold-change) values). * $p < 0.05$, ** $p < 0.01$.

Figure 5. The DMSR-12 receptor acts in non-SRW-97-expressing CEM neurons to aid in

driving the ascr#8 response. **(A)** Raw dwell times and **(B)** Log(fold-change) of *dmsr-12* knockout

and transgenic rescue (WorEx54) animals. **(C)** Expression of WorEx54 (p*dmsr-12*::*dmsr-*

12::GFP) in CEM DL and CEM DR. Anterior/Posterior and Dorsal/Ventral axes denoted. **(D)** Raw

dwell times and **(F)** Log(fold-change) of *him-5* and *dmsr-12;srw-97* animals in response to the

structurally related attracted pheromone, ascr#3. **(E)** Structural comparison of ascr#8 (top) and

ascr#3 (bottom). Error bars denote SEM. $n \geq 5$. **(A, D)** ANOVA followed by Dunnett's post-hoc

test or Friedman test followed by Dunn's post-hoc test (within strain), **(B)** ANOVA, followed by

Bonferroni post-hoc test (between strain log(fold-change) values), **(F)** Student's t-test between

strain log(fold-change) values. * $p < 0.05$.

Figure 6. Phylogenetic analysis of ascr#8-receptor candidate paralogs across the

***Caenorhabditis* genus. (A)** The phylogeny of *Cel_srw-97* reveals a *C. inopinata*-specific

amplification of *Cel_srw-98*. **(B)** The phylogeny of *Cel_dmsr-12* shows conserved counts of

orthologs across the genus. Genes for *C. elegans* denoted in purple; *C. inopinata* in pink, *C. nigoni*

in light blue, *C. briggsae* in dark blue, *C. japonica* in red, and *C. tropicalis* in dark green. Distance

reference bars (0.1) depict substitutions per site.

REFERENCES

Apirson, E. Z., and I. Ruvinsky, 2019a Coordinated behavioral and physiological responses to a

social signal are regulated by a shared neuronal circuit. Curr Biol.

Apirson, E. Z., and I. Ruvinsky, 2019b Dynamic regulation of adult-specific functions of the

nervous system by signaling from the reproductive system. Curr Biol.

Aprison, E. Z., and I. Ruvinsky, 2015 Sex Pheromones of *C. elegans* Males Prime the Female

Reproductive System and Ameliorate the Effects of Heat Stress. PLoS Genet 11**:**

e1005729.

Aprison, E. Z., and I. Ruvinsky, 2017 Counteracting Ascarosides Act through Distinct Neurons to

Determine the Sexual Identity of *C. elegans* Pheromones. Curr Biol 27**:** 2589-2599.e2583.

Artyukhin, A. B., Y. K. Zhang, A. E. Akagi, O. Panda, P. W. Sternberg *et al.*, 2018 Metabolomic

"Dark Matter" Dependent on Peroxisomal beta-Oxidation in *Caenorhabditis elegans*. J Am

Chem Soc 140**:** 2841-2852.

Bae, Y. K., H. Qin, K. M. Knobel, J. Hu, J. L. Rosenbaum *et al.*, 2006 General and cell-type

specific mechanisms target TRPP2/PKD-2 to cilia. Development 133**:** 3859-3870.

Boulin, T., J. F. Etchberger and O. Hobert, 2006 Reporter gene fusions. WormBook**:** 1-23.

Butcher, R. A., 2017 Small-molecule pheromones and hormones controlling nematode

development. Nat Chem Biol 13**:** 577-586.

Butcher, R. A., M. Fujita, F. C. Schroeder and J. Clardy, 2007 Small-molecule pheromones that

control dauer development in *Caenorhabditis elegans*. Nat Chem Biol 3**:** 420-422.

Butcher, R. A., J. R. Ragains and J. Clardy, 2009 An indole-containing dauer pheromone

component with unusual dauer inhibitory activity at higher concentrations. Org Lett 11**:**

3100-3103.

Butcher, R. A., J. R. Ragains, E. Kim and J. Clardy, 2008 A potent dauer pheromone component

in Caenorhabditis elegans that acts synergistically with other components. Proc Natl Acad

Sci U S A 105**:** 14288-14292.

Capella-Gutiérrez, S., J. M. Silla-Martínez and T. Gabaldón, 2009 trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics (Oxford,

England) 25**:** 1972-1973.

Choe, A., S. H. von Reuss, D. Kogan, R. B. Gasser, E. G. Platzer *et al.*, 2012 Ascaroside Signaling

is Widely Conserved Among Nematodes. Current Biology 22**:** 772-780.

Chute, C. D., E. M. DiLoreto, Y. K. Zhang, D. K. Reilly, D. Rayes *et al.*, 2019 Co-option of

neurotransmitter signaling for inter-organismal communication in *C. elegans*. Nat

Commun 10**:** 3186.

Cochella, L., B. Tursun, Y. W. Hsieh, S. Galindo, R. J. Johnston *et al.*, 2014 Two distinct types of

neuronal asymmetries are controlled by the *Caenorhabditis elegans* zinc finger

transcription factor die-1. Genes Dev 28**:** 34-43.

Dong, C., F. Dolke and S. H. von Reuss, 2016 Selective MS screening reveals a sex pheromone in

Caenorhabditis briggsae and species-specificity in indole ascaroside signalling. Org

Biomol Chem 14**:** 7217-7225.

Dong, C., D. K. Reilly, C. Bergame, F. Dolke, J. Srinivasan *et al.*, 2018 Comparative Ascaroside

Profiling of *Caenorhabditis* Exometabolomes Reveals Species-Specific (ω) and (ω - 2)-

Hydroxylation Downstream of Peroxisomal beta-Oxidation. J Org Chem.

Eddy, S. R., 2009 A new generation of homology search tools based on probabilistic inference.

Genome informatics. International Conference on Genome Informatics 23**:** 205-211.

Emms, D. M., and S. Kelly, 2015 OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. Genome Biology 16**:**

157.

Emms, D. M., and S. Kelly, 2019 OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biology 20**:** 238.

Fagan, K. A., J. Luo, R. C. Lagoy, F. C. Schroeder, D. R. Albrecht *et al.*, 2018 A Single-Neuron

Chemosensory Switch Determines the Valence of a Sexually Dimorphic Sensory Behavior.

Curr Biol 28**:** 902-914.e905.

Finn, R. D., P. Coggill, R. Y. Eberhardt, S. R. Eddy, J. Mistry *et al.*, 2016 The Pfam protein

families database: towards a more sustainable future. Nucleic acids research 44**:** D279-

D285.

Fraser, A. G., R. S. Kamath, P. Zipperlen, M. Martinez-Campos, M. Sohrmann *et al.*, 2000

Functional genomic analysis of *C. elegans* chromosome I by systematic RNA interference.

Nature 408**:** 325-330.

Fuxman Bass, J. I., A. M. Tamburino, A. Mori, N. Beittel, M. T. Weirauch *et al.*, 2014

Transcription factor binding to Caenorhabditis elegans first introns reveals lack of

redundancy with gene promoters. Nucleic Acids Res 42**:** 153-162.

Gene Ontology, C., 2021 The Gene Ontology resource: enriching a GOld mine. Nucleic acids

research 49**:** D325-D334.

Goodman, M. B., D. H. Hall, L. Avery and S. R. Lockery, 1998 Active currents regulate sensitivity

and dynamic range in *C. elegans* neurons. Neuron 20**:** 763-772.

Gordon, K. L., R. K. Arthur and I. Ruvinsky, 2015 Phylum-Level Conservation of Regulatory

Information in Nematodes despite Extensive Non-coding Sequence Divergence. PLoS

genetics 11**:** e1005268-e1005268.

Greene, J. S., M. Brown, M. Dobosiewicz, I. G. Ishida, E. Z. Macosko *et al.*, 2016b Balancing

selection shapes density-dependent foraging behaviour. Nature 539**:** 254-258.

Greene, J. S., M. Dobosiewicz, R. A. Butcher, P. T. McGrath and C. I. Bargmann, 2016a

Regulatory changes in two chemoreceptor genes contribute to a *Caenorhabditis elegans*

QTL for foraging behavior. Elife 5.

Harris, T. W., V. Arnaboldi, S. Cain, J. Chan, W. J. Chen *et al.*, 2020 WormBase: a modern Model

Organism Information Resource. Nucleic Acids Res 48**:** D762-d767.

Hobert, O., 2013 The neuronal genome of *Caenorhabditis elegans*. WormBook**:** 1-106.

Houck, L., C. Palmer, R. Watts, S. Arnold, P. Feldhoff *et al.*, 2007 A new vertebrate courtship

pheromone, PMF, affects female receptivity in a terrestrial salamander. Animal Behaviour

73**:** 315-320.

Howe, K. L., B. J. Bolt, S. Cain, J. Chan, W. J. Chen *et al.*, 2016 WormBase 2016: expanding to

enable helminth genomic research. Nucleic acids research 44**:** D774-D780.

Hrus, A., G. Lau, H. Hutter, S. Schenk, J. Ferralli *et al.*, 2007 *C. elegans* Agrin Is Expressed in

Pharynx, IL1 Neurons and Distal Tip Cells and Does Not Genetically Interact with Genes

Involved in Synaptogenesis or Muscle Function. PLOS ONE 2**:** e731.

Jang, Y. H., H. S. Chae and Y. J. Kim, 2017 Female-specific myoinhibitory peptide neurons

regulate mating receptivity in *Drosophila melanogaster*. Nat Commun 8**:** 1630.

Jee, C., J. Lee, J. P. Lim, D. Parry, R. O. Messing *et al.*, 2013 SEB-3, a CRF receptor-like GPCR,

regulates locomotor activity states, stress responses, and ethanol tolerance in *C. elegans*.

Genes, brain, and behavior 12**:** 10.1111/j.1601-1183X.2012.00829.x.

Kaletsky, R., V. Lakhina, R. Arey, A. Williams, J. Landis *et al.*, 2016 The C. elegans adult

neuronal IIS/FOXO transcriptome reveals adult phenotype regulators. Nature 529**:** 92-96.

Käll, L., A. Krogh and E. L. L. Sonnhammer, 2004 A combined transmembrane topology and

signal peptide prediction method. Journal of molecular biology 338**:** 1027-1036.

Kalyaanamoorthy, S., B. Q. Minh, T. K. F. Wong, A. von Haeseler and L. S. Jermiin, 2017

ModelFinder: fast model selection for accurate phylogenetic estimates. Nature methods

14**:** 587-589.

Kamath, R. S., A. G. Fraser, Y. Dong, G. Poulin, R. Durbin *et al.*, 2003 Systematic functional

analysis of the *Caenorhabditis elegans* genome using RNAi. Nature 421**:** 231-237.

Katoh, K., and D. M. Standley, 2013 MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Molecular biology and evolution 30**:** 772-780.

Kim, K., K. Sato, M. Shibuya, D. M. Zeiger, R. A. Butcher *et al.*, 2009 Two Chemoreceptors

Mediate Developmental Effects of Dauer Pheromone in *C. elegans*. Science 326**:** 994-998.

Krishnan, A., M. S. Almén, R. Fredriksson and H. B. Schiöth, 2014 Insights into the Origin of

Nematode Chemosensory GPCRs: Putative Orthologs of the Srw Family Are Found across

Several Phyla of Protostomes. PLoS ONE 9**:** e93048.

Langmead, B., and S. L. Salzberg, 2012 Fast gapped-read alignment with Bowtie 2. Nature

methods 9**:** 357-359.

Lee, H., M.-k. Choi, D. Lee, H.-s. Kim, H. Hwang *et al.*, 2012 Nictation, a dispersal behavior of the nematode Caenorhabditis elegans, is regulated by IL2 neurons. Nature Neuroscience 15**:** 107-112.

Lemoine, F., J. B. Domelevo Entfellner, E. Wilkinson, D. Correia, M. Dávila Felipe *et al.*, 2018 Renewing Felsenstein's phylogenetic bootstrap in the era of big data. Nature 556**:** 452-456.

Li, B., and C. N. Dewey, 2011 RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC bioinformatics 12**:** 323-323.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence Alignment/Map format and SAMtools. Bioinformatics (Oxford, England) 25**:** 2078-2079.

Ludewig, A. H., A. B. Artyukhin, E. Z. Aprison, P. R. Rodrigues, D. C. Pulido *et al.*, 2019 An excreted small molecule promotes *C. elegans* reproductive development and aging. Nat Chem Biol 15**:** 838-845.

Lupas, A., 1996 Prediction and analysis of coiled-coil structures. Methods in enzymology 266**:** 513-525.

Madeira, F., Y. M. Park, J. Lee, N. Buso, T. Gur *et al.*, 2019 The EMBL-EBI search and sequence analysis tools APIs in 2019. Nucleic acids research 47**:** W636-W641.

McGrath, P. T., and I. Ruvinsky, 2019 A primer on pheromone signaling in *Caenorhabditis elegans* for systems biologists. Curr Opin Syst Biol 13**:** 23-30.

McGrath, P. T., Y. Xu, M. Ailion, J. L. Garrison, R. A. Butcher *et al.*, 2011 Parallel evolution of domesticated *Caenorhabditis* species targets pheromone receptor genes. Nature 477**:** 321-325.

Narayan, A., G. Laurent and P. W. Sternberg, 2011 Transfer characteristics of a thermosensory

synapse in *Caenorhabditis elegans*. Proceedings of the National Academy of Sciences of

the United States of America 108**:** 9667-9672.

Narayan, A., V. Venkatachalam, O. Durak, D. K. Reilly, N. Bose *et al.*, 2016 Contrasting responses

within a single neuron class enable sex-specific attraction in *Caenorhabditis elegans*. Proc

Natl Acad Sci U S A 113**:** E1392-1401.

Nguyen, L.-T., H. A. Schmidt, A. von Haeseler and B. Q. Minh, 2015 IQ-TREE: a fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular

biology and evolution 32**:** 268-274.

Noble, L. M., J. Yuen, L. Stevens, N. Moya, R. Persaud *et al.*, 2021 Selfing is the safest sex for

Caenorhabditis tropicalis. Elife 10.

Park, D., I. O'Doherty, R. K. Somvanshi, A. Bethke, F. C. Schroeder *et al.*, 2012 Interaction of

structure-specific and promiscuous G-protein-coupled receptors mediates small-molecule

signaling in *Caenorhabditis elegans*. PNAS 109**:** 9917-9922.

Peden, E. M., and M. M. Barr, 2005 The KLP-6 kinesin is required for male mating behaviors and

polycystin localization in *Caenorhabditis elegans*. Curr Biol 15.

Poole, R. J., E. Bashllari, L. Cochella, E. B. Flowers and O. Hobert, 2011 A Genome-Wide RNAi

Screen for Factors Involved in Neuronal Specification in *Caenorhabditis elegans*. PLoS

Genet 7**:** e1002109.

Powell, S., D. Szklarczyk, K. Trachana, A. Roth, M. Kuhn *et al.*, 2012 eggNOG v3.0: orthologous

groups covering 1133 organisms at 41 different taxonomic ranges. Nucleic acids research

40**:** D284-D289.

Pungaliya, C., J. Srinivasan, B. W. Fox, R. U. Malik, A. H. Ludewig *et al.*, 2009 A shortcut to

identifying small molecule signals that regulate behavior and development in

Caenorhabditis elegans. Proceedings of the National Academy of Sciences of the United

States of America 106**:** 7708-7713.

Ragsdale, E. J., M. R. Muller, C. Rodelsperger and R. J. Sommer, 2013 A developmental switch

coupled to the evolution of plasticity acts through a sulfatase. Cell 155**:** 922-933.

Reilly, D. K., D. E. Lawler, D. R. Albrecht and J. Srinivasan, 2017 Using an Adapted Microfluidic

Olfactory Chip for the Imaging of Neuronal Activity in Response to Pheromones in Male

C. Elegans Head Neurons. Journal of Visualized Experiments**:** e56026.

Reilly, D. K., E. J. McGlame, E. Vandewyer, A. N. Robidoux, C. S. Muirhead *et al.*, 2021 Distinct

neuropeptide-receptor modules regulate a sex-specific behavioral response to a

pheromone. Communications Biology 4**:** 1018.

Reilly, D. K., L. J. Randle and J. Srinivasan, 2019 Evolution of hermaphroditism decreases

efficacy of Ascaroside#8-mediated mate attraction in *Caenorhabditis* nematodes.

microPublication Biology.

Robertson, H. M., and J. H. Thomas, 2006 The putative chemoreceptor families of *C. elegans*.

WormBook**:** 1-12.

Rual, J. F., J. Ceron, J. Koreth, T. Hao, A. S. Nicot *et al.*, 2004 Toward improving *Caenorhabditis*

elegans phenome mapping with an ORFeome-based RNAi library. Genome Res 14**:** 2162-

2168.

Sammut, M., S. J. Cook, K. C. Q. Nguyen, T. Felton, D. H. Hall *et al.*, 2015 Glia-derived neurons

are required for sex-specific learning in C. elegans. Nature 526**:** 385-390.

Schmitz, C., P. Kinge and H. Hutter, 2007 Axon guidance genes identified in a large-scale RNAi

screen using the RNAi-hypersensitive *Caenorhabditis elegans* strain *nre-1(hd20) lin-*

15b(hd126). Proceedings of the National Academy of Sciences 104**:** 834-839.

Schroeder, N. E., R. J. Androwski, A. Rashid, H. Lee, J. Lee *et al.*, 2013 Dauer-specific dendrite

arborization in C. elegans is regulated by KPC-1/Furin. Curr Biol 23**:** 1527-1535.

Schwarz, E. M., M. Kato and P. W. Sternberg, 2012 Functional transcriptomics of a migrating cell

in Caenorhabditis elegans. Proceedings of the National Academy of Sciences of the United

States of America 109**:** 16246-16251.

Taylor, S. R., G. Santpere, A. Weinreb, A. Barrett, M. B. Reilly *et al.*, 2021 Molecular topography

of an entire nervous system. Cell 184**:** 4329-4347.e4323.

Tenor, J. L., and A. Aballay, 2008 A conserved Toll-like receptor is required for Caenorhabditis

elegans innate immunity. EMBO Rep 9**:** 103-109.

von Reuss, S. H., N. Bose, J. Srinivasan, J. J. Yim, J. C. Judkins *et al.*, 2012 Comparative

Metabolomics Reveals Biogenesis of Ascarosides, a Modular Library of Small-Molecule

Signals in *C. elegans*. Journal of the American Chemical Society 134**:** 1817-1824.

von Reuss, S. H., and F. C. Schroeder, 2015 Combinatorial chemistry in nematodes: modular

assembly of primary metabolism-derived building blocks. Natural product reports 32**:** 994-

1006.

Wang, J., R. Kaletsky, M. Silva, A. Williams, L. A. Haas *et al.*, 2015 Cell-Specific Transcriptional

Profiling of Ciliated Sensory Neurons Reveals Regulators of Behavior and Extracellular

Vesicle Biogenesis. Curr Biol 25**:** 3232-3238.

Wang, J., M. Silva, L. A. Haas, N. S. Morsci, K. C. Q. Nguyen *et al.*, 2014 *C. elegans* Ciliated

Sensory Neurons Release Extracellular Vesicles that Function in Animal Communication.

Current Biology 24**:** 518-525.

Wang, X., J. F. Greenberg and H. M. Chamberlin, 2004 Evolution of regulatory elements

producing a conserved gene expression pattern in *Caenorhabditis*. Evolution &

Development 6**:** 237-245.

White, J. G., E. Southgate, J. N. Thomas and S. Brenner, 1986 The structure of the nervous system

of the nematode Caernorhabditis elegans. Phil Trans Royal Soc Lon 314B.

Wootton, J. C., 1994 Non-globular domains in protein sequences: automated segmentation using

complexity measures. Computers & chemistry 18**:** 269-285.

Zhang, Y. K., D. K. Reilly, J. Yu, J. Srinivasan and F. C. Schroeder, 2019 Photoaffinity probes for

nematode pheromone receptor identification. Journal of Organic & Biomolecular

Chemistry**:** 10.1039/C1039OB02099C.

Zhang, Y. K., M. A. Sanchez-Ayala, P. W. Sternberg, J. Srinivasan and F. C. Schroeder, 2017

Improved Synthesis for Modular Ascarosides Uncovers Biological Activity. Org Lett 19**:**

2837-2840.

Zwaal, R. R., J. E. Mendel, P. W. Sternberg and R. H. Plasterk, 1997 Two neuronal G proteins are

involved in chemosensation of the *Caenorhabditis* elegans Dauer-inducing pheromone.

Genetics 145**:** 715-727.

Figure - 1

Figure - 2

Figure - 3

Figure - 4

Figure - 5

Figure - 6