Signaling from the RNA sensor RIG-I is regulated by ufmylation

Daltry L. Snider[1], Moonhee Park[1], Kristen A. Murphy[1], Dia C. Beachboard[1, a], Stacy M. Horner[1,2,*]

[1] Department of Molecular Genetics & Microbiology, Duke University Medical Center, Durham, NC 27710, USA

[2] Department of Medicine, Duke University Medical Center Durham, NC 27710, USA

[a] Current address: DeSales University, Center Valley, Pennsylvania, 18034

* Correspondence: stacy.horner@duke.edu

Stacy M. Horner, Ph.D.

Duke University Medical Center

213 Research Dr., Box 3053 DUMC

Durham, NC USA 27710

Keywords

Retinoic acid-inducible gene I, RLR signaling, ubiquitin-like modifications, mitochondrial-associated ER membranes

Abstract

The RNA binding protein RIG-I is a key initiator of the antiviral innate immune response.

The signaling that mediates the antiviral response downstream of RIG-I is transduced

through the adaptor protein MAVS and results in the induction of type I and III interferons

(IFN). This signal transduction occurs at endoplasmic reticulum (ER)-mitochondrial

contact sites, to which RIG-I and other signaling proteins are recruited following their

activation. RIG-I signaling is highly regulated to prevent aberrant activation of this pathway

and dysregulated induction of IFN. Previously, we identified UFL1, the E3 ligase of the

ubiquitin-like modifier conjugation system called ufmylation, UFL1, as one of the proteins

recruited to membranes at ER-mitochondrial contact sites in response to RIG-I activation.

Here, we show that UFL1, as well as the process of ufmylation, promote IFN induction in

response to RIG-I activation. We find that following RNA virus infection, UFL1 is recruited

to the membrane targeting protein 14-3-3ε, and that this complex is then recruited to

activated RIG-I to promote downstream innate immune signaling. Importantly, we found

that 14-3-3ε has an increase in UFM1-conjugation following RIG-I activation. Additionally,

loss of cellular ufmylation prevents the interaction of 14-3-3ε with RIG-I, which abrogates

the interaction of RIG-I with MAVS and thus downstream signal transduction that induces

IFN. Our results define ufmylation as an integral regulatory component of the RIG-I

signaling pathway and as a post-translational control for IFN induction.

Significance

The viral RNA sensor RIG-I initiates the antiviral innate immune response by activating a

signaling cascade that induces interferon. Activation of the RIG-I signaling pathway is

highly regulated to quickly mount a protective immune response while preventing

dysregulation that can lead to excessive inflammation or autoimmune disorders. Here, we

characterize one such mechanism of regulation. We describe that UFL1, an E3 ligase for

the ubiquitin-like modifier conjugation system called ufmylation, is important to promote

RIG-I signaling. Using molecular approaches, we show that ufmylation promotes RIG-I

interaction with the membrane targeting protein 14-3-3ε. As such, ufmylation positively

regulates RIG-I recruitment to its signaling adaptor proteins MAVS for induction of

interferon in response to RNA virus infection.

Introduction

Detection of RNA virus infection is initiated by cellular sensors such as RIG-I. RIG-I is a pattern recognition receptor that detects unique features of viral RNA that are generally absent in cellular RNA, referred to as pathogen-associated molecular patterns (PAMPs) (1). Sensing of viral RNA PAMPs triggers RIG-I activation and induces a downstream signaling cascade that ultimately results in transcriptional induction of type I and type III interferons (IFN) and the antiviral response (2, 3). The RIG-I signaling cascade is carefully regulated by multiple mechanisms, including post-translational modifications that influence specific protein-protein interactions that can result in changes in protein localization to mediated signaling (3, 4). For example, following sensing of RNA PAMPs, RIG-I undergoes K63-linked polyubiquitination in order to transition to its fully active conformation, which promotes its interaction with the molecular trafficking protein 14-3-3ε (5–8). 14-3-3ε facilitates the recruitment of activated RIG-I from the cytosol to intracellular membranes where it interacts with MAVS (7, 9, 10), which assembles other RIG-I pathway members to transduce the signals that induce IFN (7, 11). Importantly, many RNA viruses, including influenza A virus and some flaviviruses (dengue virus, Zika virus, and West Nile virus), prevent the interaction of RIG-I with 14-3-3ε to limit IFN induction and evade the antiviral response (9, 10, 12).

In addition to RIG-I, a number of signaling proteins must be recruited to MAVS in order to propagate downstream IFN induction. Previously, we identified proteins that move to MAVS signaling sites at mitochondrial-associated endoplasmic reticulum (ER) membranes (MAM) during RNA virus infection (13, 14). These proteins likely aid in spatial organization of RIG-I pathway proteins during viral infection and include the GTPase RAB1B, which plays a role in recruiting TRAF3 to MAVS (15). In addition to RAB1B, we identified other proteins recruited to the MAM upon RIG-I signaling activation, one of which was UFL1 (referred to in our previous publication as KIAA0776) (14). UFL1 is an E3 ligase for UFM1, which is a ubiquitin-like modification of 85 amino acids. The process of ufmylation conjugates UFM1 covalently to lysine residues of target proteins through a process called ufmylation, which is similar to ubiquitination in that it also uses an E1, E2, and E3 ligase conjugation system (UBA5, UFC1, and UFL1; see Figure 2D). UFM1 is removed by the UFSP2 protease (16–20). The consequence of UFM1 addition to proteins is not fully understood, but the literature supports the idea that it can promote protein-

protein interactions to regulate a number of biological processes (21–31). Here, we

uncover a role for ufmylation in RIG-I activation. We found that the cellular proteins that

catalyze ufmylation all promote RIG-I-mediated induction of IFN. Interestingly, we found

that following RNA virus infection UFL1 interacts with both RIG-I and the molecular

trafficking protein 14-3-3ε, and that 14-3-3ε undergoes ufmylation. Further, similar to RIG-

I, UFL1 is recruited to intracellular membranes following RNA virus infection. Importantly,

loss of ufmylation prevents the interaction of 14-3-3ε with RIG-I, which results in

decreased MAVS activation and IFN induction in response to RNA virus infection. Thus,

ufmylation can regulate RIG-I activation and downstream signaling of the intracellular

innate immune system.

Results

The ufmylation activity of UFL1 promotes RIG-I signaling. Having found that the E3

ligase of ufmylation UFL1 is recruited to MAVS signaling sites at the MAM in response to

RIG-I signaling (14), we wanted to determine if UFL1 regulates RIG-I signaling. To test

this, we measured induction of the IFN-β promoter following UFL1 overexpression using

an IFN-β promoter luciferase reporter assay (32) and found that UFL1 increased activation

of the IFN-β promoter, similar to that of RIG-I expression, in a dose-dependent fashion in

response to infection with Sendai virus (SenV) (Figure 1A). SenV is a murine

paramyxovirus that specifically activates RIG-I (33). In support of UFL1 enhancing RIG-I

signaling specifically, exogenous expression of UFL1 also increased IFN-β promoter

activity in response to transfection of 293T cells with a known RIG-I immunostimulatory

RNA from hepatitis C virus (PAMP; Figure S1A) (34). However, UFL1 overexpression in

293T cells did not lead to increased induction of IFN-stimulated genes (ISG), such as

ISG56 or *ISG15,* in response to exogenous IFN-β treatment, which bypasses RIG-I

signaling and IFN induction, indicating that UFL1 primarily regulates IFN induction and not

the IFN response (Figure S1B). Next, we depleted UFL1 by siRNA in two different cell

types and measured SenV-induced activation of the RIG-I pathway. Depletion of UFL1 in

primary neonatal human dermal fibroblasts (NHDFs) reduced the SenV-mediated

induction of both *IFNB1* and *IFNL1* transcripts, as measured by RT-qPCR (Figure 1B), as

well as production of IFN-β protein, as measured by an enzyme-linked immunosorbent

assay (ELISA) (Figure 1C). Depletion of UFL1 in 293T cells resulted in decreased

phosphorylation of IRF3, a transcription factor for both type I and III IFNs, while exogenous

expression of an siRNA-resistant UFL1 restored SenV-mediated IRF3 phosphorylation

(Figure 1D).

To define the domains of UFL1 that regulate RIG-I signaling, we expressed a series of

previously described UFL1 truncation mutants and measured SenV-mediated activation

of the IFN-β promoter in a luciferase reporter assay (16). The ability of UFL1 to transfer

UFM1 to a target protein has been suggested to require the first 212 amino acids of the

protein, as this domain interacts with the E2 ligase for ufmylation, UFC1 (16). The wild-

type (WT) UFL1 (aa 1-794), as well as the C-terminal deleted mutants of UFL1, aa 1-212

and aa 1-452, which all have reported ufmylation activity (16), stimulated SenV-medicated

induction of the IFN-β promoter (Figure 1E). Interestingly, the N-terminal deleted mutant

of UFL1 aa 213-794, which does not have reported ufmylation activity, also induced

signaling, while the N-terminal deleted UFL1 mutant aa 453-794 did not (Figure 1E). To

measure UFL1 conjugation activity in our hands, we quantified the formation of higher

molecular weight UFM1-conjugates by immunoblotting. The 10 kDa UFM1 protein is

known to be conjugated to a number of proteins, including the highly abundant RPL26 (21,

28). Indeed, our analysis of the higher molecular weight UFM1-conjugates induced by

these UFL1 constructs revealed differential abundance of these UFM1-conjugates.

Specifically, UFL1 WT, aa 1-212, aa 1-452, and aa 213-794 of UFL1 all retain full

ufmylation activity, while aa 453-794 of UFL1 does not (Figure S1C). Thus, taken together,

this reveals that the ufmylation activity of UFL1 is required to promote RIG-I signaling that

results in induction of IFN.

The ufmylation machinery proteins positively regulate RIG-I signaling. Having

determined that the ufmylation activity of UFL1 is important for its role in RIG-I signaling,

we hypothesized that UFM1 and the proteins required for UFM1 conjugation would also

be required to promote this signaling. Similar to our results with UFL1, overexpression of

UFM1 increased SenV-mediated activation of the IFN-β promoter, resembling the

magnitude of induction seen with RIG-I, in a dose-dependent fashion (Figure 2A).

Conversely, the activation of the IFN-β promoter in response to SenV was significantly

abrogated in 293T cells in which UFM1 was deleted by CRISPR/Cas9, as compared to

WT 293T cells (Figure 2B). Importantly, this signaling was restored upon exogenous

expression of UFM1 (Figure 2B). The absence of UFM1 expression also prevented the

induction of IFN-β protein in response to SenV infection, as measured by ELISA (Figure

2C). The process of ufmylation has 5 steps (Figure 2D). First, UFM1 is processed to

expose the terminal glycine residue. Then, this mature UFM1 is added to the target protein

by the actions of UBA5, which acts as an E1 ligase for UFM1; UFC1, the E2 ligase; and

UFL1, the E3 ligase (19). Finally, the UFSP2 protease removes UFM1, which enables

recycling of mature UFM1 (18). We found that exogenous expression of each of the

proteins involved in UFM1 conjugation, including the UFSP2 protease, positively regulated

SenV-mediated induction of the IFN-β promoter (Figure 2E). Conversely, the induction of

IFNB1 by SenV was abrogated in 293T cells in which UFSP2 was deleted by

CRISPR/Cas9, similar to that seen upon UFM1 KO. Interestingly, we observed an

increase in UFM1-retention by an ufmylated protein and a decrease in free UFM1 in the

UFSP2 KO cells via anti-UFM1 immunoblot, suggesting UFSP2 may serve to primarily to

increase the pool of available mature UFM1 for conjugation (Figure 2F). These results

reveal that the proteins that catalyze ufmylation and the UFM1 modification itself promote

RIG-I signaling.

UFM1 is required for the RIG-I-driven transcriptional response. After establishing that

ufmylation promotes RIG-I activation, and in turn IFN expression, we next broadly

measured the impact of ufmylation upon the transcriptional response to RIG-I signaling.

Using RNA-sequencing, we analyzed gene expression in either WT or UFM1 KO 293T

cells, following mock or SenV infection (Table S1.1; Table S1.2). Gene set enrichment

analysis (Table S2.1; Table S2.2) of the transcripts significantly reduced (adjusted P<0.01)

by UFM1 KO in the absence of viral infection revealed previously described pathways

regulated by ufmylation such as cytosolic ribosomes, ribosome assembly, and

hematopoiesis (Figure S2A; Table S2.1) (21, 28, 29, 35). Following viral infection, the top

10 gene categories negatively impacted by UFM1 KO, with a darker red color indicating

more downregulation, were all related to the antiviral response, such as response to type

I IFN and defense against virus (Figure 3A; Table S2.2). Indeed, of the top 50 most

downregulated pathways impacted by UFM1 KO during infection, the majority were related

to innate immune signaling or viral replication (Table S2.2), while upregulated gene

categories were more diverse (Table S2.3; Table S2.4). Of the genes differentially

expressed during UFM1 KO in response to SenV (adjusted P<0.01), the majority are

downregulated (Figure 3B). Indeed, these downregulated genes included *IFNB1* and

IFNL1, as well as other known ISGs (in red) (36) (Figure 3B; Figure 3C). These data are

consistent with a model in which ufmylation-mediated regulation of IFN induction has
broad consequences on genes induced by the IFN response.
201
UFL1 is recruited to intracellular membranes and interacts with 14-3-3ε and RIG-I
during RNA virus infection. Following the binding of RIG-I to non-self RNA, it interacts
with several host proteins to facilitate its activation, localization to the MAM, and interaction
with MAVS. These proteins include the E3 ligases for K63-linked ubiquitin Riplet and
TRIM25 (5, 6, 37), as well as the molecular trafficking protein 14-3-3ε. In particular, 14-3-
3ε is required for RIG-I recruitment from the cytosol to MAVS signaling sites at intracellular
membranes (5–7, 13); however, the mechanism underlying how 14-3-3ε selects RIG-I as
cargo has yet to be elucidated. Using a subcellular membrane fractionation assay (38),
we confirmed that UFL1 increases its association with intracellular membranes in
response to SenV, similar to RIG-I (Figure 4A; compare fraction #1, which has Cox-I and
no GAPDH, with fractions #6-8, which are enriched for the cytosolic protein GAPDH) (7,
12). This finding is consistent with our previous report that UFL1 is recruited to the MAM
in response to either SenV or hepatitis C virus replication (14), suggesting that UFL1
recruitment occurs prior to MAVS activation, as MAVS is cleaved by the HCV NS3-NS4A
protease (39–42). As the recruitment of RIG-I to intracellular membranes is known to
require 14-3-3ε, and, as both UFL1 and UFM1 have been shown to interact with 14-3-3ε
(16), we hypothesized that UFL1 may interact with 14-3-3ε to promote the IFN induction
that we had observed in response to RNA virus infection. Thus, we first determined if the
interaction of UFL1 with 14-3-3ε is increased in response to RIG-I activation by SenV by
performing co-immunoprecipitation. We found that Myc-14-3-3ε did co-immunoprecipitate
with Flag-UFL1, as reported previously (16), and that this interaction was increased by
SenV (Figure 4B). Interestingly, the interaction of UFL1 with RIG-I also increased following
SenV, both upon over-expression and at the level of the endogenous proteins (Figure 4C;
Figure 4D). As RIG-I undergoes a series of modifications to become fully active (1, 4), we
next used a panel of RIG-I mutants to define which stage of RIG-I activation promotes
interaction with UFL1. These mutations prevent the distinct steps of RIG-I activation such
as RIG-I binding to RNA (K888/907A), interacting with TRIM25 (T55I), or ubiquitination by
Riplet and TRIM25 (K172/788R) (5, 43, 44). The interaction of UFL1 with RIG-I was
significantly impaired by each of these mutations, suggesting that UFL1 regulates RIG-I
function after it binds RNA and becomes ubiquitinated (Figure 4E). As this is the same

step of activation at which 14-3-3ε binds to RIG-I to promote its translocation to

intracellular membranes (7), we next tested the ability of UFL1 to promote RIG-I signaling

in conjunction with the RIG-I regulatory factors Riplet and 14-3-3ε (6, 7). We found that

co-expression of equal amounts of UFL1 with either Riplet or 14-3-3ε increased SenV-

mediated activation of the IFN-β promoter above that seen by the individual proteins

(Figure 4F). Intriguingly, the expression of UFL1 appears to be stabilized or enhances in

the presence of Riplet and 14-3-3ε. These data together suggest that RNA virus infection

increases the interaction of 14-3-3ε with UFL1, which then interacts with activated, K63-

ubiquitinated RIG-I to promote downstream signaling.

UFL1 interaction with RIG-I requires 14-3-3ε and UFM1. Having determined that UFL1

interacts with both activated RIG-I and 14-3-3ε following RNA virus infection, we next

defined the dynamics of this complex formation by testing two distinct models. In the first

model, UFL1 would interact first with activated RIG-I, induce its ufmylation, and then the

UFL1-RIG-I complex would interact with 14-3-3ε. In this model, depletion of 14-3-3ε or

loss of UFM1 would not be expected to change the interaction of UFL1 with RIG-I. In the

second model, UFL1 would interact first with 14-3-3ε and induces its ufmylation, or that of

another associated protein, and then the UFL1-14-3-3ε complex would interact with

activated RIG-I. In this second model, depletion of 14-3-3ε would be expected to prevent

UFL1 interaction with RIG-I, and loss of ufmylation would limit UFL1 interaction with RIG-

I but would not affect UFL1 interaction with 14-3-3ε. To elucidate these possibilities, first,

we used co-immunoprecipitation to measure the interaction of exogenously expressed

Flag-UFL1 and HA-RIG-I in SenV-infected 293T lysates that had been depleted of 14-3-

3ε or CTRL by siRNA. This revealed that formation of the SenV-activated RIG-I-UFL1

complex requires 14-3-3ε (Figure 5A). Next, we tested if ufmylation was required for

formation of the SenV-activated RIG-I-UFL1 complex by measuring this interaction in WT

or UFM1 KO 293T cells. We found that UFM1 was required for SenV-activated RIG-I-

UFL1 complex (Figure 5B). The results of these two experiments reveal that both 14-3-3ε

and UFM1 are required for UFL1 to interact with RIG-I, supporting the second model of

complex formation in which UFL1 interacts first with 14-3-3ε and catalyzes its ufmylation,

and then this complex associates with RIG-I. In support of this, we found that UFM1 was

not required for UFL1 to interact with 14-3-3ε (Figure 5C). To test if 14-3-3ε is UFM1-

conjugated , we performed an immunoprecipitation of Flag-14-3-3ε from cell extracts that

were mock or SenV-infected and expressed either HA-UFM1-WT or HA-UFM1DC3, which

lacks the terminal 3 amino acids required conjugation to target proteins (17). Importantly,

these extracts were boiled prior to immunoprecipitation to remove non-covalent

interactions. Following immunoprecipitation of Flag-14-3-3ε, immunoblotting with an anti-

HA antibody revealed a slower migrating form of 14-3-3ε approximately 15 kDa heavier

(HA+UFM1) than the predicted molecular weight of 14-3-3ε (37 kDa) (Figure 5D),

suggestive of covalent UFM1 modification. Additionally, the proportion of 14-3-3ε,

conjugated by UFM1 increases following SenV infection. Together, these data indicate

that ufmylation promotes the interaction of UFL1 with 14-3-3ε and activated RIG-I and that

UFM1 has increased conjugation to 14-3-3ε following RIG-I activation.

Ufmylation promotes RIG-I interaction with 14-3-3ε for MAVS activation. Having

found that that UFL1 requires 14-3-3ε to interact with activated RIG-I, we next tested if

UFL1 is required for the interaction of 14-3-3ε with RIG-I, which is essential for activated

RIG-I to translocate from the cytosol to intracellular membranes for interaction with MAVS

(7). We performed a co-immunoprecipitation of Flag-RIG-I and Myc-14-3-3ε from 293T

cells and found that this SenV-mediated interaction was significantly decreased upon

UFL1 depletion (Figure 6A). In addition, loss of UFM1 expression also decreased the

SenV-induced interaction of RIG-I with 14-3-3ε (Figure S3). Importantly, we also found

that UFM1 is required for the SenV-induced interaction of RIG-I with MAVS (Figure 6B)

and MAVS higher-order oligomerization, which is a hallmark of MAVS activation (45, 46)

(Figure 6C). In summary, these data reveal that UFL1 and UFM1 are required for the RIG-

I interaction with 14-3-3ε, for interaction with MAVS, and for MAVS activation by

oligomerization.

Discussion

Regulation of RIG-I activation and downstream signaling is essential for proper induction

and termination of IFN. Here, we show that both UFL1 and the process of ufmylation

promote RIG-I pathway signaling that leads to IFN induction, uncovering an important step

in the activation of the RIG-I pathway. RIG-I activation occurs upon RNA binding. Then,

RIG-I undergoes ATP hydrolysis. And interaction with K63-linked polyubiquitin chains,

both covalently and non-covalently (5, 44, 47), which promotes formation of a RIG-I

tetramer (48). This polyubiquitinated, activated RIG-I oligomer then interacts with the
membrane trafficking protein 14-3-3ε for translocation to MAVS at ER-mitochondrial
contact sites (7). We found that UFL1 is recruited to 14-3-3ε following RNA virus infection
and that ufmylation facilitates the interaction between 14-3-3ε and activated RIG-I.
Importantly, this results in increased interaction of RIG-I with MAVS and MAVS
oligomerization, ultimately promoting the downstream signal transduction which produces
IFN.
305
Ufmylation is emerging as a post-translational modification that regulates diverse
biological processes, including DNA repair, ER homeostasis, and even the replication of
hepatitis A virus (21, 22, 24, 27, 28, 30, 35, 49). In these cases, UFL1, along with the other
members of the ufmylation cascade, induce ufmylation of a target protein important for
regulating these processes. For example, both MRE11 and histone H4 are ufmylated by
UFL1 in the nucleus in response to DNA damage resulting in activation the key DNA repair
kinase ATM (22, 24). UFL1 can also act at the ER, where it plays a role in ER protein
quality control, where it ufmylates specific proteins, including ribosomal proteins RPL26,
to induce lysosomal degradation of stalled peptides and/or the ER and prevent the
unfolded protein response (27, 28, 49, 50). Hepatitis A virus translation, which occurs in
association with the ER, also requires ufmylation of RPL26 (30). Therefore, ufmylation can
regulate several aspects of translation. It is possible that ufmylation regulates translation
of certain mRNAs important for RIG-I signaling and subsequent IFN induction. However,
we identified a role for ufmylation in regulating the interaction of RIG-I with 14-3-3ε, one
of the earliest known steps of RIG-I signaling, strongly supporting a mechanism in which
following RIG-I activation, ufmylation is controlling this specific protein-protein interaction.
The mechanisms by which the process of UFM1 addition regulates interactions between
proteins or alters other aspects of protein function are largely unknown. Indeed, we found
that in both overexpression and KO conditions UFSP2, the protease that removes UFM1
from proteins (Figure 2D) (18), also promoted SenV-mediated IFN induction (Figure 2E-
F), suggesting that we do not yet have a full grasp on the ufmylation process. It is possible
that the dynamic process of ufmylation or the enhanced formation of mature UFM1
following deconjugation from targets promote RIG-I signaling independent of
deconjugation activity. Interestingly, in the UFSP2 KO cells, we observed increased
amounts of a specific higher molecular weight UFM1-conjugates by immunoblot,
suggesting that UFSP2 is important for the generation of free UFM1 for conjugation

(Figure 2F). Indeed, in support of this idea, others have shown that UFSP2 in myeloid
cells is required for influenza virus resistance in mice (31). It is also possible that UFSP2
acts on other members of the RIG-I pathway to alter their function. Future studies to define
how the process of ufmylation regulates this and other aspects of the antiviral innate
immune response will be of great interest as they may shed light broadly on how
ufmylation regulates diverse cell biological processes that alter cellular signaling.
338
The mechanisms underlying how cytoplasmic UFL1 is recruited to its protein targets that
reside in different subcellular compartments are not fully known. For example, we found
that RIG-I activation induces UFL1 translocation to intracellular membranes (Figure 3A),
and while we know that UFL1 is recruited to the MAM during infection, the mechanism by
which UFL1 becomes membrane-associated remains unknown (14). DDRGK1 (UFBP1)
may facilitate UFL1 targeting to the MAM, as DDRGK1 is localized to mitochondrial-ER
contact sites (16, 51) and in some cases it is required for UFL1 recruitment to membranes
(27, 28). Thus, both DDRGK1 and mitochondrial-ER contact sites could function as a
regulatory hub that aids in the recruitment of UFL1 and RIG-I pathway signaling proteins.
Interestingly, RAB1B, a GTPase that we found is recruited to the MAM and important for
RIG-I signaling (14, 15) is ufmylated (52, 53), which reveals that ufmylation likely regulates
a number of RIG-I pathway signaling proteins. As UFL1 contains no functional domains
common to other E3 ligases that might allow one to predict how its targets are selected
(16, 54, 55), defining the signals and features that control UFL1 localization, as well as
the target proteins ufmylated in response to RIG-I activation, will undoubtedly reveal clues
into how the process of ufmylation is activated and how specific targets are selected.
355
Our work revealed that 14-3-3ε required ufmylation to interact with activated RIG-I. The
details underlying how 14-3-3ε interacts with activated RIG-I have not been fully
elucidated, as it does not occur through the known phosphorylated amino acids on RIG-I,
the typical recruitment signal of the 14-3-3 family of proteins (7, 56, 57). Our work supports
the work of others demonstrating that 14-3-3ε interacts with UFM1 and other members of
the ufmylation pathway, and it reveals that SenV increases the proportion of ufmylated 14-
3-3ε (Figure 5D) (16). It is likely that 14-3-3ε is mono-ufmylated as the apparent molecular
weight of 14-3-3ε increased by one ufmylation group (HA+UFM1, 15 kDa; Figure 5D).
Thus, taken together with our results, this suggests that ufmylation of 14-3-3ε or a 14-3-
3ε-associated protein promotes the interaction between activated RIG-I and 14-3-3ε. In

fact, a number of 14-3-3 family proteins are post-translationally modified by
phosphorylation, acetylation, and oxidation (58). Therefore, post-translational modification
of 14-3-3ε by ufmylation could define how cargo proteins, including RIG-I, are selected.
Indeed, this mechanism could be shared with other RNA virus sensing pathways, such as
the RIG-I-like-receptor MDA5, which also interacts with a 14-3-3 protein, 14-3-3η, by an
unknown mechanism (59). Thus, ufmylation may broadly influence how 14-3-3 proteins or
other host proteins interact with each other to regulate the intracellular innate immune
response.

Overall, this work lays the groundwork for future studies to define how ufmylation of
antiviral innate immune signaling proteins regulates their function and how specific
signaling pathways are differentially activated through ufmylation. In addition, our work
adds ufmylation to the growing list of ubiquitin-like and other modifications that regulate
the intracellular innate immune response, including ISGylation, SUMOylation, FATylation,
acetylation, phosphorylation, and others (4, 60, 61) broadening our understanding of how
RIG-I signaling is activated and rapidly controlled by post-translational modifications in
response to infection, leading to greater knowledge of the exquisite regulation of these
pathways.

Materials and Methods

Cell lines, viruses, and treatments. Neonatal human dermal fibroblast (NHDF) cells and
embryonic kidney 293T cells were grown in Dulbecco's modification of Eagle's medium
(DMEM; Mediatech) supplemented with 10% fetal bovine serum (Thermo Fisher
Scientific), 1X minimum essential medium non-essential amino acids (Thermo Fisher
Scientific), and 25 mM HEPES (Thermo Fisher Scientific) (cDMEM). 293T (CRL-3216)
were obtained from American Type Culture Collection (ATCC), NHDF cells (CC-2509)
were obtained from Lonza. All cell lines were verified as mycoplasma free by the LookOut
Mycoplasma PCR detection kit (Sigma). SenV Cantell strain was obtained from Charles
River Laboratories and used at 200 hemagglutination units/mL (HAU). SenV infections
were performed in serum-free media (30 minutes to 1 hour), after which complete media
was replenished. IFN-β (PBL Assay Science) was added to cells at a concentration of 50
units/mL in cDMEM for 18 hours.

Plasmids. The following plasmids have been previously described: pEF-TAK-Flag, pEF-
BOS-Flag-RIG-I (62), pEF-TAK-Flag-Riplet (63) pIFN-β-luc (64), pCMV-Renilla
(Promega), pX459 (Addgene #62988), psPAX2 (Addgene #12260), and pMD2.G
(Addgene #12259), pEF-BOS-Flag-RIG-I T55I (65), pEF-TAK-Myc-MAVS (32).
pLJM1_Flag-UFM1 was a gift from Drs. Craig McCormick and John Rohde at Dalhouise
University. The following plasmids were generated by insertion of PCR-amplified
fragments into the NotI-to-PmeI digested pEF-TAK-Flag using InFusion cloning
(Clontech): pEF-TAK-Flag-UFL1 (GenBank: BC036379; GeneID: 23376), pEF-TAK-Flag-
UBA5 (NM_024818.6), pEF-TAK-Flag-UFC1 (NM_016406.4), pEF-TAK-Flag-UFSP2
(NM_018359.5), pEF-TAK-Flag-UFL1 1-212, pEF-TAK-Flag-UFL1 1-452, pEF-TAK-Flag-
UFL1 213-794, pEF-TAK-Flag-UFL1 453-794, and pEF-TAK-Flag-14-3-3ε. Both pEF-
TAK-Myc-14-3-3e and pEF-TAK-Myc-UFL1 were generated by insertion of PCR-amplified
fragments into the AgeI-NotI digested pEF-TAK-Myc (pEF-TAK-Myc-MAVS) by InFusion.
The pEF-TAK-HA vector was generated by PCR to replace Flag with HA, and pEF-TAK-
HA-RIG-I, and pEF-TAK-HA-UFM1 were generated by insertion of a PCR-amplified
fragment into the NotI-AgeI digested pEF-TAK-HA vector. pLEX-Flag-14-3-3e was
generated by ligation of a PCR-amplified Flag-14-3-3e into the BamHI-to-XhoI digested
pLEX using InFusion cloning. The following plasmids were generated by site-directed
mutagenesis: pEF-TAK-Flag-UFL1^{siR}, pEF-TAK-HA-UFM1DC3, pEF-BOS-Flag-RIG-I
K888/907A, and pEF-BOS-Flag-RIG-I K172/788R. To generate the CRISPR guide RNA
plasmids px459-UFM1-E2, px459-UFM1-B, and px459-UFSP2 sgRNA oligonucleotides
were annealed and inserted into the BbsI-digested pX459 (30, 66). The plasmid
sequences were verified by DNA sequencing and oligonucleotide sequences are available
upon request.

Generation of RNA PAMP. Annealed oligonucleotides containing the sequence of the
HCV 5'ppp poly-U/UC region (34) were *in vitro* transcribed using the MEGAshortscript T7
transcription kit (Ambion) followed by ethanol precipitation, with the resulting RNA
resuspended at 1 µg/µL.

Transfection. DNA transfections were performed using FuGENE6 (Promega) or TransIT-
LT1 (Mirus Bio). RNA PAMP transfections were done using the TransIT-mRNA
Transfection kit (Mirus Bio). The siRNA transfections were done using Lipofectamine
RNAiMax (Invitrogen). siRNAs directed against 14-3-3ε (Dharmacon-L-017302-02-0005),

UFL1 (Qiagen-SI04371318) or non-targeting AllStars negative control siRNA (Qiagen-

1027280) were transfected into 293T cells (25 pmol of siRNA; final concentration of 0.0125

µM) or NHDF cells (250 pmol of siRNA; final concentration of 0.25 µM). Media was

changed 4-24 hours post-transfection, and cells were incubated for 36-48 h post-

transfection prior to each experimental treatment. IFN-β-promoter luciferase assays were

performed as previously described at 18-24 hours post treatment and normalized to the

Renilla luciferase transfection control (33).

ELISA. IFN-β ELISAs were performed using Human IFN-beta DuoSet (R&D Systems)

with supernatants collected from cultured cells.

Generation of cell lines. UFM1 and UFSP2 KO 293T cells were generated by

CRISPR/Cas9, using two guides targeting exon 2 and 3 (UFM1) or a single guide targeting

exon 6 (UFSP2), similar to others, as we have done previously. Single cell clones were

validated via anti-UFM1 or anti-UFSP2 immunoblot and genomic sequencing, with one

clone used here (15, 30). 293T cell pools overexpressing Flag-14-3-3e were generated

by lentiviral transduction, as previously (36).

RNA analysis. Total cellular RNA was extracted using the RNeasy Plus mini kit (Qiagen).

RNA was then reverse transcribed using the iScript cDNA synthesis kit (BioRad), as per

the manufacturer's instructions. The resulting cDNA was diluted 1:3 in ddH2O. RT-qPCR

was performed in triplicate using the Power SYBR Green PCR master mix (Thermo-

Fisher) and QuantStudio 6 Flex RT-PCR system. Oligonucleotide sequences for qPCR

are available upon request.

RNA-seq. WT and UFM1 KO 293T cells were mock or SenV-infected (18 h) and

harvested in biological duplicate, followed by total RNA extraction via TRIzol reagent

(Thermo Fisher Scientific). Sequencing libraries were prepared using the KAPA Stranded

mRNA-Seq Kit (Roche) and sequenced on an Illumina Novaseq 6000 with 50 bp paired-

end reads (>20 million reads per sample) in an S1 flow cell by the Duke University Center

for Genomic and Computational Biology. RNA-seq data was processed using the

TrimGalore toolkit (67) which employs Cutadapt (68) to trim low-quality bases and Illumina

sequencing adapters from the 3' end of the reads. Only reads that were 20nt or longer

after trimming were kept for further analysis. Reads were mapped to the GRCh38v93

version of the human genome and transcriptome (69) using the STAR RNA-seq alignment

tool (70). Reads were kept for subsequent analysis if they mapped to a single genomic

location. Gene counts were compiled using the HTSeq tool (71). Only genes that had at

least 10 reads in any given library were used in subsequent analysis. Normalization and

differential expression was carried out using the DESeq2 (72) Bioconductor (73) package

with the R statistical programming environment. The false discovery rate was calculated

to control for multiple hypothesis testing. Gene set enrichment analysis (74) was

performed to identify gene ontology terms and pathways associated with altered gene

expression for each of the comparisons performed. All RNA-seq data are deposited in the

GEO database under GSE186287.

Immunoblotting. Cells were lysed in a modified radioimmunoprecipitation assay (RIPA)

buffer (10 mM Tris [pH 7.5], 150 mM NaCl, 0.5% sodium deoxycholate, and 1% Triton X-

100) supplemented with protease inhibitor cocktail (Sigma) and Halt Phosphatase Inhibitor

(Thermo-Fisher), and post-nuclear lysates were isolated by centrifugation. Quantified

protein (between 5 -15 µg) was resolved by SDS/PAGE, transferred to nitrocellulose or

polyvinylidene difluoride (PVDF) membranes in a 25 mM Tris-192 mM glycine-0.01% SDS

buffer. Membranes were stained with Revert 700 total protein stain (LI-COR Biosciences)

and then blocked in 3% BSA in Tris-buffered saline containing 0.01% Tween-20 (TBS-T).

After washing with PBS-T or TBS-T (for phosphoproteins) buffer, following incubation with

primary antibodies, membranes were incubated with species-specific horseradish

peroxidase-conjugated antibodies (Jackson ImmunoResearch, 1:5000) or fluorescent

secondaries (LI-COR Biosciences), followed by treatment of the membrane with Clarity

Western ECL substrate (BioRad) and imaging on a LICOR Odyssey FC. The following

antibodies were used for immunoblotting: R-anti-SenV (MBL, 1:1000), M-anti-Tubulin

(Sigma, 1:1000), R-anti-GAPDH (Cell Signaling Technology, 1:1000), R-anti-p-IRF3 (Cell

Signaling Technology, 1:1000), R-anti-IRF3 (Cell Signaling Technology, 1:1000), R-anti-

UFL1 (Novus Biologicals, 1:1000), R-anti-UFM1 (Abcam, 1:1000), anti-RIG-I (M-

AdipoGen, R-Abcam, 1:1000), R-anti-14-3-3ε (Cell Signaling Technology, 1:1000), M-anti-

Flag M2 (Sigma, 1:1000), anti-Flag-HRP (Sigma, 1:1000-1:5000), R-anti-Flag (Sigma,

1:1000), anti-HA (M- and R-Sigma, 1:1000), and anti-Myc (M-Santa Cruz or R-Cell

Signaling Technology, 1:1000).

Immunoprecipitation. Cells were lysed in RIPA buffer with or without 10% glycerol.
Quantified protein (between 100-500 µg) was incubated with protein-specific, isotype
control antibody (R-Cell Signaling Technology or M-Thermo Fisher), or anti-Flag M2
magnetic beads (Sigma), in lysis buffer either at room temperature for 2 h or at 4ºC
overnight with head over tail rotation. The lysate/antibody mixture was then incubated with
Protein G Dynabeads (Invitrogen) for 1 h. Beads were washed 3X in PBS or RIPA buffer
and eluted in 2X Laemmli Buffer (BioRad) with or without 5% 2-Mercaptoethanol at 95ºC
for 5 min. Proteins were resolved by SDS/PAGE and immunoblotting, as above. For
ufmylation immunoprecipitations, cells were lysed with NP-40 buffer (50 mM Tris [pH 8],
150 mM NaCl, 0.5% sodium deoxycholate, and 1% NP-40) supplemented with 10%
glycerol, protease and phosphatase inhibitors, as above, and N-ethylmaleimide (Sigma-
Aldrich). Post-nuclear lysates were boiled at 95°C for 5 min and incubated with anti-Flag
M2 magnetic beads, as above.

Subcellular membrane fractionation. Membrane fractionation was performed as
previously described (7, 12, 38, 75). Cells were lysed in hypotonic buffer (10 mM Tris-HCL
(pH 7.5), 10 mM KCl, and 5 mM $MgCl_2$ supplemented with protease inhibitor cocktail) for
10 minutes on ice followed by 20 passages through a 20-guage needle. Nuclei and
unbroken cells were removed by centrifugation at 1000xg for 5 min at 4ºC. The resulting
supernatants were mixed thoroughly with 72% sucrose and overlayed with 55% sucrose,
followed by 10% sucrose, all in low-salt buffer (2 nM EDTA, 20 nM HEPES (pH 8.0), 150
mM NaCl, 0.1% SDS, 1% Triton X-100). The gradients were subjected to centrifugation at
38,000 RPM in a Beckman SW41 Ti Rotor for 14 h at 4ºC. 1 mL fractions were collected
using a BioComp piston gradient fractionator and resulting fractions were divided in half
and mixed with 2 parts 100% methanol and precipitated overnight at -80ºC. Protein pellets
were collected by centrifugation and resuspended in 2X Laemmli buffer and heated for 5
min at 95ºC for immunoblot analysis. 10% pre-fractionated cells from each condition were
collected as the input.

Semi-denaturing detergent agarose gel electrophoresis. SDD-AGE was performed as
described (45, 46). Briefly, crude mitochondria (P5 fraction) were isolated from an equal
number of WT or UFM1 KO 293T cells that were mock or SenV infected (12 h),
resuspended in hypotonic buffer (10 mM Tris, pH 7.5, 10 mM KCl, 1.5 mM $MgCl_2$, and 0.5
mM EDTA). Resulting samples were split and 2X SDD-AGE sample buffer (0.5X TBE,

10% glycerol, 2% SDS, 0.2 mM Bromophenol Blue) buffer with or without 5% 2-Mercaptoethanol was added, and samples were loaded onto a vertical 1.5% agarose gel. Electrophoresis was performed with a constant voltage of 70 V at 4 °C in SDD-AGE running buffer (1X TBE and 0.1% SDS). Gels were transferred onto a nitrocellulose membrane overnight on ice at 25 V. Membranes were fixed in 0.25% glutaraldehyde in PBS and immunoblotting was performed as usual. 15% of the SDD-AGE samples were reserved for input.

Quantification of immunoblots. Immunoblots imaged using the LICOR Odyssey FC were quantified by ImageStudio software, and raw values were normalized to relevant controls for each antibody. Phosphoprotein values were normalized to Tubulin and displayed as the percentage of signal from WT. Relative membrane association of UFL1 was quantified as the ratio of UFL1 to Cox-1 in fraction 1 normalized to total protein levels of UFL1 in the input and displayed as the percentage of UFL1 membrane association normalized to mock values.

Statistical analysis. Student's unpaired t-test, one-way ANOVA, or two-way ANOVA were implemented for statistical analysis of the data followed by appropriate post-hoc test (as indicated) using GraphPad Prism software. Graphed values are presented as mean ± SD or SEM (n = 3 or as indicated); $*p \leq 0.05$, $**p \leq 0.01$, and $***p \leq 0.001$.

Acknowledgments

We thank colleagues who provided reagents (see Methods), the Duke Functional Genomics Core Facility, the Duke University Center for Genomic and Computational Biology, the Duke Genomic Analysis and Bioinformatics Shared Resource and Dr. Wei Chen for performing RNA-seq analysis, Dr. Madhuvanthi Vijayan and Boyoung Michelle Kim for generation and initial characterization of plasmids, Dr. Matthew Thompson for input in RNA-seq sample preparation and data processing, and Horner lab members for useful discussion. This work was supported by funds from Burroughs Wellcome Fund, National Institutes of Health: R21AI144380, R01AI155512, T32-CA009111, and an American Cancer Society Postdoctoral fellowship 131321-PF-17-188-01-MPC.

Author Contributions: Conceptualization: D.L.S., D.C.B., and S.M.H.; Investigation:

D.L.S, D.C.B., and M.P.; Formal analysis: D.L.S., D.C.B., K.A.M., and S.M.H.; Writing –

original draft: D.L.S. and S.M.H.; Writing – review and editing: D.C.B., D.L.S., and S.M.H.

All authors read the manuscript and provided comments. Funding acquisition: S.M.H.

Competing Interest Statement: The authors declare that they have no conflicts of

interest with the contents of this article. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institutes of

Health.

Figure Legends

Figure 1. The ufmylation activity of UFL1 promotes RIG-I signaling.

A) IFN-β-promoter reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells

expressing vector, Flag-UFL1, or Flag-RIG-I, followed by mock or SenV infection (18 h).

B) RT-qPCR analysis (rel. to *18S*) of RNA extracted from primary neonatal human

dermal fibroblasts (NHDFs) transfected with either siCTRL or siUFL1 followed by mock

or SenV infection (8 h). C) ELISA for IFN-β of supernatants harvested from NHDFs

transfected with siCTRL or siUFL1 and infected with SenV for the indicated times. D)

Immunoblot analysis of p-IRF3 following siRNA transfection along with expression of

vector or Flag-UFL1[siR], which has point mutations in the siRNA seed sequence.

Quantification of p-IRF3/Tubulin is shown on underneath. E) Relative IFN-β-promoter

reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells expressing indicated

constructs followed by mock or SenV infection (12-18 h), with results graphed as relative

SenV fold change for each. For A) mean -/+ SD, n=3 technical replicates and

representative of n=3 independent experiments. For all others, mean -/+ SEM, n=3 or

n=5 (1E) biological replicates. *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001 determined by one-

way ANOVA followed by Šidák's multiple comparisons test (C), Student's t-test (B, D), or

one-way ANOVA followed by Dunnett's multiple comparisons test (E).

Figure 2. The ufmylation machinery proteins positively regulate RIG-I signaling.

A) IFN-β-promoter reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells

expressing vector, Flag-UFM1, or Flag-RIG-I followed by mock or SenV infection (18 h)

or in B) WT or CRISPR/CAS9 UFM1 KO 293T cells transfected with vector (Vec) or

Flag-UFM1 (for KO), followed by mock or SenV infection (18 h). C) ELISA for IFN-β of

supernatants harvested from WT or CRISPR/CAS9 UFM1 KO 293T cells that were

SenV infected (18 h). D) Diagram of UFM1 conjugation. E) Relative IFN-β-promoter

reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells expressing indicated

constructs followed by mock or SenV infection (18 h), with results graphed as relative

SenV fold change for each. F) RT-qPCR analysis (rel. to *GAPDH*) of RNA extracted from

293T WT, UFM1 KO or UFSP2 KO with either mock or SenV infection (18 h). For A)

mean -/+ SD, n=3 technical replicates and representative of n=3 independent

experiments. For all others, mean -/+ SEM, n=3 biological replicates. *p ≤ 0.05, **p ≤

0.01, and ***p ≤ 0.001 determined by two-way ANOVA followed by Tukey's multiple

comparisons test (A-B), Student's t-test (C), or one-way ANOVA followed by Dunnett's

multiple comparisons test (E, F).

Figure 3. UFM1 is required for the RIG-I driven transcriptional response.

RNA-seq analysis WT or UFM1 KO 293T cells following mock or SenV infection (18 h).

A) Gene set enrichment analysis of negatively regulated differentially expressed genes

in SenV-infected 293T cells represented by normalized enrichment score (UFM1 KO /

WT). B) Volcano plot of differentially expressed genes (adj P<0.01) shown in grey, with

ISGs shown in red, in SenV-infected 293T cells (UFM1 KO / WT). C) Heatmap of the

effect of UFM1 KO on the fold change of the 50 most induced IFN and ISGs (UFM1 KO /

WT) following SenV infection (adj P<0.01).

Figure 4. UFL1 is recruited to intracellular membranes and interacts with 14-3-3ε

and RIG-I during RNA virus infection.

A) Immunoblot analysis of inputs and subcellular membrane flotation of 293T cell

extracts that were mock or SenV-infected (4 h) followed by sucrose gradient

fractionation, with fraction numbers indicated from the top of the gradient (1) to bottom

(8). Fractionation controls, GAPDH for cytosol and Cox-I for membranes, are indicated

and reveal that the membranes are localized to fraction #1. Relative quantification of the

ratio of UFL1 to a membrane marker (Cox-I) in fraction 1 normalized to total protein

levels in inputs are shown on the right. B) Immunoblot analysis of anti-Flag

immunoprecipitated extracts and inputs from 293T cells expressing Myc-14-3-3ε and

Flag-UFL1 that were mock- or SenV-infected (4 h), with relative quantification on right.

C) Immunoblot analysis of anti-Flag immunoprecipitated extracts and inputs from 293T

cells expressing Myc-UFL1 and Flag-RIG-I that were mock- or SenV-infected (4 h), with

relative quantification with IP values normalized to inputs values on right. D) Immunoblot

analysis of anti-RIG-I immunoprecipitated (or anti-IgG) extracts and inputs from 293T

cells that were mock- or SenV-infected (4 h), with relative quantification with IP values

normalized to inputs values on right. E) Immunoblot analysis of anti-Flag

immunoprecipitated extracts and inputs from 293T cells expressing Myc-UFL1 and Flag-

RIG-I constructs that were mock- or SenV-infected (4 h), with results quantified as

relative fold change (SenV to Mock) for each. F) Relative IFN-β-promoter reporter

luciferase expression (rel. to CMV-*Renilla*) from 293T cells expressing indicated

constructs followed by mock or SenV infection (18 h), with results graphed as relative

SenV fold change for each. The graphs are represented as the mean -/+ SEM, n=3 (A-B,

D-F) or n=4 (C) biological replicates and *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001

determined by Student's t-test (A-D) or one-way ANOVA followed by Dunnett's multiple

comparisons test (E-F).

Figure 5. UFL1 interaction with RIG-I requires 14-3-3ε and ufmylation.

A) Immunoblot analysis of anti-HA immunoprecipitated extracts and inputs from 293T

cells transfected with siCTRL or si14-3-3ε followed by SenV infection (4h). B)

Immunoblot of anti-HA immunoprecipitated extracts and inputs from 293T WT or UFM1

KO cells transfected with HA-RIG-I and Flag-UFL1. C) Immunoblot of anti-Flag

immunoprecipitated extracts and inputs from 293T WT or UFM1 KO cells transfected

with Flag-UFL1 and Myc-14-3-3ε. D) Immunoblot of anti-flag immunoprecipitated

extracts and inputs from either 293T-WT or 293T-Flag-14-3-3ε cells transfected with

either HA-UFM1-WT or HA-UFM1DC3. In (A-D), SenV infection was for 4 hours, and

relative quantification of the interacting protein vs. IP protein is shown underneath (A-C)

or as a graph (D), indicating the mean -/+ SEM (A, B, D), n=3 (A, B) or n=4 (D) biological

replicates. For (C) values shown are SD of IP values adjusted for input expression, with

n=2 biological replicates. *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001 determined by Student's

t-test.

Figure 6. Ufmylation promotes RIG-I interaction with 14-3-3ε for MAVS activation.

A) Immunoblot of anti-Flag immunoprecipitated extracts and inputs from 293T cells

transfected with siCTRL or siUFL1 and indicated constructs. B) Immunoblot of anti-Myc

immunoprecipitated extracts from 293T WT or UFM1 KO cells. D) 293T WT or UFM1 KO

were mock or SenV-infected (12 h). Immunoblotting shows endogenous MAVS in input

samples and MAVS aggregation from P5 fractions, in the presence or absence of

denaturing reagent (β-mercaptoethanol). SenV infection was for 4 h (A-C) or 12 h (D). In

(A-B), relative quantification of interacting protein vs. IP protein in the IP is shown

underneath; in (D) SDD-AGE MAVS values are normalized to corresponding SDS-PAGE

values and shown as the ratio of SenV/mock values. Values (A-B) or graph (C) show the

mean -/+ SEM for n=3 biological replicates. *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001

determined by Student's t-test.

Figure S1. UFL1 positively regulates IFN induction.

A) IFN-β-promoter reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells

transfected with vector (Vec) or Flag-UFL1, followed by mock or HCV PAMP RNA

transfection (24 h). B) RT-qPCR analysis (rel. to *GAPDH*) of RNA extracted from 293

cells transfected vector or Flag-UFL1 that were treated with IFN-β (18 h). C) Quantification

of immunoblots from 293T cells expressing HA-UFM1, and either vector or indicated Flag-

UFL1 represented as the ratio of UFM1 conjugates (approximately 25-50 kDa) to total

protein expression in each lane with vector set to 100. Graph indicates the mean -/+ SEM

for n=3 biological replicates. *p ≤ 0.05 and **p ≤ 0.01 as determined by two-way ANOVA

followed by Šidák's multiple comparisons test (A), Student's t-test (B) or one-way ANOVA

followed by Dunnett's multiple comparisons test (C).

Figure S2. Transcriptional response of genes negatively regulated by UFM1.

RNA-seq analysis of WT versus UFM1 KO 293T cells showing the gene set enrichment

analysis (top 10 categories) of negatively regulated differentially expressed genes

represented by normalized enrichment score to identify gene ontology terms and

pathways associated with altered gene expression for each of the comparisons

performed (adj P<0.01).

Figure S3. UFM1 is required for the interaction of RIG-I and 14-3-3ε.

Immunoblot of anti-Flag immunoprecipitated extracts and inputs from 293T WT or UFM1

KO cells with relative quantification on the right. Graph shows the mean -/+ SEM for n=3

biological replicates; ***p ≤ 0.001 determined by Student's t-test.

Dataset S1. Differential expression analysis from RNA-seq analysis for UFM1 KO /

WT 293T cells

Table S1.1: UFM1 KO / WT Mock

Table S1.2 UFM1 KO / WT SenV (18 h)

Dataset S2. Gene Set Enrichment Analysis for UFM1 KO / WT 293T cells

Table S2.1: UFM1 KO / WT Mock- negative direction

Table S2.2 UFM1 KO / WT SenV (18 h)- negative direction

Table S2.3 UFM1 KO / WT Mock- positive direction

Table S2.4 UFM1 KO / WT SenV (18 h)- negative direction

References

1. D. Thoresen, *et al.*, The molecular mechanism of RIG-I activation and signaling. *Immunol Rev* (2021) https:/doi.org/10.1111/imr.13022.

2. M. J. McFadden, N. S. Gokhale, S. M. Horner, Protect this house: cytosolic sensing of viruses. *Current Opinion in Virology* **22**, 36–43 (2017).

3. J. Rehwinkel, M. U. Gack, RIG-I-like receptors: their regulation and roles in RNA sensing. *Nature Reviews Immunology*, 1–15 (2020).

4. C. Chiang, M. U. Gack, Post-translational Control of Intracellular Pathogen Sensing Pathways. *Trends in Immunology* **38**, 39–52 (2017).

5. M. U. Gack, *et al.*, TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated antiviral activity. *Nature* **446**, 916–920 (2007).

6. H. Oshiumi, M. Matsumoto, S. Hatakeyama, T. Seya, Riplet/RNF135, a RING finger protein, ubiquitinates RIG-I to promote interferon-beta induction during the early phase of viral infection. *J. Biol. Chem.* **284**, 807–817 (2009).

7. H. M. Liu, *et al.*, The Mitochondrial Targeting Chaperone 14-3-3ε Regulates a RIG-I Translocon that Mediates Membrane Association and Innate Antiviral Immunity. *Cell Host & Microbe* **11**, 528–537 (2012).

8. H. Oshiumi, Recent Advances and Contradictions in the Study of the Individual Roles of Ubiquitin Ligases That Regulate RIG-I-Like Receptor-Mediated Antiviral Innate Immune Responses. *Front Immunol* **11**, 1296 (2020).

9. Y. K. Chan, M. U. Gack, A phosphomimetic-based mechanism of dengue virus to antagonize innate immunity. *Nature Immunology* **17**, 523–530 (2016).

10. W. Riedl, *et al.*, Zika Virus NS3 Mimics a Cellular 14-3-3-Binding Motif to Antagonize RIG-I- and MDA5-Mediated Innate Immunity. *Cell Host & Microbe* **26**, 493-503.e6 (2019).

11. C. Vazquez, S. M. Horner, MAVS Coordination of Antiviral Innate Immunity. *J Virol* **89**, 6974–6977 (2015).

12. E.-H. Tam, *et al.*, Role of the chaperone protein 14-3-3ε in the regulation of influenza A virus-activated beta interferon. *J Virol*, JVI0023121 (2021).

13. S. M. Horner, H. M. Liu, H. S. Park, J. Briley, M. Gale, Mitochondrial-associated endoplasmic reticulum membranes (MAM) form innate immune synapses and are targeted by hepatitis C virus. *Proc Natl Acad Sci U S A* **108**, 14590–14595 (2011).

14. S. M. Horner, C. Wilkins, S. Badil, J. Iskarpatyoti, M. G. Jr, Proteomic Analysis of Mitochondrial-Associated ER Membranes (MAM) during RNA Virus Infection

Reveals Dynamic Changes in Protein and Organelle Trafficking. *PLOS ONE* **10**,
e0117963 (2015).

15. D. C. Beachboard, *et al.*, The small GTPase RAB1B promotes antiviral innate
immunity by interacting with TNF receptor–associated factor 3 (TRAF3). *J. Biol.*
Chem., jbc.RA119.007917 (2019).

16. K. Tatsumi, *et al.*, A novel type of E3 ligase for the Ufm1 conjugation system. *J.*
Biol. Chem. **285**, 5417–5427 (2010).

17. M. Komatsu, *et al.*, A novel protein-conjugating system for Ufm1, a ubiquitin-fold
modifier. *EMBO J* **23**, 1977–1986 (2004).

18. S. H. Kang, *et al.*, Two novel ubiquitin-fold modifier 1 (Ufm1)-specific proteases,
UfSP1 and UfSP2. *J. Biol. Chem.* **282**, 5256–5262 (2007).

19. Y. Wei, X. Xu, UFMylation: A Unique & Fashionable Modification for Life. *Genomics*
Proteomics Bioinformatics **14**, 140–146 (2016).

20. J. Daniel, E. Liebau, The Ufm1 Cascade. *Cells* **3**, 627–638 (2014).

21. L. Wang, *et al.*, UFMylation of RPL26 links translocation-associated quality control
to endoplasmic reticulum protein homeostasis. *Cell Res*, 1–16 (2019).

22. Z. Wang, *et al.*, MRE11 UFMylation promotes ATM activation. *Nucleic Acids*
Research **47**, 4124–4135 (2019).

23. H. M. Yoo, *et al.*, Modification of ASC1 by UFM1 is crucial for ERα transactivation
and breast cancer development. *Mol. Cell* **56**, 261–274 (2014).

24. B. Qin, *et al.*, UFL1 promotes histone H4 ufmylation and ATM activation. *Nat*
Commun **10**, 1242 (2019).

25. I. A. Gak, *et al.*, UFMylation regulates translational homeostasis and cell cycle
progression. *bioRxiv*, 2020.02.03.931196 (2020).

26. R. DeJesus, *et al.*, Functional CRISPR screening identifies the ufmylation pathway
as a regulator of SQSTM1/p62. *eLife* **5**, e17290 (2016).

27. J. R. Liang, *et al.*, A Genome-wide ER-phagy Screen Highlights Key Roles of
Mitochondrial Metabolism and ER-Resident UFMylation. *Cell* **0** (2020).

28. C. P. Walczak, *et al.*, Ribosomal protein RPL26 is the principal target of
UFMylation. *PNAS* **116**, 1299–1308 (2019).

29. Y. Cai, N. Singh, H. Li, Essential role of Ufm1 conjugation in the hematopoietic
system. *Exp. Hematol.* **44**, 442–446 (2016).

30. J. Kulsuptrakul, R. Wang, N. L. Meyers, M. Ott, A. S. Puschnik, A genome-wide
CRISPR screen identifies UFMylation and TRAMP-like complexes as host factors
required for hepatitis A virus infection. *Cell Reports* **34** (2021).

31. D. R. Balce, *et al.*, UFMylation inhibits the proinflammatory capacity of interferon-γ–
activated macrophages. *PNAS* **118** (2021).

32. Y.-M. Loo, *et al.*, Viral and therapeutic control of IFN-β promoter stimulator 1 during
hepatitis C virus infection. *PNAS* **103**, 6001–6006 (2006).

33. R. Sumpter, *et al.*, Regulating intracellular antiviral defense and permissiveness to
hepatitis C virus RNA replication through a cellular RNA helicase, RIG-I. *J Virol* **79**,
2689–2699 (2005).

34. T. Saito, D. M. Owen, F. Jiang, J. Marcotrigiano, M. Gale, Innate immunity induced
by composition-dependent RIG-I recognition of hepatitis C virus RNA. *Nature* **454**,
523–527 (2008).

35. D. Simsek, *et al.*, The Mammalian Ribo-interactome Reveals Ribosome Functional
Diversity and Heterogeneity. *Cell* **169**, 1051-1065.e18 (2017).

36. M. J. McFadden, *et al.*, Post-transcriptional regulation of antiviral gene expression
by N6-methyladenosine. *Cell Reports* **34**, 108798 (2021).

37. A. Peisley, B. Wu, H. Xu, Z. J. Chen, S. Hur, Structural basis for ubiquitin-mediated
antiviral signal activation by RIG-I. *Nature* **509**, 110–114 (2014).

38. A. Mizutani, M. Fukuda, K. Ibata, Y. Shiraishi, K. Mikoshiba, SYNCRIP, a
Cytoplasmic Counterpart of Heterogeneous Nuclear Ribonucleoprotein R, Interacts
with Ubiquitous Synaptotagmin Isoforms*. *Journal of Biological Chemistry* **275**,
9823–9831 (2000).

39. E. Foy, *et al.*, Control of antiviral defenses through hepatitis C virus disruption of
retinoic acid-inducible gene-I signaling. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 2986–
2991 (2005).

40. E. Meylan, *et al.*, Cardif is an adaptor protein in the RIG-I antiviral pathway and is
targeted by hepatitis C virus. *Nature* **437**, 1167–1172 (2005).

41. X.-D. Li, L. Sun, R. B. Seth, G. Pineda, Z. J. Chen, Hepatitis C virus protease
NS3/4A cleaves mitochondrial antiviral signaling protein off the mitochondria to
evade innate immunity. *Proc Natl Acad Sci U S A* **102**, 17717–17722 (2005).

42. R. Lin, *et al.*, Dissociation of a MAVS/IPS-1/VISA/Cardif-IKKepsilon molecular
complex from the mitochondrial outer membrane by hepatitis C virus NS3-4A
proteolytic cleavage. *J Virol* **80**, 6072–6083 (2006).

43. K. Takahasi, *et al.*, Nonself RNA-Sensing Mechanism of RIG-I Helicase and
Activation of Antiviral Immune Responses. *Molecular Cell* **29**, 428–440 (2008).

44. H. Oshiumi, M. Miyashita, M. Matsumoto, T. Seya, A distinct role of Riplet-
mediated K63-Linked polyubiquitination of the RIG-I repressor domain in human
antiviral innate immune responses. *PLoS Pathog.* **9**, e1003533 (2013).

45. F. Hou, *et al.*, MAVS Forms Functional Prion-like Aggregates to Activate and
Propagate Antiviral Innate Immune Response. *Cell* **146**, 448–461 (2011).

46. N. Zamorano Cuervo, Q. Osseman, N. Grandvaux, Virus Infection Triggers MAVS
Polymers of Distinct Molecular Weight. *Viruses* **10** (2018).

47. W. Zeng, *et al.*, Reconstitution of the RIG-I pathway reveals a signaling role of
unanchored polyubiquitin chains in innate immunity. *Cell* **141**, 315–330 (2010).

48. X. Jiang, *et al.*, Ubiquitin-induced oligomerization of the RNA sensors RIG-I and
MDA5 activates antiviral innate immune response. *Immunity* **36**, 959–973 (2012).

49. L. Wang, *et al.*, UFMylation of RPL26 links translocation-associated quality control
to endoplasmic reticulum protein homeostasis. *Cell Research* **30**, 5–20 (2020).

50. Y. Zhang, M. Zhang, J. Wu, G. Lei, H. Li, Transcriptional regulation of the Ufm1
conjugation system in response to disturbance of the endoplasmic reticulum
homeostasis and inhibition of vesicle trafficking. *PLoS ONE* **7**, e48587 (2012).

51. C. Kwak, *et al.*, Contact-ID, a tool for profiling organelle contact sites, reveals
regulatory proteins of mitochondrial-associated membrane formation. *PNAS* **117**,
12109–12120 (2020).

52. Y. Merbl, P. Refour, H. Patel, M. Springer, M. W. Kirschner, Profiling of ubiquitin-
like modifications reveals features of mitotic control. *Cell* **152**, 1160–1172 (2013).

53. L. Wang, *et al.*, UFMylation of RPL26 links translocation-associated quality control
to endoplasmic reticulum protein homeostasis. *Cell Res.* **30**, 5–20 (2020).

54. M. Hochstrasser, Origin and function of ubiquitin-like proteins. *Nature* **458**, 422–
429 (2009).

55. K. N. Swatek, D. Komander, Ubiquitin modifications. *Cell Research* **26**, 399–422
(2016).

56. A. J. Muslin, J. W. Tanner, P. M. Allen, A. S. Shaw, Interaction of 14-3-3 with
Signaling Proteins Is Mediated by the Recognition of Phosphoserine. *Cell* **84**, 889–
897 (1996).

57. Y. Aghazadeh, V. Papadopoulos, The role of the 14-3-3 protein family in health,
disease, and drug development. *Drug Discovery Today* **21**, 278–287 (2016).

58.   A. Aitken, Post-translational modification of 14-3-3 isoforms and regulation of
cellular function. *Seminars in Cell & Developmental Biology* **22**, 673–680 (2011).

59.   J.-P. Lin, Y.-K. Fan, H. M. Liu, The 14-3-3η chaperone protein promotes antiviral
innate immunity via facilitating MDA5 oligomerization and intracellular
redistribution. *PLoS Pathog.* **15**, e1007582 (2019).

60.   K. Onomoto, K. Onoguchi, M. Yoneyama, Regulation of RIG-I-like receptor-
mediated signaling: interaction between host and viral factors. *Cell Mol Immunol*
18, 539–555 (2021).

61.   G. Liu, *et al.*, ISG15-dependent activation of the sensor MDA5 is antagonized by
the SARS-CoV-2 papain-like protease to evade host innate immunity. *Nature*
Microbiology, 1–12 (2021).

62.   M. Yoneyama, *et al.*, Shared and Unique Functions of the DExD/H-Box Helicases
RIG-I, MDA5, and LGP2 in Antiviral Innate Immunity. *The Journal of Immunology*
175, 2851–2858 (2005).

63.   C. Vazquez, C. Y. Tan, S. M. Horner, Hepatitis C Virus Infection Is Inhibited by a
Noncanonical Antiviral Signaling Pathway Targeted by NS3-NS4A. *Journal of*
Virology (2019) https:/doi.org/10.1128/JVI.00725-19 (February 8, 2022).

64.   B. Fredericksen, *et al.*, Activation of the Interferon-β Promoter During Hepatitis C
Virus RNA Replication. *Viral Immunology* **15**, 29–40 (2002).

65.   T. Saito, *et al.*, Regulation of innate antiviral defenses through a shared repressor
domain in RIG-I and LGP2. *PNAS* **104**, 582–587 (2007).

66.   F. A. Ran, *et al.*, Genome engineering using the CRISPR-Cas9 system. *Nat Protoc*
8, 2281–2308 (2013).

67.   Babraham Bioinformatics - Trim Galore! (October 12, 2021).

68.   M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* **17**, 10–12 (2011).

69.   P. J. Kersey, *et al.*, Ensembl Genomes: an integrative resource for genome-scale
data from non-vertebrate species. *Nucleic Acids Res* **40**, D91-97 (2012).

70.   A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–
21 (2013).

71.   HTSeq: Analysing high-throughput sequencing data with Python — HTSeq 0.13.5
documentation (October 12, 2021).

72.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

73. W. Huber, *et al.*, Orchestrating high-throughput genomic analysis with
Bioconductor. *Nat Methods* **12**, 115–121 (2015).

74. V. K. Mootha, *et al.*, PGC-1alpha-responsive genes involved in oxidative
phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* **34**,
267–273 (2003).

75. H. M. Liu, *et al.*, SYNCRIP (synaptotagmin-binding, cytoplasmic RNA-interacting
protein) is a host factor involved in hepatitis C virus RNA replication. *Virology* **386**,
249–256 (2009).

A

IFN-β promoter (RLU)

Mock / SenV

Vec   RIG-I   UFM1

Flag-UFM1 / 15
Flag-RIG-I / 100
SenV / 50
Total Protein

B

IFN-β promoter (RLU)

ns
**   *

Mock / SenV

Flag-UFM1: − − − − + +
WT   UFM1_KO

Flag-UFM1 / 15
UFM1 / 10
SenV / 50
GAPDH / 37

C

+SenV

IFN-β pg/mL

**

WT   UFM1_KO

UFM1 / 10
SenV / 50
Tubulin / 50

D

UFM1
GSC
① ↓
G
② ↓
UBA5
③ ↓
UFC1
UFL1 target ④ target
⑤ target   UFSP2
G
UFM1

E

+SenV

Relative IFN-β promoter activity

*** *** *** * ***

Vec UFL1 UFM1 UBA5 UFC1 UFSP2

Flag: UFL1 UFM1 UBA5 UFC1 UFSP2 / 75, 50, 25, 15
SenV / 50
GAPDH / 37

F

IFNB1

Relative Induction (SenV/Mock)

*** ***

WT   UFSP2_KO   UFM1_KO

SenV: − +   − +   − +
WT   UFM1_KO   UFSP2_KO

UFSP2 / 50
UFM1 / 25 (conjugated), 15 (unconjugated)
SenV / 50
Tubulin / 50

A

siCTRL | siUFL1

Myc-14-3-3ε: + + + + | + + + +
Flag-RIG-I: − − + + | − − + +
SenV: − + − + | − + − +

IP: Flag
- Myc — 37
- Flag — 100

Input
- Myc — 37
- Flag — 100
- UFL1 — 75
- SenV — 50
- Tubulin — 50

Relative IP: 100 | 32.6±5.0
***

B

| | WT | UFM1_KO |
Flag-RIG-I: + + | + +
Myc-MAVS: − + | − +
SenV: + + | + +

IP: Myc
- Flag — 100
- Myc — 75

Input
- Flag — 100
- Myc — 75
- UFM1 — 10
- Tubulin — 50
- SenV — 50

Relative IP: 100 | 37.6 ±21.3
*

C

| | WT | | | UFM1_KO | | |
SenV: − + + | − + +
β-Me: − − + | − − +

SDD-AGE
- MAVS

SDS-PAGE
- MAVS — 75
- UFM1 — 10
- SenV — 50
- Total Protein

Relative MAVS Agregation

WT | UFM1_KO
**

A

Mock / PAMP

IFN-β promoter (RLU)

***

Vec | UFL1

Flag

Total Protein

75 —

B

Vector / Flag-UFL1

Relative Induction (+IFN-β)

ISG15 — ns

ISG56 — ns

UFL1: − − + +
IFN-β: − + − +

Flag

Total Protein

75 —

C

Relative Ufmylation

Vector | WT | 1-212 | 1-452 | 213-794 | 453-794

** * * * ns

Flag-UFL1

Flag-UFL1

Vec | WT | 1-212 | 1-452 | 213-794 | 453-794

UFM1 conjugates

Flag

HA-UFM1

Total Protein

50 —
37 —
75 —
25 —
10 —

Nephron Epithelium Development
Nephron Development
Kidney Epithelium Development
Renal Tubule Development
Regulation of Platelet Activation
Negative Regulation of Coagulation
Mesonephric Tubule Morphogenesis
Cytosolic Ribosome
Cytosolic Large Ribosomal Subunit
Ras Guanyl Nucleotide Exchange Factor

Normalized Enrichment Score
$UFM1_{KO}$/WT (Mock)

-2.05
-2.10
-2.15
-2.20
-2.25