Signaling from the RNA sensor RIG-I is regulated by ufmylation

Daltry L. Snider[1], Moonhee Park[1], Kristen A. Murphy[1], Dia C. Beachboard[1, a], Stacy M. Horner[1,2,*]

[1] Department of Molecular Genetics & Microbiology, Duke University Medical Center, Durham, NC 27710, USA

[2] Department of Medicine, Duke University Medical Center Durham, NC 27710, USA

[a] Current address: DeSales University, Center Valley, Pennsylvania, 18034

* Correspondence: stacy.horner@duke.edu

Stacy M. Horner, Ph.D.

Duke University Medical Center

213 Research Dr., Box 3053 DUMC

Durham, NC USA 27710

Keywords

Retinoic acid-inducible gene I, RLR signaling, ubiquitin-like modifications, mitochondrial-associated ER membranes

Abstract

The RNA binding protein RIG-I is a key initiator of the antiviral innate immune response. The signaling that mediates the antiviral response downstream of RIG-I is transduced through the adaptor protein MAVS and results in the induction of type I and III interferons (IFN). This signal transduction occurs at endoplasmic reticulum (ER)-mitochondrial contact sites, to which RIG-I and other signaling proteins are recruited following their activation. RIG-I signaling is highly regulated to prevent aberrant activation of this pathway and dysregulated induction of IFN. Previously, we identified UFL1, the E3 ligase of the ubiquitin-like modifier conjugation system called ufmylation, UFL1, as one of the proteins recruited to membranes at ER-mitochondrial contact sites in response to RIG-I activation. Here, we show that UFL1, as well as the process of ufmylation, promote IFN induction in response to RIG-I activation. We find that following RNA virus infection, UFL1 is recruited to the membrane targeting protein 14-3-3ϵ, and that this complex is then recruited to activated RIG-I to promote downstream innate immune signaling. Importantly, loss of ufmylation prevents 14-3-3ϵ interaction with RIG-I, which abrogates the interaction of RIG-I with MAVS and thus downstream signal transduction that induces IFN. Our results define ufmylation as an integral regulatory component of the RIG-I signaling pathway and as a post-translational control for IFN induction.

Significance

The viral RNA sensor RIG-I initiates the antiviral innate immune response by activating a signaling cascade that induces interferon. Activation of the RIG-I signaling pathway is highly regulated to quickly mount a protective immune response while preventing dysregulation that can lead to excessive inflammation or autoimmune disorders. Here, we characterize one such mechanism of regulation. We describe that UFL1, an E3 ligase for the ubiquitin-like modifier conjugation system called ufmylation, is important to promote RIG-I signaling. Using molecular approaches, we show that ufmylation promotes RIG-I interaction with the membrane targeting protein 14-3-3ϵ. As such, ufmylation positively regulates RIG-I recruitment to its signaling adaptor proteins MAVS for induction of interferon in response to RNA virus infection.

Abstract

The RNA binding protein RIG-I is a key initiator of the antiviral innate immune response.

The signaling that mediates the antiviral response downstream of RIG-I is transduced

through the adaptor protein MAVS and results in the induction of type I and III interferons

(IFN). This signal transduction occurs at endoplasmic reticulum (ER)-mitochondrial

contact sites, to which RIG-I and other signaling proteins are recruited following their

activation. RIG-I signaling is highly regulated to prevent aberrant activation of this pathway

and dysregulated induction of IFN. Previously, we identified UFL1, the E3 ligase of the

ubiquitin-like modifier conjugation system called ufmylation, UFL1, as one of the proteins

recruited to membranes at ER-mitochondrial contact sites in response to RIG-I activation.

Here, we show that UFL1, as well as the process of ufmylation, promote IFN induction in

response to RIG-I activation. We find that following RNA virus infection, UFL1 is recruited

to the membrane targeting protein 14-3-3ε, and that this complex is then recruited to

activated RIG-I to promote downstream innate immune signaling. Importantly, loss of

ufmylation prevents 14-3-3ε interaction with RIG-I, which abrogates the interaction of RIG-

I with MAVS and thus downstream signal transduction that induces IFN. Our results define

ufmylation as an integral regulatory component of the RIG-I signaling pathway and as a

post-translational control for IFN induction.

Significance Statement

The viral RNA sensor RIG-I initiates the antiviral innate immune response by activating a

signaling cascade that induces interferon. Activation of the RIG-I signaling pathway is

highly regulated to quickly mount a protective immune response while preventing

dysregulation that can lead to excessive inflammation or autoimmune disorders. Here, we

characterize one such mechanism of regulation. We describe that UFL1, an E3 ligase for

the ubiquitin-like modifier conjugation system called ufmylation, is important to promote

RIG-I signaling. Using molecular approaches, we show that ufmylation promotes RIG-I

interaction with the membrane targeting protein 14-3-3ε. As such, ufmylation positively

regulates RIG-I recruitment to its signaling adaptor proteins MAVS for induction of

interferon in response to RNA virus infection.

Introduction

Detection of RNA virus infection is initiated by cellular sensors such as RIG-I. RIG-I is a

pattern recognition receptor that detects unique features of viral RNA that are generally

absent in cellular RNA, referred to as pathogen-associated molecular patterns (PAMPs) (1). Sensing of viral RNA PAMPs triggers RIG-I activation and induces a downstream signaling cascade that ultimately results in transcriptional induction of type I and type III interferons (IFN) and the antiviral response (2, 3). The RIG-I signaling cascade is carefully regulated by multiple mechanisms, including post-translational modifications that influence specific protein-protein interactions that can result in changes in protein localization to mediated signaling (3, 4). For example, following sensing of RNA PAMPs, RIG-I undergoes K63-linked polyubiquitination in order to transition to its fully active conformation, which promotes its interaction with the molecular trafficking protein 14-3-3ε (5–8). 14-3-3ε facilitates the recruitment of activated RIG-I from the cytosol to intracellular membranes where it interacts with MAVS (7, 9, 10), which assembles other RIG-I pathway members to transduce the signals that induce IFN (7, 11). Importantly, many RNA viruses, including influenza A virus and some flaviviruses (dengue virus, Zika virus, and West Nile virus), prevent the interaction of RIG-I with 14-3-3ε to limit IFN induction and evade the antiviral response (9, 10, 12).

In addition to RIG-I, a number of signaling proteins must be recruited to MAVS in order to propagate downstream IFN induction. Previously, we identified proteins that move to MAVS signaling sites at mitochondrial-associated endoplasmic reticulum (ER) membranes (MAM) during RNA virus infection (13, 14). These proteins likely aid in spatial organization of RIG-I pathway proteins during viral infection and include the GTPase RAB1B, which plays a role in recruiting TRAF3 to MAVS (15). In addition to RAB1B, we identified other proteins recruited to the MAM upon RIG-I signaling activation, one of which was UFL1 (referred to in our previous publication as KIAA0776) (14). UFL1 is an E3 ligase for UFM1, which is a ubiquitin-like modification of 85 amino acids. The process of ufmylation conjugates UFM1 covalently to lysine residues of target proteins through a process called ufmylation, which is similar to ubiquitination in that it also uses an E1, E2, and E3 ligase conjugation system (UBA5, UFC1, and UFL1; see Figure 2D). UFM1 is removed by the UFSP2 protease (16–20). The consequence of UFM1 addition to proteins is not fully understood, but the literature supports the idea that it can promote protein-protein interactions to regulate a number of biological processes (21–31). Here, we uncover a role for ufmylation in RIG-I activation. We found that the cellular proteins that catalyze ufmylation all promote RIG-I-mediated induction of IFN. Interestingly, we found that UFL1 interacts with both RIG-I and the molecular trafficking protein 14-3-3ε following

RNA virus infection. Further, similar to RIG-I, UFL1 is recruited to intracellular membranes

following RNA virus infection. Importantly, loss of ufmylation prevents the interaction of

14-3-3ε with RIG-I, which results in decreased MAVS activation and IFN induction in

response to RNA virus infection. Thus, ufmylation can regulate RIG-I activation and

downstream signaling of the intracellular innate immune system.

Results

The ufmylation activity of UFL1 promotes RIG-I signaling. Having found that the E3

ligase of ufmylation UFL1 is recruited to MAVS signaling sites at the MAM in response to

RIG-I signaling (14), we wanted to determine if UFL1 regulates RIG-I signaling. To test

this, we measured induction of the IFN-β promoter following UFL1 overexpression using

an IFN-β promoter luciferase reporter assay (32) and found that UFL1 increased activation

of the IFN-β promoter, similar to that of RIG-I expression, in a dose-dependent fashion in

response to infection with Sendai virus (SenV) (Figure 1A). SenV is a murine

paramyxovirus that specifically activates RIG-I (33). In support of UFL1 enhancing RIG-I

signaling specifically, exogenous expression of UFL1 also increased IFN-β promoter

activity in response to transfection of 293T cells with a known RIG-I immunostimulatory

RNA from hepatitis C virus (PAMP; Figure 1B) (34). However, UFL1 overexpression in

293T cells did not lead to increased induction of IFN-stimulated genes (ISG), such as

ISG56 or *ISG15,* in response to exogenous IFN-β treatment, indicating that UFL1 primarily

regulates IFN induction and not the IFN response (Figure 1C). Next, we depleted UFL1

by siRNA in two different cell types and measured SenV-induced activation of the RIG-I

pathway. Depletion of UFL1 in 293T cells resulted in decreased phosphorylation of IRF3,

a transcription factor for both type I and III IFNs, while exogenous expression of an siRNA-

resistant UFL1 restored SenV-mediated IRF3 phosphorylation (Figure 1D). Depletion of

UFL1 in primary neonatal human dermal fibroblasts (NHDFs) also reduced the SenV-

mediated induction of both *IFNB1* and *IFNL1* transcripts, as measured by RT-qPCR

(Figure 1E), as well as the production of IFN-β protein, as measured by an enzyme-linked

immunosorbent assay (ELISA) (Figure 1F).

To define the domains of UFL1 that regulate RIG-I signaling, we expressed a series of

previously described UFL1 truncation mutants (diagrammed in Figure 1H) and measured

SenV-mediated activation of the IFN-β promoter in a luciferase reporter assay (16). The

ability of UFL1 to transfer UFM1 to a target protein has been suggested to require the first

212 amino acids of the protein, as this domain interacts with the E2 ligase for ufmylation,

UFC1 (16). The wild-type (WT) UFL1 (aa 1-794), as well as the C-terminal deleted mutants

of UFL1, aa 1-212 and aa 1-452, which all have reported ufmylation activity (16),

stimulated SenV-medicated induction of the IFN-β promoter (Figure 1G). Interestingly, the

N-terminal deleted mutant of UFL1 aa 213-794, that does not have reported ufmylation

activity, also induced signaling, while the N-terminal deleted UFL1 mutant aa 453-794 did

not (Figure 1G). However, our analysis of global UFM1 conjugates by these UFL1

constructs revealed that while UFL1 WT, aa 1-212, and aa 1-452 all retain full ufmylation

activity, aa 213-794 of UFL1 retain approximately 40% ufmylation activity, while aa 453-

794 of UFL1 retain only about 20% activity (Figure 1H; Figure S1). Thus, taken together,

this reveals that the ufmylation activity of UFL1 is required to promote RIG-I signaling that

results in induction of IFN.

The ufmylation machinery proteins positively regulate RIG-I signaling. Having

determined that the ufmylation activity of UFL1 is important for its role in RIG-I signaling,

we hypothesized that UFM1 and the proteins required for UFM1 conjugation would also

be required to promote this signaling. Similar to our results with UFL1, overexpression of

UFM1 increased SenV-mediated activation of the IFN-β promoter in a dose-dependent

fashion (Figure 2A). Conversely, the activation of the IFN-β promoter in response to SenV

was significantly abrogated in 293T cells in which UFM1 was deleted by CRISPR/Cas9,

as compared to WT 293T cells (Figure 2B). Importantly, this signaling was restored upon

exogenous expression of UFM1 (Figure 2B). The absence of UFM1 expression also

prevented the induction of IFN-β protein in response to SenV infection, as measured by

ELISA (Figure 2C). The process of ufmylation has 5 steps (Figure 2D). First, UFM1 is

processed to expose the terminal glycine residue. Then, this mature UFM1 is added to the

target protein by the actions of UBA5, which acts as an E1 ligase for UFM1; UFC1, the E2

ligase; and UFL1, the E3 ligase (19). Finally, the UFSP2 protease removes UFM1, which

enables recycling of mature UFM1 (18). We found that exogenous expression of each of

the proteins involved in UFM1 conjugation, including the UFSP2 protease, positively

regulated SenV-mediated induction of the IFN-β promoter (Figure 2E). These results

reveal that the proteins that catalyze ufmylation and the UFM1 modification itself promote

RIG-I signaling.

UFM1 is required for the RIG-I-driven transcriptional response. After establishing that ufmylation promotes RIG-I activation, and in turn IFN expression, we next broadly measured the impact of ufmylation upon the transcriptional response to RIG-I signaling. Using RNA-sequencing, we analyzed gene expression in either WT or UFM1 KO 293T cells, following mock or SenV infection (Table S1.1; Table S1.2). Gene set enrichment analysis (Table S2.1; Table S2.2) of the transcripts significantly reduced (adjusted P<0.01) by UFM1 KO in the absence of viral infection revealed previously described pathways regulated by ufmylation such as cytosolic ribosomes, ribosome assembly, and hematopoiesis (Figure S2A; Table S2.1) (21, 28, 29, 35). Following viral infection, the top 10 gene categories negatively impacted by UFM1 KO, with a darker red color indicating more downregulation, were all related to the antiviral response, such as response to type I IFN and defense against virus (Figure 3A; Table S2.2). Indeed, of the top 50 most downregulated pathways impacted by UFM1 KO during infection, the majority were related to innate immune signaling or viral replication (Table S2.2), while upregulated gene categories were more diverse (Table S2.3; Table S2.4). Of the genes differentially expressed during UFM1 KO in response to SenV (adjusted P<0.01), the majority are downregulated (Figure 3B). Indeed, these downregulated genes included *IFNB1* and *IFNL1*, as well as other known ISGs (in red) (36) (Figure 3B; Figure 3C). These data are consistent with a model in which ufmylation-mediated regulation of IFN induction has broad consequences on genes induced by the IFN response.

UFL1 is recruited to intracellular membranes and interacts with 14-3-3ε and RIG-I during RNA virus infection. Following the binding of RIG-I to non-self RNA, it interacts with several host proteins to facilitate its activation, localization to the MAM, and interaction with MAVS. These proteins include the E3 ligases for K63-linked ubiquitin TRIM25 and Riplet (5, 6, 37), as well as the molecular trafficking protein 14-3-3ε. In particular, 14-3-3ε is required for RIG-I recruitment from the cytosol to MAVS signaling sites at intracellular membranes (5–7, 13); however, the mechanism underlying how 14-3-3ε selects RIG-I as cargo has yet to be elucidated. Using a subcellular membrane fractionation assay (38), we confirmed that UFL1 increases its association with intracellular membranes in response to SenV, similar to RIG-I (Figure 4A; compare fraction #1, which has Cox-I and no GAPDH, with fractions #6-8, which are enriched for the cytosolic protein GAPDH) (7, 12). This finding is consistent with our previous report that UFL1 is recruited to the MAM in response to either SenV or hepatitis C virus replication (14), suggesting that UFL1

recruitment occurs prior to MAVS activation, as MAVS is cleaved by the HCV NS3-NS4A

protease (39–42). As the recruitment of RIG-I to intracellular membranes is known to

require 14-3-3ε, and, as both UFL1 and UFM1 have been shown to interact with 14-3-3ε

(16), we hypothesized that UFL1 may interact with 14-3-3ε to promote the IFN induction

that we had observed in response to RNA virus infection. Thus, we first determined if the

interaction of UFL1 with 14-3-3ε is increased in response to RIG-I activation by SenV by

performing co-immunoprecipitation. We found that Myc-14-3-3ε did co-immunoprecipitate

with Flag-UFL1, as reported previously (16), and that this interaction was increased by

SenV (Figure 4B). Interestingly, the interaction of UFL1 with RIG-I also increased following

SenV, both upon over-expression and at the level of the endogenous proteins (Figure 4C;

Figure 4D). As RIG-I undergoes a series of modifications to become fully active (1, 4), we

next used a panel of RIG-I mutants to define which stage of RIG-I activation promotes

interaction with UFL1. These mutations prevent the distinct steps of RIG-I activation such

as RIG-I binding to RNA (K888/907A), interacting with TRIM25 (T55I), or ubiquitination by

Riplet and TRIM25 (K172/788R) (5, 43, 44). The interaction of UFL1 with RIG-I was

significantly impaired by each of these mutations, suggesting that UFL1 regulates RIG-I

function after it binds RNA and becomes ubiquitinated (Figure 4E). As this is the same

step of activation at which 14-3-3ε binds to RIG-I to promote its translocation to

intracellular membranes (7), this suggests that RNA virus infection increases the

interaction of 14-3-3ε with UFL1, which then interacts with activated, K63-ubiquitinated

RIG-I.

UFL1 interaction with RIG-I requires 14-3-3ε and UFM1. Having determined that UFL1

interacts with both activated RIG-I and 14-3-3ε following RNA virus infection, we next

defined the dynamics of this complex formation by testing two distinct models. In the first

model, UFL1 would interact first with activated RIG-I, induce its ufmylation, and then the

UFL1-RIG-I complex would interact with 14-3-3ε. In this model, depletion of 14-3-3ε or

loss of UFM1 would not be expected to change the interaction of UFL1 with RIG-I. In the

second model, UFL1 would interact first with 14-3-3ε and induces its ufmylation, or that of

another associated protein, and then the UFL1-14-3-3ε complex would interact with

activated RIG-I. In this second model, depletion of 14-3-3ε would be expected to prevent

UFL1 interaction with RIG-I, and loss of ufmylation would limit UFL1 interaction with RIG-

I but would not affect UFL1 interaction with 14-3-3ε. To elucidate these possibilities, first,

we used co-immunoprecipitation to measure the interaction of exogenously expressed
Flag-UFL1 and HA-RIG-I in SenV-infected 293T lysates that had been depleted of 14-3-
3ε or CTRL by siRNA. This revealed that formation of the SenV-activated RIG-I-UFL1
complex requires 14-3-3ε (Figure 5A). Next, we tested if ufmylation was required for
formation of the SenV-activated RIG-I-UFL1 complex by measuring this interaction in WT
or UFM1 KO 293T cells. We found that UFM1 was required for SenV-activated RIG-I-
UFL1 complex (Figure 5B). The results of these two experiments reveal that both 14-3-3ε
and UFM1 are required for UFL1 to interact with RIG-I, supporting the second model of
complex formation in which UFL1 interacts first with 14-3-3ε and catalyzes its ufmylation,
and then this complex associates with RIG-I. In support of this, we found that UFM1 was
not required for UFL1 to interact with 14-3-3ε (Figure 5C). Together, these data indicate
that ufmylation promotes the interaction of UFL1 with 14-3-3ε and activated RIG-I.
278
Ufmylation promotes RIG-I interaction with 14-3-3ε for MAVS activation. Having
found that that UFL1 requires 14-3-3ε to interact with activated RIG-I, we next tested if
UFL1 is required for the interaction of 14-3-3ε with RIG-I, which is essential for activated
RIG-I to translocate from the cytosol to intracellular membranes for interaction with MAVS
(7). We performed a co-immunoprecipitation of Flag-RIG-I and Myc-14-3-3ε from 293T
cells and found that this SenV-mediated interaction was significantly decreased upon
UFL1 depletion (Figure 6A). In addition, loss of UFM1 expression also decreased the
SenV-induced interaction of RIG-I with 14-3-3ε (Figure 6B). Importantly, we also found
that UFM1 is required for the SenV-induced interaction of RIG-I with MAVS (Figure 6C)
and MAVS higher-order oligomerization, which is a hallmark of MAVS activation (45, 46)
(Figure 6D). In summary, these data reveal that UFL1 and UFM1 are required for the RIG-
I interaction with 14-3-3ε, for interaction with MAVS, and for MAVS activation by
oligomerization.
292
293   **Discussion**
294
Regulation of RIG-I activation and downstream signaling is essential for proper induction
and termination of IFN. Here, we show that both UFL1 and the process of ufmylation
promote RIG-I pathway signaling that leads to IFN induction, uncovering an important step
in the activation of the RIG-I pathway. RIG-I activation occurs upon RNA binding. Then,

RIG-I undergoes ATP hydrolysis. and interaction with K63-linked polyubiquitin chains,
both covalently and non-covalently (5, 44, 47), which promotes formation of a RIG-I
tetramer (48). This polyubiquitinated, activated RIG-I oligomer then interacts with the
membrane trafficking protein 14-3-3ε for translocation to MAVS at ER-mitochondrial
contact sites (7). We found that UFL1 is recruited to 14-3-3ε following RNA virus infection
and that ufmylation facilitates the interaction between 14-3-3ε and activated RIG-I.
Importantly, this results in increased interaction of RIG-I with MAVS and MAVS
oligomerization, ultimately promoting the downstream signal transduction which produces
IFN.
308
Ufmylation is emerging as a post-translational modification that regulates diverse
biological processes, including DNA repair, ER homeostasis, and even the replication of
hepatitis A virus (21, 22, 24, 27, 28, 30, 35, 49). In these cases, UFL1, along with the other
members of the ufmylation cascade, induce ufmylation of a target protein important for
regulating these processes. For example, both MRE11 and histone H4 are ufmylated by
UFL1 in the nucleus in response to DNA damage resulting in activation the key DNA repair
kinase ATM (22, 24). UFL1 can also act at the ER, where it plays a role in ER protein
quality control, where it ufmylates specific proteins, including ribosomal proteins RPL26,
to induce lysosomal degradation of stalled peptides and/or the ER and prevent the
unfolded protein response (27, 28, 49, 50). Interestingly, hepatitis A virus translation,
which occurs in association with the ER, also requires ufmylation of RPL26 (30).
Therefore, ufmylation can regulate several aspects of translation. It is possible that
ufmylation regulates translation of certain mRNAs important for RIG-I signaling and
subsequent IFN induction. However, we identified a role for ufmylation in regulating the
interaction of RIG-I with 14-3-3ε, one of the earliest known steps of RIG-I signaling,
strongly supporting a mechanism in which following RIG-I activation, ufmylation is
controlling this specific protein-protein interaction. The mechanisms by which the process
of UFM1 addition regulates interactions between proteins or alters other aspects of protein
function are largely unknown. Indeed, we found that UFSP2, the protease that removes
UFM1 from proteins (18), promoted SenV-mediated IFN induction (Figure 2D), suggesting
that we do not yet have a full grasp on the ufmylation process. It is possible that the
dynamic process of ufmylation or the enhanced formation of mature UFM1 following
deconjugation from targets or promote RIG-I signaling independent of deconjugation
activity. Indeed, in support of this idea, others have shown that UFSP2 in myeloid cells is

required for influenza virus resistance in mice (31). It is also possible that UFSP2 acts on

other members of the RIG-I pathway to alter their function. Future studies to define how

the process of ufmylation regulates this and other aspects of the antiviral innate immune

response will be of great interest as they may shed light broadly on how ufmylation

regulates diverse cell biological processes that alter cellular signaling.

The mechanisms underlying how cytoplasmic UFL1 is recruited to its protein targets that

reside in different subcellular compartments are not fully known. For example, we found

that RIG-I activation induces UFL1 translocation to intracellular membranes (Figure 3A),

and while we know that UFL1 is recruited to the MAM during infection, the mechanism by

which UFL1 becomes membrane-associated remains unknown (14). DDRGK1 (UFBP1)

may facilitate UFL1 targeting to the MAM, as DDRGK1 is localized to mitochondrial-ER

contact sites (16, 51) and in some cases it is required for UFL1 recruitment to membranes

(27, 28). Thus, both DDRGK1 and mitochondrial-ER contact sites could function as a

regulatory hub that aids in the recruitment of UFL1 and RIG-I pathway signaling proteins.

Interestingly, RAB1B, a GTPase that we found is recruited to the MAM and important for

RIG-I signaling (14, 15) is ufmylated (52, 53), which reveals that ufmylation likely regulates

a number of RIG-I pathway signaling proteins. As UFL1 contains no functional domains

common to other E3 ligases that might allow one to predict how its targets are selected

(16, 54, 55), defining the signals and features that control UFL1 localization, as well as

the target proteins ufmylated in response to RIG-I activation, will undoubtedly reveal clues

into how the process of ufmylation is activated and how specific targets are selected.

Our work revealed that 14-3-3ε required ufmylation to interact with activated RIG-I. The

details underlying how 14-3-3ε interacts with activated RIG-I have not been fully

elucidated, as it does not occur through the known phosphorylated amino acids on RIG-I,

the typical recruitment signal of the 14-3-3 family of proteins (7, 56, 57). Interestingly,

others have shown that 14-3-3ε interacts with UFM1 and other members of the ufmylation

pathway (16). Thus, taken together with our results, this suggests that ufmylation of 14-3-

3ε or a 14-3-3ε-associated protein promotes the interaction between activated RIG-I and

14-3-3ε. In fact, a number of 14-3-3 family proteins are post-translationally modified by

phosphorylation, acetylation, and oxidation (58). Therefore, post-translational modification

of 14-3-3ε by ufmylation could broadly define how cargo proteins, including RIG-I, are

selected. Indeed, this mechanism could be shared with other RNA virus sensing

pathways, such as the RIG-I-like-receptor MDA5, which also interacts with a 14-3-3
protein, 14-3-3η, by an unknown mechanism (59). Thus, ufmylation may broadly influence
how 14-3-3 proteins or other host proteins interact with each other to regulate the
intracellular innate immune response.

Overall, this work lays the groundwork for future studies to define how ufmylation of
antiviral innate immune signaling proteins regulates their function and how specific
signaling pathways are differentially activated through ufmylation. In addition, our work
adds ufmylation to the growing list of ubiquitin-like and other modifications that regulate
the intracellular innate immune response, including ISGylation, SUMOylation, FATylation,
acetylation, phosphorylation, and others (4, 60, 61) broadening our understanding of how
RIG-I signaling is activated and rapidly controlled by post-translational modifications in
response to infection, leading to greater knowledge of the exquisite regulation of these
pathways.

Materials and Methods

Cell lines, viruses, and treatments. Neonatal human dermal fibroblast (NHDF) cells and
embryonic kidney 293T cells were grown in Dulbecco's modification of Eagle's medium
(DMEM; Mediatech) supplemented with 10% fetal bovine serum (Thermo Fisher
Scientific), 1X minimum essential medium non-essential amino acids (Thermo Fisher
Scientific), and 25 mM HEPES (Thermo Fisher Scientific) (cDMEM). 293T (CRL-3216)
were obtained from American Type Culture Collection (ATCC), NHDF cells (CC-2509)
were obtained from Lonza. All cell lines were verified as mycoplasma free by the LookOut
Mycoplasma PCR detection kit (Sigma). SenV Cantell strain was obtained from Charles
River Laboratories and used at 200 hemagglutination units/mL (HAU). SenV infections
were performed in serum-free media (30 minutes to 1 hour), after which complete media
was replenished. IFN-β (PBL Assay Science) was added to cells at a concentration of 50
units/mL in cDMEM for 18 hours.

Plasmids. The following plasmids have been previously described: pEF-TAK-Flag, pEF-
BOS-Flag-RIG-I (62), pIFN-β-luc (63), pCMV-Renilla (Promega), pX459 (Addgene
Plasmid #62988), pEF-BOS-Flag-RIG-I T55I (64), pEF-TAK-Myc-MAVS (32).
pLJM1_Flag-UFM1 was a gift from Dr. Craig McCormick at Dalhouise University. The

following plasmids were generated by insertion of PCR-amplified fragments into the NotI-
to-PmeI digested pEF-TAK-Flag using InFusion cloning (Clontech), all primers are
denoted in Table 1: pEF-TAK-Flag-UFL1 (GenBank: BC036379; GeneID: 23376), pEF-
TAK-Flag-UBA5 (NM_024818.6), pEF-TAK-Flag-UFC1 (NM_016406.4), pEF-TAK-Flag-
UFSP2 (NM_018359.5), pEF-TAK-Flag-UFL1 1-212, pEF-TAK-Flag-UFL1 1-452, pEF-
TAK-Flag-UFL1 213-794, and pEF-TAK-Flag-UFL1 453-794. Both pEF-TAK-Myc-14-3-3ε
and pEF-TAK-Myc-UFL1 were generated by insertion of PCR-amplified fragments into the
AgeI-NotI digested pEF-TAK-Myc (pEF-TAK-Myc-MAVS) by InFusion. The pEF-TAK-HA
vector was generated by PCR to replace Flag with HA, and pEF-TAK-HA-RIG-I was
generated by insertion of a PCR-amplified fragment into the NotI-AgeI digested pEF-TAK-
HA vector. The following plasmids were generated by site-directed mutagenesis (see
Table 1): pEF-TAK-Flag-UFL1^{siR}, pEF-BOS-Flag-RIG-I K888/907A, and pEF-BOS-Flag-
RIG-I K172/788R. To generate the CRISPR guide RNA plasmids px459-UFM1-E2 and
px459-UFM1-B, sgRNA oligonucleotides were annealed and inserted into the BbsI-
digested pX459 (30, 65). The plasmid sequences were verified by DNA sequencing and
oligonucleotide sequences are available upon request.

Generation of RNA PAMP. Annealed oligonucleotides containing the sequence of the
HCV 5'ppp poly-U/UC region (34) were *in vitro* transcribed using the MEGAshortscript T7
transcription kit (Ambion) followed by ethanol precipitation, with the resulting RNA
resuspended at 1 µg/µL.

Transfection. DNA transfections were performed using FuGENE6 (Promega) or TransIT-
LT1 (Mirus Bio). RNA PAMP transfections were done using the TransIT-mRNA
Transfection kit (Mirus Bio). The siRNA transfections were done using Lipofectamine
RNAiMax (Invitrogen). siRNAs directed against 14-3-3ε (Dharmacon-L-017302-02-0005),
UFL1 (Qiagen-SI04371318) or non-targeting AllStars negative control siRNA (Qiagen-
1027280) were transfected into 293T cells (25 pmol of siRNA; final concentration of 0.0125
µM) or NHDF cells (250 pmol of siRNA; final concentration of 0.25 µM). Media was
changed 4-24 hours post-transfection, and cells were incubated for 36-48 h post-
transfection prior to each experimental treatment. IFN-β-promoter luciferase assays were
performed as previously described at 18-24 hours post treatment and normalized to the
Renilla luciferase transfection control (33).

ELISA. IFN-β ELISAs were performed using Human IFN-beta DuoSet (R&D Systems)

with supernatants collected from cultured cells.

Generation of KO cell lines. UFM1 KO 293T cells were generated by CRISPR/Cas9,

using two guides targeting exon 2 and 3, similar to others, as we have done previously

(15, 30).

RNA analysis. Total cellular RNA was extracted using the RNeasy Plus mini kit (Qiagen).

RNA was then reverse transcribed using the iScript cDNA synthesis kit (BioRad), as per

the manufacturer's instructions. The resulting cDNA was diluted 1:3 in ddH2O. RT-qPCR

was performed in triplicate using the Power SYBR Green PCR master mix (Thermo-

Fisher) and QuantStudio 6 Flex RT-PCR system. Oligonucleotide sequences for qPCR

are available upon request.

RNA-seq. WT and UFM1 KO 293T cells were mock or SenV-infected (18 h) and

harvested in biological duplicate, followed by total RNA extraction via TRIzol reagent

(Thermo Fisher Scientific). Sequencing libraries were prepared using the KAPA Stranded

mRNA-Seq Kit (Roche) and sequenced on an Illumina Novaseq 6000 with 50 bp paired-

end reads (>20 million reads per sample) in an S1 flow cell by the Duke University Center

for Genomic and Computational Biology.

RNA-seq data was processed using the TrimGalore toolkit (66) which employs Cutadapt

(67) to trim low-quality bases and Illumina sequencing adapters from the 3' end of the

reads. Only reads that were 20nt or longer after trimming were kept for further analysis.

Reads were mapped to the GRCh38v93 version of the human genome and transcriptome

(68) using the STAR RNA-seq alignment tool (69). Reads were kept for subsequent

analysis if they mapped to a single genomic location. Gene counts were compiled using

the HTSeq tool (70). Only genes that had at least 10 reads in any given library were used

in subsequent analysis. Normalization and differential expression was carried out using

the DESeq2 (71) Bioconductor (72) package with the R statistical programming

environment. The false discovery rate was calculated to control for multiple hypothesis

testing. Gene set enrichment analysis (73) was performed to identify gene ontology terms

and pathways associated with altered gene expression for each of the comparisons

performed. All RNA-seq data are deposited in the GEO database under GSE186287.

Immunoblotting. Cells were lysed in a modified radioimmunoprecipitation assay (RIPA) buffer (10 mM Tris [pH 7.5], 150 mM NaCl, 0.5% sodium deoxycholate, and 1% Triton X-100) supplemented with protease inhibitor cocktail (Sigma) and Halt Phosphatase Inhibitor (Thermo-Fisher), and post-nuclear lysates were isolated by centrifugation. Quantified protein (between 5 -15 µg) was resolved by SDS/PAGE, transferred to nitrocellulose or polyvinylidene difluoride (PVDF) membranes in a 25 mM Tris-192 mM glycine-0.01% SDS buffer. Membranes were stained with Revert 700 total protein stain (LI-COR Biosciences) and then blocked in 3% BSA in Tris-buffered saline containing 0.01% Tween-20 (TBS-T). After washing with PBS-T or TBS-T (for phosphoproteins) buffer, following incubation with primary antibodies, membranes were incubated with species-specific horseradish peroxidase-conjugated antibodies (Jackson ImmunoResearch, 1:5000) or fluorescent secondaries (LI-COR Biosciences), followed by treatment of the membrane with Clarity Western ECL substrate (BioRad) and imaging on a LICOR Odyssey FC. The following antibodies were used for immunoblotting: R-anti-SenV (MBL, 1:1000), M-anti-Tubulin (Sigma, 1:1000), R-anti-GAPDH (Cell Signaling Technology, 1:1000), R-anti-p-IRF3 (Cell Signaling Technology, 1:1000), R-anti-IRF3 (Cell Signaling Technology, 1:1000), R-anti-UFL1 (Novus Biologicals, 1:1000), R-anti-UFM1 (Abcam, 1:1000), anti-RIG-I (M-AdipoGen, R-Abcam, 1:1000), R-anti-14-3-3ε (Cell Signaling Technology, 1:1000), M-anti-Flag M2 (Sigma, 1:1000), anti-Flag-HRP (Sigma, 1:1000-1:5000), R-anti-Flag (Sigma, 1:1000), anti-HA (M- and R-Sigma, 1:1000), and anti-Myc (M-Santa Cruz or R-Cell Signaling Technology, 1:1000).

Immunoprecipitation. Cells were lysed in RIPA buffer with or without 10% glycerol. Quantified protein (between 100-500 µg) was incubated with protein-specific, isotype control antibody (R-Cell Signaling Technology or M-Thermo Fisher), or anti-Flag M2 magnetic beads (Sigma), in lysis buffer either at room temperature for 2 h or at 4ºC overnight with head over tail rotation. The lysate/antibody mixture was then incubated with Protein G Dynabeads (Invitrogen) for 1 h. Beads were washed 3X in PBS or RIPA buffer and eluted in 2X Laemmli Buffer (BioRad) with or without 5% 2-Mercaptoethanol at 95ºC for 5 min. Proteins were resolved by SDS/PAGE and immunoblotting, as above.

Subcellular membrane fractionation. Membrane fractionation was performed as previously described (7, 12, 38, 74). Cells were lysed in hypotonic buffer (10 mM Tris-HCL (pH 7.5), 10 mM KCl, and 5 mM $MgCl_2$ supplemented with protease inhibitor cocktail) for

10 minutes on ice followed by 20 passages through a 20-guage needle. Nuclei and

unbroken cells were removed by centrifugation at 1000xg for 5 min at 4ºC. The resulting

supernatants were mixed thoroughly with 72% sucrose and overlayed with 55% sucrose,

followed by 10% sucrose, all in low-salt buffer (2 nM EDTA, 20 nM HEPES (pH 8.0), 150

mM NaCl, 0.1% SDS, 1% Triton X-100). The gradients were subjected to centrifugation at

38,000 RPM in a Beckman SW41 Ti Rotor for 14 h at 4ºC. 1 mL fractions were collected

using a BioComp piston gradient fractionator and resulting fractions were divided in half

and mixed with 2 parts 100% methanol and precipitated overnight at -80ºC. Protein pellets

were collected by centrifugation and resuspended in 2X Laemmli buffer and heated for 5

min at 95ºC for immunoblot analysis. 10% pre-fractionated cells from each condition were

collected as the input.

Semi-denaturing detergent agarose gel electrophoresis. SDD-AGE was performed as

described (45, 46). Briefly, crude mitochondria (P5 fraction) were isolated from an equal

number of WT or UFM1 KO 293T cells that were mock or SenV infected (12 h),

resuspended in hypotonic buffer (10 mM Tris, pH 7.5, 10 mM KCl, 1.5 mM $MgCl_2$, and 0.5

mM EDTA). Resulting samples were split and 2X SDD-AGE sample buffer (0.5X TBE,

10% glycerol, 2% SDS, 0.2 mM Bromophenol Blue) buffer with or without 5% 2-

Mercaptoethanol was added, and samples were loaded onto a vertical 1.5% agarose gel.

Electrophoresis was performed with a constant voltage of 70 V at 4 °C in SDD-AGE

running buffer (1X TBE and 0.1% SDS). Gels were transferred onto a nitrocellulose

membrane overnight on ice at 25 V. Membranes were fixed in 0.25% glutaraldehyde in

PBS and immunoblotting was performed as usual. 15% of the SDD-AGE samples were

reserved for input.

Quantification of immunoblots. Immunoblots imaged using the LICOR Odyssey FC

were quantified by ImageStudio software, and raw values were normalized to relevant

controls for each antibody. Phosphoprotein values were normalized to Tubulin and

displayed as the percentage of signal from WT. Relative membrane association of UFL1

was quantified as the ratio of UFL1 to Cox-1 in fraction 1 normalized to total protein levels

of UFL1 in the input and displayed as the percentage of UFL1 membrane association

normalized to mock values.

Statistical analysis. Student's unpaired t-test, one-way ANOVA, or two-way ANOVA
were implemented for statistical analysis of the data followed by appropriate post-hoc test
(as indicated) using GraphPad Prism software. Graphed values are presented as mean ±
SD or SEM (n = 3 or as indicated); *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001.

540
541 **Acknowledgments**
542

We thank colleagues who provided reagents (see Methods), the Duke Functional
Genomics Core Facility, the Duke University Center for Genomic and Computational
Biology, the Duke Genomic Analysis and Bioinformatics Shared Resource and Dr. Wei
Chen for performing RNA-seq analysis, Dr. Madhuvanthi Vijayan and Boyoung Michelle
Kim for generation and initial characterization of plasmids, Dr. Matthew Thompson for
input in RNA-seq sample preparation and data processing, and Horner lab members for
useful discussion. This work was supported by funds from Burroughs Wellcome Fund,
National Institutes of Health: R21AI144380, R01AI155512, T32-CA009111, and an
American Cancer Society Postdoctoral fellowship 131321-PF-17-188-01-MPC.

Author Contributions: Conceptualization: D.L.S., D.C.B., and S.M.H.; Investigation:
D.L.S, D.C.B., and M.P.; Formal analysis: D.L.S., D.C.B., K.A.M., and S.M.H.; Writing –
original draft: D.L.S. and S.M.H.; Writing – review and editing: D.C.B., D.L.S., and S.M.H.
All authors read the manuscript and provided comments. Funding acquisition: S.M.H.

Competing Interest Statement: The authors declare that they have no conflicts of
interest with the contents of this article. The content is solely the responsibility of the
authors and does not necessarily represent the official views of the National Institutes of
Health.

Figure Legends

Figure 1. The ufmylation activity of UFL1 promotes RIG-I signaling.

A) IFN-β-promoter reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells expressing vector, Flag-UFL1, or Flag-RIG-I, followed by mock or SenV infection (18 h) or in B) 293T cells transfected with vector (Vec) or Flag-UFL1, followed by mock or HCV PAMP RNA transfection (24 h). C) RT-qPCR analysis (rel. to *GAPDH*) of RNA extracted from 293T cells transfected vector or Flag-UFL1 that were treated with IFN-β (18 h). D) Immunoblot analysis of p-IRF3 following siRNA transfection along with expression of vector or Flag-UFL1^{siR}, which has point mutations in the siRNA seed sequence. Quantification of p-IRF3/Tubulin is shown on the right. E) RT-qPCR analysis (rel. to *18S*) of RNA extracted from primary neonatal human dermal fibroblasts (NHDFs) transfected with either siCTRL or siUFL1 followed by mock or SenV infection (8 h). F) ELISA for IFN-β of supernatants harvested from NHDFs transfected with siCTRL or siUFL1 and infected with SenV for the indicated times. G) Relative IFN-β-promoter reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells expressing indicated constructs followed by mock or SenV infection (12-18 h), with results graphed as relative SenV fold change for each. H) Diagram of UFL1 truncation constructs and corresponding ufmylation activity represented as the mean of mock and SenV values normalized to WT. For A) mean -/+ SD, n=3 technical replicates and representative of n=3 independent experiments. For all others, mean -/+ SEM, n=3 or n=5 (1G) biological replicates. *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001 determined by two-way ANOVA followed by Šidák's multiple comparisons test (B, E), Student's t-test (C, D, F), or one-way ANOVA followed by Dunnett's multiple comparisons test (G).

Figure 2. The ufmylation machinery proteins positively regulate RIG-I signaling.

A) IFN-β-promoter reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells expressing vector or Flag-UFM1, followed by mock or SenV infection (18 h) or in B) WT or CRISPR/CAS9 UFM1 KO 293T cells transfected with vector (Vec) or Flag-UFM1 (for KO), followed by mock or SenV infection (18 h). C) ELISA for IFN-β of supernatants harvested from WT or CRISPR/CAS9 UFM1 KO 293T cells that were SenV infected (18 h). D) Diagram of UFM1 conjugation. E) Relative IFN-β-promoter reporter luciferase expression (rel. to CMV-*Renilla*) from 293T cells expressing indicated constructs followed by mock or SenV infection (18 h), with results graphed as relative SenV fold change for each. Represented as mean -/+ SEM, n=3 biological replicates. *p ≤ 0.05, **p ≤ 0.01, and

***p ≤ 0.001 determined by two-way ANOVA followed by Tukey's multiple comparisons

test (A-B), Student's t-test (C), or one-way ANOVA followed by Dunnett's multiple

comparisons test (E).

Figure 3. UFM1 is required for the RIG-I driven transcriptional response.

RNA-seq analysis WT or UFM1 KO 293T cells following mock or SenV infection (18 h).

A) Gene set enrichment analysis of negatively regulated differentially expressed genes in

SenV-infected 293T cells represented by normalized enrichment score (UFM1 KO / WT).

B) Volcano plot of differentially expressed genes (adj P<0.01) shown in grey, with ISGs

shown in red, in SenV-infected 293T cells (UFM1 KO / WT). C) Heatmap of the effect of

UFM1 KO on the fold change of the 50 most induced IFN and ISGs (UFM1 KO / WT)

following SenV infection (adj P<0.01).

Figure 4. UFL1 is recruited to intracellular membranes and interacts with 14-3-3ε

and RIG-I during RNA virus infection.

A) Immunoblot analysis of inputs and subcellular membrane flotation of 293T cell extracts

that were mock or SenV-infected (4 h) followed by sucrose gradient fractionation, with

fraction numbers indicated from the top of the gradient (1) to bottom (8). Fractionation

controls, GAPDH for cytosol and Cox-I for membranes, are indicated and reveal that the

membranes are localized to fraction #1. Relative quantification of the ratio of UFL1 to a

membrane marker (Cox-I) in fraction 1 normalized to total protein levels in inputs are

shown on the right. B) Immunoblot analysis of anti-Flag immunoprecipitated extracts and

inputs from 293T cells expressing Myc-14-3-3ε and Flag-UFL1 that were mock- or SenV-

infected (4 h), with relative quantification on right. C) Immunoblot analysis of anti-Flag

immunoprecipitated extracts and inputs from 293T cells expressing Myc-UFL1 and Flag-

RIG-I that were mock- or SenV-infected (4 h), with relative quantification with IP values

normalized to inputs values on right. D) Immunoblot analysis of anti-RIG-I

immunoprecipitated (or anti-IgG) extracts and inputs from 293T cells that were mock- or

SenV-infected (4 h), with relative quantification with IP values normalized to inputs values

on right. E) Immunoblot analysis of anti-Flag immunoprecipitated extracts and inputs from

293T cells expressing Myc-UFL1 and Flag-RIG-I constructs that were mock- or SenV-

infected (4 h), with results quantified as relative fold change (SenV to Mock) for each. The

graphs are represented as the mean -/+ SEM, n=3 (A-B, D-E) or n=4 (C) biological

replicates and *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001 determined by Student's t-test (A-D)

or one-way ANOVA followed by Dunnett's multiple comparisons test (E).

Figure 5. UFL1 interaction with RIG-I requires 14-3-3ε and ufmylation.

A) Immunoblot analysis of anti-HA immunoprecipitated extracts and inputs from 293T cells

transfected with siCTRL or si14-3-3ε followed by SenV infection (4h). B) Immunoblot of

anti-HA immunoprecipitated extracts and inputs from 293T WT or UFM1 KO cells

transfected with HA-RIG-I and Flag-UFL1. C) Immunoblot of anti-Flag immunoprecipitated

extracts and inputs from 293T WT or UFM1 KO cells transfected with Flag-UFL1 and Myc-

14-3-3ε. In (A-C), SenV infection was for 4 hours, and relative quantification is shown on

the right, indicating the mean -/+ SEM (A, B), n=3 (A, B) biological replicates. For (C)

values shown are SD of IP values adjusted for input expression, with n=2 biological

replicates. *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001 determined by Student's t-test.

Figure 6. Ufmylation promotes RIG-I interaction with 14-3-3ε for MAVS activation.

A) Immunoblot of anti-Flag immunoprecipitated extracts and inputs from 293T cells

transfected with siCTRL or siUFL1 and indicated constructs. B) Immunoblot of anti-Flag

immunoprecipitated extracts and inputs from 293T WT or UFM1 KO cells. C) Immunoblot

of anti-Myc immunoprecipitated extracts from 293T WT or UFM1 KO cells. D) 293T WT or

UFM1 KO were mock or SenV-infected (12 h). Immunoblotting shows endogenous MAVS

in input samples and MAVS aggregation from P5 fractions, in the presence or absence of

denaturing reagent (β-mercaptoethanol). SenV infection was for 4 h (A-C) or 12 h (D). In

(A-C), relative quantification of indicated protein in the IP is shown on the right; in (D) SDD-

AGE MAVS values are normalized to corresponding SDS-PAGE values. Graphs show the

mean -/+ SEM for n=3 biological replicates. *p ≤ 0.05, **p ≤ 0.01, and ***p ≤ 0.001

determined by Student's t-test.

Figure S1. The domains of UFL1 that form UFM1 conjugates in cells.

Quantification of immunoblots from 293T cells expressing indicated Flag-UFL1

constructs or vector followed by mock or SenV infection (18 h) represented as the ratio

of UFM1 conjugates (approximately 25-50 kDa) to Flag-UFL1 expression in each lane

normalized to loading control with WT Flag-UFL1 mock set to 100. Graph indicates the

mean -/+ SEM for n=3 biological replicates.

Fig. S2. Transcriptional response of genes negatively regulated by UFM1.

RNA-seq analysis of WT versus UFM1 KO 293T cells showing the gene set enrichment

analysis (top 10 categories) of negatively regulated differentially expressed genes

represented by normalized enrichment score to identify gene ontology terms and

pathways associated with altered gene expression for each of the comparisons

performed (adj P<0.01).

Dataset S1. Differential expression analysis from RNA-seq analysis for UFM1 KO / WT

293T cells

Table S1.1: UFM1 KO / WT Mock

Table S1.2 UFM1 KO / WT SenV (18 h)

Dataset S2. Gene Set Enrichment Analysis for UFM1 KO / WT 293T cells

Table S2.1: UFM1 KO / WT Mock- negative direction

Table S2.2 UFM1 KO / WT SenV (18 h)- negative direction

Table S2.3 UFM1 KO / WT Mock- positive direction

Table S2.4 UFM1 KO / WT SenV (18 h)- negative direction

References

1. D. Thoresen, *et al.*, The molecular mechanism of RIG-I activation and signaling. *Immunol Rev* (2021) https:/doi.org/10.1111/imr.13022.

2. M. J. McFadden, N. S. Gokhale, S. M. Horner, Protect this house: cytosolic sensing of viruses. *Current Opinion in Virology* **22**, 36–43 (2017).

3. J. Rehwinkel, M. U. Gack, RIG-I-like receptors: their regulation and roles in RNA sensing. *Nature Reviews Immunology*, 1–15 (2020).

4. C. Chiang, M. U. Gack, Post-translational Control of Intracellular Pathogen Sensing Pathways. *Trends in Immunology* **38**, 39–52 (2017).

5. M. U. Gack, *et al.*, TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated antiviral activity. *Nature* **446**, 916–920 (2007).

6. H. Oshiumi, M. Matsumoto, S. Hatakeyama, T. Seya, Riplet/RNF135, a RING finger protein, ubiquitinates RIG-I to promote interferon-beta induction during the early phase of viral infection. *J. Biol. Chem.* **284**, 807–817 (2009).

7. H. M. Liu, *et al.*, The Mitochondrial Targeting Chaperone 14-3-3ε Regulates a RIG-I Translocon that Mediates Membrane Association and Innate Antiviral Immunity. *Cell Host & Microbe* **11**, 528–537 (2012).

8. H. Oshiumi, Recent Advances and Contradictions in the Study of the Individual Roles of Ubiquitin Ligases That Regulate RIG-I-Like Receptor-Mediated Antiviral Innate Immune Responses. *Front Immunol* **11**, 1296 (2020).

9. Y. K. Chan, M. U. Gack, A phosphomimetic-based mechanism of dengue virus to antagonize innate immunity. *Nature Immunology* **17**, 523–530 (2016).

10. W. Riedl, *et al.*, Zika Virus NS3 Mimics a Cellular 14-3-3-Binding Motif to Antagonize RIG-I- and MDA5-Mediated Innate Immunity. *Cell Host & Microbe* **26**, 493-503.e6 (2019).

11. C. Vazquez, S. M. Horner, MAVS Coordination of Antiviral Innate Immunity. *J Virol* **89**, 6974–6977 (2015).

12. E.-H. Tam, *et al.*, Role of the chaperone protein 14-3-3ε in the regulation of influenza A virus-activated beta interferon. *J Virol*, JVI0023121 (2021).

13. S. M. Horner, H. M. Liu, H. S. Park, J. Briley, M. Gale, Mitochondrial-associated endoplasmic reticulum membranes (MAM) form innate immune synapses and are targeted by hepatitis C virus. *Proc Natl Acad Sci U S A* **108**, 14590–14595 (2011).

14. S. M. Horner, C. Wilkins, S. Badil, J. Iskarpatyoti, M. G. Jr, Proteomic Analysis of Mitochondrial-Associated ER Membranes (MAM) during RNA Virus Infection Reveals Dynamic Changes in Protein and Organelle Trafficking. *PLOS ONE* **10**, e0117963 (2015).

15. D. C. Beachboard, *et al.*, The small GTPase RAB1B promotes antiviral innate immunity by
interacting with TNF receptor–associated factor 3 (TRAF3). *J. Biol. Chem.*,
jbc.RA119.007917 (2019).

16. K. Tatsumi, *et al.*, A novel type of E3 ligase for the Ufm1 conjugation system. *J. Biol.*
Chem. **285**, 5417–5427 (2010).

17. M. Komatsu, *et al.*, A novel protein-conjugating system for Ufm1, a ubiquitin-fold modifier.
EMBO J **23**, 1977–1986 (2004).

18. S. H. Kang, *et al.*, Two novel ubiquitin-fold modifier 1 (Ufm1)-specific proteases, UfSP1
and UfSP2. *J. Biol. Chem.* **282**, 5256–5262 (2007).

19. Y. Wei, X. Xu, UFMylation: A Unique & Fashionable Modification for Life. *Genomics*
Proteomics Bioinformatics **14**, 140–146 (2016).

20. J. Daniel, E. Liebau, The Ufm1 Cascade. *Cells* **3**, 627–638 (2014).

21. L. Wang, *et al.*, UFMylation of RPL26 links translocation-associated quality control to
endoplasmic reticulum protein homeostasis. *Cell Res*, 1–16 (2019).

22. Z. Wang, *et al.*, MRE11 UFMylation promotes ATM activation. *Nucleic Acids Research* **47**,
4124–4135 (2019).

23. H. M. Yoo, *et al.*, Modification of ASC1 by UFM1 is crucial for ERα transactivation and
breast cancer development. *Mol. Cell* **56**, 261–274 (2014).

24. B. Qin, *et al.*, UFL1 promotes histone H4 ufmylation and ATM activation. *Nat Commun* **10**,
1242 (2019).

25. I. A. Gak, *et al.*, UFMylation regulates translational homeostasis and cell cycle progression.
bioRxiv, 2020.02.03.931196 (2020).

26. R. DeJesus, *et al.*, Functional CRISPR screening identifies the ufmylation pathway as a
regulator of SQSTM1/p62. *eLife* **5**, e17290 (2016).

27. J. R. Liang, *et al.*, A Genome-wide ER-phagy Screen Highlights Key Roles of Mitochondrial
Metabolism and ER-Resident UFMylation. *Cell* **0** (2020).

28. C. P. Walczak, *et al.*, Ribosomal protein RPL26 is the principal target of UFMylation. *PNAS*
116, 1299–1308 (2019).

29. Y. Cai, N. Singh, H. Li, Essential role of Ufm1 conjugation in the hematopoietic system.
Exp. Hematol. **44**, 442–446 (2016).

30. J. Kulsuptrakul, R. Wang, N. L. Meyers, M. Ott, A. S. Puschnik, A genome-wide CRISPR
screen identifies UFMylation and TRAMP-like complexes as host factors required for
hepatitis A virus infection. *Cell Reports* **34** (2021).

31. D. R. Balce, *et al.*, UFMylation inhibits the proinflammatory capacity of interferon-γ–
activated macrophages. *PNAS* **118** (2021).

32. Y.-M. Loo, *et al.*, Viral and therapeutic control of IFN-β promoter stimulator 1 during
hepatitis C virus infection. *PNAS* **103**, 6001–6006 (2006).

33. R. Sumpter, *et al.*, Regulating intracellular antiviral defense and permissiveness to
hepatitis C virus RNA replication through a cellular RNA helicase, RIG-I. *J Virol* **79**, 2689–
2699 (2005).

34. T. Saito, D. M. Owen, F. Jiang, J. Marcotrigiano, M. Gale, Innate immunity induced by
composition-dependent RIG-I recognition of hepatitis C virus RNA. *Nature* **454**, 523–527
(2008).

35. D. Simsek, *et al.*, The Mammalian Ribo-interactome Reveals Ribosome Functional
Diversity and Heterogeneity. *Cell* **169**, 1051-1065.e18 (2017).

36. M. J. McFadden, *et al.*, Post-transcriptional regulation of antiviral gene expression by N6-
methyladenosine. *Cell Reports* **34**, 108798 (2021).

37. A. Peisley, B. Wu, H. Xu, Z. J. Chen, S. Hur, Structural basis for ubiquitin-mediated
antiviral signal activation by RIG-I. *Nature* **509**, 110–114 (2014).

38. A. Mizutani, M. Fukuda, K. Ibata, Y. Shiraishi, K. Mikoshiba, SYNCRIP, a Cytoplasmic
Counterpart of Heterogeneous Nuclear Ribonucleoprotein R, Interacts with Ubiquitous
Synaptotagmin Isoforms*. *Journal of Biological Chemistry* **275**, 9823–9831 (2000).

39. E. Foy, *et al.*, Control of antiviral defenses through hepatitis C virus disruption of retinoic
acid-inducible gene-I signaling. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 2986–2991 (2005).

40. E. Meylan, *et al.*, Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted
by hepatitis C virus. *Nature* **437**, 1167–1172 (2005).

41. X.-D. Li, L. Sun, R. B. Seth, G. Pineda, Z. J. Chen, Hepatitis C virus protease NS3/4A
cleaves mitochondrial antiviral signaling protein off the mitochondria to evade innate
immunity. *Proc Natl Acad Sci U S A* **102**, 17717–17722 (2005).

42. R. Lin, *et al.*, Dissociation of a MAVS/IPS-1/VISA/Cardif-IKKepsilon molecular complex
from the mitochondrial outer membrane by hepatitis C virus NS3-4A proteolytic cleavage. *J
Virol* **80**, 6072–6083 (2006).

43. K. Takahasi, *et al.*, Nonself RNA-Sensing Mechanism of RIG-I Helicase and Activation of
Antiviral Immune Responses. *Molecular Cell* **29**, 428–440 (2008).

44. H. Oshiumi, M. Miyashita, M. Matsumoto, T. Seya, A distinct role of Riplet-mediated K63-
Linked polyubiquitination of the RIG-I repressor domain in human antiviral innate immune
responses. *PLoS Pathog.* **9**, e1003533 (2013).

45. F. Hou, *et al.*, MAVS Forms Functional Prion-like Aggregates to Activate and Propagate
Antiviral Innate Immune Response. *Cell* **146**, 448–461 (2011).

46. N. Zamorano Cuervo, Q. Osseman, N. Grandvaux, Virus Infection Triggers MAVS
Polymers of Distinct Molecular Weight. *Viruses* **10** (2018).

47.   W. Zeng, *et al.*, Reconstitution of the RIG-I pathway reveals a signaling role of unanchored
788   polyubiquitin chains in innate immunity. *Cell* **141**, 315–330 (2010).

789   48.   X. Jiang, *et al.*, Ubiquitin-induced oligomerization of the RNA sensors RIG-I and MDA5
790   activates antiviral innate immune response. *Immunity* **36**, 959–973 (2012).

791   49.   L. Wang, *et al.*, UFMylation of RPL26 links translocation-associated quality control to
792   endoplasmic reticulum protein homeostasis. *Cell Research* **30**, 5–20 (2020).

793   50.   Y. Zhang, M. Zhang, J. Wu, G. Lei, H. Li, Transcriptional regulation of the Ufm1
794   conjugation system in response to disturbance of the endoplasmic reticulum homeostasis
795   and inhibition of vesicle trafficking. *PLoS ONE* **7**, e48587 (2012).

796   51.   C. Kwak, *et al.*, Contact-ID, a tool for profiling organelle contact sites, reveals regulatory
797   proteins of mitochondrial-associated membrane formation. *PNAS* **117**, 12109–12120
798   (2020).

799   52.   Y. Merbl, P. Refour, H. Patel, M. Springer, M. W. Kirschner, Profiling of ubiquitin-like
800   modifications reveals features of mitotic control. *Cell* **152**, 1160–1172 (2013).

801   53.   L. Wang, *et al.*, UFMylation of RPL26 links translocation-associated quality control to
802   endoplasmic reticulum protein homeostasis. *Cell Res.* **30**, 5–20 (2020).

803   54.   M. Hochstrasser, Origin and function of ubiquitin-like proteins. *Nature* **458**, 422–429
804   (2009).

805   55.   K. N. Swatek, D. Komander, Ubiquitin modifications. *Cell Research* **26**, 399–422 (2016).

806   56.   A. J. Muslin, J. W. Tanner, P. M. Allen, A. S. Shaw, Interaction of 14-3-3 with Signaling
807   Proteins Is Mediated by the Recognition of Phosphoserine. *Cell* **84**, 889–897 (1996).

808   57.   Y. Aghazadeh, V. Papadopoulos, The role of the 14-3-3 protein family in health, disease,
809   and drug development. *Drug Discovery Today* **21**, 278–287 (2016).

810   58.   A. Aitken, Post-translational modification of 14-3-3 isoforms and regulation of cellular
811   function. *Seminars in Cell & Developmental Biology* **22**, 673–680 (2011).

812   59.   J.-P. Lin, Y.-K. Fan, H. M. Liu, The 14-3-3η chaperone protein promotes antiviral innate
813   immunity via facilitating MDA5 oligomerization and intracellular redistribution. *PLoS*
814   *Pathog.* **15**, e1007582 (2019).

815   60.   K. Onomoto, K. Onoguchi, M. Yoneyama, Regulation of RIG-I-like receptor-mediated
816   signaling: interaction between host and viral factors. *Cell Mol Immunol* **18**, 539–555 (2021).

817   61.   G. Liu, *et al.*, ISG15-dependent activation of the sensor MDA5 is antagonized by the
818   SARS-CoV-2 papain-like protease to evade host innate immunity. *Nature Microbiology*, 1–
819   12 (2021).

820   62.   M. Yoneyama, *et al.*, Shared and Unique Functions of the DExD/H-Box Helicases RIG-I,
821   MDA5, and LGP2 in Antiviral Innate Immunity. *The Journal of Immunology* **175**, 2851–
822   2858 (2005).

63.   B. Fredericksen, *et al.*, Activation of the Interferon-β Promoter During Hepatitis C Virus
RNA Replication. *Viral Immunology* **15**, 29–40 (2002).

64.   T. Saito, *et al.*, Regulation of innate antiviral defenses through a shared repressor domain
in RIG-I and LGP2. *PNAS* **104**, 582–587 (2007).

65.   F. A. Ran, *et al.*, Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**,
2281–2308 (2013).

66.   Babraham Bioinformatics - Trim Galore! (October 12, 2021).

67.   M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**, 10–12 (2011).

68.   P. J. Kersey, *et al.*, Ensembl Genomes: an integrative resource for genome-scale data
from non-vertebrate species. *Nucleic Acids Res* **40**, D91-97 (2012).

69.   A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

70.   HTSeq: Analysing high-throughput sequencing data with Python — HTSeq 0.13.5
documentation (October 12, 2021).

71.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

72.   W. Huber, *et al.*, Orchestrating high-throughput genomic analysis with Bioconductor. *Nat
Methods* **12**, 115–121 (2015).

73.   V. K. Mootha, *et al.*, PGC-1alpha-responsive genes involved in oxidative phosphorylation
are coordinately downregulated in human diabetes. *Nat Genet* **34**, 267–273 (2003).

74.   H. M. Liu, *et al.*, SYNCRIP (synaptotagmin-binding, cytoplasmic RNA-interacting protein) is
a host factor involved in hepatitis C virus RNA replication. *Virology* **386**, 249–256 (2009).

A

IFN-β promoter (RLU)

Mock / SenV

Vector / UFM1

Flag-UFM1
SenV
Total Protein

15 / 50

B

IFN-β promoter (RLU)

Mock / SenV

Flag-UFM1: − − − − + +
WT / UFM1$_{KO}$

Flag-UFM1
UFM1
SenV
GAPDH

15 / 10 / 50 / 37

C

+SenV

IFN-β pg/mL

WT / UFM1$_{KO}$

SenV: + +
UFM1: + −

UFM1
SenV
Tubulin

10 / 50 / 50

D

① UFM1—GSC → G → G
② UBA5 ● → ③ UFC1 ●
④ UFL1 *target* → *target* ●
⑤ *target* ● UFSP2 → *target* + UFM1 ● G

E

+SenV

Relative IFN-β promoter activity (RLU)

Vec / UFL1 / UFM1 / UBA5 / UFC1 / UFSP2

*** / *** / *** / * / ***

Flag

Vec / UFL1 / UFM1 / UBA5 / UFC1 / UFSP2

Flag
75 / 15 / 50 / 25

SenV
GAPDH
50 / 37

A

Response to Type I IFN
Defense/Response to Virus
Innate Immune Response
Response to Virus
Negative Regulation of Viral Process
Cytokine-Mediated Signaling Pathway
Negative Regulation of Viral Genome Replication
Regulation of Type I IFN Production
Response to IFNγ
Defense/Response to Other Organism

Normalized Enrichment Score UFM1$_{KO}$/WT (+SenV)

-2.55 -2.60 -2.65 -2.70 -2.75 -2.80 -2.85

B

$-\log_{10}$ adj P-value

Differentially expressed genes
ISGs

Log$_2$ Fold Change UFM1$_{KO}$/ WT (+SenV)

C

APOL1	IFI6
RARRES3	CXCL10
GBP5	IL15
ODF3B	SP100
IFNB1	APOL6
TRIM22	ISG15
IFNL1	CXCL11
RSAD2	IFI35
TRANK1	SAMD9
BST2	PARP10
DDX58	UBE2L6
IFI44	OAS1
IFIT2	EPSTI1
OASL	CMPK2
IFITM3	B3GNT7
ISG20	XAF1
IFIH1	OAS3
IFIT3	PMPAIP1
GPB1	B2M
IFI44L	DDX60
HERC5	NLRC5
DHX58	CEBPD
OAS2	MX1
IFITM1	C19orf66
IFIT1	IL15RA

Log$_2$ Fold Change UFM1$_{KO}$/WT (+SenV)

-4.5 -4.0 -3.5 -3.0 -2.5 -2.0 -1.5