1

Anatomic development of the upper airway during the first five years of life: A three-dimensional imaging study

Ying Ji Chuang[1], Seong Jae Hwang[2], Kevin A. Buhr[3], Courtney A. Miller[1], Gregory. D. Avey[4],

Brad H. Story[5], and Houri K. Vorperian[1*]

[1] Vocal Tract Development Lab, Waisman Center, University of Wisconsin-Madison, Madison, Wisconsin, United States of America

[2] Department of Computer Science, University of Pittsburgh, Pittsburg, Pennsylvania, United States of America

[3] Department of Biostatistics and Medical Informatics, University of Wisconsin-Madison, Madison, Wisconsin, United States of America

[4] Department of Radiology, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, United States of America

[5] Speech, Language, and Hearing Sciences, University of Arizona, Tucson, Arizona, United States of America

*Corresponding author:

E-mail: vorperian@waisman.wisc.edu HKV

Funding: The authors are grateful for National Institutes of Health (NIH) funding support from the National Institute on Deafness and Other Communication Disorders (NIDCD) R01 DC006282 (HKV) and from the National Institute of Child Health and Human Development (NICHD) U54 HD090256 (QC). There was no involvement of the funding sources in theresearch design, data collection, analysis, interpretation, writing of this paper or submission for publication.

Abstract

Purpose. Normative data on the growth and development of the upper airway across the sexes is needed for the diagnosis and treatment of congenital and acquired respiratory anomalies and to gain insight on developmental changes in speech acoustics and disorders with craniofacial anomalies. **Methods.** The growth of the upper airway in children ages birth-to-five years, as compared to adults, was quantified using an imaging database with computed tomography studies from typically developing individuals. Methodological criteria for scan inclusion and airway measurements included: head position, histogram-based airway segmentation, anatomic landmark placement, and development of a semi-automatic centerline for data extraction. A comprehensive set of 2D and 3D supra- and sub-glottal measurements from the choanae to tracheal opening were obtained including: naso-oro-laryngo-pharynx subregion volume and length, each subregion's superior and inferior cross-sectional-area, and antero-posterior and transverse/width distances. **Results.** Growth of the upper airway during the first five years of life was more pronounced in the vertical and transverse/lateral dimensions than in the antero-posterior dimension. By age five years, females have larger pharyngeal measurement than males. Prepubertal sex-differences were identified in the subglottal region. **Conclusions.** Our findings demonstrate the importance of studying the growth of the upper airway in 3D. As the lumen length increases, its shape changes, becoming increasingly elliptical during the first five years of life. This study also emphasizes the importance of methodological considerations for both image acquisition and data extraction, as well as the use of consistent anatomic structures in defining pharyngeal regions.

Key words: upper airway; pharynx; development; volume; aerodigestive tract; vocal tract length.

Introduction

The upper airway, a virtual conduit as characterized by Marcus et al. [1], has an anatomic boundary defined by other tissues (bony, cartilaginous and soft) while serving the functions of respiration, food ingestion (mastication and deglutition), as well as vocalization/speech, hence the function-based terms '*aerodigestive tract*', '*vocal tract*', or more comprehensively the '*aerodigestive and vocal tract*'. During the course of development, especially from infancy to early childhood, the upper airway undergoes drastic changes in size, shape and mechanical properties due to the restructuring of its anatomical sub-components, such as the descent of the larynx and the hyoid bone [2-4]. The anatomic growth process persists while adapting to the various functional needs and demands during maturation. As posited by current theory on craniofacial growth, the development of the upper airway is shaped by both genetic as well as intrinsic and extrinsic epigenetic factors, such as function, mechanical forces, and trauma [5-13].

The lack of knowledge regarding the growth and development of the upper airway, defined as the air conduit from the level of the nose to the carina, was addressed in a workshop by the National Heart, Lung, and Blood Institute (NHLBI) in 2009 with a large team of clinicians and scientists from diverse fields in healthcare and the biological sciences [1]. The outcome was a comprehensive set of research guidelines on various aspects of the upper airway, each with a set of priorities relevant to clinical disorders of upper airway functions. Among the priorities was the need to study the developmental changes of the upper airway anatomy and function during childhood (neonatal to puberty) across sexes and ethnicities and to provide normative values of the upper airway. Normative data are needed to better understand common respiratory disorders such as obstructive sleep apnea syndrome (OSAS), as well as a number of other congenital and acquired respiratory anomalies [1]. Furthermore, normative data can provide additional insight

on developmental speech acoustics [14, 15], as well as speech disorders, particularly where

craniofacial anomalies are present [16, 17]. As listed in Table 1, a large number of studies have

examined the upper airway anatomy using different modalities, methodologies, airway regions,

and age ranges. Table 1 summarizes the studies to date that have examined the typical

development of the aerodigestive and vocal tract from the choanae or the soft palate superiorly to

the epiglottis or the trachea inferiorly. A subset of studies listed have factored in growth and/or

sex in their data analysis. Most studies have employed imaging to obtain quantitative

measurements, including linear, angular and/or area measurements, based on the midsagittal or

axial slices, as well as volumetric measurements. However, only a very limited number of

studies have assessed multidimensional volumetric measurements during early childhood.

Among the 34 studies listed in Table 1, only 17 studies included linear, area and volumetric

measurements, and fewer than half of those studies controlled for head position during or after

data acquisition. Of the 12 studies summarized in Table 1 that examined the pre-pubertal age-

group, the majority obtained measurements in 2D that were collected primarily from

radiographic images using mid-sagittal, axial, or coronal visualization planes; an approach

frequently used to assess the upper airway, as it is cost effective and less time-consuming to

process. However, this approach does not provide accurate representation of the complex airway

morphology, as it overlooks information of lateral dimensions [18, 19]. Two of those 12 studies

[20, 21] quantified the prepubertal airway in 3D but only Abramson et al. [20] covered the entire

prepubertal period from birth to five years and assessed sexual dimorphism. Neither of those

retrospective studies reported controlling for head position or using it as an inclusion criterion.

99   **Table 1. Summary of studies on typical upper airway development.**

Methodology		Study			Age Cohort (yrs)				Assessment/ Control		Measurement Type					Airway Region
Type	Modality	Author (year) [a]	n = Total (M/F)	Age	Pre-Pub (0-5)	Peri-Pub (6-10)	Pubertal (11-17)	Post-Pub (18+)	Sexual Dimorphism	Head Position	Dist-ance	Region Length	CSA (*)	Volume	Center-line	
2D	CT	Ronen et al. (2007)	n = 69 (38/31)	4-10, 14-19	+	+	+	+	+		YES					PNS - hyoid superior
	CT, MRI	Vorperian et al. (2009)	n = 605 (327/278)	0-19	+	+	+	+	+		YES	YES				Choanae - glottis
	MRI	Fitch & Giedd (1999)	n = 129 (76/53)	2-25	+	+	+	+	+		YES	YES				PNS - C4ai
	MRI	Litman et al. (2002)	n = 16 (9/7)	10mos-7yrs	+	+				+	YES		YES (a)			Oral-pharyngeal
	MRI	Machata et al. (2010)	n = 138 (60/78)	0-6	+	+				+	YES		YES (a)			Soft palate - tongue
	MRI	Yi et al. (2017)	n = 521 (296/225)	0-6	+	+			+		YES					Soft palate - tip of epiglottis
	MRI	Fregosi et al. (2003)	n = 18 (13/5)	7-12		+	+				YES		YES (a)	YES		Choanae - epiglottis base
	X-ray	Barbier et al. (2015)	n = 966 (494/470)	0-25	+	+	+	+	+		YES	-	-	-	-	PNS - tip of epiglottis
	X-ray	Goncalves et al. (2011)	n = 390 (195/195)	6-18		+	+	+	+		YES					Oral-pharyngeal
	X-ray	Sheng et al. (2009)	n = 239 (107/132)	7-27		+	+	+	+		YES					PNS - mandible/pharynx intersection
	X-ray	Mislik et al. (2014)	n = 880 (458/422)	6-17		+	+				YES					PNS - epiglottis base
	X-ray	Daraze et al. (2017)	n = 117 (48/69)	21-25				+	+	+	YES					PNS - tip of epiglottis
	Fluoroscopy	Rommel et al. (2003)	n = 23 (14/9)	0-4	+					+	YES	YES				Choanae - epiglottis base
3D	CBCT	Chiang et al. (2012)	n = 387 (173/214)	8-18		+	+	+	+	+	YES		YES	YES	YES	PNS - C4ai
	CBCT	Jiang et al. (2014)	n = 254 (119/135)	6-18		+	+	+	+	+	YES	YES	YES (x)	YES		Choanae - hyoid superior
	CBCT	Lenza et al. (2010)	n = 34 (14/20)	11-56			+	+	+		YES		YES (x)	YES		Choanae - tip of epiglottis
	CBCT	Schendel et al. (2012)	n = 1300 (571/729)	6-60		+	+	+	+		YES		YES (a)	YES		PNS - C4as
	CBCT	Anandarajah et al. 2017	n = 105 (44/61)	7-13		+	+						YES (a)	YES		PNS - epiglottis base
	CBCT	Masoud et al. (2020)	n = 81 (32/49)	7-17		+	+			+	YES	YES	YES (a)	YES		Choanae - epiglottis base
	CBCT	Yanagita et al. (2017)	n = 61 (0/12)	10-16		+	+			+	YES	YES	YES (a)	YES		Choanae - epiglottis base
	CBCT	Alves et al. (2012)	n = 50 (27/23)	8-10		+				+	YES		YES (a)	YES		PNS - tip of epiglottis
	CBCT	Claudino et al. (2013)	n = 54 (21/33)	13-20			+	+	+	+	YES	YES	YES (a)	YES		Choanae - hyoid inferior
	CT	Abramson et al. (2009)	n = 46 (31/15)	4mos-64yrs	+	+	+	+	+		YES		YES (a)	YES		PNS - epiglottis base
	CT	Li et al. (2011)	n = 281 (141/140)	10-18		+	+	+		+	YES	YES	YES (a)	YES		PNS - epiglottis base
	CT	Kim et al. (2011)	n = 73 (64/9)	14-72			+	+	+	+	YES		YES (a)	YES		PNS - hyoid superior
	CT	Gibelli et al. (2020)	n = 80 (40/40)	21-86				+	+	+	YES			YES		Choanae - glottis
	CT	Inamoto et al. (2015)	n = 107 (30/77)	23-77				+	+	+	YES			YES		PNS - Glottis
	CT	Shigeta et al. (2008)	n = 38 (19/19)	24-84				+	+		YES	YES		YES		PNS - tip of epiglottis
	MRI	Smitthimedhin et al. (2018)	n = 96 (53/43)	0.5-46wks	+									YES		Choanae - trachea
	MRI	Welch et al. (2002)	n = 12 (0/12)	30-64				+		+	YES		YES	YES	YES	PNS - tip of epiglottis
Other	APh	Leboulanger et al. (2011)	n = 59 (28/31)	2-18	+	+	+	+	+	+	YES					Oral-pharyngeal
	APh	Martin et al. (1997)	n = 114 (60/54)	16-74			+	+	+		YES	YES				Oral-pharyngeal
	APh, Spiro	Brooks et al. (1992)	n = 175 (77/98)	19-64				+	+		YES	YES				Airway
	APh, Spiro	Brown et al. (1986)	n = 24 (14/10)	26-42				+	+		YES	YES				Airway

Summary of studies on typical upper airway development listed per methodology in the **first** column, using measurement type/dimension: 2D

(two-dimensional), indicates that measurements were collected from radiographic images representing either the mid-sagittal, axial or coronal

visualization planes; 3D (three-dimensional), indicates that measurements were collected utilizing multiplanar visualizations (axial, coronal

and/or sagittal planes) and 3D representation of the upper airway; Other indicates that measurements were made using non-imaging techniques.

The first column also lists alphabetically study modality including imaging (CT, CBCT, MRI, X-ray) or non-imaging (Acoustic Pharyngometry

(APh), Spirometer (Spiro)) techniques. The **second** column lists alphabetically the study author(s) with year of publication in parentheses,

sample size (n=) with total male/female (M/F) numbers specified in parentheses, and age range examined. Age of study participants is also

classified using pubertal age cohorts in the **third** column, followed by assessment or control of sex-differences and head position in the **fourth**

column. The **fifth** column lists measurement reported including overall pharyngeal distance, and pharyngeal region/subregion measurements

including: length, cross-sectional area* (CSA), volume and centerline length. The **final** column lists the defined superior to inferior anatomical

boundaries in each study.

* CSA specifications include the automatic or manual calculation of the area of a plane of the pharyngeal airway in: the axial plane (a), a plane

that is orthogonal to the pharyngeal centerline (o), or a plane connecting two or more specific anatomical landmarks (x).

[a] Cited literature: [20-53]

Since the upper airway is a lumen, attention must be paid to a number of methodological

considerations, given their potential effect on various pharyngeal measurements. Methodological

procedures known to affect pharyngeal measurements include head/neck position

(flexion/extension), body position (upright/supine), and sedation [37, 54, 55]. While most studies

to date have accounted for one or more of these confounders, it is difficult to compare findings

across studies unless all confounders have been addressed. For example, Inamoto et al. [47],

reported significant sex differences between the adult male and female laryngopharynx, but

Gibelli et al. [46], who also used CT but did not control for head position, reported no sex

differences. Additionally, variations in the anatomical boundary and subregion borders of the

upper airway morphology (nasopharynx, oropharynx, and laryngopharynx/hypopharynx) as

defined by different studies, summarized in Table 1 (final column) and Table 2, further

complicates the ease and feasibility of comparing findings across studies. Thus, to ensure an

accurate and reliable assessment of the developmental changes of this cavity, it is critical to use

standardized imaging procedures, well-defined anatomical regions, and established airway data

extraction protocols.

Table 2. Summary of boundaries of the pharyngeal regions.

Main Pharyngeal Subdivisions	Anatomical boundaries	Definitions [b]
Nasopharynx	**Superior Border**	Nares (Ayappa & Rapoport, 2002) Nasal cavity (Adewale, 2009) End of nasal septum (Netter, 2019) Choanae (Arens et al., 2004; Logan et al., 2017; Schuenke et al., 2010)
	Inferior Border	Hard palate (Ayappa & Rapoport, 2002) Soft palate (Laird et al., 2019 ; Moore et al., 2006; Standring et al., 2017) Level of soft palate (Arens et al., 2004) Above soft palate (Adewale, 2009; Gu et al., 2016) Inferior/Lower border of soft palate (Netter, 2019; Logan et al., 2017)
Oropharynx	**Superior Border**	Soft palate (Ayappa & Rapoport, 2002) Uvula (Schuenke et al., 2010)
	Inferior Border	Epiglottis (Ayappa & Rapoport, 2002; Schuenke et al., 2010) Superior/Upper border of epiglottis (Gu et al., 2016; Laird et al., 2019 ; Moore et al., 2006; Netter, 2019; Standring et al., 2017) Larynx (Arens et al., 2004)
Laryngopharynx/ Hypopharynx	**Superior Border**	Posteriorlateral to the larynx (Arens et al., 2004) Base of tongue (Ayappa & Rapoport, 2002) Tip of epiglottis (Adewale, 2009) Superior/Upper border of epiglottis (Logan et al., 2017)
	Inferior Border	Larynx (Ayappa & Rapoport, 2002) Opening of esophagus (Netter, 2019) Cricoid cartilage (Schuenke et al., 2010) Inferior/Lower border of cricoid cartilage (Adewale, 2009; Gu et al., 2016; Laird et al., 2019; Logan, 2017; Moore et al., 2006; Standring et al., 2017)

133   Summary of pharyngeal regions' boundaries as defined in relevant anatomy textbooks and

134   published papers. Note the lack of consistency and/or specificity in the anatomical boundaries for

135   each region.

136   [b] Cited literature: [56-65]

This study aims to systematically study the anatomic development of the upper airway, specifically the structural changes from the choanae to the tracheal opening (inferior border of the cricoid cartilage) from birth to five years, as compared to adults. To acquire normative data of the anatomy that subserve the aerodigestive and speech functions, we used an imaging database with computed tomography (CT) studies from typically developing individuals to obtain a comprehensive set of two-dimensional (2D) and three-dimensional (3D) measurements quantifying the growth of the upper airway. Our comprehensive set of methodological criteria included control of head position, histogram-based upper airway segmentation, placement of anatomic landmarks, and development of a semi-automatic method to determine lumen centerline. In addition, to better understand the resonance/acoustic characteristics of the vocal tract, this study aimed to examine the nature of the developmental changes of the upper airway dimensions and to determine if there are sex differences in the upper airway dimensions during the pre-pubertal period. We hypothesize all pediatric airway dimensions to be substantially smaller than adult dimensions. We also hypothesize sex differences in both children and adults.

Materials and methods

a. Image acquisition/dataset

Using imaging studies performed at the University of Wisconsin Hospital and Clinics (UWHC), our Vocal Tract Development Lab (VTLab) has curated a lifespan retrospective database of more than 2000 head and neck CT scans to study the anatomic growth and development of the oral and pharyngeal structures. This database was established following approval of the University of Wisconsin-Madison Institutional Review Board (IRB) and anonymized accordingly. All CT imaging studies, performed in the supine body position, were acquired using CT scanners

manufactured by General Electric Medical Systems or Siemens and stored in Digital Imaging

and Communications in Medicine (DICOM) format. Additional details on this imaging database

and image acquisition are provided in Kelly et al. [66], Miller et al. [67, 68] and Vorperian et al.

[23].

To ensure the adequacy of imaging studies selected for this study, the VTLab imaging

database was reviewed for typically developing cases between the ages 0-5 years (pediatric) and

20-30 years (adults) who were imaged for conditions that do not affect typical growth. A total of

410 (208 Males (M), and 202 Females (F)) CT imaging studies that included 264 (161M, 103F)

pediatric scans and 146 (47M, 99F) adult scans, from 276 (143M, 133F) individuals (195 [115M,

80F] children; and 81 [28M, 53F] adults), were inspected for cases that met the following

inclusion criteria: (1) slice thickness ≤ 2.5mm, (2) 14-22cm field-of-view (FOV), (3) 512x512

matrix size, (4) no movements or dental artifacts affecting the view of pharynx structure, and (5)

neutral or flexed head position as confirmed using Miller et al.'s [68] head position classification

protocol. While all extreme flexion/extension cases were excluded, including all sedation cases,

neutral-flexed head position cases were not excluded given that the larger infant head is prone to

being flexed in the supine position. The total yield of cases that met the inclusion criteria for this

study's dataset included 61 (32M, 29F) pediatric cases from 78 imaging studies (41M, 37F), and

17 (9M, 8F) adult cases from 72 (39M, 33F) imaging studies. The individuals whose images

were used included 56 (31M, 25F) children, and 16 (8M, 8F) adults. Age specific demographics

are presented in Table 3.

Table 3. Distribution of male and female cases per age group.

Group (Age range (yr;mos))	M	F	Total
<1 (00;00 - 00;11)	4	5	9
1 (01;00 - 01;11)	4	5	9
2 (02;00 - 02;11)	6	7	13
3 (03;00 - 03;11)	9	8	17
4 (04;00 - 04;11)	6	7	13
5 (20;00 - 30;00)	8	9	17

Distribution of male (M) and female (F) cases per age group. Age groups specified in years;

months (group <1 includes cases birth (00;00) to 11 months (00;11); group 1 includes cases 1

year (01;00) to 1 year 11 months (01;11) etc., and group 5 adults ages 20-to-30 years.

b. Image reconstruction

The standard reconstruction kernel was the preferred CT reconstruction algorithm, and was

available for the majority of the imaging studies. For cases/imaging studies processed without

the standard kernel, imaging features of the standard kernel were simulated by processing the

soft kernel with an unsharp enhance filter using a kernel size of 5x5, or the by processing the

bone kernel with a low pass filter using a kernel size of 3x3. Next, the software Analyze 12.0

[69] was used to reconstruct CT images from DICOM format into 3D volume.

A histogram-based threshold method was applied to the reconstructed CT volume in

order to identify the intensity in Hounsfield Unit (HU) that allows an optimal representation of

the airway. Guided by the technique of Nakano et al. [70], per image, we used the midpoint

between the air threshold peak (-1000 HU) and soft tissue peak (+100 HU to +300 HU), as the

applied upper threshold intensity to segment the airway. The range of upper thresholds used in

this study was between -556 HU and -445 HU. The Volume Render and Volume Edit modules

were then used to visualize and segment the 3D pharynx model from the reconstructed CT

volume. Using the identified threshold value, the airway region studied was restricted inferiorly

at the first tracheal ring (lower limit of the cricoid cartilage), and superiorly at the choanae. The

resulting 3D pharynx model was saved in Analyze Object Map format [.obj].

c. Anatomic landmarks and variables

As depicted in Figs 1 and 2, and listed with descriptions in Table 4, a set of 26 anatomic

landmarks that included 20 pharyngeal, 4 maxillary, and 2 reference landmarks were manually

placed on each of the 78 3D pharynx models to quantify upper airway growth. The set of

landmarks selected were carefully determined following a thorough review of landmarks and

airway variables examined in studies to date [20, 24, 36, 47, 54, 71]. Landmark placement

entailed using the Volume Render module in Analyze 12.0 [69], to manually place each of the 26

landmarks by overlaying them on their respective CT images while using the axial, coronal and

sagittal planes to guide accuracy of landmark placement. The landmarks were similarly saved in

Analyze Object Map format. To ensure reliability in landmark placement, prior to data

collection, two researchers modeled and placed landmarks on six cases. The differences in

resulting measurements, calculated from the raters' landmarks, had an average relative error

(ARE) that was less than or equal to 5% between researchers. The landmarks were then used to

establish a data extraction protocol, described in the following section, that generates pharyngeal

cross sections perpendicular to the centerline and calculates landmark-based measurements. The

comprehensive set of 30 pharyngeal variables measured, as listed and defined in Table 5 below,

are described in the section on variable measurements.

Fig 1. Illustration of airway regions and measurements. The airway was examined using

landmark-derived planes orthogonal to the centerline, as described in text. The four airway

regions bounded by five cross sectional areas, a-to-d as depicted in the right panel, were

quantified developmentally using the following measurements: volume, region length, cross-

sectional area (CSA), anterior-posterior distance, and lateral width - as defined in Table 5. The

airway regions above the glottis (d; Table 5, definition 19), included the following pharyngeal

regions: a. *Nasopharynx* (blue; definition 1); b. *Oropharynx* (red; definition 6); c.

Laryngopharynx (cyan; definition 11); and the airway below the glottis consisted of the

subglottal region (magenta; definition 22). The pharynx (Table 5, definition 16) consisted of all

three supraglottal regions.

Fig 2. 3D airway model (choanae to trachea) of a 4-year 8-month old typically developing

male as visualized in MATLAB. Blue dots represent the 26 anatomic landmarks listed in Table

4. Centerline is shown in green, and CSAs closest to each of the anatomic landmarks are shown

in red.

Table 4. Description/definition of the 26 anatomic landmarks.

#	Description of landmarks	Landmark Name	Abbreviation
	Pharynx Landmarks		
1	The point of attachment of the vocal folds with the thyroid cartilage at the anterior commissure, 2-3 mm below the thyroid notch of the larynx.	Glottis Anterior	*(ga)*
2	In the axial plane at the level of the glottis as determined by the tear-shaped glottal area (GA), the most posterior point of the laryngo-pharynx between the lateral-most sides of the vocal folds attached to the arytenoid cartilages.	Glottis Posterior	*(gp)*
3	Most superior and posterior point of the epiglottis in the midsagittal plane.	Epiglottis Superior	*(epS)*
4	Attachment of the epiglottis with the hyoepiglottic ligament, visualized as the most anteroinferior point of contact in the midsagittal plane.	Epiglottis Base	*(epBse)*
5	The most posterior point on the pharyngeal wall at the level of the epiglottis base in the midsagittal plane.	Epiglottis Base Posterior	*(epBsePo)*
6	Most inferior point of the left piriform sinus.	Piriform Sinus Inferior Left	*(PSInL)*
7	Most inferior point of the right piriform sinus.	Piriform Sinus Inferior Right	*(PSInR)*
8	The midpoint (most 'curved' point) of the left aryepiglottic fold . Approximately halfway between the base and tip of the epiglottis.	Piriform Sinus Superior Left	*(PSSuL)*
9	The midpoint (most 'curved' point) of the right aryepiglottic fold. Approximately halfway between the base and tip of the epiglottis.	Piriform Sinus Superior Right	*(PSSuR)*
10	Most inferior point on the left vallecula.	Vallecula Inferior Left	*(VaInL)*
11	Most inferior point on the right vallecula.	Vallecula Inferior Right	*(VaInR)*
12	Most anterior point of the anterior pharyngeal wall at the level of the velum tip as visualized in the midsagittal plane.	Velum Anterior	*(VeAn)*
13	Most posterior point of the posterior pharyngeal wall at the level of the velum tip, as visualized in the midsagittal plane.	Velum Posterior	*(VePo)*
14	Midpoint between VeAn and VePo (landmarks 12 and 13 as defined above). Midpoint was calculated based on VePo and VeAn landmark coordinates.	Midpoint between VeAn and VePo	*(MidVe)*
15	The most supero-posterior point of the velum.	Velum Back	*(VeBa)*
16	The inferior tip of the velum.	Velum End	*(VeEnd)*
17	The most anterior point of the pharynx at the level of the PNS in the axial plane as guided by the midsagittal plane of the pharynx.	Nasopharynx Anterior	*(NpxAn)*
18	Most posterior point of the pharyngeal wall at the level of the PNS in the axial plane as guided by the midsagittal plane of the pharynx.	Nasopharynx Posterior	*(NpxPo)*
19	Midpoint between NpxAn and NpxPo (landmarks 17 and 18). Midpoint was calculated based on NpxAn and NpxPo landmark coordinates).	Midpoint between NpxAn and NpxPo	*(NpxMid)*
20	Landmark placed on the soft tissue between the posterior vomer bone and the nasal crest of the palatine bone.	Posterior Nasal Septum	*(NasalS)*
	Reference Landmarks		
21	The most anterior point of the Anterior Nasal Spine.	Anterior Nasal Spine	*(ANS)*
22	The most posterior point of the Posterior Nasal Spine.	Posterior Nasal Spine	*(PNS)*
	Maxilla Landmarks		
23	The most postero-inferior point of the maxillary alveolar bone in the midsagittal plane (landmark location is between the first incisors)	Alveolar bone of incisor	*(ABI)*
24	The most posterior point of the incisive canal in the midsagittal plane of the maxilla.	Posterior edge of incisive canal	*(PIC)*
25	The intersection between the transverse palatine suture and the median palatine suture.	Palatine Sutures intersection	*(PALS)*
26	Midpoint between PIC and PALS (landmarks 24 and 25) along the median palatine suture on the maxilla. Coordinates calculated based on PIC and PALS landmark coordinates x, y, z.	Maxilla midpoint	*(MMax)*

243 Description/definition of the 26 anatomic landmarks (pharynx, reference, and maxilla), listed

244 from the inferior to the superior regions of the airway as displayed in Fig 1. Landmark placement

245 entailed use of multiplanar views (at least two of the sagittal, axial, and coronal planes, or all

246   three) for accuracy. These landmarks were used in defining study variables and extracting the

247   quantitative measurements of the upper airway as specified in Table 5.

249 **Table 5. All upper airway variables examined.**

Measure-ment #	Variable Description	Variable Name	(Abbreviation)
	Nasopharynx		
1	Orthogonal volume of the region bound by the intersections of the centerline with the palatal plane (ANS-PNS) superiorly and inferiorly with the tip of the velum.	Nasopharynx Volume	Nasopharynx
2	The curvilinear segment length along the centerline of the Nasopharynx Volume.	Nasopharynx Length	NasopharynxL
3	Cross sectional area (CSA) of the superior border of Nasopharynx Volume.	Nasopharynx Area	NasopharynxArea
4	The distance between the most anterior and posterior points along the midline of the superior border of the Nasopharynx Volume.	Nasopharynx Anterior-Posterior Distance	NasopharynxAPDist
5	The distance between the most lateral left and right points along the midline of the superior border of the Nasopharynx Volume.	Nasopharynx Width	NasopharynxWidth
	Oropharynx		
6	Orthogonal volume of the region bounded superiorly by the tip of the velum, and inferiorly by the midpoint of the aryepiglottic folds.	Oropharynx Volume	Oropharynx
7	The curvilinear segment length along the centerline of the Oropharynx Volume.	Oropharynx Length	OropharynxL
8	CSA of the superior border of the Oropharynx Volume.	Oropharynx Area	OropharynxArea
9	The distance between the most anterior and posterior points along the midline of the superior border of the Oropharynx Volume.	Oropharynx Anterior-Posterior Distance	OropharynxAPDist
10	The distance between the most lateral left and right points along the midline of the superior border of the Oropharynx Volume.	Oropharynx Width	OropharynxWidth
	Laryngopharynx		
11	Orthogonal volume of the region bounded by the orthogonal planes where the centerline intersects superiorly with the tip of the midpoint of the aryepiglottic folds, and inferiorly with the glottis.	Laryngopharynx Volume	Laryngopharynx
12	The curvilinear segment length along the centerline of the Laryngopharynx Volume.	Laryngopharynx Length	LaryngopharynxL
13	CSA of the superior orthogonal border/plane of Laryngopharynx Volume.	Laryngopharynx Area	LaryngopharynxArea
14	The distance between the most anterior and posterior points along the midline of the most superior border of the Laryngopharynx Volume.	Laryngopharynx Antero-posterior Distance	LaryngopharynxAPDist
15	The distance between the most lateral left and right points along the midline of the most superior border of the Laryngopharynx Volume.	Laryngopharynx Width	LaryngopharynxWidth
	Pharynx		
16	Orthogonal volume of the supraglottal region bounded superiorly by the intersections of centerline with the palatal plane, and inferiorly by the glottis.	Total Pharynx Volume	PharynxVolume
17	The curvilinear length along the centerline of the Pharynx Volume.	Pharynx Length	PharynxLength
18	CSA of the superior border/plane of the Subglottal Volume.	Glottis Area	GlottisArea
19	The distance between the most anterior and posterior points along the midline of the most superior border of the Subglottal Volume.	Glottis Anterior-Posterior Distance	GlottisAPDist
20	The distance between the most lateral left and right points along the midline of the most superior border of the Subglottal Volume.	Glottis Width	GlottisWidth
26	The curvilinear distance extending from the posterior aspect of the maxillary incisor teeth (ABI, landmark 23) through the oral and pharyngeal cavities to the level of the glottis. (i.e. traditional VTL measures starting at incisors).	Vocal Tract Length$_i$	VTLength$_i$
27	Curvilinear length of the velum from the PNS to the tip of the velum (VeEnd).	Velum Length	Velum Length
28	Length of the left piriform sinus measured as the 3D distance from the midpoint of the left aryepiglottic fold and the most inferior aspect of the left piriform sinus.	Piriform Sinus Length Left	PSLengthLeft
29	Length of the right piriform sinus measured as the 3D distance from the midpoint of the right aryepiglottic fold and the most inferior aspect of the right piriform sinus.	Piriform Sinus Length Right	PSLengthRight
30	Average length of the piriform sinus (average of PSLengthLeft and PSLengthRight if both exist, or the length of either if one is missing).	Average Piriform Sinus Length	AveragePSLength
	Subglottal		
21	Orthogonal volume of the region bounded superiorly by the glottis, and inferiorly by the opening of the trachea.	Subglottal Volume	Subglottal
22	The curvilinear segment length along the centerline of the Subglottal Volume	Subglottal Length	SubglottalL
23	CSA of the most inferior border of the Subglottal Volume, below the cricoid bone at the level of the first tracheal ring.	Trachea Area	TracheaArea
24	The distance between the most anterior and posterior points along the midline of the most inferior border of the Subglottal Volume.	Trachea Anterior-Posterior Distance	TracheaAPDist
25	The distance between the most lateral left and right points along the midline of the most inferior border of the Subglottal Volume.	Trachea Width	TracheaWidth

250 The 30 upper airway variables examined. Measurements extracted for each region include: the

251 orthogonal volume, curvilinear/centerline volume-length, the orthogonal superior or inferior

252 cross-sectional area (CSAs) of each of the subregions, as well as the anterior-posterior distance

253 (APDist) and lateral width (Width) of each CSAs. All planar measurements are orthogonal to the

254 centerline. The sum of the nasopharynx, oropharynx and laryngopharynx subregions was used to

255 calculate pharynx volume and pharynx length (measurements 1-to-25). See text for additional

256 vocal tract (VT) measurements (measurements 26-30).

d. Pharynx centerline and data extraction protocol

A semi-automatic, centerline-based data extraction pipeline was developed to extract quantitative

measurements from the 3D pharynx in MATLAB (The MathWorks, Natick, MA). First, the

built-in marching-cube algorithm in MATLAB was used to generate 3D meshes of the pharynx

model to serve as input to the pipeline [72]. This pipeline adapted the implicit fairing diffusion

method to smooth the 3D pharynx meshes iteratively while preserving the intrinsic geometric

properties [73]. Next, a level-contour-based centroid-extraction method was applied on the

smoothed pharynx, obtaining a set of coordinates along the tubular center of the pharynx [74-76].

These coordinates were further interpolated and smoothed with the B-spline de Boor algorithm,

generating a centerline representative of the center of the airway lumen [77, 78]. This centerline

was then used as input to an in-house written script that calculated planes orthogonal (i.e.,

perpendicular) to the line segment formed by each centerline coordinate and its subsequent

centerline coordinate. Finally, the intersections between the orthogonal planes and the 3D

meshes were then extracted as boundary vertices. With the boundary vertices, cross sectional

areas (CSAs) as well as additional variable measurements were calculated along the centerline.

See Fig 2 for an illustration of the 3D pharynx model and the cross sections.

e. Variable measurements

A total of 30 airway variables, as listed and defined in Table 5, were measured by the above-

described protocol using planes orthogonal to the centerline. The variables extracted are

described below and include overall pharyngeal length and volume, modified vocal tract length

(VTLength$_i$), velum length, and piriform sinuses length measurements, as well as measurements

from the following four subregions: (i) Nasopharynx, (ii) Oropharynx, (iii) Laryngopharynx, and

(iv) Subglottal. See Fig 1. Each subregion was isolated using its respective 'landmark-derived

planes' orthogonal to the centerline using the following boundary definitions: The nasopharynx

region was defined as an orthogonal volume bound by the intersection of the centerline with the

palatal plane –formed by the anterior nasal spine (ANS) and posterior nasal spine (PNS)

landmarks– superiorly, and with the tip of the velum inferiorly. The oropharynx region was

defined as an orthogonal volume bound by the orthogonal planes at the tip of the velum

superiorly, and by the aryepiglottic fold inferiorly. The laryngopharynx region, that includes the

piriform sinuses, was defined as the orthogonal volume bound by the orthogonal planes at the

midpoint of the aryepiglottic folds superiorly (the most curved point at approximate halfway

between the base and tip of the aryepiglottic folds), and by the glottis inferiorly. The subglottal

region was bound superiorly by the inferior boundary of the laryngopharynx region, and

inferiorly by the first axial slice displaying the first tracheal ring. The first tracheal ring was used

as a guide to the inferior border of the cricoid cartilage as the unossified cricoid cartilages in

pediatric cases was difficult to delineate on the CT images [79].

Measurements extracted, as defined in Table 5, included for each region: the orthogonal

volume, curvilinear/centerline volume-length, the orthogonal superior or inferior cross-sectional

areas (CSAs, five total) of each of the subregions, as well as the anterior-posterior distance

(APDist) and lateral width (Width) of each CSAs. The sum of the nasopharynx, oropharynx and

laryngopharynx supraglottal subregions was used to calculate pharynx volume and pharynx

length (Table 5, measurements 16 and 17). Additional vocal tract (VT) measurements (Table 5,

measurements 26-to-30) included: Vocal Tract Length$_{incisor}$ (VTL$_i$); calculated as the curvilinear

distance extending from the posterior border of the maxillary incisor (seen as the most anterior

landmark in Fig 1) to the glottis, representing the vocal tract portion of the upper airway starting

at the incisor i.e. excluding the lip and teeth region. Velum length; calculated as the curvilinear

distance extending from the PNS to the tip of the velum (VeEnd). Piriform sinus length

(PSLength); left, right, and average PSLength, measured using the defined anatomical landmarks

(PSSuL/R and PSInL/R).

f. Statistical analysis

311   All statistical analyses were performed in R. A linear mixed-effect model was used to capture

312   sex-specific growth in young children and allow for developmental comparison with adult

313   pharyngeal morphology. This model, using the lmerTest package for mixed-effects in R,

314   accounted for the repeat scans included in the dataset from individuals with multiple visits. The

315   model was specified as follows:

317
$$\hat{y} = \beta_0 + \beta_1 Sex + \beta_2 Adult + \beta_3 Sex \cdot Adult + \beta_4 PediatricAge +$$

318
$$\beta_5 Sex \cdot PediatricAge + \alpha_i$$

with "Adult" a dummy variable for adult subjects, "PediatricAge" the age for non-adult subjects

(0 for adults), and α_i a random per-subject effect.

Outliers were first excluded using the model, by removing data points with residuals

exceeding 2.576 of standard deviation, as described in [23, 71]. The model was then refitted on

log-scale for each of the variables to assess for growth trends and sex differences.

Likelihood ratio test (LRT) was conducted to assess overall age effect in the first five

years of life. To assess sex-differences, Wald test was performed at three time points: age-groups

< 1 year, 5-years and adults. Tests were conducted at a nominal significance level of $\alpha = 0.05$; in

Table 6, significance at the stringent Bonferroni corrected level (<.0004) was also indicated.

Finally, using point estimate of modeled means, percent growth at age 5-years was calculated

using data at age- group < 1 year, and adults for the purpose of gaining insight on upper airway

growth type as described by Scammon [80]. Scammon determined two primary postnatal growth

types, neural and general growth types, or their combination, to characterize growth of head and

neck structures. Furthermore, he noted that while all primary growth types are characterized by a

period of rapid growth during infancy, by early childhood neural growth type achieves greater

than two-thirds of the adult size, while somatic growth type barely achieves a quarter of the adult

size [80].

Results

Measurements extracted for males and females are displayed in Figs 3A-C for each pharyngeal

subregion (naso- oro- laryngo- pharynx), Fig 3D for the entire pharynx (supraglottal region), and

Fig 3E for the subglottal region with sex-specific linear fits and confidence intervals at age <1

year, 5-years, and adults. The plots also include a second y-axis depicting the percent growth of

adult size. Statistical analysis results are also summarized numerically in Table 6. Significance at

the .05, < .01, < .001, and Bonferroni corrected < .0004 levels are marked with one, two, three

and four asterisks respectively in Table 6.

Fig 3. Measurements extracted for pharyngeal region (3A. Nasopharynx; 3B. Oropharynx;

3C. Laryngopharynx; 3D. Pharynx; and 3E. Subglottal). Measurements extracted for each

pharyngeal region (3A to 3E) is depicted in top panel image on left, with measurements as

defined in Table 5 and consisting of: Top panel; the orthogonal volume, and the

curvilinear/centerline volume-length. Bottom panel; the orthogonal cross-sectional area (CSA)

(superior 3A, or inferior 3B-3E), its anterior-posterior distance (APDist), and lateral width

(Width). Plots include measurements for male in blue filled square symbols, and for female in

red open circle symbols. Pediatric data include linear fits for males (blue solid line) and females

(red dashed line). Point estimate of modeled means and confidence intervals are plotted for adult

data, and at ages 0 and 5 years respectively for males (purple) and female (magenta). The second

Y-axis reflects the percent growth for males (blue, inwards tick orientation) and females (black,

outwards tick orientation). Significance for sex differences at birth, age five and/or adults are

denoted with single, double, and triple asterisk(s) at the interval plots using the nominal $\alpha < 0.05$,

< 0.01, and < 0.001 levels respectively; the numeric p values are displayed in Table 6.

Table 6. Likelihood ratio test and Wald test results.

Measure-ments Types	Variable	Age Effect LRT p value	Sex Effect - Wald Test									Percent Growth at Age 5	
					p value			p value			p value		
		Pediatric	M	F	Age <1	M	F	Age 5	M	F	Adult	M	F
Volume (mm³)	Nasopharynx	0.0006 ***	807.28	927.40	0.6125	2453.77	2105.76	0.5155	5221.76	4255.25	0.3591	51.96	47.51
	Oropharynx	0.0000 ****	304.42	185.39	0.1546	879.79	1338.51	0.1570	7460.16	4927.30	0.1485	22.28	47.01
	Laryngopharynx	0.0000 ****	265.04	296.85	0.6296	1064.72	1188.46	0.5845	6168.93	4215.44	0.0464 *	31.79	41.35
	PharynxVolume	0.0000 ****	1450.67	1480.53	0.9229	4597.95	5257.92	0.4566	19742.95	13573.47	0.0288 *	33.80	48.10
	Subglottal	0.0000 ****	132.25	154.83	0.3167	670.82	722.98	0.5761	5998.06	3463.51	0.0000 ****	27.99	36.94
Volume Segment Length (mm)	NasopharynxL	0.0010 **	20.68	24.28	0.0800	30.15	29.74	0.8622	35.92	32.71	0.2105	62.14	64.75
	OropharynxL	0.0004 ****	6.26	3.80	0.0635	10.35	15.14	0.0972	37.05	30.24	0.3612	13.30	42.89
	LaryngopharynxL	0.0017 **	13.11	13.32	0.8246	16.75	16.97	0.8364	27.60	22.01	0.0002 ****	25.12	41.93
	PharynxLength	0.0000 ****	40.54	42.09	0.4439	58.20	61.78	0.1530	101.65	84.68	0.0000 ****	28.90	46.23
	SubglottalL	0.0000 ****	7.47	10.07	0.0106 *	14.66	15.12	0.7570	28.00	26.44	0.5538	35.04	30.86
Area (mm²)	NasopharynxArea	0.0868	54.65	39.11	0.2446	74.51	83.92	0.6260	248.52	218.61	0.5852	14.80	34.08
	OropharynxArea	0.0434 *	56.37	53.05	0.8100	100.62	94.33	0.7651	171.08	140.44	0.3344	45.28	53.18
	LaryngopharynxArea	0.0014 **	48.66	56.11	0.5462	111.26	123.17	0.6120	344.29	276.37	0.2512	30.85	39.50
	GlottisArea	0.0000 ****	11.28	5.65	0.0598	42.56	49.90	0.6103	121.90	74.74	0.1079	41.20	74.77
	TracheaArea	0.0000 ****	23.20	24.41	0.6389	53.31	51.56	0.7178	238.62	150.92	0.0000 ****	23.37	30.49
Anterior-Posterior Distance (mm)	NasopharynxAPDist	0.2087	5.41	3.61	0.0793	5.89	6.07	0.8801	14.47	12.40	0.4132	5.37	27.93
	OropharynxAPDist	0.0094 **	6.47	7.49	0.4237	10.44	12.38	0.2730	12.82	11.55	0.4807	62.44	120.23
	LaryngopharynxAPDist	0.3702	7.22	8.23	0.3171	8.91	9.03	0.8998	14.48	12.69	0.2206	23.26	18.01
	GlottisAPDist	0.0000 ****	5.71	5.23	0.2835	9.78	10.48	0.3147	19.93	14.35	0.0000 ****	28.66	57.59
	TracheaAPDist	0.0001 ****	5.47	5.27	0.5954	7.52	6.85	0.1295	15.97	12.30	0.0000 ****	19.48	22.50
Width (mm)	NasopharynxWidth	0.0086 **	11.37	10.35	0.5211	16.12	16.78	0.7464	22.31	23.03	0.7880	43.42	50.69
	OropharynxWidth	0.0163 *	8.31	8.81	0.7408	14.65	12.61	0.3252	19.60	15.60	0.1167	56.20	55.91
	LaryngopharynxWidth	0.0000 ****	6.39	5.75	0.3324	10.35	12.00	0.1166	20.99	18.35	0.1341	27.11	49.58
	GlottisWidth	0.0000 ****	2.58	1.76	0.1098	6.23	6.73	0.7073	8.92	7.16	0.2674	57.58	92.06
	TracheaWidth	0.0000 ****	5.20	5.38	0.6007	8.52	8.75	0.6373	17.64	14.67	0.0005 ***	26.72	36.27
VT Additional Length (mm)	VTLength	0.0000 ****	78.88	76.92	0.4097	108.35	111.13	0.3312	159.81	141.01	0.0000 ****	36.42	53.38
	VelumLength	0.0000 ****	25.19	25.13	0.9533	33.61	34.40	0.5550	42.41	38.92	0.0206 *	48.85	67.22
	PSLengthLeft	0.3470	6.13	5.97	0.9279	9.01	8.26	0.6893	17.68	14.13	0.2472	24.99	28.02
	PSLengthRight	0.1376	6.01	5.79	0.8925	10.18	9.41	0.7050	20.74	15.64	0.1262	28.35	36.69
	AveragePSLength	0.0768	5.70	5.95	0.8571	9.80	8.76	0.5465	19.33	14.94	0.1376	30.09	31.31

Likelihood ratio test (LRT) results for age effect, and sex-effect using the Wald test at age <1 year, 5 years, and adults. Significant differences are denoted with asterisk (* <.05; ** <.01; *** <.001; **** <.0004 Bonferroni corrected value). Also, percent of adult size at age 5 years (final column) for each of the 30 variables as listed by measurement type (column 1), and airway sub-regions (column 2). Percentages are based on the point estimate of modelled means, see Fig 3 for 95% confidence intervals. The regions are listed

366   superior to inferior with supra-laryngeal (above glottis) measurements listed first. Refer to Fig 1 to visualize variables and subregions,

367   and Table 5 for variable definitions.

a. Age Effect

As expected, likelihood ratio test results confirmed that all airway measurements for total and subregion volume, length (including VTL_i), and width (lateral) exhibited statistically significant growth in size during the first five years of life (< 1 year to 5-years) (Table 6). Also, four of the five CSAs examined displayed significant growth in size, except for the CSA at the level of the nasopharynx ($p = 0.0868$). Similarly, the linear anterior-posterior distance (APDist) measurements displayed significant growth except for the two measurements at the level of the nasopharynx and the laryngopharynx ($p = 0.2087$ and 0.3702 respectively). Limited growth was noted for average piriform sinus length (AveragePSLength; $p = 0.0768$), but growth in Velum Length was highly significant ($p < .0001$).

Compared to the mature adult airway, both male and female pediatric upper airway dimensions by age 5-years were significantly smaller (with higher percent growth in females than males as discussed below). However, one exception was the oropharynx anterior-posterior distance (APDist) and to some extent oropharynx width, where by age 5, children had essentially attained their adult size (see Fig 3B).

b. Sex Effect

The Wald test performed on pediatric data indicated that only subglottal length (SubglottalL) showed significance ($p = .0106$) at age <1 year with females' mean length being longer than males (see Table 7 mean (s.d.); M=8.06 (3.02) mm; F=10.56 (3.17) mm). No statistical significance was detected for any other variable at age-group <1 year or age-group 5-years. However, sex differences – though not significant – were noted at age group < 1 year in the volume-length of the nasopharynx ($p = 0.08$) and the oropharynx ($p = 0.0635$) subregions with

females having longer measurement than males; also, differences in nasopharynx APDist ($p =$

0.0793) and glottis area ($p = 0.0598$) with males having larger measurements than females. By

age 5-years, insignificant differences in the volume-length of the oropharynx subregion was

noted ($p = 0.0972$) with females having slightly longer measurements than males (see Table 7;

M=10.43 (3.57) mm; F=14.28 (7.21) mm).

As for adults, sexual dimorphism was highly significant for overall pharynx length and

VTL_i ($p < .0001$; M=161.06 (7.03) mm; F=141.14 (6.53) mm), and significant for pharynx

volume ($p < .05$; M= 53.73 (16.43) cm^3; F= (44.84 (15.17) cm^3), with males having larger

measurements than females (see Tables 6 and 7). Sexual dimorphism was also present in

laryngopharynx length ($p < .0004$) and volume ($p < .05$), as well as subglottal volume (p

<.0004), with highly significant differences in Glottis APDist ($p < .0004$) superiorly and CSA at

tracheal ring inferiorly ($p <.0004$; including differences in tracheal APDist ($p <.0004$) and width

$p=.0005$) with males having larger measurements than females.

Table 7. Age-specific mean (standard deviation) of the different measurement types for

each pharyngeal region.

Types	Variables	Sex	<1 (00;00 - 00;11)	1 (01;00 - 01;11)	2 (02;00 - 02;11)	3 (03;00 - 03;11)	4 (04;00 - 04;11)	5 (20;00 - 30;00)
					Age Groups (Age range (yr;mos))			
Volume (mm³)	Nasopharynx	M	1027.75 (681.71)	1082.45 (521.43)	2184.46 (879.28)	1709.06 (629.70)	2097.12 (699.82)	5372.79 (1543.29)
		F	1022.04 (853.53)	1488.75 (335.17)	2100.80 (1073.60)	1456.65 (640.11)	2232.27 (588.07)	4483.83 (1516.60)
	Oropharynx	M	380.64 (140.79)	353.09 (162.78)	727.05 (335.33)	1021.23 (751.86)	754.96 (614.13)	8025.12 (2397.84)
		F	299.93 (250.23)	482.34 (304.17)	613.47 (280.78)	796.34 (266.50)	1359.76 (1123.66)	5179.36 (1811.89)
	Laryngopharynx	M	412.27 (154.59)	333.90 (96.72)	607.52 (405.70)	866.48 (380.82)	868.67 (317.73)	6588.71 (2894.12)
		F	328.35 (179.26)	571.61 (174.32)	666.87 (321.22)	873.74 (489.16)	1009.18 (285.87)	4262.30 (681.45)
	PharynxVolume	M	1820.66 (823.81)	1769.44 (621.23)	3519.03 (1227.26)	3596.77 (1582.71)	3720.76 (1282.02)	19986.62 (3401.72)
		F	1650.32 (963.36)	2542.70 (573.50)	3381.15 (1451.68)	3552.66 (1430.01)	4601.21 (881.81)	13925.49 (3407.88)
	Subglottal	M	181.69 (88.19)	231.53 (67.30)	316.81 (93.86)	396.33 (133.49)	617.19 (203.74)	6023.72 (1161.30)
		F	179.72 (47.69)	264.70 (34.89)	321.03 (80.81)	526.63 (107.58)	533.77 (93.47)	3572.99 (979.31)
Volume Segment Length (mm)	NasopharynxL	M	22.19 (4.35)	23.12 (5.24)	26.32 (5.36)	27.61 (5.13)	27.88 (3.48)	36.13 (4.16)
		F	22.39 (4.64)	30.18 (2.29)	29.50 (4.07)	26.26 (3.37)	29.01 (2.32)	33.03 (4.65)
	OropharynxL	M	6.71 (1.87)	7.95 (2.02)	9.34 (4.08)	9.58 (5.01)	10.43 (3.57)	38.57 (8.49)
		F	7.57 (6.03)	5.72 (3.70)	6.51 (1.57)	11.32 (3.32)	14.28 (7.21)	30.79 (6.18)
	LaryngopharynxL	M	13.52 (2.14)	14.81 (1.67)	14.14 (2.25)	15.65 (1.40)	16.56 (1.75)	27.52 (2.79)
		F	13.79 (2.92)	14.33 (2.28)	15.31 (1.62)	15.94 (2.14)	16.53 (2.69)	22.08 (1.81)
	PharynxLength	M	42.42 (5.27)	45.75 (5.42)	49.09 (4.24)	52.85 (6.31)	54.80 (4.26)	101.83 (6.48)
		F	43.59 (6.95)	50.22 (3.20)	51.32 (3.15)	53.53 (4.60)	59.81 (4.55)	84.87 (6.16)
	SubglottalL	M	8.06 (3.02)	9.74 (1.85)	12.41 (3.18)	10.96 (2.15)	14.20 (2.62)	28.17 (2.69)
		F	10.56 (3.17)	11.97 (1.97)	11.68 (2.19)	14.40 (1.91)	13.68 (2.01)	26.65 (3.68)
Area (mm²)	NasopharynxArea	M	59.01 (22.48)	58.16 (24.12)	93.06 (39.75)	60.69 (31.72)	89.03 (41.34)	256.15 (92.03)
		F	47.54 (20.78)	52.03 (18.20)	71.78 (31.60)	67.30 (34.31)	99.45 (40.13)	227.33 (69.79)
	OropharynxArea	M	67.66 (22.94)	51.51 (23.51)	118.30 (54.47)	94.23 (51.27)	87.63 (21.59)	176.34 (42.19)
		F	51.52 (32.64)	96.06 (13.02)	79.93 (40.44)	74.60 (19.79)	91.80 (14.02)	145.07 (41.00)
	LaryngopharynxArea	M	62.05 (22.29)	53.06 (19.45)	93.10 (44.07)	117.89 (49.90)	81.66 (33.04)	355.86 (78.68)
		F	55.90 (24.64)	92.19 (12.97)	103.54 (31.96)	93.93 (28.32)	123.42 (53.04)	278.90 (39.83)
	GlottisArea	M	19.29 (5.79)	14.66 (4.33)	18.23 (9.49)	30.01 (8.85)	42.59 (12.77)	128.42 (53.84)
		F	11.58 (4.61)	11.88 (10.45)	21.27 (5.55)	30.63 (7.69)	36.82 (5.11)	80.05 (27.61)
	TracheaArea	M	27.24 (10.18)	31.82 (5.90)	34.70 (7.73)	42.59 (6.76)	47.83 (6.40)	238.74 (31.36)
		F	28.22 (10.35)	30.88 (3.93)	31.71 (4.79)	45.02 (7.12)	43.96 (5.21)	152.23 (22.33)
Anterior-Posterior Distance (mm)	NasopharynxAPDist	M	5.40 (2.21)	6.25 (2.25)	7.94 (3.11)	4.83 (1.69)	6.68 (2.00)	14.62 (4.34)
		F	4.35 (1.22)	4.11 (0.87)	4.43 (1.73)	5.41 (2.37)	6.94 (3.49)	12.77 (3.50)
	OropharynxAPDist	M	7.02 (2.85)	7.59 (3.33)	9.92 (2.74)	9.81 (4.82)	9.63 (2.78)	13.28 (3.44)
		F	7.11 (1.53)	9.39 (1.60)	10.11 (1.86)	11.77 (2.36)	10.99 (3.71)	11.95 (3.33)
	LaryngopharynxAPDist	M	7.65 (1.47)	7.08 (0.74)	9.20 (2.96)	9.26 (2.20)	7.81 (1.23)	14.63 (1.80)
		F	7.52 (1.37)	9.63 (1.95)	10.32 (2.58)	8.92 (1.23)	9.04 (2.85)	12.83 (2.03)
	GlottisAPDist	M	7.09 (1.27)	6.08 (0.61)	7.03 (1.68)	8.47 (0.90)	9.60 (1.26)	19.98 (1.60)
		F	5.41 (1.20)	6.60 (1.26)	7.72 (0.69)	8.82 (0.60)	9.08 (0.55)	14.44 (1.68)
	TracheaAPDist	M	5.79 (1.07)	6.16 (0.85)	6.36 (1.23)	7.07 (0.69)	7.06 (0.72)	16.01 (1.21)
		F	5.75 (1.19)	5.45 (0.46)	5.74 (0.70)	6.81 (0.98)	6.40 (0.52)	12.33 (0.94)
Width (mm)	NasopharynxWidth	M	12.54 (4.58)	12.16 (3.97)	15.64 (2.86)	14.66 (4.48)	15.73 (3.65)	22.50 (3.02)
		F	10.90 (3.06)	12.84 (3.02)	14.23 (2.90)	13.82 (3.71)	16.99 (3.18)	23.27 (3.40)
	OropharynxWidth	M	10.06 (3.98)	8.09 (3.56)	13.82 (2.63)	12.90 (4.34)	13.51 (3.42)	19.77 (2.85)
		F	7.60 (2.29)	12.26 (1.51)	12.79 (4.53)	10.73 (4.15)	12.76 (1.20)	15.72 (2.15)
	LaryngopharynxWidth	M	8.43 (1.55)	6.29 (1.49)	8.26 (1.43)	8.41 (1.48)	10.67 (2.04)	21.08 (2.04)
		F	5.96 (1.56)	7.44 (1.65)	8.73 (1.50)	10.01 (1.24)	10.58 (2.04)	18.52 (2.68)
	GlottisWidth	M	3.60 (0.79)	3.17 (0.94)	3.58 (1.28)	4.95 (1.54)	6.10 (1.21)	9.13 (3.02)
		F	2.78 (0.47)	2.45 (1.81)	3.65 (0.67)	4.90 (1.01)	5.66 (0.76)	7.57 (2.50)
	TracheaWidth	M	5.62 (1.37)	6.23 (0.49)	6.74 (0.96)	7.31 (0.83)	8.10 (0.49)	17.66 (1.18)
		F	5.57 (1.00)	6.63 (0.52)	6.63 (0.85)	7.76 (0.41)	7.89 (0.63)	14.72 (1.27)
VT Additional Length (mm)	VTLength	M	80.97 (5.11)	87.12 (3.85)	94.53 (6.37)	98.09 (6.00)	103.03 (5.02)	161.06 (7.03)
		F	78.63 (7.79)	89.66 (4.34)	93.46 (2.80)	100.11 (5.99)	103.86 (6.34)	141.14 (6.53)
	VelumLength	M	26.49 (1.46)	26.96 (2.40)	30.32 (3.54)	29.86 (3.32)	32.97 (1.15)	42.60 (4.21)
		F	24.46 (1.72)	29.62 (1.73)	30.92 (2.29)	30.65 (2.01)	32.51 (0.83)	39.00 (2.67)
	PSLengthLeft	M	6.87 (0.85)	6.63 (2.93)	8.41 (3.31)	8.35 (2.33)	8.90 (2.99)	17.96 (5.27)
		F	5.35 (3.99)	10.73 (3.25)	8.89 (4.41)	8.46 (3.01)	8.26 (3.61)	14.43 (3.16)
	PSLengthRight	M	6.84 (1.50)	5.69 (1.55)	8.54 (3.38)	10.49 (1.85)	8.76 (3.06)	20.71 (3.23)
		F	6.47 (4.12)	10.93 (3.63)	7.65 (4.70)	9.30 (2.87)	9.34 (2.71)	15.81 (2.55)
	AveragePSLength	M	6.53 (1.21)	5.75 (1.69)	8.47 (3.31)	9.26 (1.70)	8.82 (2.82)	19.33 (3.86)
		F	5.91 (4.04)	10.83 (3.44)	8.27 (4.53)	8.58 (3.04)	8.87 (2.76)	15.12 (2.61)

Age-specific mean (standard deviation) of the different measurement types for each of the variables examined with M/F denoting the average Male/Female measurements. Age groups as specified in Table 3.

c. Percent Growth

As displayed in second y-axes of Fig 3 plots, and listed in Table 6 (final column), percent growth assessment based on modeled point estimates, revealed that by age five-years, female upper airway measurements were closer than male measurements to the adult mature size in 26 out of the 30 upper airway variables examined. (See the tabulated average data per age-group in Table 7.) Female data revealed 9 out of 30 variables to have reached over 50% of adult size nasopharynx volume-length (64.75%), nasopharynx width (50.69%), oropharynx area (53.18%), oropharynx APDist (120.23%), oropharynx width (55.91%), glottis area (74.77%), glottis APDist (57.59%), glottis width (92.05%), and VTL_i (53.38%). In contrast, males had only 5 of the 30 variables reach over 50% of the adult mature size: nasopharynx volume (51.96%), nasopharynx volume-length (62.14%), oropharynx APDist (62.44%), oropharynx width (56.20%), and glottis width (57.58%). The only 4 measurements where both male and female growth reached over 50% of their respective adult size were: nasopharynx length, oropharynx APDist, oropharynx width, and glottis width.

Discussion

This study addresses a void in normative data on the upper airway during the first five-years of life. After developing a protocol that controls for variables that can affect measurement accuracy (e.g., head position, sedation, threshold for airway segmentation), CT studies from 61 typically developing pediatric and 17 adults were used to quantify the multidimensional growth of the airway systematically with respect to age and sex. Our findings are novel in that, to our

knowledge, this is the first study that examines the birth-to 5 years age range, as compared to

adults, using a comprehensive set of 2D and 3D measurements from the choanae to below the

cricoid ring (opening to the trachea), including: supra- and sub-glottal volume and length, naso-

oro-laryngo-pharynx subregion volume and length, each subregion's superior and inferior CSA,

and their antero-posterior and transverse/width distances. Additionally, the data were collected

using a protocol that included a well-defined and established threshold for airway segmentation,

and a semi-automatic centerline that we developed for the extraction of accurate measurements

to quantify the upper airway using the natural anatomic orientation of airflow for respiration and

speech production.

The use of a centerline, as an added methodological consideration, is critical for

obtaining accurate measurements of the airway. As summarized in Table 1, two studies [35, 49]

have used a centerline to quantify the airway, but Chiang et al. [35] is the only study to date that

performed a centerline-based technique to obtain quantitative data on the growth and

development of the airway. However, their measurements stopped at the level of the epiglottis,

and they did not include the pre-pubertal age group.

The upper airway subregion dimensions are sensitive to altered head and tongue posture,

particularly for 3D assessment but also for 2D measurements as Gurani et al. [54] point out.

Given the need for a valid method to classify head position of imaging studies, our laboratory

first developed a reliable protocol that uses 14 landmarks to account for both head and neck

positions [68]. We therefore first employed this protocol for the selection of cases with a neutral

head position for inclusion in this study, then applied the centerline protocol.

Given all the methodological considerations we accounted for, the attrition rate of cases

included in this study from the imaging studies available in our database was high. We retained

only 19% of the cases reviewed. Given this rigorous approach to control for positioning and

other potential confounder, we expect our findings to reliably reflect typical airway growth.

Furthermore, we anticipate that the inclusion of additional cases in future studies, using the

above-described airway data extraction protocol, will further strengthen present findings and

observations.

a. Age Differences

Our findings reflect persistent positive increase in size for all variables examined during the first

five years of life for all measurements in all subregions as displayed in Fig 3, with the means per

age group summarized in Table 7. The age effect of the likelihood ratio test confirmed the

significant growth in size for 24 out of the 30 variables studied with some variables displaying

more rapid and extensive growth than others. Such findings provide insight on the proportional

and relational growth of upper airway dimensions with age during anatomic restructuring (e.g.,

hyo-laryngeal descent).

As expected, all pediatric airway dimensions were substantially smaller than adult

dimensions except oropharynx APDist and to some extent oropharynx width (see Fig 3).

Abramson et al. [20] also found that volume, CSA and transverse measurements –but not AP

dimensions– of the pediatric naso-oropharynx airway were significantly smaller than adult

measurements. This rapid and early maturation in oropharynx APDist dimension corresponds to

our previous research findings where growth of oral structures in the horizontal plane, in line

with neural growth, achieved most of their growth towards the adults size by age five-years [23,

66, 71]. Alternatively, though unlikely, it is possible that hypertrophy of lingual tonsils interfered

with lumen APDist measurements. As for growth in width, while not as fully developed as AP

dimensions by age 5, oropharynx lateral dimensions had reached over 55% of the adult,

suggesting that growth in this subregion undergoes a combination of neural and general growth

types, which Scammon had noted is present in the growth of structures in the neck region.

Volume, volume-length and width measurements increased with age for all pharyngeal

subregions, consistent with prior studies on upper airway development in infant and pre-pubertal

children [20, 24]. The CSA measurements in this study were extracted from anatomical

landmarks representing the superior and inferior borders of the pharyngeal subregions. This is in

contrast to the typical approach of measuring minimum or maximum CSA to examine sites of

constriction for assessment of patients' risk for OSAS [39, 40, 42, 43, 45, 49, 81]. Furthermore,

we used oblique planes – orthogonal to the centerline – which cannot be compared to studies that

used the axial plane, as in most of the above listed studies.

A factor that further complicates comparisons, including within-study cross-sectional

comparisons, is the hypertrophy of tonsils in young children that follow a lymphoid growth type.

In particular, nasopharyngeal tonsils referred to as adenoids, where hypertrophy is the highest in

4-6- year old children [82]. Keeping these issues in mind, among the five CSA measurements in

this study, the nasopharynx region was the only site that did not have a significant age-effect.

Similarly, the APDist measurements in the nasopharynx and laryngopharynx were the only sites

that did not have significant age effect between the ages < 1-year and 5-years. Such findings

could be attributed to adenoid hypertrophy, typically occurring between the ages 2-6 years, that

diminish airway dimensions [83-85]. The decrease of mean nasopharynx CSA and APDist

measurements per age group can be noted in Table 7, with changes most evident between the

ages 2-to-4 years in this study.

In all age groups, the APDist dimensions at the nasopharynx, oropharynx and trachea

(Fig 1a, b, e) were smaller than the width/transverse measurements, but larger than

width/transverse measurements at the glottis (Fig 1d, and Table 7). Such findings are in line with

Abramson et al. [20] who reported significant upper airway growth along the transverse

dimension with age, where the airway becomes more elliptical in shape. Similarly, Machata et al.

[26] using MRI studies of children ages 0-6 years, reported smaller anteroposterior dimensions

than transverse dimensions for all ages at the level of the soft palate, the base of the tongue, and

the tip of the epiglottis. In contrast, the laryngopharynx APDist dimensions (Fig 1c), were larger

than width/transverse measurements from birth to age 3, but became smaller than

width/transverse measurements at age 3 and beyond, which likely contributed to the absence of

age effect for APDist.

Changes in APDist versus width dimensions could be attributed to the cartilaginous

composition of the larynx. The laryngopharyngeal cross section in this study was designed to

capture its surrounding structures - the aryepiglottic folds on each side, the laryngeal vestibule

anteriorly, and posteriorly by arytenoid cartilages, corniculate cartilages and the interarytenoid

fold. Cartilage ossification is usually not observed until past age 20 years, and the pediatric

laryngopharynx region is often described to be "featureless" and difficult to assess using

commonly acquired medical images [79]. We employed an established method for airway

segmentation that uses image-specific airway thresholds, and therefore are confident that our

data is reflective of airway development. Since we used landmarks on the aryepiglottic folds that

connect to the piriform sinuses on each side of the cavity, aditus of larynx, the laryngopharynx

width/transverse measurement in this study excluded the piriform sinuses (see Fig 1c). The

significant age effect along this transverse dimension is therefore truly reflective of the strong

lateral growth in the laryngopharyngeal region.

The CSA and transverse width measurements at the level of the glottis are smaller than

the area and width at the first tracheal ring immediately inferior to the cricoid (i.e., subglottal

region); however, the average APDist measurement of the glottis is larger than the APDist

dimension at the level of the first tracheal ring at ages <1-year, 5-years, and in adults. This

finding is consistent with Luscan et al. [86], who concluded that "the cricoid has a round shape

regardless of the child's age." Indeed, the mean APDist and width measurements were very close

or similar for all ages at the level of the first tracheal ring proximal to the inferior border of the

cricoid, and comparable to the cricoid outlet's (to trachea) anteroposterior and transverse interior

diameters of Liu et al. [87]. While growth trends were comparable, our measurements were

closer to those of Liu et al [87] than to those of Luscan et al. [86], and indicate the importance of

methodological considerations, including having well-defined data extraction protocols such as

the determination of an appropriate threshold level (HU) to segment the airway.

Growth in VTLi was significant during the first five years of life, confirming an increase

of about 3 cm, which is consistent with VTL findings to date [23] and reflects that this modified

measure captured growth in both the oral and pharyngeal portions of the VT. The measurements

in this study were smaller than what has been reported previously, which is to be expected given

the modified length measure had an onset at the posterior margin of the incisors in lieu of the

typical anterior margin of the lips. Findings of a significant age effect on velum length were

comparable to values reported by Perry et al. [88] and Yi et al. [27]. Closer examination of the

developmental data on pharyngeal length and pharyngeal volume revealed a close relationship

particularly after about age 2. Before age 2, the growth rate was slightly more pronounced in

length than in volume, likely due to the drastic anatomic restructuring of the skeletal framework

in the region of the pharyngeal cavity, including hyo-laryngeal descent, and rapid neural growth

in length of the second cervical spine (C2) [68].

As for the piriform sinuses, our findings revealed a borderline average PSLength age

effect (p=.077) with average measurements per age-group summarized in Table 7. The pediatric

average PSLength measurements ranged from 6 to 9 mm at ages <1-year to 5-years; whereas the

adult average PSLength measurements ranged from 1.5 to 1.9 cm. While the development of

piriform sinuses has not been examined to date, and therefore comparative measurements were

not available, adult PSLength measurements were comparable to the 1.6 to 2 cm piriform sinus

depth measurements of Dang and Honda [89-91]. This similarity was despite the fact that our

PSLength measurement extended from the most inferior aspect of the piriform sinuses to the

midpoint of the aryepiglottic folds, which is beyond the arytenoid apex plane used by Dang and

Honda [89]. This could be in part due to differences in imaging modality used (CT vs MRI)

and/or segmentation thresholding levels used to obtain reliable airway measures, particularly

given the small size of this region of interest. Additional factors include methodological

differences (oblique vs, axial plane) in obtaining measurements, as well as the height of

participants, which has been shown to be related to vocal tract length [24] and pharyngeal

dimensions [47]. The piriform sinuses play an important role during swallowing by diverting

liquids around the aditus of the larynx and into the esophagus. They also affect speech acoustics

and attenuate the vocal tract resonant frequencies in adults [89, 92-94] by an estimated range of

5% of formant frequencies [89]. Thus, detailed developmental data on the piriform sinuses

would provide needed normative data and could help provide insight on pediatric dysphagia.

Furthermore, such data can be used to implement modeling [95] to systematically examine the

effect of the piriform sinuses on the resonances of the developing vocal tract, particularly given

the intriguing findings that formant frequencies reportedly remain stable during the first 24-to-36

months of life [14, 15, 96], despite documented increases in vocal tract length [2, 23, 24].

In summary, the upper airway dimensions reveal persistent growth during the first five

years of life, with some dimensions growing at a faster pace than others. Growth in the vertical

and transverse/lateral dimensions are more pronounced than growth in the AP dimension.

b. Sex Differences

As depicted in Table 6, sexual dimorphism was present in a number of supra- and sub-glottal

variables in adults. However, while there was evidence towards sexual dimorphism in all three

supra-glottal regions for a number of variables at age <1 year (specifically, larger nasopharynxL

and oropharynxL in females; also, larger nasopharynx APDist and glottis area in males), with the

larger oropharynxL persisting in females at age 5 years, none of these supra-glottal or pharyngeal

variables were significant in children.

As for the subglottal region, only subglottal volume-length displayed significant sexual

dimorphism at age < 1-year (with males shorter than females), but not at age 5-years or in adults

(see Table 6). To our knowledge, this specific subglottal volume-length measurement has not

been examined, despite its importance in procedures like tracheotomy [97], and laryngotracheal

infections/diseases including SIDS [98] where the incidence is higher in males [99]. However,

two studies have performed distance measurement in this region, specifically anterior

commissure to first tracheal ring [100], and vocal folds to the cricoid cartilage [101]. Contrary to

present findings, Khadivi and colleagues, who used laryngoscopy to collect subglottal length

data from 82 adults (57 males and 25 females), documented significant sexual dimorphism.

While our measurements for pediatric subglottal length are comparable to the normative values

reported by Sirisopana and colleagues, from the CT scans of 56 children (29 males, 27 females),

they unfortunately neither assessed for sex differences, nor reported sex-specific measurements

given their primary focus on tracheal tube design.

596   Despite methodological differences, the absence of sexual dimorphism in pediatric upper

airway data for most of the measurements analyzed in this study was consistent with past studies

[22, 24, 29]. Barbier et al. [29] did not find sex difference in pre-pubertal data but suggested that

sexual dimorphism in VTL emerged during puberty. Among studies reporting regional upper

airway normative data for the age range between 0-to-5-years, Abramson et al. [20] found no

difference in naso-pharyngeal airway size or shape between the sexes in children, but reported

longer airway length in post-pubertal males. Jeans et al. [83], using lateral cephalometric

radiographs to study the nasopharyngeal airway area – comparable to our nasopharyngeal region

–, however, reported mild decreases in nasopharyngeal area in both 3-to-5-year old males and 3-

to-6-year old females. Sex-differences in the same region using an anteroposterior distance

measure in the midsagittal plane of medical imaging studies (MRI & CT), referred to as

oropharyngeal-width, have similarly been noted to display evidence, albeit not significant,

towards sexual dimorphism in 3-to-4-year old children with males having larger width

measurements [71]. Linder-Aronson et al. [85] noted that nasopharynx airway depth/AP

dimension in males were consistently larger than females throughout ages 3-to-16 years. Sexual

dimorphism in the pharyngeal portion of the VT in ages 8-to-19 years, has also been reported by

Vorperian et al. [71], with the vertical nasopharyngeal length being longer in females than males

and the vertical posterior cavity length being longer in males than females. In contrast, Yi et al.

[27] reported no sex differences in any of their linear dimensions using MRI in infants and

children up to 72 months. Rommel et al. [34], who used curvilinear length drawn on 2D X-ray

images to assess naso-oropharynx segments, found no sex difference in children as young as 0-

to-4-years. Griscom [102] similarly found no significant sex differences in trachea dimensions

until late in adolescence. Definitive prepubertal sexual dimorphism of the pharynx thus cannot be

confirmed with studies available to date.

Detecting sex differences is a difficult task given the critical methodological

considerations outlined in our methods section and the importance of having a large number of

participants per age-group. Statistical analysis methods can overcome differences in growth rate

between males and females, such as implementing continuous-window comparisons across age

[e.g.,71]. This latter approach was particularly effective in unveiling sexual dimorphism that

does not persist during the course of development, since growth in females outpaces males

during early development, but growth in males begins to outpace females during the peripubertal

period, with sexual dimorphism emerging during puberty.

Sexual dimorphism in adults, however, was mostly present and aligned with research

findings to date in pharynx volume, pharynx length [47], VTL [71], velum length [103], glottis

APDist [47], subglottal volume [102], and tracheal dimensions [86]. The finding that subglottal

volume-length, the only measurement that displayed significant sexual dimorphism at age < 1-

year-old was not sexually dimorphic at age 5-years is not surprising, given growth rate

differences in males versus females as noted above. However, the absence of differences in

adults is likely due to both methodological differences and the limited number of adults in this

study (n=17), given our stringent inclusion criteria.

Although present findings revealed significant prepubertal sex-differences only in the

subglottal region, findings in this study and others as noted above, in the naso-oro-pharyngeal

region, provide sufficient justification to further examine this issue using a larger number of

cases particularly given the above noted critical methodological considerations, and the nuance

of growth rate differences between the two sexes. Such a conclusion is further supported by

auditory-perceptual and acoustic findings where Bloom et al. [104] reported that adults

accurately identified 3-month old infants' vocalizations as boy vs girl. The only acoustic

difference was the feature of nasality with girls' vocalizations being more nasal than boys.

Furthermore, several studies have reported sex differences in vowel formants (i.e., vocal tract

resonant frequencies) in children as young as 3 or 4 years of age [105-107].

c. Percent Growth

The growth pattern of anatomical structures in the craniofacial and upper airway region are

known to be non-uniform with the primary growth types in this region being neural,

general/somatic, and lymphoid. Since we did not include adenoid or tonsil measurements, we

will limit this discussion to the first two types. Both neural and general growth types display

rapid growth during the first few years of life. Scammon [80] summarized schematically,

however, that by age 5, the percent of the adult mature size reached was drastically different for

the neural (~80%) versus the general (< 40%) growth types. He also noted that growth in the

neck region could be a combination of both neural and general primary growth types [80]. The

final column of Table 6 presents the model-based point-estimates of percent growth of the adult

mature size by age five years for all thirty variables examined in this study.

The general finding that by age 5, female upper airway dimensions were larger than

males is not surprising since typically females have a faster growth rate during childhood and

reach the adult mature size sooner than males. Based on findings to date, structures in the upper

airway were expected to follow a mostly somatic growth type or a composite growth of somatic

and neural growth types [23, 108, 109]. Despite differences in VTL versus VTL*i* measurements

(where the onset of the former at the anterior margin of the lips and the latter is at the posterior

margin of the incisors), the general growth findings in this study are in line with the expected

growth trend indicating that VTL*i* growth type is predominantly hybrid somatic/neural in

females (53.38%) and somatic in males (36.42%) [23]. Similarly, pharynx volume, pharynx

volume-length and all other pharyngeal subdivision volume and volume-length results except for

the nasopharynx subregion confirmed the predominant somatic growth types at age 5-years, in

line with the reported growth patterns for pharyngeal cavity length and VT vertical data [23]. As

for the nasopharyngeal region, the expected hybrid somatic/neural or combination growth type,

was indeed reflected in present findings where both male and female nasopharynx volume and

volume-length measurements ranged between 47% and 65% of adult size by age 5-years. The

oropharynx AP dimension exceeded 60% for males and 85% for females, suggestive of a more

pronounced hybrid somatic/neural growth type for the males and neural growth type for the

females. These findings are in line with Vorperian et al. [23] where structures in oral region in

the horizontal plane reached maturation earlier than structures in pharyngeal region in the

vertical plane. Percent growth analysis (in Table 6, last column) also reflect the presence of sex-

specific differences in the combination of growth types for upper airway structures. Since

multiple factors contribute to growth, it is possible that such sex-specific differences further

contributed to the difficulty in detecting sexual dimorphism during early childhood.

Conclusions, Study Limitations, and Future Direction for Research

This study, using CT studies, provides data quantifying the 3D growth of the upper

airway with minimal methodological concerns for an age group with scant normative data.

Findings confirm persistent growth of the upper airway during the first five years of life with

growth in the vertical and transverse/lateral dimensions having a faster pace and greater

prominence than growth of anteroposterior dimension. Findings also reveal that at age 5, females

have larger airway dimensions than males. Such findings confirm the importance of studying

sex-specific developmental changes of the upper airway in 3D. A better understanding of

pharyngeal functions and disorders will require further, more detailed examination of the

developmental changes in pharyngeal length versus volume and the piriform sinuses, as well as

re-examination of prepubertal sex-differences.

Our painstaking efforts to optimize accurate and reliable upper airway measurements

limited the sample size. Specifically, the attrition rate of only retaining retrospective imaging

studies with a neutral head position was high given the head position protocol we applied [68].

Also, despite the imaging protocol at the University Hospital to maintain the head in midline, it

is likely that some rotation was present. However, given that the airway is functional during

imaging (breathing and swallowing), it is difficult to determine source of asymmetry as noted

with the piriform sinuses, where we resorted to averaging them in this study. Furthermore,

airway anatomic measurements can be affected by breathing phase where significant effect of

breathing phase in the oro-laryngo-pharyngeal region has been reported [34]. Such concerns,

including the above suggested assessments, could be addressed by replicating this study using a

larger sample size, and ideally increasing the age range to cover the entire developmental period

particularly ages 5-to-20 years.

Aside from using a larger sample, having relevant demographic information such as

height, weight, and race would be valuable to include. Given the retrospective nature of this

study, relevant demographic information was not available for all cases. Our imaging database

however, was representative of regional Dane County demographics, with growth between the

10th and 95th percentiles. We believe our present findings are representative of typical growth

since the natural variability in craniofacial dimensions within individual races is related to the

natural variability or variations within the different racial/ethnic groups [110].

Aside from the use of 3D anatomic landmarks and establishing standardized procedures

to minimize, if not eliminate, methodological limitations on measurement accuracy and

reliability, it is also imperative to establish standardized definitions of the pharyngeal subregions

using well-defined anatomical boundaries. This will facilitate accurate representation and

comparison of the developmental morphology of the aerodigestive and vocal tract, within and

across disciplines, including comparison between different imaging modalities. This will

undoubtedly enrich our understanding of the growth of a region that serves multiple life-

functions since each modality has its strengths and limitations. For example, while MRI provides

more accurate information on the growth of lymphoid tissue in the pharyngeal region, CBCT

could address concerns on the effect of body position and gravity on soft-tissue structures for

obtaining reliable airway dimensions. The feasibility of comparing developmental findings

across disciplines, including the individual and relational growth of structures that provide the

skeletal framework of the aerodigestive and vocal tract, will facilitate the understanding of upper

airway pathophysiology, improve surgical planning such as estimation of laryngeal mask airway

size or endotracheal tube diameter, evaluation of pharyngeal collapsibility in early childhood in

the assessment of OSAS, and other upper airway anomalies including swallowing difficulties

and speech disorders. Such information would also facilitate the advancement of developmental

models to assess various typical and atypical functions related to airflow, swallowing, and

speech production.

Acknowledgments

We gratefully acknowledge Lindell R. Gentry, MD for his assistance in establishing the Vocal

Tract Development Laboratory's imaging database. We thank Gabe Jardim for establishing a

protocol to obtain threshold for airway segmentation, and Drs. Andy Alexander and Michael

Speidel for their guidance in resolving and confirming the reliability of this protocol. We also

thank Drs. Moo K. Chung and Nagesh Adluru for their input on centerline development, Ben M.

Doherty for his guidance with the data extraction script, Sophie D. Blankenheim for assistance

with literature review and preparation of Table 1, and Dr. Jacqueline J. Houtman for comments

on an earlier version of this manuscript.

References

1.   Marcus CL, Smith RJH, Mankarious LA, Arens R, Mitchell GS, Elluru RG, et al., editors. Developmental aspects of the upper airway: Report from an NHLBI Workshop, March 5-6, 2009. Proceedings of the American Thoracic Society; 2009.

2.   Vorperian HK, Kent RD, Lindstrom MJ, Kalina CM, Gentry LR, Yandell BS. Development of vocal tract length during early childhood: A magnetic resonance imaging study. The Journal of the Acoustical Society of America. 2005;117:338-50. doi: 10.1121/1.1835958.

3.   Laitman JT, Crelin ES. Postnatal development of the basicranium and vocal tract region in man. In: Bosma J, editor. Symposium on the development of the basicranium. Bethesda, MD: DHEW Publicaiton No. 76-989, PHS-NIH; 1976. p. 206-20.

4.   Roche AF, Barkla DH. The level of the larynx during childhood. 1987:645-54.

5.   Enlow DH, Hans MG. Essentials of Facial Growth. Philadelphia: W.B. Saunders Company; 1996 Jun 1996. 303 p.

6.   Moss ML. The functional matrix hypothesis revisited. 1. The role of mechanotransduction. American Journal of Orthodontics and Dentofacial Orthopedics. 1997;112(1):8-11. doi: https://doi.org/10.1016/S0889-5406(97)70267-1.

7.   Moss ML. The functional matrix hypothesis revisited. 2. The role of an osseous connected cellular network. American Journal of Orthodontics and Dentofacial Orthopedics. 1997;112(2):221-6. doi: https://doi.org/10.1016/S0889-5406(97)70249-X.

8.   Moss ML. The functional matrix hypothesis revisited. 3. The genomic thesis. American Journal of Orthodontics and Dentofacial Orthopedics. 1997;112(3):338-42. doi: https://doi.org/10.1016/S0889-5406(97)70265-8.

9.   Moss ML. The functional matrix hypothesis revisited. 4. The epigenetic antithesis and the
resolving synthesis. American Journal of Orthodontics and Dentofacial Orthopedics.
1997;112(4):410-7. doi: https://doi.org/10.1016/S0889-5406(97)70049-0.
10.   Carlson DS. Theories of Cranofacial Growth in the Postgenomic Era. Seminars in
Orthodontics. 2005;11:172-83. Epub 2005.
11.   Standerwick RG, Roberts WE. The aponeurotic tension model of craniofacial growth in
man. Open Dent J. 2009;3:100-13. Epub 2009/07/03. doi: 10.2174/1874210600903010100.
PubMed PMID: 19572022; PubMed Central PMCID: PMC2703201.
12.   Castaldo G, Cerritelli F. Craniofacial growth: evolving paradigms. Cranio.
2015;33(1):23-31. Epub 2014/12/31. doi: 10.1179/0886963414Z.00000000042. PubMed PMID:
25547141.
13.   Lieberman DE. Epigenetic Integration, Complexity, and Evolvability of the Head. In:
Hallgrímsson B, Hall BK, editors. Epigenetics: Linking Genotype and Phenotype in
Development and Evolution: University of California Press; 2011. p. 271-89.
14.   Gilbert HR, Robb MP, Chen Y. Formant frequency development: 15 to 36 months.
Journal of Voice. 1997;11(3):260-6.
15.   Kent RD, Murray AD. Acoustic Features of Infant Vocalic Utterances at 3, 6, and 9
Months. Journal of the Acoustical Society of America. 1982;72(2):353-65. doi: Doi
10.1121/1.388089. PubMed PMID: WOS:A1982PB08400008.
16.   Bunton K, Leddy M. An evaluation of articulatory working space area in vowel
production of adults with Down syndrome. Clin Linguist Phon. 2011;25(4):321-34. Epub
2010/11/26. doi: 10.3109/02699206.2010.535647. PubMed PMID: 21091205.
17.   Kent RD, Vorperian HK. Speech Impairment in Down Syndrome: A Review. Journal of
Speech, Language, and Hearing Research. 2013;56(1):178-210. PubMed Central PMCID:
PMCNIHMSID: NIHMS 392496.
18.   Eslami E, Katz ES, Baghdady M, Abramovitch K, Masoud MI. Are three-dimensional
airway evaluations obtained through computed and cone-beam computed tomography scans
predictable from lateral cephalograms? A systematic review of evidence. Angle Orthod.
2017;87(1):159-67. Epub 2016/11/02. doi: 10.2319/032516-243.1. PubMed PMID: 27463700.
19.   Ono T, Otsuka R, Kuroda T, Honda E, Sasaki T. Effects of head and body position on
two- and three-dimensional configurations of the upper airway. Journal of Dental Research.
2000;79:1879-84. doi: 10.1177/00220345000790111101.
20.   Abramson Z, Susarla S, Troulis M, Kaban L. Age-related changes of the upper airway
assessed by 3-dimensional computed tomography. Journal of Craniofacial Surgery. 2009;20:657-
63. doi: 10.1097/SCS.0b013e318193d521.
21.   Smitthimedhin A, Whitehead M, Bigdeli M, Gustavo N, Geovanny P, Hansel O. MRI
determination of volumes for the upper airway and pharyngeal lymphoid tissue in preterm and
term infants. Clinical Imaging. 2018;50:51-6. doi: 10.1016/j.clinimag.2017.12.010.
22.   Ronen O, Malhotra A, Pillar G. Influence of Gender and Age on Upper-Airway Length
During Development. Pediatrics. 2007;120:e1028-e34.
23.   Vorperian HK, Wang S, Chung MK, Schimek EM, Durtschi RB, Kent RD, et al.
Anatomic development of the oral and pharyngeal portions of the vocal tract: An imaging study.
The Journal of the Acoustical Society of America. 2009;125:1666-78. doi: 10.1121/1.3075589.
24.   Fitch WT, Giedd J. Morphology and development of the human vocal tract: A study
using magnetic resonance imaging. The Journal of the Acoustical Society of America.
1999;106:1511-22. doi: 10.1121/1.427148. PubMed PMID: 10489707.

25.   Litman RS, Weissend EE, Shrier DA, Ward DS, editors. Morphologic Changes in the
Upper Airway of Children during Awakening from Propofol Administration. Anesthesiology;
2002.
26.   Machata AM, Kabon B, Willschke H, Prayer D, Marhofer P. Upper airway size and
configuration during propofol-based sedation for magnetic resonance imaging: An analysis of
138 infants and children. Paediatric Anaesthesia. 2010;20:994-1000. doi: 10.1111/j.1460-
9592.2010.03419.x.
27.   Yi X, Yao L, Yuan X, Wei Y, Wang Z. Linear dimensions of normal upper airway
structure by magnetic resonance imaging in Chinese Han infants and preschool children. Sleep
Medicine. 2017;37:98-104. doi: 10.1016/j.sleep.2017.06.011.
28.   Fregosi RF, Quan SF, Kaemingk KL, Morgan WJ, Goodwin JL, Cabrera R, et al. Sleep-
disordered breathing, pharyngeal size and soft tissue anatomy in children. Journal of Applied
Physiology. 2003;95:2030-8. doi: 10.1152/japplphysiol.00293.2003.
29.   Barbier G, Boë LJ, Captier G, Laboissière R. Human vocal tract growth: A longitudinal
study of the development of various anatomical structures. Proceedings of the Annual
Conference of the International Speech Communication Association, INTERSPEECH.
2015;2015-Janua:364-8.
30.   Goncalves RdC, Raveli DB, Pinto Ados S. Effects of age and gender on upper airway,
lower airway and upper lip growth. Braz Oral Res. 2011;25(3):241-7. Epub 2011/06/15. doi:
10.1590/s1806-83242011000300009. PubMed PMID: 21670855.
31.   Sheng CM, Lin LH, Su Y, Tsai HH. Developmental changes in pharyngeal airway depth
and hyoid bone position from childhood to young adulthood. Angle Orthodontist. 2009;79:484-
90. doi: 10.2319/062308-328.1.
32.   Mislik B, Hänggi MP, Signorelli L, Peltomäki TA, Patcas R. Pharyngeal airway
dimensions: A cephalometric, growth-study-based analysis of physiological variations in
children aged 6-17. European Journal of Orthodontics. 2014;36:331-9. doi: 10.1093/ejo/cjt068.
33.   Daraze A, Delatte M, Liistro G, Majzoub Z. Cephalometrics of Pharyngeal Airway Space
in Lebanese Adults. International Journal of Dentistry. 2017;2017. doi: 10.1155/2017/3959456.
34.   Rommel N, Bellon E, Hermans R, Smet M, De Meyer AM, Feenstra L, et al.
Development of the Orohypopharyngeal Cavity in Normal Infants and Young Children. Cleft
Palate-Craniofacial Journal. 2003;40:606-11. doi: 10.1597/1545-
1569(2003)040<0606:DOTOCI>2.0.CO;2.
35.   Chiang CC, Jeffres MN, Miller A, Hatcher DC. Three-dimensional airway evaluation in
387 subjects from one university orthodontic clinic using cone beam computed tomography.
Angle Orthod. 2012;82(6):985-92. Epub 2012/06/07. doi: 10.2319/122811-801.1. PubMed
PMID: 22668315.
36.   Jiang YY, Xu X, Su HL, Liu DX. Gender-related difference in the upper airway
dimensions and hyoid bone position in Chinese Han children and adolescents aged 6-18 years
using cone beam computed tomography. Acta Odontologica Scandinavica. 2014;73:391-400.
doi: 10.3109/00016357.2014.978366.
37.   Lenza MG, De MM, Dalstra M, Melsen B, Cattaneo PM. An analysis of different
approaches to the assessment of upper airway morphology: A CBCT study. Orthodontics and
Craniofacial Research. 2010;13(2):96-105. Epub 2010/05/19. doi: 10.1111/j.1601-
6343.2010.01482.x. PubMed PMID: 20477969.

38.   Schendel SA, Jacobson R, Khalessi S. Airway growth and development: A computerized
3-dimensional analysis. Journal of Oral and Maxillofacial Surgery. 2012;70:2174-83. doi:
10.1016/j.joms.2011.10.013.
39.   Anandarajah S, Dudhia R, Sandham A, Sonnesen L. Risk factors for small pharyngeal
airway dimensions in preorthodontic children: A three-dimensional study. Angle Orthod.
2017;87(1):138-46. Epub 2016/11/02. doi: 10.2319/012616-71.1. PubMed PMID: 27304232.
40.   Masoud AI, Alwadei FH, Alwadei AH, Lin EY, Viana MGC, Kusnoto B, et al.
Developing pediatric three-dimensional upper airway normative values using fixed and
interactive thresholds. Oral Radiology. 2020;36:89-99. doi: 10.1007/s11282-019-00384-3.
41.   Yanagita N, Terajima M, Kanomi R, Takahashi I. Three-dimensional analysis of
pharyngeal airway morphology in Japanese female adolescents. Orthodontic Waves.
2019;76(2):89-96. doi: 10.1016/j.odw.2017.01.004.
42.   Alves M, Franzotti ES, Baratieri C, Nunes LKF, Nojima LI, Ruellas ACO. Evaluation of
pharyngeal airway space amongst different skeletal patterns. International Journal of Oral and
Maxillofacial Surgery. 2012;41:814-9. doi: 10.1016/j.ijom.2012.01.015.
43.   Claudino LV, Mattos CT, Ruellas ACDO, Sant Anna EF. Pharyngeal airway
characterization in adolescents related to facial skeletal pattern: A preliminary study. American
Journal of Orthodontics and Dentofacial Orthopedics. 2013. doi: 10.1016/j.ajodo.2013.01.015.
PubMed PMID: 23726330.
44.   Li H, Lu X, Shi J, Shi H. Measurements of normal upper airway assessed by 3-
dimensional computed tomography in Chinese children and adolescents. International Journal of
Pediatric Otorhinolaryngology. 2011;75:1240-6. doi: 10.1016/j.ijporl.2011.06.022.
45.   Kim EJ, Choi JH, Kim YS, Kim TH, Lee SH, Lee HM, et al. Upper airway changes in
severe obstructive sleep apnea: Upper airway length and volumetric analyses using 3D MDCT.
Acta Oto-Laryngologica. 2011;131:527-32. doi: 10.3109/00016489.2010.535561.
46.   Gibelli D, Cellina M, Gibelli S, Oliva AG, Termine G, Sforza C. Three-Dimensional
Assessment of Pharyngeal Volume on Computed Tomography Scans: Applications to
Anesthesiology and Endoscopy. Journal of Craniofacial Surgery. 2020;00:1. doi:
10.1097/scs.0000000000006094.
47.   Inamoto Y, Saitoh E, Okada S, Kagaya H, Shibata S, Baba M, et al. Anatomy of the
larynx and pharynx: Effects of age, gender and height revealed by multidetector computed
tomography. Journal of Oral Rehabilitation. 2015;42:670-7. doi: 10.1111/joor.12298.
48.   Shigeta Y, Takumi O, Venturin J, Nguyen M, Clark GT, Enciso R. Gender and age-based
differences in computed-tomography measurements of the orophaynx. Oral Surg Oral Med Oral
Pathol Oral Radiol Endod. 2008;106(4).
49.   Welch KC, Foster GD, Ritter CT, Wadden TA, Arens R, Maislin G, et al. A Novel
Volumetric Magnetic Resonance Imaging Paradigm to Study Upper Airway Anatomy. Sleep.
2002;25:530-40. doi: 10.1093/sleep/25.5.530.
50.   Leboulanger N, Louis B, Fodil R, Boelle PY, Clement A, Garabedian EN, et al. Analysis
of the pharynx and the trachea by the acoustic reflection method in children: a pilot study. Respir
Physiol Neurobiol. 2011;175(2):228-33. Epub 2010/11/30. doi: 10.1016/j.resp.2010.11.008.
PubMed PMID: 21111847.
51.   Martin SE, Mathur R, Marshall I, Douglas NJ. The effect of age, sex, obesity and posture
on upper airway size. European Respiratory Journal. 1997;10:2087-90. doi:
10.1183/09031936.97.10092087.

52.   Brooks LJ, Strohl KP. Size and mechanical properties of the pharynx in healthy men and
women. American Review of Respiratory Disease. 1992;146:1394-7. doi:
10.1164/ajrccm/146.6.1394.
53.   Brown IG, Zamel N, Hoffstein V. Pharyngeal cross-sectional area in normal men and
women. Journal of Applied Physiology. 1986;61:890-5. doi: 10.1152/jappl.1986.61.3.890.
54.   Gurani SF, Di Carlo G, Cattaneo PM, Thorn JJ, Pinhol EM. Effect of Head and Tongue
Posture on the Pharyngeal Airway Dimensions and Morphology in Three-Dimensional Imaging:
a Systematic Review. Journal of Oral and Maxillofacial Research. 2016;7:1-12. doi:
10.5037/jomr.2016.7101.
55.   Di Carlo G, Gurani SF, Pinholt EM, Cattaneo PM. A new simple three-dimensional
method to characterize upper airway in orthognathic surgery patient. Dentomaxillofacial
Radiology. 2017;46. doi: 10.1259/dmfr.20170042.
56.   Ayappa I, Rapoport DM. The upper airway in sleep: Physiology of the pharynx. Sleep
Medicine Reviews. 2003;7:9-33. doi: 10.1053/smrv.2002.0238. PubMed PMID: 12586528.
57.   Adewale L. Anatomy and assessment of the pediatric airway. Paediatric Anaesthesia.
2009;19:1-8. doi: 10.1111/j.1460-9592.2009.03012.x.
58.   Netter FH. Head and Neck. Atlas of Human Anatomy. 7 ed. Philadelphia, PA: Elsevier
Inc.; 2019. p. 11-174.
59.   Arens R, Marcus CL. Pathophysiology of upper airway obstruction: A developmental
perspective. Sleep. 2004;27:997-1019. doi: 10.1093/sleep/27.5.997. PubMed PMID: 15453561.
60.   Logan BM, Reynolds PA, Rice S, Hutchings RT. Nose, oral cavity, pharynx, ear and
larynx. In: Logan BM, Reynolds PA, Rice S, editors. McMinn's Color Atlas of Head and Neck
Anatomy. 5 ed: Elsevier Ltd; 2017.
61.   Schuenke M, Schulte E, Schumacher U. Oral Cavity and Perioral Region. In: Baker EW,
editor. Head and Neck Anatomy for Dental Medicine. New York, NY: Thieme; 2010. p. 216.
62.   Laird AM, Yetkiner E, Kadioglu O, Currier GF. The Upper Airway. In: Kadioglu O,
Currier GF, editors. Craniofacial 3D Imaging: Current Concepts in Orthodontics and Oral and
Maxillofacial Surgery. Cham: Springer International Publishing; 2019. p. 97-112.
63.   Moore K, Dalley A, Agur AMR. Neck. Clinically Oriented Anatomy. 7 ed: Lippincott
Williams & Wilkins; 2006.
64.   Standring S. Gray's anatomy : the anatomical basis of clinical practice. Forty-first edition.
ed. New York: Elsevier Limited; 2016. p. xviii, 1562 pages.
65.   Gu M, McGrath CPJ, Hägg U, Wong RWK, Yang Y. Anatomy of the Upper Airway and
Its Growth in Childhood. Journal of Dentistry and Oral Biology. 2016;1.
66.   Kelly MP, Vorperian HK, Wang Y, Tillman KK, Werner HM, Chung MK, et al.
Characterizing mandibular growth using three-dimensional imaging techniques and anatomic
landmarks. Archives of Oral Biology. 2017;77:27-38. doi: 10.1016/j.archoralbio.2017.01.018.
67.   Miller CA, Hwang SJ, Cotter MM, Vorperian HK. Cervical vertebral body growth and
emergence of sexual dimorphism: a developmental study using computed tomography. Journal
of Anatomy. 2019;234:764-77. doi: 10.1111/joa.12976.
68.   Miller CA, Lee Y, Avey GD, Vorperian HK. Head position classification of medical
imaging studies: an assessment and development of a protocol. Dentomaxillofacial Radiology.
2019:20190220. doi: 10.1259/dmfr.20190220.
69.   AnalyzeDirect. Analyze 12.0. Overland Park, KS: AnalyzeDirect; 2018.

70.   Nakano H, Mishima K, Ueda Y, Matsushita A, Suga H, Miyawaki Y, et al. A new
method for determining the optimal CT threshold for extracting the upper airway.
Dentomaxillofacial Radiology. 2013;42. doi: 10.1259/dmfr/26397438.
71.   Vorperian HK, Wang S, Michael Schimek E, Durtschi RB, Kent RD, Gentry LR, et al.
Developmental sexual dimorphism of the oral and pharyngeal portions of the vocal tract: An
imaging study. Journal of Speech, Language, and Hearing Research. 2011;54:995-1010. doi:
10.1044/1092-4388(2010/10-0097).
72.   Lorensen WE, Cline HE. Marching cubes: A high resolution 3D surface construction
algorithm. Proceedings of the 14th annual conference on Computer graphics and interactive
techniques. 37422: ACM; 1987. p. 163-9.
73.   Desbrun M, Meyer M, Schröder P, Barr AH, editors. Implicit fairing of irregular meshes
using diffusion and curvature flow. Proceedings of the 26th annual conference on Computer
graphics and interactive techniques - SIGGRAPH '99; 1999.
74.   Lazarus F, Verroust A. Level set diagrams of polyhedral objects. Proceedings of the
Symposium on Solid Modeling and Applications. 1999:130-40. doi: 10.1145/304012.304025.
75.   Seo S, Chung MK, Whyms BJ, Vorperian HK, editors. Mandible shape modeling using
the second eigenfunction of the Laplace-Beltrami operator. Medical Imaging 2011: Image
Processing; 2011.
76.   Shi Y, Lai R, Krishna S, Sicotte N, Dinov I, Toga AW. Anisotropic Laplace-Beltrami
Eigenmaps: Bridging Reeb Graphs and Skeletons. Proc IEEE Comput Soc Conf Comput Vis
Pattern Recognit. 2008;2008:1-7. Epub 2008/07/15. doi: 10.1109/CVPRW.2008.4563018.
PubMed PMID: 21339850; PubMed Central PMCID: PMCPMC3041984.
77.   de Boor C. A Practical Guide to Splines. New York, NY: Springer New York; 1978.
78.   Hunyandi L. B-splines. MATLAB Central File Exchange; 2010.
79.   Hudgins PA, Siegel J, Jacobs I, Abramowsky CR. The normal pediatric larynx on CT and
MR. AJNR Am J Neuroradiol. 1997;18(2):239-45. Epub 1997/02/01. PubMed PMID: 9111658.
80.   Scammon RE. The measurement of the body in childhood. In: Harris JA, Jackson CM,
Paterson DG, Scammon RE, editors. The measurement of man. Minneapolis, MN: The
University of Minnesota Press; 1930. p. 173-215.
81.   Karia H, Shrivastav S, Karia AK. Three-dimensional evaluation of the airway spaces in
patients with and without cleft lip and palate: A digital volume tomographic study. Am J Orthod
Dentofacial Orthop. 2017;152(3):371-81. Epub 2017/09/03. doi: 10.1016/j.ajodo.2016.12.026.
PubMed PMID: 28863918.
82.   Cassano P, Gelardi M, Cassano M, Fiorella ML, Fiorella R. Adenoid tissue
rhinopharyngeal obstruction grading based on fiberendoscopic findings: a novel approach to
therapeutic management. Int J Pediatr Otorhinolaryngol. 2003;67(12):1303-9. Epub 2003/12/04.
doi: 10.1016/j.ijporl.2003.07.018. PubMed PMID: 14643473.
83.   Jeans WD, Fernando DCJ, Maw AR, Leighton BC. A longitudinal study of the growth of
the nasopharynx and its contents in normal children. The British Journal of Radiology.
1981;54(638):117-21. doi: 10.1259/0007-1285-54-638-117. PubMed PMID: 7459548.
84.   de Souza Vilella B, de Vasconcelos Vilella O, Koch HA. Growth of the nasopharynx and
adenoidal development in Brazilian subjects. Brazilian Oral Research. 2006;20:70-5. doi:
10.1590/s1806-83242006000100013.
85.   Linder-Aronson S, Stockholm BCL, London S, editors. A longitudinal study of the
development of the posterior nasopharyngeal wall between 3 and 16 years of age. European
Journal of Orthodontics; 1983.

86.   Luscan R, Leboulanger N, Fayoux P, Kerner G, Belhous K, Couloigner V, et al.
Developmental changes of upper airway dimensions in children. Pediatric Anesthesia.
2020:pan.13832. doi: 10.1111/pan.13832. PubMed PMID: 31995659.
87.   Liu S, Qi W, Zhang X, Dong Y. The development of the cricoid cartilage and its
implications for the use of endotracheal tubes in the pediatric population. Pediatric Anesthesia.
2020;30(1):63-8. doi: https://doi.org/10.1111/pan.13772.
88.   Perry JL, Kollara L, Kuehn DP, Sutton BP, Fang X. Examining age, sex, and race
characteristics of velopharyngeal structures in 4- to 9-year-old children using magnetic
resonance imaging. Cleft Palate-Craniofacial Journal. 2018;55:21-34. doi:
10.1177/1055665617718549.
89.   Dang J, Honda K. Acoustic Characteristics of the Piriform Fossa in Models and Humans.
Journal of the Acoustical Society of America. 1997;101(1):456-65. doi: 10.1121/1.417990.
90.   Story BH. Physiologically-based speech simulation using an enhanced wave-reflection
model of the vocal tract [Ph.D.]. Ann Arbor: The University of Iowa; 1995.
91.   Story BH, Titze IR, Hoffman EA. Vocal tract area functions for an adult female speaker
based on volumetric imaging. The Journal of the Acoustical Society of America.
1998;104(1):471-87.
92.   Fant G. Acoustic Theory of Speech Production. Acoustic Theory of Speech Production.
1971. doi: 10.1515/9783110873429.
93.   Baer T, Gore JC, Gracco LC, Nye PW. Analysis of Vocal Tract Shape and Dimensions
using Magnetic Resonance Imaging: Vowels. Journal of the Acoustical Society of America.
1991;90(2 Pt 1):799-828.
94.   Fujita S, Honda K. An experimental study of acoustic characteristics of hypopharyngeal
cavities using vocal tract solid models. Acoustical Science and Technology. 2005;26(4):353-7.
doi: 10.1250/ast.26.353.
95.   Story BH, Bunton K. A model of speech production based on the acoustic relativity of
the vocal tract. J Acoust Soc Am. 2019;146(4):2522. Epub 2019/11/02. doi: 10.1121/1.5127756.
PubMed PMID: 31671993.
96.   Buhr RD. The Emergence of Vowels in an Infant. Journal of Speech, Language, and
Hearing Research. 1980;23(1):73-94.
97.   Watters KF. Tracheostomy in Infants and Children. Respiratory Care. 2017;62:799-825.
98.   Thach BT. The Role of the Upper Airway in SIDS and Sudden Unexpected Infant Deaths
and the Importance of External Airway-Protective Behaviors. In: Duncan JR, Byard RW, editors.
SIDS Sudden Infant and Early Childhood Death: The Past, the Present and the Future. Adelaide
(AU)2018.
99.   Cornwell AC. Sex differences in the maturation of sleep/wake patterns in high risk for
SIDS infants. Neuropediatrics. 1993;24(1):8-14. Epub 1993/02/01. doi: 10.1055/s-2008-
1071505. PubMed PMID: 8474614.
100.   Khadivi E, Zaringhalam MA, Khazaeni K, Bakhshaee M. Distance between anterior
commissure and the first tracheal ring: An important new clinical laryngotracheal measurement.
Iranian Journal of Otorhinolaryngology. 2015;27(80):193-7. doi: 10.22038/ijorl.2015.4252.
101.   Sirisopana M, Saint-Martin C, Wang NN, Manoukian J, Nguyen LHP, Brown KA. Novel
measurements of the length of the subglottic airway in infants and young children. Anesthesia
and Analgesia. 2013;117(2):462-70. doi: 10.1213/ANE.0b013e3182991d42.
102.   Griscom NT, Wohl MEB. Dimensions of the growing trachea related to age and gender.
American Journal of Roentgenology. 1986;146(2):233-7. doi: 10.2214/ajr.146.2.233.

103.   Perry JL, Kuehn DP, Sutton BP, Gamage JK, Fang X. Anthropometric analysis of the velopharynx and related craniometric dimensions in three adult populations using MRI. Cleft Palate-Craniofacial Journal. 2016;53:e1-e13. doi: 10.1597/14-015.
104.   Bloom K, Moore-Schoenmakers K, Masataka N. Nasality of infant vocalizations determines gender bias in adult favorability ratings. Journal of Nonverbal Behavior. 1999;23(3):219-36. doi: Doi 10.1023/A:1021317310745. PubMed PMID: WOS:000083422500002.
105.   Yang S, Mu L. An investigation of the third formant of /a/ in prepubertal children. Journal of Voice. 1989;3(4):321-3. doi: https://doi.org/10.1016/S0892-1997(89)80054-2.
106.   Perry TL, Ohde RN, Ashmead DH. The acoustic bases for gender identification from children's voices. The Journal of the Acoustical Society of America. 2001;109(6):2988-98.
107.   Vorperian HK, Kent RD, Lee Y, Bolt DM. Corner vowels in males and females ages 4 to 20 years: Fundamental and F1-F4 formant frequencies. J Acoust Soc Am. 2019;146(5):3255. Epub 2019/12/05. doi: 10.1121/1.5131271. PubMed PMID: 31795713; PubMed Central PMCID: PMCPMC6850954.
108.   Buschang PH, Hinton RJ. A Gradient of Potential for Modifying Craniofacial Growth. Seminars in Orthodontics. 2005;11(4):219-26. doi: 10.1053/j.sodo.2005.07.006.
109.   Wang Y, Chung MK, Vorperian HK. Composite growth model applied to human oral and pharyngeal structures and identifying the contribution of growth types. Statistical Methods in Medical Research. 2016;25:1975-90. doi: 10.1177/0962280213508849.
110.   Durtschi RB, Chung D, Gentry LR, Chung MK, Vorperian HK. Developmental craniofacial anthropometry: Assessment of race effects. Clinical Anatomy. 2009;22(7):800-8. Epub 2009/09/16. doi: 10.1002/ca.20852. PubMed PMID: PMID: 19753647

NIHMS188027

PMCID: PMC2849302.

Supporting Information

S1 Data. Data used in this study. Variable measurements as described in Table 5.

Figure 1

M245-04-08-002

Figure 2

Figure 3A

Figure 3B

Figure 3C

Figure 3D

Figure 3E