Sign inversion in selection on ploidy

Yevgeniy Raynes and Daniel M. Weinreich

Department of Ecology, Evolution, and Organismal Biology and
Center for Computational Molecular Biology
Brown University Box G-W, Providence, RI 02912 USA

email: yevgeniy_raynes@brown.edu

Abstract

Ploidy – the number of homologous chromosome sets in a cell – is remarkably variable across the natural world, yet the evolutionary processes that have resulted in such diversity remain poorly understood. Here we use stochastic agent-based simulations to model ploidy evolution under the influence of indirect selection, i.e., selection mediated solely by statistical associations with fitness-affecting mutations. We find that in non-equilibrium asexual populations, the sign of selection on ploidy can change with population size – a phenomenon we have previously termed sign inversion. In large populations, ploidy dynamics are dominated by indirect effects of selection on beneficial mutations, which favors haploids over diploids. However, as population size declines, selection for beneficial mutations is neutralized by random genetic drift before drift can overwhelm selection against the cost of the deleterious mutational load. As a result, in small populations indirect selection is dominated by the cost of the deleterious load, which favors diploids over haploids. Our work adds to the growing body of evidence challenging established evolutionary theory that population size can affect only the efficiency, but not the sign, of natural selection.

Keywords: ploidy evolution, indirect selection, sign inversion

Introduction

Why do some species have a single set of homologous chromosomes while others have two or more? The evolutionary forces that govern transitions between haploidy, diploidy, and polyploidy and have produced the variability we see today have long been a subject of significant theoretical and empirical interest (reviewed in: 1, 2). Theories for the role that natural selection plays in ploidy evolution consider either the immediate effects of ploidy on the phenotype or longer-term consequences for the supply rate of fitness-affecting mutations and the efficacy of selection (1). When the phenotypic effects of ploidy alter the fitness of an individual, ploidy itself may be directly influenced by selection. For example, haploid yeast cells are generally smaller and have a higher surface area to volume ratio than diploid yeast cells, which may afford them higher fitness in nutrient limited conditions (2, 3, but see 4). Ploidy variants may also experience indirect selection acting through statistical associations (i.e., linkage disequilibrium) that develop with fitness-affecting mutations. Indirect selection is particularly effective in asexual populations as recombination can quickly erode linkage disequilibrium.

Most theoretical studies of indirect selection have made progress by considering the effects of selection on deleterious mutations separately from the effects of selection on beneficial mutations. Selection against deleterious mutations can favor diploids over haploids in non-equilibrium populations as diploids can mask the effects of recessive or partially dominant deleterious mutations: i.e., mutations that are fully expressed in haploids but only partially, or not at all, in diploids, where they first appear in heterozygous state (5). In other words, while in haploids such mutations lower fitness by a selection coefficient s_d, in heterozygous diploids their effect on fitness is reduced to $h_d s_d$; h_d, known as the dominance coefficient, measures the dominance of the deleterious mutation and is below 1 for any but the fully dominant mutations. However, this advantage in the fitness cost of deleterious mutations may only be a transient one. Once at mutation-selection equilibrium, diploids – with twice as many loci as haploids – end up bearing twice the cost of the deleterious mutational load (6, 7). That is, while the equilibrium mean fitness of a haploid populations is $1-U_d$ (where U_d is the deleterious mutation rate) diploids have an equilibrium mean fitness of $1-2U_d$ (6). Nevertheless, two earlier studies have shown that masking of deleterious mutations could favor diploids at equilibrium in certain limited circumstances. Kondrashov and Crow (8) found that diploid equilibrium mean fitness could exceed that of haploids under truncation selection and a relatively low h_d (below 0.25). Meanwhile, Perrot et al (9) demonstrated that diploids could invade haploids in populations with random mating between haploids and diploids, high rate of free recombination, and h_d below 0.5; later work confirmed that diploids fail to supplant haploids in populations with rare recombination (10).

Diploids may also be favored because of their increased supply of beneficial
mutations, given their 2-fold advantage in beneficial mutation rate. Specifically, if
recombination is sufficiently rare diploids may preferentially increase in frequency by
hitchhiking (11) along with those beneficial mutations that survive drift and sweep to
fixation. However, beneficial mutations with dominance coefficient, h_b, below 1 would
also be at least partially masked in diploids, reducing their fitness advantage to $h_b s_b$ and
lowering their fixation probability compared to those in haploids. This advantage in the
efficacy of selection could, in certain circumstances, favor haploids. Orr and Otto (12)
compared the rates of adaptive evolution in diploids and haploids. They found that
diploids adapt faster than haploids when the rate of adaptation is limited by the supply
of beneficial mutations – e.g., when beneficial mutations are rare or population size is
small – provided new mutations are at least partially dominant ($h_b > 0.5$). On the other
hand, as long as $h_b < 1$, diploids evolve slower than haploids when the rate of
adaptation is limited not by the supply of beneficial mutations – e.g., in large populations
in which such mutations are common – but by the strength of selection for beneficial
mutations.

In contrast to most earlier treatments, here we model the evolution of ploidy in
non-equilibrium populations with both beneficial and deleterious mutations. Using
individual-based simulations we show that, in the absence of recombination, whether
ploidy variants are favored or disfavored by indirect selection depends on population
size. In small asexual populations, selection against the deleterious mutational load
favors diploids and disfavors haploids. In large asexual populations, the cost of the
deleterious load is overwhelmed by selection for beneficial mutations, which favors
haploids over diploids.

Methods

We model asexual, clonally reproducing populations of constant size, N evolving
in discrete, non-overlapping generations. Populations are composed of genetic lineages
defined by ploidy state and the number of beneficial and deleterious mutations. Each
new beneficial mutation increases an individual's fitness by a constant s_b; a new
deleterious mutation reduces fitness by a constant s_d. We assume that fitness effects
are additive and calculate the fitness of a haploid lineage with x beneficial and y
deleterious mutations as $w_{xy} = 1+xs_b-ys_d$. As in (12) we also assume that in the absence
of recombination fitness-affecting mutations in diploids remain permanently
heterozygous (at least on the time scale of our simulations) and calculate the fitness of
a diploid lineage with x beneficial and y deleterious mutations as $w_{xy}=1+xh_b s_b-yh_d s_d$,
where h_b and h_d are the corresponding dominance coefficients.

At the outset of simulation, a single individual of a particular ploidy (either haploid
or diploid) is introduced to a population of N-1 individuals of the other ploidy. Both the

invading lineage of size 1 and the resident lineage of size N-1 bear no fitness-affecting
mutations and, thus, have an initial fitness $w = 1$. Every generation the population
reproduces according to the Wright-Fisher model (13) with the size of the lineage
carrying x beneficial mutations and y deleterious mutation in generation t+1 randomly
sampled from a multinomial distribution with expectation $Nf_t \cdot w_{xy}/\overline{w}$ where N is the size of
the population, f_t is the frequency of the lineage in generation t, w_{xy} is the fitness of the
lineage as defined above and $\overline{w}$ is the average fitness of the population. Upon
reproduction, each surviving lineage acquires a random number of beneficial and
deleterious mutations drawn from a Poisson distribution with expectation given by,
respectively, pN_iU_b and pN_iU_d. Here, p is the ploidy of the lineage, N_i is the size of the
lineage, U_b is the per-individual beneficial mutation rate, and U_d is the per-individual
deleterious mutation rate. Diploids, thus, have a 2-fold higher mutation rate than
haploids. For simulations in Fig. 2C we lowered the diploid mutation rate to that of
haploids by setting $p = 1$ for all diploid and haploid lineages.

Simulations end when the invader either fixes (reaches a frequency of 1.0) or is
eliminated from the population. We assess the fixation probably (P_{fix}) as the frequency
with which the invader supplants the resident across the replicate runs of the simulation.
Throughout, we focus on NP_{fix} - the fixation probability normalized by the neutral
expectation, 1/N. To ascertain whether an invading ploidy variant is favored or
disfavored by selection we compare its NP_{fix} to that expected of a neutral mutation, i.e.,
$NP_{fix} = 1$ (e.g., 14).

Simulation code was written in Julia 1.2 and is available upon request.

Results

We first examined the fixation probability of haploid invaders in diploid
populations. Since deleterious mutations in nature are predominantly recessive (e.g.,
15, 16, 17), we set the dominance coefficient of deleterious mutations in our simulations
to be $h_d = 0.0$. Meanwhile, we varied the dominance coefficient of new beneficial
mutations from fully recessive ($h_b = 0.0$) to fully dominant ($h_b = 1.0$). Strikingly, we found
that the sign of selection on haploids can change with population size (Fig 1A). For all
values of $h_b < 1.0$ studied, haploids are favored by selection (have $NP_{fix} > 1$) in large
populations but become disfavored (have $NP_{fix} < 1$) as population size decreases below
a critical size we call N_{crit}. We have previously designated this N-dependent switch in
the sign of selection *sign inversion* (14, 18). Fig. S1 confirms that haploids are
susceptible to sign inversion across a wide – at least over several orders of magnitude –
range of mutation rates.

Figure 1: Sign inversion in selection on haploids. A) Normalized fixation probability (NP_{fix}) of
haploid invaders (circles) in diploid populations at different h_b. Red dashed line: $P_{fix}(s = -U_d, N)$;
Teal dashed line: NP_{fix}^{hitch}. **B)** Average fitness of winning haploid populations, w. Horizontal
black lines indicate $w = 1$ and $w = 1 + s_b = 1.1$ **C)** Conditional fixation probability calculated
across simulations in which haploid invaders managed to acquire at least one beneficial
mutation. Horizontal line shows expected $P_{fix}(s_b, N)$. Parameter values: $s_b = 0.1$, $s_d = -0.1$, $U_d =$
0.01, $U_b = 0.001$. All simulation results averaged across at least 10^7 replicates.

Based on our earlier work on sign inversion (14, 18) we assumed that in small
populations haploid NP_{fix} would be reduced below the neutral expectation by selection
against the deleterious load. To test this expectation, we compared haploid NP_{fix} in
simulations to that predicted by the fitness cost of the haploid deleterious load.
Specifically, Kimura (19) showed that an allele that increases the deleterious mutation
rate by ΔU_d bears a fitness cost at equilibrium of $s = -\Delta U_d$. Given that deleterious
mutations in our simulations are permanently heterozygous and fully recessive we can
assume that the deleterious mutation rate in diploid residents is essentially zero.
Therefore, ΔU_d is simply the deleterious mutation rate of the haploid and the expected
cost of the load in haploids is given by $s = -U_d$. The probability of fixation for a mutation
of fitness effect s (in a haploid population) is given classically by $P_{fix}(s, N) = (1-e^{-2s})/(1-e^{-2Ns})$ (20). $P_{fix}(s = -U_d, N)$ is shown with the red dashed line in Fig. 1A and agrees very
well with simulations, confirming that in sufficiently small populations haploid NP_{fix} is set
by the cost of the deleterious load.

Again motivated by our earlier work, we hypothesized that in large populations
that favor haploids, haploid NP_{fix} is elevated above the neutral expectation by selection
for beneficial mutations. To confirm the influence of selection for beneficial mutations on
haploid NP_{fix} we assayed the final fitness of populations in which haploids had fixed
(Fig. 1B). We found that at smallest N, the average fitness of winning haploid
populations is, in fact, lower than the ancestral fitness, confirming that at these
population sizes, hitchhiking with beneficial mutations (which would necessitate an
average fitness of at least $1+s_b$) is not a significant influence on NP_{fix}. Rather, haploids
fix in these populations via random genetic drift. However, as Fig. 1B shows, in

sufficiently large populations haploid winners' fitness surpasses $1+s_b$ for all values of h_b
examined. This demonstrates that in large populations haploids fix almost exclusively by
hitchhiking. In fact, note that at yet larger N haploid fitness may surpass $1+2s_b$
suggesting that in those populations haploids reach fixation with multiple beneficial
mutations. For now, we leave further examination of multiple-mutation (21) dynamics in
ploidy evolution for future work.

Moreover, we found that at h_b = 0 haploid NP_{fix} could be well approximated by
the expected probability of hitchhiking to fixation with a beneficial mutation, previously
derived for a mutator allele in (14). In brief, to fix by hitchhiking an invader must first
produce a beneficial mutation, which then has to sweep to fixation. The former occurs
with probability approximately equal to the ratio of U_b to the total mutation rate, $U_b + U_d$.
Since at h_b = 0 the spread of beneficial mutations in the haploid background is
unaffected by selection on beneficial mutations in diploids, the probability of the latter for
a mutation of fitness effect s_b can be estimated with $P_{fix}(s_b, N)$ given above. The
probability of hitchhiking to fixation is the product of these two probabilities: P_{fix}^{hitch} =
$U_b/(U_b + U_d) \times P_{fix}(s_b, N)$. NP_{fix}^{hitch} is shown with a teal line in Fig. 1A.

Finally, Fig. 1A also highlights the negative relationship between haploid NP_{fix}
and h_b. As may be expected, h_b has little effect on NP_{fix} in small populations, in which
NP_{fix} is set by the deleterious load while P_{fix}^{hitch}– whether for haploids or diploids – is
miniscule. However, above N_{crit} haploids fare worse at higher h_b suggesting that
increasing dominance of beneficial mutations lowers P_{fix}^{hitch} (which dominates NP_{fix} in at
larger N). Focusing on the two components of P_{fix}^{hitch} above, we can assume that the
probability of a haploid acquiring a beneficial mutation is unaffected by h_b. However, the
likelihood of the beneficial mutation escaping drift and reaching fixation may be reduced
at higher h_b by stronger competition from beneficial mutations in the diploids. In other
words, whereas at h_b = 0, haploids with a beneficial mutation have an advantage s_b over
any diploid similarly endowed with a beneficial mutation, that advantage decreases to
$s_b-h_b s_b$ for any h_b > 0.0. (At h_b = 1.0 haploids enjoy no advantage in the strength of
selection on associated beneficial mutations; instead, diploids that benefit from a two-
fold higher U_b and a lower deleterious load are favored by selection at all N).

To test how interference from beneficial mutations in diploids reduces haploid
P_{fix}^{hitch} at h_b > 0, we assayed the probability of haploid fixation conditioned on acquiring
at least one beneficial mutation, $P_{fix|ben}$ (Fig. 1C). We found that at h_b = 0, $P_{fix|ben}$
asymptotes to the expected probability of fixation given by $P_{fix}(s_b, N)$, which shows that
beneficial mutations in haploids are unaffected by competition from fully recessive
mutations in diploids. However, at h_b > 0, $P_{fix|ben}$ declines below the expected $P_{fix}(s_b, N)$,
confirming that competition from beneficial mutations in diploid residents reduces the
probability of fixation of new beneficial mutations in haploids. Note also that $P_{fix|ben}$

declines with increasing N and does so faster at higher h_b, consistent with intensifying
interference from diploid residents, modulated by both more effective selection and the
rising supply of beneficial mutations (i.e., NU_b) in diploids.

Next, we studied the fixation probability of diploid invaders in haploid populations.
Once again, we set deleterious mutations to be fully recessive ($h_d = 0.0$), while
beneficial mutations ranged from fully recessive ($h_b = 0.0$) to fully dominant ($h_b = 1.0$).
Fig. 2A confirms that diploids are also susceptible to sign inversion. Unlike that of
haploids, though, diploid NP_{fix} has a negative slope at N_{crit}. For all studied $h_b < 1$,
diploids are favored in small populations and become disfavored as N increases. At $h_b =$
1.0 diploids are everywhere favored over haploids (have $NP_{fix} > 1$ at all N).

Our analysis of haploid NP_{fix} above suggests that in sufficiently small
populations, diploids would mostly reach fixation not by acquiring and fixing beneficial
mutations but by simply outlasting haploid residents that succumb to selection against
the load. Fig. 2A confirms that diploid NP_{fix} in small populations is well predicted simply
by the fixation probability for $s = +\Delta U_d$ – the diploid's expected fitness advantage due to
the deleterious load borne by haploids. Indeed, assaying the average fitness at the end
of simulation confirms that in sufficiently small populations diploid winners are
indistinguishable from their ancestors (Fig. 2B).

In larger populations though, in which haploid dynamics are dominated by
beneficial mutations, we expected diploid fate to be likewise determined by the
likelihood of hitchhiking with beneficial mutations of their own. Correspondingly, we
found that the fitness of diploid winners increases with N and in sufficiently large
populations rises above $1+h_b s_b$ (Fig. 2B). This suggests that in those populations
diploids also fix exclusively by hitchhiking with beneficial mutations. (Except at $h_b = 0$,
where diploid winners never exceed ancestral fitness as hitchhiking with completely
recessive beneficial mutations is impossible at any N.) For a diploid to fix by hitchhiking,
though, a beneficial mutation of effect $h_b s_b$ must sweep to fixation without interference
from beneficial mutations of larger effect s_b in haploids. Diploids possess an inherent
advantage of a two-fold higher mutation rate so a diploid invader is always more likely to
produce a beneficial mutation than any of its haploid competitors. This advantage is
expected to be most pronounced in relatively small populations in which the rate of
adaptation is limited by the supply rate of beneficial mutations (12). Our simulations
confirm that the higher U_b improves diploid probability of hitchhiking. Specifically, as
shown in Fig. 2C diploid NP_{fix} is reduced when diploid mutation rate is lowered to that of
haploids (see Methods). More importantly, Fig. 2C shows that making diploid U_b equal
to that of haploids lowers N_{crit} for sign inversion for all $0 < h_b < 1.0$, suggesting that at
some intermediate N, diploid NP_{fix} is above 1 not only due to purifying selection against
the load but also positive selection mediated by a higher U_b (note that in simulations

without deleterious mutations, diploids are only ever favored by selection at $h_b > 0.5$
(Fig. S2), consistent with (12)).

However, as long as $h_b < 1$, haploids enjoy an advantage in the strength of
selection on new beneficial mutations (i.e., beneficial mutations in haploids endow
carriers with a smaller fitness advantage $h_b s_b$ relative to s_b in diploids). This advantage
is magnified as population size increases and selection for beneficial mutations
becomes more efficient (while the supply rate of such mutations to haploids increases
as well). In sufficiently large populations selection for beneficial mutations in haploids
offsets the cost of the load and the diploids' advantage in U_b. When it does, diploid NP_{fix}
(set at that point almost exclusively by diploid NP_{fix}^{hitch}, see above) declines below 1 at
all $h_b < 1.0$ (Fig. 2A). Correspondingly, diploids fare better at higher h_b, consistent with a
reduced fitness advantage ($s_b - h_b s_b$) of beneficial mutations in haploids. And, diploid
NP_{fix} is never below 1 when beneficial mutations are fully dominant: at $h_b = 1$, haploids
have no advantage in the efficacy of selection, whereas diploids still enjoy the benefit of
a two-fold higher U_b.

Figure 2: Sign inversion in selection on diploids. A) Normalized fixation probability (NP_{fix}) of
diploid invaders (circles) in diploid populations at different h_b. Teal dashed line: $P_{fix}(s = U_d, N)$.
B) Average fitness of winning diploid populations, w. Horizontal lines indicate $w = 1$ in black
and $w = 1 + h_b s_b$ in corresponding colors. **C)** Comparison between NP_{fix} in panel A (circles
again) and diploid NP_{fix} in simulations in which diploid U was set to equal that of the haploids
(shown for convenience as dashed lines, results only available at the same N as circles).
Parameter values: $s_b=0.1$, $s_d=-0.1$, $U_d= 0.01$, $U_b = 0.001$. All simulation results averaged across
10^7 replicates.

Discussion

We simulated evolution of ploidy in population with recessive deleterious
mutations and beneficial mutations of varying dominance. We showed that in
populations with both kinds of mutations, the sign of selection on ploidy can change with
population size – a phenomenon we have previously designated sign inversion (14).
Consistent with previous expectations (5) for non-equilibrium populations, we found that

selection against recessive deleterious mutations lowers haploid fixation probability.
Indeed, in sufficiently small populations the evolutionary fate of ploidy variants is entirely
dominated by selection against the cost of the deleterious load. Even in the presence of
beneficial mutations, haploid invaders that bear the full cost of the deleterious mutations
fare worse than the neutral expectation (Fig. 1A). Meanwhile, diploid invaders that mask
deleterious mutations and, thus, can avoid the fitness cost of the load fare better than
the neutral expectation (Fig. 2A).

On the other hand, in large populations, ploidy evolution is dominated by the
indirect effects of selection on beneficial mutations. Previously, it's been shown that
selection for beneficial mutations may favor diploids – that produce twice as many
mutations as haploids – when the rate of adaptation is limited by the mutation rate (i.e.,
the product NU is small). We find that the benefit of a 2-fold higher U_b (in tandem with
the cost of haploids' load) may indeed raise diploid NP_{fix} above the neutral expectation
at some N that would otherwise favor haploids (Fig. 2C). Conversely, haploids have
been predicted to fare better than diploids if new beneficial mutations are sufficiently
common that the rate of adaptation is limited not by their supply but by their rate of
spread (12). The reason for this is that mutations in haploids have larger effects on
fitness (s_b versus $h_b s_b$) and are, thus, more likely to survive drift and reach fixation than
mutations in diploids. Our work confirms that in sufficiently large populations the
probability of hitchhiking to fixation exceeds the neutral expectation for haploids but is
below the neutral expectation for diploids. Correspondingly, in large populations
haploids fare better than neutral mutations, while diploid NP_{fix} drops below 1. The
expression for NP_{fix}^{hitch} plotted in Fig. 1A for h_b=0 illustrates why haploid NP_{fix} transitions
from below 1 to above 1 with N. Recall that NP_{fix}^{hitch} is approximately the product of two
probabilities: that of acquiring a beneficial mutation and that of the beneficial mutation
reaching fixation. The probability of generating a beneficial mutation (assuming that one
is produced) is small and well below one, given that $U_d \gg U_b$. $NP_{fix}(s_b, N)$ of a beneficial
mutation is always above the neutral expectation and is a monotonically increasing
function of N. And while it may be reduced at $h_b > 0$ by competition from beneficial
mutations in diploids, mutations in haploids should still be favored over those in diploids
by a selection coefficient of $s_b - h_b s_b$ as long as $h_b < 1.0$. Thus, when N is small, the
product of the two, NP_{fix}^{hitch}, is reduced below the neutral expectation by the low
probability of generating a beneficial mutation. However, as population size increases
above N_{crit} and selection on new beneficial mutations becomes more efficient ($NP_{fix}(s_b,$
$N)$ of new beneficial mutations rises further away from 1), NP_{fix}^{hitch} must necessarily
increase above the neutral expectation at some sufficiently large N_{crit}, lifting the total
NP_{fix} above 1 as well.

Taken together, our results demonstrate that sign inversion occurs because
ploidy variants can experience both the positive selection mediated by linkage to

beneficial mutations and the purifying selection mediated by linkage to deleterious
mutations that are overwhelmed by drift at different N. That is, haploid NP_{fix} is
dominated by selection on beneficial mutation at large N (represented by NP_{fix}^{hitch}) but as
population size decreases it is neutralized by drift (NP_{fix}^{hitch} crosses 1) before selection
against the cost of the load, which leads to sign inversion.

Previously, we have found evidence of sign inversion in four published studies of
the evolution of recombination (22), cooperation (23), variance in offspring number (24),
and the mutation rate (14). Having examined all four cases of sign inversion (18) we
discovered that all alleles susceptible to sign inversion can impart both fitness costs and
fitness benefits on their carriers. That is, similar to ploidy variants, known cases of sign
inversion exhibit variability in the sign of selection on individual carriers. We designated
such alleles as having sign-variable effects and showed that that all sign-variable effects
could be classified into either those that vary between different mutant lineages or
among individual members of a single lineage (modulated, for example, by temporal
changes in the environment or population composition) (18). Ploidy in our model
exhibits between-lineage variability. Haploid lineages that generate beneficial mutations
enjoy the indirect fitness benefit and a higher than neutral probability of fixation.
Meanwhile, lineages that generate deleterious mutations bear the indirect fitness cost
and, correspondingly, have a lower than neutral probability of fixation. More importantly,
in the same earlier study we showed that sign variability is necessary for sign inversion,
which always occurs because selection on cost-bearing and benefit-endowed carriers
exhibit differing sensitivities to drift (18). Our current work showing that sign inversion in
selection on ploidy is also driven by between-lineage sign variability corroborates the
necessity of sign variability for sign inversion and provides additional support for the
common mechanistic basis of sign inversion.

Finally, our findings may be particularly relevant for the growing number of
evolution experiments that have documented ploidy transitions in laboratory microbial
populations – those expected to be relatively far from equilibrium with access to both
beneficial and deleterious mutations. In particular, studies in *Saccharomyces cerevisiae*
have repeatedly observed shifts from haploidy to diploidy through spontaneous genome
duplication (25-29). Interestingly, most of these studies have been conducted in large
populations that, we have found, should indirectly favor haploids rather than diploids.
Accordingly, several studies have documented an inherent fitness advantage to diploidy
(25-27) that would allow diploids to fix through the action of direct selection instead.
Others, however, did not detect a significant direct advantage to diploid cells over
haploids (30, 31), suggesting that diploids might have arisen via indirect selection. Yet,
whereas the appearance of diploids in small populations of the mutation accumulation
experiment of Dickinson (30) appears consistent with our results, diploids repeatedly
overtaking large populations in the study of Gerstein et al. (32) is difficult to reconcile

with our simulations. It is possible that ploidy evolution, at least in yeast, is universally
driven by direct selection and an inherent fitness advantage will eventually be identified
in all laboratory conditions in which diploids have been shown to overtake haploids (see
also (2) for the discussion of "ploidy drive"). It is also possible, though, that experimental
populations differ in some properties that affect indirect selection on ploidy in such a
way as to indirectly favor diploids in some environments and disfavor them in others,
even at relatively large N (see also (33)). For example, our work suggests that diploids
could fare better in larger populations given a higher h_b or U_d (which would reduce
haploid P_{fix}^{hitch}), both of which remain somewhat difficult to assay experimentally. All in
all, while our work adds to our understanding of the mechanics of indirect selection on
ploidy variants, whether indirect selection can be the driving force in ploidy evolution in
laboratory or in nature remains to be determined.

Acknowledgements

Simulations were performed on the computing cluster of the Computer Science
Department at Brown University. The work was supported by National Science
Foundation Grants DEB-1556300 and 1736253.

1.   Otto SP, Gerstein AC. The evolution of haploidy and diploidy. Curr Biol.
2008;18(24):R1121-4.
2.   Gerstein AC, Sharp NP. The population genetics of ploidy change in unicellular
fungi. FEMS Microbiol Rev. 2021.
3.   Gerstein AC, McBride RM, Otto SP. Ploidy reduction in Saccharomyces
cerevisiae. Biology Letters. 2008;4(1):91-4.
4.   Mable BK. Ploidy evolution in the yeast Saccharomyces cerevisiae: a test of the
nutrient limitation hypothesis. Journal of Evolutionary Biology. 2001;14(1):157-70.
5.   Charlesworth B. When to be diploid. Nature. 1991;351(6324):273-4.
6.   Crow JF, Kimura M. Evolution in Sexual and Asexual Populations. The American
Naturalist. 1965;99(909):439-50.
7.   Haldane JBS. The Effect of Variation of Fitness. The American Naturalist.
1937;71(735):337-49.
8.   Kondrashov AS, Crow JF. Haploidy or diploidy: which is better? Nature.
1991;351:314-5.
9.   Perrot V, Richerd S, Valero M. Transition from haploidy to diploidy. Nature.
1991;351(6324):315-7.
10.   Otto SP, Goldstein DB. Recombination and the evolution of diploidy. Genetics.
1992;131(3):745-51.
11.   Maynard Smith J, Haigh J. The hitch-hiking effect of a favorable gene. Genetical
Research. 1974;23:23-5.
12.   Orr HA, Otto SP. Does diploidy increase the rate of adaptation? Genetics.
1994;136(4):1475-80.
13.   Ewens W. Mathematical Population Genetics. New York: Springer; 2004.
14.   Raynes Y, Wylie CS, Sniegowski PD, Weinreich DM. Sign of selection on
mutation rate modifiers depends on population size. Proc Natl Acad Sci U S A.
2018;115(13):3422-7.
15.   Simmons MJ, Crow JF. Mutations affecting fitness in Drosophila populations.
Annual review of genetics. 1977;11(1):49-78.
16.   Wilkie A. The molecular basis of genetic dominance. Journal of medical genetics.
1994;31(2):89-98.
17.   Peters AD, Halligan DL, Whitlock MC, Keightley PD. Dominance and
Overdominance of Mildly Deleterious Induced Mutations for Fitness Traits in
Caenorhabditis elegans. Genetics. 2003;165(2):589-99.
18.   Raynes Y, Burch CL, Weinreich DM. When good mutations go bad: how
population size can change the direction of natural selection. bioRxiv.
2021:2021.04.19.440541.
19.   Kimura M. On the evolutionary adjustment of spontaneous mutation rates.
Genetical Research. 1967;9(1):23-34.
20.   Kimura M. On The Probability Of Fixation Of Mutant Genes In A Population.
Genetics. 1962;47:713-9.
21.   Desai MM, Fisher DS, Murray AW. The speed of evolution and maintenance of
variation in asexual populations. Curr Biol. 2007;17(5):385-94.
22.   Whitlock AO, Peck KM, Azevedo RB, Burch CL. An Evolving Genetic
Architecture Interacts with Hill-Robertson Interference to Determine the Benefit of Sex.
Genetics. 2016;203(2):923-36.

23.   Nowak MA, Sasaki A, Taylor C, Fudenberg D. Emergence of cooperation and
evolutionary stability in finite populations. Nature. 2004;428(6983):646-50.
24.   Gillespie JH. Natural selection for within-generation variance in offspring number.
Genetics. 1974;76(3):601-6.
25.   Aggeli D, Li Y, Sherlock G. Changes in the distribution of fitness effects and
adaptive mutational spectra following a single first step towards adaptation. Nat
Commun. 2021;12(1):5193.
26.   Venkataram S, Dunn B, Li Y, Agarwala A, Chang J, Ebel ER, et al. Development
of a Comprehensive Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast.
Cell. 2016;166(6):1585-96 e22.
27.   Fisher KJ, Buskirk SW, Vignogna RC, Marad DA, Lang GI. Adaptive genome
duplication affects patterns of molecular evolution in Saccharomyces cerevisiae. PLOS
Genetics. 2018;14(5):e1007396.
28.   Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, Sherlock G.
Quantitative evolutionary dynamics using high-resolution lineage tracking. Nature.
2015;519(7542):181-6.
29.   Voordeckers K, Kominek J, Das A, Espinosa-Cantu A, De Maeyer D, Arslan A, et
al. Adaptation to High Ethanol Reveals Complex Evolutionary Pathways. PLoS Genet.
2015;11(11):e1005635.
30.   Dickinson WJ. Synergistic Fitness Interactions and a High Frequency of
Beneficial Changes Among Mutations Accumulated Under Relaxed Selection in
Saccharomyces cerevisiae. Genetics. 2008;178(3):1571-8.
31.   Gerstein AC, Otto SP. Cryptic fitness advantage: diploids invade haploid
populations despite lacking any apparent advantage as measured by standard fitness
assays. PLoS One. 2011;6(12):e26599.
32.   Gerstein AC, Chun H-JE, Grant A, Otto SP. Genomic Convergence toward
Diploidy in Saccharomyces cerevisiae. PLoS Genet. 2006;2(9):e145.
33.   Harari Y, Ram Y, Rappoport N, Hadany L, Kupiec M. Spontaneous Changes in
Ploidy Are Common in Yeast. Current Biology. 2018;28(6):825-35.e4.