# Sign inversion in selection on ploidy

Yevgeniy Raynes and Daniel M. Weinreich

Department of Ecology, Evolution, and Organismal Biology and
Center for Computational Molecular Biology
Brown University Box G-W, Providence, RI 02912 USA

email: yevgeniy_raynes@brown.edu

## Abstract

Ploidy – the number of homologous chromosome sets in a cell – is remarkably variable across the natural world, yet the evolutionary processes that have resulted in such diversity remain poorly understood. Here we use stochastic agent-based simulations to model ploidy evolution under the influence of indirect selection, i.e., selection mediated solely by statistical associations with fitness-affecting mutations. We find that in non-equilibrium asexual populations, the sign of selection on ploidy can change with population size – a phenomenon we have previously termed sign inversion. In large populations, ploidy dynamics are dominated by indirect effects of selection on beneficial mutations, which favors haploids over diploids. However, as population size declines, selection for beneficial mutations is neutralized by random genetic drift before drift can overwhelm selection against the cost of the deleterious mutational load. As a result, in small populations indirect selection is dominated by the cost of the deleterious load, which favors diploids over haploids. Our work adds to the growing body of evidence challenging established evolutionary theory that population size can affect only the efficiency, but not the sign, of natural selection.

## Introduction

Why do some species have a single set of homologous chromosomes while others have two or more? The evolutionary forces that govern transitions between haploidy, diploidy, and polyploidy and have produced the variability we see today have long been a subject of significant theoretical and empirical interest (reviewed in: Otto & Gerstein, 2008, Gerstein & Sharp, 2021). Theories for the role that natural selection plays in ploidy evolution consider either the immediate effects of ploidy on the phenotype or longer-term consequences for the supply rate of fitness-affecting

mutations and the efficacy of selection (Otto & Gerstein, 2008). When the phenotypic
effects of ploidy alter the fitness of an individual, ploidy itself may be directly influenced
by selection. For example, haploid yeast cells are generally smaller and have a higher
surface area to volume ratio than diploid yeast cells, which may afford them higher
fitness in nutrient limited conditions (Gerstein & Sharp, 2021, Gerstein et al., 2008, but
see Mable, 2001). Ploidy variants may also experience indirect selection acting through
statistical associations (i.e., linkage disequilibrium) that develop with fitness-affecting
mutations. Indirect selection is particularly effective in asexual populations as
recombination can quickly erode linkage disequilibrium.

Most theoretical studies of indirect selection have made progress by considering
the effects of selection on deleterious mutations separately from the effects of selection
on beneficial mutations. Selection against deleterious mutations can favor diploids over
haploids in non-equilibrium populations as diploids can mask the effects of recessive or
partially dominant deleterious mutations: i.e., mutations that are fully expressed in
haploids but only partially, or not at all, in diploids, where they first appear in
heterozygous state (Charlesworth, 1991). In other words, while in haploids such
mutations lower fitness by a selection coefficient $s_d$, in heterozygous diploids their effect
on fitness is reduced to $h_d s_d$; $h_d$, known as the dominance coefficient, measures the
dominance of the deleterious mutation and is below 1 for any but the fully dominant
mutations. However, this advantage in the fitness cost of deleterious mutations may
only be a transient one. Once at mutation-selection equilibrium, diploids – with twice as
many loci as haploids – end up bearing twice the cost of the deleterious mutational load
(Crow & Kimura, 1965, Haldane, 1937). That is, while the equilibrium mean fitness of a
haploid populations is $1-U_d$ (where $U_d$ is the deleterious mutation rate) diploids have an
equilibrium mean fitness of $1-2U_d$ (Crow & Kimura, 1965). Nevertheless, two earlier
studies have shown that masking of deleterious mutations could favor diploids at
equilibrium in certain limited circumstances. Kondrashov and Crow (Kondrashov &
Crow, 1991) found that diploid equilibrium mean fitness could exceed that of haploids
under truncation selection and a relatively low $h_d$ (below 0.25). Meanwhile, Perrot et al
(Perrot et al., 1991) demonstrated that diploids could invade haploids in populations
with random mating between haploids and diploids, high rate of free recombination, and
$h_d$ below 0.5; later work confirmed that diploids fail to supplant haploids in populations
with rare recombination (Otto & Goldstein, 1992, Otto & Marks, 1996).

Diploids may also be favored because of their increased supply of beneficial
mutations, given their 2-fold advantage in beneficial mutation rate. Specifically, if
recombination is sufficiently rare diploids may preferentially increase in frequency by
hitchhiking (Maynard Smith & Haigh, 1974) along with those beneficial mutations that
survive drift and sweep to fixation. However, beneficial mutations with dominance
coefficient, $h_b$, below 1 would also be at least partially masked in diploids, reducing their

fitness advantage to $h_bs_b$ and lowering their fixation probability compared to those in
haploids. This advantage in the efficacy of selection could, in certain circumstances,
favor haploids. Orr and Otto (Orr & Otto, 1994) compared the rates of adaptive evolution
in diploids and haploids. They found that diploids adapt faster than haploids when the
rate of adaptation is limited by the supply of beneficial mutations – e.g., when beneficial
mutations are rare or population size is small – provided new mutations are at least
partially dominant ($h_b > 0.5$). On the other hand, as long as $h_b < 1$, diploids evolve
slower than haploids when the rate of adaptation is limited not by the supply of
beneficial mutations – e.g., in large populations in which such mutations are common –
but by the strength of selection for beneficial mutations.

In contrast to most earlier treatments, here we model the evolution of ploidy in
non-equilibrium populations with both beneficial and deleterious mutations. Using
individual-based simulations we show that, in the absence of recombination, whether
ploidy variants are favored or disfavored by indirect selection depends on population
size. In small asexual populations, selection against the deleterious mutational load
favors diploids and disfavors haploids. In large asexual populations, the cost of the
deleterious load is overwhelmed by selection for beneficial mutations, which favors
haploids over diploids.

**Methods**

We model asexual, clonally reproducing populations of constant size, $N$ evolving
in discrete, non-overlapping generations. Populations are composed of genetic lineages
defined by ploidy state and the number of beneficial and deleterious mutations. Each
new beneficial mutation increases an individual's fitness by a constant $s_b$; a new
deleterious mutation reduces fitness by a constant $s_d$. We assume that fitness effects
are additive and calculate the fitness of a haploid lineage with $x$ beneficial and $y$
deleterious mutations as $w_{xy} = 1+xs_b-ys_d$. As in (Orr & Otto, 1994) we also assume that
in the absence of recombination fitness-affecting mutations in diploids remain
permanently heterozygous (at least on the time scale of our simulations) and calculate
the fitness of a diploid lineage with $x$ beneficial and $y$ deleterious mutations as
$w_{xy}=1+xh_bs_b-yh_ds_d$, where $h_b$ and $h_d$ are the corresponding dominance coefficients.

At the outset of simulation, a single individual of a particular ploidy (either haploid
or diploid) is introduced to a population of $N$-1 individuals of the other ploidy. Both the
invading lineage of size 1 and the resident lineage of size $N$-1 bear no fitness-affecting
mutations and, thus, have an initial fitness $w = 1$. Every generation the population
reproduces according to the Wright-Fisher model (Ewens, 2004) with the size of the
lineage carrying $x$ beneficial mutations and $y$ deleterious mutation in generation $t$+1
randomly sampled from a multinomial distribution with expectation $Nf_t \cdot w_{xy}/\bar{w}$ where $N$ is

the size of the population, $f_t$ is the frequency of the lineage in generation $t$, $w_{xy}$ is the
fitness of the lineage as defined above and $\bar{w}$ is the average fitness of the population.
Upon reproduction, each surviving lineage acquires a random number of beneficial and
deleterious mutations drawn from a Poisson distribution with expectation given by,
respectively, $cN_iU_b$ and $cN_iU_d$. Here, $c$ is the ploidy of the lineage (i.e., number of
chromosome copies), $N_i$ is the size of the lineage, $U_b$ is the per-individual beneficial
mutation rate, and $U_d$ is the per-individual deleterious mutation rate. Diploids, thus, have
a 2-fold higher mutation rate than haploids. For simulations in Fig. 2C we lowered the
diploid mutation rate to that of haploids by setting $c = 1$ for both diploid and haploid
lineages.

Simulations end when the invader either fixes (reaches a frequency of 1.0) or is
eliminated from the population. We assess the fixation probably ($P_{fix}$) as the frequency
with which the invader supplants the resident across the replicate runs of the simulation.
Throughout, we focus on $NP_{fix}$ - the fixation probability normalized by the neutral
expectation, $1/N$. To ascertain whether an invading ploidy variant is favored or
disfavored by selection we compare its $NP_{fix}$ to that expected of a neutral mutation, i.e.,
$NP_{fix} = 1$ (e.g., Raynes et al., 2018).

Simulation code was written in Julia 1.2 and is available at
https://github.com/yraynes/Ploidy.

## Results

We first examined the fixation probability of haploid invaders in diploid
populations. Since deleterious mutations in nature are predominantly recessive (e.g.,
Simmons & Crow, 1977, Wilkie, 1994, Peters et al., 2003), we set the dominance
coefficient of deleterious mutations in our simulations to be $h_d = 0.0$. Meanwhile, we
varied the dominance coefficient of new beneficial mutations from fully recessive ($h_b =$
0.0) to fully dominant ($h_b = 1.0$). Strikingly, we found that the sign of selection on
haploids can change with population size (Fig 1A). For all values of $h_b < 1.0$ studied,
haploids are favored by selection (have $NP_{fix} > 1$) in large populations but become
disfavored (have $NP_{fix} < 1$) as population size decreases below a critical size we call
$N_{crit}$. We have previously designated this $N$-dependent switch in the sign of selection
*sign inversion* (Raynes et al., 2018, Raynes et al., 2021). Fig. S1 confirms that haploids
are susceptible to sign inversion across a wide – at least over several orders of
magnitude – range of mutation rates.

**Figure 1: Sign inversion in selection on haploids. A)** Normalized fixation probability ($NP_{fix}$) of
haploid invaders (circles) in diploid populations at different $h_b$. Red dashed line: $P_{fix}(s = -U_d, N)$;
Teal dashed line: $NP_{fix}^{hitch}$. Horizontal black line shows the normalized fixation probability of a
neutral mutation ($NP_{fix}$ = 1). **B)** Average fitness of winning haploid populations, $w$. Horizontal
black lines indicate $w$ = 1 and $w$ = 1 + $s_b$ = 1.1 **C)** Conditional fixation probability calculated
across simulations in which haploid invaders managed to acquire at least one beneficial
mutation. Horizontal black line shows expected $P_{fix}(s_b, N)$. In all panels, parameter values are as
follows: $s_b$ = 0.1, $s_d$ = -0.1, $U_d$ = 0.01, $U_b$ = 0.001. All simulation results averaged across at least
$10^7$ replicates.

Based on our earlier work on sign inversion (Raynes et al., 2021, Raynes et al.,
2018) we assumed that in small populations haploid $NP_{fix}$ would be reduced below the
neutral expectation by selection against the deleterious load. To test this expectation,
we compared haploid $NP_{fix}$ in simulations to that predicted by the fitness cost of the
haploid deleterious load. Specifically, Kimura (Kimura, 1967) showed that an allele that
increases the deleterious mutation rate by $\Delta U_d$ bears a fitness cost at equilibrium of $s$ =
$-\Delta U_d$. Given that deleterious mutations in our simulations are permanently heterozygous
and fully recessive we can assume that the deleterious mutation rate in diploid residents
is essentially zero. Therefore, $\Delta U_d$ is simply the deleterious mutation rate of the haploid
and the expected cost of the load in haploids is given by $s$ = $-U_d$. The probability of
fixation for a mutation of fitness effect $s$ (in a haploid population) is given classically by
$P_{fix}(s, N)$ = $(1-e^{-2s})/(1-e^{-2Ns})$ (Kimura, 1962). $P_{fix}(s = -U_d, N)$ is shown with the red dashed
line in Fig. 1A and agrees very well with simulations, confirming that in sufficiently small
populations haploid $NP_{fix}$ is set by the cost of the deleterious load.

Again motivated by our earlier work, we hypothesized that in large populations
that favor haploids, haploid $NP_{fix}$ is elevated above the neutral expectation by selection
for beneficial mutations. To confirm the influence of selection for beneficial mutations on
haploid $NP_{fix}$ we assayed the final fitness of populations in which haploids had fixed
(Fig. 1B). We found that at smallest $N$, the average fitness of winning haploid

populations is, in fact, lower than the ancestral fitness, confirming that at these
population sizes, hitchhiking with beneficial mutations (which would necessitate an
average fitness of at least $1+s_b$) is not a significant influence on $NP_{fix}$. Rather, haploids
fix in these populations via random genetic drift. However, as Fig. 1B shows, in
sufficiently large populations haploid winners' fitness surpasses $1+s_b$ for all values of $h_b$
examined. This demonstrates that in large populations haploids fix almost exclusively by
hitchhiking. In fact, note that at yet larger $N$ haploid fitness may surpass $1+2s_b$
suggesting that in those populations haploids reach fixation with multiple beneficial
mutations. For now, we leave further examination of multiple-mutation (Desai et al.,
2007) dynamics in ploidy evolution for future work.

Moreover, we found that at $h_b = 0$ haploid $NP_{fix}$ could be well approximated by
the expected probability of hitchhiking to fixation with a beneficial mutation, previously
derived for a mutator allele by Raynes et al. (2018). In brief, to fix by hitchhiking an
invader must first produce a beneficial mutation, which then has to sweep to fixation.
The probability that the invader will produce a beneficial mutation is approximately equal
to the ratio of $U_b$ to the total mutation rate, $U_b + U_d$. Since at $h_b = 0$ the spread of
beneficial mutations in the haploid background is unaffected by selection on beneficial
mutations in diploids, the probability of the latter for a mutation of fitness effect $s_b$ can be
estimated with $P_{fix}(s_b, N)$ given above. The probability of hitchhiking to fixation is the
product of these two probabilities: $P_{fix}^{hitch} = U_b/(U_b + U_d) \times P_{fix}(s_b, N)$. $NP_{fix}^{hitch}$ is shown
with a teal line in Fig. 1A.

Finally, Fig. 1A also highlights the negative relationship between haploid $NP_{fix}$
and $h_b$. As may be expected, $h_b$ has little effect on $NP_{fix}$ in small populations, in which
$NP_{fix}$ is set by the deleterious load while $P_{fix}^{hitch}$ – whether for haploids or diploids – is
miniscule. However, above $N_{crit}$ haploids fare worse at higher $h_b$ suggesting that
increasing dominance of beneficial mutations lowers $P_{fix}^{hitch}$ (which dominates $NP_{fix}$ in at
larger $N$). Focusing on the two components of $P_{fix}^{hitch}$ above, we can assume that the
probability of a haploid acquiring a beneficial mutation is unaffected by $h_b$. However, the
likelihood of the beneficial mutation escaping drift and reaching fixation may be reduced
at higher $h_b$ by stronger competition from beneficial mutations in the diploids. In other
words, whereas at $h_b = 0$, haploids with a beneficial mutation have an advantage $s_b$ over
any diploid similarly endowed with a beneficial mutation, that advantage decreases to
$s_b$-$h_b s_b$ for any $h_b > 0.0$. (At $h_b = 1.0$ haploids enjoy no advantage in the strength of
selection on associated beneficial mutations; instead, diploids that benefit from a two-
fold higher $U_b$ and a lower deleterious load are favored by selection at all $N$).

To test how interference from beneficial mutations in diploids reduces haploid
$P_{fix}^{hitch}$ at $h_b > 0$, we assayed the probability of haploid fixation conditioned on acquiring
at least one beneficial mutation, $P_{fix|ben}$ (Fig. 1C). We found that at $h_b = 0$, $P_{fix|ben}$

asymptotes to the expected probability of fixation given by $P_{fix}(s_b, N)$, which shows that beneficial mutations in haploids are unaffected by competition from fully recessive mutations in diploids. However, at $h_b > 0$, $P_{fix|ben}$ declines below the expected $P_{fix}(s_b, N)$, confirming that competition from beneficial mutations in diploid residents reduces the probability of fixation of new beneficial mutations in haploids. Note also that $P_{fix|ben}$ declines with increasing $N$ and does so faster at higher $h_b$, consistent with intensifying interference from diploid residents, modulated by both more effective selection and the rising supply of beneficial mutations (i.e., $NU_b$) in diploids.

Next, we studied the fixation probability of diploid invaders in haploid populations. Once again, we set deleterious mutations to be fully recessive ($h_d = 0.0$), while beneficial mutations ranged from fully recessive ($h_b = 0.0$) to fully dominant ($h_b = 1.0$). Fig. 2A confirms that diploids are also susceptible to sign inversion. Unlike that of haploids, though, diploid $NP_{fix}$ has a negative slope at $N_{crit}$. For all studied $h_b < 1$, diploids are favored in small populations and become disfavored as $N$ increases. At $h_b = 1.0$ diploids are everywhere favored over haploids (have $NP_{fix} > 1$ at all $N$).

**Figure 2: Sign inversion in selection on diploids. A)** Normalized fixation probability ($NP_{fix}$) of diploid invaders (circles) in haploid populations at different $h_b$. Teal dashed line: $P_{fix}(s = U_d, N)$. Horizontal black line shows the normalized fixation probability of a neutral mutation ($NP_{fix} = 1$). **B)** Average fitness of winning diploid populations, $w$. Horizontal lines indicate $w = 1$ in black and $w = 1 + h_b s_b$ in corresponding colors. **C)** Comparison between $NP_{fix}$ in panel A (circles again) and diploid $NP_{fix}$ in simulations in which diploid $U$ was set to equal that of the haploids (shown for convenience as dashed lines, results only available at the same $N$ as circles). Horizontal black line: $NP_{fix} = 1$. In all panels, parameter values are as follows: $s_b = 0.1$, $s_d = -0.1$, $U_d = 0.01$, $U_b = 0.001$. All simulation results averaged across $10^7$ replicates.

Our analysis of haploid $NP_{fix}$ above suggests that in sufficiently small populations, diploids would mostly reach fixation not by acquiring and fixing beneficial

mutations but by simply outlasting haploid residents that succumb to selection against
the load. Fig. 2A confirms that diploid $NP_{fix}$ in small populations is well predicted simply
by the fixation probability for $s = +\Delta U_d$ – the diploid's expected fitness advantage due to
the deleterious load borne by haploids. Indeed, assaying the average fitness at the end
of simulation confirms that in sufficiently small populations diploid winners are
indistinguishable from their ancestors (Fig. 2B).

In larger populations though, in which haploid dynamics are dominated by
beneficial mutations, we expected diploid fate to be likewise determined by the
likelihood of hitchhiking with beneficial mutations of their own. Correspondingly, we
found that the fitness of diploid winners increases with $N$ and in sufficiently large
populations rises above $1+h_b s_b$ (Fig. 2B). This suggests that in those populations
diploids also fix exclusively by hitchhiking with beneficial mutations. (Except at $h_b =0$,
where diploid winners never exceed ancestral fitness as hitchhiking with completely
recessive beneficial mutations is impossible at any $N$.) For a diploid to fix by hitchhiking,
though, a beneficial mutation of effect $h_b s_b$ must sweep to fixation without interference
from beneficial mutations of larger effect $s_b$ in haploids. Diploids possess an inherent
advantage of a two-fold higher mutation rate so a diploid invader is always more likely to
produce a beneficial mutation than any of its haploid competitors. This advantage is
expected to be most pronounced in relatively small populations in which the rate of
adaptation is limited by the supply rate of beneficial mutations (Orr & Otto, 1994). Our
simulations confirm that the higher $U_b$ improves diploid probability of hitchhiking.
Specifically, as shown in Fig. 2C diploid $NP_{fix}$ is reduced when diploid mutation rate is
lowered to that of haploids (see Methods). More importantly, Fig. 2C shows that making
diploid $U_b$ equal to that of haploids lowers $N_{crit}$ for sign inversion for all $0 < h_b < 1.0$,
suggesting that at some intermediate $N$, diploid $NP_{fix}$ is above 1 not only due to purifying
selection against the load but also positive selection mediated by a higher $U_b$ (note that
in simulations without deleterious mutations, diploids are only ever favored by selection
at $h_b > 0.5$ (Fig. S2), consistent with (Orr & Otto, 1994)).

However, as long as $h_b < 1$, haploids enjoy an advantage in the strength of
selection on new beneficial mutations (i.e., beneficial mutations in diploids endow
carriers with a smaller fitness advantage $h_b s_b$ relative to $s_b$ in haploids). This advantage
is magnified as population size increases and selection for beneficial mutations
becomes more efficient (while the supply rate of such mutations to haploids increases
as well). In sufficiently large populations selection for beneficial mutations in haploids
offsets the cost of the load and the diploids' advantage in $U_b$. When it does, diploid $NP_{fix}$
(set at that point almost exclusively by diploid $NP_{fix}^{hitch}$, see above) declines below 1 at
all $h_b < 1.0$ (Fig. 2A). Correspondingly, diploids fare better at higher $h_b$, consistent with a
reduced fitness advantage ($s_b - h_b s_b$) of beneficial mutations in haploids. And, diploid
$NP_{fix}$ is never below 1 when beneficial mutations are fully dominant: at $h_b = 1$, haploids

have no advantage in the efficacy of selection, whereas diploids still enjoy the benefit of
a two-fold higher $U_b$.

## Discussion

We simulated evolution of ploidy in population with recessive deleterious
mutations and beneficial mutations of varying dominance. We showed that in
populations with both kinds of mutations, the sign of selection on ploidy can change with
population size – a phenomenon we have previously designated sign inversion (Raynes
et al., 2018). Consistent with previous expectations (Charlesworth, 1991) for non-
equilibrium populations, we found that selection against recessive deleterious mutations
lowers haploid fixation probability. Indeed, in sufficiently small populations the
evolutionary fate of ploidy variants is entirely dominated by selection against the cost of
the deleterious load. Even in the presence of beneficial mutations, haploid invaders that
bear the full cost of the deleterious mutations fare worse than the neutral expectation
(Fig. 1A). Meanwhile, diploid invaders that mask deleterious mutations and, thus, can
avoid the fitness cost of the load fare better than the neutral expectation (Fig. 2A).

On the other hand, in large populations, ploidy evolution is dominated by the
indirect effects of selection on beneficial mutations. Previously, it's been shown that
selection for beneficial mutations may favor diploids – that produce twice as many
mutations as haploids – when the rate of adaptation is limited by the mutation rate (i.e.,
the product $NU$ is small). We find that the benefit of a 2-fold higher $U_b$ (in tandem with
the cost of haploids' load) may indeed raise diploid $NP_{fix}$ above the neutral expectation
at some $N$ that would otherwise favor haploids (Fig. 2C). Conversely, haploids have
been predicted to fare better than diploids if new beneficial mutations are sufficiently
common that the rate of adaptation is limited not by their supply but by their rate of
spread (Orr & Otto, 1994). The reason for this is that mutations in haploids have larger
effects on fitness ($s_b$ versus $h_b s_b$) and are, thus, more likely to survive drift and reach
fixation than mutations in diploids. Our work confirms that in sufficiently large
populations the probability of hitchhiking to fixation exceeds the neutral expectation for
haploids but is below the neutral expectation for diploids. Correspondingly, in large
populations haploids fare better than neutral mutations, while diploid $NP_{fix}$ drops below
1.  The expression for $NP_{fix}^{hitch}$ plotted in Fig. 1A for $h_b=0$ illustrates why haploid $NP_{fix}$
transitions from below 1 to above 1 with $N$. Recall that $NP_{fix}^{hitch}$ is approximately the
product of two probabilities: that of acquiring a beneficial mutation and that of the
beneficial mutation reaching fixation. The probability of generating a beneficial mutation
(assuming that one is produced) is small and well below one, given that $U_d \gg U_b$.
$NP_{fix}(s_b, N)$ of a beneficial mutation is always above the neutral expectation and is a
monotonically increasing function of $N$. And while it may be reduced at $h_b > 0$ by

competition from beneficial mutations in diploids, mutations in haploids should still be
favored over those in diploids by a selection coefficient of $s_b$-$h_b s_b$ as long as $h_b$ < 1.0.
Thus, when $N$ is small, the product of the two, $NP_{fix}^{hitch}$, is reduced below the neutral
expectation by the low probability of generating a beneficial mutation. However, as
population size increases above $N_{crit}$ and selection on new beneficial mutations
becomes more efficient ($NP_{fix}(s_b, N)$ of new beneficial mutations rises further away from
1), $NP_{fix}^{hitch}$ must necessarily increase above the neutral expectation at some sufficiently
large $N_{crit}$, lifting the total $NP_{fix}$ above 1 as well.

Taken together, our results demonstrate that sign inversion occurs because
ploidy variants can experience both the positive selection mediated by linkage to
beneficial mutations and the purifying selection mediated by linkage to deleterious
mutations that are overwhelmed by drift at different $N$. That is, haploid $NP_{fix}$ is
dominated by selection on beneficial mutation at large $N$ (represented by $NP_{fix}^{hitch}$) but as
population size decreases it is neutralized by drift ($NP_{fix}^{hitch}$ crosses 1) before selection
against the cost of the load, which leads to sign inversion.

Previously, we have found evidence of sign inversion in four published studies of
the evolution of recombination (Whitlock et al., 2016), cooperation (Nowak et al., 2004),
variance in offspring number (Gillespie, 1974), and the mutation rate (Raynes et al.,
2018). Having examined all four cases of sign inversion (Raynes et al., 2021) we
discovered that all alleles susceptible to sign inversion can impart both fitness costs and
fitness benefits on their carriers. That is, similar to ploidy variants, known cases of sign
inversion exhibit variability in the sign of selection on individual carriers. We designated
such alleles as having sign-variable effects and showed that that all sign-variable effects
could be classified into either those that vary between different mutant lineages or
among individual members of a single lineage (modulated, for example, by temporal
changes in the environment or population composition) (Raynes et al., 2021). Ploidy in
our model exhibits between-lineage variability. Haploid lineages that generate beneficial
mutations enjoy the indirect fitness benefit and a higher than neutral probability of
fixation. Meanwhile, lineages that generate deleterious mutations bear the indirect
fitness cost and, correspondingly, have a lower than neutral probability of fixation. More
importantly, in the same earlier study we showed that sign variability is necessary for
sign inversion, which always occurs because selection on cost-bearing and benefit-
endowed carriers exhibit differing sensitivities to drift (Raynes et al., 2021). Our current
work showing that sign inversion in selection on ploidy is also driven by between-
lineage sign variability corroborates the necessity of sign variability for sign inversion
and provides additional support for the common mechanistic basis of sign inversion.

Finally, our findings may be particularly relevant for the growing number of
evolution experiments that have documented ploidy transitions in laboratory microbial

populations – those expected to be relatively far from equilibrium with access to both
beneficial and deleterious mutations. In particular, studies in *Saccharomyces cerevisiae*
have repeatedly observed shifts from haploidy to diploidy through spontaneous genome
duplication (Aggeli et al., 2021, Venkataram et al., 2016, Fisher et al., 2018, Levy et al.,
2015, Voordeckers et al., 2015). Interestingly, most of these studies have been
conducted in large populations that, we have found, should indirectly favor haploids
rather than diploids. Accordingly, several studies have documented an inherent fitness
advantage to diploidy (Aggeli et al., 2021, Venkataram et al., 2016, Fisher et al., 2018)
that would allow diploids to fix through the action of direct selection instead. Others,
however, did not detect a significant direct advantage to diploid cells over haploids
(Dickinson, 2008, Gerstein & Otto, 2011), suggesting that diploids might have arisen via
indirect selection. Yet, whereas the appearance of diploids in small populations of the
mutation accumulation experiment of Dickinson (Dickinson, 2008) appears consistent
with our results, diploids repeatedly overtaking large populations in the study of Gerstein
et al. (Gerstein et al., 2006) is difficult to reconcile with our simulations. It is possible that
ploidy evolution, at least in yeast, is universally driven by direct selection and an
inherent fitness advantage will eventually be identified in all laboratory conditions in
which diploids have been shown to overtake haploids (see also (Gerstein & Sharp,
2021) for the discussion of "ploidy drive"). It is also possible, though, that experimental
populations differ in some properties that affect indirect selection on ploidy in such a
way as to indirectly favor diploids in some environments and disfavor them in others,
even at relatively large $N$ (see also (Harari et al., 2018)). For example, our work
suggests that diploids could fare better in larger populations given a higher $h_b$ or $U_d$
(which would reduce haploid $P_{fix}^{hitch}$), both of which remain somewhat difficult to assay
experimentally. All in all, while our work adds to our understanding of the mechanics of
indirect selection on ploidy variants, whether indirect selection can be the driving force
in ploidy evolution in laboratory or in nature remains to be determined.

**References**

Aggeli, D., Li, Y. & Sherlock, G. 2021. Changes in the distribution of fitness effects and adaptive mutational spectra following a single first step towards adaptation. *Nature Communications* **12**: 5193.

Charlesworth, B. 1991. When to be diploid. *Nature* **351**: 273-274.

Crow, J. F. & Kimura, M. 1965. Evolution in Sexual and Asexual Populations. *The American Naturalist* **99**: 439-450.

Desai, M. M., Fisher, D. S. & Murray, A. W. 2007. The speed of evolution and maintenance of variation in asexual populations. *Curr Biol* **17**: 385-94.

Dickinson, W. J. 2008. Synergistic Fitness Interactions and a High Frequency of Beneficial Changes Among Mutations Accumulated Under Relaxed Selection in Saccharomyces cerevisiae. *Genetics* **178**: 1571-1578.

Ewens, W. 2004. *Mathematical Population Genetics.* Springer, New York.

Fisher, K. J., Buskirk, S. W., Vignogna, R. C., Marad, D. A. & Lang, G. I. 2018. Adaptive genome duplication affects patterns of molecular evolution in Saccharomyces cerevisiae. *PLOS Genetics* **14**: e1007396.

Gerstein, A. C., Chun, H.-J. E., Grant, A. & Otto, S. P. 2006. Genomic Convergence toward Diploidy in Saccharomyces cerevisiae. *PLoS Genet* **2**: e145.

Gerstein, A. C., McBride, R. M. & Otto, S. P. 2008. Ploidy reduction in Saccharomyces cerevisiae. *Biology Letters* **4**: 91-94.

Gerstein, A. C. & Otto, S. P. 2011. Cryptic fitness advantage: diploids invade haploid populations despite lacking any apparent advantage as measured by standard fitness assays. *PLoS One* **6**: e26599.

Gerstein, A. C. & Sharp, N. P. 2021. The population genetics of ploidy change in unicellular fungi. *FEMS Microbiol Rev*.

Gillespie, J. H. 1974. Natural selection for within-generation variance in offspring number. *Genetics* **76**: 601-606.

Haldane, J. B. S. 1937. The Effect of Variation of Fitness. *The American Naturalist* **71**: 337-349.

Harari, Y., Ram, Y., Rappoport, N., Hadany, L. & Kupiec, M. 2018. Spontaneous Changes in Ploidy Are Common in Yeast. *Curr Biol* **28**: 825-835 e4.

Kimura, M. 1962. On The Probability Of Fixation Of Mutant Genes In A Population. *Genetics* **47**: 713-719.

Kimura, M. 1967. On the evolutionary adjustment of spontaneous mutation rates. *Genetical Research* **9**: 23-34.

Kondrashov, A. S. & Crow, J. F. 1991. Haploidy or diploidy: which is better? *Nature* **351**: 314-315.

Levy, S. F., Blundell, J. R., Venkataram, S., Petrov, D. A., Fisher, D. S. & Sherlock, G. 2015. Quantitative evolutionary dynamics using high-resolution lineage tracking. *Nature* **519**: 181-186.

Mable, B. K. 2001. Ploidy evolution in the yeast Saccharomyces cerevisiae: a test of the nutrient limitation hypothesis. *Journal of Evolutionary Biology* **14**: 157-170.

Maynard Smith, J. & Haigh, J. 1974. The hitch-hiking effect of a favorable gene. *Genetical Research* **23**: 23-25.

Nowak, M. A., Sasaki, A., Taylor, C. & Fudenberg, D. 2004. Emergence of cooperation and evolutionary stability in finite populations. *Nature* **428**: 646-50.

Orr, H. A. & Otto, S. P. 1994. Does diploidy increase the rate of adaptation? *Genetics* **136**: 1475-80.

Otto, S. P. & Gerstein, A. C. 2008. The evolution of haploidy and diploidy. *Curr Biol* **18**: R1121-4.

Otto, S. P. & Goldstein, D. B. 1992. Recombination and the evolution of diploidy. *Genetics* **131**: 745-751.

Otto, S. P. & Marks, J. C. 1996. Mating systems and the evolutionary transition between haploidy and diploidy. *Biological Journal of the Linnean Society* **57**: 197-218.

Perrot, V., Richerd, S. & Valero, M. 1991. Transition from haploidy to diploidy. *Nature* **351**: 315-7.

Peters, A. D., Halligan, D. L., Whitlock, M. C. & Keightley, P. D. 2003. Dominance and Overdominance of Mildly Deleterious Induced Mutations for Fitness Traits in Caenorhabditis elegans. *Genetics* **165**: 589-599.

Raynes, Y., Burch, C. L. & Weinreich, D. M. 2021. When good mutations go bad: how population size can change the direction of natural selection. *bioRxiv*: 2021.04.19.440541.

Raynes, Y., Wylie, C. S., Sniegowski, P. D. & Weinreich, D. M. 2018. Sign of selection on mutation rate modifiers depends on population size. *Proc Natl Acad Sci U S A* **115**: 3422-3427.

Simmons, M. J. & Crow, J. F. 1977. Mutations affecting fitness in Drosophila populations. *Annual review of genetics* **11**: 49-78.

Venkataram, S., Dunn, B., Li, Y., Agarwala, A., Chang, J., Ebel, E. R., Geiler-Samerotte, K., Herissant, L., Blundell, J. R., Levy, S. F., Fisher, D. S., Sherlock, G. & Petrov, D. A. 2016. Development of a Comprehensive Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast. *Cell* **166**: 1585-1596 e22.

Voordeckers, K., Kominek, J., Das, A., Espinosa-Cantu, A., De Maeyer, D., Arslan, A., Van Pee, M., van der Zande, E., Meert, W., Yang, Y., Zhu, B., Marchal, K., DeLuna, A., Van Noort, V., Jelier, R. & Verstrepen, K. J. 2015. Adaptation to High Ethanol Reveals Complex Evolutionary Pathways. *PLoS Genet* **11**: e1005635.

463 Whitlock, A. O., Peck, K. M., Azevedo, R. B. & Burch, C. L. 2016. An Evolving Genetic
464    Architecture Interacts with Hill-Robertson Interference to Determine the Benefit of
465    Sex. *Genetics* **203**: 923-36.
466 Wilkie, A. 1994. The molecular basis of genetic dominance. *Journal of medical genetics*
467    **31**: 89-98.

468