Expectation violations enhance neuronal encoding of sensory information in mouse primary visual cortex

Matthew F. Tang[1,2,3 #*], Ehsan Kheradpezhouh[1,2 #], Conrad C.Y. Lee[1,2,3], J. Edwin Dickinson[4], Jason B. Mattingley[2,3,5,6], and Ehsan Arabzadeh[1,2]

1. Eccles Institute of Neuroscience, John Curtin School of Medical Research, The Australian National University, Canberra, ACT, Australia

2. Australian Research Council Centre of Excellence for Integrative Brain Function, Victoria, Australia

3. Queensland Brain Institute, The University of Queensland, Brisbane, QLD, Australia

4. School of Psychological Sciences, The University of Western Australia, Perth, WA, Australia

5. School of Psychology, The University of Queensland, Brisbane, QLD, Australia

6. Canadian Institute for Advanced Research (CIFAR), Toronto, Canada

co-first authors

* Corresponding author

Corresponding author
Matthew F. Tang
Eccles Institute of Neuroscience
John Curtin School of Medical Research
The Australian National University
Acton, ACT, 2601, Australia
Email: matthew.tang@anu.edu.au

31   **Abstract**

The response of cortical neurons to sensory stimuli is shaped both by past events (adaptation)

and the expectation of future events (prediction). Here we employed a visual stimulus

paradigm with different levels of predictability to characterise how expectation influences

orientation selectivity in the primary visual cortex (V1) of mice. We recorded neuronal activity

using two-photon calcium imaging (GCaMP6*f*) while animals viewed sequences of grating

stimuli which either varied randomly in their orientations or rotated predictably with occasional

transitions to an unexpected orientation. For single neurons and the population, there was

significant enhancement in the gain of orientation-selective responses to unexpected gratings.

This gain-enhancement for unexpected stimuli was prominent in both awake and

anaesthetised mice. We implemented a computational model to demonstrate how trial-to-trial

variability in neuronal responses were best characterised when adaptation and expectation

effects were combined.

Introduction

There is often more information in the sensory environment than the brain has the

capacity to fully process. To cope with this information overload, activity within neuronal

circuits is modulated by processes such as adaptation[1,2] attention[3,4], and prediction[5,6]. Neural

adaptation is known to improve the transmission of sensory information in circuits by

accounting for the statistics of past sensory inputs[1,7,8]. Likewise, selective attention can

enhance neural responses to task-relevant features and suppress irrelevant information[3,9]. An

influential theory of neural function argues that predictions about specific future stimuli, based

upon Bayesian inference, might similarly improve the fidelity of stimulus representations[5,6].

Based on this *predictive coding* view, the mammalian cortex is conceptualised as a predictive

machine that uses the statistical regularities of incoming sensory inputs to iteratively generate

an internal model of its external environment. Predictive coding provides a simple theoretical

view of perception which is supported by a substantial body of work in human neuroimaging

and behavioural studies[10,11]. The classic mismatch negativity effect has become a hallmark of

this literature[12,13]. When encountering an unexpected stimulus, the brain generates a

significantly larger evoked response compared with the response following an expected

stimulus[11]. Decoding of activity from electroencephalography (EEG) recordings in humans has

revealed that expectation affects the representation of visual information in the brain[14–18].

Recent work supports the idea that prediction influences single neuron responses

across a number of sensory modalities[19–24]. Theoretical models propose that higher level

processing regions generate inhibitory copies of the expected stimulus which are passed down

the cortical hierarchy to the earlier processing regions[6], where they are integrated with

incoming sensory inputs. If a stimulus is expected, the inhibitory copy should minimise the

neuronal response. By contrast, any mismatch between the expected and presented stimulus

should result in a prominent response.

Here, we tested key elements of predictive coding theory at the neuronal level in mouse

primary visual cortex (V1). We used two-photon calcium imaging in awake mice that were

exposed to sequences of oriented gratings at different levels of predictability. We characterised

how prediction affects orientation selectivity in V1 neurons, and how changes in orientation

tuning modulate the amount of information about the sensory input carried by individual

neurons and neuronal populations. We demonstrate that unexpected stimuli significantly

increase the gain of orientation selectivity without any corresponding changes to the width of

the tuning function. Such increased gain to expectation violations yields increased information

about stimulus features within single-cells and at the level of neuronal populations. This

enhanced representation of unexpected stimuli is present in both awake and anaesthetised

mice. Finally, we use a computational model to quantify the contribution of adaptation and

expectation to neuronal responses at the single trial level.

81   **Results**

We combined experimental and modelling approaches to determine how prediction

affects neuronal responses in mouse (C57BL) V1 cortical neurons to sequences of oriented

grating stimuli. We asked whether the selectivity of individual neurons changes with

expectations about the orientation of future stimuli by presenting sequences of gratings with

different levels of predictability to awake mice ($N = 3$ across 23 sessions in total, 1697

neurons) while imaging Layer 2/3 activity in V1 using two-photon excitation microscopy (Figure

1ABC, Movie 1). The stimulus sequence was adapted from the Allen Brain Institute's Brain

Observatory paradigm[25] used to quantify orientation selectivity. Each sequence consisted of a

series of full-screen gratings (0.034 c/°, 50% contrast) oriented between 0° and 150° in 30°

steps, presented at 4 Hz with no inter-stimulus interval. In the *Random* condition (Figure 1B

and C), the orientations of successive gratings were uncorrelated.

To establish predictions about stimulus orientation, in the *Rotating* condition the grating

rotated either clockwise or anti-clockwise for 5 to 9 presentations (in 30° steps), before jumping

to an unexpected random orientation. In this condition, *Expected events* were those which

constituted the rotating sequence, whereas *Unexpected events* were those in which the

stimulus jumped randomly to an unpredicted orientation. Critically, for unexpected events the

jump from the predicted orientation was to a random orientation matched to the correlation

statistics for the stimulus sequence embedded in the *Random* condition. Figure 1B and C

identify the three types of transitions within the visual stimulation protocol: *Random* transitions

(in blue), *Expected* transitions (in red) and *Unexpected* transitions (in green). Figure 1D and E

show eight example neurons imaged within a field of view, each of which exhibited a varying

degree of orientation selectivity under the *Random* condition. In line with previous work[25],

many imaged neurons showed orientation selectivity for the spatial frequency employed

(462/1697; one-way ANOVA $p < 0.05$ for orientation selectivity).

Figure 1. *Experimental procedure for testing the effects of prediction on orientation selectivity in mouse V1 neurons. (A) Apparatus for using two-photon calcium imaging in combination with visual stimulation. (B) Schematic of the Random and Rotating sequences of oriented gratings. (C) In the Random condition, the orientation of each stimulus was drawn from a pseudo-randomised distribution (uniform probability from 0 to 150° in 30° steps). In the Rotating condition, the gratings rotated clockwise (e.g., 0° -> 30° -> 60°) or anti-clockwise (e.g., 0° -> 150° -> 120°) for 5-9 presentations (red dots) before jumping to a random unexpected orientation (indicated by the green dots). (D) Mean motion-corrected two-photon image from a single session, with individual neurons highlighted in red. (E) Time course of activity in the corresponding neurons highlighted in D in response to different grating orientations from the Random condition. The tuning functions in the right panels show the average response from 0 - 1000 ms after stimulus presentation. Points are fitted with a circular Gaussian with a baseline offset. The key parameters of the fits are given as the gain (amplitude) and width (standard deviation) of the Gaussian for each neuron. Shading and error bars show ±1 standard error over trials.*

123 ***Movie 1.*** *Example sequence of gratings in the Rotating condition. Stimuli rotate in one*
124 *direction for 5 to 9 presentations before jumping to a random orientation and rotating in the*
125 *opposite direction.*
126
Prediction affects single neuron activity

We next examined how orientation selectivity of individual neurons was affected by

stimulus predictability (Figure 2). The three example neurons shown in Figure 2A all exhibit

orientation selectivity from ~85-100 ms after stimulus onset. The first neuron (top row of Figure

2A) responded maximally to gratings at 0°, with slight suppression for the more distant

orientations (60°, 90°, 120°). During presentation of the *Expected* stimulus (red trace),

modulation of neuronal activity began before the onset of the stimulus (0 ms). This pre-

stimulus modulation is due to the rotating nature of the sequence: the stimulus presented at -

500 ms was orthogonal to that presented at 0 ms. This means that in the 0° condition, the anti-

preferred stimulus (90°) was presented at -500 ms, whereas in the 90° condition, the preferred

stimulus (0°) was presented at -500 ms. The rotating nature of the stimuli during the *Expected*

sequence thus produced an idiosyncratic temporal profile in neuronal response. For this

reason, here we focus on the *Random* and *Unexpected* transitions where the stimuli presented

immediately before 0 ms were uncorrelated with the current stimulus.

Figure 2. *Expectation affects orientation-selective responses of individual V1 neurons. (A)*
Time-courses of three example neurons in response to oriented grating stimuli in the expected,
unexpected and random conditions. Each neuron is illustrated in a separate row, with the
rightmost panel showing orientation tuning curves for that neuron. The tuning is measured as
the averaged response from 250 to 1000 ms after stimulus onset (grey shading). The solid
curve is a fitted Gaussian function with a constant offset. (B) Same as in A, but shows activity
for all orientation-selective neurons (N= 462) aligned to their preferred orientation (0°) to allow
averaging. Right panel: Same as in A but showing the Gaussian tuning function for the
population response. (C) Response to the preferred orientation across the three conditions for
all orientation-selective neurons. For presentation the time-courses are smoothed with a
Gaussian with a 33.3 ms kernel. Every row represents the response of one neuron. In each
panel, neurons are sorted based on their evoked response in the Unexpected condition (most

*excited on the top). **(D)** Comparison of the response in the Unexpected and Random*
conditions at the preferred orientation. Each dot represents one neuron. Purple dots show
neurons significantly modulated by expectation (N=133); grey dots are non-modulated neurons
*(N = 329). **(E)** Time-course of orientation-selectivity (circular mean) for the Random (blue) and*
Unexpected (green) conditions. Black horizontal lines indicate timepoints with statistically
significant difference between conditions, determined using non-parametric cluster-corrected
*procedures (see Methods). **(F)** Summary statistics for fitted Gaussian parameters across the*
population for the different sequence types. The Gain is the amplitude of the Gaussian and the
*Width is the standard deviation. * indicates p < 0.05. Across all panels error bars and shading*
represent ± 1 standard error of mean.

The main effect of predictability is evident from the three example neurons illustrated in

Figure 2A. There was a systematic increase in neuronal responses to the preferred orientation,

and a decrease to the anti-preferred orientation, in the *Unexpected* (green trace) compared

with the *Random* condition (blue trace). This response profile is consistent with a positive gain

modulation for unexpected gratings. The overall population response (aligned to the preferred

orientation) showed the same pattern of results (Figure 2B), with an increased response to the

preferred stimulus in the *Unexpected* versus *Random* condition. The responses of 133/462

orientation-selective neurons (28.8%) were significantly modulated in the *Unexpected*

condition relative to the *Random* condition (t-test, $p < 0.05$). Of these, all but two (98.5%)

showed a larger response in the *Unexpected* condition (Figure 2D), and this increase in

selectivity emerged shortly after stimulus presentation (Figure 2E).

We next quantified how orientation selectivity was affected by predictability. To do this,

we fitted circular Gaussian tuning functions to separately determine the gain (amplitude) and

width (standard deviation) parameters of orientation selectivity for each neuron (Figure 2F, see

Equation 1). The gain of the tuning curve was significantly greater in the *Unexpected* condition

than in the *Random* condition ($t(961) = 34.01$, $p < 0.0001$). By contrast, there was no

difference in the width between these two conditions, (t(961) = 0.45, p = 0.65). These results

are consistent with our recent work examining how prediction affects orientation selectivity

measured non-invasively in humans[14,15]. A control condition showed these effects were not

due to the systematic rotations that followed *Unexpected* gratings (Supplementary Figure 1).

Prediction affects population coding of orientation

In our initial set of analyses, we found that expectation affected orientation selectivity in

individual V1 neurons. We next examined how enhanced orientation selectivity for unexpected

stimuli at the single-neuron level in turn shaped the information contained within the population

response. Previous human neuroimaging studies using multivariate pattern analysis have

shown that expectation affects classification accuracy of the stimulus features[14–17,26]. To

determine how these findings generalise across species, we applied a similar multivariate

pattern analysis to the neuronal population data. We used all imaged neurons (N = 1697; 23

imaging sessions), including both orientation-selective and non-orientation selective neurons to

decode the presented orientation using inverted/forward encoding modelling (see multivariate

analysis section in Methods for details). Figure 3A and B illustrate the key steps in a forward

(or inverted) encoding approach and how this method can be used to determine the amount of

orientation-selective information contained in the population activity on a trial-to-trial basis. In

line with the human work[14–17,26], in a first step the method applies an encoding model using a

subset of trials (training trials) to estimate neuronal selectivity to each orientation (Figure 3A).

Then, in a second step, it inverts these weights to reconstruct the stimulus representation from

the population response on a new set of test trials (Figure 3B).

We first applied this decoding procedure in a time-resolved manner to determine the

temporal dynamics of population-level prediction effects (Figure 3D). This showed the

decoding performance started to rise for the *Random* and *Unexpected* conditions shortly after

stimulus presentation. More importantly, greater decoding accuracy emerges for *Unexpected*

relative to *Random* stimuli from shortly after stimulus onset (~100 ms). The pre-stimulus

divergence suggests that the increase in selectivity for unexpected stimuli results from

expectations developed *before* the stimulus appears rather than from a subsequent top-down

influence. Unsurprisingly, in the Expected condition orientation information could be decoded

above chance before the stimulus appeared. This is because orientations occurring before

stimulus presentation (0 ms) were correlated with the orientation of the decoded stimulus

presented at time zero. The decoding profile for Expected stimuli also exhibits an oscillating

profile, which likely reflects a combination of three factors: oscillations in neuronal activity due

to the periodic onsets of stimuli in the presented sequences; the 30° changes in orientation

from one stimulus to the next within the rotating sequences; and the dynamics of the calcium

indicator.

Figure 3. *Expectation affects stimulus-specific information carried by neuronal population activity. (A) Schematic of training the multivariate forward orientation encoding. Example regressors for 7 training trials with different orientations. The basis functions (grey lines) in response to different orientations which produce the regressor weights. Neuronal response for 4 example neurons for the example trials. Least squared regression is applied between the regressors and response to determine selectivity. Regression coefficients (beta weights) for four example neurons for each of the regressors found from a training set of data. (B) Testing the encoding model. Activity for the four neurons in test trials. Inverting the regressor weights and multiplying them by the population responses from the four neurons produces the*

predicted orientation response from this pattern of activity. The difference between the
*predicted and presented orientation for a given stimulus is the orientation error. **(C)** Distribution*
of orientation error when encoding was performed separately on groups of 50 neurons and 500
neurons at a time (with 24 permutations of different neuronal combinations). The vector
average of these histograms was taken as the decoding accuracy for each condition. The
*coloured numbers show the vector sum for the corresponding curves. **(D)** Time-resolved*
classification from forward encoding modelling (N= 500 neurons) with 24 permutations of
*different groups of neurons. **(E)** Decoding accuracy scales with the number of neurons. The*
classifier was trained and tested on the average response from 250 to 1000 ms following
stimulus onset, with different numbers of neurons included (24 permutations of different
neurons for each population size). The coloured horizontal lines indicate statistical significance
using sign-flipped cluster permutation tests comparing Random vs. Unexpected (green line)
and Random vs. Expected (red line). In panels B to D, shading/error bars indicate ±1 standard
error of the mean across permutations.

We next examined the effect of different sized neuronal populations on decoding

accuracy (Figure 3E). To do this, we selected groups of neurons and used a 10-fold cross-

validation procedure to train and test the classifier at each time point around stimulus

presentation. This procedure was repeated 24 times with different subsets of neurons

selected. The decoding procedure was performed on the average neuronal responses from

250 to 1000 ms after stimulus onset, and different sized pools of neurons were selected (1 to

1600 neurons, in 100 logarithmically-spaced steps). This analysis again showed that the

presented orientation was decoded significantly better in the *Unexpected* than the *Random*

condition. Figure 3E illustrates that this effect emerged with population sizes of relatively few

neurons (<10). The *Expected condition* also showed greater decoding accuracy relative to the

Random condition, but this effect was smaller than in the *Unexpected* condition and did not

emerge until a population of ~100 neurons was included in the analysis.

Predictions repel perception away from the expected orientation

The analyses presented above reveal a higher gain in orientation selectivity among V1

neurons following *Unexpected* grating stimuli relative to otherwise identical gratings within

Random sequences. According to formal models of predictive coding, the magnitude of a

prediction error should be determined by the degree of surprise, with more surprising stimuli

yielding larger responses[5,6]. Consistent with these models, we have shown in human

observers that orientation-selective stimulus- evoked responses get larger as the difference

between expected and presented stimuli increases[15].

In the current study we were able to quantify the degree of prediction error in the

Rotating condition and use this index to characterise any change in orientation-selective

responses in individual V1 neurons. To do this, we grouped orientation-selective neurons (N =

462) based on their maximum orientation-selective response in the *Random* condition (Figure

4A and Supplementary Figure 2). We found that orientation selectivity was influenced by the

expected orientation, such that responses were smallest when the expected orientation was

closest to the preferred orientation. For example, as shown in Figure 4A, neurons tuned to 90°

had the lowest orientation tuning when a 90° grating was expected (darkest green line).

Orientation selectivity was reduced to a lesser degree when the surrounding orientations (60°

and 120°) were expected, suggesting that the magnitude of the prediction error affected

neuronal responses in an orientation-selective manner.

To better visualise these effects, we aligned all neurons to their preferred orientation

and replotted the data as a function of the difference between the expected orientation and the

preferred orientation (Figure 4B). To quantify these effects, we fit Gaussian curves to each

neuron's orientation selectivity for all expected orientations (Figure 4C and 4D). Both the gain

(Figure 4C, one-way ANOVA, $F(5,1835) = 3.31$, $p = 0.006$, $\eta^2 = 0.006$) and the response to the

anti-preferred orientation (Figure 4D, $F(5,1835) = 8.38$, $p < 0.001$, $\eta^2 = 0.022$) were

systematically affected by the magnitude of the violated expectation.

Figure 4. *Increase in neuronal responses to unexpected stimuli is determined by the magnitude of the prediction error.* **(A)** *Neurons tuned to each displayed orientation are affected differently when different orientations are expected. Panel A shows an example for each expected orientation using neurons selective for 90˚ gratings, as defined based on their responses in the Random condition (from 250 – 1000 ms). Responses of remaining neurons selective for the other presented orientations are shown in Supplementary Figure 2. For each unexpected stimulus in the rotating condition, we identified the difference between the orientation of the expected stimulus and the orientation of the presented unexpected stimulus. For instance, if 60° was expected but 0° was unexpectedly presented, the expectation violation would be 60°.* **(B)** *All orientation-selective neurons aligned with their preferred orientation, plotted as separate Gaussians for each difference between the expected orientation and the presented orientation (expectation violation).* **(C)** *Gain of Gaussians fitted to each neuron's response, plotted as a function of expectation violation for all orientation-selective neurons.* **(D)**

*Baseline of Gaussians fitted to each neuron's response. **(E)** Forward encoding modelling*

reveals how population representations of orientation are affected by the degree of expectation

violation. The encoding weights are shown separately here for different values of expectation

*violation. **(F)** The y axis shows the difference between the presented and decoded orientation*

(ΔPerceived orientation). The population response (filled symbols) is biased away from the

expected orientation with the largest bias at ±30°. In all panels, error bars indicate ±1 standard

error of the mean across permutations. * indicates p < 0.05.

We followed up these results by examining how population-level encoding of the

presented orientation was affected by the magnitude of the prediction error (or expectation

violation). To do this, we divided the forward encoding results (Figure 3) into separate bins

based on the difference between the expected and presented orientation (Figure 4E). We

found that the decoded orientation of the presented stimulus was biased away from the

expected orientation, with the largest effect observed for a difference of 30°. This "repulsion"

effect is reminiscent of the well-known adaptation aftereffect for oriented stimuli[27,28], in which

the largest effect typically arises when the adapting and test stimuli are separated by around

30°. In the present experiment, however, the observed repulsion effect was driven by the

expected orientation rather than the orientation of the preceding stimulus.

Computational modelling of the relative contributions of adaptation and prediction on

visual coding efficiency

Formal models of predictive coding assume that high-level cortical areas pass

predictions, which are inverse copies of the expected stimulus, to lower-level areas[5,6].

According to this framework, only a small neuronal response is required for representation if a

stimulus matches the expectation[29]. Such an account is reminiscent of the effect of adaptation

on neuronal representation, whereby an immediately preceding stimulus reduces the neuronal

response to a current stimulus without decreasing the overall amount of stimulus

information[1,8]. Indeed, a number of studies have investigated whether adaptation might be due

to prediction errors[14,30,31]. Both adaptation and prediction rely on the statistics of sensory

inputs. Adaptation exploits the recent history of stimulus presentations to alter *current* sensory

representations, whereas prediction is thought to use statistical regularities to extract *future*

patterns.

We created a simple computational model of orientation processing to better

understand how expectation interacts with adaptation to influence the neural coding of

orientation. The model is based on several tuned orientation-selective neurons (or information

channels) maximally sensitive to different orientations. The neurons respond proportionally

based on their sensitivity to the presented orientation (Figure 5). We incorporated two sources

of inhibition: adaptation (in response to a previously-presented stimulus) and expectation (in

response to a predicted future stimulus). Similar to previous work[27,28,32,33], adaptation causes

gain modulation in neuronal orientation selectivity based on the response to the preceding

stimulus (Figure 5A,B). Prediction, on the other hand, affects neuronal responses by producing

an inverse copy of the expected orientation. To account for commonly observed long-lasting

effects of gain modulation on orientation sensitivity[34,35], the model allows sensitivity to recover

gradually over a number of trials. The amount of gain modulation can be varied to increase or

decrease the influence of either adaptation or prediction.

Figure 5. *Computational model for explaining variance in neuronal response by incorporating gain modulation from prediction and adaptation effects. The model consists of a bank of six neurons maximally selective for different orientations. The model's sensitivity is affected by previous orientations in the sequence (Adaptation) as well as future predicted orientations (Expectation). These factors determine the response to the presented orientation on each trial. (A) an example sequence of trials in the rotating condition. The orientations of the preceding (mustard) and expected (pink) trials determine the adaptation and the expectation gains. (B) The adaptation gain (mustard line) is determined by the orientation of the previous stimuli. The expectation gain (pink line) is determined by the inverse copy of the response to the expected*

orientation. **(C)** *Collectively, the two gains modulate the sensitivity of the channels on the next*
trial. These weights for the different orientations are applied to the model's sensitivity channels
(black lines), which give the response (orange line) to the presented orientation (vertical
dashed line; in this case 0°). **(D)** *Dots indicate the responses of the channels, and the curves*
are fitted Gaussian functions. Fitted Gaussian values to the model's responses for the different
stimulus conditions showing gain **(E)** *and width* **(F)** *of the response.* **(G)** *An example test of*
which model parameters best match the neuronal response in mouse V1 neurons. Regressors
for two different expectation gains (0.25 and 0.75) lead to slightly different weights for 10
example trials. Warmer colours indicate higher values. These are regressed against the
response (dF/F%) of each neuron. This yields beta values for each orientation channel
(regressors) for the two different expectation gains. **(I)** *Ridge regression results when the*
model was used to predict response to the stimulus in the Expected sequence, with different
levels of modulation from adaptation and prediction. The regressor (orientation) with the
highest beta weight was chosen for each neuron (N= 226; modulated by prediction). Error
*bars indicate ±1 standard error of the mean. * indicates p < 0.05.*
366
We presented sequences of orientations to the model from both the *Random* and

Rotating conditions to determine whether it can explain the observed changes in orientation

selectivity at the single-trial level. Because there are two sources of gain (adaptation and

expectation), we assume an equilibrium of gain modulation is available to the system to allow it

to maintain population homeostasis[36]. To this end, in the initial model we implemented 0.5 a.u.

of gain available, which was varied in the two stimulus conditions. In the *Random* condition,

the adaptation gain was set to 0.5 arbitrary units (a.u.) and the expectation gain was set to 0

a.u because the stimulus sequence was completely unpredictable. In the *Rotating* condition,

by contrast, the gain for both expectation and adaptation were set to 0.25 a.u. We re-aligned

neurons (Figure 5D) to their preferred orientation and determined their response to stimuli

under different conditions by fitting the same Gaussian to the results (Figure 5E and F).

Consistent with the neuronal data (Figure 2), in the model the gain of orientation selectivity

increased in the *Unexpected* condition ($M = 0.64$, $SD = 0.05$) relative to the *Expected* ($M =$

0.59, $SD = 0.03$) and *Random* ($M = 0.55$, $SD = 0.02$) conditions. The *Unexpected* trials

resulted in greater orientation selectivity than the *Expected* trials because sensitivity to the

stimulus was reduced for a different orientation (the predicted grating orientation) than the one

that was presented (Figure 5E). As with the experimental data, the width of tuning was similar

for the *Unexpected* (M = 29.8, SD = 0.62) and *Random* (M = 30.06, SD = 0.59) conditions,

whereas the Expected condition was slightly wider (M = 32.18, SD = 0.79, Figure 5F). The

model produced a qualitative fit consistent with the effects of expectation on V1 orientation

selectivity. The modulation of stimulus selectivity is consistent with previous work which found

that uncommon stimuli result in increased stimulus-specific adaptation in auditory cortex[37], and

that the V1 population response adapts to high-level stimulus statistics in a homeostatic

manner[36].

We next determined whether the model provided a quantitative fit to the recorded

neuronal activity. To do this, we used the model to generate predictions about neuronal

responses, which we regressed against the actual data for each neuron. Specifically, for each

experimental session for the awake mice, we presented the model with the same orientation

sequence viewed by the mouse, which in turn generated a predicted response for each

simulated neuron on every trial. We used ridge regression to determine beta weights for each

of the six regressors (orientations) for the three different gain settings for each neuron.

We found that a greater proportion of the variance in the trial-to-trial activity of neurons

could be explained when the model incorporated inhibition from expectation (Figure 5I). We

presented the orientation sequences from the *Rotating* condition to the model with three

different gain responses for expectation. With no gain, only the presented stimulus determined

the response of the model. As gain was increased from 0.25 and 0.75 a.u., greater inhibition

from expectation increased the model's fit with the data (Figure 5I).

For the adaptation model, there was no significant increase in its ability to explain

neuronal activity with increasing gain (one-way ANOVA, $F(4,900) = 0.52$, $p = 0.72$). By

contrast, the explanatory power of the expectation-only model greatly increased with

increasing levels of gain (one-way ANOVA, $F(4,900) = 6.18$, $p < 0.001$). Furthermore, the

model that incorporated a moderate amount of adaptation (0.25) with varying degrees of

expectation gain best predicted the neuronal response. A 3 (Model type; Adaptation,

Expectation, Combined model) × 5 (Gain level; 0,0.2,0.4,0.6,0.8,1.0) repeated-measures

ANOVA confirmed this observation, revealing that both the type of model ($F(2,450) = 4.22$, $p =$

0.02), and the gain level ($F(4,900) = 11.55$, $p < 0.001$) significantly affected the proportion of

variance explained. These factors significantly interacted ($F(8,1800) = 2.03$, $p = 0.04$), showing

that the difference in explanatory power between the models increased with increasing gain.

Follow-up tests showed that the expectation model did not explain significantly more variance

than the adaptation model ($p = 0.13$, across all gain levels), but the combined model did ($p =$

0.004).

Predictive coding under anaesthesia

Finally, we asked whether global anaesthesia altered the influence of prediction on

orientation selectivity observed in awake animals. Previous work in humans on expectation

violations has reported larger neural responses to unexpected than to expected stimuli during

sleep[38,39], in different attention states[15,40], and when individuals were in a coma[41], vegetative

state[42–44] or under anaesthesia[45]. These findings suggest that the influence of prediction errors

on patterns of brain activity varies across different global brain states and levels of

consciousness. To address this issue at the level of individual V1 neurons, we conducted a

further experiment in which the stimulus sequences (Random versus Rotating) were displayed

to mice under urethane anaesthesia (N=3 animals). For each mouse, we ran the full stimulus

protocol with 2-4 different areas in V1 (11 in total, 576 neurons). We found 96/576 (16.6%)

neurons were orientation selective. As shown in Figure 6AB, the gain of orientation selectivity

was again significantly enhanced in the *Unexpected* relative to the *Random* condition (Figure

6C, t(95) = 5.64, p < 0.0001). As in awake animals, there was a small but non-significant

decrease in the width of the tuning curve in the *Unexpected* condition relative to the *Random*

condition (Figure 6D, $t(95) = 0.39$, $p = 0.70$).

Figure 6. *Expectations affect the gain of orientation-selective V1 neurons under anaesthesia.*
(A) Time courses for all orientation-selective neurons (N = 96) aligned to their preferred
orientation to allow averaging. Shading indicates ±1 standard error of the mean across
neurons. (B) Population orientation tuning for the three expectation conditions, averaged
across an epoch from 250 to 1,000 ms after stimulus presentation. Solid lines are fitted
Gaussian functions with a constant offset. (C) Summary statistics for the gain of the fitted
Gaussians in B. (D) Width of the fitted Gaussians in B. (E) Comparison of the "surprise" effect
(Unexpected events minus Random events) between awake and anaesthetised animals. (F)
Time course of responses to the preferred orientation of each neuron, shown separately for the
three conditions. Neurons have been sorted by their responses in the Unexpected condition.
Across panels A-E shading and error bars indicate ±1 standard error of the mean across
neurons.

Finally, for each neuron we calculated the "surprise" effect by subtracting the gain of the

Gaussian tuning curve for the *Unexpected* condition from that of the Random condition (Figure

6E). A value larger than 0 indicates that the neuron's orientation selectivity was enhanced in

the *Unexpected* condition. There was no significant difference in the magnitude of the surprise

effect in awake animals compared with those that had been anaesthetised ($t(556)=1.38$, p =

0.17), suggesting that the influence of prediction errors on orientation-selective responses in

V1 neurons is equivalent for awake and anaesthetised animals.

Discussion

Here we provided an experimental test of how neuronal representations of visual

information are affected by prediction in the primary visual cortex (V1). Awake mice viewed

streams of oriented grating stimuli in either a *Random* condition, in which there was no

correlation between the orientations of successive stimuli (*Random* stimuli), or in a *Rotating*

condition, in which grating orientations were mostly predictable from previous events within the

sequence (*Expected* stimuli), but in which an occasional random orientation appeared

unexpectedly (*Unexpected* stimuli). Expectations reliably modulated the gain of orientation

selectivity in V1 activity, both at the level of single neurons and across the population of

recorded neurons. We found that neurons tuned to an expected orientation showed a large

decrease in their response compared with those not tuned to the expected orientation. The

expectation violation response was also reliably present under anaesthesia, suggesting that

the relevant visual circuits utilise predictive patterns in the sensory input even when the animal

is unconscious. Finally, we provided a computational implementation of a predictive coding

model in V1 to better understand the interaction between adaptation and prediction. By varying

the parameters of the model, we found that the best explanation for the observed neuronal

activity relied on both inhibition from adaptation in response to immediately preceding stimulus

events, and expectations about future stimulus features.

Our model of expectation-violation responses is phenomenological, in the sense that it

describes our results in a way that is grounded in neuronal and synaptic mechanisms. This

model contrasts with more formal accounts based upon hierarchal predictive coding.

Generally, neuronal responses to violations of expectations are formulated as precision-

weighted prediction errors[45–52]. In other words, neuronal responses reflect the difference

between sensory afferents and top-down predictions that are modulated or weighted by

precision. Precision, in this context, is a prediction of predictability, as opposed to prediction of

the sensory input. In the context of precision-weighted prediction errors, we can associate

adaptation gain with the effects of predictability (i.e., precision weighting) and excitation gain

with the prediction error per se; namely, the disinhibition of stimulus-bound responses by

absent top-down predictions. This fits with predictive coding accounts of the mismatch

negativity, where the equivalent effects are sometimes discussed in terms of stimulus-specific

adaptation and a sensory memory component — which in turn correspond to predictions of

precision and stimuli, respectively[14,15].

While the notion that predictions about the future affect perception was first proposed by

Helmholtz[54], direct evidence in support of this idea at the level of individual sensory neurons

has been lacking. A number of more recent theoretical models[5,6] have proposed a 'predictive

coding' framework with the common idea that the brain inhibits sensory representations of

expected stimuli to increase coding efficiency. Although there is good evidence that predictions

affect the magnitude of neural responses measured with non-invasive, whole-brain

neuroimaging methods[10,55], few studies have addressed whether individual neuronal

responses are affected, even though this is a critical component of predictive coding models.

The current results obtained from mouse V1 neurons fit well with our previous findings in

humans, which suggest that orientation selectivity changes with expectation[14–16,26].

Specifically, and in line with the current work, forward encoding modelling of EEG activity

revealed an increase in the gain, but not the width, of orientation tuning to unexpected stimuli

in human observers[14].

Our results add to the understanding of how expectations affect the representation of

sensory information. Previous work[19,22,23] has suggested that locomotion-induced increases in

activity in primary visual cortex in mice relates to predictive coding[56,57] (but see[57] for a different

interpretation of these findings). Under the predictive coding framework, the increased activity

caused by locomotion creates an expectation that the stimulus should move and change size.

A prediction error is generated if the stimulus remains static, as is typical when measuring

orientation selectivity, or moves in an inconsistent direction. There is significantly less

locomotion-induced increase in response if the stimulus is made to move as the animal

moves[23]. Our results are consistent with these findings, but also identify an enhanced gain

mechanism reflected by a larger response to the neuron's preferred orientation.

In the human literature, expectation appears to affect sensory responses through

different neural oscillatory frequency bands[58,59]. Recordings in macaques suggest visual

information is fed forward through high-frequency gamma (60-80 Hz) oscillations, while

feedback occurs through slow theta-band (14-18 Hz) activity[60]. As the present recordings were

conducted using two-photon imaging with a relatively slow sampling rate, we are at present

unable to determine the possible role of different oscillation frequencies in the observed

expectation effects. Future work in which activity is recorded from multiple sites concurrently

using electrophysiology could help characterise the distinct contributions of top-down and

bottom-up neural signals to this expectation-induced gain modulation.

Methods

Mouse information

A total of 6 wild type mice (C57BL) were used for this study. All methods were

performed in accordance with the protocol approved by the Animal Experimentation and Ethics

Committee of the Australian National University (AEEC 2015/74; 2019/11). Mice were housed

in a ventilated and air filtered climate-controlled environment with a 12-hour light–dark (8 am

lights on, 8 pm lights off) cycle. Mice had access to food and water ad libitum. No statistical

methods were used to calculate the sample size, but these were consistent with other studies

in the field.

Expression of Ca^{2+} indicator GCaMP6f

Mice were briefly anaesthetised with isoflurane (~2% by volume in O_2) in a chamber and

moved to a thermal blanket (37°C, Physitemp Instruments) before the head was secured in a

stereotaxic frame (Stoelting, IL). Thereafter, the anaesthetic gas (isoflurane, ~2% by volume in

O_2) was passively applied through the nose mask at a flow rate of 0.6-0.8 L/min. The level of

anaesthesia was monitored by the respiratory rate, and hind paw and corneal reflexes. The

eyes were covered with a thin layer of Viscotears liquid gel (Alcon, UK). The scalp was opened

with ~5 mm rostrocaudal incision at the midline using scissors and the periosteum was gently

removed. A circular craniotomy was made over the right visual cortex (3 mm diameter; centred

2 mm lateral and 4.5 mm posterior to Bregma) with the dura left intact. A glass pipette (15-25

μm diameter at tip) containing GCaMP6f (AAV1.Syn.GCaMP6f.WPRE.SV40, Penn Vector

Core, The University of Pennsylvania, USA) was inserted into the cortex at a depth of 230-250

μm below the dura using a micromanipulator (MPC-200, Sutter Instruments, Novato, CA,

USA). GCaMP6f was injected at 4-6 sites (with four 32-nL injections per site separated by 2-5

mins; rate 92 nLs^{-1}). V1 was localised anatomically using coordinates established using

functional methods[60]. Injections were centred 2mm lateral and 4.5mm posterior to Bregma.

Injections were controlled using a Nanoject II injector (Drumont scientific, PA). After virus

injection, the craniotomy was covered with a 3mm diameter cover-glass (0.1 mm thickness,

Warner Instruments, CT). This was glued to the bone surrounding the craniotomy. Custom

made head bars were fixed to the skull over Bregma using a thin layer of cyanoacrylate

adhesive and dental acrylic. A small well was built surrounding the craniotomy window using

dental acrylic to accommodate distilled water required for the immersion lens of the 2-photon

microscope.

Ca^{2+} imaging was performed using a two-photon microscope (Thorlabs Inc., Newton,

NJ, USA) controlled by ThorImage OCT software. The visual cortex was illuminated with a

Ti:Sapphire fs-pulsed laser (Chameleon, Coherent Inc., Santa Clara, CA, USA) tuned at 920

nm. The laser was focused onto L2/3 cortex through a 16x water-immersion objective lens

(0.8NA, Nikon), and Ca^{2+} transients were obtained from neuronal populations at a resolution of

512 × 512 pixels (sampling rate, ~30 Hz). To abolish the effect of visual stimuli on the calcium

signals, we filled the gap between the objective and the well with removable adhesive (Blu-

Tack).

The obtained images were processed using the Suite2p toolbox

(https://github.com/cortex-lab/Suite2P) for motion correction and segmentation. The

surrounding neuropil signal was subtracted for each neuron's calcium traces. These corrected

traces were high-pass filtered before the median response for each neuron was subtracted to

determine dF/F.

Visual stimulus

The stimuli were displayed on a 22-inch LED monitor (resolution 1920 x 1080 pixels,

refresh rate 60 Hz) using the PsychToolbox presentation software for MATLAB[62,63]. The

mouse was placed next to the monitor, which subtended 76.8° x 43.2° (one pixel = 2.4' x 2.4')

orientated 30° from their midline. The visual stimulus sequence was based on the Allen Brain

Institute Brain Observatory paradigm used to measure orientation selectivity in mice. The

stimuli were full-screen gratings (0.034 c/°, 50% contrast) displayed for 250 ms with no inter-

stimulus blank interval giving a 4 Hz presentation rate. The spatial frequency was chosen to be

close to optimal sensitivity of neurons in V1[25]. The orientations of the gratings were equally

spaced between 0 to 150° in 30° steps so we could characterise each neuron's orientation

selective profile.

The predictability of the orientations of the gratings was varied in the two stimulus

conditions. In the *Random* condition, the orientations of the gratings were drawn from a

pseudo-random distribution with no relationship between the current orientation and the

previous orientation. In the *Rotating* condition, the orientations of the gratings rotated (in 30°

steps) either clockwise or anti-clockwise for 5 to 9 presentations, before jumping to an

unexpected random orientation, where it began rotating in the opposite direction. The *Random*

and *Rotating* conditions were presented in blocks of trials which were pseudorandomised in

time within each imaging session.

In 3 mice, we ran a total of 23 imaging sessions and collected data from 1697 neurons.

Neurons from all sessions and mice were pooled for analysis. One session (1.5 – 2 hours) was

recorded in a day from each mouse. These sessions occurred between 1 and 4 times per

week. In each session, two runs of *Rotating* and *Random* sequences were presented, and

each of these contained 1800 trials, alternating between *Rotating* and *Random*. The order of

sequences was counter-balanced across mice. For some sessions for 2 of the mice, we also

presented a *Rotating control* condition to determine whether the systematic rotational

movement after the unexpected jump affected orientation selectivity. In this condition, after the

unexpected orientation the stimulus made another jump to a random orientation before starting

to rotate in the opposite direction as the previous rotation. The number of events was

increased from 7200 (3600 x 2) in each condition to 8400 to have the same number of

unexpected trials as the original *Rotating* condition, while all other details remained identical

with the *Rotating* condition. We ran 13 sessions in these 2 mice for all three conditions to

compare the effect of the control. For all conditions, there was a balanced number of

presentations of all the orientations.

Data analysis

To determine the effect of predictability, we averaged the calcium response (dF/F%)

from 250 to 1000 ms after stimulus presentation to derive tuning curves for each condition. To

quantify how expectation affected the gain and selectivity of orientation-selective neurons we

fitted circular Gaussian distributions with a constant offset (Equation 1) using non-linear least

square regression.

605
$$G(x) = A exp - \frac{(x - \phi - j * 180)^2}{2\sigma^2} + C \qquad (1)$$

Where A is the gain (amplitude) of the Gaussian, ϕ is the preferred orientation of the neuron

(in degrees), σ is the width (in degrees) and C a constant offset to allow for baseline shifts in

the activity of the neuron. We searched for best fitting solutions with parameter j, with a search

space from -4 to +4 in integer steps.

To provide another test of how prediction affects orientation selectivity of individual

neurons, we found the circular mean[64] of the averaged orientation tuning curve across all

presentations within the condition (Figure 2E). This was done for each time point (1/sample

rate) between -500 and 2000 ms around stimulus presentation.

Multivariate encoding analysis

We used a multivariate encoding approach (forward encoding modelling) to determine

how the population activity carried information about the orientation of the presented grating on

a trial-to-trial basis. This is adapted from human neuroimaging approaches, which examine

orientation/feature selectivity from multivariate non-invasively recorded neural activity[14,15,65–68],

but is similar to encoding approaches used to describe neuronal response to sensory

stimuli[69,70]. Compared to the encoding-only, forward encoding takes the individual neuron

activity to reconstruct the stimulus representation from the population activity. The technique

goes beyond more commonly used multivariate pattern analysis procedures by producing

tuning curves showing the full representation (in both gain, width, and bias) relative to the

accuracy-only score.

The data were pooled across all experimental sessions with both orientation and non-

orientation selective neurons used. In the first instance, we examined how the number of

neurons affected decoding on a fixed time interval (250 to 1000 ms) and in the second

instance, we found the time-resolved selectivity by applying the decoding procedure at each

time point around the presentation of the stimulus (-500 to 2000 ms). A 20-fold cross-validation

procedure was used in both instances for test and training data. The procedure evenly splits

each test block to have the most even distribution of stimuli in each fold.

We used the presented orientations to construct a regression matrix with 8 regression

coefficients. This regression matrix was convolved with a tuned set of nine basis functions (half

cosine functions raised to the eighth power) centred from 0° to 160° in 20° steps. This helps

pool similar orientations and reduces overfitting[70]. This tuned regression matrix was used to

measure orientation information across trials. This was done by solving the linear Equation 2:

$$B_1 = WC_1 \hspace{4cm} (2)$$

Where B_1 (Neurons x N training trials) is the data for the training set, C_1 (8 channels x N

training trials) is the tuned channel response across the training trials, and W is the weight

matrix for the sensors to be estimated (Neurons x 8 channels). We separately estimated the

weights associated with each channel individually. W was estimated using least square

regression to solve Equation 3:

$$W = (C_1 \, C_1^T)^{-1} \, C_1^T \, B_1 \hspace{3cm} (3)$$

We removed the correlations between neurons, as these add noise to the linear equation. To

do this, we first estimated the noise correlation between neurons (which stops finding the true

solution to the equation) and removed this component through regularisation by dividing the

weights by the shrinkage matrix[68,71]. The channel response in the test set C_2 (8 channels x N

test trials) was estimated using the weights in (4) and applied to activity in B_2 (Neurons x N test

trials), as per Equation 4:

$$C_2 = (W\ W^T)\ W^T\ B_2 \tag{4}$$

To avoid overfitting, we used 10-fold cross validation, where X-1 epochs were used to

train the model, and this was then tested on the remaining (X) epoch. This process was

repeated until all epochs had served as both test and training trials. We also repeated this

procedure for each point in the epoch to determine time-resolved feature-selectivity. To re-

align the trials with the exact presented orientation, we reconstructed the item representation

by multiplying the channel weights (8 channels x time x trial) against the basis set (180

orientations × 8 channels). This resulted in an Orientations (-89° to 90°) × Trial × Time

reconstruction.

To quantify the orientation selective response, we found the vector sum of the

orientation for each trial (Figure 3) to determine the decoded orientation. The difference

between the decoded and presented orientation was the orientation error. For each condition

(and time point where applicable) we found the distribution of orientation errors and calculated

the histogram of responses.

In the temporal classification analysis, groups of 500 neurons were used in each

instance for both training and test data with the cross-validation procedure applied to each time

point around stimulus presentation. We permuted new groups of 500 neurons 24 times. Next,

we averaged the evoked activity from 250 to 1000 ms after stimulus presentation. To

determine how decoding was affected by population size, the same classification was then

used as in the previous analysis but with different numbers of neurons (2 to 1600 neurons in

100 logarithmically-spaced steps). Again, we selected different groups of neurons 24 times so

as not to skew the results by the neurons that were selected.

Statistics

Non-parametric signed permutation tests[71,72] were used to determine time resolved

differences between conditions. The sign of the data was randomly flipped (N = 5,000), with

equal probability, to create a null distribution. Cluster-based permutation testing was used to

correct for multiple comparisons over the timeseries, with a cluster-form threshold of $p<0.05$

and significance threshold of $p<0.05$.

Computational model

The analytic model is based on previous work accounting for feature (i.e. orientation,

spatial, color) adaptation based on neuronal response and human psychophysical

data[27,28,33,73,74]. The model consists of a bank of 6 orientation-selective information channels

with preferred orientations evenly spaced between 0 and 150° (in 30° steps). Each channel's

sensitivity profile is given by a Gaussian function (Equation 5).

$$G(x) \ = \ A exp - \frac{(x - \phi)^2}{2\sigma^2} \qquad (5)$$

Where A is the gain (amplitude) (set to 1 a.u.), ϕ is the channel's preferred orientation, σ is the

width of the channel (set to be 40° consistent with the neuronal data). The number of channels,

along with the width means the model is equally sensitive to all orientations. The population

response to any presented orientated stimulus is given by the sensitivity profiles of the

channels (See Supplementary Figure 3). In an unadapted state (Supplementary Figure 3A),

the model will show a maximal response around the presented orientation with the vector

average of the population response will be the presented orientation.

To account for adaptation, the gain of the information channels is reduced in inverse

proportion to their response by the previous stimulus (Supplementary Figure 3B). For instance,

if a 90° stimulus is the adapting stimulus, the sensitivity of the channels around 90° will be

maximally reduced while orthogonal channels will be unaffected. The magnitude of this

reduction (*adaptation ratio*) can be varied to allow for greater or less adaptation and was

included as a free parameter in the analysis. The adaptation aspect of the model is consistent

with previous models used to psychophysical data[27,28,33,73,74]. The new model accurately

predicts serial dependency effects (where the current orientation is biased away from the

previous orientation) seen in the neuronal data[34,35,75].

Prediction gain modulation works in a similar manner as adaptation except that the

stimulus sensitivity, rather than channel sensitivity, is modulated. Furthermore, the gain

modulation occurs before the stimulus and is for the orientation that is expected rather than

presented. The modulation of stimulus sensitivity is consistent with a previous study which

found that uncommon stimuli result in stimulus-specific adaptation in the auditory cortex[37].

Stimulus-specific adaptation has been used in modelling neuronal adaptation[36]. To model

stimulus-selective gain modulation, the tuned Gaussian function was found using Equation 1

and inverted before being applied to the channels. The amount of gain modulation by

expectation was a free parameter (*expectation gain*).

To account for long-lasting effects of gain modulation, the channel's sensitivity was

normalised by the maximum sensitivity of response on each trial. This causes the model to

have adaptation and expectation effects based on the presented orientation of at least four

stimuli back. How many n-back stimulus affect the current trials sensitivity is determined by the

modulation factor. We used this type of long-lasting gain to account for well-known effects

such as serial dependency-like which can occur with adaptation and prediction[34,35]. We

regressed the adaptation-only model against the neuronal data and found a factor of 3.0 best

fit the data which was set for other modelling experiments.

To determine the effects of the different stimulus conditions (Random, Expected and

Unexpected) on the model's channels, we presented sequences of orientations to the model

and split the responses into conditions. To allow for easier comparison, we aligned the six

orientation channels to their preferred orientation and collapsed the results across conditions.

The same effects were evident before collapsing across the channels.

Lastly, we examined how the actual neuronal responses could be predicted by the

model's predictions with different values of the free parameters. To do this, we used to model

to predict responses to the orientations presented to the mice during the session for all

stimulus conditions. For each neuron, we used the model's responses to the stimuli as

regressors to predict the neuron's response (averaged from 250 to 1000 ms) for each stimulus

condition. We iterated this procedure with different values for adaptation and expectation gain

to determine what values best predicted the data.

Data availability

The data are available at: https://osf.io/t2vb3. The code is available at:

https://github.com/MatthewFTang/PredictionOrientationSelectivityMouseV1

Acknowledgements

This work was supported by the Australian Research Council (ARC) Centre of

Excellence for Integrative Brain Function (CE140100007) to JBM and EA, a National Health

and Medical Research Council (NHMRC) Project Grant to MFT, JBM and EA (GNT1165337),

and an NHMRC Ideas Grant to EK, CL and EA (GNT1181643). MFT was supported by an

ARC DECRA fellowship (DE210100508) and the NVIDIA corporation who donated a TITAN V

GPU. CL was supported by an ARC DECRA fellowship (DE229199691). EA was supported by

an ARC Discovery Project (DP170100908). JBM was supported by an NHMRC Investigator

Grant (GNT2010141) and by the Canadian Institute for Advanced Research (CIFAR).

Author Contributions

MFT and EA conceived the experiments. EK and CL performed the experiments. MFT

and EA analysed the data. JED and MFT developed the model. MFT, EK, CYL, JED, JBM and

EA wrote the paper.

References

1.   Lundstrom, B. N. & Fairhall, A. L. Decoding stimulus variance from a distributional neural

748   code of interspike intervals. *J. Neurosci.* **26**, 9030–9037 (2006).

749   2.   Gross, C. G., Rocha-Miranda, C. E. & Bender, D. B. Visual properties of neurons in

750   inferotemporal cortex of the Macaque. *J. Neurophysiol.* **35**, 96–111 (1972).

751   3.   Treue, S. & Martínez Trujillo, J. C. Feature-based attention influences motion processing

752   gain in macaque visual cortex. *Nature* **399**, 575–579 (1999).

4.   Luck, S. J., Chelazzi, L., Hillyard, S. A. & Desimone, R. Neural mechanisms of spatial

selective attention in areas V1, V2, and V4 of macaque visual cortex. *J. Neurophysiol.*

77, 24–42 (1997).

5.   Friston, K. A theory of cortical responses. *Philos. Trans. R. Soc. B Biol. Sci.* **360**, 815–

836 (2005).

6.   Rao, R. P. N. & Ballard, D. H. Predictive coding in the visual cortex: A functional

interpretation of some extra-classical receptive-field effects. *Nat. Neurosci.* **2**, 79–87

(1999).

7.   Maravall, M., Petersen, R. S., Fairhall, A. L., Arabzadeh, E. & Diamond, M. E. Shifts in

coding properties and maintenance of information transmission during adaptation in

barrel cortex. *PLoS Biol.* **5**, 0323–0334 (2007).

8.   Adibi, M., Clifford, C. W. G. & Arabzadeh, E. Informational basis of sensory adaptation:

Entropy and single-spike efficiency in rat barrel cortex. *J. Neurosci.* **33**, 14921–14926

(2013).

9.   Bichot, N. P., Rossi, A. F. & Desimone, R. Parallel and serial neural mechanisms for

visual search in macaque area V4. *Science (80-.).* **308**, 529–534 (2005).

10.   Näätänen, R., Paavilainen, P., Rinne, T. & Alho, K. The mismatch negativity (MMN) in

basic research of central auditory processing: A review. *Clin. Neurophysiol.* **118**, 2544–

2590 (2007).

11.   Garrido, M. I., Kilner, J. M., Stephan, K. E. & Friston, K. J. The mismatch negativity: A

review of underlying mechanisms. *Clin. Neurophysiol.* **120**, 453–463 (2009).

12.   Näätänen, R., Gaillard, A. W. K. & Mäntysalo, S. Early selective-attention effect on

evoked potential reinterpreted. *Acta Psychol. (Amst).* **42**, 313–329 (1978).

13.   Näätänen, R. & Michie, P. T. Early selective-attention effects on the evoked potential: A

critical review and reinterpretation. *Biol. Psychol.* **8**, 81–136 (1979).

14.   Tang, M. F., Smout, C. A., Arabzadeh, E. & Mattingley, J. B. Prediction error and

repetition suppression have distinct effects on neural representations of visual

information. *Elife* **7**, (2018).

15.   Smout, C. A., Tang, M. F., Garrido, M. I. & Mattingley, J. B. Attention promotes the

neural encoding of prediction errors. *PLoS Biol.* **17**, 1–22 (2019).

16.   Kok, P., Mostert, P. & De Lange, F. P. Prior expectations induce prestimulus sensory

templates. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 10473–10478 (2017).

17.   King, J. R. *et al.* Single-trial decoding of auditory novelty responses facilitates the

detection of residual consciousness. *Neuroimage* **83**, 726–738 (2013).

18.   King, J. R., Gramfort, A., Schurger, A., Naccache, L. & Dehaene, S. Two distinct

dynamic modes subtend the detection of unexpected sounds. *PLoS One* **9**, e85791

(2014).

19.   Leinweber, M., Ward, D. R., Sobczak, J. M., Attinger, A. & Keller, G. B. A Sensorimotor

Circuit in Mouse Cortex for Visual Flow Predictions. *Neuron* **95**, 1420-1432.e5 (2017).

20.   Meyer, T. & Olson, C. R. Statistical learning of visual transitions in monkey

inferotemporal cortex. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 19401–19406 (2011).

21.   Schneider, D. M., Sundararajan, J. & Mooney, R. A cortical filter that learns to suppress

the acoustic consequences of movement. *Nature* **561**, 391–395 (2018).

22.   Zmarz, P. & Keller, G. B. Mismatch Receptive Fields in Mouse Visual Cortex. *Neuron* **92**,

766–772 (2016).

23.   Keller, G. B., Bonhoeffer, T. & Hübener, M. Sensorimotor Mismatch Signals in Primary

799   Visual Cortex of the Behaving Mouse. *Neuron* **74**, 809–815 (2012).

800   24.   Heindorf, M., Arber, S. & Keller, G. B. Mouse Motor Cortex Coordinates the Behavioral

801   Response to Unpredicted Sensory Feedback. *Neuron* **99**, 1040-1054.e5 (2018).

802   25.   de Vries, S. E. J. *et al.* A large-scale standardized physiological survey reveals functional

803   organization of the mouse visual cortex. *Nat. Neurosci.* **23**, 138–151 (2020).

804   26.   Kok, P., Jehee, J. F. M. & de Lange, F. P. Less Is More: Expectation Sharpens

805   Representations in the Primary Visual Cortex. *Neuron* **75**, 265–270 (2012).

806   27.   Tang, M. F., Dickinson, J. E., Visser, T. A. W. & Badcock, D. R. The broad orientation

807   dependence of the motion streak aftereffect reveals interactions between form and

808   motion neurons. *J. Vis.* **15**, (2015).

809   28.   Clifford, C. W. G. Perceptual adaptation: Motion parallels orientation. *Trends Cogn. Sci.*

810   **6**, 136–143 (2002).

811   29.   Cheadle, S., Egner, T., Wyart, V., Wu, C. & Summerfield, C. Feature expectation

812   heightens visual sensitivity during fine orientation discrimination. *J. Vis.* **15**, 14 (2015).

813   30.   Kovács, G., Kaiser, D., Kaliukhovich, D. A., Zoltán, V. & Vogels, R. Repetition probability

814   does not affect fMRI repetition suppression for objects. *J. Neurosci.* **33**, 9805–9812

815   (2013).

816   31.   Summerfield, C., Trittschuh, E. H., Monti, J. M., Mesulam, M. M. & Egner, T. Neural

817   repetition suppression reflects fulfilled perceptual expectations. *Nat. Neurosci.* **11**, 1004–

818   1006 (2008).

32. Dickinson, J. E., Harman, C., Tan, O., Almeida, R. A. & Badcock, D. R. Local contextual

interactions can result in global shape misperception. *J. Vis.* **12**, 3 (2012).

33. Westrick, Z. M., Heeger, D. J. & Landy, M. S. Pattern adaptation and normalization

reweighting. *J. Neurosci.* **36**, 9805–9816 (2016).

34. Fischer, J. & Whitney, D. Serial dependence in visual perception. *Nat. Neurosci.* **17**,

738–743 (2014).

35. Zavitz, E., Yu, H. H., Rowe, E. G., Rosa, M. G. P. & Price, N. S. C. Rapid adaptation

induces persistent biases in population codes for visual motion. *J. Neurosci.* **36**, 4579–

4590 (2016).

36. Benucci, A., Saleem, A. B. & Carandini, M. Adaptation maintains population homeostasis

in primary visual cortex. *Nat. Neurosci.* **16**, 724–729 (2013).

37. Ulanovsky, N., Las, L. & Nelken, I. Processing of low-probability sounds by cortical

neurons. *Nat. Neurosci.* **6**, 391–398 (2003).

38. Loewy, D. H., Campbell, K. B. & Bastien, C. The mismatch negativity to frequency

deviant stimuli during natural sleep. *Electroencephalogr. Clin. Neurophysiol.* **98**, 493–

501 (1996).

39. Atienza, M., L. Cantero, J. & Gómez, C. M. The mismatch negativity component reveals

the sensory memory during REM sleep in humans. *Neurosci. Lett.* **237**, 21–24 (1997).

40. Näätänen, R., Paavilainen, P., Titinen, H., Jiang, D. & Alho, K. Attention and mismatch

negativity. *Psychophysiology* vol. 30 436–450 (1993).

41. Bekinschtein, T. A. *et al.* Neural signature of the conscious processing of auditory

regularities. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 1672–1677 (2009).

42.   Perrin, F. *et al.* Brain response to one's own name in vegetative state, minimally

conscious state, and locked-in syndrome. *Arch. Neurol.* **63**, 562–569 (2006).

43.   Wijnen, V. J. M., van Boxtel, G. J. M., Eilander, H. J. & de Gelder, B. Mismatch negativity

predicts recovery from the vegetative state. *Clinical Neurophysiology* vol. 118 597–605

(2007).

44.   Melanie, B. *et al.* Preserved Feedforward But Impaired Top-Down Processes in the

Vegetative State. *Science (80-.).* **332**, 858–862 (2011).

45.   Heinke, W. *et al.* Sequential Effects of Increasing Propofol Sedation on Frontal and

Temporal Cortices as Indexed by Auditory Event-related Potentials. *Anesthesiology* **100**,

617–625 (2004).

46.   Palmer, C. E., Auksztulewicz, R., Ondobaka, S. & Kilner, J. M. Sensorimotor beta power

reflects the precision-weighting afforded to sensory prediction errors. *Neuroimage* **200**,

59–71 (2019).

47.   Sterzer, P. *et al.* The predictive coding account of psychosis. *Biol. Psychiatry* **84**, 634–

643 (2018).

48.   Haarsma, J. *et al.* Precision weighting of cortical unsigned prediction errors is mediated

by dopamine and benefits learning. *bioRxiv* 288936 (2018).

49.   Limanowski, J. (Dis-) attending to the Body: Action and Self-experience in the Active

Inference Framework. (2017).

50.   Shipp, S. Neural elements for predictive coding. *Front. Psychol.* **7**, 1792 (2016).

51.   Ainley, V., Apps, M. A. J., Fotopoulou, A. & Tsakiris, M. 'Bodily precision': a predictive

coding account of individual differences in interoceptive accuracy. *Philos. Trans. R. Soc.*

B Biol. Sci. **371**, 20160003 (2016).

52.   Auksztulewicz, R. & Friston, K. Attentional enhancement of auditory mismatch

responses: a DCM/MEG study. *Cereb. cortex* **25**, 4273–4283 (2015).

53.   Clark, A. Whatever next? Predictive brains, situated agents, and the future of cognitive

science. *Behav. Brain Sci.* **36**, 181–204 (2013).

54.   von Helmholtz, H. *Treatise on Physiological Optics Vol. III*. vol. 3 (Dover Publications,

1867).

55.   Garrido, M. I. *et al.* The functional anatomy of the MMN: A DCM study of the roving

paradigm. *Neuroimage* **42**, 936–944 (2008).

56.   Niell, C. M. & Stryker, M. P. Modulation of Visual Responses by Behavioral State in

Mouse Visual Cortex. *Neuron* **65**, 472–479 (2010).

57.   Ayaz, A., Saleem, A. B., Schölvinck, M. L. & Carandini, M. Locomotion controls spatial

integration in mouse visual cortex. *Curr. Biol.* **23**, 890–894 (2013).

58.   Summerfeld, C., Wyart, V., Johnen, V. M. & de Gardelle, V. Human scalp

electroencephalography reveals that repetiti suppression varies with expectation. *Front.*

Hum. Neurosci. **5**, 1–13 (2011).

59.   Todorovic, A., van Ede, F., Maris, E. & de Lange, F. P. Prior expectation mediates neural

adaptation to repeated sounds in the auditory cortex: An MEG study. *J. Neurosci.* **31**,

9118–9123 (2011).

60.   Bastos, A. M. *et al.* Canonical Microcircuits for Predictive Coding. *Neuron* **76**, 695–711

(2012).

61.   Zhuang, J. *et al.* An extended retinotopic map of mouse cortex. *Elife* **6**, 3 (2017).

62.   Brainard, D. H. & Vision, S. The psychophysics toolbox. *Spat. Vis.* **10**, 433–436 (1997).

63.   Pelli, D. G. & Vision, S. The VideoToolbox software for visual psychophysics:

887   Transforming numbers into movies. *Spat. Vis.* **10**, 437–442 (1997).

888   64.   Berens, P. CircStat: A MATLAB Toolbox for Circular Statistics . *J. Stat. Softw.* **31**,

889   (2009).

890   65.   Brouwer, G. J. & Heeger, D. J. Decoding and reconstructing color from responses in

891   human visual cortex. *J. Neurosci.* **29**, 13992–14003 (2009).

892   66.   Ester, E. F., Sprague, T. C. & Serences, J. T. Parietal and Frontal Cortex Encode

893   Stimulus-Specific Mnemonic Representations during Visual Working Memory. *Neuron*

894   **87**, 893–905 (2015).

895   67.   Sprague, T. C. & Serences, J. T. Attention modulates spatial priority maps in the human

896   occipital, parietal and frontal cortices. *Nat. Neurosci.* **16**, 1879–1887 (2013).

897   68.   Tang, M. F. *et al.* Neural dynamics of the attentional blink revealed by encoding

898   orientation selectivity during rapid visual presentation. *Nat. Commun.* **11**, 1–14 (2020).

899   69.   Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial

900   neural dynamics are dominated by richly varied movements. *Nat. Neurosci.* **22**, 1677–

901   1686 (2019).

902   70.   Park, I. M., Meister, M. L. R., Huk, A. C. & Pillow, J. W. Encoding and decoding in

903   parietal cortex during sensorimotor decision-making. *Nat. Neurosci.* **17**, 1395–1403

904   (2014).

905   71.   Wolff, M. J., Jochim, J., Akyürek, E. G. & Stokes, M. G. Dynamic hidden states

906   underlying working-memory-guided behavior. *Nat. Neurosci.* **20**, 864–871 (2017).

72.   Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J. M. FieldTrip: Open source software

for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput.*

Intell. Neurosci. **2011**, (2011).

73.   Dickinson, J. E., Mighall, H. K., Almeida, R. A., Bell, J. & Badcock, D. R. Rapidly

acquired shape and face aftereffects are retinotopic and local in origin. *Vision Res.* **65**,

1–11 (2012).

74.   Dickinson, J. E., Almeida, R. A., Bell, J. & Badcock, D. R. Global shape aftereffects have

a local substrate: A tilt aftereffect field. *J. Vis.* **10**, 1–12 (2010).

75.   Fritsche, M., Mostert, P. & de Lange, F. P. Opposite Effects of Recent History on

Perception and Decision. *Curr. Biol.* **27**, 590–595 (2017).

Supplementary figures

Supplementary Figure 1. *Control condition to determine whether the rotation sequence*
caused the increased gain in the Unexpected trials. ***((B).*** *Fitted gain values for each neuron*
for the three conditions. This subset of neurons showed the same effect in the original two
conditions, with an increase in gain in the Unexpected compared to Random condition (t(129 =
7.74, p < 0.001). This effect was maintained when comparing the Random to the Unexpected
control condition (t(129) = 7.81 p < 0.001). There was no significant difference between
Unexpected and Unexpected control conditions (t(129 = 1.81, p = 0.07).
929

Supplementary Figure 2. Expectations affect neurons differently depending on their preferred orientation. Each panel shows neurons tuned to different orientations, as defined by their stimulus-evoked responses in the Random condition. The different colour-coded curves show different expected orientations. Neurons show the largest decrease in response when their preferred orientation is similar to the expected orientation. Across all panels error bars indicate ±1 standard error of mean.

Supplementary Figure 3. *A simple schematic example of the model. **(A)** The model in an unadapted state, showing its response to a 110° stimulus (blueline). The green line shows the model's response to the stimulus in each channel. **(B)** Applying adaptation gain at 110° reduces the model's sensitivity to nearby, but not distant, stimuli. The model's response (red line) is reduced in magnitude relative to panel A when the same test stimuli are applied.*