NHR-23 activity is necessary for *C. elegans* developmental progression and apical extracellular matrix structure and function

Londen C. Johnson[1], An A. Vo[1], John C. Clancy[1], Krista M. Myles[1], Murugesan Pooranachithra[2], Joseph Aguilera[1], Max T. Levenson[1], Chloe Wohlenberg[1], Andreas Rechtsteiner[1], James Matthew Ragle[1], Andrew D. Chisholm[2], and Jordan D. Ward[1‡]

[1]Department of Molecular, Cell, and Developmental Biology, University of California-Santa Cruz, Santa Cruz, CA 95064, USA.

[2]Department of Cell and Developmental Biology, School of Biological Sciences, University of California San Diego, La Jolla, CA 92093, USA.

[‡]Author for correspondence.

Corresponding author email: jward2@ucsc.edu

Keywords: *C. elegans*, molting, NHR-23, nuclear hormone receptor, apical extracellular matrix, auxin-inducible degron

Running Title: NHR-23 in *C. elegans* molting

SUMMARY STATEMENT

This work shows how a *C. elegans* transcription factor controls remodeling of the apical extracellular matrix during development and in which tissues it acts.

ABSTRACT

Nematode molting is a remarkable process where animals must repeatedly build a new apical extracellular matrix (aECM) beneath a previously built aECM that is subsequently shed. The nuclear hormone receptor NHR-23/NR1F1 is an important regulator of *C. elegans* molting. NHR-23 expression oscillates in the epidermal epithelium, and soma-specific NHR-23 depletion causes severe developmental delay and death. Tissue-specific RNAi suggests that *nhr-23* acts primarily in seam and hypodermal cells. NHR-23 coordinates the expression of factors involved in molting, lipid transport/metabolism, and remodeling of the aECM. NHR-23 depletion causes dampened expression of a *nas-37* promoter reporter and a loss of reporter oscillation. The cuticle collagen ROL-6 and zona pellucida protein NOAH-1 display aberrant annular localization and severe disorganization over the seam cells following NHR-23 depletion, while the expression of the adult-specific cuticle collagen BLI-1 is diminished and frequently found in patches. Consistent with these localization defects, the cuticle barrier is severely compromised when NHR-23 is depleted. Together, this work provides insight into how NHR-23 acts in the seam and hypodermal cells to coordinate aECM regeneration during development.

INTRODUCTION

Molting is a critical developmental process required for the growth of all ecdysozoans, a clade comprising an estimated total of 4.5 million living species (Telford et al., 2008). Invertebrate molting involves conserved processes such as apical extracellular matrix (aECM) remodeling, intracellular trafficking, and oscillatory gene expression (Lažetić and Fay, 2017). Molting is also a process of interest for developing new drugs against parasitic nematodes (Ghedin et al., 2007). Parasitic nematodes cause over a billion human infections each year and loss of livestock and crops (Ward, 2015b). However, very few drugs exist, and resistance to those drugs is emerging rapidly (Ward, 2015b). While many genes involved in nematode molting have been identified in *C. elegans* (Frand et al., 2005)*,* little is known about how their gene products are coordinated to promote aECM remodeling and the generation and release of a new cuticle.

Nematodes progress through four periodic larval stages (L1-L4), prior to becoming a reproductive adult (Brenner, 1974). The end of each larval stage is punctuated by a molt that involves trafficking and secretion of aECM components, assembly of a new aECM underneath the old cuticle, followed by separation of the cuticle from the underlying aECM (apolysis) and shedding of the old cuticle (ecdysis) (Lažetić and Fay, 2017). Apolysis coincides with a sleep-like behavior called lethargus (Singh and Sulston, 1978). The cuticle is a collagenous exoskeleton secreted by hypodermal and seam epithelial cells (Page and Johnstone, 2007). The outermost layer (glycocalyx) is rich in carbohydrates and mucins (Nelson et al., 1983; Singh and Sulston, 1978). Beneath the glycocalyx is the glycolipid- and lipid-rich epicuticle which is postulated to function as a hydrophobic surface barrier (Blaxter, 1993; Blaxter et al., 1992). Underlying the epicuticle is a layer comprised mainly of cuticlin proteins. Between the epidermal membrane and the epicuticle are collagen rich layers, including a fluid-filled medial layer with collagen struts (Lažetić and Fay, 2017; Page and Johnstone, 2007). Sets of collagens oscillate in expression over the course of each molt and are classified into three groups based on temporal expression: early, intermediate, and late collagens (Johnstone and Barry, 1996; Page and Johnstone, 2007). While the specific localization of most collagens is unknown, collagens from the same group are thought to be in the

same layer (Lažetić and Fay). A transient structure, the sheath or pre-cuticle, is formed

during each larval stage and thought to pattern the cuticle (Cohen and Sundaram,

2020). Many of the components of this pre-cuticle are related to mammalian matrix

proteins. The *C. elegans* pre-cuticle contains Zona Pellucida proteins, proteins related

to small leucine-rich proteoglycans, and lipid transporters in the lipocalin family (Cohen

and Sundaram, 2020; Cohen et al., 2020; Forman-Rubinsky et al., 2017; Kelley et al.,

2015; Sapio et al., 2005; Vuong-Brender et al., 2017).

Nuclear hormone receptor (NHR) transcription factors are key regulators of molting in

insects and nematodes (King-Jones and Thummel, 2005; Taubert et al., 2011). NHRs

are characterized by a ligand-binding domain (LBD) which has the potential to bind

small molecules such as ligands and dietary-derived metabolites (Taubert et al., 2011).

NHRs have a canonical zinc finger DNA-binding domain (DBD) with an unstructured

hinge region between the DBD and LBD that is subject to post-translational regulation

(Antebi, 2015; Campbell et al., 2008; Ward et al., 2013). A single, conserved nuclear

hormone receptor, NHR-23/NR1F1 (hereafter referred to as NHR-23), an ortholog of

DHR3 in insects and RORα in mammals, is a key regulator of *C. elegans* molting. NHR-

23 is also necessary for spermatogenesis (Ragle et al., 2020; Ragle et al., 2022). *nhr-*

23 mutation or inactivation by RNAi leads to embryonic lethality, larval arrest, ecdysis

defects, and morphology defects (Frand et al., 2005; Gissendanner et al., 2004;

Kostrouchova et al., 1998; Kostrouchova et al., 2001). *nhr-23* mRNA expression

oscillates over the course of each larval stage, peaking mid-larval stage and hitting a

trough at the molt (Gissendanner et al., 2004; Kostrouchova et al., 2001). *nhr-23* is

necessary for all four larval molts and it regulates microRNAs such as *let-7* and *lin-4*

(Kostrouchova et al., 1998; Kostrouchova et al., 2001; Patel et al., 2022, Kinney et al.,

2023). *let-7* also regulates *nhr-23,* suggesting that a feedback loop might coordinate

molting with developmental timing (Patel et al., 2022). The NHR-23 insect ortholog

(DHR3) is part of the molting gene regulatory network (Lam et al., 1997; Ruaud et al.,

2010) and the mammalian ortholog (RORα) regulates circadian rhythms, lipid

metabolism, and immunity (Jetten, 2009). However, how NHR-23 regulates molting and

whether it is part of the core oscillator that promotes rhythmic gene expression over
each larval stage is poorly understood (Tsiairis and Großhans, 2021).

We show here that NHR-23 protein oscillates and rapid NHR-23 depletion via an auxin-
inducible degron causes severe developmental delay and death. Analysis of NHR-23
target genes suggests a role in coordinating aECM assembly, remodeling, and
construction of specific cuticular structures. NHR-23 depletion causes aberrant
localization of the collagens ROL-6 and BLI-1 and of the pre-cuticle factor NOAH-1.
NHR-23 activity is necessary in seam and hypodermal cells to promote molting and
timely development. Our work reveals when and where NHR-23 acts to promote
molting.

RESULTS

NHR-23 protein oscillates during development

Expression of *nhr-23* mRNA oscillates throughout each *C. elegans* larval stage
(Gissendanner et al., 2004; Hendriks et al., 2014; Kostrouchova et al., 2001; Meeuse et
al., 2020). However, mRNA expression profiles do not always correlate with protein
levels (de Sousa Abreu et al., 2009; Vogel and Marcotte, 2012). To determine whether
NHR-23 protein oscillates, we monitored the expression of an endogenously tagged
NHR-23::GFP, over 28 hours (*wrd8* allele; Ragle et al., 2020). During L1, oscillating
expression of NHR-23::GFP was observed in the nuclei of seam and hypodermal cells
(Fig. S1). Expression of NHR-23 rises and falls prior to the completion of the 1st molt
(Fig. 1A). As GFP tags can affect the expression or stability of proteins (Agbulut et al.,
2006; Baens et al., 2006), we assayed the expression of NHR-23 through western
blotting time-courses using an *nhr-23::AID*::3xFLAG* strain (*kry61* allele; Zhang et al.,
2015). These experiments confirmed that NHR-23::AID*::3xFLAG also oscillates (Fig.
1B, C), similar to our NHR-23::GFP imaging experiments (Fig. 1A). We detected three
distinct NHR-23 bands; the lowest band is consistent with the size of NHR-23b/f (Fig.
1B, C, Fig. S2). The upper two bands are larger than expected, which could reflect post-
translational modification or that NHR-23 migrates aberrantly during SDS-PAGE. All
observed NHR-23 bands oscillate, though the lowest band was expressed more

strongly than the other bands (Fig. 1B, C). The peak in NHR-23 expression is earlier in

each larval stage than previous qRT-PCR approaches suggested (Gissendanner et al.,

2004; Kostrouchova et al., 2001) and more in line with recent RNA-seq and imaging

data (Meeuse et al., 2020; Kinney et al., 2023). To understand how NHR-23 promotes

molting, it is important to clarify when it peaks in expression. We examined NHR-

23::GFP expression over the course of the 4[th] larval stage, as vulva morphology allows

for more precise sub-staging (Mok et al., 2015). NHR-23::GFP expression was first

detectable in L4.1 larvae, peaked in expression from L4.2-L4.3, and disappeared in the

vulva by L4.5 and head by L4.6 and then remained undetectable (Fig. 1D,E, S1B,C),

similar to what Kinney et al. (2023) observed with NHR-23::mScarlet.

NHR-23 depletion causes severe developmental delay

Given that NHR-23 oscillates, we tested when NHR-23 was necessary for molting

during the L1 larval stage. In a pilot experiment, we consistently found weaker depletion

phenotypes using an *nhr-23::GFP::AID*::3xFLAG* strain (Ragle et al., 2020) compared

to the *nhr-23::AID*::3xFLAG strain* (Zhang et al., 2015) (Table S1). We therefore

transitioned to using the *nhr-23::AID*::3xFLAG* strain for all experiments involving

phenotypic analysis. We first tested NHR-23::AID*::3xFLAG depletion kinetics and

found robust depletion within 15 minutes after exposure to 4 mM auxin; levels remained

low over the 16 hour timecourse (Fig. 2A).

We performed timed depletion experiments in the L1 larval stage (Fig. 2B) using an *eft-*

3p::TIR1::mRuby2 control strain (henceforth referred to as *TIR1*) and a strain that

permits somatic depletion of NHR-23 in the presence of auxin, (*eft-3p::TIR1::mRuby2;*

nhr-23::AID*::3xFLAG; henceforth referred to as *nhr-23::AID*; *TIR1*). L1s were

synchronized by starvation arrest and release onto MYOB plates. Animals were shifted

onto either 4 mM auxin or control plates every two hours and scored for molting defects

at 20 hours post-release from starvation (Fig. 2B). All *TIR1* control animals shifted onto

control or auxin and *nhr-23::AID*; *TIR1* animals shifted onto control plates reached the

L2 stage (Fig. 2C, D). In contrast, *nhr-23::AID*; *TIR1* animals shifted onto auxin

displayed multiple phenotypes including molting defects, internal vacuoles,

developmental abnormalities such as disrupted tails and viable animals with a squashed

morphology (Fig. 2C). We then quantified the molting defects, scoring animals unable to

shed their cuticles, or with morphological abnormalities (Fig. 2D). Animals shifted onto

auxin within the first four hours post-release tended to arrest as L1 larvae with few

molting or morphological defects (Fig. 2D). To determine at what stage these animals

arrested we used an *hlh-8p::GFP* promoter reporter to monitor the M cell lineage (Harfe

et al., 1998). Newly hatched L1 animals have one M cell which undergoes a

stereotypical series of divisions to produce 16 M lineage cells by the L1 molt (Fig. S3A)

(Sulston and Horvitz, 1977). *nhr-23::AID*; *TIR1* animals shifted onto auxin at 3 hours

post-release were late in L1 based on M cell number (Fig. S3B, C). When NHR-23

animals were scored 72 hours post-auxin shift we observed molting defects and

necrotic animals (Fig. 2E, S4A), suggestive of a severe developmental delay. We also

observed arrested animals with a wild-type morphology which were likely L1 larvae by

size (Fig. 2E). Shifts between 6- and 12-hours post-release resulted in increased

morphological and molting defects (Fig 2D). Animals shifted onto auxin at 9 hours

reached the end of the L1 stage by M cell number (S3B, C). Shifts at 14- and 16-hours

post-release resulted in wild-type L2 animals as judged by size (Fig. 2D).

To test whether NHR-23 was similarly required in other larval stages, we performed

depletion experiments later in development where we could use vulval morphology to

score progression through the L4 stage. We performed depletion experiments shifting

early L3 animals onto auxin and monitored development by scoring vulva morphology

23 hours later. The majority of *TIR1* animals grown on control or auxin plates and *nhr-*

23::AID; *TIR1* animals grown on control plates reached late L4, with a fraction of the

population in early or mid-L4 (Fig. 2F, S4B). In contrast, when *nhr-23::AID*; *TIR1*

animals were shifted to auxin, the vast majority of animals were early L4 larvae with a

fraction of the population remaining in L3 (Fig. 2F, S4B). Repeating these experiments

scoring later timepoints revealed that NHR-23-depleted animals were slowly

progressing through development. After 72 hours on auxin we observed adult animals

as evidenced by the presence of oocytes (Fig. 2G, S4C) while all control animals were

adults. A subset of NHR-23-depleted L4 larvae could not be precisely staged due to

aberrant vulval morphology (Fig. S4C). These animals had large vulval lumens

reminiscent of a 4.3 stage vulva (Mok et al., 2015), but we also observed adults with a

similar large vulval lumen (Fig. S4C). A fraction of *nhr-23::AID*; *TIR1* animals grown for

72 hours on auxin were dead and appeared necrotic with large fluid filled spaces within

the animal (Fig. 2G, S4C). These depletion experiments indicate that NHR-23 is

required for timely developmental progression through multiple larval stages, consistent

with previous reports of *nhr-23* inactivation by RNAi (Macneil et al., 2013; Patel et al.,

2022)

NHR-23 regulates oscillating genes involved in aECM biogenesis and regulation

of protease activity

To gain insight into how NHR-23 depletion causes developmental delay, we analyzed

the expression of *nhr-23*-regulated genes by mining an existing microarray (Kouns et

al., 2011) and oscillatory gene expression datasets (Meeuse et al., 2020) (Table S2). Of

the 265 *nhr-23*-regulated genes in the microarray dataset, 236 (89%) were oscillatory

with the bulk of genes peaking in expression between 180^{o} and 360^{o} (Fig. 3A, Table

S2); *nhr-23* mRNA peaks at 178.11^{o} (Meeuse et al., 2020). A similar trend was

observed in a recent analysis of these datasets (Tsiairis and Großhans, 2021). In

contrast, 10-20% of *C. elegans* genes oscillate in expression (Hendriks et al., 2014; Kim

et al., 2013; Meeuse et al., 2020) (Two-tailed Chi-squared test $p<0.0001$). *nhr-23*-

regulated oscillating genes were enriched in gene ontology classes such as cuticle

structure, regulation of endopeptidases, and metalloprotease activity (Table S3). These

gene ontology classes are a specific subset of the functions enriched in all oscillating

genes (Table S4). To provide biological context, we converted the 360^{o} period to

developmental time assuming a 9-hour larval stage and set the molt from the start of

lethargus (45^{o}) to the end of ecdysis (135^{o}) as in Meeuse et al. (2023). Most *nhr-23*-

regulated genes involved in aECM structure/function, cholesterol metabolism, molting

regulation, transcriptional regulation, and signal transduction had peak amplitudes

within three hours of the *nhr-23* expression peak (Fig. 3B). Genes involved in the

blocked protein unfolding response, as well as some aECM genes peaked later in each

larval stage (Fig. 3B). While the *C. elegans* genome encodes 181 collagen genes

(Teuscher et al., 2019), *nhr-23* only regulated 13 collagens including all of the furrow

collagens implicated in epidermal damage sensing; these furrow collagens are in the

"early" collagen group (*dpy-2, dpy-3, dpy-7, dpy-10;* Table S2, Fig. 3C) (Johnstone and

Barry, 1996; Page and Johnstone, 2007; Dodd et al., 2018). Other *nhr-23*-regulated

intermediate collagens have well-described roles in body morphology (*sqt-1, sqt-2, rol-*

6; Table S2, Fig. 3C) (Kramer and Johnson, 1993; Johnstone and Barry, 1996; Page

and Johnstone, 2007). There are also several sets of genes involved in aECM

biogenesis and remodeling, such as lipocalins, proteases, protease inhibitors, fibrillin,

PAN and zona pellucida (ZP) domain-containing proteins, and leucine-rich repeat

proteins (Fig. 3C). Although not enriched as a gene ontology class, NHR-23 also

regulates several transcription factors implicated in molting or energy metabolism (*peb-*

1, nhr-91, dpy-20; Table S2, Fig. 3B) (Clark et al., 1995, 20; Fernandez et al., 2004;

Kasuga et al., 2013).

To explore NHR-23 direct targets, we analyzed an NHR-23 L3 ChIP-seq dataset

(Gerstein et al., 2010). As expected, NHR-23 was enriched in the promoter region of

genes (Fig. 3D, S5, Table S5). There was also notable enrichment downstream of the

transcriptional end site (Fig. 3D, S5, Table S5). We then examined whether NHR-23

was enriched near *nhr-23*-regulated oscillating genes (Table S2). Genes that peaked in

expression in the hour following the *nhr-23* mRNA peak in expression (Fig. 3E; hour 2)

had the highest average number of NHR-23 peaks flanking and within their gene body.

The average number of NHR-23 peaks/promoter declined and few peaks were detected

flanking genes between larval stage hours 7-9 (Fig. 3E). Genes which peak in

expression close to when *nhr-23* mRNA levels peak tend to have higher NHR-23 levels

upstream, downstream, and within their gene bodies (Fig. 3D early genes).

As NHR-23-regulated oscillating genes were enriched in proteases and protease

inhibitors (Table S2), we tested how NHR-23 depletion affected expression of a *nas-37*

promoter reporter; *nas-37* is a protease implicated in *C. elegans* ecdysis (Davis et al.,

2004). We released synchronized *nhr-23::AID, TIR1* L1 larvae carrying a *nas-*

37p::GFP::PEST reporter onto control and auxin plates and monitored GFP expression

over development. Animals growing on control media displayed oscillating reporter
activity with peaks at 18 and 30 hours (Fig. 4A, B). NHR-23 depleted animals exhibited
a delayed onset of reporter expression and only a subset of animals expressed the
reporter (Fig. 4A, B). There was only one pulse of *nas-37p::GFP::PEST* expression after
which we detected no reporter activity (Fig. 4A, B). Together, these data confirm that
NHR-23 regulates *nas-37* expression and that following NHR-23 depletion, there is a
weaker pulse of target gene expression followed by a failure to express again.

NHR-23 is necessary for NOAH-1 localization to the aECM

Inactivation of the *nhr-23*-regulated predicted protease inhibitor gene *mlt-11* produces
embryonic disorganization with accompanying lethality reminiscent of *noah-1* mutants
(Ragle et al., 2022; Vuong-Brender et al., 2017). NOAH-1 is a zona pellucida domain
protein that is part of the pre-cuticular aECM in embryos and larvae, a transient
structure thought to play a role in patterning the cuticle and in embryonic elongation
(Cohen and Sundaram, 2020). We introduced an mNeonGreen::3xFLAG
(mNG::3xFLAG) tag after the ZP domain which is similar to the insertion site for a
previously generated mCherry knock-in (Fig. 5A; Vuong-Brender et al., 2017). *noah-*
1::mNG::3xFLAG(int) animals had a wild-type brood size (Fig S6A) and in *noah-*
1::mNG::3xFLAG, nhr-23::AID; TIR1 lysates there was a band of the expected full
length protein size (~150 kDa) as well as a lower ~50 kDa band (Fig. 5B). *noah-1* is only
predicted to have a single isoform so this might reflect a cleavage or degradation
product. ZP proteins are frequently cleaved after their ZP domain (Gupta, 2021; Kiefer
and Saling, 2002). While there is a predicted furin RXXR cleavage site at the start of the
CFCS domain, the smaller isoform is consistent with cleavage immediately after the ZP
domain (Fig. 5A,B). It will be valuable in the future to create isoform-specific
mNeonGreen knock-ins to determine where each localizes. NOAH-1::mNG was
observed in the cuticle, and in punctate and tubular structures in the hypodermis (Fig
5C). Hypodermal NOAH-1::mNG significantly overlapped with the lysosomal marker
NUC-1::mCherry (Fig 5C,D) (Guo et al., 2010; Clancy, Vo et al., 2023). NOAH-1::mNG
was detected in lysosomes and weakly in the aECM from L4.1-L4.4 and then reached
its maximum aECM intensity between L4.5-L4.7, localizing to thick bands reminiscent of

annuli (Fig. 5E). aECM expression decreased from L4.8 to young adulthood (Fig. 5E).

NOAH-1::mNG also localized to alae from L4.6 onwards, similar to reports for NOAH-

1::sfGFP (Katz et al., 2022).

We next examined the effect of NHR-23-depletion on NOAH-1 expression and

localization. In L4 larvae, we observed the expected vulval lumen localization and

aECM expression (Cohen et al., 2020; Katz et al., 2022), as well as localization to the

rectum and excretory duct (Fig. 5F). NHR-23 depletion did not affect NOAH-1

localization in the excretory duct or rectum (Fig. 5F). Although vulval morphology is

aberrant in NHR-23-depleted animals, NOAH-1 still localizes to cell surfaces and to

lysosomes (Fig. 5F). In mid-L4, NOAH-1 is normally localized to the cuticle and

lysosomes and is absent over the seam cells (Fig. 5F). NHR-23 depletion caused a loss

of NOAH-1::mNG localization to annuli, the formation of small irregular punctae, and

accumulation of NOAH-1::mNG over seam cells (Fig. 5F). Together, these data indicate

that NHR-23 is dispensable for NOAH-1 expression, but necessary for correct

localization in the cuticle.

NHR-23-depletion causes defects in aECM structure and function

Given the aberrant NOAH-1 localization (Fig. 5) and *nhr-23*-regulation of early and

intermediate collagens (Table S2), we next examined whether NHR-23 depletion

affected aECM structure. We used CRISPR/Cas9-mediated genome editing in *nhr-*

23::AID; TIR1 animals to introduce mNG::3xFLAG cassettes into the intermediate

collagen *rol-6* (C-terminal translational fusion) and *bli-1,* a collagen that forms struts in

the adult cuticle (internal translational fusion)(Fig. 6A). We also introduced a

3xFLAG::mNG into *bli-1* at the same location as the mNG::3xFLAG knock-in. *rol-6* is

regulated by *nhr-23* while *bli-1* is an adult-specific collagen expressed in L4 larvae, so

its expression would not be interrogated in the microarray experiment (Table S2)

(Kouns et al., 2011). We first confirmed that each knock-in had a wild-type brood size

(Fig S6A) and then tested fusion protein size by western blotting. Anti-FLAG

immunoblotting on *nhr-23::AID; TIR1, rol-6::mNG::3xFLAG* lysates detected a single

band of the expected size (~70 kDA; Fig. 6B). In *nhr-23::AID; TIR1, bli-1::3xFLAG::mNG*

there was a single band consistent with N-terminal processing of a monomer at an

RXXR furin cleavage site which would remove 10 kDa (125 kDa predicted size). We

could only detect protein by extracting soluble cuticle components. BLI-

1::3xFLAG::mNG and BLI-1::mNG::3xFLAG had identical localization to one another

(Fig. S6B), and were similar to an extrachromosomal transgenic BLI-1::GFP reporter

(Tong et al., 2009);   all subsequent experiments were performed using *bli-*

1::mNG::3xFLAG(int). Further details of BLI-1::mNG localization will be described

elsewhere (Adams et al., submitted).

We first characterized expression of each translational fusion during the 4th larval stage,

using vulval morphology to stage animals (Mok et al., 2015). ROL-6::mNG was first

detected in thin bands reminiscent of furrows with a jagged pattern over seam cells

(L4.1-L4.3; Fig. 6D). In L4.4-L4.5 some thicker aggregations began appearing and by

L4.6-L4.9 ROL-6::mNG re-localized to thicker bands reminiscent of annuli (Fig. 6D). In

L4.9, a fibrous pattern could also be observed (Fig. 6D). BLI-1::mNG expression was

not detected until L4.5 when it was detected in hypodermal cells (Fig. 6E). In L4.4-L4.8,

BLI-1::mNG was observed in a punctate localization in rows with some irregular brighter

punctae (Fig. 6E). By L4.9, BLI-1::mNG was found in rows of regularly spaced punctae

and was consistently excluded from the cuticle over seam cells throughout L4 and

adulthood (Fig. 6E).

To examine the impact of NHR-23 depletion on BLI-1 and ROL-6 localization, we shifted

nhr-23::AID; TIR1; rol-6::mNG or *nhr-23::AID; TIR1, bli-1::mNG* animals onto control or

auxin plates in early L3. We examined animals after 23 hours on control plates when

animals were mid-late L4s. We scored NHR-23-depleted animals after 47 hours on

auxin plates due to the developmental delay; this approach ensured that control- and

auxin-treated animals were stage-matched. ROL-6::mNG expression appeared

unaffected by NHR-23-depletion, but the furrow localization was irregular and thicker

and we observed gaps in the annular ROL-6::mNG over the seam cells (Fig. 6F). Many

animals had a corset of unshed cuticle to which ROL-6::mNG localized (Fig. 6G). We

next addressed whether NHR-23 depletion affected the formation of struts in the cuticle

medial layer. NHR-23 depletion caused reduced expression of BLI-1::mNG and a loss

of the punctate pattern (Fig. 6H). BLI-1::mNG weakly localized to annuli and animals

displayed bright, disorganized patches of BLI-1::mNG in the cuticle (Fig. 6H). BLI-

1::mNG expression appeared dimmed in the *nhr-23::AID; TIR1* background compared

to a wildtype background which would be consistent with low level auxin-independent

NHR-23-depletion (Fig S6B).

Given the defects in aECM structure following NHR-23-depletion, we tested whether the

epidermal barrier was intact. First, we incubated control and auxin-treated *nhr-23::AID,*

TIR1 animals with the cuticle impermeable, cell membrane permeable Hoechst 33258

dye which stains nuclei. N2 and *TIR1* animals grown on control or auxin plates and *nhr-*

23::AID, TIR1 animals grown on control plates exhibited no staining, while almost all

animals on auxin plates displayed Hoechst staining (Fig. 7A). Animals shifted onto auxin

early or late in L1 robustly expressed an *nlp-29p::GFP* reporter (Fig. 7B, Fig. S7), which

can be activated by infection, acute stress, and physical damage to the cuticle (Pujol et

al., 2008; Zugasti and Ewbank, 2009). We further tested the cuticle integrity in a hypo-

osmotic shock assay in which animals with defective cuticle barriers rapidly die following

exposure to water (Kage- Nakadai et al., 2010). *TIR1* animals grown on control media

or auxin were completely viable in water (Fig. 7C). *nhr-23::AID, TIR1* control animals

had a weakly penetrant sensitivity to hypo-osmotic shock, while the same strain grown

on auxin rapidly died following hypo-osmotic shock (Fig. 7C). These data suggest that

NHR-23 is required for the cuticle barrier establishment or maintenance.

nhr-23 is necessary in seam and hypodermal cells for larval development

Finally, we used a tissue-specific RNAi approach to determine in which cells *nhr-23* was

necessary for molting. In RNAi-proficient N2 animals, 100% of *nhr-23(RNAi)* animals

exhibited developmental delay and molting defects (Fig. 8A). *nhr-23* knockdown in

body-wall muscle and intestinal cells produced no molting defects and we observed only

mild developmental delay (Fig. 8A). Vulval precursor cell-specific *nhr-23(RNAi)* caused

moderate molting defects and developmental delay (Fig. 8A). We observed highly

penetrant developmental delay and molting defects when we performed *nhr-23* RNAi in

a strain using a *wrt-2* promoter for tissue-specific RNAi (Fig. 8A). *wrt-2* is expressed in

seam cells, rectal cells and hypodermal cells (Fig. S8A, B; Aspöck et al., 1999). Given

this result and the defective ROL-6::mNG and NOAH-1::mNG (Fig. 5,6) localization over

seam cells, we generated more tissue-restricted RNAi strains. We used a minimal seam

cell-specific enhancer with a *pes-10* minimal promoter (Ashley et al., 2021), which is

robustly expressed in seam cells with some weak hypodermal expression (Fig. S8C, D).

To test the specificity of this strain, we introduced a *his-72::mNG* knock-in. In animals

treated with control RNAi, we observed nuclear expression in seam, hypodermal,

intestinal, vulval and germline cells, as expected for a histone H3 fusion (Fig S9).

mNeonGreen RNAi caused reduced HIS-72::mNG expression in seam, hypodermal

syncytium, and intestinal nuclei. *nhr-23(RNAi)* in this *SCMp* RNAi strain phenocopied

nhr-23(RNAi) in N2 animals with almost all animals exhibiting developmental delay and

molting defects (Fig. 8A). Notably, intestine-specific *nhr-23* RNAi did not cause

phenotypes (Fig. 8A). We then constructed a hypodermis-specific RNAi strain using the

semo-1 promoter (Kaletsky et al., 2018; Köhnlein et al., 2020). *nhr-23* RNAi in this strain

produced molting defects and developmental delay, though with less penetrance

compared to our *SCMp*-specific RNAi strain (Fig 8A). To test the effect of *SCMp*-

specific *nhr-23* knockdown on the aECM, we introduced a BLI-1::mNG knock-ins in our

seam cell-specific RNAi strain. *nhr-23* depletion in this strain caused reduced BLI-1

levels relative to control RNAi (Fig. 8B). This BLI-1::mNG reduction was comparable to

both *nhr-23* RNAi in a wild-type control (Fig. 8B) and to NHR-23 protein depletion in the

soma (Fig. 6H). Together these data indicated that *nhr-23* is necessary in the seam

and hypodermal cells for timely developmental progression, completion of molting, and

aECM formation.

DISCUSSION

C. elegans molting is a powerful system to understand developmentally programmed

aECM regeneration. In this study, we determine when and where NHR-23 acts to

promote molting. NHR-23 oscillates and its depletion causes severe developmental

delay. *nhr-23*-regulated genes are enriched for cuticle components and protease

inhibitors, and NHR-23 is enriched at the transcription start and end sites of target

genes. NHR-23 depletion causes aberrant localization of the early collagen ROL-6 and
the pre-cuticle component, NOAH-1. These cuticle defects are correlated with a loss of
the cuticle permeability barrier function. Loss of NHR-23 function in L4 stages also
causes severely reduced levels of the adult collagen BLI-1, which is also mislocalized.
These cuticle defects are correlated with a loss of the cuticle barrier. Tissue-specific
RNAi suggests that *nhr-23* is necessary in seam and hypodermal cells.

NHR-23 is necessary for cuticle structure and function
NHR-23 binds more robustly in the promoter and transcription end site of genes which
peak in expression closer to the *nhr-23* mRNA peak in expression (Fig. 3). One
intriguing possibility is that the earlier NHR-23-regulated genes may be more sensitive
to NHR-23 levels. Consistent with this model, there appear to be more NHR-23 peaks
flanking and within early *nhr-23*-regulated genes (Fig. 3, Tables S2, S5). This model
would align with *E. coli* amino acid biosynthesis gene regulation where enzymes earlier
in the pathway have more responsive promoters with higher activity (Zaslaver et al.,
2004). Interestingly, the non-oscillating *nhr-23*-regulated genes had a range in number
of NHR-23 peaks. It's unclear why NHR-23 oscillation is failing to drive oscillation of
these target genes. Some possible explanations include that these genes could have
stable mRNA transcripts in comparison to the oscillating genes or there is combinatorial
gene regulation involving a non-oscillating transcription factor.

NHR-23 regulates the expression of *nas-37*, a protease implicated in molting (Fig. 4;
Frand et al., 2005). Curiously, two other targets (*noah-1, rol-6*) seemed to be
expressed at near wild-type levels but displayed aberrant localization (Fig. 5,6). These
data highlight a potential limitation of a single timepoint gene expression study. A shift in
phase of an oscillating gene without change in amplitude could create the appearance
of up- or down-regulation depending on the timepoint sampled (Tsiairis and Großhans,
2021). The *nhr-23(RNAi)* microarray is a useful starting point to understand how NHR-
23 promotes molting, but an RNA-seq timecourse on NHR-23 depleted animals may be
necessary to identify regulated genes. Such an approach was necessary to determine
how the pioneer factor, BLMP-1, promotes developmental timing (Hauser et al., 2021).

NHR-23-depletion caused aberrant localization of NOAH-1 and ROL-6, lower levels of

BLI-1, and a severe barrier defect (Fig. 5,6,7). There are numerous *nhr-23*-regulated

genes which are implicated in the epithelial barrier. The cuticle furrow formed by six *nhr-*

23-regulated collagens (*dpy-2, dpy-3, dpy-7, dpy-8, dpy-9, dpy-10*) is thought to be

monitored by a sensor which coordinates several stress responses (Dodd et al., 2018).

An RNAi screen of 91 collagens found that inactivation of only these six collagens

caused a barrier defect (Sandhu et al., 2021). Inactivation of a subset of the furrow

collagens causes permeability to Hoechst 33458 dye and elevated *nlp-29p::GFP*

reporter activity (Dodd et al., 2018), consistent with our NHR-23-depletion data (Fig. 7).

bus-8 is a predicted glycosyltransferase that plays a role in the epithelial barrier and is

also *nhr-23*-regulated and its peak expression follows that of the furrow collagens

(Table S2). Screening *nhr-23* regulated genes may reveal other genes implicated in the

epithelial barrier and the peak phase could provide insight into how this barrier is

constructed.

NHR-23-depletion causes developmental delay and failed molting

What is driving NHR-23-depleted animals to eventually attempt to molt? An NHR-23-

regulated promoter reporter (*nas-37p::GFP::PEST*) peaked at a similar time in control

and NHR-23-depleted animals (Fig. 4), but was expressed at lower levels. This data

might suggest that the low levels of NHR-23 present are sufficient to drive low levels of

target gene expression. Consistent with this idea, NHR-23 depletion or knockdown

results in developmental delay but not an arrest (Fig. 2; MacNeil et al., 2013; Patel et

al., 2022). The pre-cuticle component, NOAH-1, and early collagen, ROL-6 are both

expressed and secreted but display localization defects. One possibility is that low

sustained NHR-23-levels eventually allow the accumulation of factors that initiate

apolysis but there is insufficient expression for execution. *nas-37* is necessary for *C.*

elegans ecdysis and recombinant NAS-37 promotes the formation of retractile rings, a

structure associated with molting, in the parasitic nematode *H. contortus* (Davis et al.,

2004). Another non-exclusionary possibility is that NHR-23 controls the expression of

transcription factors in a regulatory cascade, like the gene regulatory network controlling

insect molting (King-Jones and Thummel, 2005; Thummel, 1990). *nhr-23* regulates

peb-1 and *dpy-20*, transcription factors with known roles in molting (Clark et al., 1995,

20; Fernandez et al., 2004, 1). *nhr-23* also regulates *nhr-91* (Kouns et al., 2011)(Table

S2), a nuclear hormone receptor involved in blast cell progression in response to

nutrients (Kasuga et al., 2013). A third possibility is that another timer eventually

promotes molting. The molting timer and heterochronic cycles can be uncoupled,

causing animals to attempt molting before completion of cell division and differentiation,

leading to death (Ruaud and Bessereau, 2006).

nhr-23 is necessary in seam and hypodermal cells to promote molting

Seam cell-enriched *nhr-23* RNAi caused severe developmental delay and molting

defects (Fig 8). Hypodermal-enriched *nhr-23* knock-down produced less penetrant

phenotypes (Fig. 8). The most severe defects in ROL-6 and NOAH-1 localization occur

above the seam cells (Figure 6). Determining how NHR-23 regulates BLI-1 expression

and formation of struts will provide insight into aECM assembly. *nhr-23* is expressed in

other epidermal cells that produce the pre-cuticle and cuticle, such as vulval precursor

cells, rectal cells, and excretory duct cells. NHR-23 depletion did not produce obvious

defects in localization of the pre-cuticle component NOAH-1 in these cells (Fig. 5). We

did not see phenotypes such as excretory duct or pore lumen dilation, which would be

indicative of excretory system defects (Gill et al., 2016). We observed vulval

morphology defects following NHR-23 depletion, but it is not clear whether that is due to

NHR-23 regulation of this specialized aECM or if it is the consequence of

developmental delay and failure to molt. Tissue-specific RNAi or protein depletion will

be required to determine whether NHR-23 activity is necessary in these epithelial cells,

or if it predominantly functions in seam and hypodermal cells. Our data indicate the

importance of tissue-specific RNAi strain validation. *SCMp* drove reporter expression

robustly in seam cells (Fig. S8). However, when we made a tissue-specific RNAi strain

using *SCMp* we observed depletion in seam, hypodermal, and intestinal cells (Fig. S9).

RNAi has an inherent amplification of dsRNA triggers by RNA-dependent RNA

polymerases (Zhang and Ruvkun, 2012). Weak promoter activity in undesired tissues

could reconstitute RNAi in these tissues. A systematic analysis of tissue-specific RNAi
strains is an important future direction for interpreting studies using these strains.
512

513 **Future perspectives**

514 This work highlights the power of timed protein depletion for dissecting the role of
515 oscillating developmental regulators in development. As transcription factors coordinate
516 the expression of batteries of genes in a given biological process, future work will reveal
517 how NHR-23 coordinates apical ECM remodeling, apolysis, and ecdysis.
518

MATERIALS AND METHODS

C. elegans **Strains and Culture**

C. elegans strains were cultured as originally described (Brenner, 1974), except worms
were grown on MYOB instead of NGM. MYOB was made as previously described
(Church et al., 1995). Animals were cultured at 20ºC for all assays, unless otherwise
indicated. For general strain propagation, animals were grown at 15ºC according to
standard protocols. Brood sizes were performed picking L4 larvae to individual wells of
a 6-well plate seeded with OP50 and incubating the plate at 20ºC. Animals were
transferred to new plates daily over 4 days. Two days post-transfer, the number of
hatched progeny and unfertilized eggs were scored.
529

530 **Strains used in this study:**

Strain	Construction	Genotype
CZ27591	Microinjection	*bli-1(ju1789[bli-1::3xFLAG::mNG(int)]) II*
JDW29	Ragle *et al.,* 2020	*nhr-23(wrd8[nhr-23::GFP^AID*::3xFLAG]) I*
JDW180	Microinjection	*wrdEx12[SCMp::NLS-mScarlet (dpi)-tbb-2 3'UTR + pCFJ90 (myo-2p::mCherry)]*
JDW258	Ashley *et al.,*	*wrdSi22 [eft-3p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2*

	2021	*3'UTR+SEC, I:-5.32] ; nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG)) I*
JDW259	Ashley *et al.*, 2021	*wrdSi55[eft-3p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR, I:-5.32] ; nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG)) I*
JDW322	Microinjection	*wrdEx29[wrt-2p::NLS-mScarlet (dpi)-tbb-2 3'UTR]*
JDW348	Crossing	*ieSi57 [Peft-3::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] II; frIs7 [nlp-29p::GFP + col-12p::DsRed] IV*
JDW349	Crossing	*nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG))I; ieSi57 [Peft-3::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] II; frIs7 [nlp-29p::GFP + col-12p::DsRed] IV.x*
JDW330	Crossing	*rde-1 (ne300) V.; jsTi1493 [mosL loxP mex-5p FLP sl2 mNeonGreen rpl-28p FRT GFP-HIS-58 FRT3 mosR] IV*
JDW354	Crossing	*wrdSi60[eft-3p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2:tbb-2 3'UTR, I:-5.32] I; ayIs6 [hlh-8::GFP fusion + dpy-20(+)] X*
JDW355	Crossing	*wrdSi61[eft-3p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2:tbb-2 3'UTR, I:-5.32] ; nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG)) I; ayIs6 [hlh-8::GFP fusion + dpy-20(+)] X*
JDW356	Crossing	*wrdSi59[eft-3p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR, I:-5.32], daf-16(mu86) I; ayIs6 [hlh-8::GFP fusion + dpy-20(+)] X*
JDW357	Crossing	*wrdSi59[eft-3p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR, I:-5.32], nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG)), daf-16(mu86) I; ayIs6 [hlh-8::GFP fusion + dpy-20(+)] X*
JDW370	Microinjection	*jsTi1493 {mosL::loxP::mex-5p::FLP::sl2::mNeonGreen::rpl-28p::FRT [wrdSi64(**SEC::loxP::SCMp::rde-1 CDS+3'UTR**)] FRT3::mosR} IV ; rde-1 (ne300) V*
JDW371	SEC excision of JDW370	*jsTi1493 {mosL loxP [wrdSi65(SCM**p::rde-1 CDS+3'UTR**)] FRT3::mosR} IV ; rde-1 (ne300) V*
JDW376	Clancy, Vo et al., 2023	*jsTi1493 {mosL::loxP::mex-5p::FLP::sl2::mNeonGreen::rpl-28p::FRT [wrdSi67(**SEC**::loxP::mlt-11p (-2.8 kb)::nuc-1::mCherry-tbb-2 3'UTR)] FRT3::mosR} IV*

JDW389	Microinjection	*bli-1(wrd84[bli-1::mNG::3xFLAG(int)]) II*
JDW395	Crossing	*wrdSi73[eft-3p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR] ; nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG)) I; oxIs134[Pnas-37::GFP:rODC(pest)(pWD95@90ng/µl),lin-15(+)]*
JDW459	Microinjection	*nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG))I; ieSi57 [Peft-3::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] II*
JDW460	Microinjection	*noah-1(wrd119[noah-1::internal mNeonGreen (dpi)::3xFLAG::linker])*
JDW463	Microinjection	*nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG)) I ; ieSi57 [Peft-3::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] , bli-1(wrd122[bli-1::internal linker::mNeonGreen (dpi)::3xFLAG::linker]) II*
JDW472	Microinjection	*rol-6(wrd123[rol-6::C-term mNeonGreen (dpi)::3xFLAG::linker]) I ; jsTi1493 wrdSi65[SCMmin::pes-10delta::rde-1 CDS+3'UTR in pLF3FShC]IV; rde-1(ne300) V.*
JDW475	Microinjection	*noah-1(wrd126[noah-1::internal linker::mNeonGreen (dpi)::3xFLAG::linker]) , nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG)) I ; ieSi57 [Peft-3::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] II*
JDW509	Microinjection	*jsTi1493 {mosL::loxP::mex-5p::FLP::sl2::mNeonGreen::rpl-28p::FRT [wrdSi96***(SEC::loxP::semo-1 (Y37A1B.5)p::rde-1 CDS+3'UTR)***]* *FRT3::mosR} IV ; rde-1 (ne300) V*
JDW510	SEC excision of JDW509	*jsTi1493 {mosL loxP [wrdSi97(***semo-1p::rde-1 CDS+3'UTR***)]* *FRT3::mosR} IV ; rde-1 (ne300) V*
JDW525	Microinjection	*his-72(wrd142[mNeonGreen::his-72]) III ; jsTi1493 {mosL loxP [wrdSi65(SCMp::rde-1 CDS+3'UTR)] FRT3::mosR} IV ; rde-1 (ne300) V*
JDW537	Cross of JDW376xJDW 460	*noah-1(wrd119[noah-1::internal mNeonGreen (dpi)::3xFLAG::linker]) I; jsTi1493 wrdSi67{mosL::loxP::mex-5p::FLP::sl2::mNeonGreen::rpl-28p::FRT [wrdSi67(***SEC***::loxP::mlt-11p (-2.8 kb)::nuc-1::mCherry-tbb-2 3'UTR)] FRT3::mosR} IV;*
JDW538	SEC excision of JDW537	*noah-1(wrd119[noah-1::internal mNeonGreen (dpi)::3xFLAG::linker]) I; jsTi1493 wrdSi104{mosL::loxP::mlt-11p (-2.8 kb)::nuc-1::mCherry-tbb-2 3'UTR)] FRT3::mosR} IV*

Strain		Genotype
KRY87	(Zhang et al., 2015)	*nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG))I*
KRY88	Zhang et al., 2015	*nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG))I; ieSi57 [Peft-3::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] II*
NM5179	Nonet, 2020	*jsTi1493 [mosL loxP mex-5p FLP sl2 mNeonGreen rpl-28p FRT GFP-HIS-58 FRT3 mosR] IV*

532   Strains provided by the *Caenorhabditis* Genetics Center:

Strain	Genotype
N2	Wild-type
CA1200	*ieSi57 [Peft-3::TIR1::mRuby::unc-54 3'UTR, cb-unc-119 (+)] II* (Zhang *et al.*, 2015)
CF1038	*daf-16(mu86) I* (Lin et al., 1997)
PD4666	*ayIs6 [hlh-8::GFP fusion + dpy-20(+)] X* (Harfe et al., 1998)
QK52	*rde-1(ne219) V ;* *xkIs99(wrt-2p::rde-1::unc-54 3'UTR)* (Melo and Ruvkun, 2012*)*
VP303	*rde-1(ne219) V ;* *kbIs7[nhx-2p::rde-1 + rol-6(su1006)])* (Espelt et al., 2005)
JU2039	*mfIs70* *[lin-31p::rde-1 + myo2p::GFP] IV ; rde-1(ne219) V* (Barkoulas et al., 2013)
WM118	*rde-1(ne300) V ; neIs9(neIs9 [myo-3::HA::RDE-1 + rol-6(su1006)]) X* (Watts et al., 2020)
MGH171	*alxIs9 [vha-6p::sid-1::SL2::GFP] sid-1(qt9) V* (Melo and Ruvkun, 2012)
NM5179	*jsTi1493 [mosL loxP mex-5p FLP sl2 mNeonGreen rpl-28p FRT GFP-HIS-58 FRT3 mosR] IV* (Nonet, 2020)

WM45	*rde-1(ne300) V* (Tabara et al., 1999)
IG274	*frIs7 [nlp-29p::GFP + col-12p::DsRed] IV* (Pujol et al., 2008)

Other strains

535 EG3200 *oxIs134[Pnas-37::GFP:rODC (PEST) (pWD95@90ng/µl),lin-15(+)]; lin-15(n765ts)* X (Davis et al., 2004) was provided by D. Fay (U Wyoming).

Genome editing and transgenesis

mNeonGreen::3xFLAG knock-ins into *bli-1, noah-1,* and *rol-6* and *mNeonGreen* knock-
ins into *his-72* were generated by injection of Cas9 ribonucleoprotein complexes [700
ng/µl IDT Cas9, 115 ng/µl crRNA and 250 ng/µl IDT tracrRNA] and a dsDNA repair
template (25-50 ng/ul) created by PCR amplification of a plasmid template (Paix et al.,
2014; Paix et al., 2015). The *bli-1* and *noah-1* knock-ins were internal and the
mNG::3xFLAG cassette was flanked by flexible glycine- and serine-rich linker
sequences. Generation of the *BLI-1::3xFLAG::mNG(int)* used in Fig 6C will be
described elsewhere (Adams et al submitted). The C-terminal *rol-6::mNG::3xFLAG*
fusion encoded a flexible linker between the end of *rol-6* and the start of mNG. The
noah-1::mNG::3xFLAG(int) knock-in used a pJW2332 repair template, which will be
described elsewhere; details available upon request. The PCR products were melted to
boost editing efficiency, as previously described (Ghanta and Mello, 2020). crRNAs
used are provided in Table S6. Oligonucleotides used for repair template generation
from template pJW2172 (Ashley et al., 2021) and for genotyping are provided in Table
S7. Genomic and knock-in sequences are provided in File S1. Plasmids used are
provided in Table S8. To generate JDW371 and JDW510 (seam cell and hypodermal-
specific tissue-specific RNAi) we crossed a *jsTi1493* landing pad for recombination-
mediated cassette exchange (RMCE)(Nonet, 2020) into an *rde-1(ne300)* null mutant,
generating JDW330. We used Gibson cloning to introduce a promoterless *rde-1*
genomic coding sequence+3'UTR fragment into the RMCE integration vector
pLF3FShC (Nonet, 2020), creating pJW2247. This vector can be linearized with
*Avr*II+*Bsi*WI double-digestion and promoters can be introduced by Gibson cloning

(Gibson et al., 2009). We generated a *SCMp* (seam cell-specific) and *semo-1p*
(hypodermis specific) by Gibson cloning promoters into linearized pJW2247,
constructing pJW2236 and pJW2264, respectively. These vectors were integrated into
jsTi1493 landing pad in JDW330 using RMCE, as described (Nonet, 2020). *wrt-2p* and
SCMp promoter reporters were constructed in pJW1836 (*NLS::mScarlet::tbb-2 3'*
UTR)(Ashley et al., 2021) by Gibson cloning. Plasmids were injected into N2 animals at
50 ng/μl with a pCFJ90 co-injection marker at 10 ng/μl for the *SCMp* promoter reporter
and a pRF4 co-injection marker at 50 ng/μl for the *wrt-2* promoter reporter (Frøkjaer-
Jensen et al., 2008; Mello et al., 1991). Transgenic lines propagating extrachromosomal
arrays were generated as previously described (Mello et al., 1991). Sequence files for
plasmids are provided in File S2. Primers and cloning details are available upon
request.

Auxin treatment

Control and auxin media/plates were made as described in Ragle et al. 2020. Control
media consisted of MYOB agar+0.25% ethanol. Auxin media was made by dissolving
indole 3-acetic acid (IAA; Alfa Aesar, AAA1055622) in 100% ethanol to 1.6 M and then
mixing it into melted MYOB agar at 55°C to a final concentration of 4 mM prior to
pouring plates. Control plates contained 0.25 % ethanol. Temperature of the media was
monitored with a Lasergrip 1080 infrared thermometer gun (Etekcity). Plates were
seeded with *E. coli* OP50 and incubated overnight at room temperature. Plates were
stored for up to one month at 4ºC prior to use. For most auxin treatment experiments,
animals were synchronized by alkaline bleaching
(dx.doi.org/10.17504/protocols.io.j8nlkkyxdl5r/v1). The collected eggs were incubated in
M9 buffer supplemented with 5 mg/mL cholesterol at 20ºC for 24 hours and arrested L1
larvae were released onto the indicated type of MYOB plate. For the experiment in
Table S1 a timed egg lay was performed. Twenty adult hermaphrodites animals of the
indicated genotype were picked onto control (0.25% ethanol) or auxin (4 mM IAA) plates
and allowed to lay eggs for 2 hours. The adult animals were removed and plates were
incubated for 48 hours at 25ºC.

Microscopy

Synchronized animals were collected from MYOB, control, or auxin plates by either picking or washing off plates. For washing, 1000 µl of M9 + 0.05% gelatin was added to the plate or well, agitated to suspend animals in M9+gelatin, and then transferred to a 1.5 ml tube. Animals were spun at 700xg for 1 min. The media was then aspirated off and animals were resuspended in 500µl M9 + 0.05% gelatin with 5 mM levamisole. 12 µl of animals in M9 + 0.05% with levamisole solution were placed on slides with a 2% agarose pad and secured with a coverslip. For picking, animals were transferred to a 10 µl drop of M9+5 mM levamisole on a 2% agarose pad on a slide and secured with a coverslip. Images were acquired using a Plan-Apochromat 40x/1.3 Oil DIC lens, a Plan-Apochromat 63x/1.4 Oil DIC lens, or a Plan-Apochromat 100x/1.4 OIL DIC lens on an AxioImager M2 microscope (Carl Zeiss Microscopy, LLC) equipped with a Colibri 7 LED light source and an Axiocam 506 mono camera. Acquired images were processed through Fiji software (version: 2.0.0- rc-69/1.52p) (Schindelin et al. 2012). For direct comparisons within a figure, we set the exposure conditions to avoid pixel saturation of the brightest sample and kept equivalent exposure for imaging of the other samples. Co-localization using a Manders' co-efficient (Manders et al., 1992; Manders et al. 1993) was performed as described in Clancy, Vo et al., 2023.

Phenotypic analysis

For the phenotypic analysis in Figure 2, synchronized CA1200 or KRY88 larvae were released onto MYOB plates and then shifted onto control or auxin plates every two hours up to 16 hours. Animals were collected as described in the Microscopy section and imaged by DIC microscopy to score for morphology and shedding of the L1 cuticles. For the M cell lineage experiments in Figure S3, synchronized animals of the indicated genotypes were released onto MYOB plates and shifted onto control or auxin plates at 3 hours (early shift) or 9 hours (late shift) post-release and then imaged at 24 hours post-release, as described in the Microscopy section. M cells were counted and recorded. For the analyses in Figure 2E, synchronized *nhr-23::AID, TIR1* larvae were released on auxin plates and scored for viability and molting defects 72 hours later. For the L3 shift experiments (Fig. 2F,G) synchronized animals of the indicated genotype

were grown on MYOB. At 25 hours post-release, they were shifted onto 6-well control or

auxin MYOB plates seeded with OP50. Animals were collected 23 or 72 hours later as

described in the Microscopy section and imaged by DIC microscopy using a 63x lens to

stage animals according to vulval morphology (Mok et al., 2015). For the reporter

timecourse, synchronized *nhr-23::AID, TIR1, nas-37p::GFP::PEST* animals were

released on control or auxin plates. They were then scored for GFP expression using a

PlanApo 5.0x/0.5 objective on a M165 FC stereomicroscope (Leica) equipped with an

X-cite FIRE LED lightsource (Excelitas) and long pass GFP filter set (Leica #10447407).

We scored GFP expression in the head and hypodermis and did not score rectal GFP

expression, as expression perdured in this tissue after head and hypodermal GFP

expression ceased.

Barrier assays

Hoechst 33258 staining was performed as described previously (Moribe et al., 2004),

except that we used 10 µg/ml of Hoechst 33258 as in Ward et al. (2014). Two biological

replicates were performed examining 50 animals per experiment. The number of

animals with either head or hypodermal nuclei staining with Hoechst 33258 were scored

under a 10x DIC objective. Representative images were then taken with equivalent

exposures using a 63x Oil DIC lens, as described in the imaging section. Hypo-osmotic

stress sensitivity assays were performed on L4 stage larvae as described (Ward et al.,

2014), except we used 20 µl of dH$_2$0. Each strain was assayed in triplicate.

Western Blots

Animals were synchronized by alkaline bleaching, as described in the "Auxin Treatment"

section. For the blots in Fig. 5B and 6B, 30 animals of the indicated genotype and stage

were picked into 30 µl of M9+0.05% gelatin and 10 µl of 4x Laemmli sample buffer was

added and then samples were heated at 95ºC for 5 minutes and stored at -80ºC until

they were resolved by SDS-PAGE. For the blots in Fig. 6C, synchronized animals of the

indicated genotype were collected at the L4 stage. Extracts of soluble cuticle proteins

were prepared as described (Cox *et al.*, 1981) with minor modifications. Briefly, worms

were collected using M9J buffer in a 15 ml Falcon tube and washed 3X with M9J buffer.

Worm pellets were resuspended in 5 ml of sonication buffer and incubated on ice for 10

min, sonicated (Misonix XL-2000 Series Ultrasonic Liquid Processor, 10 X 10 s pulse

with 10 s break) with 30 µl of 0.1 M PMSF. Following sonication, samples were

centrifuged at 8000 rpm for 2 min at 4°C and supernatants stored as Fraction 1 (F1;

intracellular proteins) at -20°C for the purpose of using it as loading control. Pellets were

washed 3X with sonication buffer and the resultant pellets were resuspended in 100 µl

sonication buffer, boiled at 95°C for 2 min with 1 ml of ST buffer and incubated

overnight with rotation at RT. Following incubation, samples were centrifuged at 8000

rpm for 3 min and supernatants were stored as F2 at -20°C. Subsequently, cuticle

pellets were washed 3X with 0.5 % triton X-100 and boiled at 95°C for 2 min with 300 µl

of ST buffer with 5% β-mercaptoethanol and incubated with rotation at room

temperature for 16 h. Finally, the samples were centrifuged at 8000 rpm for 3 min and

supernatants collected and stored as F3 (soluble cuticle proteins). F3 Fraction was

precipitated with methanol/chloroform and resuspended in 100 µl of rehydration buffer.

For the remaining blots, animals of the indicated genotype were washed out of wells on

a 6-well plate at the indicated timepoints with M9+0.05% gelatin (VWR, 97062-620),

transferred to a 1.5 ml tube and washed twice more with M9+0.05% gelatin, as

previously described (Vo et al., 2021). Animals were pelleted, transferred in a 100 µl

volume to a new 1.5 ml tube, and flash frozen in liquid nitrogen, then stored at -80ºC.

Prior to SDS-PAGE, samples were freeze-cracked twice in liquid nitrogen or on dry ice,

Laemmli sample buffer was added to 1X and samples were heated to 95ºC for 5

minutes. For the western blots in Figure 1C and 1D 6000 KRY88 animals per well were

transferred onto a 6-well MYOB plate seeded with OP50. For the western blot in Figure

2B, 6000 synchronized KRY88 animals were plated onto control or auxin plates. Control

animals were harvested at peak NHR-23 expression (12 hours of growth on control

plates) and auxin-treated animals were shifted onto auxin at this time and samples were

taken at the indicated time points, as described above. For the western blots in Figure

S7, 500 animals of the indicated genotype were transferred into wells of a 6-well control

or auxin plate seeded with OP50. Animals were incubated for 24 hours at 20ºC and

collected as described above.

For the blots in Figs. 1C,D, 10 µl of lysate and 7.5 µl 1:1 mix of Amersham ECL Rainbow Molecular Weight Markers (95040-114) and Precision Plus Protein Unstained Standards (#1610363) was resolved by SDS-PAGE using precast 4-20% MiniProtean TGX Stain Free Gels (Bio-Rad). The blot in Fig. S7, the lysates had more variable levels of total protein, particularly the NHR-23 depleted samples that produced an early larval arrest. Therefore, we quantified the signal of the most intense band by Stain Free imaging (Posch et al., 2013) using a Bio-Rad ChemiDoc imaging system, and ran a new gel with normalized loading. The lysates in Fig. 2A, 5B, 6B, and S7 were resolved as described above except we used a Spectra™ Multicolor Broad Range Protein Ladder (Thermo; # 26623). Proteins were transferred to a polyvinylidene difluoride membrane by semi-dry transfer with a TransBlot Turbo (Bio-Rad). In Figs 1C and 7C, total protein, pre- and post-transfer, was monitored using the stain-free fluorophore as described (Posch et al., 2013; Ward, 2015a). Membranes were washed in TBST and blocked in TBST+5% milk (TBST-M; Nestle Carnation Instant Nonfat Dry Milk, 25.6-Ounce, Amazon) for one hour at room temperature. Blots were rocked in primary antibodies in TBST-M overnight at 4ºC and then washed 4x5min with TBST. For primary antibodies conjugated with horseradish peroxidase (HRP) the blots were developed after the last TBST wash. Otherwise, blots were incubated with HRP-conjugated secondary antibodies in TBST-M for one hour at room temperature followed by 4x5min TBST washes and then developed as described below. The primary antibody used for Figures 1C,D, 2A, 5B, and 6B was horseradish peroxidase (HRP) conjugated anti-FLAG M2 (Sigma-Aldrich, A8592-5x1MG, Lot #SLCB9703) at a 1:2000 dilution. Precision Protein StrepTactin-HRP Conjugate (Bio-Rad, #1610381, Lot #64426657) was included with the primary antibody at a 1:10,000 dilution to visualize the protein size standard during blot imaging for Figs 1C,D. For the blot in Fig. 2A, 5B, 6B, we used mouse anti-alpha-Tubulin 12G10 (Developmental Studies Hybridoma Bank, the "-c" concentrated supernatant at 1:2000 (Fig 2A, S7) or 1:4000 (Fig. 5B, 6B). The secondary antibodies were Digital anti-mouse (Kindle Biosciences LLC, R1005) diluted 1:10,000 (Fig. 2A, S7) or 1:20,000 (Fig. S6B,C). Blots were incubated for 5 minutes with 1 ml of Supersignal

716 West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific, 34095) and
717 imaged using the 'chemi high-resolution' setting on a Bio-Rad ChemiDoc MP System.
718

For the blots in Fig. 6C, 20 µl was loaded on to NuPAGE 4 to 12% Protein Gels with 7
µl of 4X NuPAGE LDS sample buffer as described by the manufacturer (Invitrogen:
NP0322BOX) and run for 2 h at 100 V, transferred to 0.45 micron nitrocellulose
membrane (Bio-Rad) at 4 °C for 90 min using Bio-Rad WET transfer system. Following
transfer, membranes were blocked in Intercept blocking buffer (LI-COR: 927-60001) for
1 h. Later, the membrane was incubated overnight in 1° Ab (ChromoTek: 32F6) at
1:5,000 dilution at 4°C on rocker. Membranes were washed 3X with TBST and
incubated in 2° Ab (Sigma 12-349) at 1:100,000 dilution at RT on rocker for 1 h and
washed 3X with TBST. Blots were visualized with Supersignal West Femto detection kit
(Thermo Fisher) using a LI-COR Odyssey imager. Similarly, to test the loading control,
10 µl of F1 samples were loaded and following transfer to nitrocellulose membrane,
probed with monoclonal mouse anti-actin clone C4 (Thermofisher scientific:
ICN691001).
732

Bioinformatics
Fig. 3A was generated in R (R Core Team). Phase and amplitude in Table S2 were
converted to x and y coordinates for each gene by calculating x = amplitude *
cos(phase) and y = amplitude * sin(phase). For Fig. 3D and S5, L3 NHR-23 ChIP-Seq
data was downloaded from http://www.modencode.org, accession ID
modEncode_3837. Wig files were converted to bigwig format and loaded into the
Bioconductor package SeqPlots (Huber et al., 2015; Stempor and Ahringer, 2016). A
bed file with the start and end coordinates of all genes' CDS was generated (Wormbase
WS220) and loaded into SeqPlots. SeqPlots aligned all genes by the start and end
position and scaled the coding sequence of each gene to 2 kb. The average signal over
all aligned genes was calculated in 25 bp windows from 1 kb upstream of the start site
to 1kb downstream of the end site. Peak calls for NHR-23 in L3 (Fig 4C) were obtained
from GEO (accession numbers: GSM1183659, GSM1183660, GSM1183661,
GSM118366). Only peak calls validated by both replicates were considered. Peaks

were assigned to the following genome features: gene body (WormBase WS220), 1kb

upstream of the start site of coding genes, between 1kb and 3kb upstream of the gene

start site and between 3kb and 5kb upstream of the gene start site. The same intervals

were chosen downstream of the gene end sites. An NHR-23 peak was assigned to a

feature if it overlapped with it by at least 100bp. The peak analysis data is in Table S5.

The total number of peaks in all bins was tallied and is presented in Table S2 and in Fig.

3D. For Fig. 3D and 4E the phase for the oscillating genes in Table S1 was converted to

time during the molting cycle by assuming a 9 hour larval stage which makes each

hour=40º. We used the phasing from Meeuse et al., 2023 with lethargus starting at 45º

and ecdysis ending at 135º. The phase in hours was then plotted along the x axis and

the oscillating genes' amplitude was plotted on the y axis. The gene annotation was

based on the Concise Description and Automated Description attributes downloaded

from Wormbase.

RNAi

RNAi feeding plates were made by melting a solidified bottle of MYOB and adding

Carbenicillin (25 ug/ml) and IPTG (8mM) once cooled to 55☐. dsRNA-expressing *E.*

coli bacteria were streaked on LB plates with Ampicillin (100 µg/ml) and grown overnight

at 37☐. A single colony was picked into 25 ml LB with 100 µg/ml Ampicillin and 12.5

µg/ml Tetracycline and shaken overnight at 37☐ at 220 RPMs. The next day, the liquid

culture was pelleted and resuspended in 1.25 ml LB with 100 µg/ml Ampicillin resulting

in a 20x concentration from the original overnight liquid culture. 90 µl of the liquid

culture was spread per small RNAi plate and allowed to dry. The plates were incubated

at room temperature in the dark for 3 days. 10 adult worms from each strain were

allowed to lay eggs on plates spread with *E. coli* bacteria containing either the control

plasmid L4440 or the *nhr-23* RNAi knockdown plasmid for 1-2 hours. The adults were

removed and the eggs allowed to develop at 20☐ for 2 or 3 days depending on the

experiment. For tissue-specific RNAi experiments we used RNAi defective *rde-1* mutant

animals carrying *rde-1* transgenes to rescue RNAi in specific tissues (Qadota et al.,

2007) The one exception was strain MGH171 which used an intestinal specific rescue

of *sid-1* in an RNAi defective *sid-1* mutant (Melo and Ruvkun, 2012).

Statistical Analysis

Statistical tests and numbers of animals analyzed are detailed in figure legends.

Competing interests

The authors declare no competing or financial interests.

Acknowledgements

We thank Chris Hammell, David Matus, and Julian Ceron Madrigal for their critical reading of the manuscript. We thank David Fay, Miles Pufall, Helge Grosshans, and Ali Shariati for helpful discussions. We thank Jennifer Adams for generation of initial *rol-6* and *bli-1* mNeonGreen knock-ins. Figure S3A was created with Biorender.com.

Funding

This work was funded by the National Institutes of Health (NIH) National Institute of General Medical Sciences (NIGMS) [R00GM107345 and R01GM138701] to J.D.W. M.P. and A.D.C. were supported by NIGMS R35GM134970. Some strains were provided by the Caenorhabditis Genetics Center, which is funded by the NIH Office of Research Infrastructure Programs [P40 OD010440]. L.C.J. was funded by an NIGMS training grant [T32 GM133391]. The anti-alpha tubulin 12G10 monoclonal antibody developed by J. Frankel and E.M. Nelson of the University of Iowa was obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA 52242.

REFERENCES

Agbulut, O., Coirault, C., Niederländer, N., Huet, A., Vicart, P., Hagège, A., Puceat, M. and Menasché, P. (2006). GFP expression in muscle cells impairs actin-myosin interactions: implications for cell therapy. *Nat. Methods* **3**, 331–331.

Antebi, A. (2015). Nuclear receptor signal transduction in *C. elegans*. *WormBook* 1–49.

Ashley, G. E., Duong, T., Levenson, M. T., Martinez, M. A. Q., Johnson, L. C.,

Hibshman, J. D., Saeger, H. N., Palmisano, N. J., Doonan, R., Martinez-
Mendez, R., et al. (2021). An expanded auxin-inducible degron toolkit for
Caenorhabditis elegans. Genetics **217**, iyab006.

Aspöck, G., Kagoshima, H., Niklaus, G. and Bürglin, T. R. (1999). *Caenorhabditis*
elegans has scores of hedgehog-related genes: sequence and expression
analysis. *Genome Res.* **9**, 909–923.

Baens, M., Noels, H., Broeckx, V., Hagens, S., Fevery, S., Billiau, A. D.,
Vankelecom, H. and Marynen, P. (2006). The Dark Side of EGFP: Defective
Polyubiquitination. *PLOS ONE* **1**, e54.

Barkoulas, M., van Zon, J. S., Milloz, J., Van Oudenaarden, A. and Félix, M.-A.
(2013). Robustness and Epistasis in the *C. elegans* Vulval Signaling Network
Revealed by Pathway Dosage Modulation. *Dev. Cell* **24**, 64–75.

Baugh, L. R. and Sternberg, P. W. (2006). DAF-16/FOXO regulates transcription of
cki-1/Cip/Kip and repression of *lin-4* during *C. elegans* L1 arrest. *Curr. Biol.* **16**,
780–785.

Blaxter, M. L. (1993). Cuticle surface proteins of wild type and mutant *Caenorhabditis*
elegans. J. Biol. Chem. **268**, 6600–6609.

Blaxter, M. L., Page, A. P., Rudin, W. and Maizels, R. M. (1992). Nematode surface
coats: actively evading immunity. *Parasitol. Today Pers. Ed* **8**, 243–247.

Brenner, S. (1974). The genetics of *Caenorhabditis elegans. Genetics* **77**, 71–94.

Campbell, L. A., Faivre, E. J., Show, M. D., Ingraham, J. G., Flinders, J., Gross, J.
D. and Ingraham, H. A. (2008). Decreased Recognition of SUMO-Sensitive
Target Genes following Modification of SF-1 (NR5A1). *Mol. Cell. Biol.* **28**, 7476–
7486.

Church, D. L., Guan, K. L. and Lambie, E. J. (1995). Three genes of the MAP kinase
cascade, *mek-2*, *mpk-1/sur-1* and *let-60 ras*, are required for meiotic cell cycle
progression in *Caenorhabditis elegans. Development* **121**, 2525–2535.

Clancy, J. C., Vo, A. A., Miles, K.M., Levenson, M. T., Ragle, J. M. and Ward, J. D.
(2023). Experimental considerations for study of *C. elegans* lysosomal proteins.
G3, jkad032

Clark, D. V., Suleman, D. S., Beckenbach, K. A., Gilchrist, E. J. and Baillie, D. L.

(1995). Molecular cloning and characterization of the *dpy-20* gene of

Caenorhabditis elegans. *Mol. Gen. Genet. MGG* **247**, 367–378.

Cohen, J. D. and Sundaram, M. V. (2020). *C. elegans* Apical Extracellular Matrices

Shape Epithelia. *J. Dev. Biol.* **8**, 23.

Cohen, J. D., Sparacio, A. P., Belfi, A. C., Forman-Rubinsky, R., Hall, D. H., Maul-

Newby, H., Frand, A. R. and Sundaram, M. V. (2020). A multi-layered and

dynamic apical extracellular matrix shapes the vulva lumen in *Caenorhabditis*

elegans. *eLife* **9**, e57874.

Cox, G. N., Kusch, M. and Edgar, R. S. (1981). Cuticle of *Caenorhabditis elegans*: its

isolation and partial characterization. *J. Cell Biol.* 90, 7–17.

Davis, M. W., Birnie, A. J., Chan, A. C., Page, A. P. and Jorgensen, E. M. (2004). A

conserved metalloprotease mediates ecdysis in *Caenorhabditis elegans*.

Development **131**, 6001–6008.

de Sousa Abreu, R., Penalva, L. O., Marcotte, E. M. and Vogel, C. (2009). Global

signatures of protein and mRNA expression levels. *Mol. Biosyst.* **5**, 1512–1526.

Dodd, W., Tang, L., Lone, J.-C., Wimberly, K., Wu, C.-W., Consalvo, C., Wright, J.

E., Pujol, N. and Choe, K. P. (2018). A Damage Sensor Associated with the

Cuticle Coordinates Three Core Environmental Stress Responses in

Caenorhabditis elegans. *Genetics* **208**, 1467–1482.

Espelt, M. V., Estevez, A. Y., Yin, X. and Strange, K. (2005). Oscillatory Ca2+

signaling in the isolated *Caenorhabditis elegans* intestine: role of the inositol-

1,4,5-trisphosphate receptor and phospholipases C beta and gamma. *J. Gen.*

Physiol. **126**, 379–392.

Fernandez, A. P., Gibbons, J. and Okkema, P. G. (2004). *C. elegans peb-1* mutants

exhibit pleiotropic defects in molting, feeding, and morphology. *Dev. Biol.* **276**,

352–366.

Forman-Rubinsky, R., Cohen, J. D. and Sundaram, M. V. (2017). Lipocalins Are

Required for Apical Extracellular Matrix Organization and Remodeling in

Caenorhabditis elegans. *Genetics* **207**, 625–642.

Frand, A. R., Russel, S. and Ruvkun, G. (2005). Functional Genomic Analysis of *C.*

elegans Molting. *PLoS Biol.* **3**, e312.

Frøkjaer-Jensen, C., Davis, M. W., Hopkins, C. E., Newman, B. J., Thummel, J. M.,
Olesen, S.-P., Grunnet, M. and Jorgensen, E. M. (2008). Single-copy insertion
of transgenes in *Caenorhabditis elegans*. *Nat. Genet.* **40**, 1375–1383.

Gerstein, M. B., Lu, Z. J., Van Nostrand, E. L., Cheng, C., Arshinoff, B. I., Liu, T.,
Yip, K. Y., Robilotto, R., Rechtsteiner, A., Ikegami, K., et al. (2010).
Integrative Analysis of the *Caenorhabditis elegans* Genome by the modENCODE
Project. *Science* **330**, 1775–1787.

Ghanta, K. S. and Mello, C. C. (2020). Melting dsDNA Donor Molecules Greatly
Improves Precision Genome Editing in *Caenorhabditis elegans*. *Genetics* **216**,
643–650.

Ghedin, E., Wang, S., Spiro, D., Caler, E., Zhao, Q., Crabtree, J., Allen, J. E.,
Delcher, A. L., Guiliano, D. B., Miranda-Saavedra, D., et al. (2007). Draft
genome of the filarial nematode parasite *Brugia malayi*. *Science* **317**, 1756–
1760.

Gibson, D. G., Young, L., Chuang, R.-Y., Venter, J. C., Hutchison, C. A. and Smith,
H. O. (2009). Enzymatic assembly of DNA molecules up to several hundred
kilobases. *Nat. Methods* **6**, 343–345.

Gill, H. K., Cohen, J. D., Ayala-Figueroa, J., Forman-Rubinsky, R., Poggioli, C.,
Bickard, K., Parry, J. M., Pu, P., Hall, D. H. and Sundaram, M. V. (2016).
Integrity of Narrow Epithelial Tubes in the C. elegans Excretory System Requires
a Transient Luminal Matrix. *PLoS Genet.* **12**, e1006205.

Gissendanner, C. R., Crossgrove, K., Kraus, K. A., Maina, C. V. and Sluder, A. E.
(2004). Expression and function of conserved nuclear receptor genes in
Caenorhabditis elegans. *Dev. Biol.* **266**, 399–416.

Gupta, S. K. (2021). Human Zona Pellucida Glycoproteins: Binding Characteristics With
Human Spermatozoa and Induction of Acrosome Reaction. *Front. Cell Dev. Biol.*
9, 619868.

Guo, P., Hu, T., Zhang, J., Jiang, S. and Wang, X. (2010). Sequential action of
Caenorhabditis elegans Rab GTPases regulates phagolysosome formation
during apoptotic cell degradation. *Proc. Natl. Acad. Sci.* 107, 18016–18021.

Harfe, B. D., Gomes, A. V., Kenyon, C., Liu, J., Krause, M. and Fire, A. (1998).

Analysis of a *Caenorhabditis elegans* Twist homolog identifies conserved and divergent aspects of mesodermal patterning. *Genes Dev.* **12**, 2623–2635.

Hauser, Y. P., Meeuse, M. W. M., Gaidatzis, D. and Großhans, H. (2021). The BLMP-1 transcription factor promotes oscillatory gene expression to achieve timely molting. *bioRxiv* 2021.07.05.450828.

Hendriks, G.-J., Gaidatzis, D., Aeschimann, F. and Großhans, H. (2014). Extensive oscillatory gene expression during *C. elegans* larval development. *Mol. Cell* **53**, 380–392.

Huber, W., Carey, V. J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B. S., Bravo, H. C., Davis, S., Gatto, L., Girke, T., et al. (2015). Orchestrating high-throughput genomic analysis with Bioconductor. *Nat. Methods* **12**, 115–121.

Jetten, A. M. (2009). Retinoid-related orphan receptors (RORs): critical roles in development, immunity, circadian rhythm, and cellular metabolism. *Nucl. Recept. Signal.* **7**,.

Johnstone, I. L. and Barry, J. D. (1996). Temporal reiteration of a precise gene expression pattern during nematode development. *EMBO J.* **15**, 3633–3639.

Kage-Nakadai, E., Kobuna, H., Kimura, M., Gengyo-Ando, K., Inoue, T., Arai, H. and Mitani, S. (2010). Two very long chain fatty acid acyl-CoA synthetase genes, *acs-20* and *acs-22*, have roles in the cuticle surface barrier in *Caenorhabditis elegans*. *PLoS ONE* **5**, e8857.

Kaletsky, R., Yao, V., Williams, A., Runnels, A. M., Tadych, A., Zhou, S., Troyanskaya, O. G. and Murphy, C. T. (2018). Transcriptome analysis of adult *Caenorhabditis elegans* cells reveals tissue-specific gene and isoform expression. *PLOS Genet.* **14**, e1007559.

Kasuga, H., Fukuyama, M., Kitazawa, A., Kontani, K. and Katada, T. (2013). The microRNA miR-235 couples blast-cell quiescence to the nutritional state. *Nature* **497**, 503–506.

Katz, S. S., Barker, T. J., Maul-Newby, H. M., Sparacio, A. P., Nguyen, K. C. Q., Maybrun, C. L., Belfi, A., Cohen, J. D., Hall, D. H., Sundaram, M. V., et al. (2022). A transient apical extracellular matrix relays cytoskeletal patterns to shape permanent acellular ridges on the surface of adult *C. elegans*. *PLOS*

932   *Genet.* **18**, e1010348.

**Kelley, M., Yochem, J., Krieg, M., Calixto, A., Heiman, M. G., Kuzmanov, A., Meli,
V., Chalfie, M., Goodman, M. B., Shaham, S., et al.** (2015). FBN-1, a fibrillin-
related protein, is required for resistance of the epidermis to mechanical
deformation during *C. elegans* embryogenesis. *eLife* **4**, 1629.

Kiefer, S. M. and Saling, P. (2002). Proteolytic Processing of Human Zona Pellucida
Proteins. *Biol. Reprod.* 66, 407–414

Kim, D. hyun, Grün, D. and van Oudenaarden, A. (2013). Dampening of expression
oscillations by synchronous regulation of a microRNA and its target. *Nat. Genet.*
45, 1337–1344.

King-Jones, K. and Thummel, C. S. (2005). Nuclear receptors-a perspective from
Drosophila. *Nat. Rev. Genet.* **6**, 311–323.

**Kinney, B., Sahu, S., Stec, N., Hills-Muckey, K., Adams, D. W., Wang, J., Jaremko,
M., Joshua-Tor, L., Keil, W. and Hammell, C. M.** (2023). A circadian-like gene
network regulates heterochronic miRNA transcription in *C. elegans*. *bioRxiv*
2022.09.26.509508.

**Köhnlein, K., Urban, N., Guerrero-Gómez, D., Steinbrenner, H., Urbánek, P.,
Priebs, J., Koch, P., Kaether, C., Miranda-Vizuete, A. and Klotz, L.-O.** (2020).
A *Caenorhabditis elegans* ortholog of human selenium-binding protein 1 is a pro-
aging factor protecting against selenite toxicity. *Redox Biol.* **28**, 101323.

Kostrouchova, M., Krause, M., Kostrouch, Z. and Rall, J. E. (1998). CHR3: a
Caenorhabditis elegans orphan nuclear hormone receptor required for proper
epidermal development and molting. *Development* **125**, 1617–1626.

Kostrouchova, M., Krause, M., Kostrouch, Z. and Rall, J. E. (2001). Nuclear
hormone receptor CHR3 is a critical regulator of all four larval molts of the
nematode *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 7360–
7365.

**Kouns, N. A., Nakielna, J., Behensky, F., Krause, M. W., Kostrouch, Z. and
Kostrouchova, M.** (2011). NHR-23 dependent collagen and hedgehog-related
genes required for molting. *Biochem. Biophys. Res. Commun.* **413**, 515–520.

Kramer, J. M. and Johnson, J. J. (1993). Analysis of mutations in the sqt-1 and rol-6

collagen genes of Caenorhabditis elegans. *Genetics* **135**, 1035–1045.

Lam, G. T., Jiang, C. and Thummel, C. S. (1997). Coordination of larval and prepupal

gene expression by the DHR3 orphan receptor during *Drosophila*

metamorphosis. *Development* **124**, 1757–1769.

Lažetić, V. and Fay, D. S. (2017). Molting in *C. elegans. Worm* **6**, 1–20.

Lin, K., Dorman, J. B., Rodan, A. and Kenyon, C. (1997). *daf-16*: An HNF-3/forkhead

family member that can function to double the life-span of *Caenorhabditis*

elegans. Science **278**, 1319–1322.

Macneil, L. T., Watson, E., Arda, H. E., Zhu, L. J. and Walhout, A. J. M. (2013). Diet-

Induced Developmental Acceleration Independent of TOR and Insulin in *C.*

elegans. Cell **153**, 240–252.

Manders, E. M., Stap, J., Brakenhoff, G. J., van Driel, R. and Aten, J. A. (1992).

Dynamics of three-dimensional replication patterns during the S-phase, analysed

by double labelling of DNA and confocal microscopy. *J. Cell Sci.* 103 (Pt 3),

857–862.

Manders, E. M. M., Verbeek, F. J. and Aten, J. A. (1993). Measurement of co-

localization of objects in dual-colour confocal images. *J. Microsc.* 169, 375–382.

Meeuse, M. W., Hauser, Y. P., Morales Moya, L. J., Hendriks, G., Eglinger, J.,

Bogaarts, G., Tsiairis, C. and Großhans, H. (2020). Developmental function

and state transitions of a gene expression oscillator in *Caenorhabditis elegans*.

Mol. Syst. Biol. **16**, e9975.

Meeuse, M. W. M., Hauser, Y. P., Nahar, S., Smith, A. A. T., Braun, K., Azzi, C.,

Rempfler, M. and Großhans, H. (2023). C. elegans molting requires rhythmic

accumulation of the Grainyhead/LSF transcription factor GRH-1. *EMBO J.* n/a,

e111895.

Mello, C. C., Kramer, J. M., Stinchcomb, D. and Ambros, V. (1991). Efficient gene

transfer in *C. elegans*: extrachromosomal maintenance and integration of

transforming sequences. *EMBO J.* **10**, 3959–3970.

Melo, J. A. and Ruvkun, G. (2012). Inactivation of conserved *C. elegans* genes

engages pathogen- and xenobiotic-associated defenses. *Cell* **149**, 452–466.

Mok, D. Z. L., Sternberg, P. W. and Inoue, T. (2015). Morphologically defined sub-

stages of *C. elegans* vulval development in the fourth larval stage. *BMC Dev.*

Biol. 1–8.

Moribe, H., Yochem, J., Yamada, H., Tabuse, Y., Fujimoto, T. and Mekada, E.

(2004). Tetraspanin protein (TSP-15) is required for epidermal integrity in

Caenorhabditis elegans. *J. Cell Sci.* **117**, 5209–5220.

Nelson, F. K., Albert, P. S. and Riddle, D. L. (1983). Fine structure of the

Caenorhabditis elegans secretory-excretory system. J. Ultrastruct. Res. **82**, 156–

171.

Nonet, M. L. (2020). Efficient Transgenesis in *Caenorhabditis elegans* Using Flp

Recombinase-Mediated Cassette Exchange. *Genetics* **215**, 903–921.

Page, A. P. and Johnstone, I. L. (2007). The cuticle. *WormBook.* 1–15.

Paix, A., Wang, Y., Smith, H. E., Lee, C.-Y. S., Calidas, D., Lu, T., Smith, J.,

Schmidt, H., Krause, M. W. and Seydoux, G. (2014). Scalable and versatile

genome editing using linear DNAs with microhomology to Cas9 Sites in

Caenorhabditis elegans. Genetics **198**, 1347–1356.

Paix, A., Folkmann, A., Rasoloson, D. and Seydoux, G. (2015). High Efficiency,

Homology-Directed Genome Editing in *Caenorhabditis elegans* Using CRISPR-

Cas9 Ribonucleoprotein Complexes. *Genetics* **201**, 47–54.

Posch, A., Kohn, J., Oh, K., Hammond, M. and Liu, N. (2013). V3 stain-free workflow

for a practical, convenient, and reliable total protein loading control in western

blotting. *J. Vis. Exp.* 50948.

Pujol, N., Cypowyj, S., Ziegler, K., Millet, A., Astrain, A., Goncharov, A., Jin, Y.,

Chisholm, A. D. and Ewbank, J. J. (2008). Distinct innate immune responses to

infection and wounding in the *C. elegans* epidermis. *Curr. Biol.* **18**, 481–489.

Qadota, H., Inoue, M., Hikita, T., Köppen, M., Hardin, J. D., Amano, M., Moerman,

D. G. and Kaibuchi, K. (2007). Establishment of a tissue-specific RNAi system

in *C. elegans. Gene* 400, 166–173.

R Core Team *R: A language and environment for statistical   computing.* Vienna,

Austria: R Foundation for Statistical Computing.

Ragle, J. M., Aita, A. L., Morrison, K. N., Martinez-Mendez, R., Saeger, H. N.,

Ashley, G. A., Johnson, L. C., Schubert, K. A., Shakes, D. C. and Ward, J. D.

(2020). The conserved molting/circadian rhythm regulator NHR-23/NR1F1 serves as an essential co-regulator of *C. elegans* spermatogenesis. *Development* **147**, dev193862.

Ragle, J. M., Morrison, K. N., Vo, A. A., Johnson, Z. E., Hernandez Lopez, J., Rechtsteiner, A., Shakes, D. C. and Ward, J. D. (2022). NHR-23 and SPE-44 regulate distinct sets of genes during *C. elegans* spermatogenesis. *G3*, jkac256.

Roach, N. P., Sadowski, N., Alessi, A. F., Timp, W., Taylor, J. and Kim, J. K. (2020). The full-length transcriptome of *C. elegans* using direct RNA sequencing. *Genome Res.*

Ruaud, A.-F. and Bessereau, J.-L. (2006). Activation of nicotinic receptors uncouples a developmental timer from the molting timer in *C. elegans*. *Development* **133**, 2211–2222.

Ruaud, A.-F., Lam, G. and Thummel, C. S. (2010). The *Drosophila* nuclear receptors DHR3 and FTZ-F1 control overlapping developmental responses in late embryos. *Development* **137**, 123–131.

Sandhu, A., Badal, D., Sheokand, R., Tyagi, S. and Singh, V. (2021). Specific collagens maintain the cuticle permeability barrier in *Caenorhabditis elegans*. *Genetics* **217**,.

Sapio, M. R., Hilliard, M. A., Cermola, M., Favre, R. and Bazzicalupo, P. (2005). The Zona Pellucida domain containing proteins, CUT-1, CUT-3 and CUT-5, play essential roles in the development of the larval alae in *Caenorhabditis elegans*. *Dev. Biol.* **282**, 231–245.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods* 9, 676–682.

Schindler, A. J., Baugh, L. R. and Sherwood, D. R. (2014). Identification of Late Larval Stage Developmental Checkpoints in *Caenorhabditis elegans* Regulated by Insulin/IGF and Steroid Hormone Signaling Pathways. *PLoS Genet.* **10**, e1004426.

Singh, R. N. and Sulston, J. E. (1978). Some observations on molting in *C. elegans*. *Nematologica* **24**, 63–71.

Stempor, P. and Ahringer, J. (2016). SeqPlots - Interactive software for exploratory data analyses, pattern discovery and visualization in genomics. *Wellcome Open Res.* **1**, 14.

Sulston, J. E. and Horvitz, H. R. (1977). Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*. *Dev. Biol.* **56**, 110–156.

Tabara, H., Sarkissian, M., Kelly, W. G., Fleenor, J., Grishok, A., Timmons, L., Fire, A. and Mello, C. C. (1999). The *rde-1* gene, RNA interference, and transposon silencing in *C. elegans*. *Cell* **99**, 123–132.

Taubert, S., Ward, J. D. and Yamamoto, K. R. (2011). Nuclear hormone receptors in nematodes: Evolution and function. *Mol. Cell. Endocrinol.* **334**, 49–55.

Telford, M. J., Bourlat, S. J., Economou, A., Papillon, D. and Rota-Stabelli, O. (2008). The evolution of the Ecdysozoa. *Philos. Trans. R. Soc. B Biol. Sci.* **363**, 1529–1537.

Teuscher, A. C., Jongsma, E., Davis, M. N., Statzer, C., Gebauer, J. M., Naba, A. and Ewald, C. Y. (2019). The *in-silico* characterization of the *Caenorhabditis elegans* matrisome and proposal of a novel collagen classification. *Matrix Biol. Plus* **1**, 100001.

Thummel, C. S. (1990). Puffs and gene regulation-molecular insights into the *Drosophila* ecdysone regulatory hierarchy. *BioEssays News Rev. Mol. Cell. Dev. Biol.* **12**, 561–568.

Tong, A., Lynn, G., Ngo, V., Wong, D., Moseley, S. L., Ewbank, J. J., Goncharov, A., Wu, Y.-C., Pujol, N. and Chisholm, A. D. (2009). Negative regulation of *Caenorhabditis elegans* epidermal damage responses by death-associated protein kinase. *Proc. Natl. Acad. Sci.* 106, 1457–1461.

Tsiairis, C. and Großhans, H. (2021). Chapter Two - Gene expression oscillations in *C. elegans* underlie a new developmental clock. In *Current Topics in Developmental Biology* (ed. Jarriault, S.) and Podbilewicz, B.), pp. 19–43. Academic Press.

Vo, A. A., Levenson, M. T., Ragle, J. M. and Ward, J. D. (2021). Efficient generation of a single-copy *eft-3p::TIR1::F2A:: BFP::AID*::NLS* allele in the *C. elegans ttTi5605* insertion site through recombination-mediated cassette exchange.

MicroPublication Biol. **2021**, 10.17912/micropub.biology.000425.

Vogel, C. and Marcotte, E. M. (2012). Insights into the regulation of protein abundance

from proteomic and transcriptomic analyses. *Nat. Rev. Genet.* **13**, 227–232.

Vuong-Brender, T. T. K., Suman, S. K. and Labouesse, M. (2017). The apical ECM

preserves embryonic integrity and distributes mechanical stress during

morphogenesis. *Development* **144**, 4336–4349.

Ward, J. D., Bojanala, N., Bernal, T., Ashrafi, K., Asahina, M. and Yamamoto, K. R.

(2013). Sumoylated NHR-25/NR5A Regulates Cell Fate during *C. elegans* Vulval

Development. *PLoS Genet.* **9**, e1003992.

Ward, J. D., Mullaney, B., Schiller, B. J., He, L. D., Petnic, S. E., Couillault, C.,

Pujol, N., Bernal, T. U., Van Gilst, M. R., Ashrafi, K., et al. (2014). Defects in

the *C. elegans* acyl-CoA Synthase, *acs-3*, and Nuclear Hormone Receptor, *nhr-*

25, Cause Sensitivity to Distinct, but Overlapping Stresses. *PLoS ONE* **9**,

e92552.

Ward, J. D. (2015a). Rapid and Precise Engineering of the *Caenorhabditis elegans*

Genome with Lethal Mutation Co-Conversion and Inactivation of NHEJ Repair.

Genetics **199**, 363–377.

Ward, J. D. (2015b). Rendering the Intractable More Tractable: Tools from

Caenorhabditis elegans Ripe for Import into Parasitic Nematodes. *Genetics* **201**,

1279–1294.

Watts, J. S., Harrison, H. F., Omi, S., Guenthers, Q., Dalelio, J., Pujol, N. and

Watts, J. L. (2020). New Strains for Tissue-Specific RNAi Studies in

Caenorhabditis elegans. *G3* **10**, 4167–4176.

Zaslaver, A., Mayo, A. E., Rosenberg, R., Bashkin, P., Sberro, H., Tsalyuk, M.,

Surette, M. G. and Alon, U. (2004). Just-in-time transcription program in

metabolic pathways. *Nat. Genet.* **36**, 486–491.

Zhang, C. and Ruvkun, G. (2012). New insights into siRNA amplification and RNAi.

RNA Biol. 9, 1045–1049.

Zhang, L., Ward, J. D., Cheng, Z. and Dernburg, A. F. (2015). The auxin-inducible

degradation (AID) system enables versatile conditional protein depletion in *C.*

elegans. *Development* **142**, 4374–4384.

Zugasti, O. and Ewbank, J. J. (2009). Neuroimmune regulation of antimicrobial

peptide expression by a noncanonical TGF-β signaling pathway in

Caenorhabditis elegans epidermis. *Nat. Immunol.* **10**, 249–256.

1121

FIGURE LEGENDS

Fig. 1. NHR-23 protein oscillates during development. (A) Representative images

from a timecourse monitoring the expression of endogenous NHR-23::GFP protein Two

biological replicates were performed and the number of animals for which the image is

representative at each timepoint is provided. Scale bars=20 µm. (B and C) Anti-FLAG

immunoblot analysis of synchronized *nhr-23::AID**::3xFLAG* animals monitoring the

expression of NHR-23 3XFLAG across two timecourses, 4-28 hours (B) and 22-34

hours (C). Marker size (in kDa) is provided. Stain-free analysis, which visualizes total

protein on the membrane, is provided as a loading control. The blots are representative

of three experimental replicates. The times at which animals were observed to enter

lethargus is indicated in A-C. NHR-23::GFP protein expression in the vulva (D) and

head (E) during the L4 larval stage. Animals were staged based on vulval morphology

(Mok et al., 2015). Images are representative of 20 animals examined over 4

independent experiments. Scale bars=5 µm (D) or 20 µm (E). *nhr-*

23::GFP::AID*::3xFLAG* (A,C,E) and *nhr-23::AID**::3xFLAG* (B,C) are previously

described endogenous knock-ins that produce C-terminal translational fusions to all

known *nhr-23* isoforms (Ragle et al., 2020; Zhang et al., 2015).

1139

Fig. 2. Distinct phenotypes are observed when NHR-23 is depleted early or late in

L1 larvae. (A) Anti-FLAG immunoblot to monitor NHR-23 depletion. Synchronized *nhr-*

23::AID*::3xFLAG; TIR1* and *TIR1* animals were grown on MYOB plates for 12 hours

and then shifted onto control or auxin plates. Lysates were collected at the indicated

time points. Marker size (in kDa) is provided. An anti-alpha-tubulin immunoblot is

provided as a loading control. The blots are representative of three experimental

replicates. (B) Schematic of the experimental set-up. Synchronized L1 larvae were

grown on control media (light grey bands) and transferred to 4 mM auxin (dark grey

bands) every 2 hours. Animals remained on auxin until they were imaged at 20 hours

post-release. (C) Representative images of phenotypes observed when *nhr-*
23::AID*::3xFLAG; TIR1 animals were exposed to auxin in the first larval stage. (D)
Percent of animals of the indicated genotypes with unshed cuticles and morphological
abnormalities following exposure to auxin. The number of animals scored (n) is
provided. (E) Synchronized *nhr-23::AID; TIR1* L1 larvae were released on auxin and
scored for molting defects and death 72 hours post-release. (F) Animals of the indicated
genotype were synchronized and shifted onto auxin at 25 hours post-release and
imaged 23 hours later (48 hours post-release). Animals were staged based on vulval
morphology (Mok et al., 2015). L4 larval stages were grouped as Early L4 (4.0-4.2), Mid
L4 (4.3-4.5), and Late L4 (4.6-4.9). (G) *nhr-23::AID; TIR1* animals were treated as in (F)
and scored after 72 hours on auxin. The number of animals assayed for each genotype
and condition is indicated at the top of the graph in E-G.

1161

Fig. 3. NHR-23 regulates oscillating genes involved in aECM biogenesis. (A) Radar
chart plotting amplitude over the phase of peak expression of *nhr-23*-regulated genes
from Kouns *et al.* (2011). The dotted circles indicated amplitude, with the innermost
circle representing an amplitude of 1 and the outermost circle representing an amplitude
of 4. (B) The data from (A) was plotted as a scatter plot and functionally annotated
(Table S2). We converted the 360º oscillation to developmental timing over a 9 hour
larval stage so that 1 hour=40º. The green shaded area represents NHR-23 expression
based on RNA-seq (peak phase=178.11º, amplitude=2.11; Meeuse et al., 2020),
imaging (Figure 1A,D,E), and western blotting data (Figure 1B,C). (C) The genes in the
aECM component class were subdivided into more specific classes and plotted as in B.
(D) Average signal from NHR-23 ChIP-seq data (Gerstein et al., 2010) for *nhr-23*-
regulated oscillating genes with peak phases between 135º-254.9º (Early), 255-14.9º
(Middle), and 15º-134.9 (Late). *nhr-23* mRNA peaks in expression at 178.11º. Signal is
plotted relative to transcription start site (TSS) and transcription end site (TES). The
average NHR-23 signal for all 21,600 *C. elegans* genes (All genes) and the 20% of
genes with the lowest expression (No expression) are shown for reference. The mean
signal is plotted with a line and the mean's 95% confidence interval is indicated by the
shaded area. (E) Number of NHR-23 peaks flanking and within *nhr-23*-regulated genes.

The genes are binned by their peak phase relative to adjusted larval stage time. Non-
oscillating (Non-Osc) *nhr-23* genes are also depicted.

1182

Fig. 4. NHR-23-depletion causes reduced expression of a *nas-37p::GFP::PEST*
promoter reporter. (A) *nas-37p::GFP::PEST* expression timecourse. Synchronized
nhr-23::AID, TIR1; nas-37p::GFP::PEST L1 larvae were released on control or auxin
plates and scored for head or hypodermal GFP expression every two hours. Fifty
animals per time point were scored in two independent experiments and the percentage
of animals expressing GFP are presented. (B) Representative images of *nhr-23::AID,*
TIR1; nas-37p::GFP::PEST animals grown on control or auxin plates at the indicated
time points. Scale bars=20 µm.

1191

Fig. 5. NHR-23-depletion causes defective NOAH-1 localization. (A) Domain
prediction for NOAH-1. ZP=Zona Pellucida; CFCS=Consensus Furin Cleavage Site.
Predicted size of NOAH-1 and NOAH-1 with an internal mNeonGreen::3xFLAG is
provided in kiloDaltons (kDa). (B) Anti-FLAG immunoblots on lysates from *noah-*
1::mNG::3xFLAG(int) mid-L4 animals An age-matched WT control is included in each
experiment. Marker size (in kDa) is provided and an anti-alpha-tubulin immunoblot was
used as a loading control. (C) Representative images from a *noah-*
1::mNG::3xFLAG(int); mlt-11p::NUC-1::mCherry mid-L4 animal. Arrow indicates cuticle
localization. Scale bars=5 µm. (D) Manders' co-localization analysis of NOAH-
1::mNG::3xFLAG(int) and the lysosome marker NUC-1::mCherry. Three biological
replicates were performed analyzing a total of 29 animals. The horizontal line indicates
median. (E) NOAH-1::mNG expression timecourse through L4; animals were staged by
vulva morphology (Mok et al., 2015). 100 animals were examined over 3 independent
replicates. Scale bars=10 µm (F) Representative GFP and DIC images of the indicated
tissues of *noah-1::mNG::3xFLAG(int)*, *nhr-23::AID; TIR1,* animals grown on control or
auxin plates. NOAH-1::mNG gap over seam cells, and localization to excretory duct and
rectum are indicated by arrows. Two biological replicates were performed; the images
represent 100% of animals scored (n=51 for control and n=44 for auxin). Scale bars=20

μm for all images except the *noah-1::mNG* excretory duct, rectal epithelium, and vulva
(scale bar=5 μm for these images).

1212

Fig. 6. NHR-23-depletion causes ROL-6 and NOAH-1 localization defect and
reduced expression and mislocalization of BLI-1. (A) Cartoon of ROL-6 and BLI-1
domains with knock-in position of mNG::3xFLAG or 3xFLAG::mNG tags. Predicted
protein size is provided in kilodaltons (kDa). (B) Immunoblots on lysates from *rol-*
6::mNG::3xFLAG mid-L4 animals (C) and *bli-1::3xFLAG::mNG(int)* mid-L4 animals (C).
An age-matched WT control is included in each experiment. Marker size (in kDa) is
provided and an anti-alpha-tubulin (B) or anti-actin (C) immunoblot was used as a
loading control. For the blots in (C), the anti-mNeonGreen blot was performed on a
soluble cuticle fraction and the anti-actin control was performed on a soluble
intracellular fraction (see Methods). An arrow indicates the position of the BLI-
1::3xFLAG::mNG band. (D) ROL-6::mNG and (E) BLI-1::mNG expression timecourse
through L4; animals were staged by vulva morphology (Mok et al., 2015). 100 animals
were examined over 3 independent replicates. Scale bars=10 μm. Arrow indicates
fibrous localization pattern in D. (F) Representative images of *nhr-23::AID; TIR1, rol-*
6::mNeonGreen::3xFLAG (*rol-6::mNG*) animals grown on control or auxin plates. Arrow
indicates gap over seam cells. Images from control and auxin-treated animals are
representative of 40/40 animals observed in three independent experiments. (G)
Representative images of *nhr-23::AID; TIR1, rol-6::mNG* animals grown on auxin with a
corset phenotype. Corset is indicated by arrow. 24/40 animals displayed this phenotype.
(H) Representative images of *nhr-23::AID; TIR1, bli-1::mNeonGreen::3xFLAG* (*bli-*
1::mNG) animals grown on control or auxin plates. Three experimental replicates were
performed. Control images are representative of 42/42 animals scored. For auxin-
treated animals, 38/38 exhibited dim expression and 30/38 had patchy BLI-1::mNG
localization (indicated by arrow). Scale bars=20 μm in F-H.

1237

1238 **Fig. 7. Depletion of NHR-23 leads to a defective cuticle barrier.** (A) Permeability to
1239 Hoechst 33258 dye. Synchronized animals of the indicated genotype were shifted onto
1240 control or auxin plates at 25 hours post-release and grown for 23 hours. Animals were

washed off plates and incubated with the cuticle impermeable/membrane permeable Hoechst 33258 dye and scored for nuclear staining in body. The data are from two biological replicates. Asterisk indicates p<0.000001 in two-tailed Student's t-test. (B) Representative images of GFP expression in animals of the indicated genotype following an early and late shift to control or auxin plates. The images are representative of 50 animals observed during each of two experimental replicates. (C) *nhr-23::AID; TIR1* and *TIR1* animals were subjected to hypo-osmotic shock and scored every five minutes. Ten animals were assayed for each genotype and condition with three biological replicates.

Fig. 8. *nhr-23* is necessary in seam cells for developmental progression, molting and BLI-1 localization. (A) Tissue-specific RNAi. A timed egg lay of animals of the indicated strains was performed on control or *nhr-23* RNAi plates and plates were scored three days later. The promoters used to reconstitute RNAi, their tissue-specificity, and oscillation phase (when applicable) is provided. The number of progeny scored from two biological replicates is indicated. Developmental delay was scored as a failure to reach adulthood after 72 hours of growth. Molting defects included animals trapped in cuticles, animals dragging cuticles, and animals with cuticle corsets. (B) L2 *bli-1::mNG* (RNAi competent) and *bli-1::mNG; SCMp::rde-1; rde-1(ne300)* (*SCMp*-specific RNAi) animals were and shifted onto control or *nhr-23* RNAi plates at L2 and grown until stage L4.8. Equal exposure times were used for all images; scale bars=10 μm. Images are representative of 20 animals over two independent experimental replicates.

Figure 1

A

nhr-23::AID*::3xFLAG; TIR1

Hours on auxin: 0   0.25   0.5   12   14   16

76 kDa   anti-FLAG

52 kDa   anti-alpha-tubulin

B

Auxin Exposure Regimen

C

Wild Type (L2) | Arrest | Unshed cuticle | Morphological abnormality

Phenotypic data for nhr-23::AID; TIR1

D

Shift Hour	(n)		% Unshed cuticle		% Morphological abnormality	
	nhr-23::AID; TIR1	TIR1	nhr-23::AID; TIR1	TIR1	nhr-23::AID; TIR1	TIR1
2	40	40	0	0	5	0
4	40	40	28	0	5	0
6	40	31	40	0	18	0
8	40	35	40	0	30	0
10	31	37	58	0	42	0
12	40	20	40	0	55	0
14	39	32	5	0	0	0
16	40	36	5	0	5	0

E L1 shift on auxin+72 hr

F L3 shift on auxin+24 hr

G L3 shift on auxin+72 hr

Figure 2

Figure 3

Figure 4

Figure 5

A

■ Collagen triple helix repeat
■ Nematode cuticle collagen N-terminal domain
■ mNG::3xFLAG or 3xFLAG::mNG

ROL-6 (348 aa; 34.7 kDa)
ROL-6::mNG::3xFLAG (67 kDa)

BLI-1 (948 aa; 105.3 kDa)
BLI-1:3xFLAG::mNG (int) (135.6 kDa)

RXXR site

B

100 kDa
70 kDa
40 kDa

anti-FLAG

50 kDa

anti-alpha tubulin

WT
rol-6::mNG::3xFLAG

C

250 kDa
150 kDa
100 kDa

anti-mNG

← BLI-1::3xFLAG::mNG

50 kDa
37 kDa

anti-actin

WT
bli-1::3xFLAG::mNG(int)

D ROL-6::mNG::3xFLAG

L3
L4.1
L4.3
L4.4

L4.5
L4.6
L4.8
L4.9

E BLI-1::mNG::3xFLAG(int)

L3
L4.1
L4.3
L4.4

L4.5
L4.6
L4.8
L4.9

F

nhr-23::AID; TIR1, rol-6::mNG

Control   Auxin

G

nhr-23::AID; TIR1; rol-6::mNG

ROL-6::mNG   DIC

H

nhr-23::AID; TIR1,bli-1::mNG::3xFLAG(int)

BLI-1::mNG   DIC   BLI-1::mNG   DIC
Control   Auxin

Figure 6

Figure 7

A

Strain	Promoter for tissue-specific RNAi	Tissue-specific RNAi	Oscillation phase	RNAi	N	% molting defects	% developmental delay
N2	N/A	Whole animal	N/A	Control	362	0.00	0.83
				nhr-23	272	100.00	100.00
WM118	*myo-2p*	Body muscle	116.5°	Control	252	0.00	11.51
				nhr-23	320	0.00	7.50
VP303	*nhx-2p*	Intestine	Non-Osc	Control	317	0.00	12.62
				nhr-23	393	0.00	20.10
MGH171	*vha-6p*	Intestine	Non-Osc	Control	602	0.00	0.66
				nhr-23	573	0.00	0.70
JU2039	*lin-31p*	Vulval precursor cells	Non-Osc	Control	325	0.00	2.77
				nhr-23	367	23.43	22.07
QK52	*wrt-2p**	Seam cell, hypodermis, rectum	263.4°	Control	299	0.00	0.33
				nhr-23	198	39.90	90.40
JDW371	*SCMp†*	Seam cell, some hypodermal, intestine	281.5°**	Control	232	0.00	0.00
				nhr-23	153	98.69	99.35
JDW510	*semo-1p*	Hypodermis, intestine	139.8°	Control	259	0.00	2.30
				nhr-23	211	69.20	62.10

*wrt-2 is an nhr-23-regulated gene
† SCMp is comprised of a 573 bp enhancer from the *arf-5* (formerly named *arf-3*) intronic sequence with a *pes-10Δ* minimal promoter
**oscillation phase of *arf-5*; unclear whether the enhancer oscillates in activity

B

bli-1::mNG::3xFLAG(int) (RNAi competent)   *bli-1::mNG::3xFLAG(int)* (SCMp-specific RNAi)

BLI-1::mNG

Figure 8

A

| DIC | DIC + GFP | NHR-23::GFP |

B

| L4.7 | L4.8 | L4.9 | Young adult |

C

| L4.7 | L4.8 | L4.9 | Young adult |

Fig. S1. NHR-23::GFP expression. (A) Representative image of NHR-23 expression in *nhr-23::GFP::AID*::3xFLAG; TIR-1* L4 animals. NHR-23 expression is seen in the hypodermis and developing germline, as described in Ragle et al. 2020. White arrows point to seam cells expressing GFP and yellow arrows point to hyp cells expressing GFP. NHR-23 expression in the vulva (B) and head (C) of *nhr-23::GFP::AID*::3xFLAG* animals of the indicated stages. Images are representative of 20 animals examined over 4 independent experiments. Scale bars=5 μm (B) or 20 μm (C). Animals were staged based on vulval morphology (Mok et al., 2015).

A

B

Predicted protein isoforms expressed during larval stages

DBD **LBD**

NHR-23b/f (553/555aa; 61.3/61.5 kDa)

NHR-23c (437 aa; 48.5 kDa)

NHR-23e (361aa; 39.9 kDa)

Fig. S2. *nhr-23* isoforms detected by Nanopore direct mRNA sequencing. (A) Gene model diagrams of the *nhr-23* isoforms from the "observed isoforms" track on the ENSEMBL genome track. These are isoforms confirmed by direct mRNA sequencing (Roach et al., 2020). The top tracks are of the full *nhr-23* gene, including the *nhr-23c* isoforms, which contain a large first intron. The middle tracks are a zoomed in view starting at the *nhr-23b* transcription start site. The bottom tracks are further zoomed in to depict the alternative 3' splice site usage that generates *nhr-23b* and *nhr-23f*. **(B)** Cartoons of predicted protein isoforms expressed during larval development based on Nanopore direct mRNA sequencing data (Roach et al., 2020). All of the predicted isoforms contain a Ligand Binding Domain (LBD) while only two of the three have a DNA Binding Domain (DBD). The predicted size of each isoform is listed to the right of the structure in amino acids (aa) and kDa. The AID*::3xFLAG fusion is predicted to add 9 kDa.

Fig. S3. NHR-23 depletion causes developmental delay (A) Depiction of the M cell mesoblast cell divisions in L1 larvae created with BioRender.com. Newly hatched animals are born with a single M cell, which undergoes the indicated set of divisions to produce 16 descendent cells at the L1/L2 molt. An *hlh-8p::GFP* reporter can be used to monitor M cell divisions in living animals. (B and C) Animals of the indicated genotype were synchronized and shifted to either auxin or control plates at either 3 (early shift) or 9 hours (late shift) post-release and imaged at 24 hours post-release. Two biological replicates were performed. (B) Representative images of animals of the indicated genotypes on auxin plates. (C) M cell quantification for animals of the indicated genotype shifted to control or auxin plates. Mean and standard deviation are labeled. The number of animals assayed for each genotype and condition is indicated at the top of the graph (* two-tailed T-test p<0.0001).

A

Wild type   Molting defective   Dead/necrotic

B

24 hr on control or auxin

TIR1   *nhr-23::AID; TIR1*   *TIR1*   *nhr-23::AID; TIR1*

Control   Auxin

C

nhr-23::AID; TIR1 animals 72 hr on auxin

Adult   Indeterminable L4 with abnormal vulva lumen   Adult with abnormal vulva lumen   Dead/necrotic

Fig. S4. NHR-23 depletion causes developmental delay. (A) Representative images of synchronized *nhr-23::AID; TIR1* L1 larvae released on auxin and scored for molting defects and death/necrosis 72 hours post-release. (B) Representative images of animals of the indicated genotype that were synchronized and shifted onto auxin at 25 hours post-release and imaged 23 hours later (48 hours post-release). The position of the vulva is highlighted with a white box. (C) Representative images of *nhr-23::AID; TIR1* animals treated as in (B) and scored after 72 hours on auxin. Scale bars=20 µm. A zoomed inset is provided for the auxin-treated *nhr-23::AID, TIR1* images in B and C to highlight vulval morphology.

Fig. S5. NHR-23 enrichment around genes. Average signal over all aligned genes from two NHR-23 ChIP-seq replicates (Gerstein et al., 2010). The y-axis indicates non-normalized read coverage. The mean signal is plotted with a line and the mean's 95% confidence interval is indicated by the shaded area. TSS=transcription start site; TES=transcription end site.

A

Strain	Genotype	Brood size (±std. dev.)
N2	Wild type	290±48
JDW459	*nhr-23::AID; TIR1, rol-6::mNG::3xFLAG*	293±48
JDW463	*nhr-23::AID; TIR1, bli-1::mNG::3xFLAG(int)*	236±56
JDW475	*noah-1::mNG::3xFLAG(int), nhr-23::AID; TIR1*	246±27

B

bli-1::3xFLAG::mNG(int)   *bli-1::mNG::3xFLAG(int)*   *nhr-23::AID; TIR1, bli-1::mNG::3xFLAG(int)*

Fig. S6. Validation of *bli-1, noah-1,* and *rol-6* knock-in strains. (A) Brood size analysis of animals of the indicated genotype. (B) Representative images from L4.9 stage CZ27591 *bli-1(ju1789[bli-1::3xFLAG::mNG(int)]) II*, JDW389 *bli-1(wrd84[bli-1::mNG::3xFLAG(int)]) II*, and JDW463 *nhr-23(kry61(nhr-23::AID*-TEV-3xFLAG)) I; ieSi57 [Peft-3::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)], bli-1(wrd84[bli-1::mNG::3xFLAG(int)])* animals. Equal 500 msec exposures from the same area of animals were taken using a 100x objective. Knock-ins are all at the same location in *bli-1* (see Fig. 6A). Scale bars=10 µm. Images are representative of 40 animals examined in two independent experiments.

Fig. S7. Depletion of NHR-23 leads to a elevated *nlp-29p::GFP* reporter activity. Anti-GFP immunoblot analysis of *nlp-29::*GFP expression. Marker size (in kDa) is provided. An anti-tubulin immunoblot is provided as a loading control. The blots are representative of two experimental replicates.

Fig. S8. Expression patterns of *SCMp* and *wrt-2* promoter reporters. *wrt-2p::NLS::mScarlet* is expressed in the hypodermis and seam cells (A and B). In some animals seam cell expression is lacking. In A, arrows point to two seam cells, one with reporter expression and one lacking it. *SCMp::NLS::mScarlet* promoter reporters express robustly in the seam cells (C) with weak expression in the hypodermis (D) in a subset of animals. Two biological replicates were performed and the number of animals for which each image is representative is indicated.

Control

HIS-72::mNeonGreen

mNeonGreen
RNAi

HIS-72::mNeonGreen

Fig. S9. JDW525 animals are permissive for RNAi knockdown in seam, hypodermal, and intestinal cells. JDW 525 [*his-72(wrd142[mNeonGreen::his-72]) III ; jsTi1493 {mosL loxP [wrdSi65(SCMp::rde-1 CDS+3'UTR)] FRT3::mosR} IV ; rde-1 (ne300) V*] animals grown on control RNAi display HIS-72::mNG expression in hypodermal, intestinal, germline, vulval, and seam cells. mNeonGreen RNAi caused reduction of HIS-72::mNG expression in seam, intestinal, and syncytial hypodermal (hyp7) cells. Vulval, germline, and head hypodermal expression was unaffected. Scale bars=20μm. Data are representative of forty animals examined over two biological replicates