1 **Assessing long-read sequencing with Nanopore R9, R10 and PacBio CCS to obtain**
2 **high-quality metagenome assembled genomes from complex microbial**
3 **communities**

4 Mantas Sereika[a*], Rasmus Hansen Kirkegaard[a*], Søren Michael Karst[a], Thomas Yssing Michaelsen[a],
5 Emil Aare Sørensen[a] and Mads Albertsen[a**]
6
7 [a]Center for microbial communities, Aalborg University, Denmark
8

9 *These authors contributed equally to the paper
10 **Corresponding author ma@bio.aau.dk

Abstract

Short-read DNA sequencing has led to a massive growth of genome databases but mainly with highly

fragmented metagenome assembled genomes from environmental systems. The fragmentation is

a result of closely related species, strains, and genome repeats that cannot be resolved with short

reads. To confidently explore the functional potential of a microbial community, high-quality

reference genomes are needed. In this study, we evaluated the use of different combinations of

short (Illumina) and long-read technologies (Nanopore R9.4, R10.3, and PacBio CCS) for recovering

high-quality metagenome assembled genomes (HQ MAGs) from a complex microbial

community (anaerobic digester). Depending on the sequencing approach, 33 to 86 HQ MAGs

(encompassing up to 34 % of the assembly and 49 % of the reads) were recovered using long reads,

with Nanopore R9 featuring the lowest sequencing costs per HQ MAG recovered. PacBio CCS was

also found to be an effective platform for genome-centric metagenomics (74 HQ MAGs) and

produced HQ MAGs with the lowest fragmentation (median of 9 contigs) as a stand-alone

technology. Using PacBio CCS MAGs as reference, we show that, although a high number of high-

quality MAGs can be generated using Nanopore R9, systematic indel errors are still present, which

can lead to truncated gene calling. However, polishing the Nanopore MAGs with short-read Illumina

data, enabled recovery of MAGs with similar quality as MAGs from PacBio CCS.

Importance

Multiple approaches exists for performing genome-centric metagenomics to recover high-quality

genomes from microbial communities. The plethora of possible sequencing strategies can be

cumbersome for designing metagenomics projects, especially under limited budget conditions.

Here, we performed single sequencing runs of the same complex microbial community sample using

Illumina MiSeq, Oxford Nanopore MinION and PacBio Sequel II to assess the performance of each

platform. We show that, in general, long read sequencing significantly outperforms the short read

Illumina platform for recovering microbial genomes. In addition, we observed that the hybrid

Nanopore-Illumina approach recovers genomes of comparable quality to PacBio CCS, while

maintaining lower sequencing costs per genome recovered. For this reason, we find the

Nanopore R.9.4.1 and supplemental Illumina read polishing to be, at the moment of writing, the

most cost-effective sequencing strategy for *de novo* acquisition of high-quality genomes.

Introduction

Bacteria live in almost every environment on Earth and a recent analysis of the global microbial diversity estimated a total of 10^{12} species (1). To obtain representative genomes culturing has been used to isolate specific microbes, but the throughput is highly limited. Recently, genome recovery from metagenomes with short, high-quality reads has drastically improved genome recovery for uncultivated species (2–4), and has spurred multiple tools for automated binning of draft genomes (5–8). This has led to studies reporting thousands or even hundreds of thousands of genomes (9–13). However, the recovered metagenome assembled genomes (MAGs) are often very fragmented, incomplete, and contaminated with genome fragments from other microbes. Hence, even though the pace of genome sequencing is increasing, only a tiny fraction of the estimated microbial diversity have representative genomes (14) with 47,894 species having representative genomes in the Genome Taxonomy Database (GTDB) version 202 (15).

Recovering genomes from metagenomes using short reads is challenging due to the presence of genetic repeat elements. From an assembly point-of-view, repeat elements are classified as identical DNA segments that are not able to be spanned by the length of the sequenced reads. For example, the 16S ribosomal RNA gene is highly conserved between species, can be present in multiple copies within species, and is much longer than the typical short reads. This either breaks up the assembly graph or produce chimeric segments (16, 17). In complex microbial communities a large source of repeat elements originates from closely related microbial strains that share large parts of their genomic content. The repeat elements have a coverage that is different from the rest of the genome, and are thus often not picked up by automated binning tools (11). In addition,

genomes can also be contaminated with fragments from other organisms that just happen to

correlate with the binned genome (11, 14, 18). Tools have been developed to estimate the

completeness and the level of contamination and even "decontaminate" the bins (16, 18). However,

they rely heavily on good reference databases and assumptions about universal essential single copy

genes that are known not to be truly universal (19, 20). Maybe the only way to be certain about the

completeness of a genome is to produce closed MAGs as in the case of circular chromosomes (21).

This is especially important as repeat regions may contain biological information that is essential to

understand a microbial community (22), as have been shown for some antimicrobial resistance

genes (23).

Long-range information is, to a large degree, capable of solving the genome fragmentation caused

by repeat elements (24). Different approaches have been developed to generate such data, e.g.

mate-pair sequencing, synthetic long-read sequencing, and long-read single molecule sequencing,

such as offered by Pacific Biosciences sequencing (PacBio) and Oxford Nanopore Technologies (25–

32). All of these techniques have been used to recover genomes from metagenomes in

environments of varying complexity and recent technological breakthroughs deliver vastly

increased data yields, which makes even higher complexity samples tractable (12, 33–38). However,

long read sequencing technologies feature drawbacks of their own, as the PacBio platform is

relatively costly (39, 40), while Nanopore sequencing exhibits systematic errors in homopolymer

regions that can lead to insertions and deletions in the assemblies (41, 42).

Here, we assess the recent DNA sequencing technologies in the context of genome-centric, high-

throughput metagenomics under realistic scenarios for a typical small research project. We compare

metagenome assembly, binning, rRNA and protein gene recovery between single flow cell

sequencing runs of Illumina MiSeq and Oxford Nanopore MinION R.9.4.1 and R.10.3 (referred to as

R9 and R10 hereafter) and PacBio Sequel II (circular consensus sequencing).

Results and discussion

DNA of a microbial community from an anaerobic digester was sequenced via Illumina MiSeq (2x300

bp), PacBio CCS and the Oxford Nanopore MinION platform, using R9 and R10 chemistry flow cells

(**Fig 1a**). The reads from different sequencing platforms were then assembled using Megahit for

short reads and metaFlye for long reads. Despite being the same sample of extracted DNA, direct

comparisons are difficult as the additionally size selection of the PacBio CCS dataset both increased

the read length (**Fig 1b**) and altered the relative abundances of the species in the sample (**Fig S1a**).

The Illumina data also features variation in relative abundances (**Fig S1b**), presumably due to GC

bias (43). Only the R9 and R10 relative abundance data correlated well (**Fig S1c**), even though they

had a 2.7-fold difference in sequencing yield (**Table 1**). Hence, the comparisons are influenced by

differences in sequencing depth, read length, library preparation technique (44) and changes to

relative abundances of the community members, but still represent a realistic scenario for a typical

research project.

Table 1. General statistics for sequenced reads and assembled contigs.

Data type	Feature	Illumina	Nanopore R9	Nanopore R10	PacBio CCS
Reads	Total count	47,091,904	10,266,261	3,646,771	992,914
	Total yield (Mb)	13,285	35,237	13,009	15,309
	N50 (kb)	0.3	5.9	6.4	15.4
	Modal read accuracy (%)*	99.98	96.34	95.81	99.97
Contigs**	Total count	145,876	24,680	13,132	8,989
	Circular, > 0.5 Mb count	0	7	5	9
	Total size (Mb)	409	754	395	606
	N50 (kb)	3.5	79.9	71.7	172.5
	Reads mapped to contigs (%)	88.1	93.5	93.2	95.2

*Derived from read Phred scores.
**Assembly metrics are presented after removing < 1 kb contigs.

To assist automated contig binning, we performed Illumina sequencing of 9 additional samples from

the same anaerobic digester spread over 9 years (**Supp Table 1**) and used the coverage profiles as

input for binning. Furthermore, to evaluate the impact of micro-diversity on MAG quality, we

calculated the polymorphic site rates for each MAG as a simple proxy for presence of micro-

diversity (10).

After performing automated contig binning it is evident that micro-diversity has a large impact on

MAG fragmentation, but that long-read sequencing results in much less fragmentation (**Fig 1c-d**) of

bins at higher amounts of micro-diversity (**Fig 2a**). Despite large differences in read length for

Nanopore and PacBio CCS data (N50 read length 6 kbp vs. 15 kbp), only small differences in bin

fragmentation were observed, as compared to the Illumina-based results.

After performing bin quality estimates, the most HQ MAGs were recovered by using Nanopore

sequencing on a R9 flow cell (n=86), supplemented with Illumina read polishing, while also featuring

the second lowest laboratory costs per recovered HQ MAG (**Table 2**). The second highest number

of HQ MAGs (n=74) was achieved via PacBio CCS, which also exhibited the least HQ MAG

fragmentation in terms median contigs per bin (n=9), although the overall sequencing costs per

HQ MAG recovered were greater, compared to other long read approaches.

Table 2. Results for automated contig binning. Genome quality was defined based on the MiMAG
(14) standard using CheckM (18): HQ MAGs are defined by completeness of >90 %, <5 %
contamination, at least 18 distinct tRNA genes and the 5S, 16S, 23S rRNA genes occurring at least
once. MQ MAGs by completeness >50 % and contamination <10 %, while low quality MAGs are
defined by completeness <50 % and contamination <10 %. MAGs with contamination estimates
higher than 10 % were classified as contaminated. *Fraction of the assembled contigs (> 1kb) that
were placed into bins. **Sequencing costs refers to the expenses encountered at the time of
conducting the experiments and may differ for other research groups, depending on agreements
with sequencing service providers.

Feature	Illumina MiSeq	Nanopore R9	Nanopore R9 + Illumina	Nanopore R10	Nanopore R10 + Illumina	PacBio CCS
HQ MAGs	8	64	86	33	45	74
MQ MAGs	83	114	95	64	63	72
LQ MAGs	3	28	26	18	12	22
Contaminated MAGs	10	6	13	6	6	14
Reads binned (%)	76	86	85	86	86	83
Contigs binned (%)*	55	68	71	71	74	73
Contigs per HQ MAG (median)	184	15	16	15	18	9
Contigs binned in HQ MAGs (%)	4	22	30	22	29	34
Reads binned in HQ MAGs (%)	16	46	49	39	41	48
Sequencing costs ($)**	1,200	811	2,011	811	2,011	4,420
Cost per HQ MAG ($)	150	13	23	25	45	60

After further examining MAG quality estimates between the clustered bins (see Materials and

methods for clustering details) of different sequencing platforms, MAGs from PacBio CCS were

found to feature greater mean bin completeness estimates (**Fig 2b**) than the Illumina-only

approach (93.3 ± 6.2 vs. 89.3 ± 8.2, respectively, paired t-test p=0.0001). Furthermore, polishing of

Nanopore assemblies with Illumina reads was observed to slightly improve the mean bin

completeness for both R9 (from 91.9 ± 5.4 to 93.5 ± 4.7 %, p=0.0009) and R10 (from 91.9 ± 6.0 to

93.1 ± 5.9 %, p=0.021) chemistries. The observed increase in bin completeness from Illumina read

polishing is assumed to be caused by improved recovery of protein sequences (bin completeness

estimation is based on the presence of marker proteins (18)), due to correction of Nanopore

sequencing systematic errors, such as homopolymer truncation that can artificially induce

frameshifts, and thus introduce early stop codons in genes (45, 46).

151
Figure 1. Overview of general sequencing results from short and long read platforms. a) Read
length and estimated accuracy (inferred from Phred scores, log-scaled) distributions of the read
datasets. **b)** Contig Nx plot for the assembled metagenomes (after 1 kb length filtering). Differential
coverage plots are presented for **c)** Illumina and **d)** Nanopore R9 metagenome assemblies.

Although Illumina read polishing increases the overall completeness estimates of Nanopore-

assembled bins, the improvements are case-specific, due to differences in coverage and

homopolymer content and the degree of micro-diversity between MAGs. To investigate this further,

PacBio CCS bins were assumed as "reference quality" and compared to bins from Nanopore

sequencing with and without short read polishing (see Materials and Methods for selection criteria).

Applying Illumina read polishing was found to reduce the rate of indels (by at least 20 %) in 18 of 18

Nanopore R9 bins and 14 of 15 R10 bins, independent of coverage (**Fig 2c**). Interestingly, the

estimated reduction in mismatches (at least 20 %) was less pronounced, as it was observed in 6 of

18 Nanopore R9 bins and 8 of 15 for R10, presumably due to Nanopore R10 bins featuring overall

lower long read coverage (**Fig 2d**). Hence, even though it is possible to acquire HQ MAGs from

Nanopore-only assemblies, as demonstrated in this and other studies (47, 48), supplemental

Illumina read polishing still aids in recovering more HQ MAGs (**Table 2**) from complex microbial

communities as well as producing presumably less erroneous genome sequences.

Figure 2. Comparison of bins from different sequencing approaches. a) MAG fragmentation (log-
scaled) at different bin SNP rates in PacBio CCS MAGs. **b)** Genome bin completeness estimates for
different sequencing platforms. IL — Illumina, NP — Nanopore, PBCCS — PacBio CCS. Bin **c)** indel
and **d)** mismatch rates (log-scaled) for MAGs from Nanopore sequencing with and without Illumina
read polishing, compared to MAGs from PacBio CCS. The presented bin coverage on the x axis (log-
scaled) is for the corresponding Nanopore chemistry type. HQ MAGs are represented by circle, while
triangles denote MQ MAGs. For all figures, only the bins that were clustered together between all
the different sequencing platforms (see Materials and methods) are presented.

Reduction of indel rates in genomes is expected to yield less truncated protein sequences (49). To

investigate this we used the IDEEL test (12) that estimates protein length by comparing to known

sequences. In the MAGs using Nanopore R9 or R10 data alone, respectively 15 % and 14 % of protein

sequences were estimated to exhibit a greater than 20 % truncation, while proteins from the hybrid

Nanopore and Illumina approach were found to be of comparable length to that of PacBio CCS (3-

4 %, **Table 3**). Furthermore, Illumina-only bins were observed to feature slightly increased estimated

protein truncation levels than PacBio CCS and the hybrid approach, although the following is

expected to be influenced by the generally shorter contig lengths, leading to a higher count of

fragmented protein sequences (50).

Table 3. Protein sequence distribution (%) at different estimated truncation rates.

Query-to-Subject length ratio intervals	Illumina	Nanopore R9	Nanopore R9 + Illumina	Nanopore R10	Nanopore R10 + Illumina	PacBio CCS
1.0-0.8	90.89	85.01	96.69	85.56	96.37	96.38
0.8-0.6	3.03	4.51	1.30	4.51	1.40	1.41
0.6-0.4	2.48	4.40	0.90	4.35	1.00	0.97
0.4-0.2	2.29	4.21	0.77	3.91	0.83	0.87
0.2-0.0	1.31	1.87	0.33	1.66	0.40	0.37

A known source of errors in Nanopore-based assemblies is systematic homopolymer miscalling.

Similarly to the IDEEL test, using MAGs from PacBio CCS as a reference, Nanopore-only MAGs were

found to feature increased homopolymer errors at longer lengths, especially for cytosine and

guanine homopolymers (**Fig S3**), which coincides with read-level error rates of Nanopore

sequencing (51). As expected, sequences from Nanopore R10 featured lower rates of homopolymer

miscalling than R9 and the hybrid assemblies featured reduced error profiles, similar to that of

Illumina-only assemblies.

To reiterate, although the benefits of the hybrid Nanopore and Illumina approach have been

reported in numerous projects (52–59), our study utilized PacBio CCS as a means of establishing

reference sequences for examining the effects of short read polishing on Nanopore bins that were

recovered from a complex microbial community. Given that short read length can be a cause for

mapping issues, often leading to errors and biases in genome recovery (60–62), we observed

evidence that Illumina read polishing of Nanopore bins still significantly reduces the estimated rate

of indels, especially homopolymer errors, in a high complexity metagenome. Furthermore, although

multiple bioinformatics workflows for frame-shift correction of Nanopore assemblies without short

reads have been developed (48, 63–65), the indel corrections are not *de novo*, as the workflows are

based on comparisons to reference databases, and hence erroneous sequences of novel genes

might not get corrected or proteins with biologically occurring frame-shift mutations might be

falsely converted to a full-length state, making the method suboptimal for characterising novel

microbial species. Nonetheless, using databases to correct Nanopore sequencing errors could still

be a useful and cost-effective approach, when *de novo* recovery of novel genomic sequences is not

relevant for the research project.

Lastly, the presence of repetitive DNA sequences, including genes in multiple copies, can lead to

contig breaks in the assembly (66). Ribosomal RNA genes were chosen as an example of repeat gene

as it is also the most widely used marker gene for amplicon sequencing, phylogenetic analysis and

FISH (67–69). Hence, sufficient recovery and binning of rRNA genes is an important aspect of high

throughput genome-centric metagenomics in order to connect MAGs to additional information.

Comparing 44 MAGs captured in all assemblies (see Materials and Methods), 13 out of 44 Illumina

bins featured all rRNA genes, while it was 40 for Nanopore R9, 41 for Nanopore R10 and 41 for

PacBio CCS (**Fig S4 a-c**). In the Illumina assembly, the ratio of 16S to 23S rRNA gene counts was

0.76 (**Fig S4 d-e**) indicating either collapsed or fragmented genes as a result of micro-diversity,

whereas for long-read technologies a comparable ratio of ~1 was observed (**Supp Table 2**).

Conclusion

Long read DNA sequencing improves assembly contiguity over short read assemblies and, as a result,

produces significantly greater numbers of high-quality genomes from complex microbial

communities. Recovery of repeated rRNA genes via long read sequencing is also vastly improved

over the short read approach. Nevertheless, short read sequencing is still useful, as an economical

way of acquiring time series data for binning MAGs and it can be combined with Nanopore

sequencing to recover more high-quality genome bins and full protein sequences from

metagenomes. The PacBio CCS platform features overall superior read metrics in terms of accuracy

and length, although the relatively higher sequencing costs can act as a bottleneck for large scale

genome-centric metagenomics projects. Despite the multiple differences between Nanopore and

PacBio CCS platforms, recovering genomes from complex metagenomes via Nanopore R9

sequencing with supplemental short read Illumina polishing was, at the moment of writing, found

to be the optimal strategy, balancing laboratory costs, adequate sequencing depth and minimal

sequencing errors in the assembled genomes.

Materials and methods

Sampling

Sludge biomass was sampled from the anaerobic digester at Fredericia wastewater treatment plant (Latitude 55.552219, Longitude 9.722003) at multiple time points and stored as frozen 2 mL aliquots at -20°C.

DNA extraction

DNA was extracted from the anaerobic digester sludge using DNeasy PowerSoil Kit (QIAGEN, Germany) following the manufacturer's protocol. The extracted DNA was then size selected using the SRE XS (Circulomics, USA), according to the manufacturer's instructions.

DNA QC

DNA concentrations were determined using Qubit dsDNA HS kit and measured with a Qubit 3.0 fluorimeter (Thermo Fisher, USA). DNA size distribution was determined using an Agilent 2200 Tapestation system with genomic screentapes (Agilent Technologies, USA). DNA purity was determined using a NanoDrop One Spectrophotometer (Thermo Fisher, USA).

Nanopore DNA sequencing

Library preparation was carried out using the SQK-LSK109 kit (Oxford Nanopore Technologies, UK) following the manufacturer's protocol. The DNA libraries were sequenced using the R.9.4.1 and the R.10.3 MinION flowcells (Oxford Nanopore Technologies, UK) on a MinION Mk1B (Oxford Nanopore Technologies, UK) device. After sequencing, the MinION flowcells were washed using the Flow Cell Wash Kit (EXP-WSH002, Oxford Nanopore Technologies, UK) and the same library was loaded again to generate additional sequencing data.

Illumina DNA sequencing

The Illumina libraries were prepared using the Nextera DNA library preparation kit (Illumina, USA) following the manufacturer's protocol and sequenced using the Illumina MiSeq platform.

PacBio CCS

Size-selected DNA sample was sent out to the DNA Sequencing Center at Brigham Young University. The DNA sample was fragmented with Megaruptor (Diagenode, Belgium) to 15 kb and size-selected using the Blue Pippin (Sage Science, USA). The sample was then prepared using SMRTbell Express Template Preparation Kit 1.0 (PacBio, USA) according to manufacturers' instructions. Sequencing was performed on the Sequel II system (PacBio, USA) using the Sequel II Sequencing Kit 1.0 (PacBio, USA) with the Sequel II SMRT Cell 8M (PacBio, USA) for a 30 hour data collection time.

Data analysis

Read processing

Illumina reads were trimmed for adapters using Cutadapt (v. 1.16 (70)). The generated raw Nanopore data was basecalled in super-accurate mode with using Guppy (v. 5.0.7, https://community.nanoporetech.com/downloads) with dna_r9.4.1_450bps_sup.cfg model for R9 and dna_r10.3_450bps_sup.cfg model for R10 chemistry. Adapters for nanopore reads were removed using Porechop (v. 0.2.3 (71) and read with Phred quality scores below 7 were filtered out using Filtlong (v. 0.1.1, https://github.com/rrwick/Filtlong). The CCS tool (v. 6.0.0, https://ccs.how/) was utilized with the sub-read data from PacBio CCS to produce HiFi reads. Read statistics were acquired via NanoPlot (v. 1.24.0 (72))

Metagenome assembly and binning

Long reads were assembled using Flye (v. 2.9-b1768 (73)) with the "--meta" setting enabled and the

"--nano-hq" option for assembling Nanopore reads, whereas "--pacbio-hifi" and "--min-

overlap 7500 --read-error 0.01" options were used for assembling PacBio CCS reads, as it resulted

in more HQ MAGs than using the default settings. Polishing tools for Nanopore-based assemblies:

Minimap2 (v. 2.17 (74)), Racon (used thrice, v. 1.3.3 (75)), and Medaka (used twice, v. 1.4.1,

https://github.com/nanoporetech/medaka). Nanopore assemblies were additionally polished with

Illumina reads using Racon. The trimmed Illumina reads were also assembled using Megahit (v.

1.1.4 (76))

Automated binning was carried out using MetaBAT2 (v. 2.12.1 (77)), with "-s 500000" settings,

MaxBin2 (v. 2.2.7) and Vamb (v. 3.0.2) with "-o C --minfasta 500000" settings. Contig coverage

profiles from different sequencer data as well as 9 additional time series Illumina datasets of the

same anaerobic digester were used for generating the bins. The binning output of different tools

was then integrated and refined using DAS Tool (v. 1.1.2 (78)). CoverM (v. 0.6.1,

https://github.com/wwood/CoverM) was applied to calculate the bin coverage ("-m mean"

settings) and relative abundance ("-m relative_abundance") values.

Bin processing

The completeness and contamination of the genome bins was estimated using CheckM (v.

1.1.2 (18)). The bins were classified using GDTB-Tk (v. 1.5.0 (79), R202 database). Bin protein

sequences were predicted using Prodigal (v. 2.6.3, https://github.com/hyattpd/Prodigal) with

"-p meta" setting, while rRNA genes were predicted using Barrnap (v. 0.9,

https://github.com/tseemann/barrnap) and tRNAscan-SE (v. 2.0.5 (80)) was used for tRNA

predictions. Bin quality was determined following the Genomic Standards Consortium guidelines, wherein a MAG of high quality featured genome completeness of more than 90 %, less than 5 % contamination, at least 18 distinct tRNA genes and the 5S, 16S, 23S rRNA genes occurring at least once (14). MAGS with completeness above 50 % and contamination below 10 % were classified as medium quality, while low quality MAGs featured completeness below 50 % and contamination below 10 %. MAGs with contamination estimates higher than 10 % were classified as contaminated.

Illumina reads were mapped to the metagenome assembly using Bowtie2 (v. 2.4.2 (81)) with the "--very-sensitive-local" setting. The mapping was converted to BAM and sorted using SAMtools (v. 1.9 (82)). Single nucleotide polymorphism rate was then calculated using CMseq (v. 1.0.3 (10)) from the mapping using poly.py script with "--mincov 10 --minqual 30" settings.

Bins were clustered using dRep (v. 2.6.2 (83)) with "-comp 50 -con 10 -sa 0.95" settings. Only the bins that featured higher coverage than 10 in their respective sequencing platform and a higher Illumina read coverage than 5 for bins from the hybrid approach were included in downstream analysis. For IDEEL test (12), the predicted protein sequences from clustered bins were searched against the UniProt TrEMBL (84) database (release 2021_01) using Diamond (v. 2.0.6 (85)). Query matches, which were not present in all datasets, were omitted to reduce noise.

QUAST (v. 4.6.3 (86)) was applied on the clustered bins with less than 0.5 % SNP rate to acquire mismatch and indels metrics. Cases with Quast parameters "Genome Fraction" of less than 75 % and "Unaligned length" of more than 250 kb were omitted to reduce noise. For homopolymer analysis, the clustered bins were mapped to each other using "asm5" mode of Minimap2 and

333   Counterr (v. 0.1, https://github.com/dayzerodx/counterr) was used on the mapping files to get

334   homopolymer calling errors. For QUAST and Counterr, PacBio CCS bins were used as reference

335   sequences.

Data availability

337   The sequencing data and bins are available at the ENA with bio project ID: PRJEB48021. The code,

338   datasets used to generate the figures and additional material are available at

339   https://github.com/Serka-M/Digester-MultiSequencing.

Acknowledgements

341   We would like to acknowledge the plant operators at Fredericia wastewater treatment plant for

342   supplying the sample material.

Funding information

345   The study was funded by research grants from VILLUM FONDEN (15510) and the Poul Due Jensen

346   Foundation (Microflora Danica).

Author contributions

348   MS performed DNA extraction and size selection. RHK did Illumina and Nanopore sequencing of the

349   sample. RHK and MS performed the bioinformatics. MS, RHK and MA wrote the manuscript. SMK,

350   TYM and EAS contributed to experiment design, result interpretation and writing of the manuscript.

351   All authors reviewed the manuscript.

Conflict of interest

355   EAS, SMK, MA, RHK are also employed at DNASense ApS. TYM is employed at Lyras A/S. The

356   remaining authors declare no conflict of interest.

References

1.   Locey KJ, Lennon JT. 2016. Scaling laws predict global microbial diversity. Proceedings of the

National Academy of Sciences of the United States of America 113:5970–5.

2.   Tyson GW, Chapman J, Hugenholtz P, Allen EE, Ram RJ, Richardson PM, Solovyev V V, Rubin

EM, Rokhsar DS, Banfield JF. 2004. Community structure and metabolism through

reconstruction of microbial genomes from the environment. Nature 428:37–43.

3.   Sharon I, Morowitz MJ, Thomas BC, Costello EK, Relman DA, Banfield JF. 2013. Time series

community genomics analysis reveals rapid shifts in bacterial species, strains, and phage

during infant gut colonization. Genome research 23:111–20.

4.   Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. 2013.

Genome sequences of rare, uncultured bacteria obtained by differential coverage binning of

multiple metagenomes. Nature biotechnology 31:533–8.

5.   Wu Y, Tang Y, Tringe SG, Simmons BA, Singer SW. 2014. MaxBin: an automated binning

method to recover individual genomes from metagenomes using an expectation-

maximization algorithm. Microbiome 2:26.

6.   Kang DD, Froula J, Egan R, Wang Z. 2015. MetaBAT, an efficient tool for accurately

reconstructing single genomes from complex microbial communities. PeerJ 3:e1165.

7.   Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, Lahti L, Loman NJ,

Andersson AF, Quince C. 2014. Binning metagenomic contigs by coverage and composition.

Nature methods 11:1144–6.

8.   Miller IIJ, Rees ER, Ross J, Miller IIJ, Baxa J, Lopera J, Kerby RL, Rey FE, Kwan JC. 2019.

Autometa: automated extraction of microbial genomes from individual shotgun

metagenomes. Nucleic acids research 47:e57.

9.   Nayfach S, Shi ZJ, Seshadri R, Pollard KS, Kyrpides NC. 2019. New insights from uncultivated

genomes of the global human gut microbiome. Nature 568:505–510.

10.   Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, Beghini F, Manghi P, Tett A,

Ghensi P, Collado MC, Rice BL, DuLong C, Morgan XC, Golden CD, Quince C, Huttenhower C,

Segata N. 2019. Extensive Unexplored Human Microbiome Diversity Revealed by Over

150,000 Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. Cell

176:649-662.e20.

11.   Parks DH, Rinke C, Chuvochina M, Chaumeil P, Woodcroft BJ, Evans PN, Hugenholtz P, Tyson

GW. 2017. Recovery of nearly 8,000 metagenome-assembled genomes substantially

expands the tree of life. Nature microbiology 2:1533–1542.

12.   Stewart RD, Auffret MD, Warr A, Walker AW, Roehe R, Watson M. 2019. Compendium of

4,941 rumen metagenome-assembled genomes for rumen microbiome biology and enzyme

discovery. Nature biotechnology 37:953–961.

13.   Singleton CM, Petriglieri F, Kristensen JM, Kirkegaard RH, Michaelsen TY, Andersen MH,

Kondrotaite Z, Karst SM, Dueholm MS, Nielsen PH, Albertsen M. 2021. Connecting structure

to function with the recovery of over 1000 high-quality metagenome-assembled genomes

from activated sludge using long-read sequencing. Nat Commun 12:2009.

14.   Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-smith M, Doud D, Reddy TBK, Schulz F,

Jarett J, Rivers AR, Eloe-fadrosh EA, Tringe SG, Ivanova NN, Copeland A, Clum A, Becraft ED,

Malmstrom RR, Birren B, Podar M, Bork P, Weinstock GM, Garrity GM, Dodsworth JA,

Yooseph S, Sutton G, Glöckner FO, Gilbert JA, Nelson WC, Hallam SJ, Jungbluth SP, Ettema

TJG, Tighe S, Konstantinidis KT, Liu WT, Baker BJ, Rattei T, Eisen JA, Hedlund B, McMahon

KD, Fierer N, Knight R, Finn R, Cochrane G, Karsch-Mizrachi I, Tyson GW, Rinke C, Lapidus A,

Meyer F, Yilmaz P, Parks DH, Eren AM, Schriml L, Banfield JF, Hugenholtz P, Woyke T. 2017.

Minimum information about a single amplified genome (MISAG) and a metagenome-

assembled genome (MIMAG) of bacteria and archaea. Nature biotechnology 35:725–731.

15.   Parks DH, Chuvochina M, Chaumeil P-A, Rinke C, Mussig AJ, Hugenholtz P. 2020. A complete

domain-to-species taxonomy for Bacteria and Archaea. Nat Biotechnol 38:1079–1086.

16.   Tennessen K, Andersen E, Clingenpeel S, Rinke C, Lundberg DS, Han J, Dangl JL, Ivanova N,

Woyke T, Kyrpides N, Pati A. 2016. ProDeGe: a computational protocol for fully automated

decontamination of genomes. The ISME journal 10:269–72.

17.   Koren S, Phillippy AM. 2015. One chromosome, one contig: complete microbial genomes

from long-read sequencing and assembly. Current Opinion in Microbiology 23:110–120.

18.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes.

Genome research 25:1043–55.

19.   Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, Wilkins MJ, Wrighton KC,

419   Williams KH, Banfield JF. 2015. Unusual biology across a group comprising more than 15% of

420   domain Bacteria. Nature 523:208–11.

421   20.   Sekiguchi Y, Ohashi A, Parks DH, Yamauchi T, Tyson GW, Hugenholtz P. 2015. First genomic

422   insights into members of a candidate bacterial phylum responsible for wastewater bulking.

423   PeerJ 3:e740.

424   21.   Chen L-X, Anantharaman K, Shaiber A, Eren AM, Banfield JF. 2020. Accurate and complete

425   genomes from metagenomes. Genome Res 30:315–333.

426   22.   Treangen TJ, Salzberg SL. 2012. Repetitive DNA and next-generation sequencing:

427   computational challenges and solutions. Nat Rev Genet 13:36–46.

428   23.   Berbers B, Saltykova A, Garcia-Graells C, Philipp P, Arella F, Marchal K, Winand R, Vanneste

429   K, Roosens NHC, De Keersmaecker SCJ. 2020. Combining short and long read sequencing to

430   characterize antimicrobial resistance genes on plasmids applied to an unauthorized

431   genetically modified Bacillus. Scientific Reports 10:4310.

432   24.   Warwick-dugdale J, Solonenko N, Moore K, Chittick L, Gregory AC, Allen MJ, Sullivan MB,

433   Temperton B. 2019. Long-read viral metagenomics captures abundant and microdiverse

434   viral populations and their niche-defining genomic islands. PeerJ 1–28.

435   25.   Ng P, Tan JJS, Ooi HS, Lee YL, Chiu KP, Fullwood MJ, Srinivasan KG, Perbost C, Du L, Sung W-

436   K, Wei C-L, Ruan Y. 2006. Multiplex sequencing of paired-end ditags (MS-PET): a strategy for

437  the ultra-high-throughput analysis of transcriptomes and genomes. Nucleic acids research

438  34:e84.

439 26. Korlach J, Marks PJ, Cicero RL, Gray JJ, Murphy DL, Roitman DB, Pham TT, Otto GA, Foquet

440  M, Turner SW. 2008. Selective aluminum passivation for targeted immobilization of single

441  DNA polymerase molecules in zero-mode waveguide nanostructures. Proceedings of the

442  National Academy of Sciences of the United States of America 105:1176–81.

443 27. Jain M, Koren S, Miga KH, Quick J, Rand AC, Sasani TA, Tyson JR, Beggs AD, Dilthey AT,

444  Fiddes IT, Malla S, Marriott H, Nieto T, O'Grady J, Olsen HE, Pedersen BS, Rhie A, Richardson

445  H, Quinlan AR, Snutch TP, Tee L, Paten B, Phillippy AM, Simpson JT, Loman NJ, Loose M.

446  2018. Nanopore sequencing and assembly of a human genome with ultra-long reads.

447  Nature biotechnology 36:338–345.

448 28. Kasianowicz JJ, Brandin E, Branton D, Deamer DW. 1996. Characterization of individual

449  polynucleotide molecules using a membrane channel. Proceedings of the National Academy

450  of Sciences of the United States of America 93:13770–3.

451 29. McCoy RC, Taylor RW, Blauwkamp T a., Kelley JL, Kertesz M, Pushkarev D, Petrov D a.,

452  Fiston-Lavier A-S. 2014. Illumina TruSeq synthetic long-reads empower de novo assembly

453  and resolve complex, highly-repetitive transposable elements. PloS one 9:e106689.

454 30. Bishara A, Moss EL, Kolmogorov M, Parada AE, Weng Z, Sidow A, Dekas AE, Batzoglou S,

455  Bhatt AS. 2018. High-quality genome sequences of uncultured microbes by assembly of read

456  clouds. Nature biotechnology 36:1067–1080.

31.   Kuleshov V, Jiang C, Zhou W, Jahanbani F, Batzoglou S, Snyder M. 2016. Synthetic long-read

sequencing reveals intraspecies diversity in the human microbiome. Nature biotechnology

34:64–9.

32.   Sharon I, Kertesz M, Hug LA, Pushkarev D, Blauwkamp TA, Castelle CJ, Amirebrahimi M,

Thomas BC, Burstein D, Tringe SG, Williams KH, Banfield JF. 2015. Accurate, multi-kb reads

resolve complex populations and detect rare microorganisms. Genome research 25:534–43.

33.   White RA, Bottos EM, Roy Chowdhury T, Zucker JD, Brislawn CJ, Nicora CD, Fansler SJ,

Glaesemann KR, Glass K, Jansson JK. 2016. Moleculo Long-Read Sequencing Facilitates

Assembly and Genomic Binning from Complex Soil Metagenomes. mSystems 1:1–15.

34.   Frank JA, Pan Y, Tooming-Klunderud A, Eijsink VGH, McHardy AC, Nederbragt AJ, Pope PB.

2016. Improved metagenome assemblies and taxonomic binning using long-read circular

consensus sequence data. Scientific reports 6:25373.

35.   Daims H, Lebedeva E V, Pjevac P, Han P, Herbold C, Albertsen M, Jehmlich N, Palatinszky M,

Vierheilig J, Bulaev A, Kirkegaard RH, von Bergen M, Rattei T, Bendinger B, Nielsen PH,

Wagner M. 2015. Complete nitrification by Nitrospira bacteria. Nature 528:504–9.

36.   Bertrand D, Shaw J, Kalathiyappan M, Ng AHQ, Kumar MS, Li C, Dvornicic M, Soldo JP, Koh

JY, Tong C, Ng OT, Barkham T, Young B, Marimuthu K, Chng KR, Sikic M, Nagarajan N. 2019.

Hybrid metagenomic assembly enables high-resolution analysis of resistance determinants

and mobile elements in human microbiomes. Nature biotechnology 37:937–944.

37. Slaby BM, Hackl T, Horn H, Bayer K, Hentschel U. 2017. Metagenomic binning of a marine

sponge microbiome reveals unity in defense but metabolic specialization. The ISME journal

11:2465–2478.

38. Arumugam K, Bessarab I, Haryono MAS, Liu X, Zuniga–Montanez RE, Roy S, Qiu G, Drautz–

Moses DI, Law YY, Wuertz S, Lauro FM, Huson DH, Williams RBH. 2021. Recovery of

complete genomes and non-chromosomal replicons from activated sludge enrichment

microbial communities with long read metagenome sequencing. npj Biofilms Microbiomes

7:23.

39. Karst SM, Ziels RM, Kirkegaard RH, Sørensen EA, McDonald D, Zhu Q, Knight R, Albertsen M.

2021. High-accuracy long-read amplicon sequences using unique molecular identifiers with

Nanopore or PacBio sequencing. Nat Methods 18:165–169.

40. Rhoads A, Au KF. 2015. PacBio Sequencing and Its Applications. Genomics, Proteomics &

Bioinformatics 13:278–289.

41. Koren S, Phillippy AM, Simpson JT, Loman NJ, Loose M. 2019. Reply to "Errors in long-read

assemblies can critically affect protein prediction". Nature biotechnology 37:127–128.

42. Nicholls SM, Quick JC, Tang S, Loman NJ. 2019. Ultra-deep, long-read nanopore sequencing

of mock microbial community standards. GigaScience 8:1–9.

43. Browne PD, Nielsen TK, Kot W, Aggerholm A, Gilbert MTP, Puetz L, Rasmussen M, Zervas A,

Hansen LH. 2020. GC bias affects genomic and metagenomic reconstructions,

underrepresenting GC-poor organisms. Gigascience 9:giaa008.

44.   Wick RR, Judd LM, Wyres KL, Holt KEY 2021. 2021. Recovery of small plasmid sequences via

497   Oxford Nanopore sequencing. Microbial Genomics 7:000631.

498   45.   O'Donnell CR, Wang H, Dunbar WB. 2013. Error analysis of idealized nanopore sequencing.

499   Electrophoresis 34:2137–2144.

500   46.   Scheunert A, Dorfner M, Lingl T, Oberprieler C. 2020. Can we use it? On the utility of de

501   novo and reference-based assembly of Nanopore data for plant plastome sequencing. PLoS

502   One 15.

503   47.   Cusco A, Perez D, Viñes J, Francino O. 2020. Long-Read Metagenomics to Retrieve High-

504   Quality Metagenome-Assembled Genomes from Canine Feces. Research Square

505   https://doi.org/10.21203/rs.3.rs-60068/v1.

506   48.   Huang Y-T, Liu P-Y, Shih P-W. 2021. Homopolish: a method for the removal of systematic

507   errors in nanopore sequencing by homologous polishing. Genome Biol 22:95.

508   49.   Hu J, Ng PC. 2012. Predicting the effects of frameshifting indels. Genome Biol 13:R9.

509   50.   Korlach J, Gedman G, Kingan SB, Chin C-S, Howard JT, Audet J-N, Cantin L, Jarvis ED. 2017.

510   De novo PacBio long-read and phased avian genome assemblies correct and add to

511   reference genes generated with intermediate and short reads. Gigascience 6:1–16.

512   51.   Delahaye C, Nicolas J. 2021. Sequencing DNA with nanopores: Troubles and biases. PLoS

513   One 16:e0257521.

514   52.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial genome

515   assemblies from short and long sequencing reads. PLoS computational biology

516   13:e1005595.

517   53.   Overholt WA, Hölzer M, Geesink P, Diezel C, Marz M, Küsel K. 2020. Inclusion of Oxford

518   Nanopore long reads improves all microbial and viral metagenome-assembled genomes

519   from a complex aquifer system. Environ Microbiol 22:4000–4013.

520   54.   Liu L, Wang Y, Che Y, Chen Y, Xia Y, Luo R, Cheng SH, Zheng C, Zhang T. 2020. High-quality

521   bacterial genomes of a partial-nitrition/anammox system by an iterative hybrid assembly

522   method. Microbiome 8:155.

523   55.   Karlsson E, Lärkeryd A, Sjödin A, Forsman M, Stenberg P. 2015. Scaffolding of a bacterial

524   genome using MinION nanopore sequencing. Scientific Reports 5:11996.

525   56.   De Maio N, Shaw LP, Hubbard A, George S, Sanderson ND, Swann J, Wick R, AbuOun M,

526   Stubberfield E, Hoosdally SJ, Crook DW, Peto TEA, Sheppard AE, Bailey MJ, Read DS, Anjum

527   MF, Walker AS, Stoesser N, On Behalf Of The Rehab Consortium null. 2019. Comparison of

528   long-read sequencing technologies in the hybrid assembly of complex bacterial genomes.

529   Microb Genom 5.

530   57.   Gan HM, Tan MH, Austin CM, Sherman CDH, Wong YT, Strugnell J, Gervis M, McPherson L,

531   Miller AD. 2019. Best Foot Forward: Nanopore Long Reads, Hybrid Meta-Assembly, and

532   Haplotig Purging Optimizes the First Genome Assembly for the Southern Hemisphere

533   Blacklip Abalone (Haliotis rubra). Front Genet 10.

58.   Shin SC, Kim H, Lee JH, Kim H-W, Park J, Choi B-S, Lee S-C, Kim JH, Lee H, Kim S. 2019.

Nanopore sequencing reads improve assembly and gene annotation of the Parochlus

steinenii genome. Sci Rep 9:5095.

59.   Haghshenas E, Asghari H, Stoye J, Chauve C, Hach F. 2020. HASLR: Fast Hybrid Assembly of

Long Reads. iScience 23.

60.   Maguire F, Jia B, Gray KL, Lau WYV, Beiko RG, Brinkman FSL. 2020. Metagenome-assembled

genome binning methods with short reads disproportionately fail for plasmids and genomic

Islands. Microbial Genomics, 6:e000436.

61.   Ross MG, Russ C, Costello M, Hollinger A, Lennon NJ, Hegarty R, Nusbaum C, Jaffe DB. 2013.

Characterizing and measuring bias in sequence data. Genome Biology 14:R51.

62.   Modlin SJ, Robinhold C, Morrissey C, Mitchell SN, Ramirez-Busby SM, Shmaya T, Valafar F.

2021. Exact mapping of Illumina blind spots in the Mycobacterium tuberculosis genome

reveals platform-wide and workflow-specific biases. Microbial Genomics 7.

63.   Arumugam K, Bağcı C, Bessarab I, Beier S, Buchfink B, Górska A, Qiu G, Huson DH, Williams

RBH. 2019. Annotated bacterial chromosomes from frame-shift-corrected long-read

metagenomic data. Microbiome 7:61.

64.   Huson DH, Albrecht B, Bağcı C, Bessarab I, Górska A, Jolic D, Williams RBH. 2018. MEGAN-LR:

new algorithms allow accurate binning and easy interactive exploration of metagenomic

long reads and contigs. Biology Direct 13:6.

65.   Hackl T, Trigodet F, Eren AM, Biller SJ, Eppley JM, Luo E, Burger A, DeLong EF, Fischer MG.

2021. proovframe: frameshift-correction for long-read (meta)genomics. bioRxiv

2021.08.23.457338.

66.   Tørresen OK, Star B, Mier P, Andrade-Navarro MA, Bateman A, Jarnot P, Gruca A, Grynberg

M, Kajava AV, Promponas VJ, Anisimova M, Jakobsen KS, Linke D. 2019. Tandem repeats

lead to sequence assembly errors and impose multi-level challenges for genome and protein

databases. Nucleic Acids Research 47:10994.

67.   Karst SM, Dueholm MS, McIlroy SJ, Kirkegaard RH, Nielsen PH, Albertsen M. 2018. Retrieval

of a million high-quality, full-length microbial 16S and 18S rRNA gene sequences without

primer bias. Nat Biotechnol 36:190–195.

68.   Amann RI, Krumholz L, Stahl DA. 1990. Fluorescent-oligonucleotide probing of whole cells

for determinative, phylogenetic, and environmental studies in microbiology. J Bacteriol

172:762–770.

69.   Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig W,

Glöckner FO. 2014. The SILVA and "All-species Living Tree Project (LTP)" taxonomic

frameworks. Nucleic Acids Res 42:D643–D648.

70.   Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet.journal 17:10.

71.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Completing bacterial genome assemblies with

multiplex MinION sequencing. Microbial genomics 3:e000132.

72.   De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. 2018. NanoPack: visualizing

and processing long-read sequencing data. Bioinformatics 34:2666–2669.

73.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using

repeat graphs. Nature biotechnology 37:540–546.

74.   Li H. 2016. Minimap and miniasm: fast mapping and de novo assembly for noisy long

sequences. Bioinformatics (Oxford, England) 32:2103–10.

75.   Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo genome assembly

from long uncorrected reads. Genome research 27:737–746.

76.   Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015. MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics (Oxford, England) 31:1674–6.

77.   Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an adaptive

binning algorithm for robust and efficient genome reconstruction from metagenome

assemblies. PeerJ 7:e7359.

78.   Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield JF. 2018. Recovery

of genomes from metagenomes via a dereplication, aggregation and scoring strategy. Nat

Microbiol 3:836–843.

79.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz P.

2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises

the tree of life. Nature biotechnology 36:996–1004.

80.   Chan PP, Lowe TM. 2019. tRNAscan-SE: Searching for tRNA Genes in Genomic Sequences.

594   Methods Mol Biol 1962:1–14.

595   81.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nature methods

596   9:357–9.

597   82.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,

598   1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map

599   format and SAMtools. Bioinformatics (Oxford, England) 25:2078–9.

600   83.   Olm MR, Brown CT, Brooks B, Banfield JF. 2017. dRep: a tool for fast and accurate genomic

601   comparisons that enables improved genome recovery from metagenomes through de-

602   replication. The ISME journal 11:2864–2868.

603   84.   2017. UniProt: the universal protein knowledgebase. Nucleic Acids Res 45:D158–D169.

604   85.   Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND.

605   Nat Methods 12:59–60.

606   86.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: Quality assessment tool for

607   genome assemblies. Bioinformatics 29:1072–1075.

SUPPLEMENT

Supplemental Figure 1. Comparison of bin relative abundances between different sequencing platforms. Relative abundance values (log-scaled) are presented between the Nanopore R9 data and **a)** PacBio CCS, **b)** Illumina, **c)** Nanopore R10. Only the bins that were clustered together between different platforms are presented in the plots and are interlinked.

619
Supplemental Table 1. Overview of read datasets used in the study

Read dataset	Instrument	Yield (Gb)	Read N50 (kb)	Read count	ENA sample ID
IL-201104	Illumina HiSeq	6.2	0.15	42,727,130	ERS7673063
IL-201112	Illumina HiSeq	11.4	0.15	79,619,634	ERS7673064
IL-201301	Illumina HiSeq	7.5	0.25	31,702,618	ERS7673065
IL-201308	Illumina HiSeq	6.7	0.25	28,067,586	ERS7673066
IL-201502	Illumina HiSeq	5.3	0.25	22,351,578	ERS7673067
IL-201702	Illumina HiSeq	15.9	0.25	66,225,442	ERS7673068
IL-201705	Illumina HiSeq	4.9	0.25	20,492,240	ERS7673069
IL-201707	Illumina HiSeq	5.5	0.25	23,663,146	ERS7673070
IL-201804	Illumina MiSeq	3.2	0.3	11,981,252	ERS7673071
IL-202001	Illumina MiSeq	13.3	0.3	47,091,904	ERS7673072
PB-202001	PacBio Sequel II	15.3	15.4	992,914	ERS7673073
R9-202001	MinION Mk1B	35.2	5.9	10,266,261	ERS7673074
R10-202001	MinION Mk1B	13.0	6.4	3,646,771	ERS7673075

622
623 **Supplemental Figure 2. Bin contamination estimates.** Only the bins that were clustered together
624 between different platforms are presented in the plot.
625

626
627 **Supplemental Figure 3. Homopolymer calling estimates in metagenomes from different**
628 **sequencing platforms.** Values in the heatmap show observed homopolymer counts estimated to be
629 called correctly at given sequence length. The total count of homopolymers (called correctly and
630 incorrectly) are in brackets. Only the contigs for bins that were clustered together between different
631 platforms were used to generate values for the plot.
632

Supplemental Figure 4. Recovery of rRNA genes from complex microbial communities via different sequencing techniques. Distribution of rRNA gene counts, found in MAGs from different sequencing methods, are presented for **a)** 16S, **b)** 23S and **c)** 5S rRNA genes. Only the MAGs, which clustered together between all the different sequencing approaches, were included in the plots. **d)** Counts for total rRNA sequences that were recovered from the assembled metagenomes. **e)** Recovered rRNA gene counts, normalized to assembly size.

Supplemental Table 2. Statistics for rRNA sequence recovery from metagenome assemblies.

Feature	Illumina	Nanopore R9	Nanopore R10	PacBio CCS
Sequencing yield (Gb)	13.3	35.2	13.0	15.3
Assembly size (Gb)	0.41	0.75	0.40	0.61
16S rRNA count	360	958	528	862
16S RNA per Gb yield	27.1	27.2	40.6	56.3
16S RNA per Gb assembly	878	1,277	1,320	1,413
23S rRNA count	476	924	524	838
23S RNA per Gb yield	35.8	26.2	40.3	54.7
23S RNA per Gb assembly	1,160	1,232	1,310	1,373
5S rRNA count	360	774	432	704
5S RNA per Gb yield	27.1	22.0	33.2	46.0
5S RNA per Gb assembly	878	1,032	1,080	1,154