Regulation of defective mitochondrial DNA accumulation and transmission in *C. elegans* by the programmed cell death and aging pathways

Sagen E. Flowers[1], Rushali Kothari[1], Yamila N. Torres Cleuren[1,2], Melissa R. Alcorn[1], Chee Kiang Ewe[1], Geneva Alok[1], Pradeep M. Joshi[1], Joel H. Rothman[1]

Affiliations:

1. Department of MCD Biology and Neuroscience Research Institute, University of California Santa Barbara, CA, USA

2. Computational Biology Unit, Institute for Informatics, University of Bergen, Norway

To whom correspondence should be addressed: rothman@lifesci.ucsb.edu

Keywords: heteroplasmy, uaDf5, purifying selection, programmed cell death, aging, insulin signaling

ORCiD:

SEF - 0000-0002-7818-2188

YNTC- 0000-0003-2218-7243

MRA- 0000-0001-6284-3255

CKE - 0000-0003-1973-1308

GA- 0000-0001-6257-2344

PMJ- 0000-0002-4220-0559

JHR - 0000-0002-6844-1377

Abstract

The heteroplasmic state of eukaryotic cells allows for cryptic accumulation of defective mitochondrial genomes (mtDNA). "Purifying selection" mechanisms operate to remove such dysfunctional mtDNAs. We found that activators of programmed cell death (PCD), including the CED-3 and CSP-1 caspases, the BH3-only protein CED-13, and PCD corpse engulfment factors, are required in *C. elegans* to attenuate germline abundance of a 3.1 kb mtDNA deletion mutation, *uaDf5*, which is normally stably maintained in heteroplasmy with wildtype mtDNA. In contrast, removal of CED-4/Apaf1 or a mutation in the CED-4-interacting prodomain of CED-3, do not increase accumulation of the defective mtDNA, suggesting induction of a non-canonical germline PCD mechanism or non-apoptotic action of the CED-13/caspase axis. We also found that the abundance of germline mtDNA^{uaDf5} reproducibly increases with age of the mothers. This effect is transmitted to the offspring of mothers, with only partial intergenerational removal of the defective mtDNA. In mutants with elevated mtDNA^{uaDf5} levels, this removal is enhanced in older mothers, suggesting an age-dependent mechanism of mtDNA quality control. Indeed, we found that both steady-state and age-dependent accumulation rates of *uaDf5* are markedly decreased in long-lived, and increased in short-lived, mutants. These findings reveal that regulators of both PCD and the aging program are required for germline mtDNA quality control and its intergenerational transmission.

Introduction

Mitochondrial diseases are a group of conditions that affect mitochondrial functions in up to 1 in 4,300 people [1–3]. Generally, these diseases present as dysfunction in the tissues or organs with the most intensive energy demands, most commonly in muscle and the nervous system [4]. Many of these diseases are attributable to mutations in the mitochondrial DNA (mtDNA) or nuclear DNA (nDNA), and include those disorders with defects in mitochondrial function, dynamics, or quality control, or in which there is miscommunication between the mitochondria and the endoplasmic reticulum (ER) [3,5]. The progressive advancement of the diseased state resulting from age-dependent accumulation of mutant mtDNA is a common trait among mitochondrial diseases [6–8]. While the severity of the disease varies with the nature of the mutation, the most severe phenotypes result in childhood death, as in Leigh syndrome and MELAS [5,9]. As there are currently no pharmacological treatments for mitochondrial diseases, it is of great importance to uncover the cellular processes that underlie the regulation of mtDNA quality control.

mtDNAs show high mutation rates [10–12] and hence it is critical that cells possess mechanisms to remove detrimental mtDNA alleles, a process called purifying selection [13,14]. Defects in this process can result in mitochondrial diseases, allowing harmful mtDNA mutations to persist through the maternal germline and subsequent generations. Processes that regulate mtDNA quality control include mitochondrial fission/fusion dynamics and mitophagy [8,15–20], and the mitochondrial unfolded protein response (UPR[MT]; [21–27]. Further, it has also been found that the insulin/IGF-1 signaling pathway (IIS; [28,29]) ameliorates the fitness defects of mutant mtDNA.

One potential cellular process that could be used to eliminate defective mtDNAs is the

culling of cells bearing mtDNA mutations by programmed cell death (PCD). The mechanisms of

both developmentally controlled and genotoxicity-induced PCD have been shown to be well-

conserved across metazoans [30], and much of the machinery that choreographs this process is

directed by mitochondria [31–33]. Mitochondrial-dependent processes participating in PCD

include permeabilization of the inner mitochondrial membrane and release of mitochondrial

factors that mediate transduction of intermediary events in the cell suicide program [31–33].

Both mitochondrial function and PCD are linked to the process of organismal aging [34]. mtDNA

mutations accumulate in tissues as organisms age, and it has been suggested that this

accumulation is a major contributor to aging [35–37].

The nematode *C. elegans* provides an attractive model for exploring the potential role of

PCD in mtDNA purifying selection. The well-described, conserved PCD regulatory pathway in *C.*

elegans functions not only to eliminate 131 somatic cells during development through a rigidly

stereotyped program [38], but is also activated apparently stochastically during germline

development, resulting in the death of >95% of nuclei that would otherwise be destined to

become oocytes in the mature hermaphrodite [30,39–41]. In addition to this "physiological" PCD,

germline nuclei that have experienced genotoxic stress are eliminated through p53-dependent

apoptosis, as is also the case in somatic mammalian cells [42,43]. Thus, germline PCD allows for

selective removal of nuclei with damaged genomes, thereby preventing intergenerational

transmission of defective nuclear DNA. Given the prominent role played by mitochondria in the

PCD process, it is conceivable that mitochondrial dysfunction could trigger PCD in the germline

and, as such, might similarly provide a quality control mechanism for eliminating aberrant

mtDNA, as seen for the nuclear genome.

We report here that germline mtDNA quality control in *C. elegans* is influenced by

regulators of both PCD and the aging program. We find that pro-apoptotic regulators of germline

PCD, notably the caspases CED-3 and CSP-1, the BH3-only domain protein CED-13, and regulators

of cell corpse engulfment, reduce abundance of an mtDNA deletion and that abrogation of their

functions results in elevated levels of the defective mtDNA. Notably, however, loss of the CED-3

activator CED-4/Apaf1 [44,45] does not result in elevated levels of defective mtDNA. These

findings raise the possibilities that either the caspases and the other pro-apoptotic factors

function in mtDNA purifying selection by a non-canonical CED-4-independent cell death program,

or that these pro-apoptotic regulators function in mtDNA purifying selection through a PCD-

independent mechanism. We also report that defective mtDNA accumulates in the germline of

animals with age and that although the abundance of the defective mtDNA is reduced in

offspring, progeny of older mothers inherit higher levels of the mutant mtDNA than those from

young mothers. Intergenerational removal of the defective mtDNA appears to be enhanced in

older animals with defective mtDNA quality control. Further, we found that lifespan-extending

mutations in both the IIS pathway and the non-IIS-dependent lifespan-regulator CLK-1/MCLK1

decrease accumulation of defective mtDNA, and that short-lived mutants show elevated

accumulation, implicating molecular regulators of the aging process in mtDNA purifying

selection. Our findings reveal that the PCD machinery and the aging program contribute to the

removal of mtDNA mutations during germline development and their intergenerational

transmission.

Results

The stably maintained mtDNA deletion mutant *uaDf5* contains multiple linked mutations

resulting in aberrant proteins and shows deleterious effects on growth

To test the role of potential regulatory factors in mtDNA purifying selection, we took

advantage of *uaDf5*, a 3.1 kb mtDNA deletion mutation that removes part or all of four protein-

coding genes and seven tRNAs (Fig. 1A) [46]. Given the presumably deleterious nature of this

defective mtDNA, it was of interest to understand how it is stably transmitted despite active

purifying selection processes. While its maintenance at high levels is attributable in part to

stabilization by the mitochondrial UPR [25], *uaDf5* persists, albeit at lower levels, in animals

lacking this activity. One possible explanation for this phenomenon is that the mutant mtDNA

might be maintained in heteroplasmy with an otherwise intact mtDNA carrying a complementing

mutation. We sought to test this possibility through deep sequencing of mtDNA isolated from

the *uaDf5*-bearing strain. Comprehensive sequence analysis revealed that, in addition to the

large deletion, the strain indeed carries a second mtDNA mutation, *w47* (Fig. 1A, Suppl. Fig. 1A,

B, C). *w47* is a single base pair insertion in the *nduo-4* gene that causes a frameshift, predicted to

result in a truncated NADH dehydrogenase 4 (ND4) protein lacking 321 residues (Suppl. Fig. 1D).

ND4 is an essential transmembrane subunit within complex I of the mitochondrial respiratory

chain (MRC), which drives NADH-oxidation-dependent transport of protons across the inner

mitochondrial membrane [47–49]. While this raised the possibility of two complementing mtDNA

genomes, we found that the *w47* mutation is present at the same abundance as the *uaDf5*

deletion mutation (~75% of total mtDNA, Suppl. Fig. 1B) rather than that of the wild-type mtDNA,

strongly suggesting that it resides on the same mtDNA genome. As this second mutation cannot

explain stabilization of the defective mtDNA by *trans*-complementation of two deleterious

mutations, other mechanisms appear to promote the stable inheritance of *uaDf5*.

In addition to the aberrant protein encoded by the *w47* frameshift mutation in *nduo-4*, a

second abnormal protein is encoded by the *uaDf5* genome: one end of the deletion results in a

fusion protein comprised of the first 185 amino acids of NADH dehydrogenase 1 (ND1, a homolog

of the core MT-ND1 transmembrane subunit of complex I of the MRC [47,49,50]), and the last 81

amino acids of mitochondrial-encoded cytochrome b (CTB-1/CYTB, a transmembrane subunit of

complex III of the MRC [47,49,51]) (Suppl. Fig. 1E). It is conceivable that accumulation of these

two abnormal proteins -- the truncated ND4 and the ND1-CYTB fusion protein resulting from *w47*

and *uaDf5*, respectively – activate the UPR^{MT}, which has been shown to result in clearance of

mtDNA^{uaDf5}, dependent on the ATFS-1 transcription factor [25,27].

Animals harboring mtDNA^{uaDf5} are viable and fertile, presumably because they contain

intact wildtype mtDNA [46]. However, we found that *uaDf5*-bearing animals displayed a

significant reduction in brood size (WT 304 ± 4.8; *uaDf5* 201 ± 8.6 embryos laid, $p < 0.001$) (Fig.

1B) and a significant increase in embryonic lethality (WT 1.4 ± 0.2%; *uaDf5* 4.2 ± 0.7%, $p < 0.001$)

(Fig. 1C). Additionally, *uaDf5* animals are slow-growing, evident in both the number of hours to

reach gravidity (WT 63 ± 0.8; *uaDf5* 76 ± 1.4 hours at 20°C, $p < 0.001$) (Fig. 1D) and the stage of

development reached after 60 hours of feeding (WT: adult; *uaDf5:* mid-L4) (Fig. 1E). In contrast,

however, we were surprised to find that lifespan was not substantially affected (WT 14 ± 0.4;

uaDf5 15 ± 0.5 days) (Fig. 1F, Suppl. Fig. 2). Given the significant decline in the majority of fitness

parameters tested, we conclude that *uaDf5* is a useful tool for studying mitochondrial disease

and mechanisms underlying mtDNA quality control.

PCD regulators promote removal of mtDNA*uaDf5*

During germline development in *C. elegans*, as many as 95% of nuclei destined to become

potential oocytes are eliminated by PCD [39–41,52]. While this process has been proposed to be

stochastically determined [39,52], it has also been suggested that it may function to selectively

remove all but the most "fit" germline cells. As such, PCD could perform a role in purifying

selection in the germline, wherein potential oocytes that undergo PCD are associated with higher

levels of defective mtDNA. To test this hypothesis, we introduced *uaDf5* into various PCD mutants

and quantified abundance of the defective mtDNA by digital-droplet PCR (ddPCR; see

Supplementary Table 1 for list of mutants tested). In a wildtype genetic background, we found

that the steady-state fractional abundance of *uaDf5* in populations of 200 day 1 adults (first day

of adulthood) is highly reproducible across four separate trials, demonstrating the reliability and

robustness of the assay. Our analyses confirmed that mtDNA*uaDf5* constitutes the major molar

fraction of mtDNA in the *uaDf5*-bearing strain by a nearly 3:1 ratio (Suppl. Fig. 3).

CED-3 in *C. elegans* is the major executioner caspase in the canonical PCD pathway

[38,52,53] and is required for virtually all PCD both in the germline and the soma (Suppl. Fig. 5).

We found that two *ced-3* mutations that strongly block PCD [54] showed a significant increase in

the ratio of defective to normal mtDNA from a molar ratio of 2.9:1 for *ced-3(+)* to 3.5:1 for *ced-*

3(n717) and 4.6:1 for *ced-3(n1286)* (Fig. 2A). This effect appears to be exclusively attributable to

an increase in mtDNA*uaDf5* in the PCD-deficient strains, as the abundance of mtDNA*WT* is not

significantly different between the strains. These *ced-3(-)* mutations both localize to the p15

domain of the protease portion of CED-3 (Suppl. Fig. 4), consistent with abolition of caspase

activity. These findings implicate the CED-3 caspase and its p15 domain in mtDNA quality control.

We found that one other mutation located in the p15 domain showed only a very slight increase

that was not statistically significant (*n2454:* 2.9:1, Fig. 2B). While it is unclear why this allele

showed a weaker effect it is noteworthy that, unlike the other two mutations, which result in a

dramatic alteration of the protein, this mutation is predicted to result in a relatively modest (ala

→ thr) single amino acid substitution.

A second caspase in *C. elegans*, CSP-1, also functions, albeit less prominently, in PCD.

While loss of CSP-1 alone does not result in a strong reduction in PCD, it synergizes with loss of

CED-3 both in PCD and in other caspase-dependent processes (Suppl. Fig. 5) [55–57]. We found

that removing CSP-1 in the *csp-1(tm917)* knockout mutant results in a significant increase in

mtDNA^{uaDf5} abundance to a molar ratio of 3.9:1 (Fig. 2A). Further, we found that this mutation

enhances the effect of the *ced-3(n717)* mutation, increasing the mtDNA^{uaDf5}:mtDNA^{WT} molar

ratio from 3.5:1 to 4.7:1 (Fig. 2A). Together these findings demonstrate that caspase activity, and

possible PCD-mediated clearance, are crucial for mtDNA quality control and function in purifying

selection of defective mtDNA.

We sought to further investigate a potential role for PCD in mtDNA purifying selection by

evaluating the requirement for the pro-apoptotic factor CED-13, a BH3-only domain protein that

acts specifically in the germline to activate PCD [38,58,59]. Consistent with a requirement for PCD

in purifying selection, we found that two *ced-13* alleles result in a very substantial increase in the

mtDNA^{uaDf5}:mtDNA^{WT} molar ratio (*sv32*: 4.2:1 and *tm536*: 5.1:1) (Fig. 2A), supporting the notion

that CED-13 promotes removal of defective mtDNA in the germline. CED-13 functions in PCD by

antagonizing the function of mitochondrially localized CED-9/Bcl-2 [58,59], which normally

sequesters the apoptosome factor CED-4/APAF1 at mitochondria, thereby preventing it from

triggering autocatalytic conversion of the executioner caspase zymogen proCED-3 to its pro-

apoptotic protease structure (Suppl. Fig. 5) [60]. An equivalent action is carried out in the soma

by the BH3-only protein EGL-1 [60]. *n1950*, a gain-of-function allele of *ced-9* that blocks the

interaction of EGL-1 with CED-9 at the mitochondria, results in elimination of PCD in the soma

but not the germline [39,52,60,61]. Consistent with the lack of effect of *ced-9(n1950gf)* on

germline PCD, we found that the mtDNA^{uaDf5}:mtDNA^{WT} molar ratio was not increased in *ced-*

9(n1950gf) mutants (2.2:1) (Fig. 2B). Thus, CED-3, CSP-1, and CED-13 are required both for

germline PCD and for removal of mtDNA^{uaDf5}.

Cells that undergo PCD are cleared by the surrounding cells in the process of engulfment

and degradation, which is implemented through a set of redundant pathways that converge on

the CED-10 GTPase [62–65] (Suppl. Fig. 5). Although this engulfment process is necessary

primarily for removal of the resultant corpses, it also appears to play an active role in cell killing:

inhibition of the engulfment pathway diminishes occurrence of PCD, likely through a complex

feedback mechanism [66,67]. Further supporting a role for PCD in purifying selection, we found

that single or double mutations of several genes that promote engulfment of cell corpses result

in elevated mtDNA^{uaDf5}:mtDNA^{WT} molar ratios, ranging from 3.5:1 to 5.2:1 (4.1:1 for *ced-*

1(e1735), 4.5:1 for *ced-2(e1752)*, 5.2:1 for the *ced-1(e1735); ced-2(e1752)* double mutant, 3.6:1

for *ced-10(n1993)*, and 4.8:1 for *ced-10(n3246)*; Fig. 2A).

We tested whether *generally* increased germline PCD alters mtDNA^{uaDf5} abundance by

examining the effect of removing the caspase-related factor, CSP-2, which has been shown to

play an anti-apoptotic role through inhibition of CED-3 autoactivation in the germline (Suppl. Fig.

5) [68]. We found that a loss-of-function mutation in *csp-2*, which elevates germline PCD, did not

alter the mtDNA^{uaDf5}:mtDNA^{WT} molar ratio (2.9:1 for the *csp-2(tm3077)* knockout mutation) (Fig.

2B). This observation does not conflict with a requirement for PCD in purifying selection: a

general increase in PCD in the germline of *csp-2(-)* animals would not be expected *per se* to alter

the mechanism that *discriminates* defective from normal mtDNAs and therefore the relative

abundance of the two forms. Rather, our findings suggest that the mechanisms that recognizes

and disposes of the defective mtDNA specifically requires PCD components acting in a selective,

rather than general process (i.e., in those cells with the highest burden of the defective mtDNA).

Evidence for non-canonical action of PCD regulators in mtDNA purifying selection

The foregoing results implicate a role for the pro-apoptotic CED-3 and CSP-1 caspases,

CED-13, and the CED-1, 2, and -10 cell corpse engulfment factors in mtDNA purifying selection.

However, several observations suggest that elimination of normal, physiological germline PCD

per se is insufficient to increase accumulation of defective mtDNA, or that these factors promote

purifying selection through a PCD-independent pathway. Specifically, we observed that the

occurrence of PCD does not perfectly correlate with their effects on accumulation of mtDNA^{uaDf5}

in particular mutants.

First, we found that the *ced-3(n718)* allele lowers, rather than elevates, the abundance of

mtDNA^{uaDf5} (mtDNA^{uaDf5}:mtDNA^{WT} molar ratio of 1.8:1, Fig. 2C). This effect is likely to be

attributable to the nature of the *n718* mutation. Those *ced-3* mutations that result in increased

mtDNA^{uaDf5} levels (Fig. 2A) both alter the p15 domain, which is essential to the active caspase

function. In contrast, the *n718* mutation changes a residue in the caspase activation and

recruitment domain (CARD), located within the prodomain of the CED-3 zymogen, which is

removed upon caspase activation and affects its activation by CED-4 (Suppl. Fig. 4) [44,54]. While

ced-3(*n718)* strongly compromises PCD, this mutation might not alter CED-3 caspase function in

a way that interferes with its role in mtDNA purifying selection.

Our surprising finding that while CED-3 activity is required for mtDNA purifying selection,

a CED-3 mutation that compromises its activation by CED-4 did not elevate mtDNA^{uaDf5} levels

prompted us to investigate the requirement of CED-4 in mitochondrial purifying selection.

Consistent with the effect of the *ced-3*(*n718)* mutation, we found that eliminating the function

of the pro-apoptotic regulator CED-4, the *C. elegans* orthologue of mammalian Apaf1 and the

upstream activator of CED-3 in the canonical PCD pathway [60,69], did not result in a marked

increase in the relative abundance of mtDNA^{uaDf5}. That is, while the mtDNA^{uaDf5}:mtDNA^{WT} molar

ratio increased to 3.1:1 in the *ced-4(n1894)* mutant, the effect was not statistically significant.

Moreover, the canonical allele *ced-4(n1162)* allele similarly showed no elevation in mtDNA^{uaDf5}

(molar ratio = 2.6:1; Fig. 2B). These results suggest that CED-3 caspase functions in mitochondrial

purifying selection independently of the caspase-activating factor CED-4.

Evidence that decreased fitness, but not lifespan, is attributable to mtDNA^{uaDf5}-induced PCD

Taken together, our results implicate many PCD regulatory factors, and potentially PCD, in

the selective clearance of defective germline mtDNAs. Our additional observations suggest that

defective mtDNAs may, in fact, *trigger* elevated germline PCD, resulting in the production of

fewer mature gametes and progeny. Specifically, we found that the significant decrease in brood

size that we observed in *uaDf5*-bearing animals with a wildtype nuclear background is partially

suppressed by both *ced-3(-)* and *ced-13(-)* mutations, which prevent PCD (Fig. 2D, Suppl. Fig. 6A),

suggesting that elimination of PCD might rescue cells that would otherwise be provoked to die

as a result of accumulation of defective mtDNA. Our findings further underscore the observation

that accumulation of defective mtDNA in those animals that do survive does not affect longevity,

as we found that lifespan is unaltered in these PCD mutants even when the levels of mtDNA^{uaDf5}

are nearly doubled (Suppl. Fig. 6B, C).

Age-dependent accumulation of mtDNA^{uaDf5} in the germline

Our findings that PCD regulators are required to reduce mtDNA^{uaDf5} abundance, the

central role that mitochondria play in PCD [31–33,70], the observed decline of mitochondrial

health during the aging process [7,35–37,71–75], and the relationship between excessive PCD

and the aging phenotype [34] led us to examine the dynamics of mtDNA^{uaDf5} accumulation as

worms age. We measured the fractional abundance of *uaDf5* in adults at progressively increased

ages spanning day 1, defined as the first day of egg-laying, through day 10. Day 1 through day 4

of adulthood encompasses the time during which nearly all self-progeny are produced. After day

4, hermaphrodite sperm become depleted and the animals transition into a post-gravid,

progressively aging state [76–78]. By day 10, animals exhibit indications of advanced age. Analysis

of the abundance of mtDNA^{uaDf5} revealed a progressive increase throughout gravidity and post-

reproductive aging, with the mtDNA^{uaDf5}:mtDNA^{WT} molar ratio increasing from 2.9:1 to 5.5:1 (Fig.

3A, Suppl. Fig. 7A). This age-related accumulation of *uaDf5* in adult worms is reminiscent of the

accumulation of mtDNA mutations seen in aging mammals [35,73,79] and suggests that *uaDf5* in

C. elegans may be a useful tool for studying the role that mtDNA mutations play in aging.

Given that gametes are depleted with age, it is conceivable that the age-dependent

increase in mtDNA^{uaDf5} is attributable to accumulation in somatic mitochondria. To assess

whether the observed age-related accumulation of mtDNA^{uaDf5} occurs predominantly in the

maternal germline or in somatic cells, we analyzed animals defective in germline development

by taking advantage of the *glp-4(bn2)* mutant, which produces only a small number (~12) of

germline cells compared to that in wildtype animals (~1,500), with no known effect on somatic

gonad development [80]. In contrast to the increased mtDNA^{uaDf5} abundance with age seen at

permissive temperature (mtDNA^{uaDf5}:mtDNA^{WT} of 1.8:1 at day 1, rising to 2.6:1 at day 4, for an

overall increase by 48%), we found that *glp-4(bn2)* animals at the non-permissive temperature

showed a slight decrease in the defective mtDNA from day 1 to day 4 of adulthood (2.2:1 at day

1, dropping to 2:1 at day 4, for an overall decrease of 6.3%) (Fig. 3B). These results strongly

suggest that the observed age-dependent increase in mtDNA^{uaDf5} abundance occurs exclusively

in the germline. We found that mtDNA^{uaDf5} does eventually appear to accumulate in somatic cells

with age, as day 10 adults grown at the restrictive temperature showed a marked increase in the

mtDNA^{uaDf5}:mtDNA^{WT} molar ratio compared to day 1 adults even in the absence of a germline

(day 10 *glp-4(bn2)* molar ratio of 3.4:1, a 57% increase from day 1 levels) (Fig. 3B). We conclude

that the marked increase in mtDNA^{uaDf5} with age during the period of fecundity occurs primarily

in the germline and that the defective mtDNA accumulates in both germline and somatic tissue

during post-reproductive life.

Age-dependent increase in mtDNA^{uaDf5} burden is transmitted to progeny

As the mtDNA is inherited strictly through the maternal germline, we posited that the

age-dependent increase in the fractional abundance of germline mtDNA^{uaDf5} might be

transmitted to progeny animals. To test this hypothesis, we measured the molar ratio of

mtDNA^{uaDf5} in 200-worm populations of L1 larvae derived from day 1 – day 4 adults (Fig. 3C,

Suppl. Fig. 7B). This analysis led to two key observations: 1) the abundance of mtDNA^{uaDf5} is

reduced during transmission between mother and offspring (average decrease ranging from 6%

to 13%), presumably as a result of purifying selection, and 2) the abundance of the defective

mtDNA in the offspring correlates with the age of the mothers: the progeny of older mothers

contain a markedly higher mtDNA^{uaDf5}:mtDNA^{WT} molar ratio (4.2:1) than that of younger mothers

(2.6:1) (Fig. 3C, D). A similar trend was observed for mother-to-offspring transmission in five

mutant strains with altered levels of mtDNA^{uaDf5} (see below): in all cases, progeny contain lower

abundance of mtDNA^{uaDf5} than their mothers, and progeny of younger adults inherit a lower load

of mtDNA^{uaDf5} than progeny of older adults (Fig. 3C). These results reveal that mtDNA quality

control occurs between primordial germ cell proliferation in the female germline and L1 hatching,

i.e., during oocyte maturation, embryogenesis, or both.

319   The lifespan-determining IIS pathway regulates accumulation of mtDNA^{uaDf5}

We sought to determine whether the age-dependent accumulation of defective mtDNA

is controlled by known molecular mechanisms that drive the aging program in *C. elegans*. The

most prominent of these regulatory systems is the highly conserved insulin/IGF-1 (insulin-like

growth factor-1) pathway (IIS), which performs a pivotal regulatory function in aging and

longevity [28,82,83] (Suppl. Fig. 8). Abrogation of the IIS signaling pathways, for example, as a

result of mutations in the gene encoding the IIS receptor (DAF-2, in *C. elegans*), results in marked

slowing of the aging program and extension of lifespan in worms, flies, and mice [84–87]. The IIS

pathway also functions in a broad set of other processes including, in *C. elegans*, activation of

two stages of developmental arrest, or diapause, at the L1 larval stage and in formation of the

dispersal form, the dauer larva, as well as in the control of germline proliferation, stress

resistance, fat metabolism, and neuronal/behavioral programs [28]. It was also reported that

inhibition of the IIS pathway rescues various fitness parameters in a mtDNA mutator strain which

contains a faulty mtDNA polymerase [29], consistent with a possible role in mtDNA quality

control.

We found that two mutant alleles of *daf-2*, which reduce rates of aging and increase

lifespan, result in dramatically decreased mtDNA^{uaDf5}:mtDNA^{WT} molar ratios from 2.8:1 to as low

as 0.3:1 (0.3:1 for *daf-2(e1391)*; 0.8:1 for *daf-2(e1370)*; Fig. 4A; see Suppl. Table 2 for a list of

lifespan mutants used in the analyses). Thus, the lifespan-extending effects of *daf-2* mutations

are strongly correlated with diminished abundance of defective mtDNA, to the extent that it

becomes the minor species of mtDNA.

The DAF-2 receptor acts by antagonizing the DAF-16/FoxO transcription factor, the major

effector of IIS downstream, in response to insulin-like ligands. Thus, removal of *daf-16* function

reverses the lifespan-extending effects of *daf-2(-)* mutants. We tested whether the DAF-2 → DAF-

16 pathway similarly functions in mtDNA purifying selection. We found that eliminating DAF-16

in two *daf-16* mutants results in slightly increased, albeit not statistically significant,

mtDNA^{uaDf5}:mtDNA^{WT} molar ratios (3.3:1 for *daf-16(mu86)*; 3.5:1 for *daf-16(mgDf50)*) (Fig. 4B).

Further, we found that removal of DAF-16 in the *daf-16(mu86)* mutant suppressed the decreased

mtDNA^{uaDf5}:mtDNA^{WT} molar ratios observed in two *daf-2* mutants, from 0.3:1 for *daf-2(e1391)* to

1:1 for *daf-16(mu86); daf-2(e1391)* and from 0.6:1 for *daf-2(e1370)* to 1.8:1 for *daf-16(mu86);*

daf-2(e1370), consistent with observations reported in a recent study [88]. While the *daf-2(-)*

effect on mtDNA^{uaDf5} levels is largely dependent on DAF-16, neither double mutant restored

mtDNA^{uaDf5} levels to those seen in animals with a fully intact IIS pathway, suggesting that other

DAF-2 targets might participate in removal of defective mtDNA (Fig. 4C). We found, conversely,

that two mutations that reduce lifespan by eliminating the function of AAK-2 (AMP activated

kinase-2), a conserved factor acting in the IIS pathway [28,89], result in elevated

mtDNA^{uaDf5}:mtDNA^{WT} molar ratios as high as 4.4:1 compared to 2.8:1 in wildtype animals (3.8:1

for *aak-2(ok524)* and 4.4:1 for *aak-2(gt33)*) (Fig. 4B). These findings demonstrate that alterations

in the IIS pathway coordinately affect both lifespan and accumulation of defective mtDNA and

that the DAF-2/DAF-16/AAK-2 axis acts similarly in both processes.

Synergistic effect of multiple aging pathways on mtDNA^{uaDf5} accumulation

In addition to IIS, other molecular regulatory pathways independently contribute to the

rate of aging. These include CLK-1, a mitochondrial hydroxylase that functions in the pathway for

ubiquinone synthesis [90,91]. *clk-1* mutants with a wildtype mitochondrial genome have been

shown to contain levels of mtDNA that are elevated by 30%, perhaps as the result of a

compensatory process that increases demand on mitochondrial abundance, or the action of CLK-

1 as a regulator of mtDNA abundance in response to energy availability within the cell [92]. As

with long-lived *daf-2* mutants, we found that long-lived *clk-1(qm30)* mutants showed a greatly

diminished mtDNA^{uaDf5}:mtDNA^{WT} molar ratio of 0.3:1 (Fig. 4A), comparable to that in *daf-2*

mutants; again, mtDNA^{uaDf5} is the minor species in these animals. As the IIS pathway and CLK-1

appear to act separately in controlling lifespan, we postulated that elimination of both

mechanisms might further reduce levels of the defective mtDNA. Indeed, we found that

mtDNA^{uaDf5} was completely eliminated in *daf-2(e1391) clk-1(qm30)* double mutants, revealing a

strongly synergistic effect between the two age-determining systems (Fig. 4A). Thus, distinct

regulatory pathways for longevity modulate the abundance of mtDNA^{uaDf5} by apparently different

mechanisms and elimination of the two pathways abrogates its maintenance.

Age-dependent accumulation rate of mtDNA^{uaDf5} strongly correlates with genetically altered

rates of aging

Our findings that the steady-state abundance of mtDNA^{uaDf5} increases with maternal age

and that mutants with increased lifespan show lower levels of the defective mtDNA raised the

possibility that purifying selection is subject to the same control as aging clocks. To assess this

potential connection, we analyzed the time-dependent rates of mtDNA^{uaDf5} accumulation in

animals with genetic backgrounds that alter the aging clock. Analysis of 21 different genetic

backgrounds over the first ten days of adulthood revealed that the age-dependent progressive

accumulation of mtDNA^{uaDf5} is a consistent phenomenon (Fig. 4D). Comparison of long-lived

mutants and wildtype using a linear regression model revealed a striking positive correlation

between aging rate and age-dependent rate of accumulation of mtDNA^{uaDf5}: all long-lived

mutants in either the IIS pathway or *clk-1* accumulate mtDNA^{uaDf5} at a substantially slower rate

as they age than do wildtype animals (Fig. 4E, Suppl. Fig. 9A, B). Conversely, we analyzed six short-

lived IIS pathway mutant combinations and found that the *aak-2(ok524)* and *daf-16(mu86)* single

mutants and *daf-16(-);daf-2(-)* double mutants all showed increased rates of mtDNA^{uaDf5}

accumulation (Fig. 4E, Suppl. Fig. 9C, D). These observations suggest that in both slower-aging

and faster-aging strains, the rate of accumulation of deleterious mtDNA is a predictor of aging

rate. In the two exceptional cases, the *aak-2(gt33)* and *daf-16(mgDf50)* single mutants, we did

not observe an increased accumulation rate compared to wildtype; however, the mtDNA^{uaDf5}

levels are consistently higher than in these two mutants than in wildtype at all stages (Fig. 4E,

Suppl. Fig. 9D) and thus diminished removal of mtDNA^{uaDf5} overall correlates with decreased

lifespan in these mutants as well. The greater mtDNA^{uaDf5} accumulation rates seen in the short-

lived animals is not attributable to the higher steady-state levels *per se*, as the rates of

accumulation of defective mtDNA observed in the PCD mutants with higher mtDNA^{uaDf5} levels

show no correlation with the steady-state levels (Suppl. Fig. 9 E-H); rather these increased rates

appear specifically to be a property of the shortened lifespan mutants.

Consistent with a relationship between aging rates and accumulation of defective mtDNA,

we found that brood size is decreased and embryonic lethality is increased in short-lived *daf-16(-*

) mutants compared to those in a wildtype nuclear background, while *uaDf5* does not impact

either fitness parameter in the long-lived *clk-1* mutant (Suppl. Fig. 10). These results are

consistent with the possibility that longevity pathways modulate fitness in part by regulating

mitochondrial homeostasis.

Evidence for late adulthood-specific mechanisms for removal of mtDNA^{uaDf5}

We obtained evidence that defective mtDNA is more effectively removed in offspring of

aging adults that carry an unusually high burden of mtDNA^{uaDf5}. The offspring of day 4 adults in

those strains ("high" strains) with significantly higher steady-state fractional abundance of

mtDNA^{uaDf5} showed significantly greater rates of reduction of the defective mtDNA (reduction

from 5.4:1 to 3.3:1 in *atfs-1(et15)*, from an extraordinarily high 6.5:1 to 2.8:1 in *ced-10(n1993),*

and from 6.4:1 to 3.3:1 in *ced-13(sv32)*) compared to offspring of ("low" strain) mothers with

lower steady-state fractional abundance of the mutant mtDNA (reduction from 4.7:1 to 4.2:1 in

WT, from 3.9:1 to 2.8:1 in *ced-3(n2454),* and from 3.2:1 to 2.5:1 in *ced-4(n1162)*). Remarkably,

therefore, day 4 progeny from "high" strains actually inherit a *lower* mtDNA^{uaDf5} load than their

siblings born from day 1-3 mothers (Fig. 3C, 5A). Indeed, we found a strong correlation (r^2=0.61,

p < 0.001) between the steady-state level of mtDNA^{uaDf5} in mothers and the capacity for its

removal between mother and progeny during day 4 of adulthood (Fig. 5B). These results raise

the possibility that very high levels of mtDNA^{uaDf5} in older mothers activate additional mtDNA

purifying selection independent of the UPR^{MT} and PCD machinery, thereby ensuring that progeny

are not overloaded with defective mitochondria.

427 **Discussion**

428 We have obtained several lines of evidence indicating that regulators of PCD and the aging

429 program function in mtDNA quality control and accumulation of defective mtDNA in *C. elegans*.

430 We report eight major findings: 1) regulators of germline PCD are required for effective removal

431 of deleterious mtDNA from the germline; 2) the cell death machinery functions in either a non-

432 canonical cell death pathway or in a non-apoptotic role to mediate mitochondrial purifying

433 selection; 3) the CSP-1 caspase has as strong of an effect on mitochondrial purifying selection as

434 the major PCD regulator CED-3; 4) mtDNA^{uaDf5} progressively accumulates in the germline as

435 adults age; 5) this age-dependent accumulation of mtDNA^{uaDf5} is transmitted to progeny;

436 however, the burden of the defective mtDNA is lower in offspring than mothers suggesting

437 intergenerational purifying selection; 6) two separate aging pathways, the IIS and CLK-1

438 pathways, act synergistically to regulate mtDNA^{uaDf5} levels; longer-lived mutants show reduced

439 levels of the defective mtDNA while shorter-lived mutants show increased levels compared to

440 otherwise wildtype animals; 7) the rate of mtDNA^{uaDf5} accumulation is inversely correlated with

441 lifespan in aging mutants; 8) intergenerational removal of mtDNA^{uaDf5} occurs more effectively

442 during transmission from older mothers with high burden of the defective mtDNA.

443 Previous reports demonstrated that UPR^{MT} limits *uaDf5* clearance, and that eliminating

444 the UPR^{MT}-mediating transcription factor, ATFS-1, lowers mtDNA^{uaDf5} abundance [25,27]. Our

445 identification of a second mutation in the *uaDf5* mutant that results in premature truncation of

446 the ND4 gene product raises the possibility that expression of both the truncated ND4 and the

447 ND1-CYTB fusion protein might together activate UPR^{MT}. The possibility that it is the production

of aberrant polypeptides resulting from these mutations that trigger this response will require

analysis of additional mtDNA mutants.

Mitochondrial deletion mutant *uaDf5* as a model for mitochondrial disease

We found that *uaDf5* affects brood size, embryonic lethality, and developmental rate,

highlighting its use as a model for investigating mitochondrial diseases. The reduced brood size

in *uaDf5*-bearing animals might reflect diminished germ cell proliferation, as mitochondria have

been implicated in progression of germline maturation [93,94]. Alternatively, the defective

mtDNA might trigger hyperactivation of the germline PCD pathway that specifically removes

germ cells with the highest burden of defective mtDNA, as suggested by our results, resulting in

the survival of fewer mature oocytes. The increased embryonic lethality in the *uaDf5* strain may

be a consequence of a genetic bottleneck effect, leading to rapid differences in mtDNA allele

frequencies [95–97] and a subpopulation of oocytes containing levels of mtDNA^{uaDf5} that exceed

a threshold required for viability.

We were surprised to find, in contrast to a previous report [98], that mtDNA^{uaDf5} did not

alter lifespan. Given that mitochondrial mutations are often coupled with compensatory

mutations in the nuclear genome [99–103], one possible explanation for this discrepancy might

be that a compensatory nuclear mutation exists in the strain analyzed, diminishing the impact of

the defective mitochondrial genome. We note, however, that we backcrossed the *uaDf5* strain

extensively to the laboratory reference strain N2 prior to performing the reported analyses. It is

conceivable that although we observed no effect in the lab, *uaDf5* might alter lifespan under

natural conditions. Exposure to increased stress from growth in the wild might be less tolerated

in animals bearing mtDNA^{uaDf5}, as has been observed with other mitochondrial mutants [104],

resulting in diminished lifespan.

Caspases and cell death machinery regulate mitochondrial purifying selection

Our results lend support to the hypothesis that germline PCD mechanisms may be used

to cull germline cells with defective mtDNA (Fig. 6A). Developmentally programmed cell death

and cell death in response to genotoxic stress are mediated by caspases upon activation by the

C. elegans octameric apoptosome which is formed when the inhibition of CED-4 by CED-9 is

disrupted through binding of BH-3-only proteins EGL-1 and CED-13 in the soma and germline,

respectively. We propose that in response to mitochondrial genotoxic stress (increased

mtDNA^{uaDf5} load), caspases CED-3 and CSP-1 are activated by the BH-3 only domain protein CED-

13 thereby triggering mitochondrial purifying selection independent of CED-9 and the CED-4

apoptosome, either through germ cell PCD or a non-apoptotic role of these caspases (Fig. 6A).

Mutations that eliminate the function of caspases that act in PCD result in elevated

mtDNA^{uaDf5} levels. Two mutations affecting the p15 domain of the CED-3 caspase result in a

significantly increased molar ratio of *uaDf5*, highlighting the importance of the p15 domain in

mtDNA quality control. It is noteworthy that the two mutations that result in a more substantial

effect on mtDNA^{uaDf5} abundance affect CED-3 structure more dramatically: *ced-3(n717)* results in

a splicing error, and *ced-3(n1286)* is a nonsense mutation [54]. In contrast, *ced-3(n2454)*, which

results in a subtle (statistically insignificant) increase in mtDNA^{uaDf5}, is a substitution predicted to

impart a much weaker effect on the protein structure [54]. The effect of the *ced-3(n718)* allele,

which resides in the prodomain of CED-3, suggests that that portion of the protein may act to

inhibit mtDNA quality control. Taken together, our results suggest that CED-3 may carry out a

specialized activity in mtDNA-activated PCD. We found that a second caspase, CSP-1, which plays

a minor role in PCD [56,57], is also required in mtDNA quality control: a *csp-1(-)* knockout

mutation results in increased abundance of mtDNA^{uaDf5} and enhances the effect of *ced-3(n717)*,

suggesting that CSP-1 may play a larger role in removal of defective mtDNA than it does in other

forms of PCD. An exciting possibility is that caspases act in mtDNA quality control via a mechanism

that is distinct from their normal action in PCD. Such putative roles for caspases and other

mitochondrial factors in non-apoptotic mitochondrial quality control may have been co-opted

during metazoan evolution with the innovation of PCD.

Analysis of additional PCD components further implicate a role for germline PCD in

mitochondrial purifying selection. These include CED-13, the germline-specific PCD effector, and

components of the cell corpse engulfment pathway, which activate PCD likely through a complex

feedback mechanism that ensures cells destined to die proceed irreversibly through the process

[66,67]. In all cases, mutations in these components result in increased mtDNA^{uaDf5} levels. In

vertebrates, mitochondrial reactive oxygen species (mtROS) trigger apoptosis via the intrinsic

mitochondrial pathway [120]. Interestingly, elevated mtROS promotes longevity that is in part

dependent on the core cell death machinery in *C. elegans*, involving CED-9, CED-4, CED-3 and

CED-13, but not EGL-1 [105]. In that study, the authors reported that the protective effect of the

cell death machinery on lifespan was independent of PCD in the soma; however, germline cell

death was not characterized. Given that CED-13, and not EGL-1, is the predominant BH3-domain

protein functioning in the germline [58] and that ablation of the germline leads to extended

lifespan [106], our findings support the possibility that germline progenitors carrying defective

mitochondria selectively undergo PCD, ensuring homeostatic mtDNA copy number and health of

progeny.

A striking exception to our findings was seen with mutations that eliminate the function

of the pro-apoptotic regulator CED-4. Neither the *ced-3(n718)* mutation that disrupts the CED-3

CARD domain, which is involved in recruitment to the apoptosome by stabilizing its interaction

with CED-4 [44,107], nor two *ced-4(-)* mutations, result in increased accumulation of mtDNA^{uaDf5},

suggesting non-canonical, CED-4-independent activation of CED-3 in mitochondrial purifying

selection. It is possible that the CARD mutation (G65R) in the *ced-3(n718)* mutant [54] reduces

the fraction of CED-3 in complex with the apoptosome, which might release more of the protein

for its role in mitochondrial purifying selection. Interestingly, the CARD linker domain has been

found to have an inhibitory effect on the pro-caspase-9 zymogen [108]. *ced-3(n718)* could be

effectively acting as a gain-of-function allele in the process of purifying selection, reflected by the

lower levels of mtDNA^{uaDf5} in this mutant background.

It is noteworthy that CED-4 and its mammalian Apaf1 relatives regulate a variety of

cellular functions that are unrelated to their activities in PCD. These include cell growth control

influenced by DNA damage, centrosomal function and morphology, neuronal regeneration, and

inhibition of viral replication [109–112]. In addition, as a result of differential RNA splicing, *ced-4*

encodes proteins with opposing activities, generating both an activator and a repressor of

apoptosis [113], which further complicates analysis of its action. Thus, it is conceivable that CED-

4 might exert opposing effects on purifying selection, reflecting its pleiotropic activities in

development and confounding an unambiguous interpretation of its action in this process.

IIS and CLK-1 synergistically regulate germline accumulation of mtDNA^{uaDf5} as adults age

We found that most of the increase in the fractional abundance of mtDNA^{uaDf5} as worms

age throughout the period of self-fertility (days 1-4) occurs in the germline. However, the relative

amount of the defective mtDNA continues to increase in older animals when the germline is no

longer actively proliferating [76], suggesting that mtDNA proliferation also occurs in somatic

tissues throughout the aging process. This behavior mirrors the dynamics of mutant mtDNAs

observed in other organisms, including human, mouse, rat, and rhesus monkey [35,73,79]. While

removal of the germline in worms results in extended lifespan [114,115], it is not clear whether,

or to what extent, this increased lifespan might be attributable to accumulation of mutant

mtDNA, since the lack of germline leads to a variety of cellular responses [116,117], any of which

might lead to lifespan extension.

These studies do not reveal whether, or how, aging and accumulation of defective mtDNA

are causally linked. However, our findings that steady-state levels of mtDNA^{uaDf5} are lowered in

long-lived mutants (*daf-2* (IIS pathway; [28]) and *clk-1* (mitochondrial function; [118,119])) and

that rates of its accumulation are strongly inversely correlated with lifespan extension through

independent pathways, suggests that mtDNA purifying selection mechanisms are influenced by

aging programs (Fig. 6A). Further bolstering this potential link is our finding that short-lived

mutants (*daf-16* and *aak-2,* both involved in the IIS pathway [28]), show higher steady-state levels

of mtDNA^{uaDf5}. That this effect is greater in *aak-2* mutants than in *daf-16* mutants suggests that

the AAK-2 branch of the IIS pathway influences mtDNA quality control more significantly than

does the DAF-16 branch. One of the substrates of AAK-2 is SKN-1/Nrf2, a multifaceted

transcription factor with roles in stress response and longevity, and one of its isoforms, SKN-1a,

localizes to the mitochondrial surface [120], raising the possibility that AAK-2 might influence

clearance of defective mtDNA through SKN-1 action. It will be of interest to assess how defective

mtDNA might coordinately trigger quality control and stress-response pathways.

Our observation that IIS pathway components and CLK-1 act synergistically on the mtDNA

quality control machinery raises the possibility that these two distinct lifespan-regulating

pathways converge on a common system for removal of defective mtDNA, such as a global

mitochondrial stress response pathway. Candidates for mediating this removal process include

mitochondrial fission/fusion [17,20,121], mitophagy [18,19,122], and the UPR^{MT} [22–27], all of

which are known to act in mtDNA quality control, as well as modulation of the regulatory

pathway for PCD, as suggested by our findings.

PCD is uncoupled from aging during intergenerational mitochondrial purifying selection in

older mothers

Analysis of newly hatched L1 larvae revealed that the relative load of mtDNA^{uaDf5} is

transmitted from mother to offspring, with evidence for intergenerational purifying selection

(Fig. 6B). This finding implies that mtDNA quality control occurs between germline stem cell

expansion in the mature female germline and L1 hatching, a developmental period that spans

many potential stages at which it might occur, including germline PCD, oocyte maturation, and

the entirety of embryogenesis. It is conceivable that this selection process acts at multiple stages

throughout this developmental window and that the decreased burden of defective mtDNA in

newly hatched L1 larvae reflect the summation of a series of sequentially acting processes that

incrementally enrich for healthy mtDNA.

The efficacy of intergenerational removal of mtDNA^{uaDf5} increases in old mothers,

including in strains lacking pro-apoptotic regulators. The clearance is particularly precipitous in

strains with a very high burden of defective mtDNA as seen in the absence of CED-13, CED-10,

and ATFS-1, suggesting a critical threshold beyond which a germline PCD-independent mtDNA

quality control process may be triggered in these older mothers (Fig. 6B). One possible

explanation for this observation is that an mtDNA purifying selection mechanism that is typically

inhibited by germline PCD might be activated in older mothers. This hypothesized purifying

selection mechanism might be triggered by the unique cellular environment associated with

aging such as increased organelle or macromolecule damage. Alternatively, the effect might be

the result of an age-dependent genetic program.

While our study has uncovered new mechanisms acting in mtDNA purifying selection and

its relationship to aging, it is of note that some level of mtDNA^{uaDf5} is maintained in all but the

most extreme conditions we have observed (e.g., in the *daf-2(e1391) clk-1(qm30)* double

mutant, in which the defective mtDNA is extirpated). This finding raises the possibility that

some degree of heteroplasmy, even with defective mtDNA, is not only tolerated, but may be

adaptive by providing a degree of evolutionary plasticity. Cells might purposefully allow for

limited heteroplasmy as a way of increasing genetic heterogeneity that might prove

evolutionarily advantageous. Such heterogeneity may also be essential to allow mtDNA to co-

evolve with changes arising in the nuclear genome. It may be that a dynamic balance between

active mtDNA purifying selection, including the mechanisms identified here, and the

permissibility of limited heteroplasmy, is modulated according to environmental or

physiological demands.

Materials and Methods

Culturing of nematodes

Nematode strains were maintained on NGM plates as previously described at either 20°C or 15°C for the temperature-sensitive strains [123]. Supplementary Tables 1, 2 and 3 provide details of all strains used in this study. Strains without a JR designation were either provided by the CGC which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440) or were obtained from the Mitani lab (strains with a FX designation or JR strains containing alleles with a tm designation were generated from Mitani lab strains) [124].

Population collection by age

Upon retrieval of a stock plate for a given strain, three chunks were taken from the stock plate and placed onto three separate large NGM plates to create three biological replicates ("lines"). Each of these lines was chunked approximately each generation to fresh large NGM plates (every 3 days if maintained at 20°C or 25°C, or every four days if maintained at 15°C, being careful to not let the worms starve between chunks). After four generations of chunks, an egg prep was performed on each line (as described previously; [123]) and left to spin in M9 overnight to synchronize the hatched L1s. The next day, each egg prep was plated onto three large seeded plates at an equal density and the worms were left to grow to day 2 adults (second day of egg laying). The day 2 adult worms were egg prepped for synchronization and left to spin in M9 buffer overnight. The next day, each egg prep was plated onto five large NGM plates at equal density. Once the worms reached day 1 of adulthood (first day of egg-laying), one of the plates was used to collect 200 adult worms by picking into 400 µl of lysis buffer, and the remaining adults on the plate were egg prepped for the collection of hatched L1 larvae in 400 µl lysis buffer the following

day. The worms on the four remaining plates were transferred to a 40 μm nylon mesh filter in

order to separate the adults from the progeny, and the resulting adults were resuspended in M9

and pipetted onto fresh large NGM plates. This process was repeated for the following three days

(Day 2-4 of adulthood). Day 5-10 adults were moved to fresh NGM plates every 2nd day using a

40 μm nylon mesh filter, and the resulting day 10 adults were collected in lysis buffer.

ddPCR

The worm lysates were incubated at 65°C for 4 hours and then 95°C for 30 minutes to

deactivate the proteinase K. Each lysate was diluted; 100-fold for 200 worm adult population

lyses, 2-fold for 200 worm L1 population lyses, and 25-fold for individual adult lyses. 2 μl of the

diluted lysate was then added to 23 μl of the ddPCR reaction mixture, which contained a

primer/probe mixture and the ddPCR probe supermix with no dUTP. The primers used were:

WTF: 5'-GAGGGCCAACTATTGTTAC-3'

WTR: 5'-TGGAACAATATGAACTGGC-3'

UADF5F: 5'-CAACTTTAATTAGCGGTATCG-3'

UADF5R: 5'-TTCTACAGTGCATTGACCTA-3'

The probes used were:

WT: 5'-HEX-TTGCCGTGAGCTATTCTAGTTATTG-Iowa Black® FQ-3'

UADF5: 5'-FAM-CCATCCGTGCTAGAAGACAAAG- Iowa Black® FQ-3'

The ddPCR reactions were put on the BioRad droplet generator and the resulting droplet-

containing ddPCR mixtures were run on a BioRad thermocycler with the following cycle

parameters, with a ramp rate of 2°C/sec for each step:

1. 95°C for 5 minutes

2. 95°C for 30 seconds

3. 60°C for 2 minutes

4. Repeat steps 2 and 3 40x

5. 4°C for 5 minutes

6. 90°C for 5 minutes

After thermocycling, the ddPCR reaction plate was transferred to the BioRad droplet reader and

the Quantasoft software was used to calculate the concentration of mtDNA^{uaDf5} (FAM positive

droplets) and mtDNA^{WT} (HEX positive droplets) in each well.

Lifespan analysis

Confluent large plates were egg prepped and left to spin in M9 overnight for

synchronization. The hatched L1s were plated onto large thick plates and allowed to grow to day

2 adults before being egg prepped a second time and left to spin in M9 overnight. The next

morning, referred to as day 1 for lifespan determination, L1s were singled out onto small plates.

Once the worms started laying eggs, they were transferred each day to a fresh small plate until

egg laying ceased, after which the worms remained on the same plate unless bacterial

contamination required transfer to a fresh plate. Worms were considered dead if there was no

movement after being lightly prodded with a worm pick. Worms that died due to desiccation on

the side of the plate were excluded from analysis.

Brood size and embryonic lethality analysis

Confluent large plates were egg prepped and left to spin in M9 overnight for

synchronization. The hatched L1s were plated onto large thick plates and allowed to grow to day

2 adults before being egg prepped a second time and left to spin in M9 overnight. The next

morning, L1s were singled out onto small plates. Once the worms started laying eggs, they were

transferred each day to a fresh small plate until egg laying ceased. The day after transfer to a

fresh plate, unhatched embryos and hatched larvae on the plate from the previous day were

counted. This was done for each of the days of laying and the total of unhatched embryos and

hatched larvae from all plates from a single worm were tabulated to determine total brood size.

To determine embryonic lethality, the total number of unhatched embryos was divided by the

total brood size. Worms that died due to desiccation on the side of the plate were excluded from

analysis.

Developmental time course analysis

Confluent large plates were egg prepped and left to spin in M9 overnight for

synchronization. The hatched L1s were plated onto large thick plates and allowed to grow to day

2 adults before being egg prepped a second time and left to spin in M9 overnight. The next

morning, L1s were singled out onto small plates. The stage of the worms was assayed every 12

hours for the first 72 hours after plating. For determining the stage at 60 hours, L4 worms were

divided up into three subgroups based on morphology: young-L4, mid-L4, and late-L4; otherwise,

all other staged worms were not divided up into subgroups. Worms that died due to desiccation

on the side of the plate were excluded from analysis.

Genotyping the *w47* allele

DNA was collected from reference strain (N2) and two *uaDf5*-containing strains, LB138

and JR3630. Mitochondrial DNA was extracted and the DNA libraries were prepared using

Nextera Kit and then sequenced using an Illumina NextSeq500. Prior to alignment, reads from

fastq files were trimmed using Trimmomatic. Trimmed, pair-end reads (2x150) were then

mapped to the *C. elegans* assembly reference sequence WBcel235 using Burrows-Wheeler

Aligner (BWA) [125]. Picard Tools (http://broadinstitute.github.io/picard/) was used to mark

duplicate reads, and SAMtools [126] was used to merge, index and create pile-up format. VarScan

[127] was used to call variants, and only variants with minimum coverage of 100 and a minimum

variant frequency call of 0.01 were considered for analysis.

Statistical Analysis

Summary statistics, ANOVA, Mann-Whitney tests, and linear regression were calculated

using R v3.4.1. The details of the statistical tests are reported in the figure legends.

 Figures

 Figure 1

702
Figure 1: Analysis of the impact of mtDNA^{uaDf5} on fitness parameters. (A) Diagram of *C. elegans*

mtDNA. Black bars with arrows indicate the locations of genes and direction of transcription.

Black lines with letters indicate the locations of tRNAs. Green bars show the locations of the

mtDNA^{uaDf5} deletion as well as the linked *w47* insertion that was identified via Illumina

sequencing. **(B)** Brood size analysis of mtDNA^{uaDf5} compared to laboratory wildtype N2. **(C)**

Embryonic lethality analysis of *uaDf5* compared to laboratory wildtype N2. **(D)** Developmental

rate analysis of mtDNA^{uaDf5} compared to laboratory wildtype N2, counting how many hours it

takes for starved L1s to reach gravidity once plated on food. **(E)** Developmental rate analysis of

mtDNA^{uaDf5} compared to laboratory wildtype N2, staging worms 60 hours after synchronized,

staved L1s are plated on food. **(F)** Survival curve analysis of mtDNA^{uaDf5} compared to laboratory

wildtype N2, day 1 is defined as the day starved L1s are plated on food. Median lifespan and

statistics are presented in Supplemental Figure 2. For **B-E**, box plots show median and IQR, and

the diamond indicates the mean. Statistical analysis was performed using the Mann-Whitney

test. (*** p < 0.001, n.s. not significant).

Figure 2

Figure 2: Regulators of PCD act on mutant mtDNA. (A-C) ddPCR analysis of the steady-state molar ratio of mtDNA^{uaDf5} in 200 worm populations of day 1 adults of various PCD mutant backgrounds. **(A)** PCD mutants that result in a significant increase in the molar ratio of mtDNA^{uaDf5}. **(B)** PCD mutants that result in no statistical change in the molar ratio of mtDNA^{uaDf5}. **(C)** PCD mutant that results in a significant decrease in the molar ratio of mtDNA^{uaDf5}. **(D)** The relative brood size of the animals with and without mtDNA^{uaDf5} in the indicated mutant backgrounds. For each nuclear genotype shown, the brood size of *uaDf5*-containing worms was

normalized by dividing by the average brood size of worms containing only *WT-mtDNA*. Box plots

show the median and IQR, the diamond indicates the mean. For **A-C**, n=3 or more biological

replicates of 200 worm populations were performed for each genotype. Statistical analysis was

performed using one-way ANOVA with Dunnett's correction for multiple comparisons. For **D**,

statistical analysis was performed using the Mann-Whitney test. Error bars represent SEM. (***

$p < 0.001$, ** $p < 0.01$, * $p < 0.05$, . $p < 0.1$, n.s. not significant).

Figure 3

Figure 3: mtDNA^{uaDf5} accumulates in the germline of aging adults, and evidence of purifying

selection between mother and offspring. (A) Analysis of the molar ratio of mtDNA^{uaDf5} in aging

adults in a wildtype nuclear background. **(B)** Analysis of the percent change of *uaDf5:*WT from

day 1 (Y axis = (*uaDf5:*WT day x -*uaDf5:*WT day 1)/(*uaDf5:*WT day 1)). For *glp-1(q231ts), fem-*

3(q20ts), and *glp-4(bn2ts),* 15^{O}C is the permissive temperature (germline development occurs)

and 25^{O}C is the restrictive temperature (female germline development is inhibited). Statistical

analysis was performed using one-way ANOVA with Tukey correction for multiple comparisons.

(*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, . $p < 0.1$). Error bars represent SEM. **(C)** Analysis of the

molar ratio of mtDNA^{uaDf5} in aging adults (P0 adult) and their L1 progeny (F1 L1) in various nuclear

backgrounds shows that all strains decrease the *uaDf5* load during transmission from mother to

offspring, and that strains with significantly higher mtDNA^{uaDf5} levels (*atfs-1(et15), ced-*

10(n1993), and *ced-13(sv32)*) have a more significant removal mechanism at day 4 of adulthood.

n=3 or more replicates of 200 worm populations were performed for each timepoint. Error bars

represent SEM. Grey dashed line indicates a hypothetical threshold at which high mtDNA^{uaDf5}

burden activates enhanced intergenerational purifying selection in older mothers. **(D)** Analysis

of the measured loss of mtDNA^{uaDf5} between mother and offspring at each day of adulthood

shows that mtDNA^{uaDf5} removal occurs. n=6 replicates of 200 worm populations for each

condition.

 Figure 4

757
Figure 4: Lifespan mutants have both a lower steady-state level and accumulation rate of

mtDNA^{uaDf5}. (A-C) Analysis of molar ratio of mtDNA^{uaDf5} in day 1 adults of various mutant

backgrounds. **(A)** Analysis of steady-state mtDNA^{uaDf5} levels in long-lived mutants. **(B)** Analysis of

steady-state mtDNA^{uaDf5} levels in short-lived mutants, showing synergistic activity on mtDNA^{uaDf5}

removal capacity in the *daf-2(e1391) clk-1(qm30)* double mutant. **(C)** Analysis of steady-state

mtDNA^{uaDf5} levels in *daf-2(-)* single and *daf-16(-); daf-2(-)* double mutants, showing a partial

rescue of *daf-2(-)* phenotype by *daf-16(-)*. **(D)** Analysis of the molar ratio of mtDNA^{uaDf5} in aging

adults in 21 different nuclear backgrounds shows a consistent accumulation trend. **(E)** Summary

of the rate of increase for the lifespan regulation mutants, showing that *daf-16* rescues the *daf-*

2 accumulation rate phenotype. The normalized accumulation rate was calculated by fitting a

regression line for each trial and then dividing the slope of the regression line by the slope of the

averaged regression line found in a wildtype background. For **all**, n=3 or more replicates of 200

worm populations for each genotype and stage. For **A and E,** statistical analysis was performed

using one-way ANOVA with Tukey correction for multiple comparisons. For **C**, statistical analysis

was performed using one-way ANOVA. For **B**, statistical analysis was performed using one-way

ANOVA with Dunnett's correction for multiple comparisons. Error bars represent SEM. (*** $p <$

0.001, ** $p < 0.01$, * $p < 0.05$, . $p < 0.1$, n.s. not significant).

Figure 5

777
Figure 5: Evidence for late adulthood-specific mechanisms for removal of mtDNA^{uaDf5}

(A) Subtracting *uaDf5:*WT in progeny from day 1 adults from progeny of day 4 adults shows that

day 4 F1-L1s tend to have higher mtDNA^{uaDf5} burden than their day 1 siblings, but this is no longer

the case in nuclear backgrounds that result in a significantly higher steady-state levels of

mtDNA^{uaDf5} in the adult. n=3 or more replicates for each genotype and statistical analysis was

performed using the Mann-Whitney test. **(B)** Comparison of the molar ratio of mtDNA^{uaDf5} in day

4 adult mothers to the absolute % removal of mtDNA^{uaDf5} from mother to offspring shows a

positive correlation. (*** p < 0.001).

786
787

Figure 6

Figure 6: Regulation of mtDNA^{uaDf5} accumulation and transmission by the PCD and aging

pathways

(A) Our results suggest that CED-3 and CSP-1, which are activated by BH-3 only protein CED-13,

function cooperatively to promote mitochondrial purifying selection, independent of CED-9 and

the CED-4 apoptosome. The clearance of mtDNA^{uaDf5} may therefore involve induction of a non-

canonical germline PCD mechanism or non-apoptotic action of the CED-13/caspase axis.

Additionally, IIS and CLK-1 aging pathways act synergistically to regulate mitochondrial purifying

selection. **(B)** mtDNA^{uaDf5} (red) accumulates in the germline relative to mtDNA^{WT} (blue) as adults

age and the increased mtDNA^{uaDf5} levels are transmitted to the progeny, although the mtDNA^{uaDf5}

burden is consistently lower in progeny than mothers. This intergenerational purifying selection

is enhanced in the older mothers of mutants with high steady-state mtDNA^{uaDf5} (e.g., *ced-13*, *ced-*

10, and *atfs-1*), suggesting a threshold beyond which a germline PCD-independent mtDNA quality

control process may be initiated or enhanced in these older mothers.

Supplemental Materials

Supplemental Figure 1

Supplemental Figure 1: Characterization of the *uaDf5* allele. (A) The percentage of reads that

mapped to the *w47* insertion in *uaDf5* samples (red) and wildtype samples (blue). Error bars

represent SEM. **(B)** Table outlining the percentage of reads that mapped to the *w47* and *uaDf5*

mutations in *uaDf5* samples. The number of variant reads was determined by averaging the

mapped reads across the deleted region and subtracting that from the average number of reads

mapped to the rest of the mtDNA genome (total reads). **(C)** Diagram showing the mitochondrial

respiratory chain (MRC) machinery subunits. Blue indicates nDNA-encoded subunits, orange and

green indicate mtDNA-encoded subunits. Green indicates those subunits that are knocked out in

the *uaDf5* allele (including ND4 which is knocked out by the linked *w47* mutation). **(D)** Diagram

showing the likely effect of the *w47* mutation on ND4 protein translation. ND4 is a 409-aa long

transmembrane subunit that spans the inner mitochondrial membrane 13 times. The *w47*

mutation results in a premature stop codon at position 89, eliminating 10 of the 13 alpha-helix

membrane domains. **(E)** Diagram showing the likely effect of the *uaDf5* mutation on ND1 protein

translation. ND1 is a 291-aa long transmembrane subunit that spans the inner mitochondrial

membrane 8 times and CYTB is a 370-aa long transmembrane subunit that spans the inner

mitochondrial membrane 8 times. The *uaDf5* mutation results in a 266 amino acid long fusion

protein that connects the first 185 amino acids (and 5 subunits) of ND1 with the last 81 amino

acids (and 3 subunits) of CYTB.

Supplemental Figure 2

Supplemental Figure 2: Lifespan analysis of the impact of *uaDf5*. Lifespan analysis of N2 bearing

uaDf5 compared to wildtype N2. Day 1 is defined as the day starved L1s are plated on food. Box

plots show median and IQR, and the diamond indicates the mean. Statistical analysis was

performed using the Mann-Whitney test. (n.s. not significant).

Supplemental Figure 3

Supplemental Figure 3: Reproducibility of ddPCR measurement of *uaDf5*. Analysis of the steady-

state of *uaDf5* in a wildtype nuclear background shows highly stable steady-state levels. Trials

were done months apart on different thaws. Dots represent biological replicates. Statistical

analysis was performed using one-way ANOVA with Tukey correction for multiple comparisons.

(n.s. not significant).

Supplemental Figure 4

A

Allele	Location	Domain	Level of *uaDf5*
n1286	W428opal	p15	highest
n717	Position 403 - Exon 7 acceptor	p15	high
n2454	A466T	p15	unchanged
n718	G65R	Prodomain	low

B

Supplemental Figure 4: Analysis of the *ced-3* alleles. (A) Locations and consequences of the four

tested *ced-3* alleles, as well as the measured fractional abundance of *uaDf5*. **(B)** Diagram showing

the locations of the mutations for each allele.

Supplemental Figure 5

Supplemental Figure 5: PCD Signaling pathway. (A) Signaling pathway for programmed cell

death. The canonical pathway is shown with black arrows, the noncanonical pathway is shown

with green arrows, and the downstream engulfment pathway is shown with blue arrows.

Adapted from [38].

Supplemental Figure 6

Supplemental Figure 6: Analysis of the impact of *uaDf5* on fitness parameters in PCD mutants.

(A) Brood size analysis of PCD mutants. **(B)** Lifespan analysis shows that *uaDf5* does not affect

lifespan in both wildtype background and in PCD mutant backgrounds. **(C)** Lifespan analysis

shows that *uaDf5* does not affect lifespan in a wildtype nuclear background nor in PCD mutant

backgrounds. For **B and C,** day 1 is defined as the day starved L1s are plated on food. For **A and**

C, box plots show median and IQR, and the diamond indicates the mean. Statistical analysis was

performed using the Mann-Whitney Wilcoxon test. (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, . $p <$

0.1, n.s. not significant).

Supplemental Figure 7

Supplemental Figure 7: *uaDf5* accumulation in individual lines of adults and progeny. (A)

Analysis of *uaDf5* accumulation in individual lines of aging P0 adults shows a consistent

accumulation trend as adults age. **(B)** Analysis of *uaDf5* accumulation in individual lines of F1-L1

progeny that were born from those mothers shown in panel A show a consistent trend of progeny

born from older mothers inheriting a larger *uaDf5* load than their siblings born from younger

mothers.

Supplemental Figure 8

Supplemental Figure 8: IIS Signaling pathway. Insulin-like peptides (ILPs) bind to DAF-2 and

activate the PI3P pathway which prevents nuclear translocation of DAF-16. AAK-2 may

phosphorylate and activate DAF-16 transcriptional activity. Loss of *aak-2* or *daf-2* (highlighted in

red) reduces lifespan

    Supplemental Figure 9

Supplemental Figure 9: Analysis of the accumulation of mtDNA^{uaDf5} in PCD and lifespan

mutants. Analysis of the normalized fractional abundance *uaDf5* [(% *uaDf5* day x- % *uaDf5* day

1)* % *uaDf5* day 1] in steady-state populations, showing the accumulation rate of mtDNA^{uaDf5} as

worms age from day 1 to day 4 of adulthood. **(A)** Analysis of the accumulation of *uaDf5* in long-

lived mutants. **(B)** Analysis of the accumulation of mtDNA^{uaDf5} in short-lived mutants. **(C-D)** Linear

regression analysis of the normalized fractional abundance *uaDf5* [(% *uaDf5* day x- % *uaDf5* day

1)* % *uaDf5* day 1] in steady-state populations, showing the accumulation rate of mtDNA^{uaDf5} as

worms age from day 1 to day 4 of adulthood. **(C)** Analysis of long-lived mutants. **(D)** Analysis of

short-lived mutants. **(E)** Analysis of the accumulation of mtDNA^{uaDf5} in PCD mutants. **(F-H)** Linear

regression analysis of the normalized fractional abundance mtDNA^{uaDf5} [(% *uaDf5* day x- % *uaDf5*

day 1)* % *uaDf5* day 1] in steady-state populations, showing the accumulation rate of mtDNA^{uaDf5}

as worms age from day 1 to day 4 of adulthood. **(F)** Analysis of PCD mutants with significantly

high steady-state level of mtDNA^{uaDf5}. **(G)** Analysis of PCD mutants with significantly low steady-

state level of mtDNA^{uaDf5}. **(H)** Analysis of PCD mutants with no significant change in the steady-

state level of mtDNA$^{uaDf5.}$ For **C, D, F, G, and H** dots represent actual datapoints, lines are fitted

regression models of the data. For all, n=3 replicates or more of 200 worm populations for each

datapoint.

 Supplemental Figure 10

Supplemental Figure 10: *uaDf5* differentially impacts fitness parameters in lifespan-affecting

mutants. (A-B) Brood size analysis showing how *uaDf5* differentially impacts brood size in

lifespan mutant backgrounds. *uaDf5* has no negative impact on the long-lived mutant *clk-1* but

has a modestly larger negative impact on the short-lived mutant *daf-16* than it does in the

wildtype background. **(B)** Relative brood size of the animals with and without *uaDf5* in the

indicated mutant backgrounds. **(C-D)** Embryonic lethality analysis showing how *uaDf5*

differentially impacts embryonic lethality in lifespan mutant backgrounds. *uaDf5* has no negative

impact on the long-lived mutant *clk-1* but has a modestly larger negative impact on the short-

lived mutant *daf-16* than it does in a wildtype background. **(D)** Relative lethality of the animals

with and without *uaDf5* in the indicated mutant backgrounds. Box plots show median and IQR,

and the diamond indicates the mean. For **A and C**, statistical analysis was performed using the

Mann-Whitney test. For **B and D**, statistical analysis was performed using the Kruskal-Wallis test.

(*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, . $p < 0.1$, n.s. not significant).

Supplementary Table 1: A summary of all mutants analyzed in the PCD pathway, including their

known homologs, whether they are part of the core PCD machinery, if they are pro-apoptotic or

anti-apoptotic, whether they are mitochondrial proteins, and molecular details of the alleles

analyzed.

Gene	Homolog	Core Machinery	Role in Apoptosis	Mitocho ndrial?	Allele	Parental Strain	Molecular Nature of the Allele	Protein Change
ced-1	SCARF2	N	Pro(engulf ment)	N	*e1735*	CB3203	Substitution	Nonsense Q→Ochre
ced-2	CRK	N	Pro(engulf ment)	N	*e1752*	CB3257	Substitution	Nonsense W→Opal
ced-3	CASPASE	Y	Pro(execu tor caspase)	N	*n717*	MT1522	Substitution	Splice site C→T
					n1286	MT3002	Substitution	Nonsense W→Opal
					n718	MT1743	Substitution	Missense G→R
					n2454	MT8354	Substitution	Missense A→T
ced-4	APAF1	Y	Pro(apopt osome)	Y/N (primaril y nuclear membra ne)	*n1162*	MT2547	Substitution	Nonsense Q→Ochre
					n1894	MT5287	Not curated	Unknown
ced-9	BCL2	Y	Anti	Y	*n1950*	MT4770	Substitution	Missense G→E
ced-10	RAC	N	Pro(engulf ment)	N	*n1993*	MT5013	Substitution	Missense V→G
					n3246	MT9958	Substitution	Missense G→R
ced-13	BH3	N	Pro	N	*tm536*	FX536	523bp deletion	Only first 17bp remain
					sv32	MD792	1304bp deletion	Complete knockout, also *inx-5*
csp-1	CASPASE	N	Pro(caspa se)	N	*tm917*	JR3196	751bp deletion	Only first 59bp remain
csp-2	CASPASE	N	Anti	N	*tm307 7*	JR3397	319bp deletion	Start at position 7317, deletes parts of exon 13

								and all of 14

925 **Supplementary Table 2:** A summary of all lifespan mutants analyzed, including their known

926 homologs, cellular pathways they are known to act in, whether the mutant extends or reduces

927 lifespan, and molecular details of the alleles analyzed.

Gene	Homologs	Pathway	Allele	Parental Strain	Molecular Nature of the Allele	Protein Change	Mutant lifespan	Note
aak-2	AMPK	IIS pathway, adult lifespan, peptidyl-serine phosphorylation, regulation of protein localization	*gt33*	TG38	606bp deletion	Starts at position 3979, deletes exon 3	Decrease	-
			ok524	RB754	408bp deletion	Starts at position 4136, deletes part of exon 3	Decrease	-
clk-1	COQ7	Adult behavior, regulation of cellular metabolism	*qm30*	MQ130	590bp deletion	Starts at position 1044, deletes part of exon 4 and 5	Increase	-
daf-2	IGFR	IIS pathway, cellular response to salt, negative regulator of macromolecule metabolic processes, positive regulation of developmental growth	*e1391*	DR1574	Substitution	Missense P→L	Increase	ts
			e1370	CB1370	Substitution	Missense P→S	Increase	ts
			m41	DR1564	Substitution	Missense G→E	Increase	ts
daf-16	FOXO	IIS pathway, defense response to bacterium, regulation of cellular biosynthetic process, regulation of post-	*mu86*	CF1038	1098bp deletion	Deletes 5 exons	Decrease	-
			mgDf50	GR1307	20193bp deletion with TCTTCATTTTCAG insertion	Deletes 7 exons	Decrease	-

		embryonic development						
fem-3	novel	Male somatic sex determination, masculinization of hermaphrodite germline, positive regulation of macromolecule metabolic process	*q20*	JK816	Not curated	Unknown	Increase	Gof and ts; female germline development inhibited
glp-4	VARS	Cell fate specification, embryonic pattern specification, positive regulation of cell proliferation	*bn2*	SS104	Substitution	Missense G→D	Increase	ts; germline development inhibited

928
929

 Supplementary Table 3: *C. elegans* strains used in this study.

Strain	Genotype	Source
N2	wildtype	CGC
LB138	*him-8(e1489) IV; uaDf5/+*	CGC
JR3630	*N2; uaDf5/+ (8x backcross version of LB138, him(e1489) was outcrossed)*	This study
JR3880	*ced-3(n717) IV; uaDf5/+*	This study
JR4330	*csp-1(tm917) II; uaDf5/+*	This study
JR3977	*ced-3(n1286) IV; uaDf5/+*	This study
JR3955	*ced-13(sv32) X; uaDf5/+*	This study
JR3976	*ced-13(tm536) X; uaDf5/+*	This study
JR3926	*ced-10(n1993) IV; uaDf5/+*	This study
JR4001	*ced-10(n3246) IV; uaDf5/+*	This study
JR3972	*ced-1(e1735) I; uaDf5/+*	This study
JR3978	*ced-2(e1752) IV; uaDf5/+*	This study
JR4010	*ced-1(e1735)I; ced-2(e1752)IV; uaDf5/+*	This study
JR3938	*ced-3(n2454) IV; uaDf5/+*	This study
JR3925	*ced-4(n1162) III; uaDf5/+*	This study
JR4017	*ced-4(n1894) III; uaDf5/+*	This study
JR3986	*csp-2(tm3077) IV; uaDf5/+*	This study
JR4027	*ced-9(n1950) III; uaDf5/+*	This study
JR3983	*ced-3(n718) IV; uaDf5/+*	This study
MT1522	*ced-3(n717) IV*	CGC
FX536	*ced-13(tm536) X*	CGC
JR3966	*glp-4(bn2) I; uaDf5/+*	This study
JR3949	*fem-3(q20) IV; uaDf5/+*	This study
JR3937	*atfs-1(et15) V; uaDf5/+*	This study
JR3960	*daf-2(e1370) III; uaDf5/+*	This study
JR3963	*daf-2(e1391) III; uaDf5/+*	This study
JR3993	*clk-1(qm30) III; uaDf5/+*	This study
JR4060	*daf-2(e1391) clk-1(qm30) III; uaDf5/+*	This study
JR3995	*daf-16(mu86) I; uaDf5/+*	This study
JR3997	*daf-16(mgDf50) I; uaDf5/+*	This study
JR4011	*aak-2(ok524) X; uaDf5/+*	This study
JR4012	*aak-2(gt33) X; uaDf5/+*	This study
JR4008	*daf-16(mu86) I; daf-2(e1391) III; uaDf5/+*	This study
JR4005	*daf-16(mu86) I; daf-2(e1370) III; uaDf5/+*	This study
JR3941	*glp-1(q231) III; uaDf5/+*	This study
CF1038	*daf-16(mu86) I*	CGC
MQ130	*clk-1(qm30) III*	CGC
JR3981	*csp-1(tm917) II; ced-3(n717) IV; uaDf5/+*	This study

Acknowledgments

We would like to thank the members of the Rothman lab for their support. We thank Jen Smith

and the BNL at UCSB for providing excellent facilities which were necessary for this work. We

thank Kyle Ploense for his help in statistical analysis. Worm strains used in this work were

provided by the Mitani lab, as well as the Caenorhabditis Genetics Center (CGC), which is funded

by NIH Office of Research Infrastructure Programs Grant P40 OD010440.

Competing Interests

The authors declare no competing or financial interests.

Author contributions

Sagen E. Flowers - conceptualization, investigation, funding acquisition, writing – original draft

preparation, writing – review and editing

Rushali Kothari – investigation, writing – review and editing

Yamila N. Torres Cleuren - investigation, writing – review and editing

Melissa R. Alcorn – statistics, writing – review and editing

Chee Kiang Ewe - writing – review and editing

Geneva Alok - writing – review and editing

Pradeep M. Joshi – conceptualization, investigation, writing – review and editing

Joel H. Rothman - conceptualization, supervision, funding acquisition, writing – review and

editing

Funding

This work was supported by the grants from the National Institutes of Health (#R01HD082347,

#R01HD081266, #R01GM143771, and #R21AG068915).

Data Availability

All data generated or analyzed during this study are included in this article and accompanied

supplementary materials.

References

1.   YS N, DM T. Mitochondrial disease: genetics and management. J Neurol. J Neurol;

2016;263: 179–191. doi:10.1007/S00415-015-7884-3

2.   PF C. Mitochondrial disease in adults: what's old and what's new? EMBO Mol Med.

EMBO Mol Med; 2015;7: 1503–1512. doi:10.15252/EMMM.201505079

3.   Gorman GS, Schaefer AM, Ng Y, Gomez N, Blakely EL, Alston CL, et al. Prevalence of

nuclear and mitochondrial DNA mutations related to adult mitochondrial disease. Ann

Neurol. John Wiley & Sons, Ltd; 2015;77: 753–759. doi:10.1002/ANA.24362

4.   Ghaoui R, Sue CM. Movement disorders in mitochondrial disease. J Neurol. Dr. Dietrich

Steinkopff Verlag GmbH and Co. KG; 2018;265: 1230–1240. doi:10.1007/s00415-017-

8722-6

5.   Area-Gomez E, Schon EA. Mitochondrial genetics and disease. Journal of Child Neurology.

SAGE Publications Inc.; 2014. pp. 1208–1215. doi:10.1177/0883073814539561

6.   Dhillon VS, Fenech M. Mutations that affect mitochondrial functions and their

association with neurodegenerative diseases. Mutation Research - Reviews in Mutation

Research. 2014. pp. 1–13. doi:10.1016/j.mrrev.2013.09.001

7.   Park CB, Larsson NG. Mitochondrial DNA mutations in disease and aging. Journal of Cell

Biology. 2011. pp. 809–818. doi:10.1083/jcb.201010024

8.   Burté F, Carelli V, Chinnery PF, Yu-Wai-Man P. Disturbed mitochondrial dynamics and

neurodegenerative disorders. Nature Reviews Neurology. Nature Publishing Group;

2015. pp. 11–24. doi:10.1038/nrneurol.2014.228

9.   Schon EA, Dimauro S, Hirano M. Human mitochondrial DNA: Roles of inherited and

somatic mutations. Nature Reviews Genetics. 2012. pp. 878–890. doi:10.1038/nrg3275

10. Brown WM, George M, Wilson AC. Rapid evolution of animal mitochondrial DNA. Proc

Natl Acad Sci U S A. 1979;76: 1967–1971. doi:10.1073/pnas.76.4.1967

11. Konrad A, Thompson O, Waterston RH, Moerman DG, Keightley PD, Bergthorsson U, et

al. Mitochondrial mutation rate, spectrum and heteroplasmy in Caenorhabditis elegans

spontaneous mutation accumulation lines of differing population size. Mol Biol Evol.

Oxford University Press; 2017;34: 1319–1334. doi:10.1093/molbev/msx051

12. Denver DR, Morris K, Lynch M, Vassilieva LL, Thomas WK. High direct estimate of the

mutation rate in the mitochondrial genome of Caenorhabditis elegans. Science (80-).

2000;289: 2342–2344. doi:10.1126/science.289.5488.2342

13. Palozzi JM, Jeedigunta SP, Hurd TR. Mitochondrial DNA Purifying Selection in Mammals

and Invertebrates. Journal of Molecular Biology. Academic Press; 2018. pp. 4834–4848.

doi:10.1016/j.jmb.2018.10.019

14. Stewart JB, Freyer C, Elson JL, Larsson NG. Purifying selection of mtDNA and its

implications for understanding evolution and mitochondrial disease. Nature Reviews

Genetics. 2008. pp. 657–662. doi:10.1038/nrg2396

15. Twig G, Shirihai OS. The interplay between mitochondrial dynamics and mitophagy.

Antioxidants and Redox Signaling. 2011. pp. 1939–1951. doi:10.1089/ars.2010.3779

16. Busch KB, Kowald A, Spelbrink JN. Quality matters: How does mitochondrial network

dynamics and quality control impact on mtDNA integrity? Philosophical Transactions of

the Royal Society B: Biological Sciences. Royal Society of London; 2014.

doi:10.1098/rstb.2013.0442

17. Ni HM, Williams JA, Ding WX. Mitochondrial dynamics and mitochondrial quality control.

Redox Biology. Elsevier B.V.; 2015. pp. 6–13. doi:10.1016/j.redox.2014.11.006

18. Ashrafi G, Schwarz TL. The pathways of mitophagy for quality control and clearance of

mitochondria. Cell Death and Differentiation. 2013. pp. 31–42. doi:10.1038/cdd.2012.81

19. Youle RJ, Narendra DP. Mechanisms of mitophagy. Nat Rev Mol Cell Biol. 2011;12: 9–14.

doi:10.1038/nrm3028

20. Tilokani L, Nagashima S, Paupe V, Prudent J. Mitochondrial dynamics: Overview of

molecular mechanisms. Essays in Biochemistry. Portland Press Ltd; 2018. pp. 341–360.

doi:10.1042/EBC20170104

21. Hernando-Rodríguez B, Artal-Sanz M. Mitochondrial Quality Control Mechanisms and the

PHB (Prohibitin) Complex. Cells. MDPI AG; 2018;7: 238. doi:10.3390/cells7120238

22. Sté phane Rolland AG, Schneid S, Schwarz M, Mokranjac D, Lambie E, Conradt B.

Compromised Mitochondrial Protein Import Acts as a Signal for UPR mt. Cell Rep.

2019;28: 1659–1669. doi:10.1016/j.celrep.2019.07.049

23. Münch C. The different axes of the mammalian mitochondrial unfolded protein response.

BMC Biology. BioMed Central Ltd.; 2018. doi:10.1186/s12915-018-0548-x

24. Callegari S, Dennerlein S. Sensing the stress: A role for the UPR mt and UPR am in the

quality control of mitochondria. Frontiers in Cell and Developmental Biology. Frontiers

Media S.A.; 2018. doi:10.3389/fcell.2018.00031

1021 25. Gitschlag BL, Kirby CS, Samuels DC, Gangula RD, Mallal SA, Patel MR. Homeostatic

1022   Responses Regulate Selfish Mitochondrial Genome Dynamics in C. elegans. Cell Metab.

1023   Cell Press; 2016;24: 91–103. doi:10.1016/j.cmet.2016.06.008

1024 26. Nargund AM, Pellegrino MW, Fiorese CJ, Baker BM, Haynes CM. Mitochondrial import

1025   efficiency of ATFS-1 regulates mitochondrial UPR activation. Science (80-). American

1026   Association for the Advancement of Science; 2012;337: 587–590.

1027   doi:10.1126/science.1223560

1028 27. Lin YF, Schulz AM, Pellegrino MW, Lu Y, Shaham S, Haynes CM. Maintenance and

1029   propagation of a deleterious mitochondrial genome by the mitochondrial unfolded

1030   protein response. Nature. Nature Publishing Group; 2016;533: 416–419.

1031   doi:10.1038/nature17989

1032 28. Murphy CT, Hu PJ. Insulin/insulin-like growth factor signaling in C. elegans [Internet].

1033   WormBook : the online review of C. elegans biology. 2013. pp. 1–43.

1034   doi:10.1895/wormbook.1.164.1

1035 29. Haroon S, Li A, Weinert JL, Fritsch C, Ericson NG, Alexander-Floyd J, et al. Multiple

1036   Molecular Mechanisms Rescue mtDNA Disease in C. elegans. Cell Rep. Elsevier B.V.;

1037   2018;22: 3115–3125. doi:10.1016/j.celrep.2018.02.099

1038 30. Lord CEN, Gunawardena AHLAN. Programmed cell death in C. elegans, mammals and

1039   plants. European Journal of Cell Biology. 2012. pp. 603–613.

1040   doi:10.1016/j.ejcb.2012.02.002

1041 31. Jeong SY, Seol DW. The role of mitochondria in apoptosis. Journal of Biochemistry and

1042   Molecular Biology. 2008. pp. 11–22. doi:10.5483/bmbrep.2008.41.1.011

32.   Estaquier J, Vallette F, Vayssiere JL, Mignotte B. The mitochondrial pathways of

apoptosis. Adv Exp Med Biol. 2012;942: 157–183. doi:10.1007/978-94-007-2869-1_7

33.   Bhola PD, Letai A. Mitochondria-Judges and Executioners of Cell Death Sentences.

Molecular Cell. Cell Press; 2016. pp. 695–704. doi:10.1016/j.molcel.2016.02.019

34.   Tower J. Programmed cell death in aging. Ageing Research Reviews. Elsevier Ireland Ltd;

2015. pp. 90–100. doi:10.1016/j.arr.2015.04.002

35.   Larsson N-G. Somatic Mitochondrial DNA Mutations in Mammalian Aging. Annu Rev

Biochem. Annual Reviews; 2010;79: 683–706. doi:10.1146/annurev-biochem-060408-

093701

36.   Payne BAI, Chinnery PF. Mitochondrial dysfunction in aging: Much progress but many

unresolved questions. Biochim Biophys Acta. 2015;1847: 1347–53.

doi:10.1016/j.bbabio.2015.05.022

37.   Kauppila TES, Kauppila JHK, Larsson NG. Mammalian Mitochondria and Aging: An

Update. Cell Metabolism. Cell Press; 2017. pp. 57–71. doi:10.1016/j.cmet.2016.09.017

38.   Conradt B, Wu YC, Xue D. Programmed cell death during Caenorhabditis elegans

development. Genetics. Genetics; 2016;203: 1533–1562.

doi:10.1534/genetics.115.186247

39.   Gumienny TL, Lambie E, Hartwieg E, Horvitz HR, Hengartner MO. Genetic control of

programmed cell death in the Caenorhabditis elegans hermaphrodite germline.

Development. 1999;126: 1011–1022. doi:10.5167/uzh-1041

40.   Baum JS, St. George JP, McCall K. Programmed cell death in the germline. Seminars in

Cell and Developmental Biology. Elsevier Ltd; 2005. pp. 245–259.

doi:10.1016/j.semcdb.2004.12.008

41.   Jaramillo-Lambert A, Ellefson M, Villeneuve AM, Engebrecht JA. Differential timing of S

phases, X chromosome replication, and meiotic prophase in the C. elegans germ line. Dev

Biol. Academic Press Inc.; 2007;308: 206–221. doi:10.1016/j.ydbio.2007.05.019

42.   Hafner A, Bulyk ML, Jambhekar A, Lahav G. The multiple mechanisms that regulate p53

activity and cell fate. Nature Reviews Molecular Cell Biology. Nature Publishing Group;

2019. pp. 199–210. doi:10.1038/s41580-019-0110-x

43.   Derry WB, Putzke AP, Rothman JH. Caenorhabditis elegans p53: Role in apoptosis,

meiosis, and stress resistance. Science (80-). 2001;294: 591–595.

doi:10.1126/science.1065486

44.   Huang W, Jiang T, Choi W, Qi S, Pang Y, Hu Q, et al. Mechanistic insights into CED-4-

mediated activation of CED-3. Genes Dev. Genes Dev; 2013;27: 2039–2048.

doi:10.1101/GAD.224428.113

45.   Seshagiri S, Miller LK. Caenorhabditis elegans CED-4 stimulates CED-3 processing and

CED-3-induced. Curr Biol. Cell Press; 1997;7: 455–460. doi:10.1016/S0960-

9822(06)00216-8

46.   Tsang WY, Lemire BD. Stable heteroplasmy but differential inheritance of a large

mitochondrial DNA deletion in nematodes. Biochem Cell Biol. 2002;80: 645–654.

doi:10.1139/o02-135

47.   Lemire B. Mitochondrial genetics. WormBook. 2005; doi:10.1895/wormbook.1.25.1

48.   Van Der Bliek AM, Sedensky MM, Morgan PG. Cell biology of the mitochondrion.

Genetics. Genetics Society of America; 2017;207: 843–871.

1087   doi:10.1534/genetics.117.300262

1088 49. Sousa JS, D'Imprima E, Vonck J. Mitochondrial respiratory chain complexes. Subcellular

1089   Biochemistry. Springer New York; 2018. pp. 167–227. doi:10.1007/978-981-10-7757-9_7

1090 50. Baradaran R, Berrisford JM, Minhas GS, Sazanov LA. Crystal structure of the entire

1091   respiratory complex i. Nature. Nature; 2013;494: 443–448. doi:10.1038/nature11871

1092 51. Song Z, Laleve A, Vallières C, McGeehan JE, Lloyd RE, Meunier B. Human Mitochondrial

1093   Cytochrome b Variants Studied in Yeast: Not All Are Silent Polymorphisms. Hum Mutat.

1094   John Wiley and Sons Inc.; 2016;37: 933–941. doi:10.1002/humu.23024

1095 52. Gartner A, Boag PR, Blackwell TK. Germline survival and apoptosis. WormBook : the

1096   online review of C. elegans biology. 2008. pp. 1–20. doi:10.1895/wormbook.1.145.1

1097 53. Cohen GM. Caspases: The executioners of apoptosis. Biochemical Journal. Portland Press

1098   Ltd; 1997. pp. 1–16. doi:10.1042/bj3260001

1099 54. Shaham S, Reddien PW, Davies B, Horvitz HR. Mutational Analysis of the Caenorhabditis

1100   elegans Cell-Death Gene ced-3. Genetics. 1999;153.

1101 55. Jeong PY, Kumar A, Joshi PM, Rothman JH. Intertwined Functions of Separase and

1102   Caspase in Cell Division and Programmed Cell Death. Sci Rep. Nature Research; 2020;10.

1103   doi:10.1038/s41598-020-63081-w

1104 56. Denning DP, Hatch V, Horvitz HR. Both the Caspase CSP-1 and a Caspase-Independent

1105   Pathway Promote Programmed Cell Death in Parallel to the Canonical Pathway for

1106   Apoptosis in Caenorhabditis elegans. PLoS Genet. 2013;9.

1107   doi:10.1371/journal.pgen.1003341

1108 57. Shaham S. Identification of multiple Caenorhabditis elegans caspases and their potential

roles in proteolytic cascades. J Biol Chem. 1998;273: 35109–35117.

doi:10.1074/jbc.273.52.35109

58.   King SD, Gray CF, Song L, Nechushtai R, Gumienny TL, Mittler R, et al. The cisd gene

family regulates physiological germline apoptosis through ced-13 and the canonical cell

death pathway in Caenorhabditis elegans. Cell Death Differ. Nature Publishing Group;

2019;26: 162–178. doi:10.1038/s41418-018-0108-5

59.   Schumacher B, Schertel C, Wittenburg N, Tuck S, Mitani S, Gartner A, et al. C. elegans

ced-13 can promote apoptosis and is induced in response to DNA damage. Cell Death

Differ. Nature Publishing Group; 2005;12: 153–161. doi:10.1038/sj.cdd.4401539

60.   Fairlie WD, Perugini MA, Kvansakul M, Chen L, Huang DCS, Colman PM. CED-4 forms a 2 :

2 heterotetrameric complex with CED-9 until specifically displaced by EGL-1 or CED-13.

Cell Death Differ. 2006;13: 426–434. doi:10.1038/sj.cdd.4401762

61.   Conradt B, Xue D. Programmed cell death. WormBook : the online review of C. elegans

biology. 2005. pp. 1–13. doi:10.1895/wormbook.1.32.1

62.   Wu D, Chai Y, Zhu Z, Li W, Ou G, Li W. CED-10-WASP-Arp2/3 signaling axis regulates

apoptotic cell corpse engulfment in C. elegans. Dev Biol. Academic Press Inc.; 2017;428:

215–223. doi:10.1016/j.ydbio.2017.06.005

63.   Kinchen JM, Cabello J, Kilngele D, Wong K, Felchtinger R, Schnabel H, et al. Two pathways

converge at CED-10 to mediate actin rearrangement and corpse removal in C. elegans.

Nature. 2005;434: 93–99. doi:10.1038/nature03263

64.   Hochreiter-Hufford A, Ravichandran KS. Clearing the dead: Apoptotic cell sensing,

recognition, engulfment, and digestion. Cold Spring Harb Perspect Biol. 2013;5.

doi:10.1101/cshperspect.a008748

65.   Wang X, Yang C. Programmed cell death and clearance of cell corpses in Caenorhabditis

elegans. Cellular and Molecular Life Sciences. Birkhauser Verlag AG; 2016. pp. 2221–

2236. doi:10.1007/s00018-016-2196-z

66.   Reddien PW, Horvitz HR. THE ENGULFMENT PROCESS OF PROGRAMMED CELL DEATH IN

CAENORHABDITIS ELEGANS. Annu Rev Cell Dev Biol. Annual Reviews; 2004;20: 193–221.

doi:10.1146/annurev.cellbio.20.022003.114619

67.   Hoeppner DJ, Hengartner MO, Schnabel R. Engulfment genes cooperate with ced-3 to

promote cell death in Caenorhabditis elegans. Nature. Nature Publishing Group;

2001;412: 202–206. doi:10.1038/35084103

68.   Geng X, Zhou QH, Kage-Nakadai E, Shi Y, Yan N, Mitani S, et al. Caenorhabditis elegans

caspase homolog CSP-2 inhibits CED-3 autoactivation and apoptosis in germ cells. Cell

Death Differ. 2009;16: 1385–1394. doi:10.1038/cdd.2009.88

69.   Yang X, Chang HY, Baltimore D. Essential role of CED-4 oligomerization in CED-3

activation and apoptosis. Science (80-). American Association for the Advancement of

Science; 1998;281: 1355–1357. doi:10.1126/science.281.5381.1355

70.   Parsons MJ, Green DR. Mitochondria in cell death. Essays Biochem. 2010;47: 99–114.

doi:10.1042/BSE0470099

71.   Harman D. Free radical theory of aging. Mutat Res. 1992;275: 257–266.

doi:10.1016/0921-8734(92)90030-S

72.   HARMAN D. Aging: a theory based on free radical and radiation chemistry. J Gerontol.

1956;11: 298–300. doi:10.1093/geronj/11.3.298

73.   Szczepanowska K, Trifunovic A. Origins of mtDNA mutations in ageing. Essays in

Biochemistry. Portland Press Ltd; 2017. pp. 325–337. doi:10.1042/EBC20160090

74.   Bratic A, Larsson N-G. The role of mitochondria in aging. J Clin Invest. 2013;123: 951–957.

doi:10.1172/JCI64125

75.   Ziegler D V., Wiley CD, Velarde MC. Mitochondrial effectors of cellular senescence:

Beyond the free radical theory of aging. Aging Cell. Blackwell Publishing Ltd; 2015. pp. 1–

7. doi:10.1111/acel.12287

76.   Kimble J, Crittenden SL. Germline proliferation and its control. WormBook : the online

review of C. elegans biology. 2005. pp. 1–14. doi:10.1895/wormbook.1.13.1

77.   Yoon DS, Alfhili MA, Friend K, Lee MH. MPK-1/ERK regulatory network controls the

number of sperm by regulating timing of sperm-oocyte switch in C. elegans germline.

Biochem Biophys Res Commun. Elsevier B.V.; 2017;491: 1077–1082.

doi:10.1016/j.bbrc.2017.08.014

78.   Angeles-Albores D, Leighton DHW, Tsou T, Khaw TH, Antoshechkin I, Sternberg PW. The

Caenorhabditis elegans female-like state: Decoupling the transcriptomic effects of aging

and sperm status. G3 Genes, Genomes, Genet. Genetics Society of America; 2017;7:

2969–2977. doi:10.1534/g3.117.300080

79.   Kujoth CC, Hiona A, Pugh TD, Someya S, Panzer K, Wohlgemuth SE, et al. Medicine:

Mitochondrial DNA mutations, oxidative stress, and apoptosis in mammalian aging.

Science (80-). American Association for the Advancement of Science; 2005;309: 481–

484. doi:10.1126/science.1112125

80.   Beanan MJ, Strome S. Characterization of a germ-line proliferation mutation in C.

elegans. Development. 1992;116.

81.   Barton MK, Schedl TB, Kimble J. Gain-of-function mutations of fem-3, a sex-

determination gene in Caenorhabditis elegans. Genetics. 1987;115: 107–119.

82.   Bartke A. Insulin and aging. Cell Cycle. Taylor and Francis Inc.; 2008. pp. 3338–3343.

doi:10.4161/cc.7.21.7012

83.   Anisimov VN, Bartke A. The key role of growth hormone-insulin-IGF-1 signaling in aging

and cancer. Critical Reviews in Oncology/Hematology. 2013. pp. 201–223.

doi:10.1016/j.critrevonc.2013.01.005

84.   Piper MDW, Selman C, McElwee JJ, Partridge L. Separating cause from effect: how does

insulin/IGF signalling control lifespan in worms, flies and mice? J Intern Med. John Wiley

& Sons, Ltd; 2008;263: 179–191. doi:10.1111/J.1365-2796.2007.01906.X

85.   Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A C. elegans mutant that lives twice

as long as wild type. Nature. 1993;366: 461–464. doi:10.1038/366461a0

86.   Tatar M, Kopelman A, Epstein D, Tu M-P, Yin C-M, Garofalo RS. A Mutant Drosophila

Insulin Receptor Homolog That Extends Life-Span and Impairs Neuroendocrine Function.

Science (80-). American Association for the Advancement of Science; 2001;292: 107–

110. doi:10.1126/SCIENCE.1057987

87.   Blüher M, Kahn BB, Kahn CR. Extended Longevity in Mice Lacking the Insulin Receptor in

Adipose Tissue. Science (80-). American Association for the Advancement of Science;

2003;299: 572–574. doi:10.1126/SCIENCE.1078223

88.   Gitschlag BL, Tate AT, Patel MR. Nutrient status shapes selfish mitochondrial genome

dynamics across different levels of selection. Elife. eLife Sciences Publications Ltd;

2020;9: 1–27. doi:10.7554/ELIFE.56686

89.   Apfeld J, O'Connor G, McDonagh T, DiStefano PS, Curtis R. The AMP-activated protein

kinase AAK-2 links energy levels and insulin-like signals to lifespan in C. elegans. Genes

Dev. Cold Spring Harbor Laboratory Press; 2004;18: 3004–3009.

doi:10.1101/gad.1255404

90.   Miyadera H, Kano K, Miyoshi H, Ishii N, Hekimi S, Kita K. Quinones in long-lived clk-1

mutants of Caenorhabditis elegans. FEBS Lett. No longer published by Elsevier; 2002;512:

33–37. doi:10.1016/S0014-5793(02)02282-2

91.   Miyadera H, Amino H, Hiraishi A, Taka H, Murayama K, Miyoshi H, et al. Altered Quinone

Biosynthesis in the Long-lived clk-1Mutants of Caenorhabditis elegans. J Biol Chem.

Elsevier; 2001;276: 7713–7716. doi:10.1074/JBC.C000889200

92.   Kirby CS, Patel MR. Elevated mitochondrial DNA copy number found in ubiquinone-

deficient clk-1 mutants is not rescued by ubiquinone precursor 2-4-dihydroxybenzoate.

Mitochondrion. Elsevier B.V.; 2021;58: 38–48. doi:10.1016/j.mito.2021.02.001

93.   Charmpilas N, Tavernarakis N. Mitochondrial maturation drives germline stem cell

differentiation in Caenorhabditis elegans. Cell Death Differ. Springer Nature; 2020;27:

601–617. doi:10.1038/s41418-019-0375-9

94.   Folmes CDL, Ma H, Mitalipov S, Terzic A. Mitochondria in pluripotent stem cells:

Stemness regulators and disease targets. Current Opinion in Genetics and Development.

Elsevier Ltd; 2016. pp. 1–7. doi:10.1016/j.gde.2016.02.001

95.   Floros VI, Pyle A, DIetmann S, Wei W, Tang WWC, Irie N, et al. Segregation of

mitochondrial DNA heteroplasmy through a developmental genetic bottleneck in human

1219   embryos. Nat Cell Biol. 2018; 144–151. doi:10.1038/s41556-017-0017-8

1220 96. Zhang H, Burr SP, Chinnery PF. The mitochondrial DNA genetic bottleneck: Inheritance

1221   and beyond. Essays in Biochemistry. Portland Press Ltd; 2018. pp. 225–234.

1222   doi:10.1042/EBC20170096

1223 97. Wai T, Teoli D, Shoubridge EA. The mitochondrial DNA genetic bottleneck results from

1224   replication of a subpopulation of genomes. Nat Genet. 2008; 1484–1488.

1225   doi:10.1038/ng.258

1226 98. Liau WS, Gonzalez-Serricchio AS, Deshommes C, Chin K, LaMunyon CW. A persistent

1227   mitochondrial deletion reduces fitness and sperm performance in heteroplasmic

1228   populations of C. elegans. BMC Genet. BioMed Central; 2007;8: 1–11. doi:10.1186/1471-

1229   2156-8-8

1230 99. Zhu Z, Han X, Wang Y, Liu W, Lu Y, Xu C, et al. Identification of Specific Nuclear Genetic

1231   Loci and Genes That Interact With the Mitochondrial Genome and Contribute to

1232   Fecundity in Caenorhabditis elegans. Front Genet. Frontiers; 2019;10: 28.

1233   doi:10.3389/fgene.2019.00028

1234 100. Levin L, Blumberg A, Barshad G, Mishmar D. Mito-nuclear co-evolution: The positive and

1235   negative sides of functional ancient mutations. Front Genet. Frontiers Media S.A.;

1236   2014;5: 448. doi:10.3389/fgene.2014.00448

1237 101. Paliwal S, Fiumera AC, Fiumera HL. Mitochondrial-nuclear epistasis contributes to

1238   phenotypic variation and coadaptation in natural isolates of saccharomyces cerevisiae.

1239   Genetics. Genetics Society of America; 2014;198: 1251–1265.

1240   doi:10.1534/genetics.114.168575

102.   Zhu CT, Ingelmo P, Rand DM. G×G×E for Lifespan in Drosophila: Mitochondrial, Nuclear,

and Dietary Interactions that Modify Longevity. PLoS Genet. Public Library of Science;

2014;10. doi:10.1371/journal.pgen.1004354

103.   Meiklejohn CD, Holmbeck MA, Siddiq MA, Abt DN, Rand DM, Montooth KL. An

Incompatibility between a Mitochondrial tRNA and Its Nuclear-Encoded tRNA Synthetase

Compromises Development and Fitness in Drosophila. PLoS Genet. Public Library of

Science; 2013;9. doi:10.1371/journal.pgen.1003238

104.   Dingley SD, Polyak E, Ostrovsky J, Srinivasan S, Lee I, Rosenfeld AB, et al. Mitochondrial

DNA Variant in COX1 Subunit Significantly Alters Energy Metabolism of Geographically

Divergent Wild Isolates in Caenorhabditis elegans. J Mol Biol. 2014;426: 2199–2216.

doi:10.1016/j.jmb.2014.02.009

105.   Yee C, Yang W, Hekimi S. The intrinsic apoptosis pathway mediates the pro-longevity

response to mitochondrial ROS in C. elegans. Cell. PMC Canada manuscript submission;

2014;157: 897. doi:10.1016/J.CELL.2014.02.055

106.   H H, C K. Signals from the reproductive system regulate the lifespan of C. elegans.

Nature. Nature; 1999;399: 362–366. doi:10.1038/20694

107.   L D, CW A, S K. New insights into apoptosome structure and function. Cell Death Differ.

Cell Death Differ; 2018;25: 1194–1208. doi:10.1038/S41418-017-0025-Z

108.   S Y, X Y, JM A, JE H, SJ L, CW A. The holo-apoptosome: activation of procaspase-9 and

interactions with caspase-3. Structure. Structure; 2011;19: 1084–1096.

doi:10.1016/J.STR.2011.07.001

109.   Zermati Y, Mouhamad S, Stergiou L, Besse B, Galluzzi L, Boehrer S, et al. Nonapoptotic

Role for Apaf-1 in the DNA Damage Checkpoint. Mol Cell. Cell Press; 2007;28: 624–637.

doi:10.1016/j.molcel.2007.09.030

110.	Ferraro E, Pesaresi MG, De Zio D, Cencioni MT, Gortat A, Cozzolino M, et al. Apaf1 plays a

pro-survival role by regulating centrosome morphology and function. J Cell Sci. 2011;124:

3450–3463. doi:10.1242/jcs.086298

111.	Liu WH, Lin YL, Wang JP, Liou W, Hou RF, Wu YC, et al. Restriction of vaccinia virus

replication by a ced-3 and ced-4-dependent pathway in Caenorhabditis elegans. Proc

Natl Acad Sci U S A. National Academy of Sciences; 2006;103: 4174–4179.

doi:10.1073/pnas.0506442103

112.	Wang G, Sun L, Reina CP, Song I, Gabel C V., Driscoll M. CED-4 CARD domain residues can

modulate non-apoptotic neuronal regeneration functions independently from apoptosis.

Sci Rep. 2019;9: 13315. doi:10.1038/s41598-019-49633-9

113.	Chaudhary D, O'Rourke K, Chinnaiyan AM, Dixit VM. The Death Inhibitory Molecules CED-

9 and CED-4L Use a Common Mechanism to Inhibit the CED-3 Death Protease. J Biol

Chem. Elsevier; 1998;273: 17708–17712. doi:10.1074/JBC.273.28.17708

114.	Mack HID, Zhang P, Fonslow BR, Yates JR. The protein kinase MBK-1 contributes to

lifespan extension in daf-2 mutant and germline-deficient Caenorhabditis elegans. Aging

(Albany NY). Impact Journals LLC; 2017;9: 1414–1432. doi:10.18632/aging.101244

115.	Libina N, Berman JR, Kenyon C. Tissue-Specific Activities of C. elegans DAF-16 in the

Regulation of Lifespan. Cell. Cell Press; 2003;115: 489–502. doi:10.1016/S0092-

8674(03)00889-4

116.	Lapierre LR, Hansen M. Lessons from C. elegans: signaling pathways for longevity. Trends

Endocrinol Metab. Elsevier Current Trends; 2012;23: 637–644.

doi:10.1016/J.TEM.2012.07.007

117.   A A. Regulation of longevity by the reproductive system. Exp Gerontol. Exp Gerontol;

2013;48: 596–602. doi:10.1016/J.EXGER.2012.09.009

118.   Branicky R, Bénard C, Hekimi S. clk-1, mitochondria, and physiological rates. BioEssays.

2000. pp. 48–56. doi:10.1002/(SICI)1521-1878(200001)22:1<48::AID-BIES9>3.0.CO;2-F

119.   Hekimi S, Guarente L. Genetics and the specificity of the aging process. Science. 2003. pp.

1351–1354. doi:10.1126/science.1082358

120.   Blackwell TK, Steinbaugh MJ, Hourihan JM, Ewald CY, Isik M. SKN-1/Nrf, stress responses,

and aging in Caenorhabditis elegans. Free Radic Biol Med. Pergamon; 2015;88: 290–301.

doi:10.1016/J.FREERADBIOMED.2015.06.008

121.   Youle RJ, Van Der Bliek AM. Mitochondrial fission, fusion, and stress. Science. American

Association for the Advancement of Science; 2012. pp. 1062–1065.

doi:10.1126/science.1219855

122.   Twig G, Elorza A, Molina AJA, Mohamed H, Wikstrom JD, Walzer G, et al. Fission and

selective fusion govern mitochondrial segregation and elimination by autophagy. EMBO

J. 2008;27: 433–446. doi:10.1038/sj.emboj.7601963

123.   Stiernagle T. Maintenance of C. elegans. WormBook : the online review of C. elegans

biology. 2006. pp. 1–11. doi:10.1895/wormbook.1.101.1

124.   C. elegans Deletion Mutant Consortium. large-scale screening for targeted knockouts in

the Caenorhabditis elegans genome. G3 (Bethesda). 2012;2: 1415–25.

doi:10.1534/g3.112.003830

125.   Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform.

1308   Bioinformatics. Oxford Academic; 2009;25: 1754–1760.

1309   doi:10.1093/BIOINFORMATICS/BTP324

1310   126.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

1311   Alignment/Map format and SAMtools. Bioinformatics. Oxford Academic; 2009;25: 2078–

1312   2079. doi:10.1093/BIOINFORMATICS/BTP352

1313   127.   Koboldt DC, Chen K, Wylie T, Larson DE, McLellan MD, Mardis ER, et al. VarScan: variant

1314   detection in massively parallel sequencing of individual and pooled samples.

1315   Bioinformatics. Oxford Academic; 2009;25: 2283–2285.

1316   doi:10.1093/BIOINFORMATICS/BTP373

1317

1318