Mechanisms Governing Target Search and Binding Dynamics of Hypoxia-
Inducible Factors

Yu Chen[1,2,3], Claudia Cattoglio[1,2,3], Gina Dailey[1,3], Qiulin Zhu[1,3], Robert Tjian[1,2,3] *, and Xavier
Darzacq[1,3] *

[1] Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA.
[2] Howard Hughes Medical Institute, University of California, Berkeley, CA, USA.
[3] Li Ka Shing Center for Biomedical & Health Sciences, University of California, Berkeley, CA,
USA.

*Correspondence: R.T. (jmlim@berkeley.edu), X.D. (darzacq@berkeley.edu)

Abstract

Transcription factors (TFs) are classically attributed a modular construction, containing well-
structured sequence specific DNA-binding domains (DBDs) paired with disordered activation
domains (ADs) responsible for protein-protein interactions targeting cofactors or the core
transcription initiation machinery. However, this simple division of labor model struggles to explain
why TFs with identical DNA binding sequence specificity determined *in vitro* exhibit distinct
binding profiles *in vivo*. The family of Hypoxia-Inducible Factors (HIFs) offer a stark example:
aberrantly expressed in several cancer types, HIF-1α and HIF-2α subunit isoforms recognize the
same DNA motif *in vitro* – the hypoxia response element (HRE) – but only share a subset of their
target genes *in vivo*, while eliciting contrasting effects on cancer development and progression
under certain circumstances. To probe the mechanisms mediating isoform-specific gene
regulation, we used live cell single particle tracking (SPT) to investigate HIF nuclear dynamics
and how they change upon genetic perturbation or drug treatment. We found that HIF-α subunits
and their dimerization partner HIF-1β exhibit distinct diffusion and binding characteristics that are
exquisitely sensitive to concentration and subunit stoichiometry. Using domain-swap variants,
mutations, and a HIF-2α specific inhibitor, we found that although the DBD and dimerization
domains are important, a major determinant of chromatin binding and diffusion behavior is the
AD-containing intrinsically disordered region (IDR). Using Cut&Run and RNA-seq as orthogonal
genomic approaches we also confirmed IDR-dependent binding and activation of a specific
subset of HIF-target genes. These findings reveal a previously unappreciated role of IDRs in

regulating the TF search and binding process that contribute to functional target site selectivity on

chromatin.

Introduction

Sequence-specific transcription factors (TFs) are key frontline regulators of gene expression. Classical LexA-Gal4 domain-swap experiments in yeast presented a simple modular structure and apparent division of labor for typical TFs (Brent and Ptashne, 1985). In this textbook paradigm, the DNA-binding domain (DBD) is responsible for DNA sequence recognition and binding specificity while the activation domain (AD) is responsible for target gene transactivation that involves protein-protein interactions with co-factors, the basal transcription machinery and other ancillary factors that are generally devoid of sequence specific DNA recognition. In higher eukaryotes, each DBD class usually contains multiple closely related family members. For example, the bHLH class of TFs includes MyoD, Clock, and Max. They all recognize the same E-box DNA binding sequence motif 5'-CACGTG-3', yet each differentially regulates muscle differentiation, circadian rhythm, and cell proliferation, respectively (Kribelbauer et al., 2019). This raises the specificity paradox: how do TFs with seemingly identical DNA sequence specificity, at least as determined *in vitro*, nevertheless exhibit non-overlapping binding profiles *in vivo* and carry out distinct and even opposing functions? In general, when confronted with this conundrum, we have assumed that one or more co-factors or perhaps still to be identified "silent partner" TFs can somehow divert target recognition to a composite cis-regulatory site distinct from the canonical DNA binding site. Given the high occurrence of short binding motifs for most TFs throughout the genome, even with co-operative binding to composite sites, most potential specific binding sites nevertheless remain unoccupied as determined by genome-wide TF binding studies. What feature or motif within TFs outside of the DBD and dimerization domain may be responsible for such differential site selection has remained unclear. Thus, the simple rule of modular units with well separated divisions of labor between DBD, dimerization and transactivation may deserve a closer look. We also wondered whether quantitative single molecule dynamics measurements might reveal new aspects of TF behavior in living cells that could inform us regarding potential mechanisms influencing the target search and binding process and differential site selectivity in a native physiologically relevant context.

Here we have chosen the Hypoxia-Inducible Factors (HIFs) as a representative example to study the paradox of highly conserved DBDs carrying out distinct target site selection and to dissect potential novel features of TFs that mediate chromatin binding. HIFs are a family of α/β heterodimeric TFs stabilized under hypoxic conditions to promote angiogenesis, anaerobic metabolism, cell proliferation and "stemness" (Semenza, 2012). The oxygen-labile alpha subunits (mainly HIF-1α and HIF-2α) complex with their oxygen-stable beta partner (mainly HIF-1β) to form

a functional dimer (Figure 1A). All HIF subunit isoforms belong to the bHLH-PAS (Basic helix-

loop-helix-PER-ARNT-SIM) family, where the N-termini are structured domains containing bHLH

(DNA binding) and PAS (dimerization) domains, while the C-termini consist of intrinsically

disordered regions (IDRs) containing ADs (Figure 1A; Figure 1—figure supplement 1A). HIF-

1α/1β and HIF-2α/1β dimers share a conserved structural fold (Wu et al., 2015), recognize the

same hypoxia response element (HRE) 5'-TACGTG-3' binding motif (Schödel et al., 2011;

Wenger et al., 2005), but share only a partial overlap of target genes *in vivo* (Smythies et al.,

2019). With their own unique target gene sets, HIF-1α and -2α can exert divergent and even

contrasting functions (Keith et al., 2012). For example, while both HIF-1α and HIF-2α regulate

angiogenesis, HIF-1α specifically regulates glycolysis, apoptosis, and promotes NO production,

whereas HIF-2α binds to the *POU5F1* locus to maintain Oct4-regulated stem cell identity and

pluripotency, promotes cell cycle progression, and inhibits NO production (Keith et al., 2012, p. 1;

Smythies et al., 2019). Therefore, our current simple textbook model of exchangeable modular

TF functional units does not satisfactorily explain such isoform-specific target gene regulation.

The HIF family differential specificity paradox is even more daunting to comprehend at the level

of disease inducing mechanisms. HIFs are aberrantly upregulated and recognized as oncogenic

drivers in multiple cancers. However, in addition to their shared roles in cancer onset and

progression, HIF-1α and -2α also show many independent, sometimes even opposing roles in

specific contexts (Keith et al., 2012). For example, in clear cell renal cell carcinoma (ccRCC), HIF-

2α is the critical tumorigenic driver whereas HIF-1α, in contrast with its usual tumorigenic role, is

mostly tumor-suppressive (Raval et al., 2005; Schödel et al., 2016). The regulatory mechanism

behind such highly divergent outcomes is still largely unknown. Given such complexity, without a

deeper understanding of isoform-specific transcriptional regulation, it is hard to predict the

functional outcomes mediated by individual HIF isoforms in various cancer types or stages, which

could be a complicating factor in developing more effective HIF-targeting cancer therapeutics.

In this study, we aim to understand the molecular mechanisms mediating isoform-specific target

gene regulation at its most fundamental level – could we detect differential molecular dynamics

of distinct TF isoforms during the target search and chromatin binding process in live cells? Which

regions or domains of TFs might be responsible for such isoform specific properties? Could we

begin to discern possible mechanisms that guide TFs with highly conserved DBDs to their distinct

and specific targets beyond cognate DNA sequence recognition? Here we use HIFs as an

illustrative example, combining endogenous tagging and super-resolution single particle tracking

(SPT) (Liu et al., 2015) to study the dynamic behavior of these key gene regulators in live cells
under physiological conditions. We also dissect the contribution of different domains of HIF-α
isoforms by a series of mutation and domain swap experiments to directly test the concept of
modular functional domains. Deploying a combination of genetic and small-molecule
perturbations, we found that, although HIF DBD and dimerization are important for DNA target
acquisition, the amount of protein bound and its diffusion characteristics are mainly driven by
regions outside the DBD and dimerization domains. Finally, using genomic approaches we found
that, in concordance with our imaging results, binding strength and gene activation are IDR-
dependent for a subset of HIF target sites. Our results reveal a previously unappreciated role of
unstructured domains in the target search and binding properties of TFs to functional chromatin
sites in a live cancer cell context.
115
116 **Results**
117
Establishing a human cancer cell system for live-cell single molecule imaging of HIF
To investigate HIF dynamics, we first focused on one of the cognate dimers: HIF-2α/1β. We used
the common ccRCC line 786-O (Brodaczewska et al., 2016), derived from a VHL-deficient, H2
type primary clear cell renal cell carcinoma, wherein HIF-2α is stabilized due to an inactivating
mutation in VHL (the E3 ubiquitin ligase that targets all HIF-α isoforms for proteasomal
degradation)(Gnarra et al., 1994). The 786-O line also conveniently lacks any functional HIF-1α
due to a truncating mutation of HIF-1α (Shen et al., 2011; Swiatek et al., 2020, p.), which allows
us to study one α isoform independently from the other. Using CRISPR/Cas9-mediated genome
editing, we successfully generated several clonal lines with homozygous knock-in (KIN) of the
HaloTag (Los et al., 2008) at the N-terminus of either HIF-2α or its binding partner, HIF-1β (Figure
1B; Figure 1—figure supplement 2). Western blotting confirmed that the tagged proteins are
expressed at levels similar to wild-type (WT) in unedited cells (e.g., HIF-2α clone A31 and HIF-
1β clone A21) (Figure 1B). Confocal imaging after covalently labeling cells with a fluorescent
Halo-binding ligand (JFX646) (Grimm et al., 2021) shows the expected nuclear localization for
both Halo-HIF-2α and Halo-HIF-1β proteins (Figure 1C). In addition, we confirmed by ChIP-seq
that both tagged proteins maintain a similar genome-wide binding profile as the WT protein in
unedited cells (Figure 1—figure supplement 3A), and RNA-seq confirmed that gene expression
profiles in both edited cell lines are not significantly altered from WT unedited cells (Figure 1—
figure supplement 3B-C, with cells overexpressing WT and mutant HIF-α as controls). We have

thus established a human cancer cell system suitable for live-cell imaging of HIF-2α and HIF-1β

at endogenous expression levels.

To evaluate how HIF-2α and -1β explore the nucleus and bind DNA, we used the fast modality of

super resolution live cell single particle tracking (fSPT) that is capable of tracking rapidly diffusing

molecules. Cells with either HIF-2α or HIF-1β Halo KIN were doubly labeled with the live-cell

permeable Halo-binding JFX dyes (Figure 1D) and were imaged under highly inclined and

laminated optical sheet illumination (HiLo) (Tokunaga et al., 2008) at high frame rates (~182 Hz)

to capture the movement of single molecules in their native nuclear environment (Figure 1E).

Stroboscopic illumination at high excitation power is used to minimize motion blur, while sparse

labeling ensures only a limited number of molecules are detected at any given time in the nucleus

to minimize misconnections when computing the path of individual molecules (trajectories) (Figure

1E-F). We can then estimate relevant kinetic parameters from these trajectories, extracting

quantitative information such as diffusion coefficients and bound fraction.

fSPT detects various HIF molecular states in their native nuclear environment

To quantitatively analyze the acquired fSPT data, we used a non-parametric Bayesian "state array"

(SA) approach (Heckert et al., 2021) recently developed in our lab. Briefly, SA analyzes regular

Brownian motion with normally distributed localization error (RBME) with a two-dimensional "grid

of states" that spans a range of diffusion coefficients (first dimension) and localization error

(second dimension) magnitudes, and evaluates the occupancy (proportion) of each state in this

array using the observed fSPT trajectory data and a variational Bayesian algorithm. After

inference, the proportion of molecules as a function of diffusion coefficient is calculated by

marginalizing localization error, creating a "diffusion spectrum" that reports the underlying

molecular subpopulations according to their proportions (y-axis) and their diffusive properties

(diffusion coefficients, x-axis) (Figure 2—figure supplement 1). While our previous approach

required the assumption of a fixed and limited (3 or less) number of states for the tracked protein

(Hansen et al., 2018), this SA approach does not require *a priori* knowledge or any assumption

regarding the number of subpopulations or "states" of the tracked molecules. Also, it is more

robust for fSPT trajectories as it accounts for known experimental biases due to localization error

and fluorophore defocalization (Heckert et al., 2021). We chose to use this new method, because

as dimeric transcription factors, HIF subunits could conceivably exist in many states (e.g., bound,

moving as a monomer, moving as a dimer, and moving in a bigger complex containing co-

regulators) (Figure 2A-B) and the SA approach better suits our needs for model-free analysis

without assuming any given number of states.

We first analyzed individual cells to check cell-to-cell variation. Due to the small number of

trajectories per cell, we used a less precise version of the SA calculation, the "naïve occupation

estimate" by simply applying the RBME likelihood estimation to individual cells without refining

the posterior over diffusion coefficient and localization error (Figure 2—figure supplement 1A-B,

top) (Heckert et al., 2021). We observed rather heterogeneous results for both HIF-2α and HIF-

1β, with varied diffusion coefficient estimates from cell to cell (Figure 2A-B; Figure 2—figure

supplement 2A, clones A31 and A21). While the heterogeneity we observed was likely due to the

limited number of trajectories collected from each cell (i.e. small sample bias) rather than to a

difference in behavior of HIF proteins in each cell, our findings do indicate that a range of moving

states likely exists for both HIF proteins.

We then pooled trajectories from many cells (n ~60 from 3 biological replicates for each

experimental condition) to estimate the distribution of diffusion coefficients for the population. SA

generates a distribution of diffusion coefficient estimates that reports the fraction of stably bound

molecules while simultaneously displaying the full behavioral spectrum of the diffusing molecules

(Figure 2C; Figure 2—figure supplement 1A-B, bottom). We define the fraction with a diffusion

coefficient < 0.1 µm^2/sec as immobile and presumably chromatin-bound (see Figure 2—figure

supplement 3 and Appendix 1 for discussion of source of variations between replicates). Strikingly,

we observed a very different behavior for Halo-HIF-2α compared to Halo-HIF-1β. Whereas a large

fraction (about 40%) of Halo-HIF-2α is bound, the majority (above 70%) of Halo-HIF-1β appears

freely diffusing (Figure 2C; Figure 2—figure supplement 2B, clones A31 and A21). Also, the

overall diffusion coefficient for the Halo-HIF-1β mobile population is much larger than that of Halo-

HIF-2α. We repeated measurements in different KIN clones and confirmed the reproducibility of

these results for both Halo-HIF-2α (Figure 2—figure supplement 2A-B, clone B50) and Halo-HIF-

1β (Figure 2—figure supplement 2A-B, clone B89). Note that although we quantitatively compared

the bound fraction, due to the complexity of the composition of the moving population (i.e. multiple

states might exist but are merged into a single peak), quantitatively comparing mode (i.e. peak)

or mean diffusion coefficient may give slightly different results. Therefore, for the rest of the paper

we only present the peak diffusion coefficient in the figures but listed both peak diffusion

coefficient and mean diffusion coefficient for the moving population in Table 1.

The differences between HIF-2α and -1β seem counterintuitive at first, because one would expect
HIF-2α and HIF-1β to behave similarly since they should exist as a hetero-dimer. However, since
the endogenous HIF-1β is expressed at a much higher level than HIF-2α (Figure 1—figure
supplement 2C), the majority of HIF-1β is likely free to diffuse without HIF-2α. Of note, the
distribution plot only reflects the fraction of molecules as a function of their diffusion coefficient,
but does not report on the absolute number of molecules. Therefore, a smaller bound fraction for
Halo-HIF-1β does not mean fewer numbers of bound molecules than Halo-HIF-2α, since many
more Halo-HIF-1β molecules are present in the nucleus. Given this scenario, we hypothesized
that HIF-1β molecular dynamics and percent binding should be modulated by changing the 2α/1β
stoichiometry.
214
HIF-1β binding and diffusing dynamics can be modulated by HIF-α:β stoichiometry, and
are dependent on dimerization
To test the hypothesis that HIF-1β dynamics depends on 2α/1β ratio, we first tried to modulate its
behavior by stably over-expressing HIF-2α in the endogenously HIF-1β Halo-tagged KIN line
(Figure 2D). We found that the mobile population of Halo-HIF-1β diffuses more slowly when HIF-
2α is overexpressed, most likely due to its dimerization with the extra HIF-2α to form dimers
capable of DNA/chromatin binding. As expected, we also observed a significant increase in the
Halo-HIF-1β bound fraction (up to 50%), (Figure 2E; Figure 2—figure supplement 2C, top and
middle). To confirm that the changes in HIF-1β dynamics caused by increasing levels of HIF-2α
are dependent on hetero-dimerization, we stably overexpressed the HIF-2α R171A/V192D double
mutant (HIF-2α DM) that was previously reported to lose its dimerization capability with HIF-1β
(Wu et al., 2015). As expected, overexpression of HIF-2α DM did not increase the bound fraction
or decrease the overall diffusion coefficient of Halo-HIF-1β to the same extent seen with WT HIF-
2α overexpression (Figure 2E; Figure 2—figure supplement 2C, bottom), suggesting that the
changes we observe are dimerization-dependent.
230
We further validated our results by stably overexpressing different forms of HIF-α in the HIF-1β
Halo-tagged KIN line and treating cells with a HIF-2α-specific small molecule inhibitor, Belzutifan
(PT-2977). Belzutifan inhibits HIF-2α/1β, but not HIF-1α/1β, dimerization by specifically binding
to the dimerization domain of HIF-2α (Figure 3—figure supplement 1A), and thus has been used
as an HIF-2α inhibitor for ccRCC treatment (Wallace et al., 2016; Xu et al., 2019). We first
confirmed that Belzutifan inhibits HIF-2α transcription function in a dose-dependent manner
(Figure 3—figure supplement 1B). Importantly, Belzutifan also reduces the HIF-2α bound fraction

in the HIF-2α Halo-tagged KIN line in a similar dosage-dependent manner, again revealing the

potential of fSPT to measure TF dynamics and associated functional changes (Figure 3—figure

supplement 1C-E). We choose to use 0.2 µM Belzutifan for all subsequent experiments to

maximize its effect.

Next, we carried out a series of experiments designed to probe the consequences of swapping

different functional domains of HIF-1α and HIF-2α to determine which parts of these closely

related TFs might be involved in selective activities when paired with HIF-1β. Using the HIF-1β

Halo-tagged KIN line as the parental line, we stably overexpressed WT or chimeric HIF-α, where

we swapped the structured and disordered domains between HIF-1α and HIF-2α (Figure 3A). All

these different HIF-α variants are expressed from a relatively strong EF-1alpha promoter and are

N-terminally 3xFLAG-tagged. A construct that expresses 3xFLAG only is used as control. We

then treated these cells with either Belzutifan or DMSO control and measured Halo-HIF-1β

dynamics (Figure 3A). While 3xFLAG tag had no effect, overexpressing HIF-α, regardless of

which variant form, is able to both increase the bound fraction and reduce the overall diffusion

coefficient of HIF-1β (Figure 3B, top, and 3C, DMSO group). For cells overexpressing the α

variants that contain the HIF-2α structured domain, this effect on HIF-1β can be at least partially

reverted after Belzutifan treatment (Figure 3B-C, +HIF-2α and +HIF-2α/1α). In contrast, for cells

overexpressing the α variants that contain the HIF-1α structured domain, this effect is resistant to

Belzutifan, consistent with the subunit isoform specificity of the drug for HIF-2α (Figure 3B-C,

+HIF-1α and +HIF-1α/2α). In untransfected and 3xFLAG only overexpressing control cells,

treatment with Belzutifan only weakly reduces the HIF-1β bound fraction, again suggesting that

the majority of HIF-1β is not engaged with 2α (Figure 3B-C, parental cell and +3xF). Overall, these

results demonstrate that HIF-1β dynamics change after engagement with its α partner and can

be selectively inhibited with a specific dimerization inhibitor. The observed differences also

confirm that fSPT is a powerful platform to monitor molecular dynamic changes of TFs in living

cells thus, allowing us to gain new mechanistic insights while we introduce various perturbations,

such as subunit concentration or stoichiometry and specific mutations.

Regions outside the DBD/dimerization domain determine HIF molecular dynamics

Interestingly, comparing the effects of the four different α variants, we found that regardless of

their structured domain, those with the same C terminal IDRs behave similarly (Figure 3B-C,

middle and right). Specifically, the variants containing the HIF-2α IDR have a stronger effect on

increasing HIF-1β binding than the variants containing the HIF-1α IDR. Thus, surprisingly, it

appears that the bound fraction of HIF-1β is not determined by the HIF-α DBD, but rather by HIF-
α IDR, which we found rather counterintuitive. To confirm the importance of HIF-α IDRs in HIF
binding, we overexpressed a truncated version of either HIF-1α or -2α that contains only the N-
terminal structured region (HIF-1α NT or HIF-2α NT), which still maintains both the DBD and
dimerization capability for interacting with HIF-1β (Wu et al., 2015). Indeed, both these truncated
forms lacking the IDR/AD of HIF-α minimally affect the HIF-1β bound fraction (Figure 3—figure
supplement 2). Surprisingly, these truncated HIF-α variants also only marginally influenced the
overall HIF-1β diffusion coefficient. These results indicate that dimerization alone neither
increases HIF-1β binding nor reduces the overall diffusion coefficient of its moving population.
Instead, the extended HIF-α AD-containing IDR appears necessary to influence and direct HIF-
1β behavior.
283
To further test our hypothesis that HIF chromatin binding and the dynamics of the diffusion
population are dominated by the α subunit IDR, we switched to image the α subunit itself. We
made different forms of Halo-tagged HIF-α (WT and domain-swapped), stably but weakly
expressed them in WT 786-O cells with an L30 promoter (Figure 4A). To minimize differences in
expression levels among different HIF-α forms, we selected cells with a roughly similar
fluorescence intensity in the "cell-picking channel" (Figure 1D, left). Later we verified that the
movies captured in the "fSPT tracking channel" (Figure 1D, right) have similar localization density
in the initial 10 frames (Figure 4—figure supplement 1; see also Appendix 2 for full discussion).
We confirmed that when controlled for similar expression levels, binding and diffusion
characteristics of L30-expressed Halo-HIF-2α are very similar to the endogenous Halo-HIF-2α in
the KIN line (Figure 4—figure supplement 2), demonstrating that weak overexpression can largely
recapitulate endogenous protein behavior. Therefore, this system provides a convenient tool to
investigate the contribution of each domain of HIF-α in the target search and binding process.
297
Much like our results with endogenous HIF-1β, we observed similar behaviors of HIF-α proteins
if they contain the same IDR (Figure 4B, top and middle), while displaying distinct behaviors when
endowed with different IDR isoforms (Figure 4B, bottom). Regardless of which DBD they have,
the variants containing the HIF-2α IDR (WT HIF-2α and HIF-1α/2α) show a higher bound fraction,
compared to the ones containing HIF-1α IDR (WT HIF-1α and HIF-2α/1α) (Figure 4C). These
results suggest that indeed the disordered region on HIF-α determines how HIFs bind and diffuse
in the nucleus, and that the HIF-2α AD-containing IDR mediates more or stronger binding to
chromatin and/or some other relatively immobile components in 786-O cells.

HIF-α disordered region is necessary but not sufficient for optimal binding

The fact that the extent of binding (presumably to chromatin) of HIF proteins depends mainly on the long C-terminal IDR rather than on their DBD was unexpected. Therefore, we next examined the contribution of the HIF DBD to the bound fraction. We introduced point mutations in the DBD (HIF-2α R27E and HIF-1α R30E) that were previously reported to impair DNA binding (Michel et al., 2002; Wu et al., 2015), and expressed them in the WT 786-O cells with the same L30 promoter system (Figure 5A). Not surprisingly, DBD mutants show a reduction in the bound fraction and a concomitant increase in the diffusing fraction compared to their WT counterpart (Figure 5B-C). In agreement with the expectation that the DBD mutations should not perturb protein-protein interactions, we do not observe a significant change in the overall diffusion coefficient of the moving population. These results demonstrate that, although the AD-containing IDR is the major modulator in determining the differences in bound fraction among different HIF dimers, the DBD is also important for binding, further suggesting that the observed bound fraction likely represents chromatin/DNA binding.

We next examined whether dimerization with HIF-1β is required for HIF-α chromatin binding. Taking advantage of the same L30 weak expression system, we exogenously expressed the Halo-HIF-2α dimerization mutant (R171A/V192D), or the analogous Halo-HIF-1α dimerization mutant (R170A/V191D) in the WT 786-O cells (Figure 5D). We found that compared to the WT Halo-HIF-2α or -1α, these mutants exhibit a significantly decreased bound fraction (Figure 5E-F), demonstrating that HIF-α without -1β can no longer effectively bind to DNA/chromatin. Taken together, our results indicate that the HIF-α disordered region alone is not sufficient to maintain binding, but instead, the IDR and both the DBD and dimerization domains are also needed.

Intrinsic properties of HIF-α IDR determine the overall rate of diffusive HIF.

Interestingly, with the Halo-HIF-α dimerization mutants, we observed no obvious change in their overall diffusion coefficient in the moving population (Figure 5E-F; Table 1), indicating that losing their HIF-1β partner does not affect the overall HIF-α diffusion rate. This result suggests that it is some intrinsic property of HIF-α molecules, rather than the molecular weight of dimers versus monomers, that determines its diffusion rate and behavior. Our results suggest that while the moving population of HIF-1β alone diffuses relatively fast, the moving population of both HIF-α and HIF- α/β dimers diffuses relatively slowly. We postulate this is potentially due to the HIF-α IDR engaging in protein-protein interactions with various cofactors both when associated with

HIF-1β or when alone (Figure 8A). Indeed, this is consistent with our previous observation that

the HIF-a NT/HIF-1β dimer diffuses at a relatively fast rate, similar to HIF-1β alone which

apparently does not share this HIF-α IDR mediated capacity (Figure 3—figure supplement 2).

HIF-α IDR contributes to binding preferences and regulatory specificity of target genes.

To see how differences in HIF molecular dynamics relate to actual genome-wide binding, we

performed Cut&Run on 786-O cells stably expressing different forms of Halo-tagged HIF-α driven

by the L30 promoter (same cells we performed fSPT on in Figure 4). All these exogenously

expressed proteins contain a V5 tag at the N-terminus of the Halo-tag, which we used as the

epitope for Cut&Run antibody recognition. The Halo-HIF-2α KIN clone A31 served both as a

positive control and as a reference for endogenous HIF-2α binding, as it also contains a V5 tag

N-terminal to the Halo-tag. WT cells without genetic modification controlled for V5 antibody non-

specific binding. Interestingly, the overall genome-wide binding profiles are very similar for all HIF-

α variants we examined (Figure 6—figure supplement 1), where most of the endogenous HIF-2α

binding sites are also bound by all exogenously expressed HIF-α (-1α, -1α/2α, -2α, and -2α/1α).

Even genes that are known to be regulated by one or the other isoform (e.g, the HIF-1α responsive

gene *PGK1* and the HIF-2α responsive gene *TGFA* (Keith et al., 2012)) are bound by all α forms

at their promoters/enhancers.

However, we did find significant differential binding preferences dictated by the IDR at a subset

of HIF target sites (Figure 6A-B; Figure 6—figure supplement 2). For example, the enhancer of

HIF-2α responsive genes such as *HSD3B7* and *PLXNA2* show increased HIF-2α binding when

cells are expressing extra HIF-2α, but no change in HIF-1α binding when HIF-1α is overexpressed.

Interestingly, such an increase in binding is also seen when overexpressing HIF-1α/2α, but not

the HIF-2α/1α chimera and, importantly, results in selective gene activation (Figure 6B; Figure

6—figure supplement 2B). The opposite is true for the HIF-1α responsive *PPFIA4* promoter and

PTPRN enhancer (Figure 6A; Figure 6—figure supplement 2A): a specific increase in HIF-1α and

HIF-2α/1α binding selectively induces both genes, which are instead insensitive to elevated HIF-

2α and HIF-1α/2α levels. These genome-wide analyses confirm that IDR-guided binding by HIF-

2α versus HIF-1α can indeed differentially activate selected gene loci.

Because IDR-specific binding is not readily detectable at a genome-wide scale due to the intrinsic

limitations of ensemble measurements, we next focused on differentially regulated sites only. We

first identified sites that respond differentially to either HIF-1α or HIF-2α, and then ask what is the

binding signal strength for the HIF-1α/2α and HIF-2α/1α chimeras - do they resemble more those

of HIF-1α or HIF-2α, depending on their DBD or IDR (Figure 6C)? We define sites that show

elevated binding only when HIF-1α is overexpressed as HIF-1α responsive (Table 2), and sites

that show elevated binding only when HIF-2α is overexpressed as HIF-2α responsive (Table 3).

For HIF-1α responsive regions, we observed elevated binding when overexpressing either HIF-

1α/2α or HIF-2α/1α, compared to HIF-2α, indicating that HIF-1α DBD and IDR both contribute

binding to HIF-1α responsive regions. For HIF-2α responsive regions, binding tended to be more

elevated for HIF-1α/2α compared to HIF-2α/1α, suggesting that HIF-2α IDR might dominate

binding to HIF-2α responsive sites (Figure 6C; Figure 6—figure supplement 3).

Unlike Cut&Run data, we observe clear genome-wide IDR-dependent gene regulation for RNA-

seq performed on the same sets of cells. Specifically, cells overexpressing the HIF variants

containing the same IDR show similar overall gene expression profiles (Figure 7). For example,

genes activated by HIF-1α but not HIF-2α can also be activated by HIF-2α/1α, but not HIF-1α/2α.

Overall, our genomic results show that HIF-α IDR contributes significantly to isoform-specific

target site binding and helps determine isoform-specific target gene activation.

Discussion

Transcription factors must search, recognize and bind to their specific target sites among millions

of possible DNA sequences along chromatin to activate the correct gene. With the successful

development of X-ray crystallography and cryo-EM, mechanisms of DNA-binding specificity have

been extensively studied, primarily based on classically structured globular DNA-binding domains

of TFs. We now know that a variety of structural mechanisms are used to recognize DNA,

including formation of specific hydrogen bonds and DNA contour interactions (Rohs et al., 2010).

However, these inherent binding modalities of DBDs alone cannot explain TF binding site

selection *in vivo* in eukaryotic cells. As revealed by genome-wide *in vivo* binding assays, only a

subset of potential target sites become occupied, and this is not entirely consistent with either

DNA binding site affinity or chromatin accessibility (Behera et al., 2018; Grossman et al., 2017;

Srivastava and Mahony, 2020). On the other hand, TFs have long been recognized to also contain

long unstructured transactivation domains with simple amino acid composition (Gln-rich, acidic,

Pro-rich etc.), which often posed challenges to purification and/or crystallization of full-length TFs

(Courey and Tjian, 1988; Ma and Ptashne, 1987; Mermod et al., 1989; Tjian and Maniatis, 1994).

Recently, such intrinsically disordered regions (IDRs) were reported to play an important role in

weak and multivalent protein-protein interactions to form local small transient hubs that, when

exacerbated by overexpression, can drive phase separation. Although not structurally defined,
these interactions can still be sequence/amino acid composition selective (Chong et al., 2018;
Chong and Mir, 2021). IDRs are now proposed to have important functions in boosting gene
expression through hub or condensate formation to locally enrich for factors that are needed for
transcription (Boijja et al., 2018; Cho et al., 2018; Chong et al., 2018; Sabari et al., 2018; Wei et
al., 2020). However, few studies of IDRs have investigated their potential role in DNA binding site
search and selection. Some studies reported that for a subset of zinc finger proteins (Sp2 and
KLF3), an IDR is critical for *in vivo* binding and specificity (Burdach et al., 2014; Lim et al., 2016;
Völkel et al., 2015), and two recent studies using genomic approaches reported the IDR as a
determinant for specificity for some of the yeast transcription factors (Brodsky et al., 2020; Gera
et al., 2022).
419
Here, using advanced live cell single particle tracking, we report that TF IDRs previously
associated with ADs are, in fact, a major determinant mediating nuclear search dynamics and
chromatin binding characteristics. Employing both genetic and small molecule perturbations
together with a series of domain-swap and mutation experiments, we found that it is the AD-
associated disordered region of HIF-α rather than the intrinsic molecular weight of the TF that
dictates a relatively slow diffusion for both HIF-α monomers and HIF-α/β dimers. On the other
hand, when not engaged with HIF-α, HIF-1β diffuses rapidly as expected for an unencumbered
subunit. These results indicate that the diffusion characteristic of HIF molecules is profoundly
influenced by the properties of their disordered regions (Figure 8A). In fact, computational analysis
shows very different amino acid composition bias among HIF-1α, -2α and -1β disordered regions
(Figure 1—figure supplement 1B). Thus, it is very likely that as these molecules navigate through
the crowded nuclear environment, their distinct stretches of IDRs that also contain ADs make
differential and selective interactions with other nuclear components, resulting in distinct diffusive
behaviors. According to Coulomb's law, acidity, and thus electrostatic charge on the molecules,
contribute to the attraction or repulsion forces towards other molecules in the environment. Thus
it is also possible that due to differences in acidity, the different charges on these IDRs can cause
differential non-specific interactions with macromolecules including not only proteins, but also
DNA and RNA (Xiang et al., 2020).
438
While it is easy to conceptualize how IDRs can influence the rate of diffusion, one unexpected
result is that they also largely determine how much TFs bind to chromatin, as well as influence
their binding site preferences. Although we confirmed that the DBD and dimerization domains are

important for chromatin binding, the surprise was that our domain swap experiments clearly
demonstrated that the percentage of bound TF is mainly contributed by regions outside of the
DBD/dimerization domains. One explanation could be the differential charge propensities of the
different disordered regions (Figure 1—figure supplement 1B). For example, the HIF-2α IDR is
more positively charged and may not only slow down nuclear exploration but also stabilize
chromatin binding, possibly through stronger non-specific interactions with negatively charged
chromatin-associated RNA and/or nucleosome-free DNA regions. Besides direct chromatin
interactions, HIF-2α IDR could also increase and stabilize binding via indirect interactions with
other chromatin-bound proteins. Moreover, since different IDRs can selectively interact with other
IDRs (Chong et al., 2018; Chong and Mir, 2021), we also postulate that selective interactions with
other TFs or co-regulators may play a role in determining HIF chromatin-binding specificity. One
hint of such a "combinatorial TF selectivity mechanism" is the enrichment of different transcription
factor motifs at HIF-1α and HIF-2α binding sites. Depending on the cell type, AP-1 or FOXD2,
FOXL1 and FOXC2 were previously found at HIF-2α binding sites, while HEY1/2, ZNF263 or
SP1/2 were found at HIF-1α binding sites (Smythies et al., 2019). It was also previously reported
that while no target specificity was preserved in reporter gene assays, the N-terminal TAD of HIF-
α conferred endogenous target specificity for two of the HIF-1 unique genes examined, possibly
via specific interactions with transcriptional cofactors (Hu et al., 2007). Further Co-IP or pull-down
assay coupled with mass spectrometry (MS) will be needed to more fully dissect this type of *in*
vivo selectivity mechanism.

Given our results, we believe that the simple "division of labor" model for DBDs and IDRs is an
over-simplification which was probably only suitable for prokaryotes. Instead, eukaryotic TFs have
evolved to exploit both DBDs and IDRs for chromatin binding as well as binding site selection,
which is best suited for the eukaryotic chromatin environment. We propose that TF-chromatin
engagement (binding and site selection) comprises two components – the first being DBD-
mediated, mainly the classical motif binding force that acts directly on DNA, and the second being
IDR-mediated interactions (likely weak and multivalent) with the native chromatin environment
that include protein-protein interactions, and interactions with DNA and RNA molecules. Thus, we
want to emphasize that the term "DNA binding specificity" refers to the first interaction component
and we propose it should only be used with respect to isolated or naked DNA in eukaryotes.
Instead, "chromatin binding preferences" would be a better term to describe *in vivo* binding
preferences of eukaryotic TFs within a native chromatin environment, which not only include DBD-
DNA interactions, but also involve "micro-environment recognition" mediated by IDR-chromatin

interactions. These two forces can be additive or competitive, depending on the macromolecule

composition of each specific chromatin locus, resulting in varied TF occupancy that could not be

simply predicted from the DNA code. It seems that in the case of HIFs, for most of their binding

sites the combined DBD- and IDR-mediated forces are similar for both HIF-1α and -2α, resulting

in overlapping occupancy for different isoforms at these sites. However, for a few but striking

cases, IDR-mediated interaction dominates HIFs' binding preferences, modulating the amount of

HIF isoforms at these differentially regulated sites (Figure 8B). Interestingly, Myc, a closely related

bHLH-LZ family transcription factor, shows predominately DBD-mediated chromatin engagement,

where non-specific DBD-DNA interactions contribute most of the binding (Pellanda et al., 2021).

These different observations suggest that different eukaryotic transcription factors may exploit

DBD- and IDR-mediated interactions to different degrees, resulting in distinct chromatin

engagement mechanisms.

We note that IDR-mediated chromatin engagement could happen either before and after DNA

binding. It is possible that these IDR-mediated interactions first locate the TF to a few highly

concentrated protein hubs in the vicinity of selected genomic binding sites, and then the TF

performs a more localized target search with the DBD to find its motif, as it has been recently

proposed (Jana et al., 2021; Darzacq and Tjian, 2022; Staller, 2022). In this case, whether the TF

explores the nucleus alone to join the hubs or explores it as a larger complex containing other

TFs/coactivators, IDRs both determine the TF dynamics and significantly improves the TF on-

rate. On the other hand, it is also possible that TFs first binds to their target site via the DBD, and

then IDR-mediated interactions stabilize the binding, increasing the TF's residence time and

decreasing its off-rate. In both cases, distinct IDRs among different TF from closely related family

members can create isoform-specific binding (and thus achieve isoform-specific gene activation)

at selected genomic sites amongst all binding sites that are shared by the same family.

Finally, we have demonstrated that our fSPT platform provides a powerful tool able to resolve *in*

vivo protein dynamics that is exquisitely sensitive to concentration, subunit stoichiometry and

genetic/small molecule perturbations. This is especially important when studying TFs, where a

slight difference in expression level often generates completely different results, rendering over-

expression systems highly susceptible to artifacts. It is also worth underscoring the importance of

studying TFs in their native physiologically relevant chromatin environment, given their obligate

interactions with higher-order chromatin structures and cofactors. For example, the EPO gene is

reported to be responsive to HIF-2α but not HIF-1α in Hep3B cells (Warnecke et al., 2004) and in

murine liver (Rankin et al., 2007), however, a luciferase reporter driven by the upstream EPO

enhancer also responds strongly to HIF-1α (Varma and Cohen, 1997), which may generate

misleading results and interpretations. Our fSPT platform allows us to study transcriptional

regulation in the native chromatin context and with endogenous TF levels to obtain data with

physiological and functional relevance. Such live cell real time measurements under native cell

contexts could prove to be highly valuable, both for dissecting *in vivo* mechanisms of transcription

regulation, and for guiding the development of effective therapeutics. Our Belzutifan treatment

experiment is an example of how fSPT can reveal the mechanism of action of small molecule

inhibitors, and how it could serve as a powerful tool to screen for drugs that selectively target one

isoform versus another, using dimerization and binding readouts as indicators of efficacy and

specificity. Moreover, since our results demonstrated how IDRs can affect TF diffusion behavior,

potentially distinct dynamic features determined by a particular IDR can be exploited as a readout

for screening small molecules or peptides that target allosteric sites of TFs. Assays that can

quantitatively measure TF diffusive behavior in live cells could be transformative for advancing

drug discovery because a high throughput imaging strategy opens the door to effectively target

what has been traditionally considered "undruggable", such as most protein-protein interactions

including potentially unstructured TF activation domains.

In summary, using the HIF protein family as a case study, we uncovered a mechanism of IDR-

mediated nuclear search and differential chromatin binding leading to selective gene activation.

We expect this fundamental principle to be applicable to a broad range of TF families.

Materials and Methods

Cell culture, stable cell line construction and drug treatment

Human 786-O clear cell renal carcinoma cells were obtained from the UCB Cell Culture Facility

(RRID:SCR_017924), which were originally obtained from ATCC (#CRL-1932). The cells were

tested for mycoplasma using nuclear stain with a 100X lens on a fluorescence scope on 1/22/2020

when banked and STR was used to identify them on 1/24/2020. Cells were cultured at 37°C with

5% CO_2 in 4.5 g/L glucose DMEM (ThermoFisher, Waltham, MA, #10566016) supplemented with

10% Fetal Bovine Serum (HyClone, Logan, UT, Cat. #SH30396.03, lot #AE28209315), 1 mM

Sodium Pyruvate (ThermoFisher #11360070) and 100 U/mL Penicillin-Streptomycin

(ThermoFisher #15140122). Cells were subcultured at a ratio of 1:4 to 1:12 every 2 to 4 days for

no longer than 30 days. A regular mycoplasma testing was performed every two weeks using

PCR. Phenol red-free DMEM (ThermoFisher, #21063029) supplemented with 10% Fetal Bovine

Serum, 1 mM Sodium Pyruvate and 100 U/mL Penicillin-Streptomycin was used for imaging.

Stable cell lines expressing the exogenous gene product (Table 4) were generated by PiggyBac

transposition and antibiotic selection. The gene of interest was cloned into a PiggyBac vector

which also co-expresses a puromycin resistant gene using Gibson Assembly and confirmed by

Sanger sequencing.   Cells were transfected by nucleofection using the Lonza Cell Line

Nucleofector® Kit V (Lonza, Basel, Switzerland, #VVCA-1003) and the Amaxa Nucleofector II

device. For each transfection, cells were plated 1-2 days before nucleofection in a 15-cm dish,

and reached approximately 50-70% confluency on the day of nucleofection, which equals to

approximately 3-4 million cells.   2 µg of PiggyBac plasmid was co-transfected with 1 µg of

SuperPiggyBac transposase vector with the T-020 program according to manufacturer's protocol.

Transfected cells were cultured for 24-48 hours before changing to selection media. Cells were

then selected for 14 days with 1 µg/ml puromycin (ThermoFisher #A1113803) and stable cell lines

were maintained in selection media for up to 30 days of culturing.

For drug treatment, 100 mM Belzutifan stock solution was prepared by dissolving Belzutifan

powder (CAS No: 1672668-24-4, MedChemExpress, Monmouth Junction, NJ, Cat. #HY-125840)

in DMSO (Sigma, St. Louis, MO, #D2650), and was diluted 1:500,000 in growth media to the final

concentration of 0.2 µM. The same volume of DMSO (0.0002%) is used in the reference group

as control. Cells were treated for 24 hours in either Belzutifan or DMSO alone before imaging.

For dosage-dependent assays in Figure 3—figure supplement 1, DMSO amount was kept the

same (0.0002%) for all drug concentrations.

CRISPR/Cas9-mediated genome editing

Knock-in cell lines were generated as previously described (Hansen et al., 2017) with the following

changes.   For each editing case, we designed 3 sgRNAs using CRISPOR (Concordet and

Haeussler, 2018). For each guide/donor pair, approximately 4 million 786-O cells were

nucleofected with 3.75 µg of donor plasmid and 1.25 µg of sgRNA plasmid. 24 hours after

transfection, Venus-positive cells were sorted and cultured for another 5-7 days, then Halo-

positive cells were sorted individually into single wells of 96 well plates. Clones were expanded

and genotyped with two rounds of PCR. The first round used one primer upstream of the left

homologous arm and the other primer downstream of the right homologous arm. The second

round used either of the external primers and a corresponding internal primer located in the

HaloTag coding region. Homozygous clones with the correct genotype, including Halo-HIF-2α

KIN clone A31 and clone B50, Halo-HIF-1β KIN clone A21 and clone B89, were confirmed by

Sanger sequencing and western blotting.

Cell preparation and dye labeling for imaging

For fast SPT, cells were grown on sonicated and plasma-cleaned 25-mm circular no 1.5H

precision cover glass (Marienfeld, Germany, 0117650) in 6-well plate. At least one day before

imaging, selective medium (if used) was removed and replaced with non-selective growth medium.

On the day of imaging, cells should be less than 100% confluent. Immediately before imaging,

cells were double labeled with JFX dyes as follows: cells were first incubated for 5 min in 1 ml

growth medium containing JFX 646, at a concentration that only gives approximately 10 detected

molecules per frame in the initial frames to ensure minimum misconnection of trajectories

between detections. This concentration differs from cell line to cell line, ranging from 0.2 – 5 nM,

depending on the expression level of the Halo-fusion protein. After 5 min of incubation, medium

was removed, cells were rinsed in PBS, and incubated for 5 min in 1 ml medium containing JFX

549. The concentration of JFX 549 also varies, usually at 25x the concentration of JFX 646. After

incubation, cells were washed twice for 5 min each, a first time with 2 ml regular growth media,

and a second time with 2 ml phenol red-free growth media, with a quick PBS rinse before each

wash. After wash, coverslip was transferred to Attofluor Cell Chambers (ThermoFisher, #A7816)

with cells facing up and 1 ml phenol red-free medium added to the chamber. For Belzutifan

treatment experiments, Belzutifan or equivalent amount of DMSO was added throughout the

labeling and washing steps (except during PBS rinses), as well as in the final imaging medium,

at the indicated concentration.

Live Cell Single particle tracking

All SPT experiments were carried out on a custom-built microscope as previously described

(Hansen et al., 2017) (McSwiggen et al., 2019). In brief, a Nikon TI microscope is equipped with

a 100x/ NA 1.49 oil-immersion TIRF objective, a motorized mirror, a perfect Focus system, an

EM-CCD camera and an incubation chamber maintained with humidified atmosphere with 5%

CO_2 at 37 °C. All microscope, camera and hardware components were controlled through the

NIS-Elements software (Nikon).

During imaging, samples were excited with 561-nm laser at 1100 mW (Genesis Coherent, Santa

Clara, CA) with emission filter set to Semrock 593/40 nm band-pass filter to locate and focus the

cell nuclei, as well as to adjust laser angle to achieve highly inclined laminated optical sheet (HiLo)

illumination (Tokunaga et al., 2008). An ROI (Regions of Interest) of random size was selected to

fit into the interior of the nuclei but with maximized area. Then the emission filter was switched to

Semrock 676/37 nm bandpass filter while keeping TIRF angle, stage xyz position and ROI the

same. Movies were then taken with 633-nm laser (Genesis Coherent, Santa Clara, CA) at 1100

mW and 1 ms pulse, with camera exposure at 5.48-ms frame rate for 800-1600 frames, until

samples were completely photo-bleached. At least 20 movies (corresponding to 20 cells) were

taken for each sample as one biological replicate on a given day. A total of three biological

replicates on three separate days were collected to produce the final results (>60 cells per cell

line/condition).

SPT data processing

Raw SPT movies were processed with a publicly available single particle tracking package

(https://github.com/alecheckert/quot) to generate trajectory files (.trajs). Generally, it performs

tracking in the following steps: read a frame, find spots in the frame, localize spots to subpixel

resolution, and reconnect spots from consecutive frames into trajectories. Since a non-

photoactivatable dye was used for all SPT experiments, we labeled cells with a dye concentration

that only gives very low spot detection density, which allowed us to track spots since the first

frame. This is important because if the initial frames are filtered due to high localization density,

there might be a bias towards moving molecules, due to the bound molecules being

photobleached and diffusing molecules moving into the focal plane during the later frames.

Although we used very sparse labeling, occasionally there would be frames with high density, to

minimize misconnections due to multiple particles in close proximity, we incorporated a filtering

step where we removed frames with more than 7 detections in the following way. First, we

computed the number of detections per frame. Next, this function was smoothed with uniform

filtering with a kernel width of 21 frames. Finally, we identified frames with fewer than 7 detections

after smoothing and isolated trajectories from these frames. Specifically, the following

configuration was used for all detections and tracking: Image reading and filtering settings: start

= 0, method = "identity", chunk_size = 100; Spot detection settings: method = "llr", k = 1.0, w = 15,

t = 18; Subpixel localization settings: method = 'ls_int_gaussian', window_size = 9, sigma = 1.0,

ridge = 0.001, max_iter = 20, damp = 0.3; Tracking settings: method = 'euclidean',

max_spots_per_frame = 7, pixel_size_um = 0.16, frame_interval = 0.00548, search_radius = 1.0,

max_blinks = 0, min_I0 = 0.0, scale = 7.0.

To infer the distribution of diffusion coefficients from experimentally observed trajectories, we

used a publicly available implementation of state arrays (https://github.com/alecheckert/spagl)

(`sample_script_fss.py`), which generates the posterior mean occupations for a state array

evaluated on trajectories across all cells. In all analyses, we used the likelihood function for

regular Brownian with localization error (RBME) (Heckert et al., 2021). Settings were:

frame_interval = 0.00548, pixel_size_um = 0.16, dz = 0.7. Occupations are reported as the

mean of the posterior distribution over state occupations, marginalized on diffusion coefficient.

To generate RBME likelihood for individual cells, we used the `sample_script_by_file.py`

script in the same repository (https://github.com/alecheckert/spagl) (Heckert et al., 2021) with

the following settings: frame_interval = 0.00548, dz = 0.7, pixel_size_um=0.16,

scale_by_total_track_count = True, scale_colors_by_group = True.

Antibodies

The following antibodies were used for ChIP-seq: rabbit polyclonal anti-HIF-2α (Novus Biologicals,

Centennial, CO, #NB100-122), mouse monoclonal anti-HIF-1β (Novus Biologicals, #NB100-124),

rabbit polyclonal anti-V5 (Abcam, Cambridge, UK, #ab9116). The following antibodies were used

for Cut&Run: mouse monoclonal anti-V5 tag (ThermoFisher, # R960-25) diluted to 0.01 mg/ml,

mouse IgG (Jackson ImmunoResearch #015-000-003) diluted to 0.01 mg/ml, rabbit anti-mouse

IgG H&L (Abcam, #ab46540) diluted to 0.01 mg/ml. The following antibodies were used for

western blotting: rabbit monoclonal anti-HIF-2α (Cell Signaling, Danvers, MA, #D9E3) diluted at

1:1000, rabbit monoclonal anti-HIF-1β (Cell Signaling, #D28F3) diluted at 1:1000, mouse

monoclonal anti-V5 tag (ThermoFisher, # R960-25) diluted at 1:2500, mouse monoclonal anti-

HaloTag (Promega, Madison, WI, # G9211) diluted at 1:1000, mouse monoclonal anti-TBP

(Abcam, #ab51841) diluted at 1:2500, goat-anti-mouse-HRP (ThermoFisher, #31430) diluted at

1:2000, goat-anti-rabbit-HRP (ThermoFisher, # 31462) diluted at 1:2000.

Western blotting

All western samples were prepared as follows: cells growing in either 6-well plates or 10-cm dish

in log phase were rinsed with PBS twice and lysed on ice in 100-500 ul 2x Sample buffer (80 mM

Tris pH6.8, 2% SDS, 10% Glycerol, 0.0006% Bromophenol blue) containing 280 mM 2-

Mercaptoethanol (Sigma #M7522), 1x Aprotinin (Sigma, #A6279, diluted 1:1000), 1 mM

Benzamidine (Sigma, #B6506), 1x cOmplete™ EDTA-free Protease Inhibitor Cocktail (Sigma,

#5056489001), and 0.25 mM PMSF (Sigma #11359061001). Cell lysates were scraped and

collected into 1.5-ml Eppendorf tubes, incubated at 99 °C with constant shaking, snap frozen in

liquid nitrogen and stored at -80 °C. On the day of western blotting, samples were thawed and

centrifuged at top speed for 5 min at 4 °C. Ten to 15 µl supernatant were loaded on an 8% SDS-

Page gel, ran for 1h at 200 V and 4 °C, and transferred to 0.45-µm nitrocellulose membrane

(Fisher, # 45004031) for 2 hrs at 100V. Membranes were blocked in 10% milk in 0.1% TBS-Tween

for 1 hr at RT, and incubated overnight at 4 °C with primary antibodies diluted in 5% milk in 0.1%

TBS-Tween. After 4 x 5 min washes in 0.1% TBS-Tween, membranes were incubated at RT for

at least 1 h with secondary antibodies diluted in 5% milk in 0.1% TBS-Tween. After 4 x 5 min

washes in 0.1% TBS-Tween, membranes were incubated for 3 min in freshly made Perkin Elmer

LLC Western Lightning Plus-ECL, Enhanced Chemiluminescence Substrate (Fisher,

#509049326), and imaged with a Bio-Rad ChemiDoc imaging system (BioRad, Model No:

Universal Hood III). For reblotting, membranes were immersed in Restore™ Western Blot

Stripping Buffer (Fisher, #21059) for 15 min at RT with shaking, washed 3 x 10 min in 0.1% TBS-

Tween, followed by blocking, antibody incubation and chemiluminescence reaction as described

above.

Luciferase reporter assay

The firefly luciferase reporter gene construct was made by inserting a 3x Hypoxia Responsive

Elements (HREs) from the EPO gene enhancer (sequence:

tcgaagccctacgtgctgtctcacacagcctgtctgacctctcgacctaccggccgttcgaagccctacgtgctgtctcacacagccttct

gatctcgacctaccggccgttcgaagccctacgtgctgtctcacacagcctgtctgacctctcgacctaccggccgt) into the 5' of

the minimal TATA-box promoter in the pGL4.23 [luc2/minP] vector (Promega #E841A). A control

pHRL-TK vector (Promega #E2241) expressing Renilla luciferase with an HSV TK promoter was

used as reference to normalize luciferase activity. Cells were co-transfected with 1 ug of firefly

Luciferase vector and 0.1 ug Renilla luciferase vector by nucleofection with Lonza Cell Line

Nucleofector® Kit V (Lonza, #VVCA-1003) and the T-020 program in the Amaxa Nucleofector II

device. After nucleofection, cells were resuspended in complete growth medium, and plated into

12-well plates with Belzutifan added to various concentrations as indicated. 24 hours after

nucleofection, cells were lysed and luciferase activity was analyzed with Dual-luciferase Reporter

Assay System (Promega, #E1960) according to manufacturer's protocol. The relative luciferase

activity was calculated by normalizing firefly luciferase activity to the Renilla luciferase activity to

control for transfection efficiency.

Chromatin Immunoprecipitation and ChIP-seq library preparation

ChIP was performed as described with few modifications (Testa et al., 2005). Wild type 786-O or

endogenously tagged knock-in clones A31 (V5-Halo-HIF-2α) and A21 (V5-Halo-HIF-1β) were

expanded to two 15-cm dishes and cross-linked 5' at room temperature with 1% formaldehyde-

containing FBS-free medium; cross-linking was stopped by adding PBS-glycine (0.125 M final).

Cells were washed twice with ice-cold PBS, scraped, centrifuged for 10' and pellets were flash-

frozen. Cell pellets were thawed and resuspended in 2 ml of cell lysis buffer (5 mM PIPES, pH

8.0, 85 mM KCl, and 0.5% NP-40, 1 ml/15 cm plate) w/ protease inhibitors and incubated for 10'

on ice. Lysates were centrifuged for 10' at 4000 rpm and nuclear pellets resuspended in 6 volumes

of sonication buffer (50 mM Tris-HCl, pH 8.1, 10 mM EDTA, 0.1% SDS) w/ protease inhibitors,

incubated on ice for 10', and sonicated to obtain DNA fragments around 500 bp in length (Covaris

S220 sonicator, 20% Duty factor, 200 cycles/burst, 150 peak incident power, 10 cycles 30" on

and 30" off). Sonicated lysates were cleared by centrifugation and chromatin (400 μg per antibody)

was diluted in RIPA buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA, 0.5 mM EGTA, 1% Triton X-

100, 0.1% SDS, 0.1% Na-deoxycholate, 140 mM NaCl) w/ protease inhibitors to a final

concentration of 0.8 μg/μl, precleared with Protein G sepharose (GE Healthcare) for 2 hours at

4°C and immunoprecipitated overnight with 4 μg of specific antibodies. About 4% of the

precleared chromatin was saved as input. Immunoprecipitated DNA was purified with the Qiagen

QIAquick PCR Purification Kit, eluted in 33 μl of 0.1X TE (1 mM Tris-HCl pH 8.0, 0.01 mM EDTA)

and analyzed by qPCR together with 2% of the input chromatin prior to ChIP-seq library

preparation (SYBR® Select Master Mix for CFX, ThermoFisher). ChIP-qPCR primer sequences

were as follows:

hWISP1_positive_forward: TGAGGTCAGTGTGGTTTGGT

hWISP1_positive_reverse: ACATGGTCACGTAGCTAGCA

hWISP1_negative_forward: AGTCCCCAGCACATAGAAGG

hWISP1_negative_reverse: GGTTCTGAAGGTGACCGACT

ChIP-seq libraries were prepared using the NEBNext® Ultra™ II DNA Library Prep Kit for Illumina®

(NEB E7645) according to manufacturer instructions with a few modifications. 20 ng of ChIP input

DNA (as measured by Nanodrop) and 25 μl of the immunoprecipitated DNA were used as a

starting material and the recommended reagents' volumes were cut in half. The NEBNext Adaptor

for Illumina was diluted 1:10 in Tris/NaCl, pH 8.0 (10 mM Tris-HCl pH 8.0, 10 mM NaCl) and the

ligation step extended to 30'. After ligation, a single purification step with 0.9X volumes of

Agencourt AMPure XP PCR purification beads (Beckman Coulter A63880) was performed, eluting

DNA in 22 μl of 10 mM Tris-HCl pH 8.0. 20 μl of the eluted DNA were used for the library

enrichment step, performed with the KAPA HotStart PCR kit (Roche Diagnostics KK2502) in 50

μl of total reaction volume (10 μl 5X KAPA buffer, 1.5 μl 10 mM dNTPs, 0.5 μl 10 μM NEB

Universal PCR primer, 0.5 μl 10 μM NEB index primer, 1 μl KAPA polymerase, 16.5 μl nuclease-

free water and 20 μl sample). Samples were enriched with 9 PCR cycles (98 °C, 45”; [98 °C, 15”;

60 °C, 10”] x 9; 72 °C, 1'; 4 °C, hold), purified with 0.9 volumes of AMPure XP PCR purification

beads and eluted with 33 μl of 10 mM Tris-HCl pH 8.0. Library concentration, quality and fragment

size were assessed by Qubit fluorometric quantification (Qubit™ dsDNA HS Assay Kit,

Invitrogen™ Q32851) qPCR and Fragment analyzer™. 12 multiplexed libraries (input, HIF1-β,

HIF1-α and V5 pulldowns in WT 786O cells and A31 and A21 clones) were pooled and sequenced

in one lane on the Illumina HiSeq4000 sequencing platform (50-bp, single end-reads) at the

Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley.

ChIP-seq analysis

ChIP-seq raw reads from WT 786O cells and A31 and A21 endogenously Halo-tagged clones (12

libraries total, 1 replicate per condition) were quality-checked with FastQC 0.10.1 and aligned

onto the human genome (hg38 assembly) using Bowtie (Langmead et al., 2009), allowing for two

mismatches (-n 2) and no multiple alignments (-m 1). Peaks were called with MACS2

2.1.0.20140616 (--nomodel --extsize 300) (Zhang et al., 2008) using input DNA as a control. To

create heatmaps we used deepTools 2.4.1 (Ramírez et al., 2016). We first ran bamCoverage (--

binSize 50 --normalizeTo1x 2913022398 --extendReads 300 --ignoreDuplicates -of bigwig) and

normalized read numbers to 1x sequencing depth, obtaining read coverage per 50-bp bins across

the whole genome (bigWig files). We then used the bigWig files to compute read numbers across

6 kb centered on HIF-2α peaks called by MACS2 across all 786O cell lines, subtracted of V5

peaks called by MACS2 in WT 786O cells (computeMatrix reference-point --referencePoint=TSS

--upstream 3000 --downstream 3000 --missingDataAsZero --sortRegions=no). We sorted the

output matrices by decreasing WT 786O enrichment, calculated as the total number of reads

within a MACS2 called ChIP-seq peak. Finally, heatmaps were created with the plotHeatmap tool

(--averageTypeSummaryPlot=mean --colorMap='Blues' --sortRegions=no).

Cut&Run

Cut&Run was performed as published (Janssens and Henikoff, 2019) with the following

modifications / specifications. Around 0.5 million cells were used for each experimental condition.

Before permeabilization, proteinase inhibitor was left out from the buffer to minimize toxicity to the

cells, but was added during the permeabilization step. Digitonin was used at a final concentration

of 0.02% for Dig wash buffer and antibody buffer, as it was tested to be the minimum concentration

to fully permeabilize 786-O cells. For each cell line, a primary antibody (either mouse-anti-V5 or

mouse IgG, 1mg/ml at 1:100 dilution) was used with a 4 °C overnight incubation. A secondary

rabbit-anti-mouse IgG was used at 1:100 dilution with an hour incubation at 4 °C. For each wash,

100 µl of Dig-wash buffer was used. Chromatin digestion was done for 30 minutes on ice. DNA

was extracted with phenol / chloroform and the pellet was dissolved in 30 µl 0.1 x TE (1 mM Tris-

HCl pH 8 0.1 mM EDTA). DNA was quantified by Qubit (Qubit™ dsDNA HS Assay Kit,

InvitrogenTM Q32851) and up to 40 ng (and up to 25 µl) was used for library preparation, using

NEBNext$^®$ Ultra$^™$ II DNA Library Prep Kit for Illumina$^®$ (NEB E7645) according to manufacturer

instructions, with a few modifications. Reagents' volumes were cut in half. For end prep 20 °C 30

min followed by 50 °C 60 min was used. For adapter ligation, NEBNext Adapter was diluted 1:20

in Tris/NaCl, pH 8.0 (10 mM Tris, 10 mM NaCl), mixture was incubated for 30 min at 20 °C (instead

of 15 min). After ligation, product was size-selected with 1.75x Agencourt AMPure XP PCR

purification beads (Beckman Coulter A63880), and DNA was eluted in 15 µl 10 mM Tris-HCl pH

8.0. 13 ul of eluted DNA was used for library enrichment with 15 µl NEBNext Ultra II Q5 Master

mix, 1 µl 10 µM NEB Universal PCR primer and 1 µl 10 µM NEB Index Primer. Samples were

enriched with 11 PCR cycles (98 °C, 30"; [98 °C, 10"; 65 °C, 10"] x 11; 65 °C, 5'; 4 °C, hold). The

30 µl PCR product was size-selected with double AMPure XP cleanup (first with 24 µl (0.8x) beads

and take the supernatant, then add 12 µl beads (to final PEG/NaCl 1.2x) and discard supernatant).

Library was eluted with 15 µl 10 mM Tris-HCl pH 8.0 and quantified by Qubit (Qubit™ dsDNA HS

Assay Kit, InvitrogenTM Q32851). Libraries were sent to MedGenome Inc. (Foster City, CA, USA)

for fragment analysis, multiplexing and sequencing on the Illumina NovaSeq 6000 platform (150

bp, paired end reads).

Cut&Run data analysis

Cut&Run raw fastq reads were first quality-checked with FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc) and then aligned to the human

genome (hg38) using bowtie2 version 2.3.4.1 with options: --local --no-unal --very-sensitive --no-

mixed --no-discordant --phred33 -I 10 -X 700. Non-human chromosomes were removed, then

bam files were created using Samtools (Li et al., 2009) version 1.8. Then Sambamba (Tarasov et

al., 2015) version 0.6.6 was used to sort bam files, and filter unmapped reads. Finally, samtools

were used to index the files. For peak calling with MACS2 (--keep-dup all --max-gap 400 -g hs --

bdg -q 0.01 -f BAMPE) (Zhang et al., 2008), mouse IgG control was used as control for each cell

line. Blacklisted regions were removed from the output narrowpeak file with BEDTools (Quinlan

and Hall, 2010) version 2.28.0. For visualization of genome-wide binding strength on the hg38
genome with the Integrative Genomics Viewer (IGV) (Robinson et al., 2011; Thorvaldsdóttir et al.,
2013), individual replicates were first combined into a single bam file using Samtools version 1.8,
and then converted to bigWig output files from deepTools bamCoverage (--binSize 20 --
normalizeUsing BPM --smoothLength 60). The PCA and clustering analysis, as well as
identification of differentially bound regions by HIF-1α and HIF-2α were done with DiffBind (Stark
and Brown, 2011) version 3.0, with all three replicates as samples. The final HIF-1α and HIF-2α
responsive regions were obtained by filtering the list with P value <0.01. Heatmaps for binding
strength were generated for each individual replicate or the combined replicates, using deepTools
(Ramírez et al., 2016) version 3.5.1.
825

RNA extraction and poly-A RNA-Seq library preparation
Poly-A RNA-Seq was performed in two batches and three biological replicates per sample. In the
first batch (3 replicates each of WT 786-O cells, HIF-1b KI, HIF-2a KI, HIF-1a O.E., HIF-1a R30E,
HIF-1a DM, HIF-2a O.E., HIF-2a R27E, HIF-2a DM) total RNA was extracted with the QIAGEN
RNeasy Plus Mini Kit (74134) according to manufacturer instructions, quantified by NanoDrop
and checked for integrity by capillary electrophoresis. 1 µg of the purified RNA was added of
ERCC RNA Spike-In Mix 2 (ThermoFisher Scientific #4456740, 0.5-1 µl of a 1:100 dilution per
sample) and the total volume brought to 50 µl with UltraPure Water (ThermoFisher Scientific #
10977-023). mRNA purification, RNA fragmentation, first and second strand cDNA synthesis were
performed according to the TruSeq RNA Sample Preparation v2 Kit (Illumina RS-122-2001) using
Superscript III for reverse transcription instead of Superscript II (incubation time: 50 °C for 50 min).
cDNA was purified with AMPure XP beads and eluted in 27.5 µl 10 mM Tris-HCl pH 8.0, 25 µl of
which were transferred to a new tube and subjected to a NEBNext® Ultra™ II DNA Library Prep
Kit for Illumina (NEB #E7645), using half of the recommended reagents' volumes. The NEBNext
Adaptor for Illumina was diluted 1:5 in Tris/NaCl, pH 8.0 (10 mM Tris-HCl pH 8.0, 10 mM NaCl)
and the ligation step extended to 30'. Library enrichment was performed with the KAPA HotStart
PCR kit (Roche Diagnostics KK2502) in 50 µl of total reaction volume (10 µl 5X KAPA buffer, 1.5
µl 10 mM dNTPs, 0.5 µl 10 µM NEB Universal PCR primer, 0.5 µl 10 µM NEB index primer, 1 µl
KAPA polymerase, 16.5 µl nuclease-free water and 20 µl sample). Samples were enriched with
8 PCR cycles (98 °C, 45"; [98 °C, 15"; 60 °C, 10"] x 9; 72 °C, 1'; 4 °C, hold), purified with 0.9
volumes of AMPure XP PCR purification beads and eluted with 33 µl of 10 mM Tris-HCl pH 8.0.
Library concentration, quality and fragment size were assessed by Qubit fluorometric
quantification (Qubit™ dsDNA HS Assay Kit, InvitrogenTM Q32851) qPCR and Fragment

analyzer™. Multiplexed libraries were pooled and sequenced on the Illumina NovaSeq 6000

platform (50-100 bp, paired-end reads) at the Vincent J. Coates Genomics Sequencing

Laboratory at UC Berkeley, supported by NIH S10 OD018174 Instrumentation Grant.

In the second batch (3 replicates each of WT 786-O cells, HIF-1a O.E., HIF-2a/1a, HIF-2a O.E.,

HIF-1a/2a) total RNA was extracted with TRIzol™ (Thermo Fisher Scientific #15596026)

according to manufacturer instructions, performing an additional wash with 1 volume of chloroform

after the recommended phenol:chloroform extraction. RNA was quantified by NanoDrop and

checked for integrity by capillary electrophoresis. 5 µg of total RNA were DNase treated (Ambio

#AM1906) and 1 µg of DNase-free RNA was subject to poly-A purification and library preparation

with the NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB #E7490S) in combination with

the NEBNext Ultra II RNA Library Prep Kit for Illumina (NEB #E7770S). The NEBNext Adaptor for

Illumina was diluted 1:5 in Tris/NaCl, pH 8.0 (10 mM Tris-HCl pH 8.0, 10 mM NaCl) and the ligation

step extended to 30'. Libraries were enriched with 8 PCR cycles. Library concentration was

assessed by Qubit quantification (Qubit™ dsDNA HS Assay Kit, InvitrogenTM Q32851).

Multiplexed libraries were pooled and sequenced on the Illumina NovaSeq 6000 platform (150 bp,

paired end reads) by MedGenome Inc. (Foster City, CA, USA).

RNA-Seq analysis

RNA-Seq raw reads were quality-checked with FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc) and aligned onto the human genome

(hg38) using STAR 2.6.1d RNA-Seq aligner (Dobin et al., 2012) with the following options: --

outSJfilterReads Unique --outFilterMultimapNmax 1 --outFilterIntronMotifs RemoveNoncanonical

--outSAMstrandField intronMotif. We used Samtools 1.9 (Li et al., 2009) to convert STAR

output .sam files into .bam files, and to sort and index them. We then counted how many reads

overlapped an annotated gene (GENECODE v32 annotations) using HTSeq 0.11.0 (Anders et al.,

2014) (htseq-count --stranded=no -f bam --additional-attr=gene_name -m union), and used the

output counts files to find differentially expressed genes with DESeq2 (Love et al., 2014), run with

default parameters within the Galaxy platform (Blankenberg et al., 2010; Giardine et al., 2005;

Goecks et al., 2010). Differentially expressed genes (DEG) were called using and adjusted P-

value ≤ 0.01, a fold change $\geq$ 2 and $\geq$ 0.5 mean counts. Gene transcript levels were visualized on

the hg38 genome with the Integrative Genomics Viewer (IGV) (Robinson et al., 2011;

Thorvaldsdóttir et al., 2013) using the bigWig output files from deepTools bamCoverage (--binSize

50 --extendReads 250 –normalizeUsingRPKM).

Datasets and accession numbers

Acknowledgments

We thank Luke Lavis for providing fluorescent HaloTag ligands; the CRL Flow Cytometry Facility for assistance with cell sorting; the QB3 High Throughput Screening Facility for providing access to the Opera Phenix automated confocal microscope; Dr. Alec Heckert for sharing data analysis pipeline and usage instructions; Dr. Hatice Kaya-Okur for valuable suggestions on Cut&Run protocols. Dr. Alec Heckert, Dr. Andrew Belmont, Dr. John Lis, Dr. Jiang Xu, Dr. Max Staller, Dr. Shasha Chong, and members of Tjian/Darzacq labs for helpful discussions and critical reading of the manuscript. This work was supported by the NIH grant U54-CA231641-01 (to XD), and the Howard Hughes Medical Institute (to RT). This work used the Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley, supported by the NIH S10 OD018174 Instrumentation Grant.

Author contributions: YC designed experiments with guidance from XD and RT. YC, CC, GD and QZ performed experiments. YC drafted the original manuscript. CC, XD and RT reviewed and edited the manuscript. XD and RT supervised the project.

Competing interests: RT and XD are co-founders of Eikon Therapeutics.

References

Anders S, Pyl PT, Huber W. 2014. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**:166–169. doi:10.1093/bioinformatics/btu638

Behera V, Evans P, Face CJ, Hamagami N, Sankaranarayanan L, Keller CA, Giardine B, Tan K, Hardison RC, Shi J, Blobel GA. 2018. Exploiting genetic variation to uncover rules of transcription factor binding and chromatin accessibility. *Nat Commun* **9**:782. doi:10.1038/s41467-018-03082-6

Boijja A, Klein IA, Sabari BR, Dall'Agnese A, Coffey EL, Zamudio AV, Li CH, Shrinivas K, Manteiga JC, Hannett NM, Abraham BJ, Afeyan LK, Guo YE, Rimel JK, Fant CB, Schuijers J, Lee TI, Taatjes DJ, Young RA. 2018. Transcription factors activate genes through the phase separation capacity of their activation domains. *Cell* **175**:1842. doi:10.1016/j.cell.2018.10.042

Brent R, Ptashne M. 1985. A eukaryotic transcriptional activator bearing the DNA specificity of a
prokaryotic repressor. *Cell* **43**:729–736. doi:10.1016/0092-8674(85)90246-6
Brodaczewska KK, Szczylik C, Fiedorowicz M, Porta C, Czarnecka AM. 2016. Choosing the right
cell line for renal cell cancer research. *Mol Cancer* **15**:83. doi:10.1186/s12943-016-0565-
8
Brodsky S, Jana T, Mittelman K, Chapal M, Kumar DK, Carmi M, Barkai N. 2020. Intrinsically
Disordered Regions Direct Transcription Factor In Vivo Binding Specificity. *Mol Cell*
79:459-471.e4. doi:10.1016/j.molcel.2020.05.032
Burdach J, Funnell APW, Mak KS, Artuz CM, Wienert B, Lim WF, Tan LY, Pearson RCM, Crossley
M. 2014. Regions outside the DNA-binding domain are critical for proper in vivo
specificity of an archetypal zinc finger transcription factor. *Nucleic Acids Res* **42**:276–
289. doi:10.1093/nar/gkt895
Cho W-K, Spille J-H, Hecht M, Lee C, Li C, Grube V, Cisse II. 2018. Mediator and RNA polymerase
II clusters associate in transcription-dependent condensates. *Science* **361**:412–415.
doi:10.1126/science.aar4199
Chong S, Dugast-Darzacq C, Liu Z, Dong P, Dailey GM, Cattoglio C, Heckert A, Banala S, Lavis L,
Darzacq X, Tjian R. 2018. Imaging dynamic and selective low-complexity domain
interactions that control gene transcription. *Science* **361**:eaar2555.
doi:10.1126/science.aar2555
Chong S, Mir M. 2021. Towards Decoding the Sequence-Based Grammar Governing the
Functions of Intrinsically Disordered Protein Regions. *J Mol Biol* **433**:166724.
doi:10.1016/j.jmb.2020.11.023
Concordet J-P, Haeussler M. 2018. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome
editing experiments and screens. *Nucleic Acids Res* **46**:W242–W245.
doi:10.1093/nar/gky354
Courey AJ, Tjian R. 1988. Analysis of Sp1 in vivo reveals multiple transcriptional domains,
including a novel glutamine-rich activation motif. *Cell* **55**:887–898. doi:10.1016/0092-
8674(88)90144-4
Darzacq X, Tjian R. 2022. Weak multivalent biomolecular interactions: a strength versus
numbers tug of war with implications for phase partitioning. *RNA* **28**:48–51.
doi:10.1261/rna.079004.121
Dobin A, Davis CA, Zaleski C, Schlesinger F, Drenkow J, Chaisson M, Batut P, Jha S, Gingeras TR.
2012. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:15–21.
doi:10.1093/bioinformatics/bts635
Edgar R, Domrachev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene expression and
hybridization array data repository. *Nucleic Acids Res* **30**:207–210.
doi:10.1093/nar/30.1.207
Gaur RK. 2014. Amino acid frequency distribution among eukaryotic proteins. *IIOAB J* **5**:6–11.
Gera T, Jonas F, More R, Barkai N. 2022. Evolution of binding preferences among whole-
genome duplicated transcription factors. *eLife* **11**:e73225. doi:10.7554/eLife.73225
Gnarra JR, Tory K, Weng Y, Schmidt L, Wei MH, Li H, Latif F, Liu S, Chen F, Duh FM. 1994.
Mutations of the VHL tumour suppressor gene in renal carcinoma. *Nat Genet* **7**:85–90.
doi:10.1038/ng0594-85

Grimm JB, Xie L, Casler JC, Patel R, Tkachuk AN, Falco N, Choi H, Lippincott-Schwartz J, Brown
TA, Glick BS, Liu Z, Lavis LD. 2021. A General Method to Improve Fluorophores Using
Deuterated Auxochromes. *Jacs Au* **1**:690–696. doi:10.1021/jacsau.1c00006
Grossman SR, Zhang X, Wang L, Engreitz J, Melnikov A, Rogov P, Tewhey R, Isakova A,
Deplancke B, Bernstein BE, Mikkelsen TS, Lander ES. 2017. Systematic dissection of
genomic features determining transcription factor binding and enhancer function. *Proc
Natl Acad Sci* **114**:E1291–E1300. doi:10.1073/pnas.1621150114
Hansen AS, Pustova I, Cattoglio C, Tjian R, Darzacq X. 2017. CTCF and cohesin regulate
chromatin loop stability with distinct dynamics. *eLife* **6**:e25776. doi:10.7554/eLife.25776
Hansen AS, Woringer M, Grimm JB, Lavis LD, Tjian R, Darzacq X. 2018. Robust model-based
analysis of single-particle tracking experiments with Spot-On. *eLife* **7**:e33125.
doi:10.7554/eLife.33125
Hanson J, Yang Y, Paliwal K, Zhou Y. 2017. Improving protein disorder prediction by deep
bidirectional long short-term memory recurrent neural networks. *Bioinformatics*
33:685–692. doi:10.1093/bioinformatics/btw678
Heckert A, Dahal L, Tjian R, Darzacq X. 2021. Recovering mixtures of fast diffusing states from
short single particle trajectories. bioRxiv. doi:10.1101/2021.05.03.442482
Hu C-J, Sataur A, Wang L, Chen H, Simon MC. 2007. The N-terminal transactivation domain
confers target gene specificity of hypoxia-inducible factors HIF-1alpha and HIF-2alpha.
Mol Biol Cell **18**:4528–4542. doi:10.1091/mbc.e06-05-0419
Jana T, Brodsky S, Barkai N. 2021. Speed–Specificity Trade-Offs in the Transcription Factors
Search for Their Genomic Binding Sites. *Trends Genet* **37**:421–432.
doi:10.1016/j.tig.2020.12.001
Janssens D, Henikoff S. 2019. CUT&RUN: Targeted in situ genome-wide profiling with high
efficiency for low cell numbers. *protocols.io*. https://www.protocols.io/view/cut-amp-
run-targeted-in-situ-genome-wide-profiling-zcpf2vn
Keith B, Johnson RS, Simon MC. 2012. HIF1α and HIF2α: sibling rivalry in hypoxic tumour growth
and progression. *Nat Rev Cancer* **12**:9–22. doi:10.1038/nrc3183
Kribelbauer JF, Rastogi C, Bussemaker HJ, Mann RS. 2019. Low-Affinity Binding Sites and the
Transcription Factor Specificity Paradox in Eukaryotes. *Annu Rev Cell Dev Biol* **35**:357–
379. doi:10.1146/annurev-cellbio-100617-062719
Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. *Genome Biol* **10**:R25. doi:10.1186/gb-
2009-10-3-r25
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
Subgroup 1000 Genome Project Data Processing. 2009. The Sequence Alignment/Map
format and SAMtools. *Bioinformatics* **25**:2078–2079.
Lim WF, Burdach J, Funnell APW, Pearson RCM, Quinlan KGR, Crossley M. 2016. Directing an
artificial zinc finger protein to new targets by fusion to a non-DNA-binding domain.
Nucleic Acids Res **44**:3118–3130. doi:10.1093/nar/gkv1380
Liu Z, Lavis LD, Betzig E. 2015. Imaging Live-Cell Dynamics and Structure at the Single-Molecule
Level. *Mol Cell* **58**:644–659. doi:10.1016/j.molcel.2015.02.033
Los GV, Encell LP, McDougall MG, Hartzell DD, Karassina N, Zimprich C, Wood MG, Learish R,
Ohana RF, Urh M, Simpson D, Mendez J, Zimmerman K, Otto P, Vidugiris G, Zhu J,

Darzins A, Klaubert DH, Bulleit RF, Wood KV. 2008. HaloTag: A Novel Protein Labeling
Technology for Cell Imaging and Protein Analysis. *ACS Chem Biol* **3**:373–382.
doi:10.1021/cb800025k

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**. doi:10.1186/s13059-014-0550-8

Ma J, Ptashne M. 1987. Deletion analysis of GAL4 defines two transcriptional activating
segments. *Cell* **48**:847–853. doi:10.1016/0092-8674(87)90081-X

McSwiggen DT, Hansen AS, Teves SS, Marie-Nelly H, Hao Y, Heckert AB, Umemoto KK, Dugast-
Darzacq C, Tjian R, Darzacq X. 2019. Evidence for DNA-mediated nuclear
compartmentalization distinct from phase separation. *eLife* **8**:e47098.
doi:10.7554/eLife.47098

Mermod N, O'Neill EA, Kelly TJ, Tjian R. 1989. The proline-rich transcriptional activator of
CTF/NF-I is distinct from the replication and DNA binding domain. *Cell* **58**:741–753.
doi:10.1016/0092-8674(89)90108-6

Michel G, Minet E, Mottet D, Remacle J, Michiels C. 2002. Site-directed mutagenesis studies of
the hypoxia-inducible factor-1α DNA-binding domain. *Biochim Biophys Acta BBA - Gene*
Struct Expr **1578**:73–83. doi:10.1016/S0167-4781(02)00484-0

Pellanda P, Dalsass M, Filipuzzi M, Loffreda A, Verrecchia A, Castillo Cano V, Thabussot H, Doni
M, Morelli MJ, Soucek L, Kress T, Mazza D, Mapelli M, Beaulieu M-E, Amati B, Sabò A.
2021. Integrated requirement of non-specific and sequence-specific DNA binding in
Myc-driven transcription. *EMBO J* **40**:e105464. doi:10.15252/embj.2020105464

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinforma Oxf Engl **26**:841–842. doi:10.1093/bioinformatics/btq033

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke T.
2016. deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res **44**:W160-165. doi:10.1093/nar/gkw257

Rankin EB, Biju MP, Liu Q, Unger TL, Rha J, Johnson RS, Simon MC, Keith B, Haase VH. 2007.
Hypoxia-inducible factor-2 (HIF-2) regulates hepatic erythropoietin in vivo. *J Clin Invest*
117:1068–1077. doi:10.1172/JCI30117

Raval RR, Lau KW, Tran MGB, Sowter HM, Mandriota SJ, Li J-L, Pugh CW, Maxwell PH, Harris AL,
Ratcliffe PJ. 2005. Contrasting properties of hypoxia-inducible factor 1 (HIF-1) and HIF-2
in von Hippel-Lindau-associated renal cell carcinoma. *Mol Cell Biol* **25**:5675–5686.
doi:10.1128/MCB.25.13.5675-5686.2005

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011.
Integrative genomics viewer. *Nat Biotechnol* **29**:24–26. doi:10.1038/nbt.1754

Rohs R, Jin X, West SM, Joshi R, Honig B, Mann RS. 2010. Origins of specificity in protein-DNA
recognition. *Annu Rev Biochem* **79**:233–269. doi:10.1146/annurev-biochem-060408-
091030

Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ, Hannett NM,
Zamudio AV, Manteiga JC, Li CH, Guo YE, Day DS, Schuijers J, Vasile E, Malik S, Hnisz D,
Lee TI, Cisse II, Roeder RG, Sharp PA, Chakraborty AK, Young RA. 2018. Coactivator
condensation at super-enhancers links phase separation and gene control. *Science*
361:eaar3958. doi:10.1126/science.aar3958

Schödel J, Grampp S, Maher ER, Moch H, Ratcliffe PJ, Russo P, Mole DR. 2016. Hypoxia, Hypoxia-inducible Transcription Factors, and Renal Cancer. *Eur Urol* **69**:646–657. doi:10.1016/j.eururo.2015.08.007

Schödel J, Oikonomopoulos S, Ragoussis J, Pugh CW, Ratcliffe PJ, Mole DR. 2011. High-resolution genome-wide mapping of HIF-binding sites by ChIP-seq. *Blood* **117**:e207–e217. doi:10.1182/blood-2010-10-314427

Semenza GL. 2012. Hypoxia-Inducible Factors in Physiology and Medicine. *Cell* **148**:399–408. doi:10.1016/j.cell.2012.01.021

Shen C, Beroukhim R, Schumacher SE, Zhou J, Chang M, Signoretti S, Kaelin WG. 2011. Genetic and Functional Studies Implicate HIF1α as a 14q Kidney Cancer Suppressor Gene. *Cancer Discov* **1**:222–235. doi:10.1158/2159-8290.CD-11-0098

Smythies JA, Sun M, Masson N, Salama R, Simpson PD, Murray E, Neumann V, Cockman ME, Choudhry H, Ratcliffe PJ, Mole DR. 2019. Inherent DNA-binding specificities of the HIF-1α and HIF-2α transcription factors in chromatin. *EMBO Rep* **20**:e46401. doi:10.15252/embr.201846401

Srivastava D, Mahony S. 2020. Sequence and chromatin determinants of transcription factor binding and the establishment of cell type-specific binding patterns. *Biochim Biophys Acta Gene Regul Mech* **1863**:194443. doi:10.1016/j.bbagrm.2019.194443

Staller MV. 2022. Transcription factors perform a 2-step search of the nucleus. *Genetics* iyac111. doi:10.1093/genetics/iyac111

Stark R, Brown G. 2011. DiffBind differential binding analysis of ChIP-Seq peak data. *R Package Version* **100**.

Swiatek M, Jancewicz I, Kluebsoongnoen J, Zub R, Maassen A, Kubala S, Udomkit A, Siedlecki JA, Sarnowski TJ, Sarnowska E. 2020. Various forms of HIF-1α protein characterize the clear cell renal cell carcinoma cell lines. *IUBMB Life* **72**:1220–1232. doi:10.1002/iub.2281

Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. 2015. Sambamba: fast processing of NGS alignment formats. *Bioinformatics* **31**:2032–2034. doi:10.1093/bioinformatics/btv098

Testa A, Donati G, Yan P, Romani F, Huang TH-M, Viganò MA, Mantovani R. 2005. Chromatin immunoprecipitation (ChIP) on chip experiments uncover a widespread distribution of NF-Y binding CCAAT sites outside of core promoters. *J Biol Chem* **280**:13606–13615. doi:10.1074/jbc.M414039200

Thorvaldsdóttir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV): High-performance genomics data visualization and exploration. *Brief Bioinform* **14**:178–192. doi:10.1093/bib/bbs017

Tjian R, Maniatis T. 1994. Transcriptional activation: a complex puzzle with few easy pieces. *Cell* **77**:5–8. doi:10.1016/0092-8674(94)90227-5

Tokunaga M, Imamoto N, Sakata-Sogawa K. 2008. Highly inclined thin illumination enables clear single-molecule imaging in cells. *Nat Methods* **5**:159–161. doi:10.1038/nmeth1171

Varma S, Cohen HJ. 1997. Co-transactivation of the 3' erythropoietin hypoxia inducible enhancer by the HIF-1 protein. *Blood Cells Mol Dis* **23**:169–176. doi:10.1006/bcmd.1997.0134

Völkel S, Stielow B, Finkernagel F, Stiewe T, Nist A, Suske G. 2015. Zinc Finger Independent Genome-Wide Binding of Sp2 Potentiates Recruitment of Histone-Fold Protein Nf-y

1092   Distinguishing It from Sp1 and Sp3. *PLoS Genet* **11**:e1005102.

1093   doi:10.1371/journal.pgen.1005102

1094 Wallace EM, Rizzi JP, Han G, Wehn PM, Cao Z, Du X, Cheng T, Czerwinski RM, Dixon DD, Goggin

1095   BS, Grina JA, Halfmann MM, Maddie MA, Olive SR, Schlachter ST, Tan H, Wang B, Wang

1096   K, Xie S, Xu R, Yang H, Josey JA. 2016. A Small-Molecule Antagonist of HIF2α Is

1097   Efficacious in Preclinical Models of Renal Cell Carcinoma. *Cancer Res* **76**:5491–5500.

1098   doi:10.1158/0008-5472.CAN-16-0473

1099 Warnecke C, Zaborowska Z, Kurreck J, Erdmann VA, Frei U, Wiesener M, Eckardt K-U. 2004.

1100   Differentiating the functional role of hypoxia-inducible factor (HIF)-1alpha and HIF-

1101   2alpha (EPAS-1) by the use of RNA interference: erythropoietin is a HIF-2alpha target

1102   gene in Hep3B and Kelly cells. *FASEB J Off Publ Fed Am Soc Exp Biol* **18**:1462–1464.

1103   doi:10.1096/fj.04-1640fje

1104 Wei M-T, Chang Y-C, Shimobayashi SF, Shin Y, Strom AR, Brangwynne CP. 2020. Nucleated

1105   transcriptional condensates amplify gene expression. *Nat Cell Biol* **22**:1187–1196.

1106   doi:10.1038/s41556-020-00578-6

1107 Wenger RH, Stiehl DP, Camenisch G. 2005. Integration of oxygen signaling at the consensus

1108   HRE. *Sci Signal* **2005**:re12. doi:10.1126/stke.3062005re12

1109 Wu D, Potluri N, Lu J, Kim Y, Rastinejad F. 2015. Structural integration in hypoxia-inducible

1110   factors. *Nature* **524**:303–308. doi:10.1038/nature14883

1111 Xiang L, Chen K, Yan R, Li W, Xu K. 2020. Single-molecule displacement mapping unveils

1112   nanoscale heterogeneities in intracellular diffusivity. *Nat Methods* **17**:524–530.

1113   doi:10.1038/s41592-020-0793-0

1114 Xu R, Wang K, Rizzi JP, Huang H, Grina JA, Schlachter ST, Wang B, Wehn PM, Yang H, Dixon DD,

1115   Czerwinski RM, Du X, Ged EL, Han G, Tan H, Wong T, Xie S, Josey JA, Wallace EM. 2019.

1116   3-[(1S,2S,3R)-2,3-Difluoro-1-hydroxy-7-methylsulfonylindan-4-yl]oxy-5-

1117   fluorobenzonitrile (PT2977), a Hypoxia-Inducible Factor 2α (HIF-2α) Inhibitor for the

1118   Treatment of Clear Cell Renal Cell Carcinoma. *J Med Chem* **62**:6876–6893.

1119   doi:10.1021/acs.jmedchem.9b00719

1120 Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown

1121   M, Li W, Liu XS. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**:R137.

1122   doi:10.1186/gb-2008-9-9-r137

1123

Figure 1. Endogenous tagging of HIFs in 786-O clear cell renal cell carcinoma (ccRCC) cells for fast single particle tracking (fSPT). (**A**) Schematic showing the similar domain organizations of HIFs (top) and the HRE-bound HIF α/β dimers (bottom). Disordered regions are represented as wavy lines. (**B**) Generation of Halo-KIN clones in the HIF-1α negative 786-O ccRCC line. Top: Halo-tagging scheme of HIF-2α (left) and HIF-1β (right). Bottom: Western blot of wild-type (WT) 786-O cells and homozygously tagged knock-in clones (A31 and A21). See Figure 1—figure supplement 2 for uncropped images. (**C**) Halo-tagged HIF-2α and HIF-1β show predominant nuclear localization. Top: schematic of labeling Halo-tagged proteins in live cells with cell-permeable Halo-binding JFX646 dye. Bottom: representative images of Halo-HIF-2α (left) and

Halo-HIF-1β (right) clones labeled with 500 nM JFX646 (**D**) representative images showing the

same cell labeled with a high concentration of JFX549 dye for localizing the nucleus in one

channel (left) and labeled sparsely with JFX646 dye for tracking individual molecules in another

channel (right). Scale bar = 2 µm. (**E**) Graphical illustration of fSPT capturing trajectories of moving

particles. Top: highly inclined and laminated optical sheet illumination (HiLo). Bottom: Illumination

and camera sequence with corresponding particle position at each frame (solid magenta dots).

Particle's past positions (dashed magenta circles) are connected with dotted magenta lines to

show the particle's trajectory. Scale bar = 2 µm. (**F**) Actual data showing detection of Halo-HIF-

2α protein molecules at 5.48 ms frame rate.

Figure 2. fSPT sensitively detects molecules in a range of states. (**A-B**) Likelihood of diffusion coefficients based on a model of regular Brownian motion with localization error (RBME) (Heckert et al., 2021), for (A) Halo-HIF-2α, clone A31 and (B) Halo-HIF-1β, clone A21, with drawing illustrating bound and different hypothetical moving states: complexes, dimer and monomer. Each row represents data collected from one cell. 0.1 μm^2/sec is used as the cut-off for bound versus free. (**C**) Top: proportion of molecules as a function of their diffusion coefficients (posterior mean occupations for a state array (Heckert et al., 2021)) evaluated on trajectories across all cells measured for each KIN line (Halo-HIF-2α, clone A31 and Halo-HIF-1β, clone A21). Compared to

HIF-2α, HIF-1β has less bound fraction (grey versus yellow shaded areas) and faster diffusion

coefficient (grey versus yellow arrows). Bottom: Summary of the bound fraction (left) and peak

diffusion coefficient (right) for the two clones. Each bar represents the averaged value from three

independent measurements on different days (black dots). (**D-E**) Over-expressing HIF-2α, but

not a dimerization mutant form, in the Halo-HIF-1β knock-in (KIN) line increases HIF-1β binding

and decreases its diffusion coefficient. (D) schematic illustrating the parental Halo-HIF-1β KIN

cells (grey background) and cells stably over-expressing (O.E.) either the wild type (darker blue

background) or a dimerization mutant (DM, black crosses, lighter blue background) form of HIF-

2α. (E) Top: Proportion of molecules as a function of diffusion coefficient measured for HIF-1β in

Halo-HIF-1β KIN cells (grey) and in Halo-HIF-1β KIN cells overexpressing HIF-2α (WT, dark blue

background, or dimerization mutant (HIF-2αDM), light blue). Shaded areas indicate bound

fraction. Bottom: Bar plot of the average value (bar height) of the bound fraction (left) and peak

diffusion coefficient (right) calculated from three independent measurements (black dots) for each

condition.

Figure 3. HIF-α increases HIF-1β binding and decreases HIF-1β diffusion coefficient through dimerization, in an IDR-dependent manner. (**A**) Schematic of evaluating Halo-HIF-1β behavior with genetic and small-molecule perturbation. Parental Halo-HIF-1β knock-in (KIN) cells (grey background) and cells stably overexpressing (O.E.) either a certain form of HIF-α (WT or domain swap, HIF-1α, red, HIF-2α, orange. Disordered regions are represented as wavy lines.) (various colored background) or a 3xFLAG tag only control (blue background) are used, with and without

0.2 µM Belzutifan (HIF-2α/1β dimerization inhibitor) treatment. (**B**) Proportion of Halo-HIF-1β

molecules as a function of diffusion coefficient measured in various conditions outlined in (A) Top

row: DMSO only, showing overexpressing α subunit can change HIF-1β behavior. Cells over-

expressing the α subunit variants containing HIF-2α disordered region (orange curly line) have a

stronger effect (middle, HIF-2α and HIF-1α/2α,) compared to those containing HIF-1α disordered

region (right, HIF-1α and HIF-2α/1α). Middle and Bottom rows: proportions of HIF-1β as a function

of diffusion coefficient, measured in each of the 6 cell lines with Belzutifan treatment are compared

to the DMSO control. Changes caused by overexpressing an α subunit can be specifically

reverted by Belzutifan treatment for cell lines expressing an α subunit variant that contains the

HIF-2α structured domain (orange globule). (**C**) Summary of the average bound fractions (top)

and peak diffusion coefficient (bottom) for all 12 conditions, with black dots indicating values from

each of the three individual measurements.

Figure 4. The IDR governs HIF-α molecular dynamics and binding characteristics. (**A**) Schematic representation of different HIF (WT and domain-swapped) being weakly and stably overexpressed with an L30 promoter and tracked in WT 786-O cells. (**B**) Proportion of molecules as a function of diffusion coefficient for every tracked protein in (A). Top: overlapping distribution curves shows almost identical behavior between Halo-HIF-2α (dark blue curve) and Halo-HIF-1α/2α (light blue curve). Middle: similar behavior between Halo-HIF-1α (purple curve) and Halo-HIF-2α/1α (yellow curve), bottom: overlay of all four curves shows very different behavior between

1189   proteins containing 1α versus 2α IDR. (**C**) Bar plot comparing the average bound fraction (left)

1190   and peak diffusion coefficient (right) for cells in (B), with black dots indicating values from three

1191   independent measurements.

Figure 5. HIF-α IDR alone is not sufficient for binding. (**A-C**) Mutation in DBD reduces the bound fraction for both HIF-α isoforms. (A) Schematic representation of weakly overexpressing and tracking wild type and DBD mutant (R27E or R30E, black cross) forms of HIF-1α or -2α, using the same L30 expression system as in Figure 4. (B) Proportion of molecules as a function of diffusion coefficient for tracked protein listed in (A). (C) Bar plot summarizes the average value (bar height) of the bound fraction (left) and peak diffusion coefficient (right) of three independent measurements (black dots). (**D-F**) Mutations in the dimerization domain reduce the bound fraction for both HIF-α isoforms but do not change their diffusion coefficient. (D) Schematic representation of weakly over-expressing and tracking wild type and dimerization-mutant (DM, two black crosses) forms of Halo-HIF-1α or -2α, using the L30 expression system. (E) Proportion of

1202   molecules as a function of diffusion coefficient for tracked protein in (D). (F) Summary of the

1203   average value (bar height) of the bound fraction (left) and peak diffusion coefficient (right) for all

1204   four proteins with black dots indicating values from three independent measurements.

Figure 6. HIF-α IDR contributes to isoform-specific chromatin binding preferences and activation of isoform-specific HIF target genes. (**A-B**) Genome browser view of anti-V5 Cut&Run and RNA-seq results on WT cells, Halo-HIF-2α KIN cells and cells expressing different Halo-tagged HIF-α variants driven by an L30 promoter, showing IDR-specific regulation. (A) An example of genes that are preferentially bound (dashed box) and specifically activated by HIF-1α, as well as by HIF-

2α/1α. (B) An example of genes that are preferentially bound and specifically activated by HIF-

2α, as well as by HIF-1α/2α. (**C**) Genome-wide analysis of contribution of DBD or IDR on isoform-

specific binding with Cut&Run data. Top: Schematic showing definition of HIF-1α and HIF-2α

responsive regions. Regions that show increased binding when overexpressing Halo-tagged HIF-

1α compared to when overexpressing Halo-tagged HIF-2α are defined as HIF-1α responsive

regions (purple highlight). Regions that show increased binding when overexpressing Halo-

tagged HIF-2α compared to when overexpressing HIF-1α are defined as HIF-2α responsive

regions (green highlight). Bottom: heatmap and pile-up results (blue curves for HIF-1α responsive

regions and green curves for HIF-2α responsive regions) on binding strength at all HIF-1α

responsive regions or HIF-2α responsive regions when cells overexpressing either Halo-tagged

HIF-2α, -2α/1α, -1α, or -1α/2α. Pile-up enrichment results at all HIF binding sites (dashed grey

curves) are used as control.

1222

Figure 7. RNA-seq analysis shows HIF-α isoform-specific gene regulation is IDR-dependent. (**A-**

B) PCA analysis (A) and Clustering (B) of RNA-seq results performed on WT 786-O cells, and

cells expressing Halo-tagged HIF-α variants driven by an L30 promoter. HIF-α variants sharing

the same IDR (-1α and -2α/1α or -2α and -1α/2α) have similar gene expression profiles. (**C**)

DESeq2 output summarizing pair-wise comparison of the number of differentially expressed

genes (DEG) in WT cells and cells over-expressing various HIF-α forms.

1229

Figure 8. A model for IDR-mediated nuclear search and chromatin binding. (**A**) The HIF-α IDR
determines its slow motion of both the HIF-α monomer and HIF-α/β dimer, likely by HIF-α IDR
mediated interactions with nuclear macromolecules. For HIF-α, the IDR thus determines its slow
motion regardless of its dimerization status. For HIF-1β, dimerization slows it down due to extra
interactions (yellow and red clouds) brought by HIF-α IDR. (**B**) HIF-chromatin engagement
comprises two components: DBD- and IDR-mediated interactions. As an obligated dimer, the
DBD and the dimerization domain are both necessary for HIF binding, but the IDR determines the
bound fraction, possibly via its interaction with nearby macromolecules, including other proteins
and/or nucleic acids (DNA and/or RNA).

Figure 1—figure supplement 1. Domain analysis of HIFs. (**A**) Disorder propensity of HIF-1α, -2α and -1β, predicted by SPOT-Disorder (Hanson et al., 2017). High values indicate higher disorder propensity. (**B**) HIF-1α and HIF-2α IDRs have different amino acid composition biases. The relative enrichment for each amino acid is calculated by comparison with the average amino acid composition of non-membrane proteins in vertebrates (Gaur, 2014), and color coded, with red indicating relative high enrichment and blue indicating relative depletion. The IDR segments between the N-TAD and C-TAD are especially different between HIF-1α and HIF-2α in terms of acidity (bottom).

1248

Figure 1—figure supplement 2. Verification of endogenous tagging of HIFs in 786-O ccRCC

cells. (**A**) (**B**) (**C**) Uncropped images of western blot of wild-type (WT) 786-O cells and

homozygously tagged knock-in clones (*). See Figure 1—figure supplement 2-source data 1-6 for

raw image files. (A) HIF-2α protein is detected in WT and various KIN clones with a Rabbit

monoclonal antibody against HIF-2α (Cell Signaling, D9E3), membranes were stripped and

reblotted for TBP for loading control. (B) HIF-1β protein is detected in WT and various KIN clones

with a Rabbit monoclonal antibody against HIF-1β (Cell Signaling, D28F3), membranes were

stripped and reblotted for TBP for loading control. (C) Various HIF-2α and HIF-1β KIN clones are

loaded on the same gel and probed for tagged protein levels using an anti-Halo antibody (top);

the membrane was stripped and reblotted for TBP for loading control, then stripped again and

reblotted for V5 tag. Halo-HIF-1β is generally expressed at a much higher level than Halo-HIF-2α,

as shown by the detected Halo or V5 level.

1261

Figure 1—figure supplement 3. HIF genome-wide binding and RNA expression profiles after gene-editing and genetic modification. (**A**) HeatMap of the ChIP-seq experiments comparing

binding profiles of HIF-2α and HIF-1β in WT or KIN clones. ChIP-Seq read counts (Reads Per

Genomic Content) are plotted at MAC2-called HIF-2α peak regions (across all cell lines and

subtracted of V5 peaks called in WT 786-O cells) centered around the peak. (**B**) Principle

Component Analysis (PCA) on RNA-seq data from WT, Halo-HIF-2α and Halo-HIF-1β KIN

clones, and WT cells over-expressing various HIF-α forms driven by an L30 promoter. Only the

first principle component (largest variance) is shown for three biological replicates. Both KIN

cells (HIF-1β KIN and HIF-2α KIN) and cells overexpressing inactive HIF-α mutants (HIF-1α

R30E, HIF-1αDM, HIF-2α R27E, HIF-2αDM) are close to WT cells. In contrast, cells

overexpressing functional HIF-α (HIF-1α and HIF-2α) show farther deviations from WT cells. (**C**)

DESeq2 output summarizing the number of differentially expressed genes (DEG) in KIN cell

lines and cells over-expressing various HIF-α forms, compared to WT cells.

1275

Figure 2—figure supplement 1. State Array (SA) analysis on fSPT data. (**A**) Schematic of the

SA workflow. Top: Step 1, naïve occupation is estimated for each state in an array of states

spanning different localization errors (y axis) and diffusion coefficients (x-axis), by simply applying

the RBME likelihood estimation on trajectory data. Middle: Step 2, refining posterior over diffusion

coefficient and localization error. Bottom: Step 3, marginalizing out localization error to generate

the posterior occupation (i.e. proportion of molecules, y-axis) as a function of diffusion coefficients

(x-axis) to get the final "diffusion spectrum". (**B**) Real example of SA analysis results from each of

the three steps. Trajectory data are collected from Halo-HIF-2α KIN clone A31, the same data

presented in Figure 2C.

Figure 2—figure supplement 2. HIF-1β behavior is distinct from HIF-2α and changes as 1β-to-2α stoichiometry changes. (**A**) RBME likelihood of diffusion coefficient in individual cells for four Halo-HIF-2α or Halo-HIF-1β KIN clonal lines. (**B**) Top and middle: proportion of molecules as a function of their diffusion coefficients (posterior mean occupations for a state array) evaluated on trajectories across all cells measured for additional Halo-HIF KIN line showing reproducible results in different KIN clones (Top, clone A31 and B50 for Halo-HIF-2α KIN and middle, clone A21 and B89 for Halo-HIF-1β KIN). Data for clone A21 and A31 are replotted from Figure 2.

1292 Bottom: Summary of the bound fraction (left) and peak diffusion coefficient (right) for all four cell

1293 lines. Each bar represents the average of three independent measurements on different days

1294 (black dots). (**C**) RBME likelihood of diffusion coefficient obtained for individual cells in either Halo-

1295 HIF-1β KIN cells (top) or Halo-HIF-1β KIN cells overexpressing HIF-2α (WT, middle or DM,

1296 bottom).

1297

Figure 2—figure supplement 3. Analysis of source of variation by bootstrapping. (**A-C**) Halo-
HIF-2α trajectories imaged from KIN clone A31 are used for subsampling and SA analysis.
Sampling (with replacement) was done by the number of trajectories (sample size = 10, 25, 50,
100, 200, 400, 800, 1600, 3200 and 6400), then analyzed with SA and the estimated bound
fraction was plotted as a dot. For each sample size, 100 replicates were performed and plotted

as 100 dots with box plot on top showing variation and the brown diamond indicating the mean.

(A) The main source of variation is cell-to-cell variability, followed by day-to-day (i.e. replicate-to-

replicate) variability. Left: Each sampling was drawn from pooled trajectories of 243 movies (cells)

taken on 11 different days. As the trajectory size increases, the results converge. Middle: Each

sample was drawn from a single cell that was randomly selected from 243 cells. Increasing the

number of trajectories does not improve the variance. Right: Each sample was drawn from all the

trajectories collected on a single day that was randomly selected from the 11 different days.

Increasing the number of trajectories improves the variance but some variation remains. (B)

Different number of cells are used for sampling to evaluate variation. For each graph, each sample

was drawn from N unique cells as indicated (N = a pooled of 243 cells; or 10, 20, 40, 60, 80, 100,

120 randomly selected from the pool of 243 cells). Replicates done with more than 60 random

cells can achieve a standard deviation less than 5% when the total trajectory number is over 800.

(C) A random pool of trajectories from 3 different days are used for each draw, showing that an

estimation with data randomly collected from 3 independent replicates (i.e. days) can achieve a

standard deviation less than 5% when the total trajectory number is over 800.

Figure 3—figure supplement 1. Dosage-dependent inhibition of HIF-2α binding and activity by

Belzutifan. (**A**) Molecular structure of Belzutifan (PT-2977) and schematic illustration of its function

to specifically block dimerization between HIF-2α/1β but not HIF-1α/1β. (**B**) Luciferase assay

using an HRE-containing reporter confirms Belzutifan dosage-dependent inhibition of HIF-2α

activity in WT 786-O cells. Error bars represent SE. (**C**) Proportion of molecules as a function of

diffusion coefficient for HIF-2α measured in Halo-HIF-2α KIN line clone A31 treated with different

concentrations of Belzutifan. DMSO levels are kept the same for all conditions at 0.0002%. (**D**)

The average fraction bound decreases (top) and the average peak position diffusion coefficient

increases (bottom) at increasing doses of Belzutifan. Individual measurements are indicated as

black dots. (**E**) RBME likelihood of diffusion coefficient in individual cells for all six drug dosages.

Figure 3—figure supplement 2. Additional data for Figure 3 showing HIF-α without the IDR is

not able to increase HIF-1β binding or decrease its diffusion coefficient (**A**) Schematic of

evaluating Halo-HIF-1β behavior while overexpressing only the N-terminal structured region

(NT) of HIF-α in combination with small molecule perturbation. Parental Halo-HIF-1β KIN cells

(grey background, same as in Figure 3) and cells stably overexpressing (O.E.) either HIF-2α NT

(orange in yellow background) or HIF-1α NT (red in purple background) are used, with and

without treatment with 0.2-µM Belzutifan. (**B**) Proportion of HIF-1β molecules as a function of

diffusion coefficient measured in various cells outlined in (A). Left: treated with DMSO only,

comparing HIF-1β behavior in the parental cells (black dashed curve, same as in Figure 3) and

in cells expressing either HIF-1α NT (purple curve) or HIF-2α NT (yellow curve). Middle and

right: proportions of HIF-1β as a function of diffusion coefficient, measured in cells expressing

either HIF-1α NT (right) or HIF-2α NT (middle) with Belzutifan treatment (grey curves),

compared to the DMSO control (purple or yellow curves). (**C**) Summary of the average bound

fractions (left) and peak diffusion coefficient (right) for conditions in (B), with black dots

indicating values from each of the three individual measurements.

1343

Figure 4—figure supplement 1. Quantification of initial localization density showing we imaged cells with similar expression levels of Halo-tagged proteins. (**A**) When cells were stained with

the same dye concentration, a much higher mean density of fluorescence spots at the first 10

frames (i. e. initial localization density) was observed for proteins with higher expression levels.

(**B**) In order to achieve a similar initial localization density, a much lower dye concentration has

to be used (~1/10 of concentration used for Halo-HIF-2α is needed for Halo-HIF-1β). When a

similar dye concentration is used (0.625 and 0.3125 nM), the initial localization density for the

same protein (Halo-HIF-1β) looks similar. (**C**) Western blot analysis of expression level of

various Halo-tagged proteins in KIN cells and in cells with L30-expression system. Bulk results

are mixed-population average and do not reflect expression levels from individual cells. (**D-G**)

Analysis of initial localization density for data used in Figure 4—figure supplement 2 (D), Figure

4 (E), and Figure 5 (F and G) showing similar initial localization density for all cells imaged for

each set of experiments, verifying comparable expression levels across the samples.

Figure 4—figure supplement 2. L30 weak expression system is able to recapitulate the
endogenous protein behavior. **(A)** Schematic of the L30 overexpression system where Halo-
HIF-2α is weakly expressed with an L30 promoter in wild-type 786-O cells (left) and Knock-in
cells where HIF-2α is endogenously tagged and expressed (right). **(B)** Comparison between
endogenous Halo-HIF-2α (KIN) and Halo-HIF-2α over-expressed with L30 promoter (L30),
showing very similar behavior. Left: Proportion of Halo-HIF-2α as a function of diffusion
coefficient and the corresponding bound fraction (shaded area). Right: Bar plot summarize the
average bound fraction (left) and peak diffusion coefficient (right) with three independent
measurements (black dots).

Figure 6—figure supplement 1. Cut&Run results show no difference in overall binding profiles

between different HIF-α variants. (**A-B**) PCA (A) and clustering (B) analysis of anti-V5 Cut&Run

results performed on WT 786-O cells, Halo-HIF-1α KIN clone A31, and WT cells expressing

Halo-tagged HIF-α variant (-1α, -2α, -2α/1α, or -1α/2α) with L30 promoter. Except for WT

control, all L30-driven HIF-α variants have very similar and close to endogenous HIF-2α binding

profile. (**C**) Two examples of genome browser views of Cut&Run results showing all four HIF-α

variants bind to the same sites with similar strength, even when the target genes are only

upregulated by overexpression HIF-α (left, PGK-1) or HIF-2α (right, TGFA).

Figure 6—figure supplement 2. Additional examples showing that HIF-α IDR contributes to
isoform-specific chromatin binding preferences. (**A**) two additional examples of genes that are
activated by HIF-1α and -2α/1α overexpression, and also have elevated binding by HIF-1α and -
2α/1α at regulatory sites (dashed boxes) (**B**) two additional examples of genes that are

1378   activated by HIF-2α and -1α/2α overexpression, and also have elevated binding by HIF-2α and -

1379   1α/2α at regulatory sites (dashed boxes).

1380

1381   **Figure 6—figure supplement 3.** Results from analyzing Cut&Run individual replicates showing

1382   reproducibility (**A-C**) Same analyses as for the combined replicates in Figure 6C were performed

1383   for Cut&Run individual replicates.

1384 **Table 4.** Constructs used to generate stable cell lines

1385

Name	Prom-oter	Gene product	Short name in the paper	Appeared in
PB EF1a 3XF EX-MCS IRES Puro	EF1a	3xFLAG tag	3xF	Figure 3
PB EF1a 3XF-GDGAGLIN-hEPAS1 IRES Puro	EF1a	HIF-2α N-terminally fused with 3xFLAG tag through a short peptide linker sequence (GDGAGLIN)	HIF-2α	Figure 2 Figure 3 Figure 2—figure supplement 2
PB EF1a 3XF-GDGAGLIN-hEPAS1_R171A-V192D IRES Puro	EF1a	HIF-2α dimerization mutant (R171A-V192D) N-terminally fused with 3xFLAG tag through a short peptide linker sequence (GDGAGLIN)	HIF-2α DM	Figure 2 Figure 2—figure supplement 2
PB EF1a 3XF-EPAS1_365 IRES Puro	EF1a	The N terminal region of HIF-2α (aa 1-365) N-terminally fused with 3xFLAG tag	HIF-2α NT	Figure 3—figure supplement 2
PB EF1a 3XF-EPAS1_365-364_HIF1a IRES Puro	EF1a	HIF-2α/1α chimera protein (aa 1-365 of HIF-2α and aa 364-826 of HIF-1α) N-terminally fused with 3xFLAG tag	HIF-2α/1α	Figure 3
PB EF1a 3XF-GDGAGLIN-hHIF1a IRES Puro	EF1a	HIF-1α N-terminally fused with 3xFLAG tag through a short peptide linker sequence (GDGAGLIN)	HIF-1α	Figure 3
PB EF1a 3XF-HIF1a_363 IRES Puro	EF1a	The N terminal region of HIF-1α (aa 1-363) N-terminally fused with 3xFLAG tag	HIF-1α NT	Figure 3—figure supplement 2
PB EF1a 3XF-HIF1a_363-366_EPAS1 IRES Puro	EF1a	HIF-1α/2α chimera protein (aa 1-363 of HIF-1α and aa 366-870 of HIF-2α) N-terminally fused with 3xFLAG tag	HIF-1α/2α	Figure 3
PB L30prom V5-Halo-GDGAGLIN-hEPAS1 IRES Puro	L30	HIF-2α N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN)	Halo-HIF-2α (L30)	Figure 4 Figure 4—figure supplement 2
PB L30prom V5-Halo-GDGAGLIN-hEPAS1_365-364_HIF1a IRES Puro	L30	HIF-2α/1α chimera protein (aa 1-365 of HIF-2α and aa 364-826 of HIF-1α) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-2α/1α (L30)	Figure 4
PB L30prom V5-Halo-GDGAGLIN-hHIF1a IRES Puro	L30	HIF-1α N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN)	Halo-HIF-1α (L30)	Figure 4
I_PB L30prom V5-Halo-GDGAGLIN-HIF1A_363-366_EPAS1 IRES Puro	L30	HIF-1α/2α chimera protein (aa 1-363 of HIF-1α and aa 366-870 of HIF-2α) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-1α/2α (L30)	Figure 4
PB L30prom V5-Halo-GDGAGLIN-hEPAS1 IRES Puro_R27E	L30	HIF-2α DBD mutant (R27E) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-2α, R27E (L30)	Figure 5
PB L30prom V5-Halo-GDGAGLIN-hHIF1a IRES Puro_R30E	L30	HIF-1α DBD mutant (R30E) N-terminally fused with V5-HaloTag	Halo-HIF-1α, R30E (L30)	Figure 5

		through a short peptide linker sequence (GDGAGLIN).		
PB L30prom V5-Halo-GDGAGLIN-hEPAS1_R171A-V192D IRES Puro	L30	HIF-2α dimerization mutant (R171A-V192D) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-2α DM (L30)	Figure 5
PB L30prom V5-Halo-GDGAGLIN-hHIF1a_R170A-V191D IRES Puro	L30	HIF-1α dimerization mutant (R170A-V191D) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-1α DM (L30)	Figure 5

1386
1387

1388 **Appendix 1. Analysis of source of variations.**

1389 Since we observed huge cell-to-cell variation (Figure 2A-B), we used bootstrapping to determine

1390 the source of variation, and decide how many cells, replicates and trajectories will be needed to

1391 get a relatively good estimate for the parameters.

1392

We used Halo-HIF-2α KIN clone A31 for the bootstrapping, as this is the cell line where the

most data were collected. We used 243 movies (each movie corresponds to one cell) collected

from 11 different replicates (each replicate was done on a different day) as the pool to draw

sample from. We performed sampling in three different ways. First, we pooled all trajectories

from all cells and randomly drawn different number of trajectories (i.e. sample size n= 10, 25,

50, 100, 200, 400, 800, 1600, 3200 and 6400) from the pool with replacement. After each draw,

the resulting trajectories were used for SA analysis to generate the diffusion spectrum and then

the fraction bound was calculated. We performed 100 draws for each of the sampling size and

plotted box plot on the resulting bound fraction estimation to see variation. This way, we can

estimate if we randomly sample n trajectories 100 times, how variable the result will be. The

result was as expected- the more trajectories we draw, the less variable the estimation is

between the 100 times of drawing when looking at the bound fraction (Figure 2—figure

supplement 3A, left).

1406

Second, instead of sampling a mixture of trajectories from all 243 cells, we randomly picked N

number of unique cells (N = 1, 10, 20, 40, 60, 80, 100, 120) from the 243 cells, pooled the

trajectories from these selected cells, and sampled from these pooled trajectories. The sample

size and number of draws were kept the same. This way, we can compare if we randomly take

only N cells from the population (243 cells) to estimate the bound fraction, do that 100 times for

each sample size, how bad the variation is compared to if we take all 243 cells. This will be a

good estimation of how many cells we need to pick. Certainly, small number of cells gives

smaller number of total trajectories and if we draw large number of trajectories from them (such

as n = 6400) it will be over-sampled. Our result shows that when we pick as many as 60 cells,

the difference of estimated bound fraction from these 60 cells compared to when estimated from

all 243 cells can be less than 5% when sampling more than 800 trajectories (Figure 2—figure

supplement 3A, middle; and 3B).

1419

Third, since different replicates were performed on different days, we use day as the unit to

group trajectories. We randomly picked 1 day (i.e. 1 replicate), combined all trajectories from

Halo-HIF-2α KIN clone A31 from that day, which normally is ~20 cells, and use this mixture of

trajectories for sampling. This way, we can compare how variable our estimation will be from

replicate to replicate. Indeed 1 replicate is not good enough to give good estimation as the

variation is still high even if we sample large number of trajectories (Figure 2—figure

supplement 3A, right). However, if we randomly pick 3 days instead of 1, we reduce the

variance, and the estimated bound fraction from these 3 days compared to when estimated from

all 243 cells from 11 days can be less than 5% when sampling more than 800 trajectories

(Figure 2—figure supplement 3C).

1430

1431 From our bootstrapping results, we think an estimation of bound fraction from 3 replicates with

1432 20 cells each replicate (and usually we can collect total more than 2500 trajectories from 3

1433 replicates) can be good enough, at least for comparing different conditions such as WT versus

1434 mutants, where a difference of 10% bound fraction was usually observed.

Appendix 2. Using fluorescence intensity as approximation for protein expression level.

Since we have shown that stoichiometry may affect protein behavior, to compare the molecular dynamics across different HIF-α forms that were exogenously expressed, it is important to control for similar expression levels. Unfortunately, we observed variegated expression levels with the same L30 promoter even with stable cell lines. As a result, for exogenously expressed Halo-proteins in Figure 4 and Figure 5, we used fluorescence intensity as approximation for protein expression levels to guide our selection of which cell to image.

During imaging, cells were doubly labeled- one channel with excess JFX549 dye (500 nM), which was used for localizing cells and define ROI, the other channel with limited JFX646 dye for fSPT (Figure 1D). The expression level for the Halo-tagged protein can be estimated by either looking at fluorescence intensity from the JFX549 channel, or looking at the initial localization density (i.e. average number of molecules detected per ROI) from the JFX646 channel. For quantification of initial localization density, we only averaged the first 10 frames of each movie, because the initial frames have minimum impact from photobleaching.

For Halo-HIF-1β and Halo-HIF-2α KIN lines, because the endogenous HIF-1β expresses at a much higher concentration than HIF-2α, when cells were fully labelled with JFX549, the fluorescence level for Halo-HIF-1β KIN cells were much higher than Halo-HIF-2α KIN cells. Similarly, when stained with same concentration of JFX646 for these two cell lines, the overall Halo-HIF-1β localization density was much higher compared to Halo-HIF-2α, consistent with much higher expression levels (Figure 4—figure supplement 1A). However, for SPT, a low localization density is needed to minimize tracking error. To keep localization density low, for Figure 2, Halo-HIF-1β had to be stained at ~1/10 of the JFX646 concentration used for Halo-HIF-2α (Figure 4—figure supplement 1B, 0.3125 or 0.625 nM instead of 5 nM) to achieve ideal localization density that is similar to Halo-HIF-2α (orange dots). This shows that for different expression levels, an order-of-magnitude difference in dye concentration needs to be used to get similar localization density. In addition, we found that while 5 nM gives very high localization density, when stained at a similarly low concentration (0.3125 vs 0.625 nM), the resulting localization density for Halo-HIF-1β was also roughly the same. This shows that with similar dye concentration, localization density can be a rough approximation for protein levels.

We then utilize the localization density as a quality control for experiments where expression level needs to be kept similarly. For the L30 driven Halo-HIF-α expression, the variation of

expression level is big from cell to cell and we are able to find cells with similar expression

levels (close to endogenous) across different HIF-α variants. Thus, we used the JFX549

channel to pick cells with similar fluorescence intensity (usually the lowest intensity, which is the

closest to endogenous expression level) for imaging, after cells were fully labeled with JFX549.

Because these cells were also stained with similar level of JFX646 dye for fSPT when a set of

experiments with multiple conditions were compared, cells picked for similar JFX 549 intensity

usually have similar localization densities in the JFX 646 channel. As confirmed by

quantification in Figure 4—figure supplement 1D-G, we did have very close localization density

for different conditions in each set of experiments performed, indicating that for each

comparison, the expression level across different conditions were kept roughly similar.