Mechanisms Governing Target Search and Binding Dynamics of Hypoxia-Inducible Factors

Yu Chen[1,2,3], Claudia Cattoglio[1,2,3], Gina Dailey[1,3], Qiulin Zhu[1,3], Robert Tjian[1,2,3] *, and Xavier Darzacq[1,3] *

[1] Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA.

[2] Howard Hughes Medical Institute, University of California, Berkeley, CA, USA.

[3] Li Ka Shing Center for Biomedical & Health Sciences, University of California, Berkeley, CA, USA.

*Correspondence: R.T. (jmlim@berkeley.edu), X.D. (darzacq@berkeley.edu)

Abstract

Transcription factors (TFs) are classically attributed a modular construction, containing well-structured sequence specific DNA-binding domains (DBDs) paired with disordered activation domains (ADs) responsible for protein-protein interactions targeting cofactors or the core transcription initiation machinery. However, this simple division of labor model struggles to explain why TFs with identical DNA binding sequence specificity determined *in vitro* exhibit distinct non-overlapping binding profiles *in vivo*. The family of Hypoxia-Inducible Factors (HIFs) offer a stark example: aberrantly expressed in several cancer types, HIF-1α and HIF-2α subunit isoforms recognize the same DNA motif *in vitro* – the hypoxia response element (HRE) – but only share a subset of their target genes *in vivo*, while eliciting contrasting effects on cancer development and progression under certain circumstances. To probe the mechanisms mediating isoform-specific gene regulation, we used live cell single particle tracking (SPT) to investigate HIF nuclear dynamics and how they change upon genetic perturbation or drug treatment. We found that HIF-α subunits and their dimerization partner HIF-1β exhibit distinct diffusion and binding characteristics that are exquisitely sensitive to concentration and subunit stoichiometry. Using domain-swap variants, mutations, and a HIF-2α specific inhibitor, we found that although the DBD and dimerization domains are important, a major determinant of chromatin binding and diffusion behavior is dictated by the AD-containing intrinsically disordered regions. These findings reveal a previously unappreciated role of IDRs in regulating the TF search process that may play a role in selective functional target site binding on chromatin.

Introduction

Sequence-specific transcription factors (TFs) are key frontline regulators of gene expression. Classical LexA-Gal4 domain-swap experiments in yeast presented a simple modular structure and apparent division of labor for typical TFs (Brent and Ptashne, 1985). In this textbook paradigm, the DNA-binding domain (DBD) is responsible for DNA sequence recognition and binding specificity while the activation domain (AD) is responsible for target gene transactivation that involves protein-protein interactions with co-factors, the basal transcription machinery and other ancillary factors that are generally devoid of sequence specific DNA recognition. In higher eukaryotes, each DBD class usually contains multiple closely related family members. For example, the bHLH class of TFs includes MyoD, Clock, and Max. They all recognize the same E-box DNA binding sequence motif 5'-CACGTG-3', yet each differentially regulates muscle differentiation, circadian rhythm, and cell proliferation, respectively (Kribelbauer et al., 2019). This raises the specificity paradox: how do TFs with seemingly identical DNA sequence specificity, at least as determined *in vitro*, nevertheless exhibit non-overlapping binding profiles *in vivo* and carry out distinct and even opposing functions? In general, when confronted with this conundrum, we have assumed that one or more co-factors or perhaps still to be identified "silent partner" TFs can somehow divert target recognition to a composite cis-regulatory site distinct from the canonical DNA binding site. Given the high occurrence of short binding motifs for most TFs throughout the genome, even with co-operative binding to composite sites, most potential specific binding sites nevertheless remain unoccupied as determined by genome-wide TF binding studies. What feature or motif within TFs outside of the DBD and dimerization domain may be responsible for such differential site selection has remained unclear. Thus, the simple rule of modular units with well separated divisions of labor between DBD, dimerization and transactivation may deserve a closer look. We also wondered whether quantitative single molecule dynamics measurements might reveal new aspects of TF behavior in living cells that could inform us regarding potential mechanisms influencing the target search process and differential site selectivity in a native physiologically relevant context.

Here we have chosen the Hypoxia-Inducible Factors (HIFs) as a representative example to study the paradox of highly conserved DBDs carrying out distinct target site selection and to dissect potential novel features of TFs that mediate chromatin binding. HIFs are a family of α/β heterodimeric TFs stabilized under hypoxic conditions to promote angiogenesis, anaerobic metabolism, cell proliferation and "stemness" (Semenza, 2012). The oxygen-labile alpha subunits (mainly HIF-1α and HIF-2α) complex with their oxygen-stable beta partner (mainly HIF-1β) to form

a functional dimer (Fig. 1A). All HIF subunit isoforms belong to the bHLH-PAS (Basic helix-loop-

helix-PER-ARNT-SIM) family, where the N-termini are structured domains containing bHLH (DNA

binding) and PAS (dimerization) domains, while the C-termini consist of intrinsically disordered

regions (IDRs) containing ADs (Fig. 1A and Fig. S1). HIF-1α/1β and HIF-2α/1β dimers share a

conserved structural fold (Wu et al., 2015), recognize the same hypoxia response element (HRE)

5'-TACGTG-3' binding motif (Schödel et al., 2011; Wenger et al., 2005), but share only a partial

overlap of target genes *in vivo* (Smythies et al., 2019). With their own unique target gene sets,

HIF-1α and -2α can exert divergent and even contrasting functions (Keith et al., 2012). For

example, while both HIF-1α and HIF-2α regulate angiogenesis, HIF-1α specifically regulates

glycolysis, apoptosis, and promotes NO production, whereas HIF-2α binds to the *POU5F1* locus

to maintain Oct4-regulated stem cell identity and pluripotency, promotes cell cycle progression,

and inhibits NO production (Keith et al., 2012; Smythies et al., 2019). Therefore, our current simple

textbook model of exchangeable modular TF functional units does not satisfactorily explain such

isoform-specific target gene regulation.

The HIF family differential specificity paradox is even more daunting to comprehend at the level

of disease inducing mechanisms. HIFs are aberrantly upregulated and recognized as oncogenic

drivers in multiple cancers. However, in addition to their shared roles in cancer onset and

progression, HIF-1α and -2α also show many independent, sometimes even opposing roles in

specific contexts (Keith et al., 2012). For example, in clear cell renal cell carcinoma (ccRCC), HIF-

2α is the critical tumorigenic driver whereas HIF-1α, in contrast with its usual tumorigenic role, is

mostly tumor-suppressive (Raval et al., 2005; Schödel et al., 2016). The regulatory mechanism

behind such highly divergent outcomes is still largely unknown. Given such complexity, without a

deeper understanding of isoform-specific transcriptional regulation, it is hard to predict the

functional outcomes mediated by individual HIF isoform in various cancer types or stages, which

could be a complicating factor in developing more effective HIF-targeting cancer therapeutics.

In this study, we aim to understand the molecular mechanisms mediating isoform-specific target

gene regulation at its most fundamental level – could we detect differential molecular dynamics

of distinct TF isoforms during the target search and chromatin binding process in live cells? Which

regions or domains of TFs might be responsible for such isoform specific properties? Could we

begin to discern possible mechanisms that guide TFs with highly conserved DBDs to their distinct

and specific targets beyond cognate DNA sequence recognition? Here we use HIFs as an

illustrative example, combining endogenous tagging and super-resolution single particle tracking

(SPT) (Liu et al., 2015) to study the dynamic behavior of these key gene regulators in live cells
under physiological conditions. We also dissect the contribution of different domains of HIF-α
isoforms by a series of mutation and domain swap experiments to directly test the concept of
modular functional domains. Deploying a combination of genetic and small-molecule
perturbations, we found that, although HIF DBD and dimerization are important for DNA target
acquisition, the amount of protein bound and its diffusion characteristics are mainly driven by
regions outside the DBD and dimerization domains. Our results reveal a previously unappreciated
role of unstructured domains in the target search and binding properties of TFs to functional
chromatin sites in live cancer cell context.
111
112   **Results**
113
Establishing a human cancer cell system for live-cell single molecule imaging of HIF
To investigate HIF dynamics, we first focused on one of the cognate dimers: HIF-2α/1β. We used
the common ccRCC line 786-O (Brodaczewska et al., 2016), derived from a VHL-deficient, H2
type primary clear cell renal cell carcinoma, wherein HIF-2α is stabilized due to an inactivating
mutation in VHL (the E3 ubiquitin ligase that targets all HIF-α isoforms for proteasomal
degradation)(Gnarra et al., 1994). The 786-O line also conveniently lacks any functional HIF-1α
due to a truncating mutation of HIF-1α (Shen et al., 2011; Swiatek et al., 2020), which allows us
to study one α isoform independently from the other. Using CRISPR/Cas9-mediated genome
editing, we successfully generated several clonal lines with homozygous knock-in (KIN) of the
HaloTag (Los et al., 2008) at the N-terminus of either HIF-2α or its binding partner, HIF-1β (Fig.
1B and Fig. S2, A-C). Western blotting confirmed that the tagged proteins are expressed at levels
similar to wild-type (WT) in unedited cells (e.g., HIF-2α clone A31 and HIF-1β clone A21) (Fig.
1B). Confocal imaging after covalently labeling cells with a fluorescent Halo-binding ligand
(JFX646) (Grimm et al., 2021) shows the expected nuclear localization for both Halo-HIF-2α and
Halo-HIF-1β proteins (Fig. 1C). In addition, using ChIP-seq, we confirmed that both tagged
proteins maintain a similar genome-wide binding profile as the WT protein in unedited cells (Fig.
S2D). We have thus established a human cancer cell system suitable for live-cell imaging on HIF-
2α and HIF-1β at endogenous expression levels.
132
To evaluate how HIF-2α and -1β explore the nucleus and bind DNA, we used the fast modality of
super resolution live cell single particle tracking (fSPT) that is capable of tracking rapidly diffusing
molecules. Cells with either HIF-2α or HIF-1β Halo KIN were doubly labeled with the live-cell

permeable Halo-binding JFX dyes (Fig. 1D) and were imaged under highly inclined and laminated
optical sheet illumination (HiLo) (Tokunaga et al., 2008) at high frame rates to capture the
movement of single molecules in their native nuclear environment (Fig. 1E). Stroboscopic
illumination at high excitation power is used to minimize motion blur, while sparse labeling ensures
only a limited number of molecules are detected at any given time in the nucleus to minimize
misconnections when computing the path of individual molecules (trajectories) (Fig. 1E and F).
We can then estimate relevant kinetic parameters from these trajectories, extracting quantitative
information such as diffusion coefficients and bound fraction.
144
fSPT detects various HIF molecular states in their native nuclear environment
To quantitatively analyze the acquired fSPT data, we used a non-parametric Bayesian approach
(Heckert et al., 2021) recently developed in our lab. We chose to use this new method, because
as dimeric transcription factors, HIF subunits could conceivably exist in multiple states (e.g.,
bound, moving as a monomer, moving as a dimer, and moving in a bigger complex containing
co-regulators) (Fig. 2A and 2B). One feature distinct from our previous approach, which required
the assumption of a fixed and limited (3 or less) number of states for the tracked protein (Hansen
et al., 2018), our new data analysis method does not require *a priori* knowledge of how many
underlying states exist for each tracked protein. Instead, it estimates the constituents in a range
of dynamic states with diffusion rates from 0.01 to 100 $\mu m^2 s^{-1}$ while accounting for known
experimental biases due to localization error and fluorophore defocalization (Heckert et al., 2021).
After aggregating all trajectories from all the cells examined, we generate a distribution of diffusion
coefficient estimates that reports the fraction of stably bound molecules while simultaneously
displaying the full behavioral spectrum of the diffusing molecules (Heckert et al., 2021) (Fig. 2C).
159
We observed highly heterogeneous results for both HIF-2α and HIF-1β, with varied diffusion
coefficient estimates from cell to cell, when applying the likelihood estimation to individual cells
(Fig. 2A and 2B, Fig. S3A, clones A31 and A21). These results indicate that for both HIF proteins,
a range of moving states likely exists. We then pooled trajectories from many cells (n ~60) to
estimate the distribution of diffusion coefficients for the population. Strikingly, we observed a very
different behavior for Halo-HIF-2α compared to Halo-HIF-1β. Whereas a large fraction (about 40%)
of Halo-HIF-2α is immobile and presumably chromatin-bound (defined as the fraction with a
diffusion coefficient < 0.1 μm^2/sec), the majority (above 70%) of Halo-HIF-1β appears freely
diffusing (Fig. 2C and S3B, clones A31 and A21). Also, the overall diffusion coefficient for the
Halo-HIF-1β mobile population is much larger than that of Halo-HIF-2α. We repeated

measurements in different KIN clones and confirmed the reproducibility of these results for both

Halo-HIF-2α (Fig. S3A-B, clone B50) and Halo-HIF-1β (Fig. S3A-B, clone B89). The differences

between HIF-2α and -1β seem counterintuitive at first, because one would expect HIF-2α and

HIF-1β to behave similarly since they should exist as a hetero-dimer. However, since the

endogenous HIF-1β is expressed at a much higher level than HIF-2α (Fig. S2C), the majority of

HIF-1β is free to diffuse without HIF-2α. Of note, the distribution plot only reflects the fraction of

molecules as a function of their diffusion coefficient, but does not report on the absolute number

of molecules. Therefore, a smaller bound fraction for Halo-HIF-1β does not mean fewer numbers

of bound molecules than Halo-HIF-2α, since many more Halo-HIF-1β molecules are present in

the nucleus. Given this scenario, we hypothesized that HIF-1β molecular dynamics and percent

binding should be modulated by changing the 2α/1β stoichiometry.

HIF-1β binding and diffusing dynamics can be modulated by HIF-α:β stoichiometry, and

are dependent on dimerization

To test the hypothesis that HIF-1β dynamics depends on 2α/1β ratio, we first tried to modulate its

behavior by stably over-expressing HIF-2α in the endogenously HIF-1β Halo-tagged KIN line (Fig.

2D). We found that the mobile population of Halo-HIF-1β diffuses more slowly when HIF-2α is

overexpressed, most likely due to its dimerization with the extra HIF-2α to form dimers capable of

DNA/chromatin binding. As expected, we also observed a significant increase in the Halo-HIF-1β

bound fraction (up to 50%), (Fig. 2E, Fig. S3C top and middle). To confirm that the changes in

HIF-1β dynamics caused by increasing levels of HIF-2α are dependent on hetero-dimerization,

we stably overexpressed the HIF-2α R171A/V192D double mutant (HIF-2α DM) that was

previously reported to lose its dimerization capability with HIF-1β (Wu et al., 2015). As expected,

overexpression of HIF-2α DM did not increase the bound fraction or decrease the overall diffusion

speed of Halo-HIF-1β to the same extent seen with WT HIF-2α overexpression (Fig. 2E, Fig. S3C

bottom), suggesting that the changes we observe are dimerization-dependent.

We further validated our results by stably overexpressing different forms of HIF-α in the HIF-1β

Halo-tagged KIN line and treating cells with a HIF-2α-specific small molecule inhibitor, Belzutifan

(PT-2977). Belzutifan inhibits HIF-2α/1β, but not HIF-1α/1β, dimerization by specifically binding

to the dimerization domain of HIF-2α (Fig. S4A), and thus has been used as an HIF-2α inhibitor

for ccRCC treatment (Wallace et al., 2016; Xu et al., 2019). We first confirmed that Belzutifan

inhibits HIF-2α transcription function in a dose-dependent manner (Fig. S4B). Importantly,

Belzutifan also reduces the HIF-2α bound fraction in the HIF-2α Halo-tagged KIN line in a similar

dosage-dependent manner, again revealing the potential of fSPT to measure TF dynamics and
associated functional changes (Fig. S4, C-E). We choose to use 0.2 µM Belzutifan for all
subsequent experiments to maximize its effect.
207
Next, we carried out a series of experiments designed to probe the consequences of swapping
different functional domains of HIF-1α and HIF-2α to determine which parts of these closely
related TFs might be involved in selective activities when paired with HIF-1β. Using the HIF-1β
Halo-tagged KIN line as the parental line, we stably overexpressed WT or chimeric HIF-α, where
we swapped the structured and disordered domains between HIF-1α and HIF-2α (Fig. 3A). All
these different HIF-α variants are expressed from a relatively strong EF-1alpha promoter and are
N-terminally 3xFLAG-tagged. A construct that expresses 3xFLAG only is used as control. We
then treated these cells with either Belzutifan or DMSO control and measured Halo-HIF-1β
dynamics (Fig. 3A). While 3xFLAG tag had no effect, overexpressing HIF-α, regardless of which
variant form, is able to both increase the bound fraction and reduce the overall diffusion speed of
HIF-1β (Fig. 3B top, and 3C, DMSO group). For cells overexpressing the α variants that contain
the HIF-2α structured domain, this effect on HIF-1β can be at least partially reverted after
Belzutifan treatment (Fig. 3B and 3C, +HIF-2α and +HIF-2α/1α). In contrast, for cells
overexpressing the α variants that contain the HIF-1α structured domain, this effect is resistant to
Belzutifan, consistent with the subunit isoform specificity of the drug for HIF-2α (Fig. 3B and 3C,
+HIF-1α and +HIF-1α/2α). In untransfected and 3xFLAG only overexpressing control cells,
treatment with Belzutifan only weakly reduces the HIF-1β bound fraction, again suggesting that
the majority of HIF-1β is not engaged with 2α (Fig. 3B and 3C, parental cell and +3xF). Overall,
these results demonstrate that HIF-1β dynamics change after engagement with its α partner and
can be selectively inhibited with a specific dimerization inhibitor. The observed differences also
confirm that fSPT is a powerful platform to monitor molecular dynamic changes of TFs in living
cells thus, allowing us to gain new mechanistic insights while we introduce various perturbations,
such as subunit concentration or stoichiometry and specific mutations.
231

Regions outside the DBD/dimerization domain determine HIF molecular dynamics
Interestingly, comparing the effects of the four different α variants, we found that regardless of
their structured domain, those with the same C terminal IDRs behave similarly (Fig 3B and 3C,
middle and right). Specifically, the variants containing the HIF-2α IDR have a stronger effect on
increasing HIF-1β binding than the variants containing the HIF-1α IDR. Thus, surprisingly, it
appears that the bound fraction of HIF-1β is not determined by the HIF-α DBD, but rather by HIF-

α IDR, which we found rather counterintuitive. To confirm the importance of HIF-α IDRs in HIF

binding, we overexpressed a truncated version of either HIF-1α or 2α that contains only the N-

terminal structured region (HIF-1α NT or HIF-2α NT), which still maintains both the DBD and

dimerization capability for interacting with HIF-1β (Wu et al., 2015). Indeed, both these truncated

forms lacking the IDR/AD of HIF-α minimally affect the HIF-1β bound fraction (Fig. S5).

Surprisingly, these truncated HIF-α variants also only marginally influenced the overall HIF-1β

diffusion speed. These results indicate that dimerization alone neither increases HIF-1β binding

nor reduces the overall diffusion speed of its moving population. Instead, the extended HIF-α AD-

containing IDR is necessary to influence and direct HIF-1β behavior.

To further test our hypothesis that HIF chromatin binding and the dynamics of the diffusion

population are dominated by the α subunit IDR, we switched to image the α subunit itself. We

made different forms of Halo-tagged HIF-α (WT and domain-swapped), stably but weakly

expressed them in WT 786-O cells with an L30 promoter (Fig. 4A). We first confirmed that binding

and diffusion characteristics of L30-expressed Halo-HIF-2α are very similar to the endogenous

Halo-HIF-2α in the KIN line (Fig. S6), demonstrating that weak overexpression can largely

recapitulate endogenous protein behavior. Therefore, this system provides a convenient tool to

investigate the contribution of each domain of HIF-α in the target search and binding process.

Much like our results with endogenous HIF-1β, we observed similar behaviors of HIF-α proteins

if they contain the same IDR (Fig. 4B top and middle), while displaying distinct behaviors when

endowed with different IDR isoforms (Fig. 4B bottom). Regardless of which DBD they have, the

variants containing the HIF-2α IDR (WT HIF-2α and HIF-1α/2α) show a higher bound fraction,

compared to the ones containing HIF-1α IDR (WT HIF-1α and HIF-2α/1α) (Fig. 4C). These results

suggest that indeed the disordered region on HIF-α determines how HIFs bind and diffuse in the

nucleus, and that the HIF-2α AD-containing IDR mediates HIF binding to chromatin and/or some

other relatively immobile components in 786-O cells.

HIF-α disordered region is necessary but not sufficient for optimal binding

The fact that the extent of binding (presumably to chromatin) of HIF proteins depends mainly on

the long C-terminal IDR rather than on their DBD was unexpected. Therefore, we next examined

the contribution of the HIF DBD to the bound fraction. We introduced point mutations in the DBD

(HIF-2α R27E and HIF-1α R30E) that were previously reported to impair DNA binding (Michel et

al., 2002; Wu et al., 2015), and expressed them in the WT 786-O cells with the same L30 promoter

system (Fig. 5A). Not surprisingly, DBD mutants show a reduction in the bound fraction and a

concomitant increase in the diffusing fraction compared to their WT counterpart (Fig. 5B and 5C).

In agreement with the expectation that the DBD mutations should not perturb protein-protein

interactions, we do not observe a significant change in the overall speed of the diffusing population.

These results demonstrate that, although the AD-containing IDR is the major modulator in

determining the bound fraction, the DBD is also important for binding, further suggesting that the

observed bound fraction likely represents chromatin/DNA binding.

We next examined whether dimerization with HIF-1β is required for HIF-α chromatin binding.

Taking advantage of the same L30 weak expression system, we exogenously expressed the

Halo-HIF-2α dimerization mutant (R171A/V192D), or the analogous Halo-HIF-1α dimerization

mutant (R170A/V191D) in the WT 786-O cells (Fig. 5D). We found that compared to the WT Halo-

HIF-2α or -1α, these mutants exhibit a significantly decreased bound fraction (Fig. 5E and 5F),

demonstrating that HIF-α without -1β can no longer effectively bind to DNA/chromatin. Taken

together, our results indicate that the HIF-α disordered region alone is not sufficient to maintain

binding, but instead, the IDR and both the DBD and dimerization domains are also needed.

Intrinsic properties of HIF-α IDR determine the overall speed of diffusive HIF.

Interestingly, with the Halo-HIF-α dimerization mutants, we observed no obvious change in their

overall diffusion coefficient in the moving population (Fig. 5E), indicating that losing their HIF-1β

partner does not affect the overall HIF-α diffusion speed. This result suggests that it is some

intrinsic property of HIF-α molecules, rather than the molecular weight of dimers versus

monomers, that determines its diffusion speed and behavior. Our results suggest that while the

moving population of HIF-1β alone diffuses relatively fast, the moving population of both HIF-α

and HIF- α/β dimers diffuses relatively slowly. We postulate this is potentially due to the HIF-α

IDR engaging in protein-protein interactions with various cofactors both when associated with

HIF-1β or when alone. (Fig. 6A). Indeed, this is consistent with our previous observation that the

HIF-a NT/HIF-1β dimer diffuses at a relatively fast speed, similar to HIF-1β alone which apparently

does not share this HIF-α IDR mediated capacity (Fig. S5).

Discussion

Transcription factors must search, recognize and bind to their specific target sites among millions

of possible DNA sequences along chromatin to activate the correct gene. With the successful

development of X-ray crystallography and cryo-EM, mechanisms of DNA-binding specificity have

been extensively studied, primarily based on classically structured globular DNA-binding domains

of TFs. We now know that a variety of structural mechanisms are used to recognize DNA,
including formation of specific hydrogen bonds and DNA contour interactions (Rohs et al., 2010).
However, these inherent binding modalities of DBDs alone cannot explain TF binding site
selection *in vivo*. As revealed by genome-wide *in vivo* binding assays, only a subset of potential
target sites become occupied, and this is not entirely consistent with either DNA binding site
affinity or chromatin accessibility (Behera et al., 2018; Grossman et al., 2017; Srivastava and
Mahony, 2020). On the other hand, TFs have long been recognized to also contain long
unstructured transactivation domains with simple amino acid composition (Gln-rich, acidic, Pro-
rich etc.), which often posed challenges to purification and/or crystallization of full-length TFs
(Courey and Tjian, 1988; Ma and Ptashne, 1987; Mermod et al., 1989; Tjian and Maniatis, 1994).
Recently, such intrinsically disordered regions (IDRs) were reported to play an important role in
weak and multivalent protein-protein interactions to form local small transient hubs that, when
exacerbated by overexpression, can drive phase separation. Although not structurally defined,
these interactions can still be sequence/amino acid composition selective (Chong et al., 2018;
Chong and Mir, 2021). IDRs are now proposed to have important functions in boosting gene
expression through hub or condensate formation to locally enrich for factors that are needed for
transcription (Boijja et al., 2018; Cho et al., 2018; Chong et al., 2018; Sabari et al., 2018; Wei et
al., 2020). However, few studies of IDRs have investigated their potential role in DNA binding site
search and selection. Some studies reported that for a subset of zinc finger proteins (Sp2 and
KLF3), an IDR is critical for *in vivo* binding and specificity (Burdach et al., 2014; Lim et al., 2016;
Völkel et al., 2015), and another recent study using genomic approaches reports the IDR as a
determinant for specificity of the yeast bZIP TF Yap1 (Brodsky et al., 2020).

Here, using advanced live cell single particle tracking, we report that TF IDRs previously
associated with ADs are, in fact, a major determinant mediating nuclear search dynamics and
chromatin binding characteristics. Employing both genetic and small molecule perturbations
together with a series of domain-swap and mutation experiments, we found that it is the AD
associated disordered region of HIF-α rather than the intrinsic molecular weight of the TF that
dictates a relatively slow diffusion for both HIF-α monomers and HIF-α/β dimers. On the other
hand, when not engaged with HIF-α, HIF-1β diffuses rapidly as expected for an unencumbered
subunit. These results indicate that the diffusion characteristic of HIF molecules is profoundly
influenced by the properties of their disordered regions (Fig. 6A). In fact, computational analysis
shows very different amino acid composition bias among HIF-1α, -2α and -1β disordered regions
(Fig. S1B). Thus, it is very likely that as these molecules navigate through the crowded nuclear

environment, their distinct stretches of IDRs that also contain ADs make differential and selective
interactions with other nuclear components, resulting in distinct diffusive behaviors. It is also
possible that due to differences in acidity, the different charges on these IDRs can cause
differential interactions with macromolecules including not only proteins, but also DNA and RNA
(Xiang et al., 2020).

While it is easy to conceptualize how IDRs can influence the speed of diffusion, one unexpected
result is that they also largely determine how much and with what differential specificity TFs bind
to chromatin. Although we confirmed that the DBD and dimerization domains are important for
binding, the surprise was that our domain swap experiments clearly demonstrated that the
percentage of bound TF is mainly contributed by regions outside of the DBD/dimerization domains.
One explanation could be the differential charge propensities of the different disordered regions
(Fig. S1B). For example, the HIF-2α IDR is more positively charged and may not only slow down
nuclear exploration but also stabilize chromatin binding, possibly through stronger interactions
with negatively charged chromatin-associated RNA and/or nucleosome-free DNA regions.
Besides direct chromatin interactions, HIF-2α IDR could also increase and stabilize binding via
indirect interactions with other chromatin-bound proteins. Moreover, since different IDRs can
selectively interact with other IDRs (Chong et al., 2018; Chong and Mir, 2021), we also postulate
that selective interactions with other TFs or co-regulators may play a role in determining HIF
genome-binding specificity. One hint of such a "combinatorial TF selectivity mechanism" is that
HIF-2α binding sites were frequently found adjacent to AP1/Fox binding sites, while HIF-1α
binding sites were usually found next to HEY/SP1 binding sites (Smythies et al., 2019). It was
also previously reported that while no target specificity was preserved in reporter gene assays,
the N-terminal TAD of HIF-α conferred endogenous target specificity for two of the HIF-1 unique
genes examined, possibly via specific interactions with transcriptional cofactors (Hu et al., 2007).
Further Co-IP or pull-down assay coupled with mass spectrometry (MS) will be needed to more
fully dissect this type of *in vivo* selectivity mechanism.

Interestingly, Myc, a closely related bHLH-LZ family transcription factor, shows
predominately DBD-mediated chromatin engagement, where non-specific DBD-DNA
interactions contribute most of the binding (Pellanda et al., 2021). This non-specific
interaction agrees well with the 1D sliding mode of TF-DNA interaction (Normanno et al.,
2015). In contrast, for HIF, we observed an additional mode of TF-chromatin interaction

that is modulated by an IDR and likely involves protein-protein interactions. These
differences indicate that eukaryotic transcription factors may exploit distinct mechanisms
when engaging with chromatin, some mainly involving classical DBD-DNA interactions
while others rely on regions outside of DBD to contribute to their binding and potential
target selection.

Finally, we have shown that our fSPT platform provides a powerful tool able to resolve *in vivo*
protein dynamics that is exquisitely sensitive to concentration, subunit stoichiometry and
genetic/small molecule perturbations. This is especially important when studying TFs, where a
slight difference in expression level often generates completely different results, rendering over-
expression systems highly susceptible to artifacts. It is also worth underscoring the importance of
studying TFs in their native physiologically relevant chromatin environment, given their obligate
interactions with higher-order chromatin structures and cofactors. For example, the EPO gene is
reported to be responsive to HIF-2α but not HIF-1α in Hep3B cells (Warnecke et al., 2004) and in
murine liver (Rankin et al., 2007), however, a luciferase reporter driven by the upstream EPO
enhancer also responds strongly to HIF-1α (Varma and Cohen, 1997), which may generate
misleading results and interpretations. Our fSPT platform allows us to study transcriptional
regulation in the native chromatin context and with endogenous TF levels to obtain data with
physiological and functional relevance. Such live cell real time measurements under native cell
contexts could prove to be highly valuable, both for dissecting *in vivo* mechanisms of transcription
regulation, and for guiding the development of effective therapeutics. Our Belzutifan treatment
experiment is an example of how fSPT can reveal the mechanism of action of small molecule
inhibitors, and how it could serve as a powerful tool to screen for drugs that selectively target one
isoform versus another, using dimerization and binding readouts as indicators of efficacy and
specificity. Moreover, since our results demonstrated how IDRs can affect TF diffusion behavior,
potentially distinct dynamic features determined by a particular IDR can be exploited as a readout
for screening small molecules or peptides that target allosteric sites of TFs. Assays that can
quantitatively measure TF diffusive behavior in live cells could be transformative for advancing
drug discovery because a high throughput imaging strategy opens the door to effectively target
what has been traditionally considered "undruggable", such as most protein-protein interactions
including potentially unstructured TF activation domains.

In summary, using the HIF protein family as a case study, we uncovered a mechanism of IDR-
mediated nuclear search and differential chromatin binding. We expect this fundamental principle
to be applicable to a broad range of TF families.
408
409 **Materials and Methods**
410
Cell culture, stable cell line construction and drug treatment
Human 786-O clear cell renal carcinoma cells were obtained from the Cell Culture Facility at the
University of California at Berkeley, and were cultured at 37°C with 5% CO_2 in 4.5 g/L glucose
DMEM (ThermoFisher, Waltham, MA, #10566016) supplemented with 10% Fetal Bovine Serum
(HyClone, Logan, UT, Cat. #SH30396.03, lot #AE28209315), 1 mM Sodium Pyruvate
(ThermoFisher #11360070) and 100 U/mL Penicillin-Streptomycin (ThermoFisher #15140122).
Cells were subcultured at a ratio of 1:4 to 1:12 every 2 to 4 days for no longer than 30 days.
Phenol red-free DMEM (ThermoFisher, #21063029) supplemented with 10% Fetal Bovine Serum,
1 mM Sodium Pyruvate and 100 U/mL Penicillin-Streptomycin was used for imaging.
420
Stable cell lines expressing the exogenous gene product (supplement table 1) were generated by
PiggyBac transposition and antibiotic selection. The gene of interest was cloned into a PiggyBac
vector which also co-expresses a puromycin resistant gene using Gibson Assembly and
confirmed by Sanger sequencing. Cells were transfected by nucleofection using the Lonza Cell
Line Nucleofector® Kit V (Lonza, Basel, Switzerland, #VVCA-1003) and the Amaxa Nucleofector
II device. For each transfection, cells were plated 1-2 days before nucleofection in a 15-cm dish,
and reached approximately 50-70% confluency on the day of nucleofection, which equals to
approximately 3-4 million cells. 2 µg of PiggyBac plasmid was co-transfected with 1 µg of
SuperPiggyBac transposase vector with the T-020 program according to manufacturer's protocol.
Transfected cells were cultured for 24-48 hours before changing to selection media. Cells were
then selected for 14 days with 1 µg/ml puromycin (ThermoFisher #A1113803) and stable cell lines
were maintained in selection media for up to 30 days of culturing.
433
For drug treatment, 100 mM Belzutifan stock solution was prepared by dissolving Belzutifan
power (CAS No: 1672668-24-4, MedChemExpress, Monmouth Junction, NJ, Cat. #HY-125840)
in DMSO (Sigma, St. Louis, MO, #D2650), and was diluted 1:500,000 in growth media to the final
concentration of 0.2 µM. The same volume of DMSO (0.0002%) is used in the reference group
as control. Cells were treated for 24 hours in either Belzutifan or DMSO alone before imaging.

For dosage-dependent assays in Figure S4, DMSO amount was kept the same (0.0002%) for all
drug concentrations.
441

CRISPR/Cas9-mediated genome editing
Knock-in cell lines were generated as previously described (Hansen et al., 2017) with the following
changes. For each editing case, we designed 3 sgRNAs using CRISPOR (Concordet and
Haeussler, 2018). For each guide/donor pair, approximately 4 million 786-O cells were
nucleofected with 3.75 μg of donor plasmid and 1.25 μg of sgRNA plasmid. 24 hours after
transfection, Venus-positive cells were sorted and cultured for another 5-7 days, then Halo-
positive cells were sorted individually into single wells of 96 well plates. Clones were expanded
and genotyped with two rounds of PCR. The first round used one primer upstream of the left
homologous arm and the other primer downstream of the right homologous arm. The second
round used either of the external primers and a corresponding internal primer located in the
HaloTag coding region. Homozygous clones with the correct genotype, including Halo-HIF-2α
KIN clone A31 and clone B50, Halo-HIF-1β KIN clone A21 and clone B89, were confirmed by
Sanger sequencing and western blotting.
455

Cell preparation and dye labeling for imaging
For fast SPT, cells were grown on sonicated and plasma-cleaned 25-mm circular no 1.5H
precision cover glass (Marienfeld, Germany, 0117650) in 6-well plate. At least one day before
imaging, selective medium (if used) was removed and replaced with non-selective growth medium.
On the day of imaging, cells should be less than 100% confluent. Immediately before imaging,
cells were double labeled with JFX dyes as follows: cells were first incubated for 5 min in 1 ml
growth medium containing JFX 646, at a concentration that only gives approximately 10 detected
molecules per frame in the initial frames to ensure minimum misconnection of trajectories
between detections. This concentration differs from cell line to cell line, ranging from 0.2 – 5 nM,
depending on the expression level of the Halo-fusion protein. After 5 min of incubation, medium
was removed, cells were rinsed in PBS, and incubated for 5 min in 1 ml medium containing JFX
549. The concentration of JFX 549 also varies, usually at 25x the concentration of JFX 646. After
incubation, cells were washed twice for 5 min each, a first time with 2 ml regular growth media,
and a second time with 2 ml phenol red-free growth media, with a quick PBS rinse before each
wash. After wash, coverslip was transferred to Attofluor Cell Chambers (ThermoFisher, #A7816)
with cells facing up and 1 ml phenol red-free medium added to the chamber. For Belzutifan
treatment experiments, Belzutifan or equivalent amount of DMSO was added throughout the

labeling and washing steps (except during PBS rinses), as well as in the final imaging medium,

at the indicated concentration.

Live Cell Single particle tracking

All SPT experiments were carried out on a custom-built microscope as previously described

(Hansen et al., 2017) (McSwiggen et al., 2019). In brief, a Nikon TI microscope is equipped with

a 100x/ NA 1.49 oil-immersion TIRF objective, a motorized mirror, a perfect Focus system, an

EM-CCD camera and an incubation chamber maintained with humidified atmosphere with 5%

CO_2 at 37 °C. All microscope, camera and hardware components were controlled through the

NIS-Elements software (Nikon).

During imaging, samples were excited with 561-nm laser at 1100 mW (Genesis Coherent, Santa

Clara, CA) with emission filter set to Semrock 593/40 nm band-pass filter to locate and focus the

cell nuclei, as well as to adjust laser angle to achieve highly inclined laminated optical sheet (HiLo)

illumination (Tokunaga et al., 2008). An ROI (Regions of Interest) of random size was selected to

fit into the interior of the nuclei but with maximized area. Then the emission filter was switched to

Semrock 676/37 nm bandpass filter while keeping TIRF angle, stage xyz position and ROI the

same. Movies were then taken with 633-nm laser (Genesis Coherent, Santa Clara, CA) at 1100

mW and 1 ms pulse, with camera exposure at 5.48-ms frame rate for 800-1600 frames, until

samples were completely photo-bleached. At least 20 movies (corresponding to 20 cells) were

taken for each sample as one biological replicate on a given day. A total of three biological

replicates on three separate days were collected to produce the final results (>60 cells per cell

line/condition).

SPT data processing

Raw SPT movies were processed with a publicly available single particle tracking package

(https://github.com/alecheckert/quot) to generate trajectory files (.trajs). Generally, it performs

tracking in the following steps: read a frame, find spots in the frame, localize spots to subpixel

resolution, and reconnect spots from consecutive frames into trajectories. Since a non-

photoactivatable dye was used for all SPT experiments, we labeled cells with a dye concentration

that only gives very low spot detection density, which allowed us to track spots since the first

frame. This is important because if the initial frames are filtered due to high localization density,

there might be a bias towards moving molecules, due to the bound molecules being

photobleached and diffusing molecules moving into the focal plane during the later frames.

Although we used very sparse labeling, occasionally there would be frames with high density, to
minimize misconnections due to multiple particles in close proximity, we incorporated a filtering
step where we removed frames with more than 7 detections in the following way. First, we
computed the number of detections per frame. Next, this function was smoothed with uniform
filtering with a kernel width of 21 frames. Finally, we identified frames with fewer than 7 detections
after smoothing and isolated trajectories from these frames. Specifically, the following
configuration was used for all detections and tracking: Image reading and filtering settings: start
= 0, method = "identity", chunk_size = 100; Spot detection settings: method = "llr", k = 1.0, w = 15,
t = 18; Subpixel localization settings: method = 'ls_int_gaussian', window_size = 9, sigma = 1.0,
ridge = 0.001, max_iter = 20, damp = 0.3; Tracking settings: method = 'euclidean',
max_spots_per_frame = 7, pixel_size_um = 0.16, frame_interval = 0.00548, search_radius = 1.0,
max_blinks = 0, min_I0 = 0.0, scale = 7.0.
519
To infer the distribution of diffusion coefficients from experimentally observed trajectories, we
used a publicly available implementation of state arrays (https://github.com/alecheckert/spagl)
(`sample_script_fss.py`), which generates the posterior mean occupations for a state array
evaluated on trajectories across all cells. In all analyses, we used the likelihood function for
regular Brownian with localization error (RBME) (Heckert et al., 2021). Settings were:
frame_interval = 0.00548, pixel_size_um = 0.16, dz = 0.7. Occupations are reported as the
mean of the posterior distribution over state occupations, marginalized on diffusion coefficient.
527
528 To generate RBME likelihood for individual cells, we used the `sample_script_by_file.py`
529 script in the same repository (https://github.com/alecheckert/spagl) (Heckert et al., 2021) with
530 the following settings: frame_interval = 0.00548, dz = 0.7, pixel_size_um=0.16,
531 scale_by_total_track_count = True, scale_colors_by_group = True.
532

Antibodies

The following antibodies were used for ChIP-seq: rabbit polyclonal anti-HIF-2α (Novus Biologicals,
Centennial, CO, #NB100-122), mouse monoclonal anti-HIF-1β (Novus Biologicals, #NB100-124),
rabbit polyclonal anti-V5 (Abcam, Cambridge, UK, #ab9116). The following antibodies were used
for western blotting: rabbit monoclonal anti-HIF-2α (Cell Signaling, Danvers, MA, #D9E3) diluted
at 1:1000, rabbit monoclonal anti-HIF-1β (Cell Signaling, #D28F3) diluted at 1:1000, mouse
monoclonal anti-V5 tag (ThermoFisher, # R960-25) diluted at 1:2500, mouse monoclonal anti-
HaloTag (Promega, Madison, WI, # G9211) diluted at 1:1000, mouse monoclonal anti-TBP

(Abcam, #ab51841) diluted at 1:2500, goat-anti-mouse-HRP (ThermoFisher, #31430) diluted at

1:2000, goat-anti-rabbit-HRP (ThermoFisher, # 31462) diluted at 1:2000.

Western blotting

All western samples were prepared as follows: cells growing in either 6-well plates or 10-cm dish

in log phase were rinsed with PBS twice and lysed on ice in 100-500 ul 2x Sample buffer (80 mM

Tris pH6.8, 2% SDS, 10% Glycerol, 0.0006% Bromophenol blue) containing 280 mM 2-

Mercaptoethanol (Sigma #M7522), 1x Aprotinin (Sigma, #A6279, diluted 1:1000), 1 mM

Benzamidine (Sigma, #B6506), 1x cOmplete™ EDTA-free Protease Inhibitor Cocktail (Sigma,

#5056489001), and 0.25 mM PMSF (Sigma #11359061001). Cell lysates were scraped and

collected into 1.5-ml Eppendorf tubes, incubated at 99 °C with constant shaking, snap frozen in

liquid nitrogen and stored at -80 °C. On the day of western blotting, samples were thawed and

centrifuged at top speed for 5 min at 4 °C. Ten to 15 µl supernatant were loaded on an 8% SDS-

Page gel, ran for 1h at 200 V and 4 °C, and transferred to 0.45-µm nitrocellulose membrane

(Fisher, # 45004031) for 2 hrs at 100V. Membranes were blocked in 10% milk in 0.1% TBS-Tween

for 1 hr at RT, and incubated overnight at 4 °C with primary antibodies diluted in 5% milk in 0.1%

TBS-Tween. After 4 x 5 min washes in 0.1% TBS-Tween, membranes were incubated at RT for

at least 1 h with secondary antibodies diluted in 5% milk in 0.1% TBS-Tween. After 4 x 5 min

washes in 0.1% TBS-Tween, membranes were incubated for 3 min in freshly made Perkin Elmer

LLC Western Lightning Plus-ECL, Enhanced Chemiluminescence Substrate (Fisher,

#509049326), and imaged with a Bio-Rad ChemiDoc imaging system (BioRad, Model No:

Universal Hood III). For reblotting, membranes were immersed in Restore™ Western Blot

Stripping Buffer (Fisher, #21059) for 15 min at RT with shaking, washed 3 x 10 min in 0.1% TBS-

Tween, followed by blocking, antibody incubation and chemiluminescence reaction as described

above.

Chromatin Immunoprecipitation and ChIP-seq library preparation

ChIP was performed as described with few modifications (Testa et al., 2005). Wild type 786-O or

endogenously tagged knock-in clones A31 (V5-Halo-HIF-2α) and A21 (V5-Halo-HIF-1β) were

expanded to two 15-cm dishes and cross-linked 5' at room temperature with 1% formaldehyde-

containing FBS-free medium; cross-linking was stopped by adding PBS-glycine (0.125 M final).

Cells were washed twice with ice-cold PBS, scraped, centrifuged for 10' and pellets were flash-

frozen. Cell pellets were thawed and resuspended in 2 ml of cell lysis buffer (5 mM PIPES, pH

8.0, 85 mM KCl, and 0.5% NP-40, 1 ml/15 cm plate) w/ protease inhibitors and incubated for 10'

on ice. Lysates were centrifuged for 10' at 4000 rpm and nuclear pellets resuspended in 6 volumes

of sonication buffer (50 mM Tris-HCl, pH 8.1, 10 mM EDTA, 0.1% SDS) w/ protease inhibitors,

incubated on ice for 10', and sonicated to obtain DNA fragments around 500 bp in length (Covaris

S220 sonicator, 20% Duty factor, 200 cycles/burst, 150 peak incident power, 10 cycles 30" on

and 30" off). Sonicated lysates were cleared by centrifugation and chromatin (400 μg per antibody)

was diluted in RIPA buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA, 0.5 mM EGTA, 1% Triton X-

100, 0.1% SDS, 0.1% Na-deoxycholate, 140 mM NaCl) w/ protease inhibitors to a final

concentration of 0.8 μg/μl, precleared with Protein G sepharose (GE Healthcare) for 2 hours at

4°C and immunoprecipitated overnight with 4 μg of specific antibodies. About 4% of the

precleared chromatin was saved as input. Immunoprecipitated DNA was purified with the Qiagen

QIAquick PCR Purification Kit, eluted in 33 μl of 0.1X TE (1 mM Tris-HCl pH 8.0, 0.01 mM EDTA)

and analyzed by qPCR together with 2% of the input chromatin prior to ChIP-seq library

preparation (SYBR® Select Master Mix for CFX, ThermoFisher). ChIP-qPCR primer sequences

were as follows:

hWISP1_positive_forward: TGAGGTCAGTGTGGTTTGGT

hWISP1_positive_reverse: ACATGGTCACGTAGCTAGCA

hWISP1_negative_forward: AGTCCCCAGCACATAGAAGG

hWISP1_negative_reverse: GGTTCTGAAGGTGACCGACT

ChIP-seq libraries were prepared using the NEBNext® Ultra™ II DNA Library Prep Kit for Illumina®

(NEB E7645) according to manufacturer instructions with a few modifications. 20 ng of ChIP input

DNA (as measured by Nanodrop) and 25 μl of the immunoprecipitated DNA were used as a

starting material and the recommended reagents' volumes were cut in half. The NEBNext Adaptor

for Illumina was diluted 1:10 in Tris/NaCl, pH 8.0 (10 mM Tris-HCl pH 8.0, 10 mM NaCl) and the

ligation step extended to 30'. After ligation, a single purification step with 0.9X volumes of

Agencourt AMPure XP PCR purification beads (Beckman Coulter A63880) was performed, eluting

DNA in 22 μl of 10 mM Tris-HCl pH 8.0. 20 μl of the eluted DNA were used for the library

enrichment step, performed with the KAPA HotStart PCR kit (Roche Diagnostics KK2502) in 50

μl of total reaction volume (10 μl 5X KAPA buffer, 1.5 μl 10 mM dNTPs, 0.5 μl 10 μM NEB

Universal PCR primer, 0.5 μl 10 μM NEB index primer, 1 μl KAPA polymerase, 16.5 μl nuclease-

free water and 20 μl sample). Samples were enriched with 9 PCR cycles (98 °C, 45"; [98 °C, 15";

60 °C, 10"] x 9; 72 °C, 1'; 4 °C, hold), purified with 0.9 volumes of AMPure XP PCR purification

beads and eluted with 33 μl of 10 mM Tris-HCl pH 8.0. Library concentration, quality and fragment

size were assessed by Qubit fluorometric quantification (Qubit™ dsDNA HS Assay Kit,

Invitrogen^TM Q32851) qPCR and Fragment analyzer™. 12 multiplexed libraries (input, HIF1-β,

HIF1-α and V5 pulldowns in WT 786O cells and A31 and A21 clones) were pooled and sequenced

in one lane on the Illumina HiSeq4000 sequencing platform (50-bp, single end-reads) at the

Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley.

ChIP-seq analysis

ChIP-seq raw reads from WT 786O cells and A31 and A21 endogenously Halo-tagged clones (12

libraries total, 1 replicate per condition) were quality-checked with FastQC and aligned onto the

human genome (hg38 assembly) using Bowtie (Langmead et al., 2009), allowing for two

mismatches (-n 2) and no multiple alignments (-m 1). Peaks were called with MACS2 (--nomodel

--extsize 300) (Zhang et al., 2008) using input DNA as a control. To create heatmaps we used

deepTools (version 2.4.1) (Ramírez et al., 2016). We first ran bamCoverage (--binSize 50 --

normalizeTo1x 2913022398 --extendReads 300 --ignoreDuplicates -of bigwig) and normalized

read numbers to 1x sequencing depth, obtaining read coverage per 50-bp bins across the whole

genome (bigWig files). We then used the bigWig files to compute read numbers across 6 kb

centered on HIF-2α peaks called by MACS2 across all 786O cell lines, subtracted of V5 peaks

called by MACS2 in WT 786O cells (computeMatrix reference-point --referencePoint=TSS --

upstream 3000 --downstream 3000 --missingDataAsZero --sortRegions=no). We sorted the

output matrices by decreasing WT 786O enrichment, calculated as the total number of reads

within a MACS2 called ChIP-seq peak. Finally, heatmaps were created with the plotHeatmap tool

(--averageTypeSummaryPlot=mean --colorMap='Blues' --sortRegions=no).

Luciferase reporter assay

The firefly luciferase reporter gene construct was made by inserting a 3x Hypoxia Responsive

Elements (HREs) from the EPO gene enhancer (sequence:

tcgaagccctacgtgctgtctcacacagcctgtctgacctctcgacctaccggccgttcgaagccctacgtgctgtctcacacagccttct

gatctcgacctaccggccgttcgaagccctacgtgctgtctcacacagcctgtctgacctctcgacctaccggccgt) into the 5' of

the minimal TATA-box promoter in the pGL4.23 [luc2/minP] vector (Promega #E841A). A control

pHRL-TK vector (Promega #E2241) expressing *Renilla* luciferase with an HSV TK promoter was

used as reference to normalize luciferase activity. Cells were co-transfected with 1 ug of firefly

Luciferase vector and 0.1 ug *Renilla* luciferase vector by nucleofection with Lonza Cell Line

Nucleofector® Kit V (Lonza, #VVCA-1003) and the T-020 program in the Amaxa Nucleofector II

device. After nucleofection, cells were resuspended in complete growth medium, and plated into

12-well plates with Belzutifan added to various concentrations as indicated. 24 hours after

nucleofection, cells were lysed and luciferase activity was analyzed with Dual-luciferase Reporter

Assay System (Promega, #E1960) according to manufacturer's protocol. The relative luciferase

activity was calculated by normalizing firefly luciferase activity to the *Renilla* luciferase activity to

control for transfection efficiency.

Datasets and accession numbers

The ChIP-seq data generated in this publication have been deposited in NCBI's Gene Expression

Omnibus (Edgar et al., 2002) and are accessible through GEO Series accession number

GSE183900. SPT raw data are accessible through DOI: 10.5281/zenodo.5559234.

Acknowledgments

We thank Luke Lavis for providing fluorescent HaloTag ligands; the CRL Flow Cytometry Facility

for assistance with cell sorting; the QB3 High Throughput Screening Facility for providing access

to Opera Phenix automated confocal microscope; Dr. Alec Heckert for sharing data analysis

pipeline and usage instructions; Dr. Alec Heckert, Dr. Andrew Belmont, Dr. Max Staller, Dr.

Shasha Chong, Dr. Jiang Xu, and members of Tjian/Darzacq labs for helpful discussions and

critical reading of the manuscript. This work was supported by the NIH grant U54-CA231641-01

(to XD), and the Howard Hughes Medical Institute (to RT). This work used the Vincent J. Coates

Genomics Sequencing Laboratory at UC Berkeley, supported by NIH S10 OD018174

Instrumentation Grant.

Author contributions: YC designed experiments with guidance from XD and RT. YC, CC, GD

and QZ performed experiments. YC drafted the original manuscript. CC, XD and RT reviewed

and edited the manuscript. XD and RT supervised the project.

Competing interests: RT and XD are co-founders of Eikon Therapeutics.

References

Behera V, Evans P, Face CJ, Hamagami N, Sankaranarayanan L, Keller CA, Giardine B, Tan K,
Hardison RC, Shi J, Blobel GA. 2018. Exploiting genetic variation to uncover rules of
transcription factor binding and chromatin accessibility. *Nat Commun* **9**:782.
doi:10.1038/s41467-018-03082-6
Boijja A, Klein IA, Sabari BR, Dall'Agnese A, Coffey EL, Zamudio AV, Li CH, Shrinivas K, Manteiga
JC, Hannett NM, Abraham BJ, Afeyan LK, Guo YE, Rimel JK, Fant CB, Schuijers J, Lee TI,

Taatjes DJ, Young RA. 2018. Transcription factors activate genes through the phase
separation capacity of their activation domains. *Cell* **175**:1842.
doi:10.1016/j.cell.2018.10.042
Brent R, Ptashne M. 1985. A eukaryotic transcriptional activator bearing the DNA specificity of a
prokaryotic repressor. *Cell* **43**:729–736. doi:10.1016/0092-8674(85)90246-6
Brodaczewska KK, Szczylik C, Fiedorowicz M, Porta C, Czarnecka AM. 2016. Choosing the right
cell line for renal cell cancer research. *Mol Cancer* **15**:83. doi:10.1186/s12943-016-0565-
8
Brodsky S, Jana T, Mittelman K, Chapal M, Kumar DK, Carmi M, Barkai N. 2020. Intrinsically
Disordered Regions Direct Transcription Factor In Vivo Binding Specificity. *Mol Cell*
79:459-471.e4. doi:10.1016/j.molcel.2020.05.032
Burdach J, Funnell APW, Mak KS, Artuz CM, Wienert B, Lim WF, Tan LY, Pearson RCM, Crossley
M. 2014. Regions outside the DNA-binding domain are critical for proper in vivo
specificity of an archetypal zinc finger transcription factor. *Nucleic Acids Res* **42**:276–
289. doi:10.1093/nar/gkt895
Cho W-K, Spille J-H, Hecht M, Lee C, Li C, Grube V, Cisse II. 2018. Mediator and RNA polymerase
II clusters associate in transcription-dependent condensates. *Science* **361**:412–415.
doi:10.1126/science.aar4199
Chong S, Dugast-Darzacq C, Liu Z, Dong P, Dailey GM, Cattoglio C, Heckert A, Banala S, Lavis L,
Darzacq X, Tjian R. 2018. Imaging dynamic and selective low-complexity domain
interactions that control gene transcription. *Science* **361**:eaar2555.
doi:10.1126/science.aar2555
Chong S, Mir M. 2021. Towards Decoding the Sequence-Based Grammar Governing the
Functions of Intrinsically Disordered Protein Regions. *J Mol Biol* **433**:166724.
doi:10.1016/j.jmb.2020.11.023
Concordet J-P, Haeussler M. 2018. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome
editing experiments and screens. *Nucleic Acids Res* **46**:W242–W245.
doi:10.1093/nar/gky354
Courey AJ, Tjian R. 1988. Analysis of Sp1 in vivo reveals multiple transcriptional domains,
including a novel glutamine-rich activation motif. *Cell* **55**:887–898. doi:10.1016/0092-
8674(88)90144-4
Edgar R, Domrachev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene expression and
hybridization array data repository. *Nucleic Acids Res* **30**:207–210.
doi:10.1093/nar/30.1.207
Gaur RK. 2014. Amino acid frequency distribution among eukaryotic proteins. *IIOAB J* **5**:6–11.
Gnarra JR, Tory K, Weng Y, Schmidt L, Wei MH, Li H, Latif F, Liu S, Chen F, Duh FM. 1994.
Mutations of the VHL tumour suppressor gene in renal carcinoma. *Nat Genet* **7**:85–90.
doi:10.1038/ng0594-85
Grimm JB, Xie L, Casler JC, Patel R, Tkachuk AN, Falco N, Choi H, Lippincott-Schwartz J, Brown
TA, Glick BS, Liu Z, Lavis LD. 2021. A General Method to Improve Fluorophores Using
Deuterated Auxochromes. *Jacs Au* **1**:690–696. doi:10.1021/jacsau.1c00006
Grossman SR, Zhang X, Wang L, Engreitz J, Melnikov A, Rogov P, Tewhey R, Isakova A,
Deplancke B, Bernstein BE, Mikkelsen TS, Lander ES. 2017. Systematic dissection of

genomic features determining transcription factor binding and enhancer function. *Proc*
Natl Acad Sci **114**:E1291–E1300. doi:10.1073/pnas.1621150114

Hansen AS, Pustova I, Cattoglio C, Tjian R, Darzacq X. 2017. CTCF and cohesin regulate
chromatin loop stability with distinct dynamics. *eLife* **6**:e25776. doi:10.7554/eLife.25776

Hansen AS, Woringer M, Grimm JB, Lavis LD, Tjian R, Darzacq X. 2018. Robust model-based
analysis of single-particle tracking experiments with Spot-On. *eLife* **7**:e33125.
doi:10.7554/eLife.33125

Hanson J, Yang Y, Paliwal K, Zhou Y. 2017. Improving protein disorder prediction by deep
bidirectional long short-term memory recurrent neural networks. *Bioinformatics*
33:685–692. doi:10.1093/bioinformatics/btw678

Heckert A, Dahal L, Tjian R, Darzacq X. 2021. Recovering mixtures of fast diffusing states from
short single particle trajectories. bioRxiv. doi:10.1101/2021.05.03.442482

Hu C-J, Sataur A, Wang L, Chen H, Simon MC. 2007. The N-terminal transactivation domain
confers target gene specificity of hypoxia-inducible factors HIF-1alpha and HIF-2alpha.
Mol Biol Cell **18**:4528–4542. doi:10.1091/mbc.e06-05-0419

Keith B, Johnson RS, Simon MC. 2012. HIF1α and HIF2α: sibling rivalry in hypoxic tumour growth
and progression. *Nat Rev Cancer* **12**:9–22. doi:10.1038/nrc3183

Kribelbauer JF, Rastogi C, Bussemaker HJ, Mann RS. 2019. Low-Affinity Binding Sites and the
Transcription Factor Specificity Paradox in Eukaryotes. *Annu Rev Cell Dev Biol* **35**:357–
379. doi:10.1146/annurev-cellbio-100617-062719

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. *Genome Biol* **10**:R25. doi:10.1186/gb-
2009-10-3-r25

Lim WF, Burdach J, Funnell APW, Pearson RCM, Quinlan KGR, Crossley M. 2016. Directing an
artificial zinc finger protein to new targets by fusion to a non-DNA-binding domain.
Nucleic Acids Res **44**:3118–3130. doi:10.1093/nar/gkv1380

Liu Z, Lavis LD, Betzig E. 2015. Imaging Live-Cell Dynamics and Structure at the Single-Molecule
Level. *Mol Cell* **58**:644–659. doi:10.1016/j.molcel.2015.02.033

Los GV, Encell LP, McDougall MG, Hartzell DD, Karassina N, Zimprich C, Wood MG, Learish R,
Ohana RF, Urh M, Simpson D, Mendez J, Zimmerman K, Otto P, Vidugiris G, Zhu J,
Darzins A, Klaubert DH, Bulleit RF, Wood KV. 2008. HaloTag: A Novel Protein Labeling
Technology for Cell Imaging and Protein Analysis. *ACS Chem Biol* **3**:373–382.
doi:10.1021/cb800025k

Ma J, Ptashne M. 1987. Deletion analysis of GAL4 defines two transcriptional activating
segments. *Cell* **48**:847–853. doi:10.1016/0092-8674(87)90081-X

McSwiggen DT, Hansen AS, Teves SS, Marie-Nelly H, Hao Y, Heckert AB, Umemoto KK, Dugast-
Darzacq C, Tjian R, Darzacq X. 2019. Evidence for DNA-mediated nuclear
compartmentalization distinct from phase separation. *eLife* **8**:e47098.
doi:10.7554/eLife.47098

Mermod N, O'Neill EA, Kelly TJ, Tjian R. 1989. The proline-rich transcriptional activator of
CTF/NF-I is distinct from the replication and DNA binding domain. *Cell* **58**:741–753.
doi:10.1016/0092-8674(89)90108-6

Michel G, Minet E, Mottet D, Remacle J, Michiels C. 2002. Site-directed mutagenesis studies of
the hypoxia-inducible factor-1α DNA-binding domain. *Biochim Biophys Acta BBA - Gene*
Struct Expr **1578**:73–83. doi:10.1016/S0167-4781(02)00484-0

Normanno D, Boudarène L, Dugast-Darzacq C, Chen J, Richter C, Proux F, Bénichou O, Voituriez
R, Darzacq X, Dahan M. 2015. Probing the target search of DNA-binding proteins in
mammalian cells using TetR as model searcher. *Nat Commun* **6**:7357.
doi:10.1038/ncomms8357

Pellanda P, Dalsass M, Filipuzzi M, Loffreda A, Verrecchia A, Castillo Cano V, Thabussot H, Doni
M, Morelli MJ, Soucek L, Kress T, Mazza D, Mapelli M, Beaulieu M-E, Amati B, Sabò A.
2021. Integrated requirement of non-specific and sequence-specific DNA binding in
Myc-driven transcription. *EMBO J* **40**:e105464. doi:10.15252/embj.2020105464

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke T.
2016. deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res **44**:W160-165. doi:10.1093/nar/gkw257

Rankin EB, Biju MP, Liu Q, Unger TL, Rha J, Johnson RS, Simon MC, Keith B, Haase VH. 2007.
Hypoxia-inducible factor-2 (HIF-2) regulates hepatic erythropoietin in vivo. *J Clin Invest*
117:1068–1077. doi:10.1172/JCI30117

Raval RR, Lau KW, Tran MGB, Sowter HM, Mandriota SJ, Li J-L, Pugh CW, Maxwell PH, Harris AL,
Ratcliffe PJ. 2005. Contrasting properties of hypoxia-inducible factor 1 (HIF-1) and HIF-2
in von Hippel-Lindau-associated renal cell carcinoma. *Mol Cell Biol* **25**:5675–5686.
doi:10.1128/MCB.25.13.5675-5686.2005

Rohs R, Jin X, West SM, Joshi R, Honig B, Mann RS. 2010. Origins of specificity in protein-DNA
recognition. *Annu Rev Biochem* **79**:233–269. doi:10.1146/annurev-biochem-060408-
091030

Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ, Hannett NM,
Zamudio AV, Manteiga JC, Li CH, Guo YE, Day DS, Schuijers J, Vasile E, Malik S, Hnisz D,
Lee TI, Cisse II, Roeder RG, Sharp PA, Chakraborty AK, Young RA. 2018. Coactivator
condensation at super-enhancers links phase separation and gene control. *Science*
361:eaar3958. doi:10.1126/science.aar3958

Schödel J, Grampp S, Maher ER, Moch H, Ratcliffe PJ, Russo P, Mole DR. 2016. Hypoxia,
Hypoxia-inducible Transcription Factors, and Renal Cancer. *Eur Urol* **69**:646–657.
doi:10.1016/j.eururo.2015.08.007

Schödel J, Oikonomopoulos S, Ragoussis J, Pugh CW, Ratcliffe PJ, Mole DR. 2011. High-
resolution genome-wide mapping of HIF-binding sites by ChIP-seq. *Blood* **117**:e207–
e217. doi:10.1182/blood-2010-10-314427

Semenza GL. 2012. Hypoxia-Inducible Factors in Physiology and Medicine. *Cell* **148**:399–408.
doi:10.1016/j.cell.2012.01.021

Shen C, Beroukhim R, Schumacher SE, Zhou J, Chang M, Signoretti S, Kaelin WG. 2011. Genetic
and Functional Studies Implicate HIF1α as a 14q Kidney Cancer Suppressor Gene. *Cancer*
Discov **1**:222–235. doi:10.1158/2159-8290.CD-11-0098

Smythies JA, Sun M, Masson N, Salama R, Simpson PD, Murray E, Neumann V, Cockman ME,
Choudhry H, Ratcliffe PJ, Mole DR. 2019. Inherent DNA-binding specificities of the HIF-
1α and HIF-2α transcription factors in chromatin. *EMBO Rep* **20**:e46401.
doi:10.15252/embr.201846401

Srivastava D, Mahony S. 2020. Sequence and chromatin determinants of transcription factor
binding and the establishment of cell type-specific binding patterns. *Biochim Biophys*
Acta Gene Regul Mech **1863**:194443. doi:10.1016/j.bbagrm.2019.194443
Swiatek M, Jancewicz I, Kluebsoongnoen J, Zub R, Maassen A, Kubala S, Udomkit A, Siedlecki JA,
Sarnowski TJ, Sarnowska E. 2020. Various forms of HIF-1α protein characterize the clear
cell renal cell carcinoma cell lines. *IUBMB Life* **72**:1220–1232. doi:10.1002/iub.2281
Testa A, Donati G, Yan P, Romani F, Huang TH-M, Viganò MA, Mantovani R. 2005. Chromatin
immunoprecipitation (ChIP) on chip experiments uncover a widespread distribution of
NF-Y binding CCAAT sites outside of core promoters. *J Biol Chem* **280**:13606–13615.
doi:10.1074/jbc.M414039200
Tjian R, Maniatis T. 1994. Transcriptional activation: a complex puzzle with few easy pieces. *Cell*
77:5–8. doi:10.1016/0092-8674(94)90227-5
Tokunaga M, Imamoto N, Sakata-Sogawa K. 2008. Highly inclined thin illumination enables clear
single-molecule imaging in cells. *Nat Methods* **5**:159–161. doi:10.1038/nmeth1171
Varma S, Cohen HJ. 1997. Co-transactivation of the 3' erythropoietin hypoxia inducible
enhancer by the HIF-1 protein. *Blood Cells Mol Dis* **23**:169–176.
doi:10.1006/bcmd.1997.0134
Völkel S, Stielow B, Finkernagel F, Stiewe T, Nist A, Suske G. 2015. Zinc Finger Independent
Genome-Wide Binding of Sp2 Potentiates Recruitment of Histone-Fold Protein Nf-y
Distinguishing It from Sp1 and Sp3. *PLoS Genet* **11**:e1005102.
doi:10.1371/journal.pgen.1005102
Wallace EM, Rizzi JP, Han G, Wehn PM, Cao Z, Du X, Cheng T, Czerwinski RM, Dixon DD, Goggin
BS, Grina JA, Halfmann MM, Maddie MA, Olive SR, Schlachter ST, Tan H, Wang B, Wang
K, Xie S, Xu R, Yang H, Josey JA. 2016. A Small-Molecule Antagonist of HIF2α Is
Efficacious in Preclinical Models of Renal Cell Carcinoma. *Cancer Res* **76**:5491–5500.
doi:10.1158/0008-5472.CAN-16-0473
Warnecke C, Zaborowska Z, Kurreck J, Erdmann VA, Frei U, Wiesener M, Eckardt K-U. 2004.
Differentiating the functional role of hypoxia-inducible factor (HIF)-1alpha and HIF-
2alpha (EPAS-1) by the use of RNA interference: erythropoietin is a HIF-2alpha target
gene in Hep3B and Kelly cells. *FASEB J Off Publ Fed Am Soc Exp Biol* **18**:1462–1464.
doi:10.1096/fj.04-1640fje
Wei M-T, Chang Y-C, Shimobayashi SF, Shin Y, Strom AR, Brangwynne CP. 2020. Nucleated
transcriptional condensates amplify gene expression. *Nat Cell Biol* **22**:1187–1196.
doi:10.1038/s41556-020-00578-6
Wenger RH, Stiehl DP, Camenisch G. 2005. Integration of oxygen signaling at the consensus
HRE. *Sci Signal* **2005**:re12. doi:10.1126/stke.3062005re12
Wu D, Potluri N, Lu J, Kim Y, Rastinejad F. 2015. Structural integration in hypoxia-inducible
factors. *Nature* **524**:303–308. doi:10.1038/nature14883
Xiang L, Chen K, Yan R, Li W, Xu K. 2020. Single-molecule displacement mapping unveils
nanoscale heterogeneities in intracellular diffusivity. *Nat Methods* **17**:524–530.
doi:10.1038/s41592-020-0793-0
Xu R, Wang K, Rizzi JP, Huang H, Grina JA, Schlachter ST, Wang B, Wehn PM, Yang H, Dixon DD,
Czerwinski RM, Du X, Ged EL, Han G, Tan H, Wong T, Xie S, Josey JA, Wallace EM. 2019.
3-[(1S,2S,3R)-2,3-Difluoro-1-hydroxy-7-methylsulfonylindan-4-yl]oxy-5-

fluorobenzonitrile (PT2977), a Hypoxia-Inducible Factor 2α (HIF-2α) Inhibitor for the
Treatment of Clear Cell Renal Cell Carcinoma. *J Med Chem* **62**:6876–6893.
doi:10.1021/acs.jmedchem.9b00719
Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown
M, Li W, Liu XS. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**:R137.
doi:10.1186/gb-2008-9-9-r137
855

Fig 1. Endogenous tagging of HIFs in 786-O clear cell renal cell carcinoma (ccRCC) cells for fast single particle tracking (fSPT). (A) Schematic showing the similar domain organizations of HIFs (top) and the HRE-bound HIF α/β dimers (bottom). Disordered regions are represented as wavy lines. **(B)** Generation of Halo-KIN clones in the HIF-1α negative 786-O ccRCC line. Top: Halo-tagging scheme of HIF-2α (left) and HIF-1β (right). Bottom: Western blot of wild-type (WT) 786-O cells and homozygously tagged knock-in clones (A31 and A21). See supplemental figure 2 for uncropped images. **(C)** Halo-tagged HIF-2α and HIF-1β show predominant nuclear localization. Top: schematic of labeling Halo-tagged proteins in live cells with cell-permeable Halo-binding JFX646 dye. Bottom: representative images of Halo-HIF-2α (left) and Halo-HIF-1β (right)

clones labeled with 500 nM JFX646 **(D)** representative images showing the same cell labeled with

a high concentration of JFX549 dye for localizing the nucleus in one channel (left) and labeled

sparsely with JFX646 dye for tracking individual molecules in another channel (right). **(E)**

Graphical illustration of fSPT capturing trajectories of moving particles. Top: highly inclined and

laminated optical sheet illumination (HiLo). Bottom: Illumination and camera sequence with

corresponding particle position at each frame (solid magenta dots). Particle's past positions

(dashed magenta circles) are connected with dotted magenta lines to show the particle's

trajectory. **(F)** Actual data showing detection of Halo-HIF-2α protein molecules at 5.48 ms frame

rate.

 Fig 2. fSPT sensitively detects molecules in a range of states. (A-B) Likelihood of diffusion

coefficients based on a model of regular Brownian motion with localization error (RBME) (Heckert

et al., 2021), for (A) Halo-HIF-2α, clone A31 and (B) Halo-HIF-1β, clone A21, with drawing

illustrating bound and different hypothetical moving states: complexes, dimer and monomer. Each

row represents data collected from one cell. 0.1 μm^2/sec is used as the cut-off for bound versus

free. **(C)** Left: proportion of molecules as a function of their diffusion coefficients (posterior mean

occupations for a state array (Heckert et al., 2021)) evaluated on trajectories across all cells

measured for each KIN line (Halo-HIF-2α, clone A31 and Halo-HIF-1β, clone A21). Compared to

HIF-2α, HIF-1β has less bound fraction (grey versus yellow shaded areas) and faster diffusion

coefficient (grey versus yellow arrows). Right: Summary of the bound fraction for the two clones.

Each bar represents the averaged value from three independent measurements on different days

(black dots). **(D-E)** Over-expressing HIF-2α, but not a dimerization mutant form, in the Halo-HIF-

1β knock-in (KIN) line increases HIF-1β binding and decreases its diffusion coefficient. (D)

schematic illustrating the parental Halo-HIF-1β KIN cells (grey background) and cells stably over-

expressing (O.E.) either the wild type (darker blue background) or a dimerization mutant (DM,

black crosses, lighter blue background) form of HIF-2α. (E) Left: Proportion of molecules as a

function of diffusion coefficient measured for HIF-1β in Halo-HIF-1β KIN cells (grey) and in Halo-

HIF-1β KIN cells overexpressing HIF-2α (WT, dark blue background, or dimerization mutant (HIF-

2αDM), light blue). Shaded areas indicate bound fraction. Right: Bar plot of the average value

(bar height) of the bound fraction calculated from three independent measurements (black dots)

for each condition.

Fig 3. HIF-α increases HIF-1β binding and decreases HIF-1β diffusion coefficient through dimerization, in an IDR-dependent manner. (A) Schematic of evaluating Halo-HIF-1β behavior with genetic and small-molecule perturbation. Parental Halo-HIF-1β knock-in (KIN) cells (grey background) and cells stably overexpressing (O.E.) either a certain form of HIF-α (WT or domain swap, HIF-1α, red, HIF-2α, orange. Disordered regions are represented as wavy lines.) (various colored background) or a 3xFLAG tag only control (blue background) are used, with and without 0.2 μM Belzutifan (HIF-2α/1β dimerization inhibitor) treatment. **(B)** Proportion of Halo-HIF-1β molecules as a function of diffusion coefficient measured in various conditions outlined in (A) Top row: DMSO only, showing overexpressing α subunit can change HIF-1β behavior. Cells over-

expressing the α subunit variants containing HIF-2α disordered region (orange curly line) have a

stronger effect (middle, HIF-2α and HIF-1α/2α,) compared to those containing HIF-1α disordered

region (right, HIF-1α and HIF-2α/1α). Middle and Bottom rows: proportions of HIF-1β as a function

of diffusion coefficient, measured in each of the 6 cell lines with Belzutifan treatment are compared

to the DMSO control. Changes caused by overexpressing an α subunit can be specifically

reverted by Belzutifan treatment for cell lines expressing an α subunit variant that contains the

HIF-2α structured domain (orange globule). **(C)** Summary of the average bound fractions for all

12 conditions, with black dots indicating values from each of the three individual measurements.

 Fig 4. The IDR governs HIF-α molecular dynamics and binding characteristics. (A)
Schematic representation of different HIF (WT and domain-swapped) being weakly and stably
overexpressed with an L30 promoter and tracked in WT 786-O cells. **(B)** Proportion of molecules
as a function of diffusion coefficient for every tracked protein in (A). Top: overlapping distribution
curves shows almost identical behavior between Halo-HIF-2α (dark blue curve) and Halo-HIF-
1α/2α (light blue curve). Middle: similar behavior between Halo-HIF-1α (purple curve) and Halo-

HIF-2α/1α (yellow curve), bottom: overlay of all four curves shows very different behavior between

proteins containing 1α versus 2α IDR. **(C)** Bar plot comparing the average bound fraction for cells

in (B), with black dots indicating values from three independent measurements.

Fig 5. HIF-α IDR alone is not sufficient for binding. (A-C) Mutation in DBD reduces the bound fraction for both HIF-α isoforms. (A) Schematic representation of weakly overexpressing and tracking wild type and DBD mutant (R27E or R30E, black cross) forms of HIF-1α or 2α, using the same L30 expression system as in Fig 5. (B) Proportion of molecules as a function of diffusion coefficient for tracked protein listed in (A). (C) Bar plot summarizes the average bound fraction (height of the bar) of three independent measurements (black dots). **(D-F)** Mutations in the dimerization domain reduce the bound fraction for both HIF-α isoforms but do not change their diffusion coefficient. (D) Schematic representation of weakly over-expressing and tracking wild type and dimerization-mutant (DM, two black crosses) forms of Halo-HIF-1α or -2α, using the L30 expression system. (E) Proportion of molecules as a function of diffusion coefficient for tracked

931 protein in (D). (F) Summary of the average bound fraction for all four proteins with black dots

932 indicating values from three independent measurements.

Fig 6. A model for IDR-mediated nuclear search and chromatin binding. (A) The HIF-α IDR determines its slow motion of both the HIF-α monomer and HIF-α/β dimer, likely by HIF-α IDR mediated interactions with nuclear macromolecules. For HIF-α, the IDR thus determines its slow motion regardless of its dimerization status. For HIF-1β, dimerization slows it down due to extra interactions (yellow and red clouds) brought by HIF-α IDR. **(B)** As an obligated dimer, the DBD and the dimerization domain are both necessary for HIF binding, but the IDR determines the degree of binding, possibly via its interaction with nearby macromolecules, including other proteins and/or nucleic acids (DNA and/or RNA).

Fig S1. Domain analysis of HIFs. (A) Disorder propensity of HIF-1α, -2α and -1β, predicted by SPOT-Disorder (Hanson et al., 2017). High values indicate higher disorder propensity. **(B)** HIF-1α and HIF-2α IDRs have different amino acid composition biases. The relative enrichment for each amino acid is calculated by comparison with the average amino acid composition of non-membrane proteins in vertebrates (Gaur, 2014), and color coded, with red indicating relative high enrichment and blue indicating relative depletion. The IDR segments between the N-TAD and C-TAD are especially different between HIF-1α and HIF-2α in terms of acidity (bottom).

Fig S2. Verification of endogenous tagging of HIFs in 786-O ccRCC cells. (A) (B) (C)
Uncropped images of western blot of wild-type (WT) 786-O cells and homozygously tagged

knock-in clones (*). (A) HIF-2α protein is detected in WT and various KIN clones with a Rabbit
monoclonal antibody against HIF-2α (Cell Signaling, D9E3), membranes were stripped and
reblotted for TBP for loading control. (B) HIF-1β protein is detected in WT and various KIN clones
with a Rabbit monoclonal antibody against HIF-1β (Cell Signaling, D28F3), membranes were
stripped and reblotted for TBP for loading control. (C) Various HIF-2α and HIF-1β KIN clones are
loaded on the same gel and probed for tagged protein levels using an anti-Halo antibody (top);
the membrane was stripped and reblotted for TBP for loading control, then stripped again and
reblotted for V5 tag. Halo-HIF-1β is generally expressed at a much higher level than Halo-HIF-2α,
as shown by the detected Halo or V5 level. **(D)** HeatMap of the ChIP-seq experiments comparing
binding profiles of HIF-2α and HIF-1β in WT or KIN clones. ChIP-Seq read counts (Reads Per
Genomic Content) are plotted at MAC2-called HIF-2α peak regions (across all cell lines and
subtracted of V5 peaks called in WT 786-O cells) centered around the peak.

Fig S3. HIF-1β behavior is distinct from HIF-2α and changes as 1β-to-2α stoichiometry changes. (A) RBME likelihood of diffusion coefficient in individual cells for four Halo-HIF-2α or Halo-HIF-1β KIN clonal lines. **(B)** Top and middle: proportion of molecules as a function of their diffusion coefficients (posterior mean occupations for a state array) evaluated on trajectories across all cells measured for additional Halo-HIF KIN line showing reproducible results in different KIN clones (Top, clone A31 and B50 for Halo-HIF-2α KIN and middle, clone A21 and B89 for Halo-HIF-1β KIN). Data for clone A21 and A31 are replotted from Figure 2. Bottom: Summary of

the bound fraction for all four cell lines. Each bar represents the average of three independent

measurements on different days (black dots). **(C)** RBME likelihood of diffusion coefficient obtained

for individual cells in either Halo-HIF-1β KIN cells (top) or Halo-HIF-1β KIN cells overexpressing

HIF-2α (WT, middle or DM, bottom).

Figure S4. Dosage-dependent inhibition of HIF-2α binding and activity by Belzutifan. (A)

Molecular structure of Belzutifan (PT-2977) and schematic illustration of its function to specifically

block dimerization between HIF-2α/1β but not HIF-1α/1β. **(B)** Luciferase assay using an HRE-

containing reporter confirms Belzutifan dosage-dependent inhibition of HIF-2α activity. Error bars

represent SE. **(C)** Proportion of molecules as a function of diffusion coefficient for HIF-2α

measured in Halo-HIF-2α KIN line clone A31 treated with different concentrations of Belzutifan.

DMSO levels are kept the same for all conditions at 0.0002%. **(D)** The average fraction bound

decreases (top) and the average peak position diffusion coefficient increases (bottom) at

increasing doses of Belzutifan. Individual measurements are indicated as black dots. **(E)** RBME

likelihood of diffusion coefficient in individual cells for all six drug dosages.

Figure S5. Additional data for figure 3 showing HIF-α without the IDR is not able to

increase HIF-1β binding or decrease its diffusion coefficient (A) Schematic of evaluating

Halo-HIF-1β behavior while overexpressing only the N-terminal structured region (NT) of HIF-α

in combination with small molecule perturbation. Parental Halo-HIF-1β KIN cells (grey

background, same as in figure 3) and cells stably overexpressing (O.E.) either HIF-2α NT

(orange in yellow background) or HIF-1α NT (red in purple background) are used, with and

without treatment with 0.2-µM Belzutifan. **(B)** Proportion of HIF-1β molecules as a function of

diffusion coefficient measured in various cells outlined in (A). Left: treated with DMSO only,

comparing HIF-1β behavior in the parental cells (black dashed curve, same as in figure 3) and

in cells expressing either HIF-1α NT (purple curve) or HIF-2α NT (yellow curve). Middle and

right: proportions of HIF-1β as a function of diffusion coefficient, measured in cells expressing

either HIF-1α NT (right) or HIF-2α NT (middle) with Belzutifan treatment (grey curves),

compared to the DMSO control (purple or yellow curves). **(C)** Summary of the average bound

fractions for conditions in (B), with black dots indicating values from each of the three individual

measurements.

Figure S6. L30 weak expression system is able to recapitulate the endogenous protein

behavior. (A) Schematic of the L30 overexpression system where Halo-HIF-2α is weakly

expressed with an L30 promoter in wild-type 786-O cells (left) and Knock-in cells where HIF-2α

is endogenously tagged and expressed (right). **(B)** Comparison between endogenous Halo-HIF-

2α (KIN) and Halo-HIF-2α over-expressed with L30 promoter (L30), showing very similar

behavior. Left: Proportion of Halo-HIF-2α as a function of diffusion coefficient and the

corresponding bound fraction (shaded area). Right: Bar plot summarize the average bound

fraction with three independent measurements (black dots).

Supplementary Table 1. Constructs for stable cell lines

Name	promoter	Gene product	Short name in the paper	Appeared in
PB EF1a 3XF EX-MCS IRES Puro	EF1a	3xFLAG tag	3xF	Fig. 3
PB EF1a 3XF-GDGAGLIN-hEPAS1 IRES Puro	EF1a	HIF-2α N-terminally fused with 3xFLAG tag through a short peptide linker sequence (GDGAGLIN)	HIF-2α	Fig. 2 Fig. 3 Fig. S3
PB EF1a 3XF-GDGAGLIN-hEPAS1_R171A-V192D IRES Puro	EF1a	HIF-2α dimerization mutant (R171A-V192D) N-terminally fused with 3xFLAG tag through a short peptide linker sequence (GDGAGLIN)	HIF-2α DM	Fig. 2 Fig. S3
PB EF1a 3XF-EPAS1_365 IRES Puro	EF1a	The N terminal region of HIF-2α (aa 1-365) N-terminally fused with 3xFLAG tag	HIF-2α NT	Fig. S5
PB EF1a 3XF-EPAS1_365-364_HIF1a IRES Puro	EF1a	HIF-2α/1α chimera protein (aa 1-365 of HIF-2α and aa 364-826 of HIF-1α) N-terminally fused with 3xFLAG tag	HIF-2α/1α	Fig. 3
PB EF1a 3XF-GDGAGLIN-hHIF1a IRES Puro	EF1a	HIF-1α N-terminally fused with 3xFLAG tag through a short peptide linker sequence (GDGAGLIN)	HIF-1α	Fig. 3
PB EF1a 3XF-HIF1a_363 IRES Puro	EF1a	The N terminal region of HIF-1α (aa 1-363) N-terminally fused with 3xFLAG tag	HIF-1α NT	Fig. S5
PB EF1a 3XF-HIF1a_363-366_EPAS1 IRES Puro	EF1a	HIF-1α/2α chimera protein (aa 1-363 of HIF-1α and aa 366-870 of HIF-2α) N-terminally fused with 3xFLAG tag	HIF-1α/2α	Fig. 3
PB L30prom V5-Halo-GDGAGLIN-hEPAS1 IRES Puro	L30	HIF-2α N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN)	Halo-HIF-2α (L30)	Fig. 4 Fig. S6
PB L30prom V5-Halo-GDGAGLIN-hEPAS1_365-364_HIF1a IRES Puro	L30	HIF-2α/1α chimera protein (aa 1-365 of HIF-2α and aa 364-826 of HIF-1α) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-2α/1α (L30)	Fig. 4
PB L30prom V5-Halo-GDGAGLIN-hHIF1a IRES Puro	L30	HIF-1α N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN)	Halo-HIF-1α (L30)	Fig. 4
I_PB L30prom V5-Halo-GDGAGLIN-	L30	HIF-1α/2α chimera protein (aa 1-363 of HIF-1α and aa 366-	Halo-HIF-1α/2α (L30)	Fig. 4

HIF1A_363-366_EPAS1 IRES Puro		870 of HIF-2α) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).		
PB L30prom V5-Halo-GDGAGLIN-hEPAS1 IRES Puro_R27E	L30	HIF-2α DBD mutant (R27E) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-2α, R27E (L30)	Fig. 5
PB L30prom V5-Halo-GDGAGLIN-hHIF1a IRES Puro_R30E	L30	HIF-1α DBD mutant (R30E) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-1α, R30E (L30)	Fig. 5
PB L30prom V5-Halo-GDGAGLIN-hEPAS1_R171A-V192D IRES Puro	L30	HIF-2α dimerization mutant (R171A-V192D) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-2α DM (L30)	Fig. 5
PB L30prom V5-Halo-GDGAGLIN-hHIF1a_R170A-V191D IRES Puro	L30	HIF-1α dimerization mutant (R170A-V191D) N-terminally fused with V5-HaloTag through a short peptide linker sequence (GDGAGLIN).	Halo-HIF-1α DM (L30)	Fig. 5

1009